Effects of multiple cell regulators on curli gene expression in

Escherichia coli

Maryia Ratnikava[1], Olga Lamprecht[1], Victor Sourjik[1]*

[1]Max Planck Institute for Terrestrial Microbiology and Center for Synthetic Microbiology (SYNMIKRO),

Karl-von-Frisch Strasse 14, 35043 Marburg, Germany

*Corresponding author: Victor Sourjik, victor.sourjik@synmikro.mpi-marburg.mpg.de

Abstract

Curli amyloid fibers is the key protenacious component of extracellular matrix in *Escherichia coli*. Regulation of curli expression is highly complex and depends on the cellular response to diverse environmental conditions. Although many genetic determinants of curli production are known, previous studies were performed under diverse conditions and using different strains of *E. coli* or *Salmonella*. Furthermore, the origin of bimodal curli expression in bacterial population remains unknown. Here, we systematically investigated the role of multiple cell factors in expression of curli structural genes *csgBA* in unstructured planktonic *E. coli* culture. We observed that multiple regulators involved in regulation of stress response, cell motility, cell physiology and metabolism, as well as in formation of extracellular matrix and maintenance of DNA architecture modulated expression of curli by either promoting or repressing activity of *csgBA*. We further elucidated which regulators act upstream of the master transcription factor CsgD and which are crucial for bimodality of curli gene expression. We also investigated the impact of individual diguanylate cyclases on the regulation of the *csgBA* activity. This study provides an overview of regulation of curli gene expression in planktonic *E. coli* culture in the absence of any microenvironmental gradients.

26 **Keywords:** curli, transcriptional control, c-di-GMP, motility, RpoS, bimodality

Introduction

Most bacteria are able to produce extracellular matrix that promotes formation of multicellular

structures, giving protection against adverse environmental factors [1-3]. Curli are the major

protenacious component of the extracellular matrix produced by *Escherichia coli* [4-8]. They are

highly stable to degradation, thus supporting environmental cell persistence and cell stress survival

[5, 6, 9]. The formation of curli is known to be triggered by a combination of suboptimal growth

conditions and environmental stresses, such as slow growth, nutrient starvation, low temperatures,

low osmolarity etc [2, 10-15].

In *E. coli*, genes required for curli formation are organized in two oppositely transcribed

operons *csgBAC* and *csgDEFG* [2, 12, 15]. They are separated by a 754 bp long intergenic non-

coding region, which is one of the largest and most heavily regulated in *E. coli* [5, 16], with curli

expression dependent on the cellular response to multiple environmental conditions [8, 11, 13, 15,

17]. The *csgBAC* operon controls production of curli fibers, while the *csgDEFG* operon encodes a

master curli regulator CsgD and accessory proteins required for assembly and secretion of the curli

structural subunits CsgB and CsgA [5, 9, 18, 19]. CsgD activates expression of curli structural

genes as well as its own expression by binding to multiple sites within the *csg* region [9, 20]. The

expression of curli genes depends on the general stress response regulator RpoS and it is further

known to be affected by multiple transcriptional factors, small RNAs (sRNAs) and second

messenger bis-(3′–5′)-cyclic di-GMP (c-di-GMP) [2, 5, 14, 15, 17, 21-31]. A number of

transcription factors have been reported to bind within the *csg* region and collaboratively modulate

expression of curli genes from σ^{70}-and σ^{S}-dependent promoters [16, 23, 27, 32, 33], including those

associated with metabolic transitions, two-component signaling and flagellar gene expression, as

well as regulators maintaining DNA architecture (IHF and H-NS) [2, 8, 16, 33-37].

Intracellular c-di-GMP concentration, another key regulator of curli gene expression, is itself

under complex control by a set of diguanylate cyclases (DGCs) and phosphodiesterases (PDEs) [29,

38, 39]. In total, 14 DGCs have been identified in non-pathogenic *E. coli* strains, including two

degenerate proteins that lack the catalytic function [39, 40]. Curli gene expression is primarily

modulated by DgcE and DgcM [28, 29, 31, 41, 42]. Elevated c-di-GMP levels promote expression

of curli by derepressing MlrA, the transcriptional activator of *csgD* [28, 31]. Other active DGCs,

including DgcC, DgcP, and the degenerate DGC CsrD, have been shown to further regulate CsgD

activity in *E. coli* [39].

Expression of curli within a population is restricted to a group of cells, resulting in formation

of curli-producing and non-producing cells [43-47]. Heterogeneous production of curli in

macrocolony and submerged biofilms could be driven by presence of microenvironmental and

metabolic gradients [43, 48-50]. However, bimodal curli expression is also observed in an isogenic

well-stirred planktonic culture of *E. coli* [43, 46], suggesting that bimodality is an inherent property

of curli gene activation.

Here, we systematically assessed how multiple cellular regulators and DGCs contribute to

expression of the *csgBA* genes in unstructured planktonic *E. coli* culture grown under continuous

mixing and thus in the absence of any gradients or biofilm formation. We confirmed involvement of

several known curli regulators and demonstrated impact of several regulators that were not

previously shown to control curli expression. In contrast, other previously proposed regulators of

curli expression showed no impact under our growth conditions. Furthermore, we classified which

regulators act upstream of *csgD* and uncovered a set of regulators crucial for bimodality of

expression of curli structural genes in absence of microenvironmental gradients. Our study provides

a systematic overview of curli regulation in unstructured planktonic *E. coli* culture.

Materials and methods

Bacterial strains and plasmids

All strains and plasmids used in this study are listed in S1 Table. An RpoS$^+$ derivative of *E. coli* W3110 [51] that was engineered to encode a chromosomal transcriptional superfolder green fluorescent protein (sfGFP) reporter downstream of the *csgA* gene (VS1146) [44] was used as the wild-type (WT) strain.

Majority of gene deletions were obtained with the help of P1 phage transduction as described in [52] with some modifications. Strains of the Keio collection [53] were used as donors for the transfer of the kanamycin resistance cassette, which disrupted the coding sequence leaving intact only start codon and last 21 nucleotides (except for *dgcO* deletion, where first 342 and last 21 nucleotides were retained). Deletion of *dgcN* led to the in-frame addition of the 240 nucleotides encoding for a part of Tn5 transposase to the upstream located *yfiR* gene. In the case of *csgD*, the deletion strain was constructed by lambda red recombination method, with first 200 nucleotides and last 30 nucleotides being kept to avoid possible disruption of the common curli gene regulatory region. In the strains that showed differences in curli expression from WT (except for Δ*ihfA*), the kanamycin resistance cassette was eliminated using FLP recombinase as described in [54] to confirm absence of polar effects of the cassette [55]. Obtained gene deletions were confirmed by PCR with primers flanking the corresponding genes followed by Senger sequencing.

For complementation analysis of curli-expression deficient strains, *csgD* gene was cloned into the pTrc99A vector by restriction-ligation [56] and transformed in cells using TSS method as described in [57] with modifications.

Growth conditions of cultures

To analyze *csgBA* expression levels and fraction of curli positive cells by flow cytometry approach, *E. coli* cultures were inoculated from lysogeny broth (LB) agar plates (10 g tryptone, 5 g yeast extract, 5 g NaCl, 15g Bacto Agar per liter; pH 7.0) supplemented with 50 µg/ml kanamycin where

necessary and grown in 5 ml of LB medium at 37°C at 200 rpm in 100 ml flasks in a rotary shaker

overnight (14-16 h). Obtained cultures were diluted 1:100 in 5 ml of tryptone broth (TB) medium

(10 g tryptone, 5 g NaCl per liter; pH 7.0) and grown at 30°C in a rotary shaker at 200 rpm in 100

ml flasks for a time required to detect maximum *csgBA* expression levels (appr. 12-34 h depending

on the strain).

To analyze *csgBA* expression levels and growth curves (OD_{600}) by plate reader analysis,

E. coli cultures were inoculated from LB agar plates supplemented with 50 µg/ml kanamycin where

necessary and grown in liquid LB at 37°C at 200 rpm in 100 ml flasks in a rotary shaker overnight

(14-16 h). Obtained cultures were diluted 1:100 in TB medium and incubated at 30°C in INFINITE

200 PROplate reader (Tecan Group Ltd., Switzerland) for 24-25 h.

For complementation analysis, overnight cultures were inoculated from LB agar plates

supplemented with 100 µg/ml ampicillin and grown in 5 ml of TB supplemented with 100 µg/ml

ampicillin and indicated levels of inducer isopropyl β-d-1-thiogalactopyranoside (IPTG) at 30°C at

200 rpm in 100 ml flasks in a rotary shaker overnight (18-26 h).

Flow cytometry analysis

Flow cytometry method was used for sfGFP expression measurements. Aliquots of 40-70 µl of

bacterial cultures were mixed with 2 ml of PBS (0.14 mM NaCl, 2.7 mM KCl, 1.5 mM KH_2PO_4,

8.1 mM Na_2HPO_4 per liter; pH 7.0) and obtained samples were subjected to the flow cytometry

analysis within next 30 min. Before measurements the probes were vigorously vortexed to disrupt

cell clumps and then immediately subjected to BD LSRFortessa Sorp cell analyzer (BD

Biosciences, Germany) using 488-nm laser. 50 000 individual cells were analyzed in each

experimental run. Recorded events were further filtered to exclude cell duplets and debris with the

help of forward scatter (FSC) and side scatter (SSC) plots (SSC-H vs. SSC-W and FSC-A vs. SSC-

A). Data were analyzed using FlowJo software version v10.7.1 (FlowJo LLC, Ashland, OR, US).

The gate used to differentiate subpopulation of negative cells from curli-positive was defined as

corresponding fraction of curli-negative cells in W3110 RpoS+ strain carrying no *csgBA* genomic

reporter.

Statistical analysis

Expression of *csgBA* genes and fraction of curli-positive cells were statistically analyzed using

ordinary one-way ANOVA followed by Dunnett's test to run multiple comparisons between WT

and deletion strains. The significance level (α) was set to 0.05. The adjusted p-value between 0.01 and

0.05 is indicated with one asterisk (*), between 0.01 and 0.001 with two asterisks (**), between 0.001 and

0.0001 with three asterisks (***), $p < 0.0001$ with four asterisks (****). The statistical analysis was

performed in GraphPad Prism version 10.0.0 for Windows (GraphPad Software, Boston, Massachusetts

USA).

Results

Deletions of multiple regulator genes affect expression of curli fibers

In order to systematically investigate the role of different cellular factors in the regulation of curli

structural genes in *E. coli*, we utilized a recently described strain that encodes a chromosomal GFP

reporter as a part of the *csgBA* operon [44, 46]. We first constructed a library of 32 mutants in this

strain, each carrying a deletion of a single gene selected based on its known involvement into

extracellular matrix formation, control of cell stress response, metabolic transitions as well as

maintenance of DNA architecture and cross-regulation between motility and curli production. We

next measured activity of the *csgBA* reporter in planktonic cultures of these mutants using flow

cytometry. Consistent with previous findings, expression of the *csgBA* genes was bimodal, with

only a fraction of cells (~80%) activating GFP reporter in the overnight culture of the wild-type

(WT) strain background (Fig. 1A) [46]. Half of the obtained deletion mutants showed the *csgBA*

expression similar to that of WT (Fig. 1B-C and Fig. S1). Notably, these also included genes

encoding regulators that were previously suggested by *in vivo* and *in vitro* experiments to bind to

the *csg* intergenic region and/or modulate curli expression, either positively (*basR, bolA, rcdA, rstA*)

[16, 58-60] or negatively (*btsR, mqsR, cpxR, rstA*) [16, 33, 35, 36, 61-63]. Similarly, deletion of

ariR, *fnr*, *phoP* and *gadE* had no impact on the average *csgBA* expression levels (Fig. 1B), although

they led to a moderate decrease of number of curli-expressing cells, whereas *gadE* – to a slight

increase (Fig. S2).

Four mutants (Δ*hns*, Δ*arcA*, Δ*fliA* and Δ*rcsB*) showed increased activity of the

csgBA transcriptional reporter as well as the number of curli-positive cells, with the effects of *hns*

and *arcA* deletions being most pronounced (Fig. 1B, D and Fig. S2 and S3A).

Moderate reduction of the *csgBA* expression and the number of curli-expressing cells was

observed in mutants lacking genes that encode stress response genes UspE and IraP, indicating

positive but auxiliary involvement of these proteins in regulation of curli structural genes in a well-

shaken *E. coli* planktonic culture (Fig. 1B, E and Fig. S2 and S3B). In contrast to the positive

impact of *fliA* deletion, deletion of *fliZ* showed decreased activity of the *csgBA* reporter (Fig. 1B,

S2, S3C). Deletion of genes encoding global regulators of carbon metabolism *cra* and *crp* and the

regulator of iron homeostasis *fur* severely reduced expression of the *csgBA* operon (Fig. 1B, F and

Fig. S2 and S3C).

Finally, expression of the *csgBA* genes was completely abolished by the removal of

regulators known to directly bind to the *csg* intergenic region and be required for the activation of

curli gene expression (*rpoS, mlrA, csgD, ompR* and *ihfA*)(Fig. 1B and Fig. S2 and S3D) [8]. Similar

abolishment was observed upon disruption of *hfq* gene encoding a chaperone for sRNAs (Fig. 1B,

G and Fig. S2).

Effects of deletions of individual regulator genes on cell growth and dynamics of *csgBA* activation

In order to further explore how removal of individual regulators influences the dynamics of curli gene expression throughout the growth phase, we monitored cell growth and activity of the *csgBA* operon over time using plate reader. The observed *csgBA* expression levels were generally consistent with those measured in the batch culture experiment using flow cytometry (Fig. 2 and Fig. S4, S5, S6). An exception was *arcA* mutant, which showed either same as WT or less activation of curli fibers in the plate reader (Fig. 2B and Fig. S6B), likely because of its slower growth (Fig. 2A and Fig. S6A). Nonetheless, the deletion of *arcA*, and even more so of *hns* led to activation of *csgBA* expression at lower cell density, consistent with derepression of curli genes in these deletion strains. In contrast, removal of IraP – positive regulator of RpoS – delayed curli activation (Fig. 2 and Fig. S6). These observations were further confirmed by measuring reporter expression in corresponding strains grown in well-shaken planktonic cultures using flow cytometry (Fig. S7). Also in these measurements, curli gene expression in Δ*hns* strain was heterogeneous but not as clearly bimodal as in other strains.

Complementation of curli-expression deficient strains by ectopic expression of *csgD*

We further investigated whether the negative impact of dene deletions on the *csgBA* expression could be rescued by the expression of their immediate upstream regulator CsgD from a plasmid under control of the IPTG-inducible promoter. As expected, the expression of *csgD* at higher induction levels resulted in almost all cells expressing curli in the WT strain (Fig.3A). Restoration of the *csgBA* expression was observed in *fur*, *crp, cra* and *hfq* deletion strains, with formation of a well-distinguishable subpopulation of curli-positive cells at intermediate induction levels (Fig. 3B). Expression of CsgD in Δ*hfq* background also resulted in the increased number of curli-expressing cells, but with lower expression levels, and hence lesser separation between curli-positive and -negative cells. Our results thus suggest that these four regulators are primarily required for

regulation upstream of *csgD* and are not essential for bimodal expression of curli fibers. CsgD could

also activate *csgBA* expression in *rpoS* deletion strain (Fig. 3C), which confirms that the expression

of curli structural genes can be induced from a vegetative promoter [12, 20, 32, 64-66]. Reduced

separation between curli-positive and -negative cell subpopulations in this background at

intermediate levels of CsgD induction indicates that the RpoS-dependent expression contributes to

the bimodality of curli gene expression.

In contrast, strains carrying deletions of *ihfA*, *ompR* or *mlrA* genes demonstrated only partial

recovery of curli expression, without formation of a distinct positive population even at high IPTG

induction levels (Fig. 3D). This suggests that these genes directly contribute to the regulation of

curli gene expression downstream of CsgD and might modulate bimodal activation of the

csgBA expression. Notably, although complementation by CsgD from the plasmid was better in the

csgD deletion strain (lacking 421 nucleotides of the coding sequence (see Materials and Methods

for details)), it failed to fully restore the *csgBA* expression, indicating that the native regulation of

csgD and/or gene regulatory elements within its coding sequence might be required to fully induce

expression of curli structural genes.

Effects of deletions of individual DGC genes on curli expression

We next investigated how disruption of individual DGC-encoding genes affects *csgBA* expression

in unstructured planktonic *E. coli* culture. Consistent with our previous study [46], deletions of the

known curli regulators *dgcE* and *dgcM* nearly abolished curli expression (Fig. 4A and Fig. S8).

Although removal of the majority of DGCs had no pronounced effects on the *csgBA* expression

(Fig. 4A and Fig. S9A, S10A, S11), deletions of *dgcO*, *csrD* and *dgcC* genes resulted in markedly

lower activity of the *csgBA* reporter and reduced number of cells expressing curli (Fig. 4A-D and

Fig. S9A, S12), suggesting that only these DGCs contribute to curli regulation in *E. coli* in addition

to DgcE and DgcM under conditions used in this study. Deletion of *dgcI* gene had a more complex

effect, leading to decrease in the number of curli-expressing cells (Fig. S9A), as could be expected

from removal of a DGC, but also resulting in a fraction of positive cells with highly elevated

expression (Fig. 4B), and to transiently higher overall expression level of curli reporter in cell

population in the early growth phase (Fig. 4C, D and Fig. S12).

Since c-di-GMP is known to influence expression of the *csgBA* operon through the

DgcM/PdeR-dependent *csgD* activation, we further measured reporter activity in triple mutants

lacking DgcM/PdeR module and one of the individual DGCs (Fig. 4E). In agreement to our

previous findings, activity of the *csgBA* operon was lower in a mutant lacking DgcM/PdeR

regulatory module compared to WT but similarly bimodal (Fig. 4A, E, F, H, I and Fig. S13) [46].

Consistent with the expected dependence of c-di-GMP regulation on the DgcM/PdeR module,

deletions of other cyclase genes had no effect in this background (Fig. 4E and Fig. S9B, S10B,

S14), including the deletions of *dgcO*, *dgcI* and *dgcC*. In contrast, deletion of the *csrD* gene

encoding a degenerate DGC led to a much stronger decrease of the reporter activity in the

dgcM/pdeR knock-out background compared to the WT background (Fig. 4A, B, E-I and Fig. S9B,

S15). This suggests that the expression of curli structural genes exhibits stronger dependence on

CsrD in the absence of the local c-di-GMP regulatory module.

Discussion

E. coli and other enterobacteria are known to express curli fibers in response to various

environmental stresses. Multiple cellular factors were described to affect curli gene expression in

structured biofilms communities, but the interpretation of previous studies was often complicated

by environmental gradients in such communities. In this study, we systematically investigated the

role of multiple cellular factors in regulation of curli gene expression under simpler conditions, in

E. coli planktonic culture grown under continuous mixing.

Out of 32 tested strains, 21 individual gene deletions were found to modulate the *csgBA*

expression levels and/or the number of curli-expressing cells in our experiments. Curli expression

was elevated in several deletions strains, including those lacking components of the Rcs and

ArcA/B signaling systems. The repression of curli expression by these pathways is in a good

agreement with previous studies, with RcsB known to negatively regulate *csgD* expression [26, 30,

67-70], and ArcA known to repress transcription of *rpoS* [22, 70, 71]. An even more dramatic

enhancement of curli gene expression was observed upon deletion of *hns*, confirming importance of

this DNA-organizing factor as a curli repressor in *E. coli* [32]. Interestingly, H-NS was shown to

activate curli expression in *S. typhimurium* but to repress in *E. coli*, by directly binding to the *csg*

region and also by negatively affecting stability of *rpoS* mRNA [16, 24, 65, 72, 73].

In contrast, two stress regulators, the universal stress factor UspE and the RpoS-stabilizing

factor IraP [74-77] showed positive impact on curli gene expression. The effect of IraP is likely

mediated by the modulation of RpoS stability under starvation conditions [74]. Consistent with that,

the deletion of *iraP* led to the delayed activation of the *csgBA* genes. We further observed the

involvement of the iron-binding global transcriptional regulator Fur in positive curli regulation,

which may be related to the previously observed transcriptional upregulation of *csgD* under iron

limitation [78].

We further observed opposite effects of *fliZ* and *fliA* gene deletions, with curli gene

expression being elevated in Δ*fliA* strain but lowered in Δ*fliZ* strain. The former effect was expected

and it could be explained by the FliA-dependent activation of expression of *pdeH,* the major PDE of

E. coli [79] that inhibits curli gene expression by lowering global levels of c-di-GMP [80].

However, FliZ was previously show to antagonize the activity of RpoS by binding within RpoS-

dependent promoters [37, 79]. Why under our conditions FliZ has a positive rather than negative

impact on curli expression remains to be investigated, but it might be due to its interplay with other

regulators.

Dramatic reduction of curli expression was observed in the absence of global transcriptional

factors Cra and CRP that are activated at low-nutrient conditions [11], thus confirming their

importance for initiation of curli expression [8, 81-84]. Similar abortion of curli expression was

observed upon deletion of *hfq*. Hfq is a chaperone that promotes pairing between sRNAs and their

target mRNA, thus helping to regulate mRNA stability in response to different types of cell stress

[85, 86]. The effect of *hfq* on curli gene expression is likely mediated by altered activity of sRNAs

that are known to modulate curli expression, either directly by binding to the *csg* region [25, 26, 30,

69, 87, 88] or indirectly, through regulation of other cell factors that affect curli production,

including RpoS [89-91]. Consistent with our observation, it has been previously shown that deletion

of *hfq* leads to decreased *csgD* transcript levels in *S. enterica* [92]. Importantly, the defect of curli

expression in all of *crp*, *cra, hfq* as well as *fur* deletion strains could be complemented by

expressing CsgD from an inducible plasmid, confirming that these regulators act upstream of CsgD.

However, the complementation was only partial in the case of *hfq* strain, indicating that a CsgD-

independent impact of sRNA regulation on *csgBA* reporter activity is possibly mediated by RpoS.

Indeed, although ectopic CsgD expression restored activity of the *csgBA* genes in the absence of

rpoS, the maximal level of expression was also lower and similar to that in the complemented *hfq*

strain. Furthermore, it demonstrated that expression of curli structural genes can occur

independently of RpoS, which is consistent with existence of RpoD-dependent promoter upstream

of the *csgBAC* operon [12, 20, 32, 64-66]. Another similarity between the *rpoS* and *hfq* strains was

much reduced bimodality of curli gene expression, indicating that it is at least partly RpoS-

dependent.

The impact on bimodality was even more pronounced for *ompR*, *mlrA* and *ihfA* genes,

whose loss led to the complete abolishment of the *csgBA* activity and could not be fully

compensated by ectopically induced CsgD. The corresponding regulators are known to regulate

transcription of *csgD* and have multiple binding sites within the *csg* intergenic region [16, 23, 33,

65, 93]. Our results indicate that they may also directly activate transcription of curli structural

genes, and that this direct regulation plays an important role in the establishment of *csgBA*

bimodality, possibly by stabilizing the positive expression state. On the other hand, the negative

expression state may be stabilized due to the activity of H-NS, which also has multiple binding sites

within the *csg* region [16, 65]. Indeed, the *csg* intergenic region has strong inherent curvature and

high AT content [94], which aids binding H-NS and IHF. This could induce sharp DNA bending

[16] and, in turn, affect promoter recognition by RpoS [95] and/or accessibility of binding sites for

transcription factors [96]. Indeed, it was shown that OmpR can bind simultaneously with IHF to

positively regulate curli expression, whereas H-NS and IHF act in competition [16]. However,

further experiments are required to elucidate how DNA curvature and the interplay between H-NS

and IHF may determine the bimodality of curli expression.

A number of transcriptional factors did not show any involvement into regulation of curli

fibers expression. These included LeuO, RelA, PdeL, PhoP, ZraP and QseB, but also factors that

were previously described as putative curli regulators (BolA) [60] or proposed to bind to *csg*

intergenic region and affect curli expression in either positive (BasR, RcdA) [58, 59], negative

(BtsR, MqsR) [61, 63] or dual (RstA) [16] manner. Although this discrepancy might be due to

higher complexity of curli regulation in structured communities or differences in growth conditions,

our experiments make it clear that these factors are not essential for curli regulation.

In addition to transcriptional regulators, several DGCs also showed effects on curli

expression in unstructured planktonic *E. coli* culture. Deletion of *dgcO* and *dgcC* resulted in

decreased levels of the *csgBA* expression, dependent on the presence of DgcM/PdeR regulatory

module. A previous study indicated that this effect might be mediated by the interaction of DgcO

with PdeR [29], but the underlying regulatory mechanism remains to be understood. Deletion of

another DGC gene, *dgcI*, had a more complex effect, reducing the number of curli-positive cells at

the late stage of growth – as might be expected from a DGC gene deletion – but increasing the

expression level of curli during entry into the stationary phase. This indicates that DgcI might act

not only as a cyclase but also through a different regulatory mechanism.

Finally, we found that CsrD protein with degenerate diagunilate activity [40] has a positive

effect on curli expression that is particularly pronounced in the absence of the DgcM/PdeR module.

CsrD was reported to enable the RNaseE-mediated degradation of sRNAs CsrB and CsrC [97, 98],

which are known to suppress *csgD* [89]. Moreover, interaction between CsrD and DgcM was

previously observed *in vitro* [29], raising a possibility that it might be sequestered by the DgcM/PdeR complex.

Overall, this work provides a systematic analysis of regulation of curli fiber expression in unstructured planktonic *E. coli* culture and demonstrates the role of several transcription factors and chromosome-shaping proteins in bimodal activation of curli expression.

Acknowledgements

We thank Anna Pashchenko for help with generating DGC deletion strains, Silvia González Sierra for advice on flow cytometry and Julian Pietsch for suggestions regarding data analysis and fruitful discussions. This work was supported by the Max Planck Society.

Data availability

All data needed to evaluate the conclusions are present in the manuscript and/or supporting information.

Conflict of interest

The authors do not declare any conflict of interest.

Figure 1. Curli gene expression responds differently to removal of individual genes. *E. coli* cells carrying genomic transcriptional reporter of the *csgBA* operon were grown in flasks with tryptone broth (TB) at 30°C under constant shaking until they reach maximal *csgBA* reporter levels and then subjected to the flow cytometry analysis. **(A)** Distribution of single-cell fluorescence levels in the wild-type strain (WT) strain. **(B)** Activity of the *csgBA* transcriptional reporter in library of single deletion mutants compared to WT. WT is shown in black color and deletion strains with

enhanced curli expression are indicated in green, with not affected – in blue, reduced – in brown,

strongly impaired – in purple and entirely abolished – in red. Error bars indicate standard error of

the mean (SEM) of at least 3 biological replicates. * at p = 0.01-0.05, ** at p = 0.01-0.001, *** at p

= 0.001-0.0001, **** at p < 0.0001. **(C-G)** Distribution of single-cell fluorescence in indicated

deletion strains with not affected **(C)**, enhanced **(D)**, reduced **(E)**, strongly impaired **(F)** and entirely

abolished **(G)** *csgBA* expression. Fraction of positive cells in the population (mean of at least 3

biological replicates ± SD) is indicated for each strain. Note that the scale in the *y* axes is different

for individual strains to improve readability.

Figure 2. Removal of individual genes alters curli activation pattern. *E. coli* planktonic cultures carrying genomic transcriptional reporter of the *csgBA* operon were grown in a plate reader in TB at 30˚C with shaking. Optical density (OD_{600}) **(A)** and relative fluorescence (absolute fluorescence/OD_{600}) **(B)** of WT and individual deletions strains during their growth in a plate reader. Shown graphs are representative of at least 3 independent replicates (other replicates are shown in Figure S6)

Figure 3. Expression of CsgD from a plasmid restores activity of *csgBA* operon to a different

extent. *E. coli* cells carrying genomic transcriptional reporter of the *csgBA* operon and one of the

indicated gene deletions were transformed with either empty pTrc99a plasmid (control) or with

pTrc99a plasmid carrying *csgD* gene (pTrc99a::*csgD*). Expression from the vector was induced

with 10, 50 and 100 μM IPTG. Bacteria were grown as well-shaked planktonic cultures in TB until

stationary phase (OD$_{600}$ ≥ 1.1) and cultures were subjected to the flow cytometry analysis.

Distribution of single-cell fluorescence levels is shown. Note that the scale in the *y* axes is different

for individual strains to improve readability.

Figure 4. c-di-GMP regulatory network regulates curli expression in addition to DgcM/PdeR

module. Activity of the *csgBA* transcriptional reporter (**A, E**) and distribution of single-cell

fluorescence levels (**B, F, G**) in deletion strains lacking individual DGCs in WT background (**A, B**)

and in absence of DgcM/PdeR regulatory module (**E-G**). WT is shown in black color, strain lacking

DgcM/PdeR regulatory module – in grey and deletion strains with not affected curli expression – in

blue, with affected – in different colors. Error bars indicate SEM of at least 3 biological replicates.

* at p = 0.01-0.05, ** at p = 0.01-0.001, *** at p = 0.001-0.0001, **** at p < 0.0001. *E. coli* cells

carrying genomic transcriptional reporter of the *csgBA* operon were grown in flasks with TB at

30°C under constant shaking until they reach maximal *csgBA* reporter levels and then subjected to

the flow cytometry analysis. Fraction of positive cells in the population (mean of at least 3

biological replicates ± SD) is indicated for each strain. Note that the scale in the *y* axes is different

for individual strains to improve readability. **(C, H)** Optical density (OD$_{600}$) and **(D, I)** relative

fluorescence (absolute fluorescence/OD$_{600}$) of WT and mutant strains lacking indicated DGCs

during the growth in a plate reader (other replicates are shown in Figure S12, S13 and S14). *E. coli*

cells carrying genomic transcriptional reporter of the *csgBA* operon were grown in TB at 30˚C

under constant shaking.

References

402
403
1.   Zogaj X, Bokranz W, Nimtz M, Römling U. Production of cellulose and curli fimbriae by
members of the family Enterobacteriaceae isolated from the human gastrointestinal tract. Infect
Immun. 2003;71(7):4151-8. doi: 10.1128/iai.71.7.4151-4158.2003. PubMed PMID: 12819107;
PubMed Central PMCID: PMCPMC162016.
2.   Hufnagel DA, Depas WH, Chapman MR. The Biology of the *Escherichia coli* Extracellular
Matrix. Microbiol Spectr. 2015;3(3). Epub 2015/07/18. doi: 10.1128/microbiolspec.MB-0014-2014.
PubMed PMID: 26185090; PubMed Central PMCID: PMCPMC4507285.
3.   Brombacher E, Dorel C, Zehnder AJB, Landini P. The curli biosynthesis regulator CsgD co-
ordinates the expression of both positive and negative determinants for biofilm formation in
Escherichia coli. Microbiology (Reading). 2003;149(Pt 10):2847-57. Epub 2003/10/03. doi:
10.1099/mic.0.26306-0. PubMed PMID: 14523117.
4.   McCrate OA, Zhou X, Reichhardt C, Cegelski L. Sum of the Parts: Composition and
Architecture of the Bacterial Extracellular Matrix. Journal of Molecular Biology.
2013;425(22):4286-94. doi: https://doi.org/10.1016/j.jmb.2013.06.022.
5.   Evans ML, Chapman MR. Curli biogenesis: order out of disorder. Biochim Biophys Acta.
2014;1843(8):1551-8. Epub 2013/10/02. doi: 10.1016/j.bbamcr.2013.09.010. PubMed PMID:
24080089; PubMed Central PMCID: PMCPMC4243835.
6.   Blanco LP, Evans ML, Smith DR, Badtke MP, Chapman MR. Diversity, biogenesis and
function of microbial amyloids. Trends Microbiol. 2012;20(2):66-73. Epub 20111223. doi:
10.1016/j.tim.2011.11.005. PubMed PMID: 22197327; PubMed Central PMCID: PMCPMC3278576.
7.   Pham CL, Kwan AH, Sunde M. Functional amyloid: widespread in Nature, diverse in
purpose. Essays Biochem. 2014;56:207-19. doi: 10.1042/bse0560207. PubMed PMID: 25131597.
8.   Smith DR, Price JE, Burby PE, Blanco LP, Chamberlain J, Chapman MR. The Production of
Curli Amyloid Fibers Is Deeply Integrated into the Biology of *Escherichia coli*. Biomolecules.
2017;7(4). Epub 2017/11/01. doi: 10.3390/biom7040075. PubMed PMID: 29088115; PubMed
Central PMCID: PMCPMC5745457.
9.   Van Gerven N, Klein RD, Hultgren SJ, Remaut H. Bacterial amyloid formation: structural
insights into curli biogensis. Trends Microbiol. 2015;23(11):693-706. Epub 20151001. doi:
10.1016/j.tim.2015.07.010. PubMed PMID: 26439293; PubMed Central PMCID: PMCPMC4636965.
10.   Gerstel U, Römling U. Oxygen tension and nutrient starvation are major signals that
regulate *agfD* promoter activity and expression of the multicellular morphotype in *Salmonella*
typhimurium. Environ Microbiol. 2001;3(10):638-48. Epub 2001/11/28. doi: 10.1046/j.1462-
2920.2001.00235.x. PubMed PMID: 11722544.
11.   Rossi E, Paroni M, Landini P. Biofilm and motility in response to environmental and host-
related signals in Gram negative opportunistic pathogens. J Appl Microbiol. 2018. Epub 20180828.
doi: 10.1111/jam.14089. PubMed PMID: 30153375.
12.   Römling U, Rohde M, Olsén A, Normark S, Reinköster J. AgfD, the checkpoint of
multicellular and aggregative behaviour in *Salmonella* typhimurium regulates at least two
independent pathways. Mol Microbiol. 2000;36(1):10-23. Epub 2000/04/12. doi: 10.1046/j.1365-
2958.2000.01822.x. PubMed PMID: 10760159.
13.   Landini P. Cross-talk mechanisms in biofilm formation and responses to environmental and
physiological stress in *Escherichia coli*. Res Microbiol. 2009;160(4):259-66. Epub 20090405. doi:
10.1016/j.resmic.2009.03.001. PubMed PMID: 19345733.
14.   Ghaz-Jahanian MA, Khodaparastan F, Berenjian A, Jafarizadeh-Malmiri H. Influence of small
RNAs on biofilm formation process in bacteria. Mol Biotechnol. 2013;55(3):288-97. doi:
10.1007/s12033-013-9700-6. PubMed PMID: 24062263.

15.   Barnhart MM, Chapman MR. Curli biogenesis and function. Annu Rev Microbiol.
2006;60:131-47. Epub 2006/05/18. doi: 10.1146/annurev.micro.60.080805.142106. PubMed
PMID: 16704339; PubMed Central PMCID: PMCPMC2838481.
16.   Ogasawara H, Yamada K, Kori A, Yamamoto K, Ishihama A. Regulation of the *Escherichia coli*
csgD promoter: interplay between five transcription factors. Microbiology (Reading). 2010;156(Pt
8):2470-83. Epub 20100513. doi: 10.1099/mic.0.039131-0. PubMed PMID: 20466769.
17.   Gerstel U, Römling U. The *csgD* promoter, a control unit for biofilm formation in *Salmonella*
typhimurium. Res Microbiol. 2003;154(10):659-67. Epub 2003/12/04. doi:
10.1016/j.resmic.2003.08.005. PubMed PMID: 14643403.
18.   Ke PC, Zhou R, Serpell LC, Riek R, Knowles TPJ, Lashuel HA, et al. Half a century of amyloids:
past, present and future. Chem Soc Rev. 2020;49(15):5473-509. Epub 20200707. doi:
10.1039/c9cs00199a. PubMed PMID: 32632432; PubMed Central PMCID: PMCPMC7445747.
19.   Jain N, Chapman MR. Bacterial functional amyloids: Order from disorder. Biochim Biophys
Acta Proteins Proteom. 2019;1867(10):954-60. Epub 20190610. doi:
10.1016/j.bbapap.2019.05.010. PubMed PMID: 31195143; PubMed Central PMCID:
PMCPMC6661199.
20.   Zakikhany K, Harrington CR, Nimtz M, Hinton JC, Römling U. Unphosphorylated CsgD
controls biofilm formation in *Salmonella enterica* serovar Typhimurium. Mol Microbiol.
2010;77(3):771-86. Epub 2010/06/16. doi: 10.1111/j.1365-2958.2010.07247.x. PubMed PMID:
20545866.
21.   Chambers JR, Sauer K. Small RNAs and their role in biofilm formation. Trends Microbiol.
2013;21(1):39-49. Epub 20121120. doi: 10.1016/j.tim.2012.10.008. PubMed PMID: 23178000;
PubMed Central PMCID: PMCPMC3752386.
22.   Mika F, Hengge R. Small Regulatory RNAs in the Control of Motility and Biofilm Formation
in *E. coli* and *Salmonella*. Int J Mol Sci. 2013;14(3):4560-79. Epub 20130226. doi:
10.3390/ijms14034560. PubMed PMID: 23443158; PubMed Central PMCID: PMCPMC3634460.
23.   Ogasawara H, Yamamoto K, Ishihama A. Regulatory role of MlrA in transcription activation
of *csgD*, the master regulator of biofilm formation in *Escherichia coli*. FEMS Microbiol Lett.
2010;312(2):160-8. Epub 20100927. doi: 10.1111/j.1574-6968.2010.02112.x. PubMed PMID:
20874755.
24.   Mika F, Hengge R. Small RNAs in the control of RpoS, CsgD, and biofilm architecture of
Escherichia coli. RNA Biol. 2014;11(5):494-507. Epub 20140425. doi: 10.4161/rna.28867. PubMed
PMID: 25028968; PubMed Central PMCID: PMCPMC4152358.
25.   Andreassen PR, Pettersen JS, Szczerba M, Valentin-Hansen P, Møller-Jensen J, Jørgensen
MG. sRNA-dependent control of curli biosynthesis in *Escherichia coli*: McaS directs endonucleolytic
cleavage of *csgD* mRNA. Nucleic Acids Res. 2018;46(13):6746-60. doi: 10.1093/nar/gky479.
PubMed PMID: 29905843; PubMed Central PMCID: PMCPMC6061853.
26.   Jørgensen MG, Nielsen JS, Boysen A, Franch T, Møller-Jensen J, Valentin-Hansen P. Small
regulatory RNAs control the multi-cellular adhesive lifestyle of *Escherichia coli*. Mol Microbiol.
2012;84(1):36-50. Epub 20120118. doi: 10.1111/j.1365-2958.2012.07976.x. PubMed PMID:
22250746.
27.   Simm R, Ahmad I, Rhen M, Le Guyon S, Römling U. Regulation of biofilm formation in
Salmonella enterica serovar Typhimurium. Future Microbiol. 2014;9(11):1261-82. doi:
10.2217/fmb.14.88. PubMed PMID: 25437188.
28.   Lindenberg S, Klauck G, Pesavento C, Klauck E, Hengge R. The EAL domain protein YciR acts
as a trigger enzyme in a c-di-GMP signalling cascade in *E. coli* biofilm control. Embo j.
2013;32(14):2001-14. Epub 20130524. doi: 10.1038/emboj.2013.120. PubMed PMID: 23708798;
PubMed Central PMCID: PMCPMC3715855.

29.   Sarenko O, Klauck G, Wilke FM, Pfiffer V, Richter AM, Herbst S, et al. More than Enzymes
That Make or Break Cyclic Di-GMP-Local Signaling in the Interactome of GGDEF/EAL Domain
Proteins of *Escherichia coli*. mBio. 2017;8(5). Epub 20171010. doi: 10.1128/mBio.01639-17.
PubMed PMID: 29018125; PubMed Central PMCID: PMCPMC5635695.
30.   Mika F, Busse S, Possling A, Berkholz J, Tschowri N, Sommerfeldt N, et al. Targeting of *csgD*
by the small regulatory RNA RprA links stationary phase, biofilm formation and cell envelope stress
in *Escherichia coli*. Mol Microbiol. 2012;84(1):51-65. Epub 20120222. doi: 10.1111/j.1365-
2958.2012.08002.x. PubMed PMID: 22356413; PubMed Central PMCID: PMCPMC3465796.
31.   Junkermeier EH, Hengge R. Local signaling enhances output specificity of bacterial c-di-
GMP signaling networks. Microlife. 2023;4:uqad026. Epub 20230511. doi:
10.1093/femsml/uqad026. PubMed PMID: 37251514; PubMed Central PMCID:
PMCPMC10211494.
32.   Arnqvist A, Olsén A, Normark S. Sigma S-dependent growth-phase induction of the *csgBA*
promoter in *Escherichia coli* can be achieved in vivo by sigma 70 in the absence of the nucleoid-
associated protein H-NS. Mol Microbiol. 1994;13(6):1021-32. doi: 10.1111/j.1365-
2958.1994.tb00493.x. PubMed PMID: 7854117.
33.   Jubelin G, Vianney A, Beloin C, Ghigo JM, Lazzaroni JC, Lejeune P, Dorel C. CpxR/OmpR
interplay regulates curli gene expression in response to osmolarity in *Escherichia coli*. J Bacteriol.
2005;187(6):2038-49. doi: 10.1128/jb.187.6.2038-2049.2005. PubMed PMID: 15743952; PubMed
Central PMCID: PMCPMC1064031.
34.   Prüß BM. Involvement of Two-Component Signaling on Bacterial Motility and Biofilm
Development. J Bacteriol. 2017;199(18). Epub 20170822. doi: 10.1128/jb.00259-17. PubMed
PMID: 28533218; PubMed Central PMCID: PMCPMC5573070.
35.   Dorel C, Vidal O, Prigent-Combaret C, Vallet I, Lejeune P. Involvement of the Cpx signal
transduction pathway of *E. coli* in biofilm formation. FEMS Microbiol Lett. 1999;178(1):169-75. doi:
10.1111/j.1574-6968.1999.tb13774.x. PubMed PMID: 10483736.
36.   Prigent-Combaret C, Brombacher E, Vidal O, Ambert A, Lejeune P, Landini P, Dorel C.
Complex regulatory network controls initial adhesion and biofilm formation in *Escherichia coli* via
regulation of the *csgD* gene. J Bacteriol. 2001;183(24):7213-23. doi: 10.1128/jb.183.24.7213-
7223.2001. PubMed PMID: 11717281; PubMed Central PMCID: PMCPMC95571.
37.   Pesavento C, Hengge R. The global repressor FliZ antagonizes gene expression by σS-
containing RNA polymerase due to overlapping DNA binding specificity. Nucleic Acids Res.
2012;40(11):4783-93. Epub 2012/02/11. doi: 10.1093/nar/gks055. PubMed PMID: 22323519;
PubMed Central PMCID: PMCPMC3367168.
38.   Hengge R. Principles of c-di-GMP signalling in bacteria. Nature Reviews Microbiology.
2009;7(4):263-73. doi: 10.1038/nrmicro2109.
39.   Sommerfeldt N, Possling A, Becker G, Pesavento C, Tschowri N, Hengge R. Gene expression
patterns and differential input into curli fimbriae regulation of all GGDEF/EAL domain proteins in
Escherichia coli. Microbiology (Reading). 2009;155(Pt 4):1318-31. doi: 10.1099/mic.0.024257-0.
PubMed PMID: 19332833.
40.   Hengge R, Galperin Michael Y, Ghigo J-M, Gomelsky M, Green J, Hughes Kelly T, et al.
Systematic Nomenclature for GGDEF and EAL Domain-Containing Cyclic Di-GMP Turnover Proteins
of *Escherichia coli*. Journal of Bacteriology. 2015;198(1):7-11. doi: 10.1128/jb.00424-15.
41.   Weber H, Pesavento C, Possling A, Tischendorf G, Hengge R. Cyclic-di-GMP-mediated
signalling within the sigma network of *Escherichia coli*. Mol Microbiol. 2006;62(4):1014-34. Epub
20060929. doi: 10.1111/j.1365-2958.2006.05440.x. PubMed PMID: 17010156.
42.   Richter AM, Possling A, Malysheva N, Yousef KP, Herbst S, von Kleist M, Hengge R. Local c-
di-GMP Signaling in the Control of Synthesis of the *E. coli* Biofilm Exopolysaccharide pEtN-

Cellulose. J Mol Biol. 2020;432(16):4576-95. Epub 2020/06/14. doi: 10.1016/j.jmb.2020.06.006.
PubMed PMID: 32534064; PubMed Central PMCID: PMCPMC7397504.
43.　Serra DO, Hengge R. A c-di-GMP-Based Switch Controls Local Heterogeneity of Extracellular
Matrix Synthesis which Is Crucial for Integrity and Morphogenesis of *Escherichia coli* Macrocolony
Biofilms. J Mol Biol. 2019;431(23):4775-93. Epub 2019/04/08. doi: 10.1016/j.jmb.2019.04.001.
PubMed PMID: 30954572.
44.　Besharova O, Suchanek VM, Hartmann R, Drescher K, Sourjik V. Diversification of Gene
Expression during Formation of Static Submerged Biofilms by *Escherichia coli*. Front Microbiol.
2016;7:1568. Epub 20161005. doi: 10.3389/fmicb.2016.01568. PubMed PMID: 27761132; PubMed
Central PMCID: PMCPMC5050211.
45.　DePas WH, Hufnagel DA, Lee JS, Blanco LP, Bernstein HC, Fisher ST, et al. Iron induces
bimodal population development by *Escherichia coli*. Proceedings of the National Academy of
Sciences. 2013;110(7):2629-34. doi: 10.1073/pnas.1218703110.
46.　Lamprecht O, Ratnikava M, Jacek P, Kaganovitch E, Buettner N, Fritz K, et al. Regulation by
cyclic di-GMP attenuates dynamics and enhances robustness of bimodal curli gene activation in
Escherichia coli. PLOS Genetics. 2023;19(5):e1010750. doi: 10.1371/journal.pgen.1010750.
47.　Grantcharova N, Peters V, Monteiro C, Zakikhany K, Römling U. Bistable expression of CsgD
in biofilm development of *Salmonella enterica* serovar typhimurium. J Bacteriol. 2010;192(2):456-
66. Epub 20091106. doi: 10.1128/jb.01826-08. PubMed PMID: 19897646; PubMed Central PMCID:
PMCPMC2805326.
48.　Serra DO, Hengge R. Stress responses go three dimensional - the spatial order of
physiological differentiation in bacterial macrocolony biofilms. Environ Microbiol.
2014;16(6):1455-71. Epub 20140505. doi: 10.1111/1462-2920.12483. PubMed PMID: 24725389;
PubMed Central PMCID: PMCPMC4238805.
49.　Serra DO, Klauck G, Hengge R. Vertical stratification of matrix production is essential for
physical integrity and architecture of macrocolony biofilms of *Escherichia coli*. Environ Microbiol.
2015;17(12):5073-88. Epub 20150910. doi: 10.1111/1462-2920.12991. PubMed PMID: 26234179;
PubMed Central PMCID: PMCPMC5014196.
50.　Klauck G, Serra DO, Possling A, Hengge R. Spatial organization of different sigma factor
activities and c-di-GMP signalling within the three-dimensional landscape of a bacterial biofilm.
Open Biol. 2018;8(8). doi: 10.1098/rsob.180066. PubMed PMID: 30135237; PubMed Central
PMCID: PMCPMC6119863.
51.　Serra DO, Richter AM, Klauck G, Mika F, Hengge R. Microanatomy at cellular resolution and
spatial order of physiological differentiation in a bacterial biofilm. mBio. 2013;4(2):e00103-13.
Epub 2013/03/21. doi: 10.1128/mBio.00103-13. PubMed PMID: 23512962; PubMed Central
PMCID: PMCPMC3604763.
52.　Thomason LC, Costantino N, Court DL. *E. coli* genome manipulation by P1 transduction.
Curr Protoc Mol Biol. 2007;Chapter 1:1.17.1-1..8. Epub 2008/02/12. doi:
10.1002/0471142727.mb0117s79. PubMed PMID: 18265391.
53.　Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of *Escherichia
coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. Mol Syst Biol.
2006;2:2006.0008. Epub 2006/06/02. doi: 10.1038/msb4100050. PubMed PMID: 16738554;
PubMed Central PMCID: PMCPMC1681482.
54.　Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli* K-
12 using PCR products. Proc Natl Acad Sci U S A. 2000;97(12):6640-5. Epub 2000/06/01. doi:
10.1073/pnas.120163297. PubMed PMID: 10829079; PubMed Central PMCID: PMCPMC18686.
55.　Cherepanov PP, Wackernagel W. Gene disruption in *Escherichia coli*: TcR and KmR cassettes
with the option of Flp-catalyzed excision of the antibiotic-resistance determinant. Gene.

1995;158(1):9-14. Epub 1995/05/26. doi: 10.1016/0378-1119(95)00193-a. PubMed PMID:
7789817.
56.   Amann E, Ochs B, Abel KJ. Tightly regulated tac promoter vectors useful for the expression
of unfused and fused proteins in *Escherichia coli*. Gene. 1988;69(2):301-15. Epub 1988/09/30. doi:
10.1016/0378-1119(88)90440-4. PubMed PMID: 3069586.
57.   Chung CT, Niemela SL, Miller RH. One-step preparation of competent *Escherichia coli*:
transformation and storage of bacterial cells in the same solution. Proc Natl Acad Sci U S A.
1989;86(7):2172-5. Epub 1989/04/01. doi: 10.1073/pnas.86.7.2172. PubMed PMID: 2648393;
PubMed Central PMCID: PMCPMC286873.
58.   Ogasawara H, Shinohara S, Yamamoto K, Ishihama A. Novel regulation targets of the metal-
response BasS-BasR two-component system of *Escherichia coli*. Microbiology (Reading).
2012;158(Pt 6):1482-92. Epub 20120322. doi: 10.1099/mic.0.057745-0. PubMed PMID: 22442305.
59.   Shimada T, Katayama Y, Kawakita S, Ogasawara H, Nakano M, Yamamoto K, Ishihama A. A
novel regulator RcdA of the *csgD* gene encoding the master regulator of biofilm formation in
Escherichia coli. Microbiologyopen. 2012;1(4):381-94. Epub 20121008. doi: 10.1002/mbo3.42.
PubMed PMID: 23233451; PubMed Central PMCID: PMCPMC3535384.
60.   Azam MW, Zuberi A, Khan AU. *bolA* gene involved in curli amyloids and fimbriae production
in *E. coli*: exploring pathways to inhibit biofilm and amyloid formation. J Biol Res (Thessalon).
2020;27:10. Epub 20200617. doi: 10.1186/s40709-020-00120-7. PubMed PMID: 32566535;
PubMed Central PMCID: PMCPMC7301969.
61.   Ogasawara H, Ishizuka T, Yamaji K, Kato Y, Shimada T, Ishihama A. Regulatory role of
pyruvate-sensing BtsSR in biofilm formation by *Escherichia coli* K-12. FEMS Microbiol Lett.
2019;366(24). Epub 2019/12/14. doi: 10.1093/femsle/fnz251. PubMed PMID: 31834370.
62.   Ogasawara H, Hasegawa A, Kanda E, Miki T, Yamamoto K, Ishihama A. Genomic SELEX
search for target promoters under the control of the PhoQP-RstBA signal relay cascade. J Bacteriol.
2007;189(13):4791-9. Epub 20070427. doi: 10.1128/jb.00319-07. PubMed PMID: 17468243;
PubMed Central PMCID: PMCPMC1913430.
63.   Soo VW, Wood TK. Antitoxin MqsA represses curli formation through the master biofilm
regulator CsgD. Sci Rep. 2013;3:3186. Epub 2013/11/12. doi: 10.1038/srep03186. PubMed PMID:
24212724; PubMed Central PMCID: PMCPMC4894380.
64.   Brown PK, Dozois CM, Nickerson CA, Zuppardo A, Terlonge J, Curtiss R, 3rd. MlrA, a novel
regulator of curli (AgF) and extracellular matrix synthesis by *Escherichia coli* and *Salmonella*
enterica serovar Typhimurium. Mol Microbiol. 2001;41(2):349-63. Epub 2001/08/08. doi:
10.1046/j.1365-2958.2001.02529.x. PubMed PMID: 11489123.
65.   Gerstel U, Park C, Römling U. Complex regulation of *csgD* promoter activity by global
regulatory proteins. Mol Microbiol. 2003;49(3):639-54. Epub 2003/07/17. doi: 10.1046/j.1365-
2958.2003.03594.x. PubMed PMID: 12864849.
66.   Weber H, Polen T, Heuveling J, Wendisch VF, Hengge R. Genome-wide analysis of the
general stress response network in *Escherichia coli*: sigmaS-dependent genes, promoters, and
sigma factor selectivity. J Bacteriol. 2005;187(5):1591-603. Epub 2005/02/18. doi:
10.1128/jb.187.5.1591-1603.2005. PubMed PMID: 15716429; PubMed Central PMCID:
PMCPMC1063999.
67.   Ferrières L, Clarke DJ. The RcsC sensor kinase is required for normal biofilm formation in
Escherichia coli K-12 and controls the expression of a regulon in response to growth on a solid
surface. Mol Microbiol. 2003;50(5):1665-82. doi: 10.1046/j.1365-2958.2003.03815.x. PubMed
PMID: 14651646.
68.   Vianney A, Jubelin G, Renault S, Dorel C, Lejeune P, Lazzaroni JC. *Escherichia coli tol* and *rcs*
genes participate in the complex network affecting curli synthesis. Microbiology (Reading).
2005;151(Pt 7):2487-97. doi: 10.1099/mic.0.27913-0. PubMed PMID: 16000739.

69.   Boehm A, Vogel J. The *csgD* mRNA as a hub for signal integration via multiple small RNAs.
Mol Microbiol. 2012;84(1):1-5. Epub 20120313. doi: 10.1111/j.1365-2958.2012.08033.x. PubMed
PMID: 22414234.
70.   Mika F, Hengge R. A two-component phosphotransfer network involving ArcB, ArcA, and
RssB coordinates synthesis and proteolysis of sigmaS (RpoS) in *E. coli*. Genes Dev.
2005;19(22):2770-81. doi: 10.1101/gad.353705. PubMed PMID: 16291649; PubMed Central
PMCID: PMCPMC1283968.
71.   Hengge R. Linking bacterial growth, survival, and multicellularity - small signaling molecules
as triggers and drivers. Curr Opin Microbiol. 2020;55:57-66. Epub 20200331. doi:
10.1016/j.mib.2020.02.007. PubMed PMID: 32244175.
72.   Olsén A, Arnqvist A, Hammar M, Sukupolvi S, Normark S. The RpoS sigma factor relieves H-
NS-mediated transcriptional repression of *csgA*, the subunit gene of fibronectin-binding curli in
Escherichia coli. Mol Microbiol. 1993;7(4):523-36. Epub 1993/02/01. doi: 10.1111/j.1365-
2958.1993.tb01143.x. PubMed PMID: 8459772.
73.   Zhou Y, Gottesman S. Modes of regulation of RpoS by H-NS. J Bacteriol. 2006;188(19):7022-
5. doi: 10.1128/jb.00687-06. PubMed PMID: 16980505; PubMed Central PMCID:
PMCPMC1595528.
74.   Bougdour A, Wickner S, Gottesman S. Modulating RssB activity: IraP, a novel regulator of
sigma(S) stability in *Escherichia coli*. Genes Dev. 2006;20(7):884-97. Epub 2006/04/08. doi:
10.1101/gad.1400306. PubMed PMID: 16600914; PubMed Central PMCID: PMCPMC1472289.
75.   Bougdour A, Cunning C, Baptiste PJ, Elliott T, Gottesman S. Multiple pathways for
regulation of sigmaS (RpoS) stability in *Escherichia coli* via the action of multiple anti-adaptors. Mol
Microbiol. 2008;68(2):298-313. doi: 10.1111/j.1365-2958.2008.06146.x. PubMed PMID: 18383615.
76.   Girard ME, Gopalkrishnan S, Grace ED, Halliday JA, Gourse RL, Herman C. DksA and ppGpp
Regulate the σ(S) Stress Response by Activating Promoters for the Small RNA DsrA and the Anti-
Adapter Protein IraP. J Bacteriol. 2018;200(2). Epub 20171220. doi: 10.1128/jb.00463-17. PubMed
PMID: 29061665; PubMed Central PMCID: PMCPMC5738727.
77.   Bougdour A, Gottesman S. ppGpp regulation of RpoS degradation via anti-adaptor protein
IraP. Proc Natl Acad Sci U S A. 2007;104(31):12896-901. Epub 20070719. doi:
10.1073/pnas.0705561104. PubMed PMID: 17640895; PubMed Central PMCID: PMCPMC1937563.
78.   Baez A, Sharma AK, Bryukhanov A, Anderson ED, Rudack L, Olivares-Hernández R, et al. Iron
availability enhances the cellular energetics of aerobic *Escherichia coli* cultures while upregulating
anaerobic respiratory chains. N Biotechnol. 2022;71:11-20. Epub 20220628. doi:
10.1016/j.nbt.2022.06.004. PubMed PMID: 35777694; PubMed Central PMCID: PMCPMC9444934.
79.   Pesavento C, Becker G, Sommerfeldt N, Possling A, Tschowri N, Mehlis A, Hengge R. Inverse
regulatory coordination of motility and curli-mediated adhesion in *Escherichia coli*. Genes Dev.
2008;22(17):2434-46. doi: 10.1101/gad.475808. PubMed PMID: 18765794; PubMed Central
PMCID: PMCPMC2532929.
80.   Wang L, Malengo G, Sanches-Medeiros A, Chen X, Teteneva N, Sierra SG, et al. A toolbox of
FRET-based c-di-GMP biosensors and its FRET-To-Sort application for genome-wide mapping of the
second messenger regulatory network. bioRxiv. 2024:2024.08.21.609041. doi:
10.1101/2024.08.21.609041.
81.   Ramseier TM. Cra and the control of carbon flux via metabolic pathways. Res Microbiol.
1996;147(6-7):489-93. doi: 10.1016/0923-2508(96)84003-4. PubMed PMID: 9084760.
82.   Reshamwala SM, Noronha SB. Biofilm formation in *Escherichia coli cra* mutants is impaired
due to down-regulation of curli biosynthesis. Arch Microbiol. 2011;193(10):711-22. Epub
20110511. doi: 10.1007/s00203-011-0708-7. PubMed PMID: 21559929.
83.   Amores GR, de Las Heras A, Sanches-Medeiros A, Elfick A, Silva-Rocha R. Systematic
identification of novel regulatory interactions controlling biofilm formation in the bacterium

Escherichia coli. Sci Rep. 2017;7(1):16768. Epub 2017/12/03. doi: 10.1038/s41598-017-17114-6.
PubMed PMID: 29196655; PubMed Central PMCID: PMCPMC5711951.

84.   Zheng D, Constantinidou C, Hobman JL, Minchin SD. Identification of the CRP regulon using
in vitro and in vivo transcriptional profiling. Nucleic Acids Res. 2004;32(19):5874-93. Epub
20041101. doi: 10.1093/nar/gkh908. PubMed PMID: 15520470; PubMed Central PMCID:
PMCPMC528793.

85.   Guisbert E, Rhodius VA, Ahuja N, Witkin E, Gross CA. Hfq modulates the sigmaE-mediated
envelope stress response and the sigma32-mediated cytoplasmic stress response in *Escherichia*
coli. J Bacteriol. 2007;189(5):1963-73. Epub 20061208. doi: 10.1128/jb.01243-06. PubMed PMID:
17158661; PubMed Central PMCID: PMCPMC1855744.

86.   Santiago-Frangos A, Woodson SA. Hfq chaperone brings speed dating to bacterial sRNA.
Wiley Interdiscip Rev RNA. 2018;9(4):e1475. Epub 20180406. doi: 10.1002/wrna.1475. PubMed
PMID: 29633565; PubMed Central PMCID: PMCPMC6002925.

87.   Bordeau V, Felden B. Curli synthesis and biofilm formation in enteric bacteria are
controlled by a dynamic small RNA module made up of a pseudoknot assisted by an RNA
chaperone. Nucleic Acids Res. 2014;42(7):4682-96. Epub 20140201. doi: 10.1093/nar/gku098.
PubMed PMID: 24489123; PubMed Central PMCID: PMCPMC3985669.

88.   Holmqvist E, Reimegård J, Sterk M, Grantcharova N, Römling U, Wagner EG. Two antisense
RNAs target the transcriptional regulator CsgD to inhibit curli synthesis. Embo j. 2010;29(11):1840-
50. Epub 20100420. doi: 10.1038/emboj.2010.73. PubMed PMID: 20407422; PubMed Central
PMCID: PMCPMC2885931.

89.   Bak G, Lee J, Suk S, Kim D, Young Lee J, Kim KS, et al. Identification of novel sRNAs involved
in biofilm formation, motility, and fimbriae formation in *Escherichia coli*. Sci Rep. 2015;5:15287.
Epub 20151015. doi: 10.1038/srep15287. PubMed PMID: 26469694; PubMed Central PMCID:
PMCPMC4606813.

90.   Rodgers ML, O'Brien B, Woodson SA. Small RNAs and Hfq capture unfolded RNA target
sites during transcription. Mol Cell. 2023;83(9):1489-501.e5. Epub 20230427. doi:
10.1016/j.molcel.2023.04.003. PubMed PMID: 37116495; PubMed Central PMCID:
PMCPMC10176597.

91.   Majdalani N, Vanderpool CK, Gottesman S. Bacterial small RNA regulators. Crit Rev
Biochem Mol Biol. 2005;40(2):93-113. doi: 10.1080/10409230590918702. PubMed PMID:
15814430.

92.   Monteiro C, Papenfort K, Hentrich K, Ahmad I, Le Guyon S, Reimann R, et al. Hfq and Hfq-
dependent small RNAs are major contributors to multicellular development in *Salmonella enterica*
serovar Typhimurium. RNA Biol. 2012;9(4):489-502. Epub 20120216. doi: 10.4161/rna.19682.
PubMed PMID: 22336758.

93.   Gerstel U, Kolb A, Römling U. Regulatory components at the *csgD* promoter--additional
roles for OmpR and integration host factor and role of the 5' untranslated region. FEMS Microbiol
Lett. 2006;261(1):109-17. Epub 2006/07/18. doi: 10.1111/j.1574-6968.2006.00332.x. PubMed
PMID: 16842367.

94.   Pedersen AG, Jensen LJ, Brunak S, Staerfeldt HH, Ussery DW. A DNA structural atlas for
Escherichia coli. J Mol Biol. 2000;299(4):907-30. doi: 10.1006/jmbi.2000.3787. PubMed PMID:
10843847.

95.   Maciag A, Peano C, Pietrelli A, Egli T, De Bellis G, Landini P. In vitro transcription profiling of
the σS subunit of bacterial RNA polymerase: re-definition of the σS regulon and identification of
σS-specific promoter sequence elements. Nucleic Acids Res. 2011;39(13):5338-55. Epub 20110311.
doi: 10.1093/nar/gkr129. PubMed PMID: 21398637; PubMed Central PMCID: PMCPMC3141248.

96. Yilmaz C, Schnetz K. High Abundance of Transcription Regulators Compacts the Nucleoid in
Escherichia coli. J Bacteriol. 2022;204(6):e0002622. Epub 20220518. doi: 10.1128/jb.00026-22.
PubMed PMID: 35583339; PubMed Central PMCID: PMCPMC9210969.
97. Suzuki K, Babitzke P, Kushner SR, Romeo T. Identification of a novel regulatory protein
(CsrD) that targets the global regulatory RNAs CsrB and CsrC for degradation by RNase E. Genes
Dev. 2006;20(18):2605-17. doi: 10.1101/gad.1461606. PubMed PMID: 16980588; PubMed Central
PMCID: PMCPMC1578682.
98. Jonas K, Tomenius H, Römling U, Georgellis D, Melefors O. Identification of YhdA as a
regulator of the *Escherichia coli* carbon storage regulation system. FEMS Microbiol Lett.
2006;264(2):232-7. doi: 10.1111/j.1574-6968.2006.00457.x. PubMed PMID: 17064377.
749