Alcohol disrupts long-term potentiation at hippocampus-medium spiny neuron synapses in the

medial shell of the nucleus accumbens

Ashley E. Copenhaver, Ph.D.[1], Jalane R. Campbell[1], Tara A. LeGates, Ph.D.[1,2]

[1]Department of Biological Sciences, University of Maryland, Baltimore County (UMBC)

[2]Department of Pharmacology and Physiology, University of Maryland, School of Medicine

Corresponding author:

Tara LeGates, Ph.D.

1000 Hilltop Circle, Interdisciplinary Life Sciences Building 315

Baltimore, MD 21250

tlegates@umbc.edu

Support: We would like to thank Dr. Yaohui Zhu and Kaela Befano for their assistance and Dr.

Brian Mathur for his technical support and thoughtful comments on the manuscript. This work was

supported by NSF IOS2402645, T32GM144876-02, the Meyerhoff program at UMBC, and start-

up funds from UMBC.

ABSTRACT

Background: Chronic alcohol exposure is a major driver of alcohol use disorders (AUD), in

part through its ability to induce maladaptive plasticity within neural circuits that regulate reward,

motivation, and affect. Excitatory projections from the hippocampus (Hipp) to the nucleus

accumbens (NAc) play a pivotal role in regulating reward-related behaviors, and this pathway

serves as a key locus for establishing associations between rewarding stimuli and related

contextual information. Regulation of the strength of Hipp-NAc synapses is critical for supporting

these behaviors, and aberrant Hipp-NAc plasticity is associated with anhedonia and disrupted

reward learning.

Methods: To examine acute ethanol effects, we used whole-cell electrophysiology to record

Hipp-NAc synaptic plasticity in acute brain slices in the presence or absence of 50mM ethanol.

To examine the effects of chronic ethanol administration, mice were exposed to ethanol vapor in

a 3-week chronic intermittent ethanol (CIE) paradigm. Slices from ethanol and air exposed mice

were used for whole-cell electrophysiology to examine Hipp-NAc synaptic plasticity.

Results: Here, we demonstrate that acute ethanol application to *ex vivo* brain slices prevents

long-term potentiation (LTP) at Hipp-NAc synapses, without altering presynaptic release

probability. Furthermore, chronic intermittent exposure to ethanol abolishes LTP at these

synapses, even during abstinence, indicating persistent synaptic dysfunction.

Conclusions: Together, our findings demonstrate that ethanol has immediate and long-lasting

effects on Hipp-NAc plasticity. Given the behavioral relevance of these synapses, this work has

important implications for the mechanisms underlying ethanol-dependent effects on reward

processing and negative affective states associated with AUD.

KEYWORDS hippocampus, nucleus accumbens, synaptic plasticity, chronic intermittent ethanol,

reward

INTRODUCTION

Alcohol exerts both immediate and long-lasting effects on motivated behaviors, cognitive

function, and emotional regulation (Koob and Volkow, 2010). These effects arise primarily from

alcohol's direct modulation of synaptic transmission and its influence over plasticity within key

brain reward circuits (Lovinger and Abrahao, 2018). Although heightened plasticity in reward

pathways may initially promote alcohol-seeking behaviors, persistent changes in circuit activation

can ultimately impair the brain's ability to respond to natural rewards and trigger prolonged

activation of stress-related neural systems. As a result, disruption of normal reward processing

and persistent stress signaling are thought to underlie the negative emotional states frequently

associated with chronic alcohol use and dependence. AUD with comorbid anhedonia (loss of

pleasure derived from previously pleasurable activities) is associated with increased risk of

relapse and poorer clinical outcomes, presenting a major obstacle to recovery (Garfield, Lubman

and Yücel, 2014; Destoop *et al.*, 2019).

The nucleus accumbens (NAc) is a key mediator of motivated behaviors where both stress

and rewarding experiences modulate synaptic function and plasticity to drive alterations in

behavior (Carlezon and Thomas, 2009; Stuber, Britt and Bonci, 2012; Russo and Nestler, 2013).

Excitatory input from several key brain regions (e.g., hippocampus, prefrontal cortex, thalamus,

amygdala) has a crucial role in regulating NAc neuron activity to effectively encode reward-

associated information and modulate motivated behaviors (O'Donnell and Grace, 1995; Goto and

Grace, 2008; Calhoon and O'Donnell, 2013). Alterations in the strength and plasticity of these

synapses as well as changes in excitability of NAc neurons have been associated with substance

use disorders and depression, further demonstrating the key role of NAc activity in modulating

motivated behaviors and emotional states (Volkow and Morales, 2015; Scofield *et al.*, 2016;

Francis and Lobo, 2017). For instance, chronic ethanol exposure enhances excitability and

increases NMDAR activity in NAc neurons, consistently leading to increased volitional ethanol

intake while also disrupting N-methyl-D-aspartate receptor (NMDAR)-dependent long-term

depression (LTD) (Renteria *et al.*, 2017; Kircher *et al.*, 2019). This combination of effects mirrors

the consequences of chronic stress, which similarly induces anhedonia and disrupts NMDAR-

dependent LTD within the NAc (Lim *et al.*, 2012). Together, this suggests that ethanol and stress

may converge on overlapping mechanisms within the NAc, particularly through alterations in

excitatory synaptic plasticity, to promote anhedonia and dysregulated reward processing.

Projections from the hippocampus (Hipp) provide a majority of the excitatory input to the NAc,

critically modulating neuronal activity and gating the flow of information through the region

(O'Donnell and Grace, 1995; Wilson and Kawaguchi, 1996; Stern, Jaeger and Wilson, 1998; Goto

and O'Donnell, 2001). Hipp-NAc connectivity has been tied to several behaviors related to reward

and motivation in humans and rodents including positive reinforcement learning, substance

abuse, positive affect, as well as spatial- and contextual-based reward learning (Floresco and

Phillips, 1999; Ito *et al.*, 2008; Britt *et al.*, 2012; Davidow *et al.*, 2016; LeGates *et al.*, 2018; Sjulson

et al., 2018; Trouche *et al.*, 2019; Zhou *et al.*, 2019; Heller *et al.*, 2020; Huntley *et al.*, 2020; Sosa,

Joo and Frank, 2020; Barnstedt, Mocellin and Remy, 2024). Our previous work revealed that

activity-dependent strengthening of Hipp-NAc synapses drives reward-related behaviors while

chronic stress disrupts this plasticity and induces anhedonia (LeGates *et al.*, 2018). Hipp and NAc

activity also influences alcohol-related behaviors, where reduced Hipp activity causes an increase

in alcohol consumption, and chemogenetic activation of Hipp-NAc synapses attenuates excessive

drinking (Griffin *et al.*, 2023). Furthermore, chronic alcohol exposure induces pre- and

postsynaptic changes at Hipp-NAc synapses and abolishes activity-dependent LTD (Kircher *et*

al., 2019), again highlighting the similarly powerful effects of chronic stress and chronic alcohol

on synaptic function.

Focusing on medium spiny neurons (MSNs), which represent ~95% of the neuron population

in the NAc, we recently showed that LTP at Hipp-NAc synapses relies on NMDAR and L-type

voltage gated calcium channels (L-type VGCCs) in males and females, respectively (Copenhaver

and LeGates, 2024). Given the previously established ability of ethanol to inhibit NMDAR function

within the NAc and its ability to inhibit L-type VGCCs (Lovinger, White and Weight, 1989; Zucca

and Valenzuela, 2010; Mah, Fleck and Lindsley, 2011; Lee, Yeh and Yeh, 2022), we sought to

determine whether ethanol exposure influenced activity-dependent potentiation of Hipp-NAc

synapses similarly in both sexes. Here, we show that both acute *in vitro* and chronic *in vivo*

exposure to ethanol prevents LTP at Hipp-NAc synapses in males and females. Given the link

between Hipp-NAc potentiation in mediating reward-related behaviors, ethanol's effects on these

synapses may be a key mechanism driving maladaptive reward-related behaviors observed in

response to alcohol use.

MATERIALS AND METHODS

Male and female C57BL/6J mice (8-10 weeks) were group housed (2-5 mice per cage) in a

12:12LD cycle with food and water ad libitum. We did not track the estrous cycle in female mice.

All experiments were performed in accordance with the regulations set forth by the Institutional

Animal Care and Use Committee at the University of Maryland, Baltimore County.

Ethanol Vapor Chamber Exposure

Male and female mice (8-10 weeks at the start of the experiment) were placed in acrylic

inhalation chambers (60 × 59 × 35 cm) in their home cages and exposed to chronic intermittent

ethanol (CIE) vapor or air 14 hours/day (ZT11-1), 4 days a week, for a total of 3 weeks.

Compressed air (10–20 psi) was passed through a tube submerged in 100% ethanol to volatilize.

Control vapor chambers delivered only air with no ethanol vapor. Vapor chamber ethanol

concentrations were monitored daily, and air flow was adjusted to produce ethanol concentrations

within 1.8-2.0% ethanol content as measured by a digital alcohol breath tester (BACTrack). After

four days in the inhalation chambers, mice underwent a 72-h forced abstinence period. Whole

cell electrophysiology was performed on week 4 after the 72-h abstinence period.

Brain Slice Preparation

Acute parasagittal slices (lateral 0.36-0.72) containing the fornix and nucleus accumbens

were prepared for whole-cell patch-clamp electrophysiology. Animals were deeply anesthetized

with isoflurane, decapitated, and brains were quickly dissected and submerged in ice-cold,

bubbled (carbogen: 95% O_2/5% CO_2) N-methyl-D-glucamine (NMDG) recovery solution

containing the following (in mM): 93 NMDG, 2.5 KCl, 1.2 NaH_2PO_4, 11 glucose, 25 $NaHCO_3$, 1.2

$MgCl_2$, and 2.4 $CaCl_2$, pH=7.3-7.4, osmolarity=300-310 mOsm. Using a vibratome (VT1000S,

Leica Microsystems), parasagittal slices (400 μm) were cut in cold, oxygenated NMDG. Slices

were transferred to 32-34°C NMDG for 7-12 minutes to recover and were then transferred to

room-temperature artificial cerebrospinal fluid (aCSF) containing the following (in mM): 120 NaCl,

3 KCl, 1.0 NaH_2PO_4, 20 glucose, 25 $NaHCO_3$, 1.5 $MgCl_2 \cdot 7H_2O$, and 2.5 $CaCl_2$, pH=7.3-7.4. Slices

were allowed to recover for 1-hour at room-temperature before beginning electrophysiological

recordings.

Whole-Cell Recordings

Slices were placed in a submersion-type recording chamber and superfused with room-

temperature aCSF (flow rate 0.5-1mL/min). Cells were visualized using a 60x water immersion

objective on a Nikon Eclipse FN-1 or Scientifica microscope. Whole-cell patch-clamp recordings

were performed using an Axopatch 200B amplifier (Axon Instruments, Molecular Devices) and a

Digidata 1550B digitizer (Axon Instruments). Patch pipettes (4-8MΩ) were made from borosilicate

glass (World Precision Instruments) using a Sutter Instruments P-97 model puller.

All recordings were performed in voltage-clamp conditions from MSNs in the NAc medial shell.

A bipolar stimulating electrode (FHC) was placed in the fornix to electrically stimulate hippocampal

axons while recording evoked excitatory postsynaptic currents (EPSC) (Figure 1a). Patch pipettes

were filled with a solution containing 130 mM K-gluconate, 5 mM KCl, 2 mM MgCl6-H2O, 10 mM

HEPES, 4 mM Mg-ATP, 0.3 mM Na2-GTP, 10 mM Na2-phosphocreatine, and 1 mM EGTA;

pH=7.3-7.4; osmolarity=285-295mOsm. EPSCs were recorded at -70mV from paired pulses

(100ms apart) elicited every 10s. After obtaining a stable 5-minute baseline EPSC recording, high-

frequency stimulation (HFS: four trains of 100Hz stimulation for 1s with 15s between trains while

holding the cell at -40mV) was applied, followed by a 30-minute recording of EPSCs. One cell

was recorded per slice. For acute ethanol experiments, aCSF containing 50mM ethanol (Sigma)

was washed onto slices for at least 20 minutes prior to recording, and experiments were done in

the presence of ethanol. We used the following exclusion criteria to eliminate unhealthy cells and

unreliable recordings: 1) We only proceeded with experiments on cells with series resistances

<10MΩ, 2) Cells were excluded if their series resistance changed by >20% (comparing the

resistance at the beginning and end of the experiment), 3) Cells in poor health or poor recording

status were excluded (i.e. cell partially or fully sealed up, a decrease in holding current >100pA

that is consistent with the cell dying, an increase in the variability of timing of EPSCs (jitter) post

HFS, and/or an increase in response failure rate to > 50%).

Data analysis

To analyze data from HFS experiments, EPSC amplitudes were normalized to the average

baseline amplitude. Normalized EPSC amplitudes were averaged over the last 5 minutes of the

recording (minutes 25-30 post HFS) to examine the direction and magnitude of plasticity. Paired

pulse ratio (PPR) was measured by calculating 5-minute averages of EPSC amplitudes from our

paired pulses ($EPSC_1$ and $EPSC_2$) and dividing $EPSC_2$ by $EPSC_1$ (ie: $EPSC_2/EPSC_1$ = PPR).

Statistical tests were performed using GraphPad Prism 9/10 software. A p-value of <0.05 was

considered statistically significant, and exact p-values can be found in the results section. For

acute ethanol experiments, n represents the number of cells/slices. For chronic ethanol

experiments, n represents the number of mice. For line graphs, points represent the mean and

error bars indicate SEM. Summary data plots display one point for each cell, with each point

representing the average EPSC amplitude from the last 5 minutes of recording. For bar graphs,

the bar represents the mean and error bars indicate SEM with individual data points overlayed.

Statistical significance (p<0.05) was assessed by Two-way ANOVA (sex x treatment) with Šídák's

multiple comparisons. A three-way ANOVA (sex x treatment x time) was used to analyze PPR

data before and after HFS.

RESULTS

Acute ethanol application to *ex vivo* slices prevents Hipp-NAc LTP

To determine whether ethanol acutely influences plasticity at Hipp-NAc synapses, we

recorded HFS-induced LTP in the presence and absence of 50mM ethanol (Figure 1a). We found

that, in the absence of ethanol, HFS induced LTP at Hipp-NAc synapses slices taken from male

and female mice with no difference in magnitude between the sexes (Figure 1b-c), similar to our

previous published results (LeGates *et al.*, 2018; Copenhaver and LeGates, 2024). However,

pretreatment with 50mM ethanol prevented LTP induction in slices taken from males and females

(Figure 1b-c; Sex x Treatment Interaction: $F_{(1,16)}$ = 0.1512, p=0.7025; Main effect of Sex: $F_{(1,16)}$ =

0.01263, p=0.9119; Main effect of treatment: $F_{(1,16)}$ = 19.24, p=0.0005; Šídák's multiple

comparisons post hoc: $p_{(Male:control\ vs\ EtOH)}$ = 0.0175, $p_{(Female:control\ vs\ EtOH)}$ = 0.0107). We observed no

difference in paired pulse ratio (PPR) at baseline (Figure 1d; Sex x Treatment Interaction: $F_{(1,16)}$ =

0.003732, p=0.9520; Main effect of Sex: $F_{(1,16)}$ = 1.066, p=0.3173; Main effect of treatment: $F_{(1,16)}$

= 0.2307, p=0.6375) suggesting that application of ethanol did not alter basal release probability.

Furthermore, no change in PPR was observed following HFS irrespective of whether ethanol was

present (Figure 1b, bottom; Time x Sex x Treatment Interaction: $F_{(2.590,\ 41.45)}$ = 0.5932, p=0.5992;

Time x Treatment Interaction: $F_{(2.590,41.45)}$ = 1.814, p=0.1661; Time x Sex Interaction: $F_{(2.590,\ 41.45)}$ =

0.3843, p=0.7359), suggesting that HFS did not elicit changes in presynaptic neurotransmitter

release probability.

CIE disrupts Hipp-NAc LTP

To determine whether chronic exposure to ethanol leads to changes in Hipp-NAc plasticity,

we used CIE to ethanol vapor for controlled delivery and dosing of ethanol over a 3-week period

(Figure 2a). Control mice were exposed to air. Following at least 72 hours of forced abstinence,

slices from CIE-exposed mice demonstrated a deficit in HFS-induced LTP relative to air-exposed

controls, which showed typical LTP (Figure 2b-c; Sex x Treatment Interaction: $F_{(1,20)}$ = 0.03737,

p=0.8487; Main effect of Sex: $F_{(1,20)}$ = 0.1118, p=0.7415; Main effect of treatment: $F_{(1,20)}$ = 12.53,

p=0.002; Šídák's multiple comparisons post hoc: $p_{(Male:control\ vs\ EtOH)}$ = 0.0444, $p_{(Female:control\ vs\ EtOH)}$ =

0.0390). This effect was consistent in both male and female mice. Comparison of baseline PPRs

revealed no difference between CIE mice and air exposed controls (Figure 2d; Sex x Treatment

Interaction: $F_{(1,20)}$ = 0.2254, p=0.6401; Main effect of Sex: $F_{(1,20)}$ = 0.0008649, p=0.9768; Main

effect of treatment: $F_{(1,20)}$ = 1.146, p=0.2971), suggesting that CIE did not induce changes in

release probability. Similar to the data shown in Figure 1, HFS did not induce changes in PPR

regardless of ethanol exposure or sex (Figure 2b bottom; Time x Sex x Treatment Interaction:

$F_{(6,20)}$ = 1.600, p=0.1529; Time x Treatment Interaction: $F_{(6,20)}$ = 0.6492, p=0.6906; Time x Sex

Interaction: $F_{(6,20)}$ = 1.067, p=3862), demonstrating a lack of effect on presynaptic function.

DISCUSSION

Our data demonstrate that acute application of ethanol to *ex vivo* brain slices disrupts HFS-

induced LTP at Hipp-NAc synapses. We also found that chronic exposure to ethanol *in vivo*

interfered with HFS-induced LTP at Hipp-NAc synapses, an effect observed after a 72-hour

abstinence period when ethanol was no longer present. Neither treatment led to a change in basal

release probability, suggesting that ethanol may be acting postsynaptically to interfere with LTP.

Given the previously described role for Hipp-NAc synaptic strength in mediating reward-related

behaviors, ethanol's impact on Hipp-NAc plasticity has important implications for understanding

the neurobiological basis for disrupted reward processing and negative mood states associated

with alcohol use disorders.

Alcohol influences affective states and cognitive function

Acute alcohol intoxication and persistent alcohol consumption have extensive effects on

cognitive processing, executive function, and memory, while also influencing affective state and

emotion. Chronic alcohol use contributes to the emergence of negative affective symptoms,

including depression-related features such as anhedonia and sleep disturbances (McHugh and

Weiss, 2019). Additionally, individuals with co-occurring alcohol use disorder (AUD) and major

depressive disorder (MDD) are at elevated risk for relapse and tend to exhibit more severe

symptoms (Garfield, Lubman and Yücel, 2014; Destoop *et al.*, 2019). These behavioral results

suggest key interplay between AUD and MDD that may contribute to the development and

severity of these disorders.

Bidirectional regulation of the strength of Hipp-NAc connectivity plays a central role in

mediating reward-based learning and memory, as well as regulating affective state and has been

implicated in substance use and depression (Davidow *et al.*, 2016; LeGates *et al.*, 2018; Sjulson

et al., 2018; Trouche *et al.*, 2019; Heller *et al.*, 2020; Huntley *et al.*, 2020; Griffin *et al.*, 2023). We

previously showed that LTP of Hipp-NAc synapses facilitates reward-related behaviors while

disruptions in LTP is associated with anhedonia (LeGates *et al.*, 2018). Here, we show that both

acute and chronic ethanol exposure also interferes with LTP at these synapses, mirroring the

immediate and lasting effects of ethanol on synaptic plasticity previously observed within the Hipp

and at other synapses in the NAc. Given the role of Hipp-NAc synapses in mediating reward-

related behaviors, regulating affective state, and enduring alterations in response to ethanol, this

connection may represent a key site of information convergence, providing an opportunity to

identify mechanistic relationships among alcohol, mood, and cognitive processes.

Alcohol-dependent effects on synaptic function and plasticity

Alcohol has a wide range of effects on the brain, but a key factor that may underlie the ability

of alcohol to influence cognitive function and emotional state across several temporal domains is

its ability to impact synaptic function and plasticity in several relevant brain regions including the

Hipp and NAc (Zorumski, Mennerick and Izumi, 2014; Ewin *et al.*, 2019; Van Skike, Goodlett and

Matthews, 2019; Bach, Morgan, *et al.*, 2021). At intoxicating concentrations (~ 40–60 mM), acute

ethanol application can reversibly inhibit both NMDAR responses and LTP in the Hipp (Lovinger,

White and Weight, 1990; Zorumski, Mennerick and Izumi, 2014) as well as NMDAR-dependent

LTD of Hipp-NAc synapses (Kircher *et al.*, 2019). However, ethanol-mediated inhibition of

NMDARs is partial at these concentrations, and partial inhibition by selective NMDAR antagonists

does not fully block LTP (Schummers, Bentz and Browning, 1997; Izumi *et al.*, 2005). This would

suggest that NMDAR blockade does not completely explain ethanol's effect on LTP, implicating

potential involvement of additional signaling pathways. In line with this, ethanol has also been

found to block a form of NMDAR-independent LTP that is dependent on L-type VGCCs (Izumi *et*

al., 2005). We previously demonstrated Hipp-NAc LTP is mediated by sex-specific mechanisms:

in males, LTP is NMDAR-dependent while in females, LTP is NMDAR-independent and instead

relies on L-type VGCCs (Copenhaver and LeGates, 2024). Dual-effects of ethanol on NMDARs

and L-type VGCCs, as observed in the Hipp (Chandler, Harris and Crews, 1998; Izumi *et al.*,

2005), may therefore explain the deficits in LTP we observed in both males and females at Hipp-

NAc synapses.

Chronic ethanol, especially with binge-like or high-dose patterns, consistently diminishes the

ability of hippocampal neurons to elicit LTP, an effect that can persist for days to months after

ethanol withdrawal (Roberto *et al.*, 2002; Zorumski, Mennerick and Izumi, 2014). Similar effects

on plasticity have been observed in the NAc shell where CIE-induced disruption of NMDAR-

dependent LTD persists for two weeks following the last exposure (Renteria, Buske and Morrisett,

2018). Changes in glutamatergic and GABAergic receptor function and expression, which can

lead to overall disruptions in excitation/inhibition balance, are thought to be key factors mediating

the effects of chronic ethanol on synaptic plasticity (Nelson, Ur and Gruol, 2005; Qiang, Denny

and Ticku, 2007; Obara *et al.*, 2009). Additionally, chronic ethanol exposure induces

hyperexcitability, neuron loss, and morphological alterations in the Hipp (Stragier *et al.*, 2015;

Bach, Ewin, *et al.*, 2021), which may serve as additional factors that ultimately contribute to

alterations in synaptic plasticity and produce long-lasting changes that persist into abstinence.

Our data demonstrate that chronic ethanol exposure disrupts LTP at Hipp-NAc synapses

(Figure 2b-c). The absence of ethanol during recordings from slices of CIE-exposed mice

indicates that ethanol produces long-lasting alterations in synaptic plasticity that persist into

abstinence, consistent with previously reported effects described above. The plasticity deficits we

observed are also supported by recent work demonstrating chronic ethanol exposure induces

disruption to NMDAR-dependent LTD of Hipp-NAc synapses (Kircher *et al.*, 2019). This effect

coincided with synaptic alterations consistent with increased presynaptic release probability and

postsynaptic incorporation of Ca^{2+}-permeable α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic

acid receptors (AMPARs), implicating both presynaptic and postsynaptic mechanisms in ethanol-

related plasticity disruptions (Kircher *et al.*, 2019). In contrast, we found no evidence of ethanol-

induced changes in release probability, as measured by paired-pulse ratio (PPR) (Figure 1d and

2d) suggesting ethanol is likely acting primarily through postsynaptic mechanisms to interfere with

LTP. The differences in experimental outcomes may also be due to differences in study design

including the length of exposure time (3 weeks versus ~10 weeks), the method of administration

in the previous study involving CIE and two-bottle choice, or the length of the abstinence period

(72 versus 24 hours).

As discussed above, the mechanisms underlying LTP at Hipp-NAc synapses are sex-specific,

so while previously described alterations in NMDAR function and expression may explain the

effects of ethanol on LTP in males, this does not fully explain the LTP deficits we observed in

females. A considerable amount of the previous work that has provided the mechanistic insight

into ethanol's influence on synaptic function and plasticity has relied on data collected from males.

Growing evidence of sex differences in synaptic function suggests that focusing on only one sex

may obscure key molecular mechanisms underlying alcohol's effects. These synaptic differences

may be hidden, producing similar behaviors across sexes, or manifest overtly in the well-

documented disparities between males and females in alcohol sensitivity and vulnerability to AUD

(Erol and Karpyak, 2015; Flores-Bonilla and Richardson, 2020). Taken together, the ability of

ethanol to disrupt plasticity of synapses that rely on inherent sex-specific mechanisms highlights

the complexity of mechanistically dissecting its effects on synaptic function. These findings also

point to the potential for sex-specific synaptic adaptations that may contribute to differential

vulnerability and dependence.

Stress and alcohol

The ethanol-induced effects on Hipp-NAc LTP observed in this study closely resemble those

we reported previously in response to chronic stress (LeGates *et al.*, 2018). In many respects,

responses to alcohol closely resemble stress responses. For instance, both acute alcohol or

stress exposure activate the hypothalamic-pituitary-adrenal (HPA) axis, whereas repeated or

chronic exposure leads to dysregulation of HPA axis function (Richardson *et al.*, 2008; Stephens

and Wand, 2012). Chronic stress is a common precipitator of several reward-related psychiatric

disorders including depression and substance use disorders, and there is a high rate of

comorbidity between depression and AUD that is associated with greater severity and poorer

outcomes (McHugh and Weiss, 2019). Stress-induced changes in excitatory synaptic function

within key circuits involved in reward and emotion regulation are thought to be a crucial factor

underlying the pathophysiology of depression (Thompson *et al.*, 2015), and there is evidence to

suggest that drugs of misuse can elicit changes in circuit function similar to those observed in

response to stress (Saal *et al.*, 2003). These overlapping neurobiological responses highlight the

potential importance of interactions between stress, alcohol, and neural circuit function in

depression and AUD pathophysiology, though further research is needed to fully delineate

mechanisms and relevant circuits.

Hipp-NAc synapses are critical mediators of reward-related behaviors that are both stress and

alcohol sensitive (LeGates *et al.*, 2018; Kircher *et al.*, 2019; Griffin *et al.*, 2023; Lucantonio *et al.*,

2025). Activation and potentiation of these synapses is rewarding and attenuates excessive

alcohol consumption (Britt *et al.*, 2012; LeGates *et al.*, 2018; Griffin *et al.*, 2023) while chronic

stress as well as acute and chronic alcohol exposure disrupt activity-dependent synaptic plasticity

and is associated with anhedonia (LeGates *et al.*, 2018; Kircher *et al.*, 2019) (and Figures 1 and

2). Together, this underscores the crucial interplay between Hipp-NAc circuit function and

motivated behaviors, highlighting how alcohol's influence can rapidly alter behavior and induce

lasting disruptions that contribute to dependence and comorbid mental health disorders. Together,

these insights position the Hipp-NAc pathway as a critical synaptic locus underlying alcohol

dependence and depression, and a promising target for therapeutic intervention.

References

Bach, E.C., Morgan, J.W., *et al.* (2021) "Chronic Ethanol Exposures Leads to a Negative

Affective State in Female Rats That Is Accompanied by a Paradoxical Decrease in Ventral

Hippocampus Excitability," *Frontiers in Neuroscience*, 15. Available at:

https://www.frontiersin.org/articles/10.3389/fnins.2021.669075 (Accessed: November 28, 2023).

Bach, E.C., Ewin, S.E., *et al.* (2021) "Chronic intermittent ethanol promotes ventral subiculum

hyperexcitability via increases in extrinsic basolateral amygdala input and local network activity,"

Scientific Reports, 11(1), p. 8749. Available at: https://doi.org/10.1038/s41598-021-87899-0.

Barnstedt, O., Mocellin, P. and Remy, S. (2024) "A hippocampus-accumbens code guides goal-

directed appetitive behavior," *Nature Communications*, 15(1), p. 3196. Available at:

https://doi.org/10.1038/s41467-024-47361-x.

Britt, J.P. *et al.* (2012) "Synaptic and Behavioral Profile of Multiple Glutamatergic Inputs to the

Nucleus Accumbens," *Neuron*, 76, pp. 790–803.

Calhoon, G.G. and O'Donnell, P. (2013) "Closing the Gate in the Limbic Striatum: Prefrontal

Suppression of Hippocampal and Thalamic Inputs," *Neuron*, 78(1), pp. 181–190. Available at:

https://doi.org/10.1016/j.neuron.2013.01.032.

Carlezon, W.A. and Thomas, M.J. (2009) "Biological substrates of reward and aversion: A

nucleus accumbens activity hypothesis," *Neuropharmacology*, 56(SUPPL. 1), pp. 122–132.

Available at: https://doi.org/10.1016/j.neuropharm.2008.06.075.

Chandler, L.J., Harris, R.A. and Crews, F.T. (1998) "Ethanol tolerance and synaptic plasticity,"

Trends in Pharmacological Sciences, 19(12), pp. 491–495. Available at:

https://doi.org/10.1016/s0165-6147(98)01268-1.

Copenhaver, A.E. and LeGates, T.A. (2024) "Sex-Specific Mechanisms Underlie Long-Term

Potentiation at Hippocampus→Medium Spiny Neuron Synapses in the Medial Shell of the

Nucleus Accumbens," *Journal of Neuroscience*, 44(27). Available at:

https://doi.org/10.1523/JNEUROSCI.0100-24.2024.

Davidow, J.Y. *et al.* (2016) "An Upside to Reward Sensitivity: The Hippocampus Supports

Enhanced Reinforcement Learning in Adolescence," *Neuron*, 92(1), pp. 93–99. Available at:

https://doi.org/10.1016/j.neuron.2016.08.031.

Destoop, M. *et al.* (2019) "Addiction, Anhedonia, and Comorbid Mood Disorder. A Narrative

Review," *Frontiers in Psychiatry*, 10, p. 311. Available at:

https://doi.org/10.3389/fpsyt.2019.00311.

Erol, A. and Karpyak, V.M. (2015) "Sex and gender-related differences in alcohol use and its

consequences: Contemporary knowledge and future research considerations," *Drug and*

Alcohol Dependence, 156, pp. 1–13. Available at:

https://doi.org/10.1016/j.drugalcdep.2015.08.023.

Ewin, S.E. *et al.* (2019) "Chronic Intermittent Ethanol Exposure Selectively Increases Synaptic

Excitability in the Ventral Domain of the Rat Hippocampus," *Neuroscience*, 398, pp. 144–157.

Available at: https://doi.org/10.1016/j.neuroscience.2018.11.028.

Flores-Bonilla, A. and Richardson, H.N. (2020) "Sex Differences in the Neurobiology of Alcohol

Use Disorder," *Alcohol Research : Current Reviews*, 40(2), p. 04. Available at:

https://doi.org/10.35946/arcr.v40.2.04.

Floresco, S.B. and Phillips, A.G. (1999) "Dopamine and hippocampal input to the nucleus

accumbens play an essential role in the search for food in an unpredictable environment,"

Psychobiology, 27(2), pp. 277–286. Available at: https://doi.org/10.3758/BF03332121.

Francis, T.C. and Lobo, M.K. (2017) "Emerging Role for Nucleus Accumbens Medium Spiny

Neuron Subtypes in Depression," *Biological Psychiatry*, 81(8), pp. 645–653. Available at:

https://doi.org/10.1016/j.biopsych.2016.09.007.

Garfield, J.B.B., Lubman, D.I. and Yücel, M. (2014) "Anhedonia in substance use disorders: A

systematic review of its nature, course and clinical correlates," *Australian & New Zealand*

Journal of Psychiatry, 48(1), pp. 36–51. Available at:

https://doi.org/10.1177/0004867413508455.

Goto, Y. and Grace, A.A. (2008) "Limbic and cortical information processing in the nucleus

accumbens," *Trends in Neurosciences*, 31(11), pp. 552–558. Available at:

https://doi.org/10.1016/j.tins.2008.08.002.

Goto, Y. and O'Donnell, P. (2001) "Network synchrony in the nucleus accumbens in vivo," *The*

Journal of Neuroscience: The Official Journal of the Society for Neuroscience, 21(12), pp.

4498–4504. Available at: https://doi.org/10.1523/JNEUROSCI.21-12-04498.2001.

Griffin, W.C. *et al.* (2023) "Alcohol dependence and the ventral hippocampal influence on

alcohol drinking in male mice," *Alcohol*, 106, pp. 44–54. Available at:

https://doi.org/10.1016/j.alcohol.2022.10.004.

Heller, A.S. *et al.* (2020) "Association between real-world experiential diversity and positive

affect relates to hippocampal-striatal functional connectivity.," *Nature neuroscience*, 23(7), pp.

800–804. Available at: https://doi.org/10.1038/s41593-020-0636-4.

Huntley, E.D. *et al.* (2020) "Adolescent substance use and functional connectivity between the

ventral striatum and hippocampus," *Behavioural Brain Research*, 390, p. 112678. Available at:

https://doi.org/10.1016/j.bbr.2020.112678.

Ito, R. *et al.* (2008) "Functional interaction between the hippocampus and nucleus accumbens

shell is necessary for the acquisition of appetitive spatial context conditioning," *Journal of*

Neuroscience, 28(27), pp. 6950–6959. Available at: https://doi.org/10.1523/JNEUROSCI.1615-

08.2008.

Izumi, Y. *et al.* (2005) "Acute effects of ethanol on hippocampal long-term potentiation and long-

term depression are mediated by different mechanisms," *Neuroscience*, 136(2), pp. 509–517.

Available at: https://doi.org/10.1016/j.neuroscience.2005.08.002.

Kircher, D.M. *et al.* (2019) "Ethanol Experience Enhances Glutamatergic Ventral Hippocampal

Inputs to D1 Receptor-Expressing Medium Spiny Neurons in the Nucleus Accumbens Shell,"

Journal of Neuroscience, 39(13), pp. 2459–2469. Available at:

https://doi.org/10.1523/JNEUROSCI.3051-18.2019.

Koob, G.F. and Volkow, N.D. (2010) "Neurocircuitry of Addiction," *Neuropsychopharmacology*,

35(1), pp. 217–238. Available at: https://doi.org/10.1038/npp.2009.110.

Lee, S.M., Yeh, P.W.L. and Yeh, H.H. (2022) "L-Type Calcium Channels Contribute to Ethanol-

Induced Aberrant Tangential Migration of Primordial Cortical GABAergic Interneurons in the

Embryonic Medial Prefrontal Cortex," *eNeuro*, 9(1). Available at:

https://doi.org/10.1523/ENEURO.0359-21.2021.

LeGates, T.A. *et al.* (2018) "Reward behaviour is regulated by the strength of hippocampus–

nucleus accumbens synapses," *Nature*, 564(7735), pp. 258–262. Available at:

https://doi.org/10.1038/s41586-018-0740-8.

Lim, B.K. *et al.* (2012) "Anhedonia requires MC4R-mediated synaptic adaptations in nucleus

accumbens," *Nature*, 487(7406), pp. 183–189. Available at: https://doi.org/10.1038/nature11160.

Lovinger, D.M. and Abrahao, K.P. (2018) "Synaptic plasticity mechanisms common to learning

and alcohol use disorder," *Learning & Memory (Cold Spring Harbor, N.Y.)*, 25(9), pp. 425–434.

Available at: https://doi.org/10.1101/lm.046722.117.

Lovinger, D.M., White, G. and Weight, F.F. (1989) "Ethanol Inhibits NMDA-Activated Ion Current

in Hippocampal Neurons," *Science*, 243(4899), pp. 1721–1724.

Lovinger, D.M., White, G. and Weight, F.F. (1990) "NMDA receptor-mediated synaptic excitation

selectively inhibited by ethanol in hippocampal slice from adult rat," *Journal of Neuroscience*,

10(4), pp. 1372–1379. Available at: https://doi.org/10.1523/JNEUROSCI.10-04-01372.1990.

Lucantonio, F. *et al.* (2025) "Ketamine rescues anhedonia by cell-type- and input-specific

adaptations in the nucleus accumbens," *Neuron*, p. S0896627325001394. Available at:

https://doi.org/10.1016/j.neuron.2025.02.021.

Mah, S.J., Fleck, M.W. and Lindsley, T.A. (2011) "ETHANOL ALTERS CALCIUM SIGNALING IN

AXONAL GROWTH CONES," *Neuroscience*, 189, pp. 384–396. Available at:

https://doi.org/10.1016/j.neuroscience.2011.05.042.

McHugh, R.K. and Weiss, R.D. (2019) "Alcohol Use Disorder and Depressive Disorders,"

Alcohol Research : Current Reviews, 40(1), p. arcr.v40.1.01. Available at:

https://doi.org/10.35946/arcr.v40.1.01.

Nelson, T.E., Ur, C.L. and Gruol, D.L. (2005) "Chronic intermittent ethanol exposure enhances

NMDA-receptor-mediated synaptic responses and NMDA receptor expression in hippocampal

CA1 region," *Brain Research*, 1048(1), pp. 69–79. Available at:

https://doi.org/10.1016/j.brainres.2005.04.041.

Obara, I. *et al.* (2009) "Differential Effects of Chronic Ethanol Consumption and Withdrawal on Homer/Glutamate Receptor Expression in Subregions of the Accumbens and Amygdala of P Rats," *Alcoholism, clinical and experimental research*, 33(11), pp. 1924–1934. Available at: https://doi.org/10.1111/j.1530-0277.2009.01030.x.

O'Donnell, P. and Grace, A.A. (1995) "Synaptic interactions among excitatory afferents to nucleus accumbens neurons: Hippocampal gating of prefrontal cortical input," *Journal of Neuroscience*, 15(5 I), pp. 3622–3639. Available at: https://doi.org/10.1523/jneurosci.15-05-03622.1995.

Qiang, M., Denny, A.D. and Ticku, M.K. (2007) "Chronic intermittent ethanol treatment selectively alters N-methyl-D-aspartate receptor subunit surface expression in cultured cortical neurons," *Molecular Pharmacology*, 72(1), pp. 95–102. Available at: https://doi.org/10.1124/mol.106.033043.

Renteria, R. *et al.* (2017) "Selective alterations of NMDAR function and plasticity in D1 and D2 medium spiny neurons in the nucleus accumbens shell following chronic intermittent ethanol exposure," *Neuropharmacology*, 112(Pt A), pp. 164–171. Available at: https://doi.org/10.1016/j.neuropharm.2016.03.004.

Renteria, R., Buske, T.R. and Morrisett, R.A. (2018) "Long-term subregion-specific encoding of enhanced ethanol intake by D1DR medium spiny neurons of the nucleus accumbens," *Addiction Biology*, 23(2), pp. 689–698. Available at: https://doi.org/10.1111/adb.12526.

Richardson, H.N. *et al.* (2008) "Alcohol self-administration acutely stimulates the hypothalamic-pituitary-adrenal axis, but alcohol dependence leads to a dampened neuroendocrine state," *The European Journal of Neuroscience*, 28(8), pp. 1641–1653. Available at: https://doi.org/10.1111/j.1460-9568.2008.06455.x.

Roberto, M. *et al.* (2002) "Long-Term Potentiation in the Rat Hippocampus Is Reversibly

Depressed by Chronic Intermittent Ethanol Exposure," *Journal of Neurophysiology*, 87(5), pp.

2385–2397. Available at: https://doi.org/10.1152/jn.2002.87.5.2385.

Russo, S.J. and Nestler, E.J. (2013) "The brain reward circuitry in mood disorders," *Nature*

Reviews Neuroscience, 14(9), pp. 609–625. Available at: https://doi.org/10.1038/nrn3381.

Saal, D. *et al.* (2003) "Drugs of Abuse and Stress Trigger a Common Synaptic Adaptation in

Dopamine Neurons," *Neuron*, 37(4), pp. 577–582. Available at: https://doi.org/10.1016/S0896-

6273(03)00021-7.

Schummers, J., Bentz, S. and Browning, M.D. (1997) "Ethanol's inhibition of LTP may not be

mediated solely via direct effects on the NMDA receptor," *Alcoholism, Clinical and Experimental*

Research, 21(3), pp. 404–408. Available at: https://doi.org/10.1111/j.1530-0277.1997.tb03783.x.

Scofield, M.D. *et al.* (2016) "The nucleus accumbens: Mechanisms of addiction across drug

classes reflect the importance of glutamate homeostasis," *Pharmacological Reviews*, 68(3), pp.

816–871. Available at: https://doi.org/10.1124/pr.116.012484.

Sjulson, L. *et al.* (2018) "Cocaine Place Conditioning Strengthens Location-Specific

Hippocampal Coupling to the Nucleus Accumbens," *Neuron*, 98(5), pp. 926-934.e5. Available at:

https://doi.org/10.1016/j.neuron.2018.04.015.

Sosa, M., Joo, H.R. and Frank, L.M. (2020) "Dorsal and Ventral Hippocampal Sharp-Wave

Ripples Activate Distinct Nucleus Accumbens Networks," *Neuron*, 105(4), pp. 725-741.e8.

Available at: https://doi.org/10.1016/j.neuron.2019.11.022.

Stephens, M.A.C. and Wand, G. (2012) "Stress and the HPA axis: role of glucocorticoids in

alcohol dependence," *Alcohol Research: Current Reviews*, 34(4), pp. 468–483.

Stern, E.A., Jaeger, D. and Wilson, C.J. (1998) "Membrane potential synchrony of

simultaneously recorded striatal spiny neuronsin vivo," *Nature*, 394(6692), pp. 475–478.

Available at: https://doi.org/10.1038/28848.

Stragier, E. *et al.* (2015) "Brain plasticity and cognitive functions after ethanol consumption in

C57BL/6J mice," *Translational Psychiatry*, 5(12), pp. e696–e696. Available at:

https://doi.org/10.1038/tp.2015.183.

Stuber, G.D., Britt, J.P. and Bonci, A. (2012) "Optogenetic modulation of neural circuits that

underlie reward seeking," *Biological Psychiatry*, 71(12), pp. 1061–1067. Available at:

https://doi.org/10.1016/j.biopsych.2011.11.010.

Thompson, S.M. *et al.* (2015) "An excitatory synapse hypothesis of depression," *Trends in*

Neurosciences, 38(5), pp. 279–294. Available at: https://doi.org/10.1016/j.tins.2015.03.003.

Trouche, S. *et al.* (2019) "A Hippocampus-Accumbens Tripartite Neuronal Motif Guides

Appetitive Memory in Space," *Cell*, 176(6), pp. 1393-1406.e16. Available at:

https://doi.org/10.1016/j.cell.2018.12.037.

Van Skike, C.E., Goodlett, C. and Matthews, D.B. (2019) "Acute alcohol and cognition:

Remembering what it causes us to forget," *Alcohol*, 79, pp. 105–125. Available at:

https://doi.org/10.1016/j.alcohol.2019.03.006.

Volkow, N.D. and Morales, M. (2015) "The Brain on Drugs: From Reward to Addiction," *Cell*,

162(4), pp. 712–725. Available at: https://doi.org/10.1016/j.cell.2015.07.046.

Wilson, C.J. and Kawaguchi, Y. (1996) "The origins of two-state spontaneous membrane

potential fluctuations of neostriatal spiny neurons," *Journal of Neuroscience*, 16(7), pp. 2397–

2410. Available at: https://doi.org/10.1523/jneurosci.16-07-02397.1996.

Zhou, Y. *et al.* (2019) "A ventral CA1 to nucleus accumbens core engram circuit mediates

conditioned place preference for cocaine," *Nature Neuroscience*, 22(12), pp. 1986–1999.

Available at: https://doi.org/10.1038/s41593-019-0524-y.

Zorumski, C.F., Mennerick, S. and Izumi, Y. (2014) "Acute and Chronic Effects of Ethanol on

Learning-Related Synaptic Plasticity," *Alcohol (Fayetteville, N.Y.)*, 48(1), pp. 1–17. Available at:

https://doi.org/10.1016/j.alcohol.2013.09.045.

Zucca, S. and Valenzuela, C.F. (2010) "Low Concentrations of Alcohol Inhibit BDNF-Dependent

GABAergic Plasticity via L-type Ca2+ Channel Inhibition in Developing CA3 Hippocampal

Pyramidal Neurons," *The Journal of Neuroscience*, 30(19), pp. 6776–6781. Available at:

https://doi.org/10.1523/JNEUROSCI.5405-09.2010.

Figure Legends

Figure 1: Ethanol interferes with LTP at Hipp-NAc synapses a) Schematic of recording strategy

with stimulating electrode in the fornix and recording electrode in the NAc medial shell to record

Hipp-evoked EPSCs from MSNs. The illustrated parasagittal section represents a slice from

lateral +0.48mm. ac-anterior commissure f-fornix. b) Pretreatment of slices with 50mM ethanol

prevents HFS-induced LTP at Hipp-NAc synapses in males and females (top). Paired-pulse ratios

were not altered by HFS or ethanol (bottom). c) Summary data from the last 5 minutes of recording

with representative traces below. Gray traces indicate baseline and black/blue indicate post-HFS.

d) No differences in paired-pulse ratios recorded in the presence and absence of ethanol with

representative traces shown below. n= $6_{control\ male}$, $4_{control\ female}$, $5_{ethanol\ male}$, $5_{ethanol\ female}$ cells. Scale

bar for representative traces is 20pA/10ms. * p<0.05, Scale bar for representative traces =

20pA/10ms.

Figure 2: Chronic intermittent ethanol exposure disrupts LTP at Hipp-NAc synapses. a) Schematic

of experimental design to record Hipp-evoked EPSCs in the NAc shell of mice exposed to CIE or

air. ac-anterior commissure f-fornix. b) Slices taken from male and female mice exposed to CIE

show disrupted LTP as compared to air-exposed controls (top). Paired-pulse ratios were not

altered by HFS or ethanol (bottom). c) Summary data from the last 5 minutes of recording with

representative traces below. Gray traces indicate baseline and black/blue indicate post-HFS. d)

No differences in paired-pulse ratios between cells recorded from air- and CIE-exposed mice with

representative traces shown below. n= $7_{control\ male}$, $6_{control\ female}$, $6_{ethanol\ male}$, $5_{ethanol\ female}$ mice * p<0.05,

Scale bar for representative traces is 20pA/10ms.

Figure 1

Figure 2