Alcohol disrupts long-term potentiation at hippocampus-medium spiny neuron synapses in the

medial shell of the nucleus accumbens

Ashley E. Copenhaver, Ph.D.[1], Jalane R. Campbell[1], Tara A. LeGates, Ph.D.[1,2]

[1]Department of Biological Sciences, University of Maryland, Baltimore County (UMBC)

[2]Department of Pharmacology and Physiology, University of Maryland, School of Medicine

Corresponding author:

Tara LeGates, Ph.D.

1000 Hilltop Circle, Interdisciplinary Life Sciences Building 315

Baltimore, MD 21250

tlegates@umbc.edu

Support: We would like to thank Dr. Yaohui Zhu and Kaela Befano for their assistance and Dr.

Brian Mathur for his technical support and thoughtful comments on the manuscript. This work was

supported by NSF IOS2402645, T32GM144876-02, the Meyerhoff program at UMBC, and start-

up funds from UMBC.

ABSTRACT

Background: Chronic alcohol exposure is a major driver of alcohol use disorders (AUD), in part through its ability to induce maladaptive plasticity within neural circuits that regulate reward, motivation, and affect. Excitatory projections from the hippocampus (Hipp) to the nucleus accumbens (NAc) play a pivotal role in regulating reward-related behaviors, and this pathway serves as a key locus for establishing associations between rewarding stimuli and related contextual information. Regulation of the strength of Hipp-NAc synapses is critical for supporting these behaviors, and aberrant Hipp-NAc plasticity is associated with anhedonia and disrupted reward learning.

Methods: To examine acute ethanol effects, we used whole-cell electrophysiology to record Hipp-NAc synaptic plasticity in acute brain slices in the presence or absence of 50mM ethanol. To examine the effects of chronic ethanol administration, mice were exposed to ethanol vapor in a 3-week chronic intermittent ethanol (CIE) paradigm. Slices from ethanol and air exposed mice were used for whole-cell electrophysiology to record Hipp-NAc synaptic plasticity.

Results: Here, we demonstrate that acute ethanol application to *ex vivo* brain slices prevents long-term potentiation (LTP) at Hipp-NAc synapses, without altering presynaptic release probability. Furthermore, chronic intermittent exposure to ethanol abolishes LTP at these synapses, even during abstinence, indicating persistent synaptic dysfunction.

Conclusions: Together, our findings demonstrate that ethanol has immediate and long-lasting effects on Hipp-NAc plasticity. Given the behavioral relevance of these synapses, this work has important implications for the mechanisms underlying ethanol-dependent effects on reward processing and negative affective states associated with AUD.

KEYWORDS hippocampus, nucleus accumbens, synaptic plasticity, chronic intermittent ethanol, reward

INTRODUCTION

Alcohol exerts both immediate and long-lasting effects on motivated behaviors, cognitive function, and emotional regulation (Koob and Volkow, 2010). These effects arise primarily from alcohol's direct modulation of synaptic transmission and its influence over plasticity within key brain reward circuits (Lovinger and Abrahao, 2018). Although heightened plasticity in reward pathways may initially promote alcohol-seeking behaviors, persistent changes in circuit activation can ultimately impair the brain's ability to respond to natural rewards and trigger prolonged activation of stress-related neural systems. As a result, disruption of normal reward processing and persistent stress signaling are thought to underlie the negative emotional states frequently associated with chronic alcohol use and dependence. AUD with comorbid anhedonia (loss of pleasure derived from previously pleasurable activities) is associated with increased risk of relapse and poorer clinical outcomes, presenting a major obstacle to recovery (Garfield, Lubman and Yücel, 2014; Destoop *et al.*, 2019).

The nucleus accumbens (NAc) is a key mediator of motivated behaviors where both stress and rewarding experiences modulate synaptic function and plasticity to drive alterations in behavior (Carlezon and Thomas, 2009; Stuber, Britt and Bonci, 2012; Russo and Nestler, 2013). Excitatory input from several key brain regions (e.g., hippocampus, prefrontal cortex, thalamus, amygdala) has a crucial role in regulating NAc neuron activity to effectively encode reward-associated information and modulate motivated behaviors (O'Donnell and Grace, 1995; Goto and Grace, 2008; Calhoon and O'Donnell, 2013). Alterations in the strength and plasticity of these synapses as well as changes in excitability of NAc neurons have been associated with substance use disorders and depression, further demonstrating the key role of NAc activity in modulating motivated behaviors and emotional states (Volkow and Morales, 2015; Scofield *et al.*, 2016;

Francis and Lobo, 2017). For instance, chronic ethanol exposure enhances excitability and

increases NMDAR activity in NAc neurons, consistently leading to increased volitional ethanol

intake while also disrupting N-methyl-D-aspartate receptor (NMDAR)-dependent long-term

depression (LTD) (Renteria *et al.*, 2017; Kircher *et al.*, 2019). This combination of effects mirrors

the consequences of chronic stress, which similarly induces anhedonia and disrupts NMDAR-

dependent LTD within the NAc (Lim *et al.*, 2012). Together, this suggests that ethanol and stress

may converge on overlapping mechanisms within the NAc, particularly through alterations in

excitatory synaptic plasticity, to promote anhedonia and dysregulated reward processing.

Projections from the hippocampus (Hipp) provide a majority of the excitatory input to the NAc,

critically modulating neuronal activity and gating the flow of information through the region

(O'Donnell and Grace, 1995; Wilson and Kawaguchi, 1996; Stern, Jaeger and Wilson, 1998; Goto

and O'Donnell, 2001). Hipp-NAc connectivity has been tied to several behaviors related to reward

and motivation in humans and rodents including positive reinforcement learning, substance

abuse, positive affect, as well as spatial- and contextual-based reward learning (Floresco and

Phillips, 1999; Ito *et al.*, 2008; Britt *et al.*, 2012; Davidow *et al.*, 2016; LeGates *et al.*, 2018; Sjulson

et al., 2018; Trouche *et al.*, 2019; Zhou *et al.*, 2019; Heller *et al.*, 2020; Huntley *et al.*, 2020; Sosa,

Joo and Frank, 2020; Barnstedt, Mocellin and Remy, 2024). Our previous work revealed that

activity-dependent strengthening of Hipp-NAc synapses drives reward-related behaviors while

chronic stress disrupts this plasticity and induces anhedonia (LeGates *et al.*, 2018). Hipp and NAc

activity also influences alcohol-related behaviors, where reduced Hipp activity causes an increase

in alcohol consumption, and chemogenetic activation of Hipp-NAc synapses attenuates excessive

drinking (Griffin *et al.*, 2023). Furthermore, chronic alcohol exposure induces pre- and

postsynaptic changes at Hipp-NAc synapses and abolishes activity-dependent LTD (Kircher *et

al.*, 2019), again highlighting the similarly powerful effects of chronic stress and chronic alcohol

on synaptic function.

Focusing on medium spiny neurons (MSNs), which represent ~95% of the neuron population

in the NAc, we recently showed that LTP at Hipp-MSN synapses relies on NMDAR and L-type

voltage gated calcium channels (L-type VGCCs) in males and females, respectively (Copenhaver

and LeGates, 2024). Given the previously established ability of ethanol to inhibit NMDAR function

within the NAc and its ability to inhibit L-type VGCCs (Lovinger, White and Weight, 1989; Zucca

and Valenzuela, 2010; Mah, Fleck and Lindsley, 2011; Lee, Yeh and Yeh, 2022), we sought to

determine whether ethanol exposure influenced activity-dependent potentiation of Hipp-MSN

synapses. Here, we show that both acute *in vitro* and chronic *in vivo* exposure to ethanol prevents

LTP at Hipp-MSN synapses in males and females. Given the link between Hipp-NAc potentiation

in mediating reward-related behaviors, ethanol's effects on these synapses may be a key

mechanism driving maladaptive reward-related behaviors observed in response to alcohol use.

MATERIALS AND METHODS

Male and female C57BL/6J mice (8-10 weeks) were group housed (2-5 mice per cage) in a

12:12LD cycle with food and water ad libitum. We did not track the estrous cycle in female mice.

All experiments were performed in accordance with the regulations set forth by the Institutional

Animal Care and Use Committee at the University of Maryland, Baltimore County.

Ethanol Vapor Chamber Exposure

Male and female mice (8-10 weeks at the start of the experiment) were placed in acrylic

inhalation chambers (60 × 59 × 35 cm) in their home cages and exposed to chronic intermittent

ethanol (CIE) vapor or air 14 hours/day (ZT11-1), 4 days a week, for a total of 3 weeks.

Compressed air (10–20 psi) was passed through a tube submerged in 100% ethanol to volatilize.

Control vapor chambers delivered only air with no ethanol vapor. Vapor chamber ethanol

concentrations were monitored daily, and air flow was adjusted to produce ethanol concentrations

within 1.8–2.0% ethanol content as measured by a digital alcohol breath tester (BACTrack). These

conditions produce stable blood ethanol concentrations in C57BL/6J mice ranging from 100 to

200 mg/dl (Becker and Lopez, 2004). After four days in the inhalation chambers, mice underwent

a 72-h forced abstinence period. Whole cell electrophysiology was performed on week 4 after the

72-h abstinence period.

Brain Slice Preparation

Acute parasagittal slices (lateral 0.36-0.72) containing the fornix and nucleus accumbens

were prepared for whole-cell patch-clamp electrophysiology. Animals were deeply anesthetized

with isoflurane, decapitated, and brains were quickly dissected and submerged in ice-cold,

bubbled (carbogen: 95% O_2/5% CO_2) N-methyl-D-glucamine (NMDG) recovery solution

containing the following (in mM): 93 NMDG, 2.5 KCl, 1.2 NaH_2PO_4, 11 glucose, 25 $NaHCO_3$, 1.2

$MgCl_2$, and 2.4 $CaCl_2$, pH=7.3-7.4, osmolarity=300-310 mOsm. Using a vibratome (VT1000S,

Leica Microsystems), parasagittal slices (400 µm) were cut in cold, oxygenated NMDG. Slices

were transferred to 32-34°C NMDG for 7-12 minutes to recover and were then transferred to

room-temperature artificial cerebrospinal fluid (aCSF) containing the following (in mM): 120 NaCl,

3 KCl, 1.0 NaH_2PO_4, 20 glucose, 25 $NaHCO_3$, 1.5 $MgCl_2 \cdot 7H_2O$, and 2.5 $CaCl_2$, pH=7.3-7.4. Slices

were allowed to recover for 1-hour at room-temperature before beginning electrophysiological

recordings.

Whole-Cell Recordings

Slices were placed in a submersion-type recording chamber and superfused with room-

temperature aCSF (flow rate 0.5-1mL/min). Cells were visualized using a 60x water immersion

objective on a Nikon Eclipse FN-1 or Scientifica microscope. Whole-cell patch-clamp recordings

were performed using an Axopatch 200B amplifier (Axon Instruments, Molecular Devices) and a

Digidata 1550B digitizer (Axon Instruments). Patch pipettes (4-8MΩ) were made from borosilicate

glass (World Precision Instruments) using a Sutter Instruments P-97 model puller.

All recordings were performed in voltage-clamp conditions from MSNs in the NAc medial shell.

A bipolar stimulating electrode (FHC) was placed in the fornix to electrically stimulate hippocampal

axons while recording evoked excitatory postsynaptic currents (EPSC). Patch pipettes were filled

with a solution containing 130 mM K-gluconate, 5 mM KCl, 2 mM MgCl6-H2O, 10 mM HEPES, 4

mM Mg-ATP, 0.3 mM Na2-GTP, 10 mM Na2-phosphocreatine, and 1 mM EGTA; pH=7.3-7.4;

osmolarity=285-295mOsm. EPSCs were recorded from paired pulses (100ms apart) elicited

every 10s. After obtaining a stable 5-minute baseline EPSC recording, high-frequency stimulation

(HFS: four trains of 100Hz stimulation for 1s with 15s between trains while holding the cell at -

40mV) was applied, followed by a 30-minute recording of EPSCs. For acute ethanol experiments,

aCSF containing 50mM ethanol (Sigma) was washed onto slices for at least 20 minutes prior to

recording, and experiments were done in the presence of ethanol. We used the following

exclusion criteria to eliminate unhealthy cells and unreliable recordings: 1) We only proceeded

with experiments on cells with series resistances <10MΩ, 2) Cells were excluded if their series

resistance changed by >20% (comparing the resistance at the beginning and end of the

experiment), 3) Cells in poor health or poor recording status were excluded (i.e. cell partially or

fully sealed up, a decrease in holding current >100pA that is consistent with the cell dying, an

increase in jitter post HFS, and/or an increase in response failure rate to > 50%).

Data analysis

Statistical tests were performed using GraphPad Prism 9/10 software. A p-value of <0.05 was

considered statistically significant, and exact p-values can be found in the figure captions. For line

graphs, points represent the mean and error bars indicate SEM. Summary data plots display one

point for each cell, with each point representing the average EPSC amplitude from the last 5

minutes of recording. For bar graphs, the bar represents the mean and error bars indicate SEM

with individual data points overlayed. Statistical significance (p<0.05) was assessed by Two-way

ANOVA (sex x treatment) with Šídák's multiple comparisons. A three-way ANOVA (sex x treatment

x time) was used to analyze PPR data before and after HFS.

RESULTS

Acute ethanol application to *ex vivo* slices prevents Hipp-MSN LTP

To determine whether ethanol acutely influences plasticity at Hipp-MSN synapses, we

recorded HFS-induced LTP in the presence and absence of 50mM ethanol. We found that, in the

absence of ethanol, HFS induced LTP at Hipp-MSN synapses slices taken from male and female

mice with no difference in magnitude between the sexes (Figure 1a-b), similar to our previous

published results (LeGates *et al.*, 2018; Copenhaver and LeGates, 2024). However, pretreatment

with 50mM ethanol prevented LTP induction in slices taken from males and females (Figure 1a-

b; Sex x Treatment Interaction: $F_{(1,16)}$ = 0.1512, p=0.7025; Main effect of Sex: $F_{(1,16)}$ = 0.01263,

p=0.9119; Main effect of treatment: $F_{(1,16)}$ = 19.24, p=0.0005; Šídák's multiple comparisons post

hoc: $p_{(Male:control\ vs\ EtOH)}$ = 0.0175, $p_{(Female:control\ vs\ EtOH)}$ = 0.0107). We observed no difference in paired

pulse ratio (PPR) at baseline (Figure 1c; Sex x Treatment Interaction: $F_{(1,16)}$ = 0.003732, p=0.9520;

Main effect of Sex: $F_{(1,16)}$ = 1.066, p=0.3173; Main effect of treatment: $F_{(1,16)}$ = 0.2307, p=0.6375)

suggesting that application of ethanol did not alter basal release probability. Furthermore, no

change in PPR was observed following HFS irrespective of whether ethanol was present (Figure

1a, bottom; Time x Sex x Treatment Interaction: $F_{(2.590,\ 41.45)}$ = 0.5932, p=0.5992; Time x Treatment

Interaction: $F_{(2.590,41.45)}$ = 1.814, p=0.1661; Time x Sex Interaction: $F_{(2.590,\ 41.45)}$ = 0.3843, p=0.7359),

suggesting that HFS did not elicit changes in presynaptic neurotransmitter release probability.

CIE disrupts Hipp-MSN LTP

To determine whether chronic exposure to ethanol leads to changes in Hipp-MSN plasticity, we used CIE to ethanol vapor for controlled delivery and dosing of ethanol over a 3-week period. Control mice were exposed to air. Following at least 72 hours of forced abstinence, slices from CIE-exposed mice demonstrated a deficit in HFS-induced LTP relative to air-exposed controls, which showed typical LTP (Figure 2a-b; Sex x Treatment Interaction: $F_{(1,20)}$ = 0.03737, p=0.8487; Main effect of Sex: $F_{(1,20)}$ = 0.1118, p=0.7415; Main effect of treatment: $F_{(1,20)}$ = 12.53, p=0.002; Šídák's multiple comparisons post hoc: $p_{(Male:control \, vs \, EtOH)}$ = 0.0444, $p_{(Female:control \, vs \, EtOH)}$ = 0.0390). This effect was consistent in both male and female mice. Comparison of baseline PPRs revealed no difference between CIE mice and air exposed controls (Figure 2c; Sex x Treatment Interaction: $F_{(1,20)}$ = 0.2254, p=0.6401; Main effect of Sex: $F_{(1,20)}$ = 0.0008649, p=0.9768; Main effect of treatment: $F_{(1,20)}$ = 1.146, p=0.2971), suggesting that CIE did not induce changes in release probability. Similar to the data shown in Figure 1, HFS did not induce changes in PPR regardless of ethanol exposure or sex (Figure 2a bottom; Time x Sex x Treatment Interaction: $F_{(6,20)}$ = 1.600, p=0.1529; Time x Treatment Interaction: $F_{(6,20)}$ = 0.6492, p=0.6906; Time x Sex Interaction: $F_{(6,20)}$ = 1.067, p=3862), demonstrating a lack of effect on presynaptic function.

DISCUSSION

Our data demonstrate that acute application of ethanol to *ex vivo* brain slices disrupts HFS-induced LTP at Hipp-MSN synapses. We also found that chronic exposure to ethanol *in vivo* interfered with HFS-induced LTP at Hipp-MSN synapses, an effect observed after a 72-hour abstinence period when ethanol was no longer present. Neither treatment led to a change in basal release probability, suggesting that ethanol may be acting postsynaptically to interfere with LTP. Given the previously described role for Hipp-NAc synaptic strength in mediating reward-related

211   behaviors, ethanol's impact on Hipp-MSN plasticity has important implications for understanding

212   the neurobiological basis for disrupted reward processing and negative mood states associated

213   with alcohol use disorders.

Alcohol influences affective states and cognitive function

Acute alcohol intoxication and persistent alcohol consumption have extensive effects on

cognitive processing, executive function, and memory, while also influencing affective state and

emotional processing. Many of these alcohol-dependent changes in behavior are closely linked

to altered activity within key brain regions such as the Hipp and NAc. (Zorumski, Mennerick and

Izumi, 2014; Ewin *et al.*, 2019; Van Skike, Goodlett and Matthews, 2019; Bach *et al.*, 2021).

Hippocampal input to the NAc plays a central role in mediating reward-based learning and

memory, as well as regulating affective state. Furthermore, the bidirectional regulation of Hipp-

NAc connectivity is a critical regulator of reward-related behaviors and has been implicated in

substance use and depression (Davidow *et al.*, 2016; LeGates *et al.*, 2018; Sjulson *et al.*, 2018;

Trouche *et al.*, 2019; Heller *et al.*, 2020; Huntley *et al.*, 2020; Griffin *et al.*, 2023). We previously

showed that LTP of Hipp-NAc synapses facilitates reward-related behaviors while disruptions in

LTP is associated with anhedonia (LeGates *et al.*, 2018). Here, we show that both acute and

chronic ethanol exposure interfere with this behaviorally relevant LTP, mirroring the immediate

and lasting effects of ethanol on synaptic plasticity observed within the Hipp and at other synapses

in the NAc.

Alcohol-dependent effects on synaptic function

The ability of alcohol to exert effects across several temporal domains is likely due to its ability

to impact synaptic function and plasticity in several brain regions that regulate cognitive function

and emotional state such as the Hipp and NAc. At intoxicating concentrations (~ 40–60 mM),

ethanol can reversibly inhibit both NMDAR responses and LTP in the Hipp (Chandler, Harris and

Crews, 1998). However, the ethanol-mediated inhibition of NMDARs is partial at these

concentrations, and partial inhibition by selective NMDAR antagonists does not fully block LTP

(Schummers, Bentz and Browning, 1997; Izumi *et al.*, 2005). This would suggest that blockade of

NMDARs may not completely explain ethanol's effect on LTP and may involve alterations of

additional pathways. For instance, ethanol has also been found to block NMDAR-independent

LTP that instead relies on L-type VGCCs (Izumi *et al.*, 2005). Chronic ethanol, especially with

binge-like or high-dose patterns, consistently diminishes the ability of hippocampal neurons to

elicit LTP, an effect that can persist for days to months after ethanol withdrawal, though some

recovery is possible with prolonged abstinence (Roberto *et al.*, 2002). This is thought to be driven

by alterations in NMDAR-mediated responses and receptor expression as well as altered

$GABA_A$ receptor function, leading to an overall disruption in the balance required for normal

synaptic plasticity (Nelson, Ur and Gruol, 2005). Depending on the exposure model, chronic

ethanol can also induce changes in signaling pathways (like MAPK) and neurotrophins (BDNF,

NGF) as well as neuron loss and morphological alterations in the Hipp (Stragier *et al.*, 2015).

In the NAc shell, ethanol dose-dependently inhibits NMDAR-dependent LTD through a

mechanism involving GluN2B-containing NMDARs and Dopamine-1 receptor signaling while

chronic ethanol reduces NMDAR function and depresses NMDAR-dependent LTP in the NAc

(Jeanes, Buske and Morrisett, 2014). Post-withdrawal, GluN2B and mGluR1 expression

increases, altering synaptic signaling (Renteria *et al.*, 2017). Chronic intermittent ethanol (CIE)

exposure followed by withdrawal has time-dependent effects on synaptic plasticity: 24-hours after

the last ethanol exposure LTD converts to NMDAR-dependent LTP while 72-hours of abstinence

abolishes both LTD and LTP (Renteria *et al.*, 2017). These neuroadaptations promote aberrant

learning, reinforcing alcohol-seeking behaviors and contributing to dependence.

Our data demonstrate that acute ethanol treatment applied to slices and CIE *in vivo* prevent

LTP at Hipp-MSN synapses. The absence of ethanol during recordings from slices of CIE-

exposed mice suggests that ethanol has both acute and longer lasting effects on plasticity that

persist into abstinence. We previously found that the mechanisms underlying Hipp-MSN plasticity

are sex-specific where LTP in males is NMDAR-dependent while in females, LTP is dependent

on L-type VGCCs. Ethanol's ability to prevent LTP in both sexes may therefore be due to

combined effects on NMDARs and L-type VGCCs, in line with prior observations in the Hipp

(Chandler, Harris and Crews, 1998; Izumi *et al.*, 2005). Previous work also found that acute and

chronic ethanol treatment can interfere with NMDAR-dependent LTD of Hipp-MSN synapses

(Kircher *et al.*, 2019), providing further support that ethanol may be altering NMDAR function at

Hipp-NAc synapses. We did not observe ethanol-induced changes in release probability as

measured by PPR (Figure 1c and 2c). This differs from previous work where chronic ethanol

exposure through vapor and two-bottle choice induced presynaptic changes increasing Hipp input

to the NAc as well as postsynaptic changes involving preferential insertion of Ca^{2+}-permeable α-

amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors (AMPARs) (Kircher *et al.*, 2019).

The variations in results between the two studies may be due to the differences in the length of

exposure time (3 weeks versus ~10 weeks) or the method of administration involving CIE and

two-bottle choice.

Stress and alcohol

The ethanol-induced effects on Hipp-NAc LTP observed in this study closely resemble those

we reported previously in response to chronic stress (LeGates *et al.*, 2018). In many respects,

responses to alcohol closely resemble stress responses. For instance, both acute alcohol or

stress exposure activate the hypothalamic-pituitary-adrenal (HPA) axis, whereas repeated or

chronic exposure leads to dysregulation of HPA axis function (Richardson *et al.*, 2008; Stephens

and Wand, 2012). Chronic stress is a common precipitator of several reward-related psychiatric

disorders including depression and substance use disorders, and there is a high rate of

comorbidity between depression and AUD that is associated with greater severity and poorer

outcomes (McHugh and Weiss, 2019). Stress-induced changes in excitatory synaptic function

within key circuits involved in reward and emotion regulation are thought to be a crucial factor

underlying the pathophysiology of depression (Thompson *et al.*, 2015), and there is evidence to

suggest that drugs of misuse can elicit changes in circuit function similar to those observed in

response to stress (Saal *et al.*, 2003). These overlapping neurobiological responses highlight the

potential importance of interactions between stress, alcohol, and neural circuit function in

depression and AUD pathophysiology, though further research is needed to fully delineate

mechanisms and relevant circuits.

Hipp-NAc synapses are critical mediators of reward-related behaviors that are both stress and

alcohol sensitive (LeGates *et al.*, 2018; Kircher *et al.*, 2019; Griffin *et al.*, 2023; Lucantonio *et al.*,

2025). Activation and potentiation of these synapses is rewarding and attenuates excessive

alcohol consumption (Britt *et al.*, 2012; LeGates *et al.*, 2018; Griffin *et al.*, 2023) while chronic

stress as well as acute and chronic alcohol exposure disrupt activity-dependent synaptic plasticity

and is associated with anhedonia (LeGates *et al.*, 2018; Kircher *et al.*, 2019) (and Figures 1 and

2). Together, this underscores the crucial interplay between Hipp-NAc circuit function and

motivated behaviors, highlighting how alcohol's influence can rapidly alter behavior and induce

lasting disruptions that contribute to dependence and comorbid mental health disorders. Together,

these insights position the Hipp-NAc pathway as a critical synaptic locus underlying alcohol

dependence and depression, and a promising target for therapeutic intervention.

References

Bach, E.C. *et al.* (2021) 'Chronic Ethanol Exposures Leads to a Negative Affective State in

Female Rats That Is Accompanied by a Paradoxical Decrease in Ventral Hippocampus

Excitability', *Frontiers in Neuroscience*, 15. Available at:

https://www.frontiersin.org/articles/10.3389/fnins.2021.669075 (Accessed: 28 November 2023).

Barnstedt, O., Mocellin, P. and Remy, S. (2024) 'A hippocampus-accumbens code guides goal-

directed appetitive behavior', *Nature Communications*, 15(1), p. 3196. Available at:

https://doi.org/10.1038/s41467-024-47361-x.

Becker, H.C. and Lopez, M.F. (2004) 'Increased ethanol drinking after repeated chronic ethanol

exposure and withdrawal experience in C57BL/6 mice', *Alcoholism, Clinical and Experimental*

Research, 28(12), pp. 1829–1838. Available at:

https://doi.org/10.1097/01.alc.0000149977.95306.3a.

Britt, J.P. *et al.* (2012) 'Synaptic and Behavioral Profile of Multiple Glutamatergic Inputs to the

Nucleus Accumbens', *Neuron*, 76, pp. 790–803.

Calhoon, G.G. and O'Donnell, P. (2013) 'Closing the Gate in the Limbic Striatum: Prefrontal

Suppression of Hippocampal and Thalamic Inputs', *Neuron*, 78(1), pp. 181–190. Available at:

https://doi.org/10.1016/j.neuron.2013.01.032.

Carlezon, W.A. and Thomas, M.J. (2009) 'Biological substrates of reward and aversion: A

nucleus accumbens activity hypothesis', *Neuropharmacology*, 56(SUPPL. 1), pp. 122–132.

Available at: https://doi.org/10.1016/j.neuropharm.2008.06.075.

Chandler, L.J., Harris, R.A. and Crews, F.T. (1998) 'Ethanol tolerance and synaptic plasticity',

Trends in Pharmacological Sciences, 19(12), pp. 491–495. Available at:

https://doi.org/10.1016/s0165-6147(98)01268-1.

Copenhaver, A.E. and LeGates, T.A. (2024) 'Sex-Specific Mechanisms Underlie Long-Term

Potentiation at Hippocampus→Medium Spiny Neuron Synapses in the Medial Shell of the

Nucleus Accumbens', *Journal of Neuroscience*, 44(27). Available at:

https://doi.org/10.1523/JNEUROSCI.0100-24.2024.

Davidow, J.Y. *et al.* (2016) 'An Upside to Reward Sensitivity: The Hippocampus Supports

Enhanced Reinforcement Learning in Adolescence', *Neuron*, 92(1), pp. 93–99. Available at:

https://doi.org/10.1016/j.neuron.2016.08.031.

Destoop, M. *et al.* (2019) 'Addiction, Anhedonia, and Comorbid Mood Disorder. A Narrative

Review', *Frontiers in Psychiatry*, 10, p. 311. Available at:

https://doi.org/10.3389/fpsyt.2019.00311.

Ewin, S.E. *et al.* (2019) 'Chronic Intermittent Ethanol Exposure Selectively Increases Synaptic

Excitability in the Ventral Domain of the Rat Hippocampus', *Neuroscience*, 398, pp. 144–157.

Available at: https://doi.org/10.1016/j.neuroscience.2018.11.028.

Floresco, S.B. and Phillips, A.G. (1999) 'Dopamine and hippocampal input to the nucleus

accumbens play an essential role in the search for food in an unpredictable environment',

Psychobiology, 27(2), pp. 277–286. Available at: https://doi.org/10.3758/BF03332121.

Francis, T.C. and Lobo, M.K. (2017) 'Emerging Role for Nucleus Accumbens Medium Spiny

Neuron Subtypes in Depression', *Biological Psychiatry*, 81(8), pp. 645–653. Available at:

https://doi.org/10.1016/j.biopsych.2016.09.007.

Garfield, J.B.B., Lubman, D.I. and Yücel, M. (2014) 'Anhedonia in substance use disorders: A

systematic review of its nature, course and clinical correlates', *Australian & New Zealand*

Journal of Psychiatry, 48(1), pp. 36–51. Available at:

https://doi.org/10.1177/0004867413508455.

Goto, Y. and Grace, A.A. (2008) 'Limbic and cortical information processing in the nucleus

accumbens', *Trends in Neurosciences*, 31(11), pp. 552–558. Available at:

https://doi.org/10.1016/j.tins.2008.08.002.

Goto, Y. and O'Donnell, P. (2001) 'Network synchrony in the nucleus accumbens in vivo', *The*

Journal of Neuroscience: The Official Journal of the Society for Neuroscience, 21(12), pp.

4498–4504. Available at: https://doi.org/10.1523/JNEUROSCI.21-12-04498.2001.

Griffin, W.C. *et al.* (2023) 'Alcohol dependence and the ventral hippocampal influence on alcohol

drinking in male mice', *Alcohol*, 106, pp. 44–54. Available at:

https://doi.org/10.1016/j.alcohol.2022.10.004.

Heller, A.S. *et al.* (2020) 'Association between real-world experiential diversity and positive

affect relates to hippocampal-striatal functional connectivity.', *Nature neuroscience*, 23(7), pp.

800–804. Available at: https://doi.org/10.1038/s41593-020-0636-4.

Huntley, E.D. *et al.* (2020) 'Adolescent substance use and functional connectivity between the

ventral striatum and hippocampus', *Behavioural Brain Research*, 390, p. 112678. Available at:

https://doi.org/10.1016/j.bbr.2020.112678.

Ito, R. *et al.* (2008) 'Functional interaction between the hippocampus and nucleus accumbens

shell is necessary for the acquisition of appetitive spatial context conditioning', *Journal of*

Neuroscience, 28(27), pp. 6950–6959. Available at: https://doi.org/10.1523/JNEUROSCI.1615-

08.2008.

Izumi, Y. *et al.* (2005) 'Acute effects of ethanol on hippocampal long-term potentiation and long-term depression are mediated by different mechanisms', *Neuroscience*, 136(2), pp. 509–517. Available at: https://doi.org/10.1016/j.neuroscience.2005.08.002.

Jeanes, Z.M., Buske, T.R. and Morrisett, R.A. (2014) 'Cell type-specific synaptic encoding of ethanol exposure in the nucleus accumbens shell', *Neuroscience*, 277, pp. 184–195. Available at: https://doi.org/10.1016/j.neuroscience.2014.06.063.

Kircher, D.M. *et al.* (2019) 'Ethanol Experience Enhances Glutamatergic Ventral Hippocampal Inputs to D1 Receptor-Expressing Medium Spiny Neurons in the Nucleus Accumbens Shell', *Journal of Neuroscience*, 39(13), pp. 2459–2469. Available at: https://doi.org/10.1523/JNEUROSCI.3051-18.2019.

Koob, G.F. and Volkow, N.D. (2010) 'Neurocircuitry of Addiction', *Neuropsychopharmacology*, 35(1), pp. 217–238. Available at: https://doi.org/10.1038/npp.2009.110.

Lee, S.M., Yeh, P.W.L. and Yeh, H.H. (2022) 'L-Type Calcium Channels Contribute to Ethanol-Induced Aberrant Tangential Migration of Primordial Cortical GABAergic Interneurons in the Embryonic Medial Prefrontal Cortex', *eNeuro*, 9(1). Available at: https://doi.org/10.1523/ENEURO.0359-21.2021.

LeGates, T.A. *et al.* (2018) 'Reward behaviour is regulated by the strength of hippocampus–nucleus accumbens synapses', *Nature*, 564(7735), pp. 258–262. Available at: https://doi.org/10.1038/s41586-018-0740-8.

Lim, B.K. *et al.* (2012) 'Anhedonia requires MC4R-mediated synaptic adaptations in nucleus accumbens', *Nature*, 487(7406), pp. 183–189. Available at: https://doi.org/10.1038/nature11160.

Lovinger, D.M. and Abrahao, K.P. (2018) 'Synaptic plasticity mechanisms common to learning

and alcohol use disorder', *Learning & Memory (Cold Spring Harbor, N.Y.)*, 25(9), pp. 425–434.

Available at: https://doi.org/10.1101/lm.046722.117.

Lovinger, D.M., White, G. and Weight, F.F. (1989) 'Ethanol Inhibits NMDA-Activated Ion Current

in Hippocampal Neurons', *Science*, 243(4899), pp. 1721–1724.

Lucantonio, F. *et al.* (2025) 'Ketamine rescues anhedonia by cell-type- and input-specific

adaptations in the nucleus accumbens', *Neuron*, p. S0896627325001394. Available at:

https://doi.org/10.1016/j.neuron.2025.02.021.

Mah, S.J., Fleck, M.W. and Lindsley, T.A. (2011) 'ETHANOL ALTERS CALCIUM SIGNALING IN

AXONAL GROWTH CONES', *Neuroscience*, 189, pp. 384–396. Available at:

https://doi.org/10.1016/j.neuroscience.2011.05.042.

McHugh, R.K. and Weiss, R.D. (2019) 'Alcohol Use Disorder and Depressive Disorders',

Alcohol Research : Current Reviews, 40(1), p. arcr.v40.1.01. Available at:

https://doi.org/10.35946/arcr.v40.1.01.

Nelson, T.E., Ur, C.L. and Gruol, D.L. (2005) 'Chronic intermittent ethanol exposure enhances

NMDA-receptor-mediated synaptic responses and NMDA receptor expression in hippocampal

CA1 region', *Brain Research*, 1048(1), pp. 69–79. Available at:

https://doi.org/10.1016/j.brainres.2005.04.041.

O'Donnell, P. and Grace, A.A. (1995) 'Synaptic interactions among excitatory afferents to

nucleus accumbens neurons: Hippocampal gating of prefrontal cortical input', *Journal of*

Neuroscience, 15(5 I), pp. 3622–3639. Available at: https://doi.org/10.1523/jneurosci.15-05-

03622.1995.

Renteria, R. *et al.* (2017) 'Selective alterations of NMDAR function and plasticity in D1 and D2

medium spiny neurons in the nucleus accumbens shell following chronic intermittent ethanol

exposure', *Neuropharmacology*, 112(Pt A), pp. 164–171. Available at:

https://doi.org/10.1016/j.neuropharm.2016.03.004.

Richardson, H.N. *et al.* (2008) 'Alcohol self-administration acutely stimulates the hypothalamic-

pituitary-adrenal axis, but alcohol dependence leads to a dampened neuroendocrine state', *The*

European Journal of Neuroscience, 28(8), pp. 1641–1653. Available at:

https://doi.org/10.1111/j.1460-9568.2008.06455.x.

Roberto, M. *et al.* (2002) 'Long-Term Potentiation in the Rat Hippocampus Is Reversibly

Depressed by Chronic Intermittent Ethanol Exposure', *Journal of Neurophysiology*, 87(5), pp.

2385–2397. Available at: https://doi.org/10.1152/jn.2002.87.5.2385.

Russo, S.J. and Nestler, E.J. (2013) 'The brain reward circuitry in mood disorders', *Nature*

Reviews Neuroscience, 14(9), pp. 609–625. Available at: https://doi.org/10.1038/nrn3381.

Saal, D. *et al.* (2003) 'Drugs of Abuse and Stress Trigger a Common Synaptic Adaptation in

Dopamine Neurons', *Neuron*, 37(4), pp. 577–582. Available at: https://doi.org/10.1016/S0896-

6273(03)00021-7.

Schummers, J., Bentz, S. and Browning, M.D. (1997) 'Ethanol's inhibition of LTP may not be

mediated solely via direct effects on the NMDA receptor', *Alcoholism, Clinical and Experimental*

Research, 21(3), pp. 404–408. Available at: https://doi.org/10.1111/j.1530-0277.1997.tb03783.x.

Scofield, M.D. *et al.* (2016) 'The nucleus accumbens: Mechanisms of addiction across drug

classes reflect the importance of glutamate homeostasis', *Pharmacological Reviews*, 68(3), pp.

816–871. Available at: https://doi.org/10.1124/pr.116.012484.

Sjulson, L. *et al.* (2018) 'Cocaine Place Conditioning Strengthens Location-Specific

Hippocampal Coupling to the Nucleus Accumbens', *Neuron*, 98(5), pp. 926-934.e5. Available at:

https://doi.org/10.1016/j.neuron.2018.04.015.

Sosa, M., Joo, H.R. and Frank, L.M. (2020) 'Dorsal and Ventral Hippocampal Sharp-Wave

Ripples Activate Distinct Nucleus Accumbens Networks', *Neuron*, 105(4), pp. 725-741.e8.

Available at: https://doi.org/10.1016/j.neuron.2019.11.022.

Stephens, M.A.C. and Wand, G. (2012) 'Stress and the HPA axis: role of glucocorticoids in

alcohol dependence', *Alcohol Research: Current Reviews*, 34(4), pp. 468–483.

Stern, E.A., Jaeger, D. and Wilson, C.J. (1998) 'Membrane potential synchrony of

simultaneously recorded striatal spiny neuronsin vivo', *Nature*, 394(6692), pp. 475–478.

Available at: https://doi.org/10.1038/28848.

Stragier, E. *et al.* (2015) 'Brain plasticity and cognitive functions after ethanol consumption in

C57BL/6J mice', *Translational Psychiatry*, 5(12), pp. e696–e696. Available at:

https://doi.org/10.1038/tp.2015.183.

Stuber, G.D., Britt, J.P. and Bonci, A. (2012) 'Optogenetic modulation of neural circuits that

underlie reward seeking', *Biological Psychiatry*, 71(12), pp. 1061–1067. Available at:

https://doi.org/10.1016/j.biopsych.2011.11.010.

Thompson, S.M. *et al.* (2015) 'An excitatory synapse hypothesis of depression', *Trends in*

Neurosciences, 38(5), pp. 279–294. Available at: https://doi.org/10.1016/j.tins.2015.03.003.

Trouche, S. *et al.* (2019) 'A Hippocampus-Accumbens Tripartite Neuronal Motif Guides

Appetitive Memory in Space', *Cell*, 176(6), pp. 1393-1406.e16. Available at:

https://doi.org/10.1016/j.cell.2018.12.037.

Van Skike, C.E., Goodlett, C. and Matthews, D.B. (2019) 'Acute alcohol and cognition:

Remembering what it causes us to forget', *Alcohol*, 79, pp. 105–125. Available at:

https://doi.org/10.1016/j.alcohol.2019.03.006.

Volkow, N.D. and Morales, M. (2015) 'The Brain on Drugs: From Reward to Addiction', *Cell*,

162(4), pp. 712–725. Available at: https://doi.org/10.1016/j.cell.2015.07.046.

Wilson, C.J. and Kawaguchi, Y. (1996) 'The origins of two-state spontaneous membrane

potential fluctuations of neostriatal spiny neurons', *Journal of Neuroscience*, 16(7), pp. 2397–

2410. Available at: https://doi.org/10.1523/jneurosci.16-07-02397.1996.

Zhou, Y. *et al.* (2019) 'A ventral CA1 to nucleus accumbens core engram circuit mediates

conditioned place preference for cocaine', *Nature Neuroscience*, 22(12), pp. 1986–1999.

Available at: https://doi.org/10.1038/s41593-019-0524-y.

Zorumski, C.F., Mennerick, S. and Izumi, Y. (2014) 'Acute and Chronic Effects of Ethanol on

Learning-Related Synaptic Plasticity', *Alcohol (Fayetteville, N.Y.)*, 48(1), pp. 1–17. Available at:

https://doi.org/10.1016/j.alcohol.2013.09.045.

Zucca, S. and Valenzuela, C.F. (2010) 'Low Concentrations of Alcohol Inhibit BDNF-Dependent

GABAergic Plasticity via L-type Ca2+ Channel Inhibition in Developing CA3 Hippocampal

Pyramidal Neurons', *The Journal of Neuroscience*, 30(19), pp. 6776–6781. Available at:

https://doi.org/10.1523/JNEUROSCI.5405-09.2010.

Figure Legends

Figure 1: Ethanol interferes with LTP at Hipp-MSN synapses a) Pretreatment of slices with 50mM ethanol prevents HFS-induced LTP at Hipp-MSN synapses in males and females (top). Paired-pulse ratios were not altered by HFS or ethanol (bottom). b) Summary data from the last 5 minutes of recording with representative traces below. c) No differences in paired-pulse ratios recorded in the presence and absence of ethanol with representative traces shown below. Scale bar for representative traces is 20pA/10ms. * $p < 0.05$, Scale bar for representative traces = 20pA/10ms.

Figure 2: Chronic intermittent ethanol exposure disrupts LTP at Hipp-MSN synapses. a) Slices taken from male and female mice exposed to CIE show disrupted LTP as compared to air-exposed controls (top). Paired-pulse ratios were not altered by HFS or ethanol (bottom). b) Summary data from the last 5 minutes of recording with representative traces below. c) No differences in paired-pulse ratios between cells recorded from air- and CIE-exposed mice with representative traces shown below. * $p < 0.05$, Scale bar for representative traces is 20pA/10ms.

Figure 1

Figure 2