Twitching motility suppressors reveal a role for FimX in type IV pilus extension dynamics

Nathan Roberge[1], Nathan Yuen[1], Hanjeong Harvey[1], Taylor J. Ellison[2], Courtney K. Ellison[2],

and Lori L. Burrows[1]*

[1] Department of Biochemistry and Biomedical Sciences, and the Michael G. DeGroote Institute

for Infectious Disease Research, McMaster University, Hamilton, ON, Canada, L8S4K1

[2] Department of Microbiology, University of Georgia, Athens, GA, 30602

*Corresponding author

lori.burrows@mcmaster.ca

Contributions

NR: Conceptualization, Investigation, Writing – original draft, Writing – review and editing

NY: Investigation, Writing – review and editing

HH: Resources, Writing – review and editing

TJE: Investigation, Writing – review and editing

CKE: Conceptualization, Investigation, Writing – review and editing

LLB: Conceptualization, Funding acquisition, Writing – original draft, Writing – review and
editing

Abstract

In *Pseudomonas aeruginosa,* retractable protein filaments called type IV pili (T4P)

facilitate surface adherence, sensing, and directional movement known as twitching motility.

T4P are necessary for the bacteria to engage in surface-associated behaviors, including

establishing acute infections. Pilus extension is driven by the hexameric ATPase, PilB, at the

base of the T4P nanomachine in coordination with various protein regulatory effectors. The

cyclic-di-GMP binding protein, FimX, works with PilB to mediate normal extension processes,

though how this effector controls pilus assembly remains unclear. To explore the role of FimX in

T4P function, we leveraged the significant Δ*fimX* twitching motility deficit to screen for mutants

capable of overcoming this phenotype. We identified suppressor mutations that increase

twitching in Δ*fimX* background, mapping primarily to cyclic-AMP homeostatic machinery or to

PilB, the FimX target. Distinct suppressor mutations in PilB increased ATP hydrolysis *in vitro*

and this activity was subject to modulation by FimX. Using microscopy to monitor the extension

dynamics of fluorescently labelled T4P, we showed that Δ*fimX* mutants produce slow-to-extend,

short pili, a phenotype that is rescued by mutations enhancing PilB ATP hydrolysis and/or re-

introduction of FimX. Together, these data implicate FimX as a regulator of PilB enzymatic

function, potentially enabling *P. aeruginosa* to fine-tune pilus extension dynamics in response to

environmental cues.

Summary

Type IV pili enable *Pseudomonas aeruginosa* to attach to surfaces, move (twitch), and

form biofilms. Pilus extension is powered by the motor protein PilB, which is regulated by other

factors, including FimX, a protein that binds cyclic-di-GMP. Although FimX is important for

twitching, how it influences PilB was unclear. We deleted *fimX*, which severely reduces motility,

and searched for mutants that regained movement. We identified two types: some had mutations

in PilB that increased its ATPase activity, allowing it to function without FimX, while others

affected the cyclic-AMP signaling pathway and increased overall production of pilus

components, showing that motility can also be improved through changes in quantity versus

quality. Our results suggest that FimX normally fine-tunes PilB enzymatic activity, enabling

dynamic control of pilus extension in response to surface signals. This work helps explain how

P. aeruginosa adapts to different environments, a process crucial for infection and biofilm

development.

Introduction

   Pseudomonas aeruginosa deploys type IV pili (T4P) to adhere to and spread across

surfaces (1, 2). Directed bacterial movement, or twitching motility, is powered by repeated

cycles of protein filament extension, surface attachment, and retraction to pull cells along a

surface (3–5). Pilus function is coordinated through the action of four distinct protein

subcomplexes which together comprise the T4P nanomachine (Fig 1A) (6). T4P also allow cells

to sense surface contact and assist in the transition from planktonic to sessile lifestyles (5, 7–9),

facilitating microcolony formation, and eventually, biofilm production (10, 11). Engagement of

the extracellular filament with a surface induces an as-yet undefined signal that is propagated to

the inner membrane embedded Pil-Chp regulatory network, which in turn stimulates CyaB-

dependent synthesis of the secondary messenger molecule, cyclic adenosine monophosphate

(cAMP) (Fig 1A) (5, 7, 12). High intracellular cAMP activates the transcription factor Vfr to

modulate the expression of over 300 genes, including those encoding T4P machinery

components (13), with cAMP homeostasis maintained by the action of the CpdA

phosphodiesterase (14–16). The cascade of secondary messenger signaling culminates in

elevated cyclic diguanylate monophosphate (cdGMP) (16), which in many bacteria is associated

with sessile phenotypes such as biofilm formation (17). This increase in cdGMP potentially

occurs through stimulation of T4P-associated diguanylate cyclases (DGCs) like SadC (18, 19),

and/or through surface sensing by pilus tip-associated proteins (6, 16, 20, 21).

Pilus extension is an essential first step in filament biogenesis. Extended pili can interact

with external substrates, ultimately triggering downstream signaling and facilitating motility (2,

8, 22, 23). Extension is driven by the hexameric ATPase PilB (1, 24–26) in conjunction with

regulatory effectors PilZ (27, 28) and the cdGMP-binding protein FimX (27, 29–31) (Fig 1A),

both of which are necessary for normal pilus assembly. The role of FimX in pilus extension is

not well understood, with studies of *P. aeruginosa* FimX and its homologues in other species

implicating a variety of disparate functions (27, 30, 32, 33). In *P. aeruginosa*, FimX interaction

with PilB is contingent upon cdGMP binding by FimX's EVL motif in the C-terminal EAL

domain. *P. aeruginosa* FimX mutants that are unable to bind the secondary messenger are

defective in surface piliation (30, 31), twitching motility (29–31), and ATPase localization (27,

30). Some of these phenotypes can be overcome through suppressor mutations that increase

cellular cdGMP levels. Those studies suggest that FimX is responsible for maintaining PilB

functional efficiency over broad concentrations of cdGMP (31). On the other hand, more recent

work showed selective trafficking of the PilB-FimX subcomplex to the leading poles of

twitching cells. This localization promotes biased pilus extension at a single pole, resulting in

increased Pil-Chp network signaling at the same pole. The positive feedback loop was proposed

to ensure directional movement of the rod-shaped cells in response to PilB localization. Further,

PilB polar localization was dependent upon FimX and various cytoplasmic components of the

Pil-Chp network, suggesting that FimX may also be a cdGMP-binding polar localization factor

that coordinates optimal PilB subcellular distribution (8, 34).

How then, do these proposed functions intersect and what is the role of FimX in *P.*

aeruginosa pilus function and/or signaling? To shed light on how FimX controls pilus function,

we leveraged the Δ*fimX* twitching motility deficit to select for suppressor mutants capable of

overcoming loss of its input. Our screen revealed two classes of mutants with increased motility

in the absence of FimX. These mutations were primarily in the Pil-Chp network responsible for

modulating cAMP levels or PilB, the interaction partner of FimX. We focused on the PilB

mutations, which notably were not clustered but rather mapped to various regions of the protein.

We characterized two distinct examples and demonstrated that each significantly increased PilB

ATP hydrolysis *in vitro*. Phenotypically, this elevated activity translated to increased rates of

extension, resulting in longer pili. Both ATPase activity and pilus extension rates were subject to

FimX-dependent modulation. Therefore, FimX modulates PilB enzymatic efficiency to produce

longer pili suited for twitching motility.

Results

Suppressor mutations restore motility in Δ*fimX* mutants

FimX is a regulator of pilus extension in *P. aeruginosa*, though its exact mechanism is

not well understood (29–31). Consistent with previous work, we confirmed that a PAO1 deletion

mutant of *fimX* had significantly reduced twitching motility, 5-10% of the wild type (Fig 1B).

However, this strain still twitched when compared to a non-motile major pilin (*pilA*::Tn5) mutant

and remained susceptible to the pilus-targeting bacteriophage, PO4 (Fig 1C). FimX mutants did

not produce detectable surface pili in sheared protein fractions (Fig S1), further supporting a

defect in pilus extension. This phenotype is distinct from that of mutants lacking another PilB

regulator, the putative chaperone, PilZ (27, 28), which do not twitch and are resistant to pilus-

specific phages (Fig 1B and C).

We reasoned that we could leverage the Δ*fimX* pilus extension deficit to screen for

suppressor mutants capable of restoring twitching motility, to inform on the function of FimX.

After allowing *fimX* mutants to twitch at the sub-agar interface for an extended period (48-72 h),

we isolated highly motile flares that emerged from the small primary twitching zone (Fig 1D).

These cells were collected, isolated for single colonies, and after confirming their increased

motility compared to the Δ*fimX* parent, were sequenced to identify candidate suppressor

mutations. The mutations mapped to relevant proteins including the ATPase PilB, alignment

subcomplex component PilO (35, 36), members of the Pil-Chp network (PilJ and FimL) (7), and

CpdA (additional mutants are listed in Table S1). Each mutation increased motility to varying

degrees (Fig 1E), though not to wild-type levels, but did not impact growth (Fig S2) or phage

susceptibility (Fig S3). Complementing the suppressor mutants with *fimX* further increased

twitching motility to levels similar to or greater than WT, except for PilO P168L, suggesting it

has a *fimX* independent motility phenotype (Fig S4A). In contrast, complementation of the

suppressors with a defective version of FimX – where its cdGMP-binding EVL motif was

mutated to AAA – did not further increase twitching motility, consistent with previous reports

that this non-functional variant is incapable of PilB binding (30). Diminished twitching motility

in the WT strain when FimX AAA was expressed in trans is likely due to formation of FimX-

FimX AAA heterodimers, supported by their pairwise interactions in a bacterial 2-hybrid assay

(Fig S4B).

Δ*fimX* suppressor mutations do not directly influence cdGMP metabolism

Previous efforts to isolate Δ*fimX* suppressors revealed several mutations which collectively upregulated intracellular cdGMP levels (31). To test whether our mutants shared this phenotype, we assessed intracellular levels of cdGMP using a *pcdrA*:Lux cassette-based reporter system in liquid cultures (20). Under these conditions, none of our mutants demonstrated higher cdGMP levels over time when compared to control cells expressing the well-characterized diguanylate cyclase (DGC), SadC (Fig 2A). To directly test the impact of cdGMP on Δ*fimX* suppression, we overexpressed DGCs from *P. aeruginosa* (*sadC*) or *Escherichia coli* (*ydeH*) in WT or Δ*fimX* backgrounds and tested twitching motility. In all cases, cells expressing the DGCs showed significantly reduced twitching motility compared to the vector control (Fig 2B). Together, these data suggest that suppression of the Δ*fimX* motility defect in PAO1 is not related to changes in cdGMP levels.

Elevated cAMP levels increase motility in the Δ*fimX* background

Since several of our motile suppressor mutations mapped to the Pil-Chp network or CpdA, we hypothesized that changes to cAMP homeostasis (5, 7) increased twitching in the absence of *fimX*. We therefore assessed intracellular levels of cAMP in both liquid cultures and surface-adapted cellular populations using a *PaQa*:YFP reporter plasmid system (37). WT cells had significantly higher levels of cAMP when the cells were grown on a surface versus in liquid, but elevated cAMP-dependent signal was lost in a *pilA* mutant (Fig 3A), consistent with T4P being a major input to cAMP generation in *P. aeruginosa* (38). Deletion of CyaB or expression of a constitutively active CyaB R456L *in trans* further reduced or increased cAMP levels,

respectively (15, 39, 40). In these assay conditions, ΔfimX had lower than WT levels of cAMP

and was unable to respond to surface contact. The ΔfimX suppressor mutants PilJ T570I, FimL

L115F, and CpdA V258G all had significantly elevated cAMP levels. In contrast, neither the

PilB nor PilO P168L suppressor mutants had significant increases in cAMP levels compared to

the ΔfimX parent. These data suggested that there were at least two routes to increase twitching

in the *fimX* background, one of which was independent of cAMP levels.

We next asked whether high cAMP levels were responsible for increased ΔfimX

twitching. Expressing CyaB from a multicopy plasmid in the ΔfimX background was sufficient to

significantly increase twitching motility compared to the empty vector control (Fig 3B), while

expression of an inactive CyaB variant (Fig S5A) (41) was not. This increased twitching

phenotype was also observed in a double mutant lacking the Pil-Chp response regulators PilG

and PilH (Fig S5B), bypassing Pil-Chp-dependent CyaB activation (34). Together, these data

indicate that elevated cAMP levels due to Pil-Chp or CpdA mutations increase ΔfimX motility.

Strains with high intracellular cAMP in turn had increased Vfr-dependent expression (13) of an

mRuby3 fluorescent protein reporter integrated upstream of the T4P *pilMNOPQ* operon (Fig

3C). Integration of the reporter did not impact phage susceptibility of these strains (Fig S6)

indicating that pilus function is maintained. Consistent with this increased expression, levels of

PilO were also elevated (Fig 3D) suggesting that more T4P machines are expressed in these

backgrounds.

Notably, the PilB and PilO suppressors had no increase in Vfr-dependent transcription or

abundance of T4P components (Fig 2C and D) compared to ΔfimX, consistent with low cAMP

levels (Fig 3A). Another cAMP-dependent trait, type II secretion system-dependent secretion of

proteases generating zones of clearing on skim-milk plates (42) was also unchanged (Fig S7A

and B), collectively suggesting that the cAMP levels in those mutants were not significantly

impacted (13, 15), and that ΔfimX suppression occurs through non-cAMP related mechanisms in

these backgrounds.

Determinants for *pilB* Δ1bp function

The suppressors in the non-cAMP related class predominantly mapped to the pilus

extension ATPase, PilB. To probe this second set, we focused first on a mutant in which a single

adenine was deleted in the last codon of the PilB open reading frame (*pilB* Δ1bp). This deletion

produced a frameshift that extended the coding sequence by an additional 21 bases, or seven

amino acids, to the next stop codon (Fig 4A). When modelled in Alphfold3, the extended

segments were predicted to be solvent-exposed, protruding from the base of the hexamer and

away from its putative interaction interfaces with additional T4P components (Fig S8). To find

the minimum length of this C-terminal extension that was sufficient to increase motility in

ΔfimX, we generated single codon truncations from the 3′ end of the *pilB* Δ1bp sequence (Fig

4A). Each variant complemented motility in both a *pilB*::Tn5 background (Fig S9A), and in the

WT (Fig S9B), confirming that they produced a functional gene product with no dominant

negative effect through mixed oligomerization with WT PilB. In a strain lacking both *pilB* and

fimX, all *pilB* Δ1bp truncations tested significantly increased twitching motility relative to the

control complemented with WT PilB (Fig 3B). Notably, longer extensions rescued motility to a

greater degree than shorter ones.

Interestingly, the shortest *pilB* Δ1bp-21 truncation was identical in length to WT PilB, yet

only the *pilB* Δ1bp-21 sequence significantly increased twitching motility in the double mutant

(Fig 4B). Only the C-terminal amino acid differs between the two, with a valine substituted for

an aspartate in the *pilB* Δ1bp-21 sequence. PilB D566 may therefore play an important role in

modulating the function of the enzyme. The longer *pilB* Δ1bp sequences that support increased

motility have additional positively charged residues (Fig 4A), suggesting that loss of negative

charge at the C-terminus enhances PilB function. Since the above studies were done with

multicopy plasmids, we next introduced single chromosomal point mutants at the PilB C-

terminus in a Δ*fimX* background to assess recovery of twitching motility. A PilB D566V

mutation significantly increased twitching motility compared to the Δ*fimX* parent (Fig 4C). We

also observed a significant increase in twitching when the C-terminal amino acid was charge-

swapped to a lysine, but not when we introduced glutamate (Fig 4C). The second-last amino acid

in the native PilB sequence is a positively charged lysine, and we hypothesized that shortening

the WT sequence by a single amino acid to expose the positive charge would also restore

motility in a Δ*fimX* mutant. However, substituting D566 with a stop codon did not significantly

increase motility relative to the control (Fig 4C). These data suggest that PilB C-terminal charge,

in addition to C-terminal length, is an important regulator of PilB function.

Determinants for PilB T430P function

We next examined a second PilB suppressor (PilB T430P) which mapped to a site distal

to the C-terminus (Fig S8). A multiple protein sequence alignment of full-length PilB

homologues from various species revealed that the region surrounding T430 was well conserved,

although T430 itself was not (Fig 4D). We therefore reasoned that this amino acid was less

important to overall PilB function or potentially a species-specific regulatory site. We made

individual substitutions at T430, choosing amino acids found at this position in the other

homologues. All mutant constructs complemented twitching motility in a *pilB*::Tn5 background

and the WT (Fig S10). In a *pilB fimX* double mutant, only the original T430P suppressor

supported twitching motility (Fig 4E). Therefore, disruption of local PilB secondary structure,

rather than specific side-chain chemistry, is likely responsible for increased motility.

An Alphafold3 model of a PilB monomer predicted that residue T430 is located in the

second turn of a short α-helix (Fig 4D and S8). While T430 is not well conserved, other amino

acids in this helical segment, such as an adjacent R431, are. Alignment of the model onto the

crystal structure of hexameric PilB from *Geobacter metallireducens* bound to a non-hydrolysable

ATP analogue shows that the equivalent N429 (T430) and R430 (R431) residues are positioned

at the protomer-protomer interface or jutting into the lumen of the torus, depending on the phase

of the ATP hydrolysis cycle (Fig S8). We hypothesized that this secondary structural element

could be involved in inter-subunit communication and that the introduction of a helix-breaking

proline would alter these functional contacts. Supporting this idea, mutation of the adjacent

residues to proline, L429P or R431P, significantly increased twitching motility in *fimX pilB* (Fig

4F). In contrast, R431A failed to increase motility in the double mutant, thus disruption of the α-

helical segment rather than loss of this sidechain chemistry was likely responsible for altering

PilB function. Interestingly, in the *pilB*::Tn5 or WT backgrounds where FimX was present (Fig

S11A and B), twitching motility of PilB R431A and R431P variants was significantly decreased

compared to control (Fig S11A). Therefore, the presence or absence of FimX modulates the

function of these PilB variants.

Increased twitching of suppressors is due to enhanced PilB ATPase activity

We next investigated how the *pilB* Δ1bp and PilB T430P mutations restored pilus

function in a Δ*fimX* background. Other PilB regulatory effectors, such as PilZ, are thought to

serve as molecular chaperones and *pilZ* mutants lack detectable intracellular PilB on Western

blots (Fig 5A). This is consistent with results from previous studies of the *Xanthomonas citri*

PilZ homologue (27). The *pilZ* mutants are also non-motile and phage resistant (Fig 1B and C),

in keeping with their negligible levels of PilB. In contrast, PilB was still detectable in Δ*fimX* or

FimX AAA backgrounds, although its levels were approximately half of WT (Fig 5A). Neither

suppressor increased levels of PilB compared to those in Δ*fimX* or FimX AAA. Since FimX is

important for PilB polar localization (8, 27, 30), the reduction in PilB levels in *fimX* mutants may

be a consequence of fewer interactions between the ATPase and T4P machinery, making it more

susceptible to degradation.

Since PilB is an ATPase used to power pilus extension, we reasoned that the mutations

might increase enzymatic activity. PilB has two N-terminal domains (N1D and N2D) connected

by a flexible linker to the C-terminal domain (Figure S8) (43). The N1D can be removed without

disrupting hexamer formation to improve stability and solubility for *in vitro* studies (27, 44–48).

We purified ΔN1D-PilB variants to assess ATP hydrolysis efficiency *in vitro* (Fig S12). Both

PilB suppressors hydrolyzed significantly more ATP than the WT and catalytically inactive

Walker A mutant (PilB K332A) (1) controls over time (Fig 5B), as well as following overnight

incubation (Fig 5C). Despite mapping to disparate sites in the protein, both PilB suppressors

upregulated enzymatic activity and likely compensate for loss of FimX through this shared

mechanism. Interestingly, when we added FimX and cdGMP to the reaction, we saw a decrease

in ATP hydrolysis for *pilB* Δ1bp but an increase for PilB T430P (Fig 5D). While this result is

difficult to interpret, it suggests that FimX can modulate PilB enzymatic activity. In a genetic

context, *pilB* Δ1bp and PilB T430P mutations integrated into the chromosome restored twitching

motility to at least WT levels in the presence of FimX (Fig 5E), but not FimX AAA. Taken

together, this indicates that these PilB suppressor mutants can continue to respond to cdGMP-

bound FimX input and regulation.

FimX promotes pilus extension to produce longer pili

287   To test the impact of increased ATP hydrolysis by PilB on pilus assembly, we imaged

pilus dynamics using fluorescence microscopy (Fig 6A). The major pilin subunit, PilA, was first

labelled with a fluorescent maleimide dye at A86C as previously described (4, 49). All PilA

A86C mutants remained motile with slightly reduced twitching compared to the wild type (Fig

S13). WT cells produced T4P at the cell poles as expected, with some cells producing pili at both

poles (Fig 6A). Notably, the Δ*fimX* mutant produced very short pili that were difficult to detect,

often presenting only as bright polar puncta which we scored as a polar accumulation of labelled

PilA or very short T4P, below the limit of detection. In addition, fewer cells were piliated

compared to WT (Fig 6B and C). We chose to quantify pili using the Δ*fimX pilB* Δ1bp mutant as

this strain showed the most robust twitching phenotype of our suppressors. Furthermore, in the

presence of FimX, the *pilB* Δ1bp mutation alone had greater than WT levels of twitching,

providing an optimal background to study FimX's functional contribution. In the Δ*fimX pilB*

Δ1bp background, piliation was restored on many cells (Fig 6B and C), with longer pili

compared to the Δ*fimX* mutant (Fig 6A). The percentage of piliated cells in the Δ*fimX pilB* Δ1bp

mutant was similar to WT (Fig 6B and C), but the average maximum pilus length was

significantly shorter (Fig 6D), likely explaining its comparatively reduced motility (Fig 5E).

Maximum pilus length was significantly increased compared to the Δ*fimX* parent, indicating that

the *pilB* Δ1bp mutation caused cells to produce longer pili, presumably through enhanced ATP

hydrolysis. The *pilB* Δ1bp alone background produced the longest pili (Fig 6A and D) and had

the highest percentage of piliated cells (Fig 6B and C), consistent with this strain being the most

motile (Fig 5E). Longer and long-lived pili in PAO1 (~2-3 μm) comprise a rare subpopulation

(4). We therefore anticipate that more Δ*fimX* cells are likely piliated, but since the maximum

pilus length in these mutants is ~0.75 μm, most pili may not be detectable.

The *pilB* Δ1bp mutants afforded an opportunity to quantify pilus dynamics in the

presence or absence of FimX. We quantified the rates of pilus extension and retraction over time

in the Δ*fimX pilB* Δ1bp versus *pilB* Δ1bp to assess FimX's contribution to pilus dynamics. The

FimX-expressing strain had a significantly increased average rate of pilus extension (Fig 6E)

while the average rate of retraction was not significantly different, supporting FimX's role as a

modulator of PilB-dependent pilus extension. These data suggest that FimX directly increases

pilus extension efficiency, likely by acting on PilB.

In *P. aeruginosa* and other bacterial species, fluorescently labelled FimX fusions display

defects in polar localization when PilB interaction is impaired (27, 30, 33). Consistent with these

other studies, most Δ*fimX* mutant cells complemented with an N-terminal fusion of mNeonGreen

(mNGr) to FimX showed bright puncta localized to a single cell pole (Fig 7A), with a smaller

population of cells displaying bipolarly-distributed puncta (Fig 7B). The WT FimX fusion also

complemented twitching motility (Fig S14A), while cells expressing an empty vector control had

no discernable puncta (Fig S14B). The number of cells with identifiable polar puncta was

decreased by ~20-50% when *pilB* was deleted or if a mNGr-FimX AAA fusion was used (Fig

7B). Importantly, these data indicate that FimX localization patterns in our strains are consistent

with previous reports (8, 27, 30) and are contingent on interaction with PilB.

PilB enhancing mutations are uncommon

Given the relative ease of selection for *fimX* suppressor mutations that mapped to PilB

and improved twitching motility, we wondered if similar PilB mutations were common in

sequenced strains of *P. aeruginosa*. We compared 417 PilB orthologue sequences from the

Pseudomonas.com database to identify comparable mutations. Of these, 297 were the same

length as PAO1 PilB at 566 residues, and 82 were longer. Of those sequences, 72 were longer by

one residue, 8 by two residues, 1 by four residues, and 1 by 15 residues (Fig S15A). Strikingly,

despite these differences in length, the C-terminal residue of all sequences was an aspartic acid.

We also assessed the local sequence surrounding PilB T430 for helix-disrupting residues which

may confer a similar effect to T430P. We examined the region +/- three amino acids from the

residue aligned with PAO1 PilB T430 in a full-length protein sequence alignment, to capture the

most likely positions to influence stability of a 3-turn α-helix. Of the 417 sequences, 4 had a

proline residue and 8 had a glycine within this window, accounting for less than 1% and 2% of

the total sequences respectively (Fig S15B). This analysis of *P. aeruginosa* sequences suggests

that PilB mutations similar to the ones identified in this screen are uncommon.

Discussion

The reason that optimal PilB function in *P. aeruginosa* needs the regulatory input of

FimX remains poorly understood (30, 31). To address this knowledge gap, we screened for

suppressors capable of overcoming the Δ*fimX* motility deficit. We identified several suppressor

mutants, most of which mapped to the Pil-Chp network or to PilB (Table S1). A previous Tn-Seq

screen used to isolate suppressors in *P. aeruginosa* strain PA103 implicated FimX as a sensor of

the secondary messenger cdGMP. In that strain, Δ*fimX* suppressors with high levels of cdGMP

had increased motility, leading the authors to posit that FimX senses internal cdGMP to tune

pilus assembly efficiency over broad concentrations of the molecule (31). Unexpectedly, we

found no suppressors in cdGMP homeostatic machinery, nor did any of our mutations increase

cdGMP synthesis (Fig 2A). Further, exogenous expression of DGCs also failed to alleviate the

Δ*fimX* motility defect in PAO1 (Fig 2B), unlike in PA103 Δ*fimX* where expression of the

unrelated DGC PleD from *Caulobacter crescentus* was sufficient to restore motility and surface

piliation. While these phenotypic discrepancies may be strain-specific, overexpression of other

DGCs in PAO1 was reported to suppress twitching motility, though surface piliation was not

significantly impacted (31). Why these strains behave differently in response to cdGMP

production is not yet clear.

Instead, we found that increases in the secondary messenger cAMP increased twitching

in Δ*fimX*, though this is likely an indirect mechanism. Increasing the levels of cAMP drives

transcription of the Vfr regulon, which includes T4P components. Mutations that promote

accumulation of cAMP lead to the expression of additional pilus machines. Given that Δ*fimX*

mutants retain the ability to twitch, albeit poorly, these high cAMP mutants likely produce an

abundance of sub-optimally functioning T4P machines which cumulatively improve the cells'

ability to twitch (Fig 8). A similar pattern of compensation via elevated cAMP production was

recently reported for suppressors of another modulator of pilus function in strain PA14 (50).

We also isolated a suppressor that mapped to PilO (PilO P168L), a member of the T4P

alignment subcomplex (35). Notably, this mutant was the only one tested that was not

complemented to at least WT twitching by FimX expression (Fig S4). Therefore, the twitching

recovery mechanism for this mutant is likely independent of the FimX-PilB extension pathway.

PilO interacts with regulatory components such as SadC (18) and may connect pilus function to

environmental signals.

The suppressor mutations in PilB provided a useful tool to study FimX-dependent

physiology. Despite mapping to separate sites (Fig S8), both mutations increased ATP hydrolysis

(Fig 5A and B) suggesting that one of FimX's roles may be to enhance PilB enzymatic activity.

For the *pilB* Δ1bp mutant, mechanistic clues come from structural work on other extension

ATPases. The C-terminal solvent-exposed face of the *G. metallireducens* PilB hexamer has

dispersed negative surface charge (25, 26). The extended Δ1bp C-termini, which include several

positively charged residues, are predicted to protrude into this environment (Fig S8). Increasing

positive charge, or removing the native C-terminal negative charge, could promote electrostatic

stabilization of the hexamer, or facilitate the PilB conformational changes necessary for substrate

hydrolysis.

Analysis of the PilB T430P mutant uncovered a novel regulatory region in the PilB

multimer. Disruption of the short helix containing T430 restored twitching motility in a Δ*fimX*

background. The adjacent R431 residue is highly conserved in PilB homologues, and an R431P

variant increased twitching in Δ*fimX* but interestingly, significantly attenuated motility in the

presence of FimX (Fig S11). These PilB mutants therefore continue to interact with and respond

to FimX, though resulting in different functional outcomes. This secondary structural element

may represent an intramolecular regulatory site involved in FimX recognition or perhaps is

involved in communicating conformational changes induced through FimX interaction. More

detailed biochemical studies are still required to unravel these possibilities.

In the absence of FimX, twitching is diminished. Using fluorescence microscopy, we

showed that the pili of *fimX* mutants are significantly shorter than WT (Fig 6D). This result is

consistent with low levels of surface pili in sheared protein fractions (Fig S1). The Δ*fimX pilB*

Δ1bp suppressor mutant had increased PilB ATP hydrolysis and produced longer pili,

presumably through increased PilA polymerization. These data suggest that twitching motility in

PAO1 is maximized when cells make longer pili, as WT and *pilB* Δ1bp strains generated the

longest pili and largest twitching zones. When comparing Δ*fimX pilB* Δ1bp and *pilB* Δ1bp, we

saw a significant increase in the rate of pilus extension, but not retraction (Fig 6E). Our data

therefore support a model where FimX-PilB interaction stimulates pilus extension, potentially by

stabilizing or promoting the PilB conformational changes necessary to drive pilus assembly

within the stator (Fig 8). In *X. citri*, the addition of FimX stimulated ATP hydrolysis by a PilB-

PilZ complex *in vitro* (27). We also showed that FimX influenced *P. aeruginosa* ΔN1D-PilB

function *in vitro* (Fig 5D), implying that the two proteins interact. In other bacteria, FimX is

thought to engage the extension machinery through PilZ (27, 33, 51, 52). Similar FimX-PilZ

interactions have not been detected in *P. aeruginosa* (53), however, leaving definition of the

specific interaction interfaces for future studies.

In *P. aeruginosa* and *Xanthomonas spp.*, FimX has also been proposed to function as a

PilB polar localization factor (8, 27, 30), rather than as a chaperone like PilZ. In *P. aeruginosa*,

FimX shows polar localization defects in *pilB* backgrounds or when it is unable to bind to

cdGMP (Fig 7), suggesting that both cdGMP-bound FimX and PilB are required for polar co-

localization (30). The mechanism behind polar trafficking is poorly understood, but recent

evidence suggested that the Pil-Chp response regulator PilG was essential for proper PilB polar

localization (8, 34). Consistent with these observations, a recent study in *Acinetobacter baylyi*

demonstrated that its FimX homologue controls T4P localization through interactions with

components of the Pil-Chp pathway, but that this localization is independent of cdGMP binding

(32). How those factors are targeted to the cell poles in *P. aeruginosa* remains an open question.

PilB and FimX may first need to form a complex which is then recognized by additional factors

such as PilG. Alternatively, FimX interaction may bias PilB polar occupancy or keep it

sequestered, ensuring that the leading pole is primed for multiple rounds of pilus assembly (Fig

8). In this scenario, PilB and FimX polar retention is mutually contingent, but FimX instead

functions to reduce disengagement of PilB from active T4P machinery.

In other bacteria such as *Vibrio* or *Neisseria* species, pili are used predominantly for

surface adhesion (54, 55), cell-cell contact (56), DNA uptake (57, 58), or biofilm formation (55).

These organisms lack obvious FimX homologues. Instead, the N-terminal domain of extension

ATPases in these species binds directly to cdGMP, and nucleotide binding is essential for

function (54, 59–61). Pilus extension is therefore optimal at high intracellular cdGMP

concentrations, but this condition suppresses twitching motility. In *P. aeruginosa,* nucleotide

binding to FimX, which then dictates PilB localization and function, may serve as a functional

alternative. As a high affinity sensor of cdGMP (29, 33), FimX may drive optimal pilus

extension rates to enable multiple dynamic cycles of pilus assembly, and therefore facilitate

twitching, in response to nucleotide availability. This organization might also explain why

mutations which enhance PilB function are rare in *P. aeruginosa*, as these may dysregulate pilus

functional responses to signaling cues.

Materials and Methods

Bacterial strains and growth conditions

All strains and plasmids used in this study are outlined in Table S2. Bacterial strains were grown aerobically at 37 °C in Lysogeny Broth (LB) media (10 g/L tryptone, 5 g/L yeast extract, and 5 g/L NaCl), supplemented with the appropriate concentration of antibiotics where necessary unless otherwise specified. Antibiotics were used at the following concentrations: 15 µg/mL or 30 µg/mL of gentamicin for *E. coli* or *P. aeruginosa* respectively unless otherwise indicated. 50 µg/mL or 200 µg/mL kanamycin for *E. coli* or *P. aeruginosa* respectively. 100 µg/mL of ampicillin or 200 µg/mL of carbenicillin for *E. coli* or *P. aeruginosa* respectively. Plasmid constructs were generated using *E. coli* DH5α before introduction into the appropriate cellular background by heat shock (*E. coli*) or electroporation (*P. aeruginosa* or *E. coli*) where appropriate.

Plasmid and strain construction

Fragments of interest were PCR amplified from isolated mPAO1 genomic DNA. All PCR primers are listed in Table S3. For point mutations, two separate PCR amplified DNA fragments were stitched together through a second single overlap extension PCR, with each DNA overhang including the mutation of interest introduced during primer synthesis. For plasmid construction, the in-frame genetic insert and corresponding vector were digested with the appropriate FastDigest restriction enzymes (ThermoFisher Scientific) as per manufacturer instruction and ligated with T4 DNA ligase. Ligated products were transformed via heat shock into *E. coli* DH5α and selected on 1.5% agar LB media plates supplemented with the appropriate

antibiotic. Following selection, single colonies were harvested, and the plasmid was mini

prepped using a GeneJet plasmid prep kit (ThermoFisher Scientific) for sequence confirmation

(Plasmidsaurus).

Genetic manipulation of mPAO1 chromosomal DNA was performed by double

homologous recombination as previously described (62). In brief, for genetic deletions, ~500 bp

upstream and downstream of the desired target sequence were PCR amplified and purified from

isolated PAO1 genomic DNA. Upstream and downstream sequences were engineered to contain

~50-100 bases of the 5′ or 3′ ends of the gene deletion target to ensure minimal disruption of

adjacent genes. Upstream and downstream fragments were digested with the appropriate

restriction enzymes along with the suicide vector backbone pEX18Gm. Digested fragments were

ligated using T4 ligase and transformed into via heat shock into *E. coli* DH5α and selected for on

1.5% agar LB media plates supplemented with gentamicin. Purified and sequenced plasmids

were then transformed into *E. coli* SM10 (λ*pir*). A single colony of SM10 harbouring the

plasmid was grown overnight in 3 mL of LB media supplemented with gentamicin. The

following day, the SM10 culture was mixed with the appropriate strain of mPAO1 overnight

culture in a 1:1 ratio (by volume), pelleted by centrifugation, the resulting cell pellet was

resuspended in 50 µL of LB and then spotted on 1.5% agar LB plates dried in a biological safety

cabinet for 20-30 min. The spot was allowed to dry at room temperature before being incubated

at 37 °C overnight. The following day, the cell spot was scraped into fresh LB media and plated

onto premixed *Pseudomonas* isolation agar (Difco) supplemented with 100 µg/mL of

gentamicin. The plates were incubated again at 37 °C overnight. Single colonies were then re-

streaked again on *Pseudomonas* isolation agar plates supplemented with 100 µg/mL of

gentamicin and incubated overnight at 37 °C. The next day, cells were streaked in a zig-zag

pattern on 10% LB (1 g/L tryptone, 0.5 g/L yeast extract) with no added NaCl, 1.5% agar, and

5% sucrose media plates and incubated overnight at 37 °C. Single colonies from the sucrose

media selection plate were patched again onto a fresh sucrose media plate as well as an LB

media plate supplemented with gentamicin. Patches which grew on sucrose media but not LB

with gentamicin were assessed for chromosomal deletion via colony PCR.

For chromosomal knock-in of a point mutation, the same protocol as outlined above was

performed only differing at initial construct design. Knock-in constructs were instead generated

~500 bp upstream or downstream of the mutation site through single overlap extension PCR.

Overlap extension primers were also designed to silently introduce or remove a restriction cut

site in the genome where applicable to facilitate downstream quality assurance. The remaining

procedure was then carried out as described.

Twitching motility assays

Bacterial strains were streaked onto 1.5% agar LB media plates supplemented with

gentamicin where applicable and grown overnight at 37 °C. The following day, cells were picked

using a sterile pipette tip and stabbed down to the bottom layer plastic of 1% agar LB media

tissue culture treated plates, supplemented with antibiotic or arabinose where applicable, dried

for 30 min in a biological safety cabinet. Plates were incubated overnight at 37 °C. The

following day, the agar was carefully removed with a pipette tip and discarded. The twitching

zones adhered to the plate were stained with 0.1% crystal violet dye for a minimum of 10 min,

washed twice with DI water, and left to air dry overnight at room temperature before imaging

with a scanner. Twitching zone areas were then digitally traced using ImageJ, and either

duplicates or triplicates were averaged.

Phage-susceptibility assays

Bacterial strains were inoculated into 3 mL of LB media liquid cultures and grown

overnight at 37 °C. The next morning, 5 µL of overnight culture was sub-cultured into 3 mL of

fresh LB media and grown at 37 ºC to mid-logarithmic phase ($OD_{600} \sim 0.5$-1.0). Cells were

normalized to an OD_{600} of 0.1 in LB. For plate-based assays, 10 µL of bacterial suspension was

mixed with 5 µL of serially diluted PO4 phage stock (3e5 PFU/mL original titer). 10 µL of cell-

serial phage dilution was spotted onto 1.5% agar LB media dried for 30 min in a biological

safety cabinet. The plate was incubated overnight at room temperature. The following day, the

plate was incubated again at 37 °C for 2-3 hours before imaging with a scanner.

For liquid-based assays, 2 µL of OD_{600} normalized cell suspension was added to 200 µL

of LB media total in a flat bottom 96 well plate. To each well, 1 µL of 3e5 PFU/mL PO4 phage

stock was also added. The plate lid was sealed with masking tape and then incubated overnight

in a 37 °C shaking plate reader, with OD_{600} readings taken every 30 min for 16-18 hours. Curves

were normalized to LB media blank wells in the same plate.

Sheared surface protein preparation

Surface pilin and flagellin protein levels were assessed as previously described (63).

Briefly, strains were streaked in grid-like pattern on a 1.5% agar LB media plate dried for 30

minutes in a biological safety cabinet. The plate was incubated at 37 ºC overnight. The next

morning, cells were gently scrapped off the surface using a combination of a glass coverslip and

sterile inoculation loop into 1.2 mL of sterile 1X PBS. Surface appendages were sheared by

vortexing the suspension for 30 s. The homogenized suspension was transferred to a 1.5 mL

Eppendorf tube and pelleted by centrifugation at maximum speed for 30 min. The supernatant

was decanted into a fresh tube and 0.1 volume of 5M NaCl and 30% (w/v) polyethylene glycol

(MW range 8k) was added. The tubes were then incubated on ice for 90 min, inverting the tubes

periodically. The precipitated protein was collected by centrifugation again for 30 min at

maximum speed. The supernatant was decanted, and the precipitated protein resuspended in 50

µL of 1X SDS loading dye (62.5 mM Tris-HCl pH 6.8, 0.5% (w/v) SDS, 5% (v/v) β-

mercaptoethanol, 10% (v/v) glycerol, 0.25% (w/v) bromophenol blue), and boiled for 10 min.

The tubes were centrifuged for another 5 min at maximum speed and separated on a 15% SDS-

polyacrylamide gel. Proteins were visualized with Coomassie blue staining.

Bacterial 2-hybrid assay

Single colonies of *E. coli* BTH101 co-transformed with both T18/T25 tagged target

proteins were inoculated into 3 mL LB media overnight cultures supplemented with kanamycin,

ampicillin, and 0.5 mM IPTG. The following morning, the cultures were normalized to an OD_{600}

of 0.1 in fresh LB and 4 µL was spotted onto MacConkey agar plates (40 g/L, Difco)

supplemented with kanamycin, ampicillin, 0.5 mM IPTG, and 1% maltose pre-dried for 40 min

in a biological safety cabinet. The spots were allowed to dry at room temperature before

incubation at 30 ºC in a standing incubator. The plates were imaged after 24 hours.

Selecting ΔfimX twitching suppressor mutants

Motility suppressors were isolated through the standard twitching assay described with

some alterations. ΔfimX bacteria were first streaked for single colonies on an LB agar plate.

Using a sterile pipette tip, a single colony was stab-inoculated into a standard twitching plate ~4-

5 times in a row. The tip was discarded, and a new colony was then stab inoculated in the same

manner. This was repeated for a total of 35-40 individual twitching zones on a single assay plate.

The plate was sealed with parafilm and incubated at 37 ºC for 48-72 hours. Following

incubation, the top agar layer was carefully removed and discarded. Flares emerging from small

twitching zones were collected with a sterile cotton swab and streaked again for single colonies.

Single mutant colonies were reassessed for twitching motility. Those with increased twitching

relative to the ΔfimX strain were collected and their genomic DNA isolated using a Wizard

genomic DNA isolation kit (Promega) as per manufacturer's instructions. Isolated chromosomal

DNA was sent for Illumina Sequencing (SeqCenter) and mutations were identified using BreSeq

(64). Where appropriate, candidate mutations were reintroduced into the chromosome as

described above.

cdGMP reporter assay

The cdGMP reporter assay was conducted as previously described (20) with some

alterations. Briefly, cells transformed with the reporter construct (*pcdrA*:LuX-PMS402) were

grown in LB media liquid cultures supplemented with kanamycin (as well as gentamicin for the

high control carrying *sadC*-pBADGr) overnight at 37 ºC. Overnight cultures were diluted (1:25)

in 3 mL of fresh LB media with antibiotics and grown at 37 ºC to mid-logarithmic phase.

Cultures were then normalized to an OD_{600} of 0.1 and 2 µL of suspension was added to 200 µL

of LB media total supplemented with antibiotics in a black flat clear-bottom 96 well plate. The

plate was sealed with masking tape and then incubated overnight in a 37 °C shaking plate reader,

with chemiluminescence and OD_{600} readings taken every 15 min for 16-18 hours. Relative

luminescence signal was normalized to OD_{600} reading at the same timepoint.

cAMP reporter assay

The cAMP reporter assay was conducted as previously described (37) with some

alterations. Briefly, cells transformed with the reporter construct (*rpoD*:mKate/*PaQa*:YFP –

O'Toole Lab) were grown in LB media liquid cultures supplemented with carbenicillin (as well

as gentamicin for the high cAMP control carrying CyaB R456L-pBADGr) overnight at 37 ºC.

Overnights were sub-cultured 1:25 in 5 mL of M8 minimal media containing 0.2% glucose,

0.5% casamino acids, and 1 mM $MgSO_4$ supplemented with carbenicillin to an OD_{600} of 0.2-0.4.

At this point, 200 µL of liquid culture was spread onto 1% agar M8 media plates dried for 20

min in a biological safety cabinet. Liquid cultures and plates were then incubated for an

additional 5 hours at 37 ºC. After the incubation, 1 mL of sterile PBS was added to each plate

and cells were scraped up with a sterile inoculation loop. 100 µL of cell suspension was taken

from each plate and normalized to an OD_{600} of ~0.2 in sterile 1X PBS. The liquid cultures were

also normalized to an OD_{600} of ~0.2 in sterile 1X PBS. 200 µL of normalized cell suspensions

were then added to a black flat clear bottom 96 well plate (Corning). mKate (Ex: 588/Em: 635)

and YFP (Ex: 500/Em: 541) fluorescence were measured, and the signal was normalized by

dividing YFP/mKate for both surface-adapted and liquid samples.

Vfr reporter assay

For *vfr* promoter activity detection, cells harbouring a gene encoding mRuby3

chromosomally integrated upstream of *pilM* and under control of the *pilMNOPQ* promoter were

streaked onto 1.5% agar LB media plates and incubated overnight at 37 ºC. The next morning,

cells were harvested into sterile 1X PBS and normalized to an OD_{600} of ~0.1. 200 µL of cell

suspension was then added to a black flat clear bottom 96 well plate. mRuby3 (Ex: 558/Em: 600)

fluorescence and exact OD_{600} were then measured, and the signal was normalized by dividing

mRuby3/OD_{600}.

Western blot analysis

Whole cell lysates were first separated on a 15% or 12% SDS-polyacrylamide gels for

PilA and PilB respectively. Samples were transferred to a nitrocellulose membrane at 225 mA

for 1 hour. The membranes were blocked with 5% (w/v) skim milk (Bioshop) dissolved in 1X

Tris-buffered saline (TBS: 20 mM Tris-HCl pH 7.6, 150 mM NaCl) overnight or for 2 hours for

PilA or PilB respectively. Primary incubation was performed as previously described using

rabbit antisera (65). The membranes were washed 4 times for 5 min with 1X TBS before

incubation using alkaline phosphatase-conjugated goat-anti rabbit IgG secondary antibody

(BioRad) in a 1:3500 dilution. The membranes were washed 4 times with 1X TBS and developed

using alkaline phosphatase developing reagent according to manufacturer's protocol, before

imaging the membranes in an Azure Biosystems 400 transilluminator.

Skim milk T2SS secretion assay

Liquid cultures were grown overnight in LB media at 37 ºC. The next morning, each culture was normalized to an OD_{600} of 0.2 and spotted onto tryptic soy agar plates supplemented with 1.5% skim milk powder (Bioshop) dried for 30 min in a biological safety cabinet. The spots were allowed to dry at room temperature before incubation overnight at 37 ºC. The resulting zones of clearance were then imaged with a scanner and the zone areas were quantified digitally using ImageJ.

Protein modelling and sequence alignments

Alphafold3 was used to predict protein structures and generate predicted aligned error plots using the unmodified default settings (66). All models and plots were visualized in UCSF ChimeraX (67). Full length PilB sequences were aligned using Geneious Prime and visualized using ESPript3 (68).

Protein purification

N-terminally truncated PilB (ΔN1D-PilB: M1-D180) and full length FimX were subcloned into pET28b with an N-terminal His_6-tag for inducible expression. PilB was purified as previously described with some alterations. Briefly, ΔN1D-PilB expressing overnight cultures of BL21-CodonPlus (DE3)-RIL were sub-cultured (1:100) in 1 L of LB supplemented with kanamycin and incubated at 37 ºC with shaking until mid-logarithmic phase (~0.4-0.8 OD_{600}). Cells were induced with 0.5 mM IPTG overnight at 18ºC with shaking. The next morning, cells

were harvested by centrifugation at 3000 xg for 20 min. Cell pellets were resuspended in lysis

buffer (50 mM HEPES pH 7.3, 250 mM NaCl, 10 mM imidazole) and incubated on ice with

20mg of lysozyme for 15 min with periodic inversion. Cells were then lysed using sonication and

the supernatant clarified through centrifugation at 31000 xg for 30 min. Supernatant was then

incubated with ~5 mL of Ni^{2+}-NTA resin (Millipore) pre-equilibrated with lysis-buffer for 1 hour

on ice with gentle shaking. The resin was washed with 3 separate 30 mM imidazole washes (15

mL) before elution with 250 mM imidazole. The elution fraction was dialyzed overnight at 4 ºC

into 7.3 mM HEPES pH 7.3, 200 mM NaCl and concentrated using an Amicon concentrator

(Millipore-Sigma). Protein was then aliquoted, flash frozen in liquid nitrogen, and stored at -

80ºC. FimX was expressed as described above and purified as previously described (30), except

for the final purification step involving separation by size-exclusion chromatography (Superdex

200, Cytiva) equilibrated using 50 mM Tris-HCl pH 8.0, 250 mM NaCl, and 5 mM β-ME.

Protein-containing fractions were determined using SDS-PAGE and then pooled for

concentration using an Amicon concentrator.

ATPase activity assay

All ATPase assays contained equimolar amount of ΔN1D-PilB or mutants (100 nM) to

ensure valid comparison. PilB reaction buffer was prepared containing 50 mM HEPES pH 7.3,

200 mM NaCl, 5 mM $MgCl_2$, 1 µM $ZnCl_2$, and 1 mM ATP. Where applicable, the reaction

buffer was supplemented with 1 µM cdGMP, and/or 200 nM of purified FimX. Aliquots of

ΔN1D-PilB was thawed and the tube was centrifuged at maximum speed for 10 min at 4 ºC to

pellet insoluble precipitate. The remaining supernatant was transferred to a fresh tube and kept

on ice. ATPase reactions were performed in a clear flat-bottom 96 well plate (Corning) in a total

volume of 100 µL at 37 ºC in a standing incubator. Following incubation and at the appropriate

time point, 50 µL of reaction was mixed with 100 µL of BioMol green reagent (Enzo

Lifesciences) incubated at room temperature for 15 min at which point absorbance at 620 nm

was read with a spectrophotometer. Each ΔN1D-PilB reaction series was normalized to the

timepoint at 0 min to ensure consistent background signal reduction. For overnight incubations,

the 96-well plate lid was sealed with masking tape, and the reaction was performed as described

with samples being normalized to an equivalent reaction well lacking PilB.

Fluorescence microscopy experiments

T4P labeling experiments were performed using a cysteine knock-in method described

previously, with some species-specific changes (4, 69). 100 µl of overnight cultures of *P.*

aeruginosa were spotted onto prewarmed LB agar plates and grown upright at 37 °C for 4-5

hours. Plate-grown cells were then scraped off of plates using a P200 pipette tip and gently

resuspended into 100 µl of fresh LB. AlexaFluor488-maleimide (Sigma) was added to the cell

suspension for a final concentration of 50 ng/µl and incubated for 45-60 min in the dark at room

temperature. Cells were washed once with 200 µl fresh LB using centrifugation settings of

18,000 x g for 1 min. Cell were resuspended into 200 µl of fresh LB. mNeonGreen-FimX strains

were prepared exactly the same way except with gentamycin supplemented in the medium and

with no AlexaFluor488-maleimide labeling steps. All imaging was performed under 1% agarose

pads made with PBS solution. Cell bodies were imaged using phase-contrast microscopy on a

Nikon Ti2-E microscope using a Plan Apo 100X oil immersion objective, a GFP/FITC/Cy2 filter

set for pili and mNeonGreen-FimX, a Hamamatsu ORCA-Fusion Gen-III cCMOS camera, and

Nikon NIS Elements Imaging Software. Measurements of pili were quantified manually using

Fiji. Notably, populations of *P. aeruginosa* containing labeled T4P exhibit variability in labeling

efficiency for unknown reasons. Therefore, T4P fluorescence signal contrast was adjusted so that

T4P were clearly visible for each strain.

Acknowledgements

We thank Dr. George O'Toole and Dr. Christopher Geiger for the cAMP reporter

plasmid and their help in optimizing the detection assay. We thank Dr. Lynne Howell and Ian

Yen for the *ydeH*-pBADGr plasmid. We also thank Dr. Sara Andres, Caitlin Doubleday, and the

Andres lab for access to equipment and assistance in purifying all proteins used in this study. We

thank the members of the Burrows lab for their helpful discussions and insights. This work was

supported by a Canadian Institutes of Health Research Project grant (PJT-156080) to LLB and a

National Institutes of Health grant (R35GM150916) to CKE. LLB holds a Tier 1 Canada

Research Chair in Microbe-Surface Interactions (CRC 2021–00103). CKE is a Damon Runyon-

Marilyn and Scott Urdang Breakthrough Scientist supported by the Damon Runyon Cancer

Research Foundation (DFS6023). NR holds an NSERC CGS-D award. NY was funded through

an NSERC USRA.

 Figure Captions

 Figure 1: Suppressor mutations overcome Δ*fimX* twitching motility defects. (A) The type IV
 pilus nanomachine comprising four distinct protein subcomplexes (motor, alignment, secretin
 subcomplexes, and pilus fiber) and associated inner-membrane Pil-Chp signaling network
 responsible for cAMP generation. PilB in association with regulatory effectors PilZ and FimX
 drives pilus extension. **(B)** Quantification of sub-agar stab twitching motility zones for extension
 regulatory effector mutants. Representative crystal violet-stained twitching zones are shown to
 the left. X-axis depicts a $\log_2$ scale. Bars represent the means of triplicate samples from three
 independent experiments ± SD. **(C)** Representative plaque assay demonstrating 10-fold serial
 dilutions of bacteriophage PO4 on extension regulatory effector mutants. **(D)** Cartoon
 representation of the Δ*fimX* suppressor mutant screen setup. A representative motile flare is
 shown to the right. Image border = 1 cm. **(E)** Quantification of sub-agar stab twitching motility
 zones for selected Δ*fimX* suppressor mutants. Representative twitching zones are shown to the
 left. Data are plotted as a fold change of the Δ*fimX* parent strain. Bars represent the means of
 triplicate samples from three independent experiments ± SD. All scale bars = 1 cm. *******: $p \leq$
 0.001 (Two-tailed parametric *t*-test).

 Figure 2: FimX suppressor mutations do not impact intracellular cdGMP levels. (A) Time-
 course curves of relative luminesence signal of a *pcdrA*:LuX cassette cdGMP reporter
 normalized to OD_{600}. Each timepoint is the mean of triplicate samples from three independent
 experiments ± SD. **EV**: empty pMS402 vector, ***sadC***: *sadC* in pBADGr vector. **(B)**
 Quantification of sub-agar stab twitching motility zones for strains overexpressing diguanylate
 cyclases (DGCs). Representative crystal violet-stained twitching zones are shown to the left.
 DGC expression was induced with 0.1% arabinose. Bars represent the means of triplicate
 samples from three independent experiments ± SD. X-axis depicts a $\log_2$ scale. Scale bar = 1 cm.
 EV: empty pBADGr vector, ***sadC***: *sadC* in pBADGr vector, ***ydeH***: *ydeH* in pBADGr vector.
 ***: $0.001 \geq p$ (Two-tailed parametric *t*-test).

 Figure 3: A subset of Δ*fimX* twitching suppressors elevate cAMP levels and upregulate
 associated phenotypes. (A) Plasmid-based cAMP reporter assay quantifying cAMP-dependent
 YFP expression (*PaQa*) normalized to constitutive mKate expression (*rpoD*). Bars represent the
 means of duplicate samples from three independent experiments ± SD. **(B)** Quantification of sub-
 agar stab twitching motility zones for strains overexpressing the adenylate cyclase CyaB or a
 catalytically inactive mutant. Representative crystal violet-stained twitching zones are shown to
 the left. Expression was induced with 0.1% arabinose. Bars represent the means of triplicate
 samples from three independent experiments ± SD. Scale bar = 1 cm. **EV**: empty pHERD30T
 vector, **His$_6$-CyaB (D234S)**: N-terminally hexa-histidine tagged CyaB or CyaB D234S in
 pHERD30T vector. **(C)** Vfr-dependent expression of mRuby3 chromosomally inserted upstream
 of the native alignment complex *pilMNOPQ* operon. Bars represent the means of duplicate
 samples from three independent experiments ± SD. Cartoon schematic of the mRuby3
 expression system is shown above the graph. **(D)** Representative PilO Western blot. Sample

loading was normalized to a non-specific band shown below. Molecular weight is indicated to
the left. Blot is representative of three independent experiments. **CyaB R456L**: CyaB R456L in
pBADGr vector. *****: $0.05 \geq p \geq 0.01$; ******: $0.01 \geq p \geq 0.001$; *******: $0.001 \geq p$ (Two-tailed parametric
t-test).

Figure 4: PilB mutations increase twitching motility in the absence of *fimX*. (A) Cartoon
schematic of deleted base pair A1697 at the PilB C-terminus, shown in red, and resulting
extended sequence. The *pilB* Δ1bp truncation sequences are shown below. **(B)** Quantification of
twitching zone areas of Δ*fimX pilB*::Tn5 double mutants overexpressing *pilB* Δ1bp 3′
truncations. Representative crystal violet stained twitching zones are shown to the left. PilB
expression was induced with 0.05% arabinose. Bars represent the means of duplicate samples
from three independent experiments ± SD. **(C)** Quantification of twitching zone areas of Δ*fimX*
PilB D566X point mutants chromosomal knock-ins. Representative twitching zones are shown to
the left. Bars represent the means of triplicate samples from three independent experiments ± SD.
(D) Full-length multiple sequence alignment of PilB homologues from various species
highlighting the sequence surrounding PAO1 PilB T430, labelled in red. The Alphafold3 model
of the predicted secondary structure in this region is shown to the left. **(E)** Quantification of
twitching zone areas of *fimX pilB* double mutants overexpressing PilB T430X mutants.
Representative twitching zones are shown to the left. PilB expression was induced with 0.05%
arabinose. Bars represent the means of duplicate samples from three independent experiments ±
SD. **(F)** Quantification of twitching zone areas of *fimX pilB*::Tn5 double mutants overexpressing
PilB constructs with point mutations in positions adjacent to T430. Representative twitching
zones are shown to the left. PilB expression was induced with 0.05% arabinose. Bars represent
the means of triplicate samples from three independent experiments ± SD. **EV**: empty
pHERD30T vector, ***pilB* Δ1bp-3 to 21**: *pilB* Δ1bp-3 to 21 in pHERD30T vector, **PilB**
L429/T430/R431X: indicated PilB substitution in pHERD30T vector. All scale bars = 1 cm. **ns**:
$p \geq 0.05$; *****: $0.05 \geq p \geq 0.01$; ******: $0.01 \geq p \geq 0.001$; *******: $0.001 \geq p$ (Two-tailed parametric *t*-test).

Figure 5: PilB suppressor mutations increase ATP hydrolysis. (A) Representative PilB
Western blot. Sample loading was normalized to a non-specific band shown below. Molecular
weight is indicated to the left. Blot is representative of three independent experiments.
Quantification using BioMol green reagent of free phosphate released over a time course **(B)** or
following overnight incubation **(C)** by ΔN1D-His$_6$-PilB-catalyzed ATP hydrolysis. PilB K332A,
a non-catalytically active Walker mutant, was used as a negative control. **(D)** Quantification of
free phosphate release by ΔN1D- His$_6$-PilB-catalyzed ATP hydrolysis in the presence of cdGMP
containing reaction buffer (+Buffer) or with cdGMP and FimX (+FimX) following overnight
incubation. **(E)** Quantification of twitching zone areas of Δ*fimX* or FimX AAA mutants with or
without individual PilB suppressor mutation chromosomal knock-ins. Representative crystal
violet stained twitching zones are shown to the left. Bars represent the means of triplicate
samples from three independent experiments ± SD. Scale bar = 1 cm. **ns**: $p \geq 0.05$; *****: $0.05 \geq p \geq$
0.01; ******: $0.01 \geq p \geq 0.001$; *******: $0.001 \geq p$ (Two-tailed parametric *t*-test).

Figure 6: FimX mutants produce short pili due to slow extension rates. (A) Representative images of PilA A86C type IV pili labelled with maleimide-Alexafluor 488 with background fluorescence subtracted. White arrows indicate visible pili. All scale bars = 2 μm. Number of cells producing visible PilA A86C type IV pili expressed as a percentage of total cells **(B)** or normalized relative to the WT arbitrarily set to 1 **(C)**. Bars represent the means ± SD of three independent experiments (WT *n* = 193, Δ*fimX n* = 181, Δ*fimX pilB* Δ1bp *n* = 210, *pilB* Δ1bp *n* = 197 total cells). Technical replicates are indicated by colour. **(D)** Maximum PilA A86C pilus length. Each sample depicts the means ± SD of three independent experiments, each point represents a single PilA A86C pilus length (WT, Δ*fimX pilB* Δ1bp, and *pilB* Δ1bp *n* = 60, Δ*fimX n* = 45 total pili). Technical replicates are indicated by colour. **(E)** Rate of PilA A86C pilus extension or retraction. Each sample depicts the means ± SD of three independent experiments, each point represents the rate of extension and/or retraction of a single PilA A86C pilus (Δ*fimX pilB* Δ1bp *n* = 46, *pilB* Δ1bp *n* = 44 total extension and Δ*fimX pilB* Δ1bp *n* = 40, *pilB* Δ1bp *n* = 41 total retraction events). **ns**: $p \geq 0.05$; *****: $0.05 \geq p \geq 0.01$; ******: $0.01 \geq p \geq 0.001$ (Two-tailed parametric *t*-test).

Figure 7: FimX polar localization is disrupted in the absence of PilB binding. (A) Representative images of cells expressing an mNeonGreen N-terminal FimX or FimX AAA fusion with background fluorescence subtracted. All scale bars = 2 μm. **(B)** Quantification of cells expressing an mNeonGreen-N-terminal FimX or FimX AAA fusion scored for bipolar, unipolar, or no discernable puncta and expressed as a percentage of total cells. Bars represent the means ± SD of three independent experiments (Δ*fimX* + mNGr-FimX *n*= 216, Δ*fimX pilB*::Tn5 + mNGr-FimX *n*= 213, Δ*fimX* + mNGr-FimX AAA *n*= 235, Δ*fimX pilB*::Tn5 + mNGr-FimX AAA *n*= 206, total cells). **mNGr-FimX** or **mNGr-FimX AAA**: mNeonGreen N-terminal FimX or FimX AAA fusion in pHERD30T vector. Expression was not induced with arabinose.

Figure 8: Model of Δ*fimX* suppression pathways. Suppression of the Δ*fimX* twitching motility deficit can occur through mutations which elevate intracellular levels of the secondary messenger molecule cAMP. More cAMP generation leads to increased Vfr-dependent expression of T4P components, which assemble into more machines, increasing twitching motility. This mechanism is not FimX-specific. Alternatively, mutations that increase PilB ATP hydrolysis lead to production of longer pili. This overcomes the short-pilus defect observed in *fimX* mutants. FimX in complex with PilB enhances the rate of PilB-dependent pilus extension to produce longer pili optimized for twitching motility. FimX polar localization to T4P machines is contingent on PilB binding, therefore FimX influences its polar trafficking and/or sequestration.

References

1. Chiang, P., Sampaleanu, L. M., Ayers, M., Pahuta, M., Howell, P. L., and Burrows, L. L. Y. 2008 Functional role of conserved residues in the characteristic secretion NTPase motifs of the *Pseudomonas aeruginosa* type IV pilus motor proteins PilB, PilT and PilU. *Microbiology*. **154**, 114–126

2. Burrows, L. L. (2012) *Pseudomonas aeruginosa* twitching motility: type IV pili in action. *Annu. Rev. Microbiol.* **66**, 493–520

3. Jakovljevic, V., Leonardy, S., Hoppert, M., and Søgaard-Andersen, L. (2008) PilB and PilT Are ATPases Acting Antagonistically in Type IV Pilus Function in *Myxococcus xanthus*. *J. Bacteriol.* **190**, 2411–2421

4. Koch, M. D., Fei, C., Wingreen, N. S., Shaevitz, J. W., and Gitai, Z. (2021) Competitive binding of independent extension and retraction motors explains the quantitative dynamics of type IV pili. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2014926118

5. Persat, A., Inclan, Y. F., Engel, J. N., Stone, H. A., and Gitai, Z. (2015) Type IV pili mechanochemically regulate virulence factors in *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci.* **112**, 7563–7568

6. Guo, S., Chang, Y., Brun, Y. V., Howell, P. L., Burrows, L. L., and Liu, J. (2024) PilY1 regulates the dynamic architecture of the type IV pilus machine in *Pseudomonas aeruginosa*. *Nat. Commun.* **15**, 9382

7. Inclan, Y. F., Persat, A., Greninger, A., Von Dollen, J., Johnson, J., Krogan, N., Gitai, Z., and Engel, J. N. (2016) A scaffold protein connects type IV pili with the Chp chemosensory system to mediate activation of virulence signaling in *Pseudomonas aeruginosa*. *Mol. Microbiol.* **101**, 590–605

8. Kühn, M. J., Talà, L., Inclan, Y. F., Patino, R., Pierrat, X., Vos, I., Al-Mayyah, Z., Macmillan, H., Negrete, J., Engel, J. N., and Persat, A. (2021) Mechanotaxis directs *Pseudomonas aeruginosa* twitching motility. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2101759118

9. Laventie, B.-J., Sangermani, M., Estermann, F., Manfredi, P., Planes, R., Hug, I., Jaeger, T., Meunier, E., Broz, P., and Jenal, U. (2019) A Surface-Induced Asymmetric Program Promotes Tissue Colonization by *Pseudomonas aeruginosa*. *Cell Host Microbe*. **25**, 140-152.e6

10. Comolli, J. C., Hauser, A. R., Waite, L., Whitchurch, C. B., Mattick, J. S., and Engel, J. N. (1999) *Pseudomonas aeruginosa* Gene Products PilT and PilU Are Required for Cytotoxicity In Vitro and Virulence in a Mouse Model of Acute Pneumonia. *Infect. Immun.* **67**, 3625–3630

11. O'Toole, G. A., and Kolter, R. (1998) Flagellar and twitching motility are necessary for *Pseudomonas aeruginosa* biofilm development. *Mol. Microbiol.* **30**, 295–304

12. Fulcher, N. B., Holliday, P. M., Klem, E., Cann, M. J., and Wolfgang, M. C. (2010) The *Pseudomonas aeruginosa* Chp Chemosensory System Regulates Intracellular cAMP Levels by Modulating Adenylate Cyclase Activity. *Mol. Microbiol.* **76**, 889–904

13. Wolfgang, M. C., Lee, V. T., Gilmore, M. E., and Lory, S. (2003) Coordinate regulation of bacterial virulence genes by a novel adenylate cyclase-dependent signaling pathway. *Dev. Cell.* **4**, 253–263

14. Kim, H.-S., Kim, S.-M., Lee, H.-J., Park, S.-J., and Lee, K.-H. (2009) Expression of the
cpdA gene, encoding a 3',5'-cyclic AMP (cAMP) phosphodiesterase, is positively
regulated by the cAMP-cAMP receptor protein complex. *J. Bacteriol.* **191**, 922–930

15. Fuchs, E. L., Brutinel, E. D., Klem, E. R., Fehr, A. R., Yahr, T. L., and Wolfgang, M. C.
(2010) In Vitro and In Vivo Characterization of the *Pseudomonas aeruginosa* Cyclic AMP
(cAMP) Phosphodiesterase CpdA, Required for cAMP Homeostasis and Virulence Factor
Regulation. *J. Bacteriol.* **192**, 2779–2790

16. Luo, Y., Zhao, K., Baker, A. E., Kuchma, S. L., Coggan, K. A., Wolfgang, M. C., Wong,
G. C. L., and O'Toole, G. A. (2015) A Hierarchical Cascade of Second Messengers
Regulates *Pseudomonas aeruginosa* Surface Behaviors. *mBio*. **6**, e02456-14

17. Jenal, U., Reinders, A., and Lori, C. (2017) Cyclic di-GMP: second messenger
extraordinaire. *Nat. Rev. Microbiol.* **15**, 271–284

18. Webster, S. S., Lee, C. K., Schmidt, W. C., Wong, G. C. L., and O'Toole, G. A. (2021)
Interaction between the type 4 pili machinery and a diguanylate cyclase fine-tune c-di-GMP
levels during early biofilm formation. *Proc. Natl. Acad. Sci.* **118**, e2105566118

19. Merritt, J. H., Ha, D.-G., Cowles, K. N., Lu, W., Morales, D. K., Rabinowitz, J., Gitai, Z.,
and O'Toole, G. A. (2010) Specific Control of *Pseudomonas aeruginosa* Surface-
Associated Behaviors by Two c-di-GMP Diguanylate Cyclases. *mBio*. **1**,
10.1128/mbio.00183-10

20. Marko, V. A., Kilmury, S. L. N., MacNeil, L. T., and Burrows, L. L. (2018) *Pseudomonas
aeruginosa* type IV minor pilins and PilY1 regulate virulence by modulating FimS-AlgR
activity. *PLOS Pathog.* **14**, e1007074

21. Webster, S. S., Mathelié-Guinlet, M., Verissimo, A. F., Schultz, D., Viljoen, A., Lee, C. K.,
Schmidt, W. C., Wong, G. C. L., Dufrêne, Y. F., and O'Toole, G. A. (2022) Force-Induced
Changes of PilY1 Drive Surface Sensing by *Pseudomonas aeruginosa*. *mBio*. **13**, e03754-
21

22. Harvey, H., Bondy-Denomy, J., Marquis, H., Sztanko, K. M., Davidson, A. R., and
Burrows, L. L. (2018) *Pseudomonas aeruginosa* defends against phages through type IV
pilus glycosylation. *Nat. Microbiol.* **3**, 47–52

23. Koch, M. D., Black, M. E., Han, E., Shaevitz, J. W., and Gitai, Z. (2022) *Pseudomonas
aeruginosa* distinguishes surfaces by stiffness using retraction of type IV pili. *Proc. Natl.
Acad. Sci.* **119**, e2119434119

24. Collins, R., Karuppiah, V., Siebert, C. A., Dajani, R., Thistlethwaite, A., and Derrick, J. P.
(2018) Structural cycle of the *Thermus thermophilus* PilF ATPase: the powering of type
IVa pilus assembly. *Sci. Rep.* **8**, 14022

25. McCallum, M., Tammam, S., Khan, A., Burrows, L. L., and Howell, P. L. (2017) The
molecular mechanism of the type IVa pilus motors. *Nat. Commun.* **8**, 15091

26. McCallum, M., Burrows, L. L., and Howell, P. L. (2019) The Dynamic Structures of the
Type IV Pilus. *Microbiol. Spectr.* 10.1128/microbiolspec.PSIB-0006-2018

27. Llontop, E. E., Cenens, W., Favaro, D. C., Sgro, G. G., Salinas, R. K., Guzzo, C. R., and
Farah, C. S. (2021) The PilB-PilZ-FimX regulatory complex of the Type IV pilus from
Xanthomonas citri. *PLOS Pathog.* **17**, e1009808

28. Alm, R. A., Bodero, A. J., Free, P. D., and Mattick, J. S. (1996) Identification of a novel
gene, *pilZ*, essential for type 4 fimbrial biogenesis in *Pseudomonas aeruginosa*. *J.
Bacteriol.* **178**, 46–53

29. Kazmierczak, B. I., Lebron, M. B., and Murray, T. S. (2006) Analysis of FimX, a
phosphodiesterase that governs twitching motility in *Pseudomonas aeruginosa*. *Mol.*
Microbiol. **60**, 1026–1043

30. Jain, R., Sliusarenko, O., and Kazmierczak, B. I. (2017) Interaction of the cyclic-di-GMP
binding protein FimX and the Type 4 pilus assembly ATPase promotes pilus assembly.
PLOS Pathog. **13**, e1006594

31. Jain, R., Behrens, A.-J., Kaever, V., and Kazmierczak, B. I. (2012) Type IV Pilus Assembly
in *Pseudomonas aeruginosa* over a Broad Range of Cyclic di-GMP Concentrations. *J.*
Bacteriol. **194**, 4285–4294

32. Ellison, T. J., Yen, I. Y., Howell, P. L., and Ellison, C. K. (2025) FimX regulates type IV
pilus localization via the Pil-Chp chemosensory system in *Acinetobacter baylyi*.
10.1101/2025.05.21.655318

33. Wei, C., Wang, S., Liu, P., Cheng, S.-T., Qian, G., Wang, S., Fu, Y., Qian, W., and Sun, W.
(2021) The PdeK-PdeR two-component system promotes unipolar localization of FimX and
pilus extension in *Xanthomonas oryzae* pv. *oryzicola*. *Sci. Signal.* **14**, eabi9589

34. Kühn, M. J., Macmillan, H., Talà, L., Inclan, Y., Patino, R., Pierrat, X., Al-Mayyah, Z.,
Engel, J. N., and Persat, A. (2023) Two antagonistic response regulators control
Pseudomonas aeruginosa polarization during mechanotaxis. *EMBO J.* **42**, e112165

35. Leighton, T. L., Yong, D. H., Howell, P. L., and Burrows, L. L. (2016) Type IV Pilus
Alignment Subcomplex Proteins PilN and PilO Form Homo- and Heterodimers *in Vivo*. *J.*
Biol. Chem. **291**, 19923–19938

36. Tammam, S., Sampaleanu, L. M., Koo, J., Sundaram, P., Ayers, M., Chong, P. A., Forman-
Kay, J. D., Burrows, L. L., and Howell, P. L. (2011) Characterization of the PilN, PilO and
PilP type IVa pilus subcomplex. *Mol. Microbiol.* **82**, 1496–1514

37. Geiger, C. J., and O'Toole, G. A. Evidence for the Type IV Pilus Retraction Motor PilT as
a Component of the Surface Sensing System in *Pseudomonas aeruginosa*. *J. Bacteriol.* **205**,
e00179-23

38. Kuchma, S. L., and O'Toole, G. A. (2022) Surface-Induced cAMP Signaling Requires
Multiple Features of the *Pseudomonas aeruginosa* Type IV Pili. *J. Bacteriol.* **204**, e00186-
22

39. Buensuceso, R. N. C., Nguyen, Y., Zhang, K., Daniel-Ivad, M., Sugiman-Marangos, S. N.,
Fleetwood, A. D., Zhulin, I. B., Junop, M. S., Howell, P. L., and Burrows, L. L. (2016) The
Conserved Tetratricopeptide Repeat-Containing C-Terminal Domain of *Pseudomonas*
aeruginosa FimV Is Required for Its Cyclic AMP-Dependent and -Independent Functions.
J. Bacteriol. **198**, 2263–2274

40. Buensuceso, R. N. C., Daniel-Ivad, M., Kilmury, S. L. N., Leighton, T. L., Harvey, H.,
Howell, P. L., and Burrows, L. L. (2017) Cyclic AMP-Independent Control of Twitching
Motility in *Pseudomonas aeruginosa*. *J. Bacteriol.* **199**, e00188-17

41. Topal, H., Fulcher, N. B., Bitterman, J., Salazar, E., Buck, J., Levin, L. R., Cann, M. J.,
Wolfgang, M. C., and Steegborn, C. (2012) Crystal Structure and Regulation Mechanisms
of the CyaB Adenylyl Cyclase from the Human Pathogen *Pseudomonas aeruginosa*. *J.*
Mol. Biol. **416**, 271–286

42. Zhang, Y., Faucher, F., Zhang, W., Wang, S., Neville, N., Poole, K., Zheng, J., and Jia, Z.
(2018) Structure-guided disruption of the pseudopilus tip complex inhibits the Type II
secretion in *Pseudomonas aeruginosa*. *PLoS Pathog.* **14**, e1007343

43. Roberge, N. A., and Burrows, L. L. (2024) Building permits-control of type IV pilus
assembly by PilB and its cofactors. *J. Bacteriol.* **206**, e0035924
44. Dye, K. J., Vogelaar, N. J., Sobrado, P., and Yang, Z. (2021) High-Throughput Screen for
Inhibitors of the Type IV Pilus Assembly ATPase PilB. *mSphere*. **6**, e00129-21
45. McDonald-Ramos, J. S., Hicklin, I. K., Yang, Z., and Brown, A. M. (2024) Identification of
small molecule inhibitors of the *Chloracidobacterium thermophilum* Type IV Pilus protein
PilB by ensemble virtual screening. *Arch. Biochem. Biophys.* 10.1016/j.abb.2024.110127
46. Black, W. P., Wang, L., Jing, X., Saldaña, R. C., Li, F., Scharf, B. E., Schubot, F. D., and
Yang, Z. (2017) The type IV pilus assembly ATPase PilB functions as a signaling protein
to regulate exopolysaccharide production in *Myxococcus xanthus*. *Sci. Rep.* **7**, 7263
47. Mancl, J. M., Black, W. P., Robinson, H., Yang, Z., and Schubot, F. D. (2016) Crystal
Structure of a Type IV Pilus Assembly ATPase: Insights into the Molecular Mechanism of
PilB from *Thermus thermophilus*. *Struct. Lond. Engl. 1993*. **24**, 1886–1897
48. Solanki, V., Kapoor, S., and Thakur, K. G. (2018) Structural insights into the mechanism of
Type IVa pilus extension and retraction ATPase motors. *FEBS J.* **285**, 3402–3421
49. Thongchol, J., Yu, Z., Harb, L., Lin, Y., Koch, M., Theodore, M., Narsaria, U., Shaevitz, J.,
Gitai, Z., Wu, Y., Zhang, J., and Zeng, L. (2024) Removal of *Pseudomonas* type IV pili by
a small RNA virus. *Science*. **384**, eadl0635
50. Qaderi, I., Chan, I., Harvey, H., and Burrows, L. L. (2025) Structural conservation and
functional role of TfpY-like proteins in type IV pilus assembly. *J. Bacteriol.* **207**, e00343-
24
51. Yang, F., Tian, F., Li, X., Fan, S., Chen, H., Wu, M., Yang, C.-H., and He, C. (2014) The
degenerate EAL-GGDEF domain protein Filp functions as a cyclic di-GMP receptor and
specifically interacts with the PilZ-domain protein PXO_02715 to regulate virulence in
Xanthomonas oryzae pv. *oryzae*. *Mol. Plant-Microbe Interact. MPMI*. **27**, 578–589
52. Chin, K.-H., Kuo, W.-T., Yu, Y.-J., Liao, Y.-T., Yang, M.-T., and Chou, S.-H. (2012)
Structural polymorphism of c-di-GMP bound to an EAL domain and in complex with a
type II PilZ-domain protein. *Acta Crystallogr. D Biol. Crystallogr.* **68**, 1380–1392
53. Qi, Y., Xu, L., Dong, X., Yau, Y. H., Ho, C. L., Koh, S. L., Shochat, S. G., Chou, S.-H.,
Tang, K., and Liang, Z.-X. (2012) Functional Divergence of FimX in PilZ Binding and
Type IV Pilus Regulation. *J. Bacteriol.* **194**, 5922–5931
54. Wang, Y.-C., Chin, K.-H., Tu, Z.-L., He, J., Jones, C. J., Sanchez, D. Z., Yildiz, F. H.,
Galperin, M. Y., and Chou, S.-H. (2016) Nucleotide binding by the widespread high-
affinity cyclic di-GMP receptor MshEN domain. *Nat. Commun.* **7**, 12481
55. Floyd, K. A., Lee, C. K., Xian, W., Nametalla, M., Valentine, A., Crair, B., Zhu, S.,
Hughes, H. Q., Chlebek, J. L., Wu, D. C., Hwan Park, J., Farhat, A. M., Lomba, C. J.,
Ellison, C. K., Brun, Y. V., Campos-Gomez, J., Dalia, A. B., Liu, J., Biais, N., Wong, G. C.
L., and Yildiz, F. H. (2020) c-di-GMP modulates type IV MSHA pilus retraction and
surface attachment in *Vibrio cholerae*. *Nat. Commun.* **11**, 1549
56. Aubey, F., Corre, J.-P., Kong, Y., Xu, X., Obino, D., Goussard, S., Lapeyrere, C.,
Souphron, J., Couturier, C., Renard, S., and Duménil, G. (2019) Inhibitors of the *Neisseria
meningitidis* PilF ATPase provoke type IV pilus disassembly. *Proc. Natl. Acad. Sci.* **116**,
8481–8486
57. Hughes, H. Q., Christman, N. D., Dalia, T. N., Ellison, C. K., and Dalia, A. B. (2022) The
PilT retraction ATPase promotes both extension and retraction of the MSHA type IVa pilus
in *Vibrio cholerae*. *PLOS Genet.* **18**, e1010561

58. Ellison, C. K., Dalia, T. N., Klancher, C. A., Shaevitz, J. W., Gitai, Z., and Dalia, A. B. (2021) *Acinetobacter baylyi* regulates type IV pilus synthesis by employing two extension motors and a motor protein inhibitor. *Nat. Commun.* **12**, 3744

59. Keller, H., Kruse, K., Averhoff, B., Duchardt-Ferner, E., and Wöhnert, J. (2019) NMR resonance assignments for the GSPII-C domain of the PilF ATPase from *Thermus thermophilus* in complex with c-di-GMP. *Biomol. NMR Assign.* **13**, 361–366

60. Dye, K. J., Salar, S., Allen, U., Smith, W., and Yang, Z. *Myxococcus xanthus* PilB interacts with c-di-GMP and modulates motility and biofilm formation. *J. Bacteriol.* **205**, e00221-23

61. Dye, K. J., and Yang, Z. (2020) Cyclic-di-GMP and ADP bind to separate domains of PilB as mutual allosteric effectors. *Biochem. J.* **477**, 213–226

62. Hoang, T. T., Karkhoff-Schweizer, R. R., Kutchma, A. J., and Schweizer, H. P. (1998) A broad-host-range Flp-*FRT* recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene*. **212**, 77–86

63. Kilmury, S. L. N., and Burrows, L. L. (2016) Type IV pilins regulate their own expression via direct intramembrane interactions with the sensor kinase PilS. *Proc. Natl. Acad. Sci.* **113**, 6017–6022

64. Deatherage, D. E., and Barrick, J. E. (2014) Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using breseq. *Methods Mol. Biol. Clifton NJ*. **1151**, 165–188

65. Ayers, M., Sampaleanu, L. M., Tammam, S., Koo, J., Harvey, H., Howell, P. L., and Burrows, L. L. (2009) PilM/N/O/P Proteins Form an Inner Membrane Complex That Affects the Stability of the *Pseudomonas aeruginosa* Type IV Pilus Secretin. *J. Mol. Biol.* **394**, 128–142

66. Abramson, J., Adler, J., Dunger, J., Evans, R., Green, T., Pritzel, A., Ronneberger, O., Willmore, L., Ballard, A. J., Bambrick, J., Bodenstein, S. W., Evans, D. A., Hung, C.-C., O'Neill, M., Reiman, D., Tunyasuvunakool, K., Wu, Z., Žemgulytė, A., Arvaniti, E., Beattie, C., Bertolli, O., Bridgland, A., Cherepanov, A., Congreve, M., Cowen-Rivers, A. I., Cowie, A., Figurnov, M., Fuchs, F. B., Gladman, H., Jain, R., Khan, Y. A., Low, C. M. R., Perlin, K., Potapenko, A., Savy, P., Singh, S., Stecula, A., Thillaisundaram, A., Tong, C., Yakneen, S., Zhong, E. D., Zielinski, M., Žídek, A., Bapst, V., Kohli, P., Jaderberg, M., Hassabis, D., and Jumper, J. M. (2024) Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature*. **630**, 493–500

67. Meng, E. C., Goddard, T. D., Pettersen, E. F., Couch, G. S., Pearson, Z. J., Morris, J. H., and Ferrin, T. E. (2023) UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci. Publ. Protein Soc.* **32**, e4792

68. Robert, X., and Gouet, P. (2014) Deciphering key features in protein structures with the new ENDscript server. *Nucleic Acids Res.* **42**, W320-324

69. Ellison, C. K., Kan, J., Dillard, R. S., Kysela, D. T., Ducret, A., Berne, C., Hampton, C. M., Ke, Z., Wright, E. R., Biais, N., Dalia, A. B., and Brun, Y. V. (2017) Obstruction of pilus retraction stimulates bacterial surface sensing. *Science*. **358**, 535–538

SI Figure Captions

Figure S1: Extension regulatory effector mutants produce no detectable surface pili. Representative SDS-PAGE of sheared surface proteins. Major pilin subunits (PilA) are indicated. Sample loading was normalized to flagellin (FliC) levels.

Figure S2: Mutations which suppress the *fimX* twitching deficit do not confer a growth benefit. Bacterial growth curves across 18 hours in LB media. Points represent the means of triplicate samples from three independent experiments ± SD.

Figure S3: Motility suppressor mutants remain susceptible to pilus-targeting phage PO4. Bacterial growth curves across 18 hours in LB media with PO4 phage-challenge. Points represent the means of triplicate samples from three independent experiments ± SD.

Figure S4: FimX complementation increases twitching motility in original Δ*fimX* suppressor isolates. (A) Quantification of sub-agar stab twitching motility zones for Δ*fimX* twitching suppressor mutants complemented with FimX variants. Representative crystal violet-stained twitching zones are shown to the left. Scale bar = 1 cm. Bars represent the means of triplicate samples from three independent experiments ± SD. **(B)** Representative colonies showing pairwise interactions (pink) between FimX and FimX AAA mutant. PilZ was used as a known non-interacting negative control with FimX (52). Untagged T18/T25 plasmid and PilS-T18/PilS-T25 homodimers (63) were used as negative and positive controls respectively. **EV**: empty pHERD30T vector, ***fimX***: *fimX* in pHERD30T vector, **FimX AAA**: FimX AAA in pHERD30T vector.

Figure S5: Expression of CyaB *in trans* bypasses the Pil-Chp network to increase twitching motility. (A) Quantification of sub-agar stab twitching motility zones for a Δ*cyaB* mutant complemented with CyaB or a catalytically inactive mutant (D234S). Representative CyaB or CyaB D234S-expressing BTH101 colonies are shown in the inset to the right. Pink colour results from increased cAMP levels. Representative crystal violet-stained twitching zones are shown to the left. Bars represent the means of triplicate samples from three independent experiments ± SD. **(B)** Quantification of sub-agar stab twitching motility zones for a Δ*pilGH* double mutant complemented with CyaB. Representative twitching zones are shown to the left. Bars represent the means of triplicate samples from three independent experiments ± SD. All scale bars = 1 cm. *******: $p \leq 0.001$ (Two-tailed parametric *t*-test). **EV**: empty pHERD30T vector, **His$_6$-CyaB (D234S)**: N-terminally hexa-histidine tagged CyaB or CyaB D234S in pHERD30T vector.

Figure S6: Mutants with an mRuby3 cassette insertion upstream of the *pilMNOPQ* operon remain sensitive to pilus-targeting phage PO4. Bacterial growth curves across 10 hours in LB media with PO4 phage-challenge. Points represent the means of triplicate samples from two independent experiments ± SD. **CyaB R456L**: CyaB R456L in pBADGr vector.

Figure S7: Type II secretion system-dependent clearing on skim milk agar plates for original *fimX* suppressor isolates. (A) Representative colonies and surrounding zones of clearance for the Δ*fimX* twitching suppressor mutants. Scale bar = 1 cm. **(B)** Quantification of the colony area subtracted from the skim milk clearance area. Bars represent the means of duplicate samples from three independent experiments ± SD. *****: $0.05 \geq p \geq 0.01$; *******: $0.001 \geq p$ (Two-tailed parametric *t*-test).

Figure S8: Alphafold3-predicted models of PilB supressor mutants *pilB* Δ1bp and PilB T430P. (A) Model of *P. aeruginosa* PilB (PilB^{Pa}) monomer. The N1D and linker region (dark grey), N2D (light blue), and CTD (white) are indicated. The *pilB* Δ1bp C-terminal extension is shown in dark blue. PilB T430 side chain is shown in stick and in orange. The model PAE plot is shown to the right. **(B)** *pilB* Δ1bp full length homohexameric Alphafold3 predicted model is shown below with the extended sequence residues highligted in dark blue. The predicted PilC interface is shown on the bottom. The N1Ds and linker regions for each monomer are hidden. The model on the bottom has the two front monomers hidden for clarity. **(C)** X-ray crystal structure of hexameric PilB from *G. metallireducens* (5TSH) with sequence aligned residues of interest N429 (PilB^{Pa} T430) and R430 (PilB^{Pa} R431) highlighted in orange and magenta, respectively. C2-symetric monmers are highlited in the same colours. Dashed lines indicate approximate protomer interfaces.

Figure S9: *pilB* Δ1bp constructs complement twitching motility to at least WT levels. (A) Quantification of sub-agar stab twitching zone areas of a *pilB*::Tn5 mutant overexpressing *pilB* Δ1bp 3′ truncations. Representative crystal violet-stained twitching zones are shown to the left. PilB expression was induced with 0.05% arabinose. Bars represent the means of duplicate samples from three independent experiments ± SD. **(B)** Quantification of sub-agar stab twitching zone areas of WT cells overexpressing *pilB* Δ1bp 3′ truncations. Representative twitching zones are shown to the left. PilB expression was induced with 0.05% arabinose. Bars represent the means of duplicate samples from three independent experiments ± SD. **EV**: empty pHERD30T vector, ***pilB* Δ1bp-3 to 21**: *pilB* Δ1bp-3 to 21 in pHERD30T vector. All scale bars = 1 cm. **ns**: $p \geq 0.05$; ******: $0.01 \geq p \geq 0.001$ (Two-tailed parametric *t*-test).

Figure S10: PilB T430X constructs complement twitching motility to at least WT levels. (A) Quantification of sub-agar stab twitching zone areas of a *pilB*::Tn5 mutant overexpressing T430X mutants. Representative crystal violet-stained twitching zones are shown to the left. PilB

expression was induced with 0.05% arabinose. Bars represent the means of duplicate samples from three independent experiments ± SD. **(B)** Quantification of sub-agar stab twitching zone areas of WT cells overexpressing T430X mutants. Representative twitching zones are shown to the left. PilB expression was induced with 0.05% arabinose. Bars represent the means of duplicate samples from three independent experiments ± SD. **PilB T430X**: indicated PilB substitution in pHERD30T vector. All scale bars = 1 cm. **ns**: $p \geq 0.05$; *****: $0.05 \geq p \geq 0.01$ (Two-tailed parametric t-test).

Figure S11: PilB T429/431X constructs complement twitching motility. (A) Quantification of sub-agar stab twitching zone areas of a *pilB*::Tn5 mutant overexpressing T429/431X mutants. Representative crystal violet-stained twitching zones are shown to the left. PilB expression was induced with 0.05% arabinose. Bars represent the means of duplicate samples from three independent experiments ± SD. **(B)** Quantification of sub-agar stab twitching zone areas of WT cells overexpressing T429/431X mutants. Representative twitching zones are shown to the left. PilB expression was induced with 0.05% arabinose. Bars represent the means of duplicate samples from three independent experiments ± SD. **PilB T429/431X**: indicated PilB substitution in pHERD30T vector. All scale bars = 1 cm. **ns**: $p \geq 0.05$; *******: $0.001 \geq p$ (Two-tailed parametric t-test).

Figure S12: Purified ΔN1D-His$_6$-PilB and FimX protein samples. SDS-PAGE of representative protein samples used in the study. Samples are diluted to the same concentrations added to the reaction mixture preparation.

Figure S13: PilA A86C mutants retain twitching motility. Quantification of sub-agar stab twitching zone areas of PilA A86C mutants. Representative crystal violet stained twitching zones are shown to the left. Bars represent the means of triplicate samples from three independent experiments ± SD. Scale bar = 1 cm.

Figure S14: Vector control assays for FimX localization studies. (A) Quantification of sub-agar stab twitching zone areas of Δ*fimX* mutants complemented with N-terminal fusion of mNeonGreen (mNGr) to FimX. Representative crystal violet stained twitching zones are shown to the left. Expression was not induced with arabinose. Bars represent the means of triplicate samples from three independent experiments ± SD. Scale bar = 1 cm. **EV**: empty pHERD30T, **mNGr-FimX (AAA)**: N-terminal fusion of mNeonGreen to FimX or FimX AAA in pHERD30T vector. **(B)** Representative images of Δ*fimX* mutant cells carrying empty pHERD30T with background fluorescence subtracted. Scale bar = 2 μm. *******: $0.001 \geq p$ (Two-tailed parametric t-test).

1168 **Figure S15: PilB mutations contributing to enhanced twitching motility in Δ*fimX* are rare.**

1169 **(A)** Total protein sequence length of PilB orthologues in *P. aeruginosa* from the

1170 *Pseudomonas*.com database. Sequences which are the same length as PAO1 PilB (566 residues)

1171 are indicated in blue. **(B)** Percentage of sequences with an identifiable α-helix-disrupting residue

1172 ± three amino acids from the position which aligns with PAO1 PilB T430.

1173

1174

1175

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

Fig 7

Fig 8