Exploring Hypoxia-Related Genes as Prognostic Indicators in Lung Adenocarcinoma

Bandar Alghamdi[1] and Sonia Rocha[2]*

[1]IKing Fahad Army Forces Hospital, cardiology and cardiac surgery center, Jeddah, Saudi Arabia.

[2]Department of Biochemistry, Cell and Systems Biology, Institute of Systems Molecular and Integrative Biology, University of Liverpool. Liverpool United Kingdom.

*correspondence to: Sonia Rocha srocha@liverpool.ac.uk

Abstract

Background: Lung adenocarcinoma (LUAD) is a leading cause of cancer-related mortality, with hypoxia contributing to tumor progression and treatment resistance. Identifying hypoxia-related biomarkers could enhance prognosis and therapeutic strategies for LUAD. **Methods:** This study aimed to identify hypoxia-related differentially expressed genes (HRDEGs) in lung adenocarcinoma (LUAD) through differential expression analysis. Functional analysis and protein-protein interaction (PPI) network construction were performed to explore the biological roles and interactions of these genes. Kaplan-Meier survival analysis, univariate Cox regression, and Lasso regression were used to identify key genes associated with survival. Multivariate Cox regression was then conducted to assess independent prognostic factors. **Results:** This analysis revealed 283 upregulated HRDEGs and 322 downregulated HRDEGs in LUAD. Functional enrichment analysis indicated that the upregulated genes were primarily involved in cancer-related and cellular signaling pathways, while downregulated genes were associated with immunity-related pathways. We further identified 201 common upregulated hub genes (including *MMP9, CDH1, HSP90AB1, SOX2, CDKN2A, SPP1, EZH2*) and 224 common downregulated hub genes (such as *IL6, TNF, IL1B, JUN, CCL2, TLR4, FOS, PTGS2*). Kaplan-Meier survival analysis, univariate Cox regression, and Lasso regression led to the identification of 17 key genes (*ADRB2, ALDH2, CAT, CCNE1, MAP3K8, DSG2, EIF6, ABCB1, PIK3R1, RAD51, SFTPD, SOD3, CLEC3B, ADAM12, EXO1, FBLN5,* and *IGF2BP3*) associated with patient survival. Finally, multivariate Cox regression analysis identified *DSG2, EIF6,* and *EXO1* as independent prognostic factors for LUAD, highlighting their potential as biomarkers for prognosis and therapeutic targets in lung cancer. **Conclusion:** In conclusion, *DSG2, EIF6,* and *EXO1* were identified as key hypoxia-related genes in lung adenocarcinoma. These genes were

found to be independent prognostic factors, highlighting their potential as biomarkers for

predicting patient survival and guiding future therapeutic approaches.

Keyword: Lung Adenocarcinoma, Hypoxia-Related Differentially Expressed Genes, immunity,

independent prognostic factors

Authors summary:

This study investigates the role of hypoxia-related genes in lung adenocarcinoma (LUAD), a major cause of cancer mortality. By analyzing gene expression data, this study identified 283 upregulated and 322 downregulated hypoxia-related differentially expressed genes (HRDEGs). Functional enrichment analysis revealed that the upregulated genes are largely involved in cancer progression and cell signaling, while the downregulated ones are linked to immune responses. A set of key hub genes was identified through protein-protein interaction analysis. Using survival analysis tool, 17 genes significantly associated with patient outcomes in two different patient cohorts. Among these, confirmed three genes—DSG2, EIF6, and EXO1—were confirmed as independent prognostic markers. These findings suggest that these genes may serve as valuable biomarkers for predicting survival and developing targeted therapies in LUAD, offering new avenues for improving patient management in hypoxia-driven tumors.

Introduction

Lung Adenocarcinoma and Hypoxia Lung cancer remains one of the most common and deadly cancers worldwide, with non-small cell lung cancer (NSCLC) being the dominant histological subtype (1). Lung adenocarcinoma (LUAD) is the most prevalent histological subtype of NSCLC and represents a major clinical and biological focus within this disease category due to its distinct molecular and genetic characteristics. Although advances in molecular profiling and targeted therapies have improved the management of NSCLC, the prognosis for patients with LUAD remains poor, particularly in advanced stages (2). The overall five-year survival rate for LUAD patients is low, making it essential to identify novel biomarkers for early diagnosis, prognosis, and therapeutic targeting (3). One of the critical microenvironments influencing cancer progression, metastasis, and resistance to therapy is tumor hypoxia. Hypoxia refers to the state of low oxygen availability within the tumor, often resulting from an imbalance between the rapid growth of tumor cells and the insufficient development of blood vessels to supply oxygen (4). Hypoxia is a hallmark feature of many solid tumors, including lung adenocarcinoma, and it is associated with aggressive tumor behavior, poor prognosis, and resistance to conventional therapies (5). The development of therapeutic strategies targeting hypoxic tumor cells remains an area of active research.

The Role of Hypoxia in Tumorigenesis Hypoxia plays a crucial role in driving various aspects of tumor biology, such as tumor angiogenesis, metabolic reprogramming, and epithelial-to-mesenchymal transition (EMT) (6). It activates a range of adaptive cellular responses through hypoxia-inducible factors (HIFs), primarily HIF-1α and HIF-2α. These transcription factors regulate the expression of genes that mediate critical processes such as angiogenesis (formation of new blood vessels), glycolysis, metastasis, and drug resistance (7, 8). Tumor cells exposed to

hypoxic conditions undergo metabolic shifts that favor anaerobic glycolysis, which allows them

to survive and proliferate despite low oxygen availability. The HIF signaling pathway is essential

for the regulation of genes involved in cellular survival, metabolism, and tumor progression (9).

Among the most well-known hypoxia-regulated genes are vascular endothelial growth factor

(VEGF), glucose transporter proteins (e.g., GLUT1), and matrix metalloproteinases (e.g.,

MMPs) (10). These genes promote tumor growth and metastasis by facilitating the adaptation of

tumor cells to low-oxygen conditions. Additionally, hypoxia is implicated in the regulation of

immune evasion and resistance to chemotherapies and radiation therapy, further complicating

treatment strategies (11). Therefore, the identification of key hypoxia-related genes and their role

in LUAD progression and prognosis is crucial for improving the clinical management of lung

adenocarcinoma.

The link between hypoxia and poor prognosis in LUAD has been well established through

various studies. Tumors with hypoxic regions have been shown to have a higher potential for

metastasis, increased tumor aggressiveness, and reduced sensitivity to treatment (12). Patients

with high expression levels of hypoxia-related genes often exhibit shorter overall survival and

progression-free survival compared to those with low expression levels of these genes

(13).Several studies have sought to identify specific hypoxia-related biomarkers that could serve

as predictors of patient outcomes in LUAD (14). However, the heterogeneity of LUAD, coupled

with the complexity of hypoxia-induced changes in gene expression, presents challenges in

pinpointing specific prognostic genes. Additionally, the tumor microenvironment in LUAD is

influenced by various factors, such as genetic mutations, immune responses, and stromal

interactions, which may confound the impact of hypoxia on survival (15, 16).

In this context, a comprehensive analysis of hypoxia-related genes and their potential prognostic

value in LUAD is crucial. Identifying genes that are consistently associated with poor prognosis,

regardless of other clinical variables, will allow for the development of better prognostic models.

These models could guide personalized treatment strategies, such as targeting hypoxic regions of

the tumor with hypoxia-activated prodrugs or improving the efficacy of therapies through the

modulation of the HIF pathway.

This study aims to systematically investigate hypoxia-related genes in LUAD and evaluate their

potential prognostic significance. We used a combination of computational and statistical

approaches to analyze large-scale genomic datasets. By identifying key hypoxia-associated

genes, this work seeks to find independent prognostic biomarkers and provide insights into

hypoxia-driven mechanisms underlying disease progression in LUAD.

Methods

Data downloading and processing

The analysis utilized two distinct datasets: the TCGA LUAD cohort and GSE18842. We downloaded the TCGA LUAD cohort form the cBioPortal (17). The TCGA LUAD cohort comprised 517 tumor samples and 59 control samples, representing lung adenocarcinoma (LUAD) patients and normal controls, respectively (https://www.cbioportal.org/study/summary?id=luad_tcga). Clinical data (https://www.cbioportal.org/study/summary?id=luad_tcga) from this cohort were incorporated for Kaplan-Meier survival analysis, univariate regression analysis, and multivariate regression analysis to investigate the relationship between gene expression and survival outcomes. For the GSE18842 dataset, which contains 46 tumor samples and 45 control samples. data normalization was performed using a log2 transformation. To account for multiple probes corresponding to individual genes, the data was averaged, resulting in a single expression value for each gene. The GSE18842 dataset is publicly available and can be accessed at https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE18842.

Differentially expressed genes identifications

In this study, we utilized multiple datasets to identify differentially expressed genes (DEGs) associated with tumorigenesis. To identify DEGs between tumor and normal samples, we employed the LIMMA package, applying a stringent cutoff of $|log2FC| > 0.585$ and an adjusted p-value < 0.05. Following DEG identification, we specifically targeted hypoxia-related differentially expressed genes (HRDEGs), which were selected through a Venn diagram approach. This method allowed us to cross-reference the LUAD DEGs and further refine the set of genes related to hypoxia, thus providing a focused analysis of the genes potentially involved in the tumor microenvironment's hypoxic conditions.

Curation of hypoxia related genes and identification of hypoxia related differentially expressed genes (HRDEGs)

To curate a comprehensive list of hypoxia-related genes, we integrated data from several established resources. First, we retrieved hypoxia-related genes from Gene Set Enrichment Analysis (GSEA) (https://www.gsea-msigdb.org/gsea/index.jsp), focusing on the following gene sets: GOBPHYPOXIAINDUCIBLEFACTOR1ALPHASIGNALINGPATHWAY, GOBPINTRINSICAPOPTOTICSIGNALINGPATHWAYINRESPONSETOHYPOXIA, GOBPNEGATIVEREGULATIONOFCELLULARRESPONSETOHYPOXIA, GOBPNEGATIVEREGULATIONOFHYPOXIAINDUCEDINTRINSICAPOPTOTICSIGNALI NGPATHWAY,GOBPREGULATIONOFCELLULARRESPONSETOHYPOXIA, HALLMARKHYPOXIA, and REACTOMEREGULATIONOFGENEEXPRESSIONBYHYPOXIAINDUCIBLEFACTOR.

We then performed a search on OMIM (https://www.ncbi.nlm.nih.gov/omim/?term=HYPOXIA) using the term "HYPOXIA" to identify genes associated with hypoxia-related genetic disorders. Additionally, we retrieved genes from the hsa04066:HIF-1 signaling pathway in the KEGG database (https://www.genome.jp/kegg/). Finally, we obtained genes from GeneCards (https://www.genecards.org/), selecting those with a score greater than or equal to 1. After compiling the data from these sources, we performed deduplication to ensure that only unique genes were included. This resulted in a final list of 3292unique genes, which are tabulated in **Supplementary Table S1**.

Functional enrichment analysis of DEGs

We used DAVID v6.8 (https://david.ncifcrf.gov/) (18) to perform functional annotation of the

identified genes. This platform allowed us to categorize the genes into various biological

contexts, such as significantly enriched KEGG pathways, biological processes, cellular

components, and molecular functions. These functional analyses provided valuable insights into

the roles and interactions of the genes in key cellular mechanisms. To assess the statistical

significance of the enrichment, we applied a false discovery rate (FDR) threshold of < 0.05,

calculated using the Benjamini–Hochberg method (19), ensuring that only robust and

biologically relevant findings were included.

165 Exploration of hub HRDEGs

We constructed Protein-Protein Interaction (PPI) networks for the differentially expressed genes

(DEGs) using STRING v11.0 (https://string-db.org/) (20)). Both upregulated and downregulated

genes were entered into the STRING search module to generate the interaction networks. To

identify key hub genes within these networks, we utilized the cytoHubba plugin in Cytoscape

(https://cytoscape.org/) (21),which ranks genes based on their interaction degree. A medium

interaction score threshold of ≥0.40 was applied to build the PPI network, and genes with an

interaction degree of ≥10 were selected for further analysis. This approach enabled us to focus on

the most significant hub genes, highlighting those with strong interactions relevant to lung

cancer.

175 Exploration of survival associated hub HRDEGs

We retrieved clinical data from TCGA LUAD to perform an analysis of overall survival (OS) in

relation to gene expression. Patients were categorized into two groups based on their gene

expression levels, either above or below the median, and survival rates between these groups

were compared. To further explore survival differences, patients were also stratified into risk

groups based on their gene expression profiles. The log-rank test and Kaplan-Meier survival

curves were utilized to evaluate and visualize differences in survival outcomes. For a more

detailed survival analysis, we used the R package "survival" (22), performing both univariate and

multivariate Cox regression analyses to assess the independent effects of genes on survival.

Additionally, we applied LASSO-Cox regression via the "glmnet" package (23), incorporating

10-fold cross-validation to optimize the selection of penalty parameters and minimize the risk of

overfitting. This comprehensive approach allowed us to identify significant survival-associated

genes while controlling for potential biases and overfitting, ensuring robust and reliable findings.

Validation of Prognostic Signature in an Independent Cohort

To validate the prognostic relevance of the identified three-gene signature, we utilized the

independent LUAD dataset GSE13213 via the SurvExpress platform (24). SurvExpress is an

online tool that allows assessment of gene expression-based prognostic models across

independent cohorts. The expression levels of gene signature were used to calculate a risk score

for each patient, and patients were stratified into high-risk and low-risk groups based on the

median risk score. Survival differences between the groups were evaluated using Kaplan-Meier

survival analysis and the Log-Rank test**,** and hazard ratios (HRs) with 95% confidence intervals

were computed to assess prognostic significance. Additionally, expression differences of gene

signature between the risk groups were compared to confirm consistency with the primary cohort

findings.

Statistical analysis and overall methodology

All statistical analyses were performed using R software (version 4.0.1). Differentially expressed genes (DEGs) were identified using a cutoff threshold of |log$_2$ fold change (log$_2$FC)|> 0.585 and an adjusted p-value < 0.05. Multiple testing correction was conducted using the Benjamini–Hochberg false discovery rate (FDR) method, with an FDR threshold of < 0.05 applied to control for false positives (19). The overall methodology provided in the Figure 1.

209 **Figure 1.** The overall methodology of the study.

Result and discussion

Identification of hypoxia related differentially expressed genes (HRDEGs) in Lung adenocarcinoma

We explored differentially expressed genes (DEGs) in the TCGA LUAD cohort (**Supplementary Table S2**) and identified additional DEGs from the GSE18842 dataset (**Supplementary Table S3**). Volcano plots depicting the results of both analyses are shown in **Figure 2A** and **Figure 2B**, respectively. These plots provide a visual representation of the relationship between fold changes (logFC) and adjusted p-values, helping to identify genes with significant differential expression in lung adenocarcinoma (LUAD). Next, we compared the identified DEGs with hypoxia-related genes to determine the hypoxia-related differentially expressed genes (HRDEGs). This comparison revealed a total of 283 upregulated HRDEGs and 322 downregulated HRDEGs in LUAD, which are displayed in **Figure 2C-D** and detailed in **Supplementary Table S4**. These findings indicate that specific hypoxia-related genes are significantly altered in LUAD, providing potential insights into the molecular mechanisms underlying the disease. The top ten upregulated and downregulated HRDEGs are listed in Table 1. Among the top upregulated HRDEGs are *CYP24A1, CA9, MMP13, SPP1, COL10A1, MMP1, EEF1A2, GREM1, PITX1,* and *MMP12*. These genes are associated with various biological processes such as cell metabolism, matrix remodeling, and tumor progression. On the other hand, the top downregulated HRDEGs include *SFTPC, SLC6A4, AGER, FABP4, SCGB1A1, VEGFD, HBB, SFTPA2, AQP4,* and *CLEC3B*, which are involved in processes like lung surfactant production, cell signaling, and vasculature maintenance.

Figure 2. Volcano plots and Venn diagrams depicting DEGs and HRDEGs in LUAD.A. Volcano plot of DEGs in the TCGA LUAD cohort. **B.** Volcano plot of DEGs in the GSE18842 dataset, illustrating differential gene expression in LUAD. **C.** Venn diagram showing the overlap of upregulated HRDEGs in LUAD. **D**. Venn diagram illustrating the overlap of downregulated HRDEGs in LUAD.TCGA UP: TCGA-LUAD upregulated; TCGA DOWN: TCGA-LUAD downregulated; HRG: hypoxia-related genes

Table 1. The top ten upregulated and downregulated HRDEGs in TCGA and GSE18842

Regulatory status	Gene ID	Gene Symbol	TCGA		GSE18842		Name of the genes
			logFC	Adjusted P Value	logFC	Adjusted P Value	
Upregulated	1591	CYP24A1	5.211	8.81E-33	1.293	9.12E-06	cytochrome P450 family 24 subfamily A member 1
	768	CA9	4.929	2.12E-27	1.749	3.30E-13	carbonic anhydrase 9
	4322	MMP13	4.885	2.07E-29	1.825	1.41E-07	matrix metallopeptidase 13
	6696	SPP1	4.739	1.41E-49	2.540	2.92E-20	secreted phosphoprotein 1
	1300	COL10A1	4.653	6.9E-60	3.547	8.02E-21	collagen type X alpha 1 chain
	4312	MMP1	4.651	1.47E-33	5.110	5.85E-21	matrix metallopeptidase 1
	1917	EEF1A2	4.297	1.69E-18	0.989	2.14E-05	eukaryotic translation elongation factor 1 alpha 2
	26585	GREM1	4.257	1.7E-37	4.666	8.94E-24	"gremlin 1, DAN family BMP antagonist"
	5307	PITX1	4.233	9.46E-32	1.629	6.43E-17	paired like homeodomain 1
	4321	MMP12	4.085	1.72E-29	5.626	7.05E-29	matrix metallopeptidase 12
Downregulated	6440	SFTPC	-8.560	3.4E-38	-6.0378	1.01E-25	surfactant protein C
	6532	SLC6A4	-7.971	7.8E-90	-1.9304	1.49E-15	solute carrier family 6 member 4
	177	AGER	-6.765	2.62E-72	-4.9568	1.01E-44	advanced glycosylation end-product specific receptor
	2167	FABP4	-6.160	9.71E-77	-3.4366	5.77E-26	fatty acid binding protein 4
	7356	SCGB1A1	-5.786	2.46E-22	-3.9076	3.06E-08	secretoglobin family 1A member 1
	2277	VEGFD	-5.008	2.11E-46	-4.4324	4.79E-40	vascular endothelial growth factor D
	3043	HBB	-4.916	1.54E-51	-2.5255	1.07E-25	hemoglobin subunit beta
	729238	SFTPA2	-4.863	6.09E-23	-3.7907	6.99E-10	surfactant protein A2
	361	AQP4	-4.688	1.29E-27	-3.9426	9.98E-26	aquaporin 4
	7123	CLEC3B	-4.241	2.27E-72	-4.4159	8.41E-38	C-type lectin domain family 3 member B

HRDEGs are associated with cellular processes such as cell cycle and metabolism.

Figure 3 illustrates the pathways significantly enriched by the upregulated genes, with a false

discovery rate (FDR) < 0.05. These pathways highlight key biological processes and molecular

mechanisms potentially associated with tumorigenesis (**Supplementary Table S5**). Notably, the

cell cycle (hsa04110), biosynthesis of amino acids (hsa01230), and carbon metabolism

(hsa01200) are central to cellular growth and energy production. The p53 signaling pathway

(hsa04115) and cellular senescence (hsa04218) are also implicated, reflecting the regulation of

cell death and aging processes. Other enriched pathways, such as glycolysis/gluconeogenesis

(hsa00010) and HIF-1 signaling (hsa04066), suggest alterations in metabolic pathways and

hypoxic responses often observed in cancer. Additionally, pathways related to cancer

progression, such as central carbon metabolism in cancer (hsa05230), pathways in cancer

(hsa05200), and microRNAs in cancer (hsa05206), are also prominently featured, highlighting

the complex molecular interactions in cancer biology. Other enriched pathways include DNA

replication (hsa03030), mismatch repair (hsa03430), and glutathione metabolism (hsa00480),

which are critical for genome stability and oxidative stress responses. The statistical analysis of

upregulated pathways is tabulated in Supplementary Table S5. These results emphasize the

multifaceted nature of the upregulated genes and their involvement in various cellular functions

relevant to cancer development.

Figure 3. Significantly enriched pathways associated with the upregulated genes (FDR < 0.05). DEG upregulated were analyzed for functional pathways using DAVID.

The top enriched pathway, the Cell Cycle (**Figure 4**), with a highly significant p-value of 4.51E-06. This pathway is critical for regulating cell division and proliferation. The genes enriched within this pathway, including *PCNA, MCM7, CDKN2A, PRKDC, HDAC1, CDC6,* and others such as *CDK4, CHEK1,* and *E2F3*, play pivotal roles in cell cycle aberration. These genes are integral to maintaining proper cell cycle regulation, and their upregulation suggests a potential disruption in normal cell division processes, which is commonly associated with tumorigenesis. The enrichment of these genes further underscores the importance of the cell cycle in hypoxic lung cancer biology.

Figure 4: KEGG pathway illustrating the Cell Cycle pathway from DAVID, with upregulated genes marked by red stars.

Similar to the Cell Cycle pathway, another significantly enriched pathway is the HIF-1 signaling pathway (hsa04066), with genes such as *LDHA, EGLN3, SLC2A1, EIF4EBP1, PGK1, TIMP1, ENO1, ALDOA, GAPDH, PFKP,* and *PDK1* being upregulated (**Figure 5**). The upregulation of these genes suggests a critical role of hypoxia in lung cancer, as these genes are involved in metabolic adaptation to low oxygen conditions (25). This highlights the importance of hypoxia-

288 induced signaling in promoting tumor survival and progression through altered cellular

289 metabolism.

Figure 5: KEGG pathway illustrating the HIF-1 signaling pathway from DAVID, with upregulated genes marked by red stars. Upregulated genes, such as *LDHA, EGLN3, SLC2A1, EIF4EBP1, PGK1, TIMP1, ENO1, ALDOA, GAPDH, PFKP, and PDK1*, are shown with red stars, indicating their involvement in metabolic reprogramming and cellular adaptation to low oxygen conditions.

In addition to the upregulated genes, we observed that the downregulated genes are

predominantly associated with immune-related pathways and signaling processes. As illustrated

in **Figure 6**, several immune-related pathways are significantly affected, indicating a potential

suppression of immune responses in lung cancer. Key pathways include the TNF signaling

pathway (hsa04668), Cytokine-cytokine receptor interaction (hsa04060), and IL-17 signaling

pathway (hsa04657), all of which play essential roles in immune activation and inflammation.

Other important pathways such as Toll-like receptor signaling (hsa04620), JAK-STAT signaling

(hsa04630), and Natural killer cell-mediated cytotoxicity (hsa04650) are also downregulated,

suggesting that the immune system's ability to recognize and eliminate cancer cells may be

impaired. The statistical analysis of upregulated pathways is tabulated in **Supplementary Table**

S6. These findings suggests a potential immune evasion mechanism in lung cancer, where the

downregulation of key immune pathways may allow tumors to escape immune surveillance and

progress unchecked.

Figure 6: Immune-associated pathways enriched in the downregulated genes. This figure illustrates the major immune-related pathways that are significantly downregulated in lung cancer.

Explorations of hub HRDEGs

Next, we conducted a Protein-Protein Interaction (PPI) analysis by inputting all the hypoxia-related differentially expressed genes (DEGs) into the STRING database. The primary PPI network consisted of 600 nodes and 9,281 edges, with an average node degree of 30.9 (**Figure 7**). The PPI enrichment p-value was found to be < 1.0e-16, indicating a highly significant

interaction among the hypoxia-related genes. Based on a degree of interaction greater than 10,

we identified the hub genes within the network. After identifying the hub genes based on the

degree of interaction, we found that the top hub genes include *GAPDH, IL6, TNF, IL1B, MMP9,*

JUN, CCL2, TLR4, FOS, PTGS2, PECAM1, CDH1, HSP90AB1, PTPRC, ICAM1, FGF2,

PPARG, ITGAM, CXCL12, and KDR. The list of all interacted genes and their degree of

interaction provided in the **Supplementary Table S7**. These genes play pivotal roles in key

biological processes such as inflammation, immune response, cell signaling, and tumor

progression. **Figure 8A** illustrates the interaction network of the top 50 hub genes an,

highlighting their interconnections and potential involvement in the molecular pathways that

drive lung cancer progression. We further identified 201 common upregulated hub genes and 224

common downregulated hub genes, which were shared across the datasets. These genes represent

key molecular players in the tumorigenic processes of lung cancer, with their dysregulation

potentially contributing to cancer progression. The results of this analysis are presented in

Figure 8B and are detailed in **Supplementary Table S7**.

Figure 7. Primary Protein-Protein Interaction (PPI) network of hypoxia-related differentially expressed genes (DEGs) generated using STRING. The network consists of 600 nodes and 9,281 edges, with an average node degree of 30.9.

The top upregulated hub genes identified in this analysis include *MMP9, CDH1, HSP90AB1, SOX2, CDKN2A, SPP1, EZH2, TIMP1, CXCL10, COL1A1, HDAC1, BRCA1,* and *CDK4*. These genes are involved in critical processes such as tumor invasion, immune modulation, cell cycle regulation, and epigenetic changes. The top downregulated hub genes identified in this analysis *include IL6, TNF, IL1B, JUN, CCL2, TLR4, FOS, PTGS2, PECAM1, PTPRC, ICAM1, FGF2,*

PPARG, ITGAM, CXCL12, and *KDR*. These genes are primarily involved in immune response, inflammation, and cellular signaling pathways.

Figure 8. Protein-Protein Interaction (PPI) Network of Hub Genes. A. Interaction network of the top 50 hub genes identified from the PPI analysis. B. Common hub genes shared between the upregulated and downregulated gene sets.

Exploration of independent prognostic HRDEGs

We first conducted Kaplan-Meier survival analysis on all hub genes and identified 174 significant genes that were associated with survival (**Supplementary Table S8**). The following seven genes (Identified by Univariate regression and LASO)—*ADAM12* (HR=1.39, P=0.02), *CCNE1* (HR=1.43, P=0.01), *DSG2* (HR=1.58, P=0.002), *EIF6*(HR=1.49, P=0.008), EXO1

(HR=1.54, P=0.003), *IGF2BP3* (HR=1.6, P=0.001), and *RAD51* (HR=1.37, P=0.03)—are

identified as upregulated hub genes in lung cancer (**Figure 9A**).

The upregulated group, categorized based on median expression levels, is associated with worse

survival outcomes. Increased expression of these genes is correlated with poorer prognosis,

suggesting that they may play a critical role in tumor progression and could serve as potential

biomarkers for identifying high-risk patients with lung cancer. The following ten genes

((Identified by Univariate regression and LASO))—*ABCB1*(HR=0.71, P=0.02),

ADRB2(HR=0.67, P=0.008), *ALDH2*(HR=0.66, P=0.006), *CAT*(HR=0.73, P=0.03),

CLEC3B(HR=0.67, P=0.006), *FBLN5*(HR=0.64, P=0.003), *MAP3K8*(HR=0.67, P=0.006),

PIK3R1(HR=0.64, P=0.003), *SFTPD*(HR=0.68, P=0.01), and *SOD3*(HR=0.75, P=0.04)—are

identified as downregulated hub genes in lung cancer.

A

Figure 9. Kaplan-Meier Survival Analysis of Upregulated and Downregulated Hub Genes in Lung Cancer. A. Kaplan-Meier survival analysis of the group of upregulated hub genes, showing their significant association with survival in lung cancer patients. (B) Kaplan-Meier survival analysis of the group of downregulated hub genes, demonstrating their significant impact on survival outcomes.

To refine this list, we performed univariate regression analysis, which revealed 106 genes with statistically significant associations (**Supplementary Table S9**). Subsequently, we applied Lasso regression to the consistently upregulated and downregulated genes to identify key genes with

389   non-zero coefficients. This analysis led to the identification of 17 key genes: *ADRB2, ALDH2,*

390   *CAT, CCNE1, MAP3K8, DSG2, EIF6, ABCB1, PIK3R1, RAD51, SFTPD, SOD3, CLEC3B,*

391   *ADAM12, EXO1, FBLN5,* and *IGF2BP3*. The univariate regression analysis and LASSO

392   regression analysis depicted in the **Figure 10** and **Figure 11**, respectively.

Figure 10: The results of univariate Cox regression analysis for hub genes in lung cancer patients. Each gene's hazard ratio (HR) with 95% confidence interval (CI) is displayed, indicating the association between gene expression and patient survival.

Figure 11: Lasso Regression Analysis of Key Genes. Identification of key genes through Lasso regression. (A) Lasso coefficients for each gene selected through Lasso regression. The plot shows the evolution of the coefficients for each gene as the regularization penalty (lambda) increases. (B) Log(lambda) vs. coefficients plot demonstrating the relationship between the log-transformed lambda values and the corresponding coefficients.

Furthermore, a multivariate Cox regression analysis was conducted to identify independent prognostic factors. The results indicated that *DSG2, EIF6,* and *EXO1* emerged as independent prognostic factors (**Figure 12**). The multivariate model demonstrated a significant fit, with the following statistics: Concordance = 0.657 (SE = 0.024), Likelihood ratio test = 50.39 on 17

410   degrees of freedom (df), p = 4e-05, Wald test = 48.47 on 17 df (p = 7e-05), and Score (logrank)

411   test = 48.73 on 17 df (p = 7e-05). *DSG2* (HR = 1.238, p = 0.012), *EIF6* (HR = 1.387, p = 0.046),

412   and *EXO1* (HR = 1.257, p = 0.047) all have hazard ratios greater than 1, indicating that higher

413   expression of these genes is associated with poorer survival outcomes.

Figure 12: Multivariate Cox regression analysis identifying independent prognostic factors.

Validation of prognostic value in an independent cohort

To validate the prognostic relevance of *DSG2, EIF6*, and *EXO1* in our risk model, we utilized the independent dataset GSE13213 through the SurvExpress platform (24). The analysis revealed that all three genes maintained a significant association with patient prognosis in this external cohort. Specifically, DSG2 (coefficient: 0.2795, p-value: 0.05), EIF6 (coefficient: 0.8356, p-value: 0.019), and EXO1 (coefficient: 0.7033, p-value: 0.001) were all positively correlated with poor survival outcomes. The overall model performance was statistically robust, as indicated by a Log-Rank $p < 0.001$ and a hazard ratio (HR) of 3.19, underscoring the model's strong prognostic power in stratifying patients.

Further stratification of the cohort into high-risk and low-risk groups based on the risk scores revealed a significant survival difference, with patients in the high-risk group exhibiting markedly shorter survival times compared to those in the low-risk group (**Figure 13A**). Moreover, expression analysis confirmed that DSG2, EIF6, and EXO1 were significantly upregulated in the high-risk group, consistent with our primary findings (**Figure 13B**). These validation results provide strong support for the clinical applicability of the proposed gene signature and reinforce the role of these three genes as potential biomarkers for risk stratification and prognosis in lung cancer.

Figure 13. Validation of Prognostic Gene Signature Using Independent Cohort (GSE13213). A: Survival Analysis Based on Risk Groups in GSE13213 Dataset utilizing SurvExpress platform. B: Expression Levels of DSG2, EIF6, and EXO1 in High- and Low-Risk Groups.

Discussion

Lung adenocarcinoma remains one of the leading causes of cancer-related mortality worldwide, with hypoxia recognized as a key contributor to tumor progression, metastasis, and therapy resistance. In this study, we comprehensively analyzed hypoxia-related differentially expressed genes (HRDEGs) in LUAD and identified 283 upregulated and 322 downregulated HRDEGs, reflecting the broad impact of hypoxic signaling on multiple cellular processes. Functional enrichment and protein-protein interaction analyses revealed that upregulated genes were primarily involved in cancer-related and cellular signaling pathways, whereas downregulated

genes were enriched in immunity-associated pathways, highlighting the dual role of hypoxia in

promoting tumor growth while modulating immune responses.

The enrichment of cell cycle–related genes, including *PCNA, MCM7, CDKN2A, PRKDC,*

HDAC1, CDC6, CDK4, CHEK1, and *E2F3*, highlights a pronounced dysregulation of cell cycle

control mechanisms in lung cancer. These genes are central to DNA replication, cell cycle

progression, and checkpoint regulation, ensuring genomic integrity under normal physiological

conditions (26-28). Their coordinated upregulation suggests enhanced proliferative capacity and

impaired checkpoint surveillance, which are hallmark features of tumorigenesis. In particular,

aberrant activation of CDKs and E2F transcription factors can drive uncontrolled entry into the S

phase (29), while altered expression of DNA damage response genes such as *CHEK1* and

PRKDC may allow cancer cells to bypass critical repair checkpoints, thereby promoting genomic

instability (30, 31). Collectively, these findings underscore the pivotal role of cell cycle

dysregulation in lung cancer development and progression.

In addition to cell cycle alterations, the HIF-1 signaling pathway (hsa04066) emerged as another

significantly enriched pathway, with notable upregulation of genes including *LDHA, EGLN3,*

SLC2A1, EIF4EBP1, PGK1, TIMP1, ENO1, ALDOA, GAPDH, PFKP, and *PDK1*. These genes

are key mediators of cellular adaptation to hypoxic conditions and are closely associated with

metabolic reprogramming in cancer. The elevated expression of glycolytic enzymes and glucose

transporters, such as *SLC2A1* and *LDHA*, reflects a shift toward aerobic glycolysis (the Warburg

effect), enabling tumor cells to sustain rapid growth in oxygen-deprived microenvironments (32).

Taken together, the concurrent activation of cell cycle and HIF-1 signaling pathways suggests a

coordinated mechanism by which lung cancer cells promote unchecked proliferation while

adapting to hypoxic stress. These pathways may act synergistically to facilitate tumor

progression and aggressiveness, highlighting their potential as therapeutic targets and prognostic

biomarkers in lung cancer.

Beyond the activation of oncogenic and metabolic programs, lung cancer was characterized by a

marked suppression of immune-associated pathways, reflecting the establishment of an

immunosuppressive tumor microenvironment (33). The coordinated downregulation of signaling

cascades involved in cytokine communication, inflammatory responses, and immune cell

activation indicates a weakened immune network within tumors (34). Furthermore, attenuation

of pathways governing innate immune sensing and effector mechanisms suggests reduced

recognition and clearance of malignant cells by immune surveillance systems (35). Such

widespread impairment of immune signaling is consistent with tumor-driven immune evasion, in

which cancer cells actively modulate immune pathways to avoid immune-mediated destruction

(36). This immune suppression may not only facilitate tumor progression but also contribute to

resistance to immune-based therapies, underscoring the critical role of immune pathway

dysregulation in lung cancer pathogenesis (34). Interestingly, we have previously shown that

HIF-1 suppresses NF-κB signalling (37), one of the pathways found to be significantly

downregulated in the analysis of lung adenocarcinoma presented here.

Moreover, the top upregulated hub genes, including, *MMP9* and *SPP1* play key roles in

extracellular matrix remodeling and metastasis (38, 39), while *CDH1* is important for

maintaining cell-cell adhesion (40). Other upregulated hub genes, including *SOX2* and *EZH2*

regulate stem cell-like properties and epigenetic modifications, contributing to cancer stem cell

maintenance and progression (41, 42). *CDK4* and *CDKN2A* are crucial regulators of the cell

cycle, promoting cell division and tumor growth (43, 44). *BRCA1* and *HDAC*1 are involved in

DNA repair and chromatin remodeling (45, 46), while *CXCL10* and *TIMP1* are associated with

immune responses (47, 48). The upregulation of these hub genes highlights their critical roles in

tumor progression and their potential as therapeutic targets in lung cancer. On the other hand, the

downregulation of IL6, TNF, and IL1B suggests impaired inflammatory signaling (49), while

TLR4, FOS, and *ICAM1* are involved in immune cell activation and adhesion (50, 51), indicating

potential immune evasion mechanisms. *PECAM1* and *KDR* are associated with endothelial cell

function and angiogenesis, both crucial for tumor vasculature (51, 52). The downregulation of

PPARG, *ITGAM*, and *CXCL12* further highlights disruptions in immune regulation and cell

migration (53-55). Overall, the downregulation of these hub genes suggests a weakened immune

response, contributing to tumor progression and immune evasion in lung cancer.

Some of the survival associated genes are involved in the several cancers. For example, Li et al.

reported that both the transcriptional and post-transcriptional expression levels of *ADAM12* are

significantly elevated in tumor tissues compared to normal tissues. Furthermore, increased

ADAM12 expression was found to be associated with poor prognosis in patients with lung

adenocarcinoma (56). Ma et al. found that *CCNE1* is upregulated in LUAD and that its

knockdown suppresses tumor growth and metastasis. High *CCNE1* expression correlates with

advanced stage, larger tumors, lymph node involvement, and poor prognosis (57). The

downregulated group, categorized based on median expression levels, is associated with worse

survival outcomes. Lower expression of these genes correlates with poorer prognosis, suggesting

that their downregulation may contribute to tumor progression and aggressive cancer behavior

(58, 59). These findings highlight the potential role of these downregulated genes in worsening

survival outcomes and underscore their relevance as prognostic biomarkers for lung cancer.

These genes, selected through Lasso, represent the most influential biomarkers for survival

prediction, highlighting their potential roles in prognosis and therapeutic targeting.

By integrating survival analysis and regression modeling, we further identified *DSG2, EIF6,* and

EXO1 as independent prognostic factors, establishing a novel hypoxia-related gene signature that

may inform both patient prognosis and therapeutic strategies in LUAD. Previous studies reported

that independent prognostic factors are involved in the several cancer pathogeneses. Gong et al.

reported that elevated *DSG2* expression is significantly associated with poor prognosis and

reduced immune cell infiltration in various cancers (60). Zhu et al. found that overexpression of

eIF6 regulates its downstream target, cell division control protein, contributing to oncogenesis.

They also highlighted the potential of eIF6 as a diagnostic biomarker for cancer (61). Liu et al.

identified *EXO1* as a significant biomarker with both prognostic and diagnostic value across

multiple cancers, highlighting its potential as a therapeutic target and its role in

immunomodulation within the tumor microenvironment (62, 63).These results further support

the robustness of *DSG2, EIF6,* and *EXO1* as key genes in the prediction of survival, making

them critical targets for future therapeutic strategies and prognostic modeling.

While numerous studies have reported associations between hypoxia-related genes and LUAD

prognosis, our study identifies a novel three-gene signature comprising *DSG2, EIF6, and EXO1,*

which has not been previously reported as a prognostic panel. This signature captures distinct

aspects of hypoxia-driven tumor biology: DSG2 regulates cell adhesion and tumor progression

(64), EIF6 controls translation initiation supporting rapid tumor growth under hypoxia (65), and

EXO1 mediates DNA repair under replication stress, which is often exacerbated in hypoxic

tumors (66). Importantly, these genes were validated in an independent cohort and may have

clinical implications, as hypoxia influences the tumor microenvironment and response to

immune checkpoint inhibitors. Thus, the *DSG2-EIF6-EXO1* signature not only provides robust

prognostic stratification but also offers mechanistic and potentially therapeutic insight,

distinguishing it from previously reported hypoxia-related prognostic markers.

Conclusions

In conclusion, our analysis identifies *DSG2, EIF6,* and *EXO1* as key hypoxia-related genes with

significant prognostic value in lung adenocarcinoma. These genes, through their involvement in

hypoxia-related pathways, may play critical roles in tumor progression and patient survival. The

findings suggest that *DSG2, EIF6,* and *EXO1* could serve as potential biomarkers for predicting

patient outcomes and may offer novel therapeutic targets for improving prognosis in lung

adenocarcinoma. Further validation of these genes in larger cohorts and clinical settings is

necessary to confirm their utility in clinical practice.

Acknowledgements

We thank all the members of the scientific community for making their datasets public. This
work was funded by a Wellcome Trust grant to SR (206293/Z/17/Z).

References

1.   Zhang Y, Vaccarella S, Morgan E, Li M, Etxeberria J, Chokunonga E, et al. Global variations in lung
cancer incidence by histological subtype in 2020: a population-based study. Lancet Oncol.
2023;24(11):1206-18.
2.   Seguin L, Durandy M, Feral CC. Lung Adenocarcinoma Tumor Origin: A Guide for Personalized
Medicine. Cancers (Basel). 2022;14(7).
3.   Zhang C, Liu Y, Lu Y, Chen Z, Liu Y, Mao Q, et al. Identification of potential biomarkers for lung
adenocarcinoma: a study based on bioinformatics analysis combined with validation experiments. Front
Oncol. 2024;14:1425895.
4.   Emami Nejad A, Najafgholian S, Rostami A, Sistani A, Shojaeifar S, Esparvarinha M, et al. The role
of hypoxia in the tumor microenvironment and development of cancer stem cell: a novel approach to
developing treatment. Cancer Cell Int. 2021;21(1):62.
5.   Ciepla J, Smolarczyk R. Tumor hypoxia unveiled: insights into microenvironment, detection tools
and emerging therapies. Clin Exp Med. 2024;24(1):235.
6.   Petrova V, Annicchiarico-Petruzzelli M, Melino G, Amelio I. The hypoxic tumour
microenvironment. Oncogenesis. 2018;7(1):10.

7.   Child F, Frost J, Shakir D, Wilson JW, Rocha S. Transcription| regulation of gene transcription by
hypoxia-inducible factor 1α. 2021.
8.   Bae T, Hallis SP, Kwak MK. Hypoxia, oxidative stress, and the interplay of HIFs and NRF2 signaling
in cancer. Exp Mol Med. 2024;56(3):501-14.
9.   Bui BP, Nguyen PL, Lee K, Cho J. Hypoxia-Inducible Factor-1: A Novel Therapeutic Target for the
Management of Cancer, Drug Resistance, and Cancer-Related Pain. Cancers (Basel). 2022;14(24).
10.   Gonzalez-Avila G, Sommer B, Flores-Soto E, Aquino-Galvez A. Hypoxic Effects on Matrix
Metalloproteinases' Expression in the Tumor Microenvironment and Therapeutic Perspectives. Int J Mol
Sci. 2023;24(23).
11.   Roy S, Kumaravel S, Sharma A, Duran CL, Bayless KJ, Chakraborty S. Hypoxic tumor
microenvironment: Implications for cancer therapy. Exp Biol Med (Maywood). 2020;245(13):1073-86.
12.   Chen Z, Han F, Du Y, Shi H, Zhou W. Hypoxic microenvironment in cancer: molecular
mechanisms and therapeutic interventions. Signal Transduct Target Ther. 2023;8(1):70.
13.   Matic N, Pettersson L, Sellebjerg F, Lindberg L, Roberg K, Wiechec E. Prognostic value of hypoxia-
responsive gene expression profile in patients diagnosed with head and neck squamous cell carcinoma.
Transl Oncol. 2024;39:101841.
14.   Li Y, Huang H, Jiang M, Yu N, Ye X, Huang Z, et al. Identification and validation of a hypoxia-
immune signature for overall survival prediction in lung adenocarcinoma. Front Genet. 2022;13:975279.
15.   Rahal Z, El Darzi R, Moghaddam SJ, Cascone T, Kadara H. Tumour and microenvironment
crosstalk in NSCLC progression and response to therapy. Nat Rev Clin Oncol. 2025;22(7):463-82.
16.   Shi R, Sun J, Zhou H, Hu T, Gao Z, Wang X, et al. Hypoxia within tumor microenvironment
characterizes distinct genomic patterns and aids molecular subtyping for guiding individualized
immunotherapy. Journal of Big Data. 2024;11(1):81.
17.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio cancer genomics
portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov.
2012;2(5):401-4.
18.   Huang da W, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the
comprehensive functional analysis of large gene lists. Nucleic Acids Res. 2009;37(1):1-13.
19.   Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B (Methodological).
1995;57(1):289-300.
20.   Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al. STRING v11: protein-
protein association networks with increased coverage, supporting functional discovery in genome-wide
experimental datasets. Nucleic Acids Res. 2019;47(D1):D607-D13.
21.   Chin CH, Chen SH, Wu HH, Ho CW, Ko MT, Lin CY. cytoHubba: identifying hub objects and sub-
networks from complex interactome. BMC Syst Biol. 2014;8 Suppl 4(Suppl 4):S11.
22.   Therneau T. A package for survival analysis in R. (95).
23.   Friedman JH, Hastie T, Tibshirani R. Regularization Paths for Generalized Linear Models via
Coordinate Descent. Journal of Statistical Software. 2010;33(1):1 - 22.
24.   Aguirre-Gamboa R, Gomez-Rueda H, Martinez-Ledesma E, Martinez-Torteya A, Chacolla-
Huaringa R, Rodriguez-Barrientos A, et al. SurvExpress: an online biomarker validation tool and database
for cancer gene expression data using survival analysis. PLoS One. 2013;8(9):e74250.
25.   Zhao Y, Xing C, Deng Y, Ye C, Peng H. HIF-1alpha signaling: Essential roles in tumorigenesis and
implications in targeted therapies. Genes Dis. 2024;11(1):234-51.
26.   Barnum KJ, O'Connell MJ. Cell cycle regulation by checkpoints. Methods Mol Biol. 2014;1170:29-
40.
27.   Pedroza-Garcia JA, Xiang Y, De Veylder L. Cell cycle checkpoint control in response to DNA
damage by environmental stresses. Plant J. 2022;109(3):490-507.

28.   Rankin BD, Rankin S. The MCM2-7 Complex: Roles beyond DNA Unwinding. Biology (Basel).
2024;13(4).
29.   Bertoli C, Herlihy AE, Pennycook BR, Kriston-Vizi J, de Bruin RAM. Sustained E2F-Dependent
Transcription Is a Key Mechanism to Prevent Replication-Stress-Induced DNA Damage. Cell Rep.
2016;15(7):1412-22.
30.   Zhang Y, Wen GM, Wu CA, Jing ZL, Li DZ, Liu GL, et al. PRKDC is a prognostic marker for poor
survival in gastric cancer patients and regulates DNA damage response. Pathol Res Pract.
2019;215(8):152509.
31.   Klomp JE, Lee YS, Goodwin CM, Papke B, Klomp JA, Waters AM, et al. CHK1 protects oncogenic
KRAS-expressing cells from DNA damage and is a target for pancreatic cancer treatment. Cell Rep.
2021;37(9):110060.
32.   Ooi AT, Gomperts BN. Molecular Pathways: Targeting Cellular Energy Metabolism in Cancer via
Inhibition of SLC2A1 and LDHA. Clin Cancer Res. 2015;21(11):2440-4.
33.   Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. Cell. 2011;144(5):646-74.
34.   Chen DS, Mellman I. Oncology meets immunology: the cancer-immunity cycle. Immunity.
2013;39(1):1-10.
35.   Schreiber RD, Old LJ, Smyth MJ. Cancer immunoediting: integrating immunity's roles in cancer
suppression and promotion. Science. 2011;331(6024):1565-70.
36.   Herbst RS, Morgensztern D, Boshoff C. The biology and management of non-small cell lung
cancer. Nature. 2018;553(7689):446-54.
37.   Bandarra D, Biddlestone J, Mudie S, Muller HA, Rocha S. HIF-1alpha restricts NF-kappaB-
dependent gene expression to control innate immunity signals. Dis Model Mech. 2015;8(2):169-81.
38.   Pang X, Xie R, Zhang Z, Liu Q, Wu S, Cui Y. Identification of SPP1 as an Extracellular Matrix
Signature for Metastatic Castration-Resistant Prostate Cancer. Front Oncol. 2019;9:924.
39.   Napoli S, Scuderi C, Gattuso G, Bella VD, Candido S, Basile MS, et al. Functional Roles of Matrix
Metalloproteinases and Their Inhibitors in Melanoma. Cells. 2020;9(5).
40.   Bhattacharyya S, Mote RD, Freimer JW, Tiwari M, Bansi Singh S, Arumugam S, et al. Cell-cell
adhesions in embryonic stem cells regulate the stability and transcriptional activity of beta-catenin. FEBS
Lett. 2022;596(13):1647-60.
41.   French R, Pauklin S. Epigenetic regulation of cancer stem cell formation and maintenance. Int J
Cancer. 2021;148(12):2884-97.
42.   Mirzaei S, Paskeh MDA, Entezari M, Mirmazloomi SR, Hassanpoor A, Aboutalebi M, et al. SOX2
function in cancers: Association with growth, invasion, stemness and therapy response. Biomed
Pharmacother. 2022;156:113860.
43.   Baker SJ, Poulikakos PI, Irie HY, Parekh S, Reddy EP. CDK4: a master regulator of the cell cycle
and its role in cancer. Genes Cancer. 2022;13:21-45.
44.   Choi SA, Moon YJ, Koh EJ, Phi JH, Lee JY, Kim KH, et al. Cyclin-Dependent Kinase Inhibitor 2A is a
Key Regulator of Cell Cycle Arrest and Senescence in Endothelial Colony-Forming Cells in Moyamoya
Disease. J Korean Neurosurg Soc. 2023;66(6):642-51.
45.   Yarden RI, Brody LC. BRCA1 interacts with components of the histone deacetylase complex. Proc
Natl Acad Sci U S A. 1999;96(9):4983-8.
46.   Bhaskara S. Histone deacetylases 1 and 2 regulate DNA replication and DNA repair: potential
targets for genome stability-mechanism-based therapeutics for a subset of cancers. Cell Cycle.
2015;14(12):1779-85.
47.   Allen JR, Ge L, Huang Y, Brauer R, Parimon T, Cassel SL, et al. TIMP-1 Promotes the Immune
Response in Influenza-Induced Acute Lung Injury. Lung. 2018;196(6):737-43.
48.   Elemam NM, Talaat IM, Maghazachi AA. CXCL10 Chemokine: A Critical Player in RNA and DNA
Viral Infections. Viruses. 2022;14(11).

49. Schultheiss C, Willscher E, Paschold L, Gottschick C, Klee B, Henkes SS, et al. The IL-1beta, IL-6,
and TNF cytokine triad is associated with post-acute sequelae of COVID-19. Cell Rep Med.
2022;3(6):100663.
50. Zheng M, Ambesi A, McKeown-Longo PJ. Role of TLR4 Receptor Complex in the Regulation of the
Innate Immune Response by Fibronectin. Cells. 2020;9(1).
51. Guerra-Espinosa C, Jimenez-Fernandez M, Sanchez-Madrid F, Serrador JM. ICAMs in Immunity,
Intercellular Adhesion and Communication. Cells. 2024;13(4).
52. del Puerto-Nevado L, Rojo F, Zazo S, Carames C, Rubio G, Vega R, et al. Active angiogenesis in
metastatic renal cell carcinoma predicts clinical benefit to sunitinib-based therapy. Br J Cancer.
2014;110(11):2700-7.
53. Angeli V, Hammad H, Staels B, Capron M, Lambrecht BN, Trottein F. Peroxisome proliferator-
activated receptor gamma inhibits the migration of dendritic cells: consequences for the immune
response. J Immunol. 2003;170(10):5295-301.
54. Guyon A. CXCL12 chemokine and its receptors as major players in the interactions between
immune and nervous systems. Front Cell Neurosci. 2014;8:65.
55. Huang H, Wang Q, Ma L, Wu Y. ITGAM: A Pivotal Regulator in Macrophage Dynamics and
Cardiac Function During Sepsis-Induced Cardiomyopathy. Cureus. 2024;16(4):e59342.
56. Li K, Quan L, Huang F, Li Y, Shen Z. ADAM12 promotes the resistance of lung adenocarcinoma
cells to EGFR-TKI and regulates the immune microenvironment by activating PI3K/Akt/mTOR and RAS
signaling pathways. Int Immunopharmacol. 2023;122:110580.
57. Ma G, Yang L, Dong J, Zhang L. CCNE1 Promotes Progression and is Associated with Poor
Prognosis in Lung Adenocarcinoma. Curr Pharm Biotechnol. 2022;23(9):1168-78.
58. Cizkova M, Vacher S, Meseure D, Trassard M, Susini A, Mlcuchova D, et al. PIK3R1
underexpression is an independent prognostic marker in breast cancer. BMC Cancer. 2013;13:545.
59. Wei X, Chen L, Yang A, Lv Z, Xiong M, Shan C. ADRB2 is a potential protective gene in breast
cancer by regulating tumor immune microenvironment. Transl Cancer Res. 2021;10(12):5280-94.
60. Zhang G, Chen Z, Wang Y, Huang A, Nie F, Gao L, et al. Up-regulated DSG2 promotes tumor
growth and reduces immune infiltration in cervical cancer. Pathol Res Pract. 2024;262:155554.
61. Zhu W, Li GX, Chen HL, Liu XY. The role of eukaryotic translation initiation factor 6 in tumors.
Oncol Lett. 2017;14(1):3-9.
62. Yang G, Dong K, Zhang Z, Zhang E, Liang B, Chen X, et al. EXO1 Plays a Carcinogenic Role in
Hepatocellular Carcinoma and is related to the regulation of FOXP3. J Cancer. 2020;11(16):4917-32.
63. Liu Z, Huang S, Luo R, Shi X, Xiu M, Wang Y, et al. EXO1's pan-cancer roles: diagnostic,
prognostic, and immunological analyses through bioinformatics. Discov Oncol. 2025;16(1):310.
64. Myo Min KK, Ffrench CB, McClure BJ, Ortiz M, Dorward EL, Samuel MS, et al. Desmoglein-2 as a
cancer modulator: friend or foe? Front Oncol. 2023;13:1327478.
65. Sun L, Liu S, Wang X, Zheng X, Chen Y, Shen H. eIF6 promotes the malignant progression of
human hepatocellular carcinoma via the mTOR signaling pathway. J Transl Med. 2021;19(1):216.
66. Garcia-Rodriguez N, Dominguez-Garcia I, Dominguez-Perez MDC, Huertas P. EXO1 and DNA2-
mediated ssDNA gap expansion is essential for ATR activation and to maintain viability in BRCA1-
deficient cells. Nucleic Acids Res. 2024;52(11):6376-91.

Supplementary Information

Supplementary Tables S1-S9

Supplementary Table S1. List of collected hypoxia related genes.

Supplementary Table S2. List of DEGs in TCGA LUAD.

Supplementary Table S3. List of DEGs in GSE18842.

Supplementary Table S4. List of hypoxia related DEGs in the both datasets.

Supplementary Table S5. Enriched pathways associated with the upregulated genes (FDR <
0.05).

Supplementary Table S6. Enriched pathways associated with the downregulated genes (FDR <
0.05).

Supplementary Table S7. Rank of the PPI network by Degree method.

Supplementary Table S8. Significant Genes Identified in Kaplan-Meier Survival Analysis.

Supplementary Table S9. Significant Genes Identified in Univariate regression Analysis.