Bending friction: a new mechanism of dissipation within DNA explains its slow looping dynamics.

Georgii Pobegalov and Maxim Molodtsov[*]

The Francis Crick Institute, London, United Kingdom

Dept of Physics & Astronomy, University College London, United Kingdom

Bhavin S. Khatri[†]

The Francis Crick Institute, London, United Kingdom and

Department of Life Sciences, Imperial College London, United Kingdom

Abstract

DNA bending and looping is crucial for gene expression, packaging, and chromatin organisation, as well as the design of artificial nanomaterials and devices. But what determines how quickly DNA bends? While DNA's static flexibility is well-characterised by its persistence length, we lack an understanding of how quickly DNA responds to mechanical forces: remarkably current semiflexible polymer theory based on solvent dissipation underestimates spontaneous looping times by $\sim$1000-fold. By analysing fluctuations of DNA several kilobases long and developing new theory for bending dissipation in semiflexible polymers, we show DNA bending dynamics cannot be explained by solvent friction alone and requires significant contributions from intramolecular friction. The theory defines a new material constant of DNA — the bending friction, which we determine to be $\zeta_B = 241 \pm 17$ μg nm^3/ms. Strikingly, our measurement does not depend on the buffer ionic conditions. We predict bending friction will dominate DNA dynamics between $\approx$ 50 nm and 420 nm and significantly longer under external force. We show that mean first passage time calculations are greatly simplified when bending friction dominates and so using this constant, with no fitting parameters, we accurately predict the slow experimental spontaneous looping times. Our discovery of significant bending dissipation is unexpected as DNA has no obvious large ($> k_B T$) internal energy barriers. The salt-independence of this dissipation also rules out long range electrostatic interactions as its origins. Instead our findings point to a complex local energy landscape for bending and a potential previously unappreciated role of water binding DNA constraining its local mobility. Our findings radically change our understanding of DNA dynamics and reveal DNA as a viscoelastic semiflexible polymer with dramatically slower dynamics compared to an ideal elastic rod. This work establishes bending friction as a fundamental material property that must underpin any model of DNA dynamics in biology, physics, and nanotechnology.

I. INTRODUCTION

DNA is one of biology's most important polymers, storing the instructions for life. Beyond information storage, its mechanical properties play a fundamental role in numerous processes in molecular biology, from transcription initiation [7, 41] and chromatin packing around nucleosomes [45], to genome organisation in eukaryotes [24, 37, 50]. These processes often involve molecular motors that manipulate, bend and loop DNA, where there is evidence that the local bending properties of DNA may be coded through sequence in the genome[3]. The spontaneous bending and looping of DNA is also a crucial step in high throughout sequencing through the process of bridge amplification [5], which forms the basis of the recent advances in genomics and its industry and the emergence of DNA as a means high volume data storage [2]. In addition, our ability to artificially construct DNA nanostructures and develop bio-inspired nanomachines [21, 25, 47] depends crucially on understanding its mechanical properties.

While DNA's bending elasticity determines its equilibrium shape under mechanical constraints or forces, we lack a comparable understanding of an equally important mechanical property: how rapidly DNA changes shape under external forces or thermal fluctuations. The prevailing framework for DNA dynamics, semiflexible polymer theory — that assumes dissipation that limits dynamics arises solely from Stokes friction with the solvent[27, 36] — severely underestimates spontaneous loop-closure times for short DNA ($\sim$100nm contour length or 300bp) by a remarkable 1000-fold [32, 51]. Therefore, we lack a theoretical framework to explain these slow looping dynamics and so are unable to answer the simple question: what determines how quickly DNA bends.

However, Kuhn's theorem has long predicted that for short flexible chains, intramolecular friction — arising from internal energy barriers — must dominate

* Corresponding author: m.molodtsov@ucl.ac.uk

† Corresponding author: bkhatri@imperial.ac.uk

polymer dissipation[9, 13, 28, 31, 38, 39]. de Gennes illustrated this with a striking thought experiment: if thermal energy $k_B T$ is much smaller than internal energy barriers $\Delta U^{\ddagger}$, a single polymer chain would enter a frozen, glassy state[13]. Lacking the obvious internal rotation barriers of flexible polymers, DNA is often treated as a simple elastic rod. Yet, its complex molecular structure creates a potentially intricate energy landscape for bending, shaped by atomic interactions within DNA and with surrounding water (Fig.1). Whether these processes within DNA are sufficient to explain empirically measured loop-closure times is unknown.

By analysing DNA fluctuations across a range of lengths and tensions, and developing a new first-principles theory for bending dissipation in semiflexible polymers under tension, we reveal clear signatures of intramolecular friction that rule out a purely solvent-based origin. What emerges is a new material constant of DNA — the bending friction — describing the local dissipation within DNA itself as it bends. Strikingly, we find this constant is independent of the buffer's ionic strength, which suggests its origin lies in local energetic interactions within DNA and potentially water binding DNA, which would be a new unappreciated role for water modulating its bending dynamics. Using our measurement of this constant, we show that we can fit measured loop-closure times with no fitting parameters. This demonstrates that DNA's slow looping dynamics is explained by intramolecular bending friction and radically changes our picture of DNA at these length scales: from an ideal elastic rod to a viscoelastic semiflexible polymer with dramatically slower dynamics.

FIG. 1. **The origins of bending friction within DNA.** a) an ideal elastic rod with no internal structure has a smooth quadratic elastic energy $U(\theta)$ as function of bend angle θ, where the curvature is controlled by the bending elasticity. Rod friction ζ_s arises from viscosity with solvent, determined by the timescale τ of water hydrogen bond lifetimes. Grey dot moving to black dot over time indicates a change in bend angle with red arrow indicating retarding frictional force. b) DNA segment: unlike an idealised elastic rod has a complex internal molecular structure. As DNA bends, complex changes in the relative positions of all the atoms, gives a corresponding complex change in the energy of DNA due to numerous interactions (red dotted lines). In addition, specific binding of water molecules to nucleotides, will necessitate unbinding and binding of water as DNA bends. These factors could give rise to rough energy landscape (red) with local energy barriers, where $\Delta U^{\ddagger}/k_B T$ is the roughness of the energy landscape — on the scale of thermal energy $k_B T$ — to DNA bending due to local interactions in DNA, and ζ_B is the emergent bending friction of DNA due to local barrier hopping on timescale $\tau^{\ddagger}$.

II. EXTRACTING DNA FRICTION FROM SINGLE-MOLECULE FLUCTUATIONS

To determine the role of bending friction in DNA dynamics, single DNA molecules were tethered between two beads each in optical traps and stretched to various mean forces F (1pN$\leq F \leq$ 5pN) by appropriately choosing the distance between the trap centres Fig.2a. This range ensures DNA is highly stretched ($F \gg k_B T / \ell_P \approx 0.08$ pN) while avoiding backbone extension. At each force, we recorded the positions of both beads, $x_1(t)$ and $x_2(t)$, at 78 kHz for 6 seconds. DNA length fluctuations are calculated as $\ell(t) = x_2(t) - x_1(t) + X - D$, where X is the separation between trap centres and D the diameter of the beads. Example time series of $\ell(t)$ are shown in Fig.2b as a function of increasing force. To specifically isolate dissipative dynamics, we calculated the velocity power spectral density (VPSD, $P_v(\omega)$), derived from the length fluctuations as $P_v = \omega^2 P_\ell$ (Fig.2c), where $P_\ell(\omega)$ is the regular power spectrum. This transformation weights higher frequencies, providing a direct measure of the power dissipated unit frequency — which increases at higher frequencies — and thus a more reliable signature of DNA friction (Fig.S1).

The velocity power spectrum is a function of two main force-dependent viscoelastic parameters of DNA: the end-to-end elasticity $\kappa_{DNA}(F)$ — which is the well-characterised entropic elasticity often described by a Worm-like chain (WLC) model [8, 35] — and the end-to-end friction of DNA $\zeta_{DNA}(F)$. Our data in Fig.2c shows that the velocity power spectrum is highly sensitive to changes in the average tension applied to DNA. As we demonstrate in Fig.S2 the maximum value of the VPSD is a direct read-out of the inverse of the total friction constant of the system. Therefore, even before fitting any model to the VPSD, we can see that the friction of DNA is increasing with increasing tension, which as we argue below cannot be explained by solvent friction. Fitting to the VPSD at each force in

FIG. 2. **Measuring the friction response of DNA as a function of force.** a) DNA is tethered between two beads at a mean force F using dual optical trap. Time-series of the position of each trap, $x_1(t)$ and $x_2(t)$, is recorded, where total friction of system is given by $\zeta = \zeta_{DNA} + \frac{1}{2}(\zeta_0 + \zeta_{12})$ (Supplementary Information:Eqn.S17), where ζ_{DNA} is the friction of DNA, ζ_0 friction of each bead and ζ_{12} the friction due to the hydrodynamic interactions between the beads. b) Example time series of length $\ell(t)$ for 2.7kbp (903 nm) DNA as a function of force, showing increased extension for increasing forces. c) the velocity power spectrum (VPSD) of the fluctuations in b) : decrease in peak height with increasing tension can only be explained by increasing friction of DNA (Supplementary Information: section S3). d) Total effective friction of the trap DNA system (red pentagrams) as a function of force. DNA friction increases above background hydrodynamic friction (grey hexagrams — measured without DNA — and grey solid line, fit with hydrodynamic theory between two beads). This contrasts to expectation if solvent friction were only source of dissipation (pink dashed line: $\zeta_s = \frac{2\pi\eta L}{\ln(L/w)}$ [16]; $L = 902.6$ nm, $w = 2.4$ nm[34]).

Fig.2c, using a model of fluctuations which includes appropriates high frequency corrections to Brownian motion (Supplementary Information: Eqn.S18), we determine the combined friction constant of the bead, trap and DNA, $\zeta(F)$, as a function of force shown in Fig.2d, as solid squares.

Our data show that the friction constant with DNA tethered between beads increases approximately linearly with increasing force over and above the back-

ground bead friction (grey hexagrams with solid grey line fit to Oseen hydrody-namic theory [17] (Supplementary Information Eqn.S11)). This increase cannot be attributed to the solvent friction of DNA: at high stretch ($F \gg k_B T / \ell_P \approx 0.08$pN) solvent dissipation gives an approximately constant friction of $\zeta_s \approx 0.00092 \mu$g/ms for 2.7kb long DNA (dashed pink line in Fig.2d, where we assume DNA is a rod of length $L \approx 900$nm, and width $w = 2.4$ nm[34]). Thus, contrary to the predictions based on the solvent friction alone, our data in Fig.2d show that friction increases with force and reaches approximately 0.0035μg/ms, above the background hydro-dynamic friction, at $F = 5$ pN, which is almost four-times the expected solvent friction. In the next section, we develop a theory of bending dissipation in DNA to explain these observations.

Our fitting also yields the elastic constant $\kappa(F)$, which matches the entropic elasticity expected of a WLC at high stretch (Fig.S3). The fit of the VPSD data also accounts for the effective frequency-dependent mass of water entrained around each bead, which should be independent of force, which we verify in Fig.S4.

III. FORCE-DEPENDENT FRICTION IS EXPLAINED BY BENDING DISSIPATION WITHIN DNA

Because solvent friction alone could not explain the linear increase in friction of DNA with force, we sought to rationalise our findings by modifying the standard model used to describe DNA dynamics when highly stretched at a tension F ($F \gg k_B T / \ell_P \approx 0.08$pN). Semiflexible polymer dynamics describes DNA as an idealised elastic rod with space curve $\boldsymbol{R}(s, t)$ — where s is the backbone co-ordinate of the polymer — whose shape fluctuates dynamically due to Brownian motion and friction with the solvent. We propose that due to internal frictional processes within DNA there will be local bending dissipation proportional to the rate of change of local curvature, described by the Rayleigh dissipation function $\mathcal{D} =$

116 $\frac{1}{2}\zeta_B(\partial_t\partial_s^2\boldsymbol{R})^2$, where ζ_B, is the bending friction constant[23, 43]. Calculating the

117 local forces due to this dissipation gives rise to a modified equation for semiflexible

118 polymer dynamics under tension, which we call the Dissipative Worm-Like Chain

119 (DWLC):

$$\tilde{\zeta}_s\frac{\partial\boldsymbol{R}(s,t)}{\partial t} + \zeta_B\frac{\partial}{\partial t}\frac{\partial^4\boldsymbol{R}(s,t)}{\partial s^4} - F\frac{\partial^2\boldsymbol{R}(s,t)}{\partial s^2} + \kappa_B\frac{\partial^4\boldsymbol{R}(s,t)}{\partial s^4} = \boldsymbol{f}(s,t), \qquad (1)$$

where $\tilde{\zeta}_s$ is a solvent friction per unit length, and $\kappa_B = k_B T \ell_p$ is the bending elastic

constant and $\boldsymbol{f}(s,t)$ is a temporally white, but spatially coloured, noise term as is

necessary for a continuum description of internal friction[29] (see Supplementary

Information section S2 for details).

Our aim is to evaluate the additional, or excess, end-to-end friction of the

chain, as a function of tension F, due to bending dissipation, over and above

hydrodynamic solvent contributions or Stokes' friction of DNA. We do this by

analysing Eqn.1 using standard normal mode analysis to give the mode relaxation

time for the n^{th} mode:

$$\tau_n = \frac{\tilde{\zeta}_s + \zeta_B q_n^4}{\kappa_B q_n^4 + F q_n^2}. \qquad (2)$$

129 where q_n is its wavenumber. We then evaluate the relevant 4^{th} order moments

130 using Wick's theorem[17] to calculate the autocorrelation function of the end-to-

131 end distance ΔR of the chain:

$$\rho_{\Delta R}(t) = \langle\Delta R(t)\Delta R(0)\rangle - \langle\Delta R(t)\rangle\langle\Delta R(0)\rangle = (Lk_BT)^2\sum_{n=0}^{\infty}\frac{q_n^4}{\kappa_n^2}e^{-\frac{2t}{\tau_n}}, \qquad (3)$$

132 where mode elasticity is given by $\kappa_n = (\kappa_B q_n^4 + F q_n^2)L$ and mode friction by

133 $\zeta_n = (\tilde{\zeta}_s + \zeta_B q_n^4)L$. The end-to-end excess friction is then calculated from the

134 derivative of the autocorrelation function [18]:

$$\frac{k_B T}{\zeta_{\Delta R}} = \left| \frac{\mathrm{d}}{\mathrm{d}t} \rho_{\Delta R}(t) \right|_{t=0} = 2(L k_B T)^2 \sum_{n=0}^{\infty} \frac{q_n^4}{\kappa_n \zeta_n}. \tag{4}$$

135 We then take the continuum limit with $q_n \to q$, with mode spacing $\delta q = \pi / L$
136 to convert the summation to an integral to give the following, valid for forces
137 $F > 0.08 \mathrm{pN}$, in the limit of the chain at high force/stretch (full derivation given
138 in Supplementary Information section S2):

$$\Delta \zeta_{DNA}(F) = \Delta \zeta_{\Delta R}(F) = \frac{\sqrt{2} \tilde{\zeta}_s^{1/4} \zeta_B^{3/4}}{k_B T L} F. \tag{5}$$

This shows that explicit inclusion of bending friction in the equations of motion
leads to the observed linear increase of the friction coefficient with external force
(Fig.2d). Note that the friction of the dissipative WLC is inversely proportional to
the contour length L, which is a classic signature of internal dissipation in polymers
[13, 29]. Thus, DNA friction that takes into account internal bending dissipation
increases with force, and it does so more significantly for shorter DNA.

IV. A NEW MATERIAL CONSTANT OF DNA

Having established the theoretical description of bending dissipation using the
DWLC model, and parameterised by the bending friction constant, we can now
infer its value from the experimental data. Since bending dissipation in DNA
gives rise to a friction inversely proportional to contour length, we measured a
number of single molecules of DNA for lengths between 2.7kbp and 8.8kbp ($\approx$
900nm to 3000nm). To extract the DNA friction from the measurements, we
determined the excess DNA friction, $\Delta \zeta_{DNA}(F)$, as the residual friction above the
background friction, due to the hydrodynamic interactions between beads (grey
pentagrams in Fig.2d) and the hydrodynamic (Stokes) solvent friction of DNA

with a fitting parameter to account for the complex hydrodynamic interactions between the beads and DNA (Fig.S6). The data is then fit using Eqn.5, which has a single fitting parameter ζ_B.

The resulting measurements of the excess friction for all DNA lengths (various colours), with the effective background bead and DNA solvent friction removed are shown in Fig.3. The fitting of Eqn.5 to the experimental data shows that our DWLC model accounts for the main unexpected feature of the data: a linear increase of the excess DNA friction with increasing force. In addition, Fig.3 shows that as the contour length decreases, the excess friction increases, which is in qualitative agreement with Eqn.5. In Fig.S7, we additionally plot the normalised friction $L\Delta\zeta_{DNA}$ as a function of force, which shows excellent quantitative agreement with $\Delta\zeta_{DNA} \sim \frac{1}{L}$. This is in clear contrast to solvent DNA friction, which would increase approximately linearly with increasing length (Fig.S5). Thus, the experimentally observed excess friction is fully explained by the bending friction of DNA.

Furthermore, to test whether DNA bending friction depends on long-range electrostatic interactions within DNA, we compared experiments performed at physiological salt concentrations (in PBS buffer) with experiments carried out under very low salt conditions (1 mM NaCl, in 1mM Hepes buffer; see Methods). This comparison (closed and open data circles, respectively, in Fig.3) shows no dependence of the bending friction on the salt concentration (Fig.S7). As a control, for each molecule in these experiments, we also performed force-extension measurements and extracted their persistence lengths. As expected and in quantitative agreement with previous studies[4, 10, 22, 52], we observed an increase in the persistence length for the low salt buffer (Supplementary Information: Fig.S12).

Our experiments show that bending friction is a material constant of DNA independent of the buffer conditions. This suggests that the origin of the excess DNA friction we measure is due to local interactions within DNA, rather than

183 long-range electrostatic interactions. Therefore, we combine fits to data across
184 both buffer conditions, accounting for finite-size effects [48] (Fig.S7), to give an
185 empirical value of $\zeta_B = 241 \pm 17\mu$g nm^3/ms (s.e.m).

FIG. 3. **The end-to-end friction of DNA.** The measured excess end-to-end friction of DNA as a function of force showing linearity and increasing friction for decreasing contour length as predicted by Eqn.5. Closed circles are in normal PBS buffer and open circles are in the low salt Hepes buffer. The data is fit using Eqn.5 with single fitting parameter ζ_B, with $\tilde{\zeta}_s = 2\pi\eta/\ln\left(L/w\right)$ with fits shown with solid lines (PBS) or dashed lines (Hepes). The number of measurements on independent single molecules at each length and for each buffer condition are given in Methods (Table.II). The shaded region indicate the forces for which we would expect the linear dependence to be invalid (i.e. when at low stretch when $F \ll k_B T/\ell_P \approx 0.08$ pN); in fact as $F \to 0$, we would expect the excess friction due to bending dissipation to tend to a constant. These measurements cannot be due to solvent dissipation, as we would expect solvent friction to be approximately independent of force at high stretch and increase with increasing contour length of DNA (Fig.S5).

 V.   THE DYNAMICS OF DNA LOOPING ON SMALL LENGTH SCALES

FIG. 4. **DNA dynamics on small length scales.** a) Relative contribution of bending and solvent friction to total mode friction for a given wavelength. b) Plot of mean first passage times (MFPT) for DNA loop closure as a function of contour length. The predictions based on the theory developed in this paper for circular loops (Eqn.S3) and minimum energy tear-drop loops (Supplementary Information: section S5), where bending friction and bending energy dominate, are shown as dashed blue line and solid blue line, respectively. While the prediction based on a WLC with only solvent dissipation [27] is shown by the dashed pink line. Experimental measurements of MFPT using cyclisation assays are shown by open circles[32] and open pentagrams [51].

We have determined a new fundamental material constant for DNA, ζ_B which
determines how DNA locally dissipates energy as it bends and loops. With our
measurement of ζ_B we can now quantify the relative contributions of solvent and
bending friction for any DNA length at zero force. The normal mode analysis
(Eqn.2) shows that the critical length scale at which solvent and bending contri-
butions to the friction are equal is $L^* \approx 750$nm (Fig.4a) (Supplementary Informa-
tion: section S4), which is equivalent to a DNA length of approximately 2.2kbp.
This figure also shows that for DNA contour lengths $L \leq 420$nm, solvent friction

is 10× smaller than bending friction; in other words we can model the sponta-

neous dynamics of DNA, considering only the contributions from bending friction,

for contour lengths up to about $9\ell_p$, or 1.2kbp. For very small contour lengths

less than persistence length, DNA kinking likely plays a significant role in DNA

flexibility and dynamics[53]. Therefore, we expect bending friction dominates for

DNA lengths $\ell_P \leq L \leq 9\ell_P$ and have significant contributions up to $15\ell_P \to 20\ell_P$

(750nm $\to$1μm, or 2.2kbp $\to$ 3kbp) at zero force. As we show below, the stochas-

tic dynamics of DNA is greatly simplified in the regime where bending friction

dominates.

From this zero force limit, as tension increases and the entropy of the chain

decreases, we expect in general that the bending friction contribution will increase,

and therefore be more important at even longer length scales. Eqn.5, demonstrates

this explicitly once the chain is at high stretch ($F \gg k_B T/\ell_P \approx 0.08$ pN). In cell

nuclei DNA can be locally subjected to forces up to 20 pN [42, 44], at which

we calculate bending friction will dominate up to $\approx 1.2\mu$m and with significant

contributions up to $\approx 12\mu$m (Fig.S8). At lower forces and stretch, we expect

the bending friction contribution to decrease monotonically to the zero force limit

described by the mode analysis above.

When bending friction dominates the dynamics, we can calculate a new useful

quantity — the persistence time of DNA — which is the average time it takes

for DNA to completely change shape due to random thermal fluctuations. In the

absence of external force and in the limit that bending friction dominates, this is

simply given by the mode relaxation time, since all modes relax with the same

characteristic time τ_n independent of mode number n:

$$\tau_P = \frac{\zeta_B}{\kappa_B}. \tag{6}$$

219 Given the value of ζ_B that we have determined, and assuming $\kappa_B = k_B T \ell_P \approx$

220 200pNnm^2 this gives $\tau_P = 1.2$ms. Thus, DNA segments of lengths 750nm (1.2kbp)

or shorter completely change their shape under thermal fluctuations in approximately 1.2 ms. Furthermore, as we will see below, the persistence time of DNA is directly proportional to the time it takes DNA to spontaneously make a loop.

A. Comparison to experimentally measured DNA looping times

Cyclisation assays have become the standard way to estimate the rate of loop closure of DNA [32, 51], the characterisation of which is important to understanding the dynamics of number of DNA looping processes in molecular biology. Current theory to calculate mean first passage times are based on semiflexible polymer dynamics without bending friction [27, 36] and are consequentially complex calculations as they require consideration of the different time scales of each of the modes contributing to the polymer dynamics. However, when bending friction dominates solvent friction and in the limit that bending energy dominates the chain entropy, we can make the assumption that the DNA conformation is roughly an arc of a circle with curvature $\Gamma = 1/R$, where R is the radius of this circle. The resulting stochastic dynamics is then given by the standard Smoluchowski equation for a single degree of freedom, the curvature Γ, with effective friction $\zeta_\Gamma = L\zeta_B$ and bending energy $U(\Gamma) = \frac{L}{2}\kappa_B\Gamma^2$. Using the standard flux-over population method and Kramer's approximations (see Supplementary Information— section S5 — for details) we find the mean first passage time (MFPT) τ^* for loop closure is:

$$\tau^* = \frac{\zeta_B}{\kappa_B}\sqrt{\frac{L}{2\pi\ell_p}}e^{\frac{2\pi^2\ell_p}{L}},\tag{7}$$

which is a pleasingly simple closed-form expression, compared to the theory of Jun et. al [27] for a WLC where only solvent dissipation is considered. The ratio of bending friction to bending elasticity is simply the persistence time of DNA (Eqn.6), which means the loop closure time is proportional to the persistence time of DNA conformations.

However, experiments measuring the time for loop closure of DNA [26], suggest that it is most likely to adopt — at least initially — a tear-drop conformation, where the ends come together at a non-zero angle θ between the end tangent vectors (Fig.4b) . Previous theoretical work, shows the minimum energy conformation corresponds to $\theta \approx 100°$ [54] and an overall energy which is roughly 70% the energy of a circular loop [1, 54]. In Supplementary Information (section S5), we estimate the time for closure of a tear-drop loop, by solving the Euler-Lagrange equations for a rod with free ends to calculate its curvature as a function of backbone coordinate. We then make the assumption that the MFPT will be dominated by a portion of the chain around its midpoint where the curvature is highest, which are indicated by the red circles in Fig.4b.

In Fig.4b, we compare the circular and tear-drop predictions of loop-closure times — based on bending dissipation with bending friction constant we measured experimentally — against theory assuming only solvent friction (Jun et al. [27]) and experimentally available looping times from Vafabakhsh & Ha [51] and Le & Kim [32], for DNA lengths between $60 - 70$nm. Our theory agrees with the data, which lies between the lower bound set by the tear-drop loop closure theory and upper bound by the circular loop closure theory; in contrast, we see that the theory with only solvent friction massively underestimates the looping time by approximately 1,000-fold. This is a remarkable agreement between experimental looping times and the prediction of bending dissipation within DNA given no fitting parameters.

VI.   DISCUSSION

A key question remains: what is the underlying origin of bending friction within DNA? Our finding that the bending friction constant is insensitive to salt concentration, and the strength of electrostatic screening, suggests long-range electro-

statics, such as repulsion from the negatively charged backbone, plays a weak role. We instead attribute the origin to local complex energy barriers from the complex atomic interactions within each base pair (Fig.1) as well as potentially complex interactions with water. The latter contributions may come from water binding in the minor groove [30, 46] or specific hydrogen bonds, such as adenine–thymine bridges predicted to stabilize DNA by ≈ 3 $k_B T$ [33]. This previously unanticipated role of water in bending friction would carry important implications for the proper choice of water models to obtain realistic explicit solvent molecular dynamics simulations of DNA, as well as implicit solvent and coarse-grained simulations of DNA, such as oxDNA[40].

Our work identifies a new material constant for local bending dissipation within DNA which we have demonstrated dominates looping and bending dynamics from 50nm to 420nm (150 bp to 1.2 kbp), with significant contributions up to $\approx 1 \mu$m (3 kbp) and potentially an order of magnitude longer under physiological tensions of ~ 20pN found in the cell[42, 44].This has broad implications for the trade-off between the maximum speed versus energy demands of biological processes where proteins, enzymes, or motors manipulate, bend or loop DNA. Examples at these length scales in biology include gene expression[11, 19, 20] and loop initiation and extrusion in replisomes, SMCs or telomerase complexes[12, 14]. Our calculations in Fig.4b show that for DNA between 50nm and 420nm loop-closure times vary by many orders of magnitude, ranging from roughly 3 hours to 10 ms. This suggests spontaneous loop formation for small length DNAs is likely inefficient and therefore requires significant energy input to overcome bending dissipation, whereas for longer DNA, looping may occur spontaneously. These same trade-offs will play a role in the design, construction and optimisation of nanomaterials and devices[47] — for example strand-displacement artificial DNA motors[49, 55] — and particularly in the engineering of bio-inspired systems that mimic the non-equilibrium nature of molecular biology[21], where dynamics will be a key consideration.

Overall our findings show that DNA — despite lacking any obvious large internal energy barriers — nonetheless has significant intramolecular dissipative forces described by a new physical principle of bending dissipation in semiflexible polymers, which explain the puzzle of its very slow loop-closure times. This sets-forth a new paradigm for DNA dynamics — the dissipative worm-like chain (DWLC) — and calls for investigation of signatures of bending dissipation in other semiflexible polymers such as actin and microtubules. This work establishes bending friction as a fundamental material property that must underpin any model of DNA dynamics in molecular biology, DNA-based condensates[15] and in material science for its implications for defining new engineering principles of DNA nanodevices and nanomaterials[21, 25, 47].

ACKNOWLEDGEMENTS

We would like to thank Ard Louis, Agnes Noy, Tanniemola Liverpool and Frank Uhlmann, for various discussions and suggestions on the text and figures of the manuscript. We thank Holly Folkard-Tapp for her advice on the design of the graphical elements.

AUTHOR CONTRIBUTIONS:

GP performed the experiments. MM contributed to study design, supervised and provided funding for the experiments. BSK conceived and designed the study, and performed the theoretical calculations. BSK and MM wrote the manuscript with input from GP.

Appendix A: Methods

1. DNA functionalisation

To facilitate attachment to beads, shorter DNA molecules (2.7 kbp and 4.5 kbp) were labelled with a single biotin and a single digoxigenin at either end of the DNA molecule. For longer molecules (6.5 kbp and 8.8 kbp) both DNA ends were labelled with biotins. Restriction enzymes, DNA polymerases and buffers were purchased from New England Biolabs (NEB). Oligonucleotides were purchased from Merck. All PCR reactions were performed using Phusion High-Fidelity DNA Polymerase in a Phusion HF Buffer (NEB).

a.   2,655 bp biotin-digoxigenin DNA

2,655 bp DNA fragment was synthesised by PCR using Lambda-phage DNA (NEB) as a template, a 5'-biotin containing primer (LHD2_RT7bio) and a primer containing XbaI restriction site (LHD2_FXbaI). PCR reaction was purified using a PCR Cleanup Kit (Monarch T1030S, NEB) and the product was further digested with XbaI restriction enzyme (NEB) in a Standard Taq Reaction Buffer (NEB) for 40 minutes at 37 °C to generate a 4-nt single-stranded overhang. XbaI was inactivated by incubating the reaction for 20 minutes at 65 °C and the second end of the DNA was labelled with digoxigenin by end filling reaction using Taq DNA polymerase (NEB) and a mixture of dATP, dCTP, dGTP (Promega) and dUTP-digoxigenin (Jena Bioscience) performed for 30 min at 72 °C. The final product was purified using Micro Bio-Spin P30 spin column (Bio-Rad).

 b. 4,500 bp biotin-digoxigenin DNA

4,500 bp DNA fragment was synthesised by PCR using Lambda-phage DNA
(NEB) as a template, a 5'-digoxigenin containing primer (LHT1_Fdig) and a 5'-
biotin containing primer (4.5kb_rev_bio). PCR reaction was purified using a PCR
Cleanup Kit (Monarch T1030S, NEB).

c. 6,473 bp biotin-biotin DNA

9,847 bp plasmid DNA (8x601-pKYB1, a gift from Graeme King) was double
digested with XbaI and SpeI restriction enzymes (NEB) for 1 hour at 37 °C in
rCutSmart buffer (NEB), followed by inactivation for 20 minutes at 80 °C. This
resulted in 2 fragments (6,473 bp and 3,374 bp long) containing 4-nt single-stranded
ends. DNA ends were further biotinylated by addition of Klenow Fragment ($3´{\rightarrow}5´$
exo-) of DNA Pol I (NEB), a mixture of dATP, dTTP, dGTP (Promega) and
dCTP-biotin (Jena Bioscience) and incubation for 25 minutes at 37 °C, followed
by inactivation for 20 minutes at 75 °C. DNA was cleaned-up from unincorporated
nucleotides using Micro Bio-Spin P30 spin column (Bio-Rad). After tethering
DNA to the beads only 6,473 bp-long molecules were selected by force-extension
analysis.

d. 8,815 bp biotin-biotin DNA

8,815 bp plasmid DNA (2x601-pKYB1, a gift from Graeme King) was digested
with XbaI (NEB) in the Standard Taq Reaction Buffer (NEB) for 45 minutes at
37 °C to generate a 4-nt single-stranded overhangs. DNA ends were biotinylated
by end filling reaction using Taq DNA polymerase (NEB) and a mixture of dATP,
dTTP, dGTP (Promega) and dCTP-biotin (Jena Bioscience) performed for 30 min
at 72 °C. The final product was purified using the Micro Bio-Spin P30 spin column

(Bio-Rad).

4.5kb_rev_bio	Biotin-GTAAAAGCTCTTGGATTCCTGAAAC
LHT1_Fdig	Digoxigenin-CTGTTACAGGTCACTAATACCATC
LHD2_FXbaI	CACTCTAGAGTGTTTGATCCATTCTTTGGGAC
LHD2_RT7bio	Biotin-TAATACGACTCACTATAGGGTTTCCAGCATAAGCGGCTACATG

TABLE I. List of oligonucleotides

2. Single-molecule DNA tethering

Experiments were performed using commercial dual-trap optical tweezers (C-
trap, Lumicks) combined with a multi-channel microfluidic laminar flow cell (u-
Flux, Lumicks). Before experiments, the flow cell was passivated by incubating it
with 0.5% Pluronic F-127 (Merck) diluted in PBS (Phosphate Buffered Saline) for
at least 30 minutes. Pluronic was washed by flowing at least 1 mL of PBS through
each channel of the flow cell. Individual DNA molecules were tethered between
2 polystyrene beads of 2 μm diameter. Shorter digoxigenin-biotin labelled DNA
molecules (2.7 and 4.5 kbp) were first coupled to Anti-digoxigenin-coated beads
(DIGP-20-2, Spherotech): 5 microliters of beads stock solution (0.1% w/v) were
gently mixed with 5 microliters of DNA diluted down to 30 pM in PBS and in-
cubated for 10 min. The mixture was further diluted by adding 300 microliters
of PBS and loaded into the first channel of the flow cell. The second channel was
loaded with PBS and third channel contained Streptavidin-coated beads (SVP-20-
5, Spherotech) diluted down to 0.002% w/v in PBS. 4th and 5th channels of the
flow-cell contained a low salt buffer (Hepes pH 7.5 1mM, NaCl 1mM). To form
a DNA tether, first a bead-DNA complex and a Streptavidin-coated bead were
consecutively trapped in two optical traps in the first and the third channels re-
spectively. The beads were further moved into the second channel where the free
biotinylated end of DNA was attached to a Streptavidin-coated bead. The flow

cell was extensively washed with PBS keeping only the 2nd channel open to re-move any free-floating beads, after which the flow was stopped and measurements started. For longer biotinylated DNA molecules (6.5 and 8.8 kbp) the first chan-nel contained Streptavidin-coated beads (SVP-20-5, Spherotech) diluted down to 0.002% w/v in PBS, second channel contained DNA diluted down to 5 pM in PBS, and the third channel was loaded with PBS. Two beads were trapped in the first channel and transferred into the second channel to form a DNA tether, which was further transferred into the 3rd channel and the flow was stopped prior to measurements. For measurements in the low salt buffer, after DNA fluctuations were measured in PBS, the 4th and 5th channels were extensively washed with the Hepes buffer (2 minutes at the pressure of 1 Bar). After this the flow was stopped and the DNA tether was transferred into the 5th channel where measurements were repeated. Not all DNA tethers lasted long enough to be measured in both PBS and Hepes buffers and hence in Table.II the number of single molecules for PBS and Hepes differ for each length. After molecule detachment, time series of bead positions were collected at approximately the same trap separations as the force measurements with DNA.

3.   Force-extension and force-dependent DNA fluctuation measurements

After the DNA tether was formed and placed in a channel containing either PBS or a low salt buffer, a force-extension curve was recorded by increasing the beads separation at the rate of 40 nm per second while simultaneously recording the distance between the beads and the force in the optical traps until the DNA tension reached 10 pN.

After that the DNA molecule was fully relaxed by bringing the beads closer to each other. Subsequently DNA was stretched to a required force and the fluctua-tions of each bead were recorded simultaneously at $f_s = 78.125\text{kHz}$ for a period of

₄₁₂ 6 seconds. These measurements were performed at DNA tensions varying between ₄₁₃ 1pN and 5pN with 0.5 pN incremental steps, with 3 repeats per each force.

₄₁₄ Prior to the measurements the laser power was adjusted to reach the nominal ₄₁₅ trap stiffness of $\kappa_1 = \kappa_2 \approx 0.05$pN for each optical trap.

₄₁₆ Data were acquired using a custom-made automation script. High-frequency ₄₁₇ force data and the distance between the beads measured at ≈ 50Hz were saved as ₄₁₈ an h5 file and further processed using Matlab.

₄₁₉ The final numbers of molecules used for each length and each buffer condition ₄₂₀ are as in Table.II

Length	n
2.7kbp	(20,6)
4.5kbp	(21,18)
6.5kbp	(14,10)
8.8kbp	(13,8)

TABLE II. Number of single molecules used at each length and each buffer condition. The first number in the brackets are the number of single molecules used in PBS buffer and the second number the number used in the Hepes buffer. Each single molecule experiment consisted of measurements in PBS followed by Hepes; however, the Hepes stage did not complete for all single molecules and so the 2nd number is typically less than the first.

₄₂₁ For each single molecule experiment, we corrected forces according to Supple- ₄₂₂ mentary Information (section S6) to account for small positive forces — in the ₄₂₃ absence of DNA — between beads due to the cross-talk between traps at small ₄₂₄ trap separations.

₄₂₅ **4. Data processing**

Raw data in the form of force signals from both traps were used to extract the DNA length fluctuations as follows: at each time point the length of the DNA

molecule was calculated as

$$\ell(t) = D - 2b + x_2(t) - x_1(t) = \langle \ell \rangle + \frac{\langle F_1 \rangle + F_1(t)}{\kappa_1} + \frac{\langle F_2 \rangle - F_2(t)}{\kappa_2}$$

where D is the distance between trap centres, b the radius of each bead, $\langle F_2 \rangle =$
$-\langle F_1 \rangle = F$ are the mean force in each trap averaged over the measurement period
(6 seconds), where F is the magnitude of the desired force, κ_1, κ_2 are the trap
stiffness values for trap 1 and trap 2, and are nominally equal and $\langle \ell \rangle$ is the
mean DNA length, calculated as the distance between the beads averaged over
the measurement period (6 seconds) with $2b$ subtracted, and F_1, and F_2 are the
instantaneous forces on the beads in trap 1 and trap 2.

The power spectrum $P_\ell(\omega)$ of the time-series of DNA length $\ell(t)$ is then calcu-
lated by standard Fast Fourier Transform (FFT) in MatLab,

$$P_\ell(\omega) = \frac{1}{T} \mathsf{FFT}\{\ell(t)\} \times \mathsf{FFT}^*\{\ell(t)\}$$

where * indicates complex conjugate and $T = 5000\mathrm{msec}$ is the maximum obser-
vation time in the data. The power spectrum is then block-averaged on a log-
frequency scale [6] with $n = 25$ blocks, to reduce the noise on the raw power
spectrum estimate. The zero-frequency (DC) component is removed and only fre-
quencies less than the Nyquist frequency ($f_s/2$) are retained. The velocity power
spectrum is then calculated by $P_v(\omega) = \omega^2 P_\ell(\omega)$.

As described in the main text, the velocity power spectrum is then fit with
Eqn.S18 (Supplementary Information) at each force F to obtain $\zeta(F)$, $\kappa(F)$ and
$m_{eff}(F)$.

---

444 [1] J. F. Allemand, S. Cocco, N. Douarche, and G. Lia. Loops in DNA: An overview
445   of experimental and theoretical approaches. *The European Physical Journal E,*

446   19(3):293–302, 2006.

[2] Daniella Bar-Lev, Omer Sabary, and Eitan Yaakobi. The zettabyte era is in our DNA. *Nature Computational Science*, 4(11):813–817, 2024.

[3] Aakash Basu, Dmitriy G. Bobrovnikov, Zan Qureshi, Tunc Kayikcioglu, Thuy T. M. Ngo, Anand Ranjan, Sebastian Eustermann, Basilio Cieza, Michael T. Morgan, Miroslav Hejna, H. Tomas Rube, Karl-Peter Hopfner, Cynthia Wolberger, Jun S. Song, and Taekjip Ha. Measuring DNA mechanics on the genome scale. *Nature*, 589(7842):462–467, 2021.

[4] Christoph G. Baumann, Steven B. Smith, Victor A. Bloomfield, and Carlos Busta- mante. Ionic effects on the elasticity of single DNA molecules. *Proceedings of the National Academy of Sciences*, 94(12):6185–6190, 1997.

[5] David R. Bentley, Shankar Balasubramanian, Harold P. Swerdlow, Geoffrey P. Smith, John Milton, Clive G. Brown, Kevin P. Hall, Dirk J. Evers, Colin L. Barnes, Helen R. Bignell, Jonathan M. Boutell, Jason Bryant, Richard J. Carter, R. Keira Cheetham, Anthony J. Cox, Darren J. Ellis, Michael R. Flatbush, Niall A. Gormley, Sean J. Humphray, Leslie J. Irving, Mirian S. Karbelashvili, Scott M. Kirk, Heng Li, Xiaohai Liu, Klaus S. Maisinger, Lisa J. Murray, Bojan Obradovic, Tobias Ost, Michael L. Parkinson, Mark R. Pratt, Isabelle M. J. Rasolonjatovo, Mark T. Reed, Roberto Rigatti, Chiara Rodighiero, Mark T. Ross, Andrea Sabot, Subra- manian V. Sankar, Aylwyn Scally, Gary P. Schroth, Mark E. Smith, Vincent P. Smith, Anastassia Spiridou, Peta E. Torrance, Svilen S. Tzonev, Eric H. Vermaas, Klaudia Walter, Xiaolin Wu, Lu Zhang, Mohammed D. Alam, Carole Anastasi, Ify C. Aniebo, David M. D. Bailey, Iain R. Bancarz, Saibal Banerjee, Selena G. Barbour, Primo A. Baybayan, Vincent A. Benoit, Kevin F. Benson, Claire Be- vis, Phillip J. Black, Asha Boodhun, Joe S. Brennan, John A. Bridgham, Rob C. Brown, Andrew A. Brown, Dale H. Buermann, Abass A. Bundu, James C. Burrows, Nigel P. Carter, Nestor Castillo, Maria Chiara E. Catenazzi, Simon Chang, R. Neil

Cooley, Natasha R. Crake, Olubunmi O. Dada, Konstantinos D. Diakoumakos, Belen Dominguez-Fernandez, David J. Earnshaw, Ugonna C. Egbujor, David W. Elmore, Sergey S. Etchin, Mark R. Ewan, Milan Fedurco, Louise J. Fraser, Karin V. Fuentes Fajardo, W. Scott Furey, David George, Kimberley J. Gietzen, Colin P. Goddard, George S. Golda, Philip A. Granieri, David E. Green, David L. Gustafson, Nancy F. Hansen, Kevin Harnish, Christian D. Haudenschild, Narinder I. Heyer, Matthew M. Hims, Johnny T. Ho, Adrian M. Horgan, Katya Hoschler, Steve Hurwitz, Denis V. Ivanov, Maria Q. Johnson, Terena James, T. A. Huw Jones, Gyoung-Dong Kang, Tzvetana H. Kerelska, Alan D. Kersey, Irina Khrebtukova, Alex P. Kindwall, Zoya Kingsbury, Paula I. Kokko-Gonzales, Anil Kumar, Marc A. Laurent, Cynthia T. Lawley, Sarah E. Lee, Xavier Lee, Arnold K. Liao, Jennifer A. Loch, Mitch Lok, Shujun Luo, Radhika M. Mammen, John W. Martin, Patrick G. McCauley, Paul McNitt, Parul Mehta, Keith W. Moon, Joe W. Mullens, Taksina Newington, Zemin Ning, Bee Ling Ng, Sonia M. Novo, Michael J. O'Neill, Mark A. Osborne, Andrew Osnowski, Omead Ostadan, Lambros L. Paraschos, Lea Pickering, Andrew C. Pike, Alger C. Pike, D. Chris Pinkard, Daniel P. Pliskin, Joe Podhasky, Victor J. Quijano, Come Raczy, Vicki H. Rae, Stephen R. Rawlings, Ana Chiva Rodriguez, Phyllida M. Roe, John Rogers, Maria C. Rogert Bacigalupo, Nikolai Romanov, Anthony Romieu, Rithy K. Roth, Natalie J. Rourke, Silke T. Ruediger, Eli Rusman, Raquel M. Sanches-Kuiper, Martin R. Schenker, Josefina M. Seoane, Richard J. Shaw, Mitch K. Shiver, Steven W. Short, Ning L. Sizto, Johannes P. Sluis, Melanie A. Smith, Jean Ernest Sohna Sohna, Eric J. Spence, Kim Stevens, Neil Sutton, Lukasz Szajkowski, Carolyn L. Tregidgo, Gerardo Turcatti, Stephanie vandeVondele, Yuli Verhovsky, Selene M. Virk, Suzanne Wakelin, Gregory C. Walcott, Jingwen Wang, Graham J. Worsley, Juying Yan, Ling Yau, Mike Zuerlein, Jane Rogers, James C. Mullikin, Matthew E. Hurles, Nick J. McCooke, John S. West, Frank L. Oaks, Peter L. Lundberg, David Klenerman, Richard Durbin, and

Anthony J. Smith. Accurate whole human genome sequencing using reversible terminator chemistry. *Nature*, 456(7218):53–59, 2008.

[6] K. Berg-Sorensen and H. Flyvbjerg. Power spectrum analysis for optical tweezers. *Rev. Sci. Ins.*, 75:594–612, 2004.

[7] Elizabeth M. Blackwood and James T. Kadonaga. Going the Distance: A Current View of Enhancer Action. *Science*, 281(5373):60–63, 1998.

[8] C Bustamante, JF Marko, ED Siggia, and S Smith. Entropic elasticity of lambda-phage DNA. *Science*, 265(5178):1599–1600, 1994.

[9] R. Cerf. Possible existence of two classes of conformational changes in linear polymeric chain molecules. *Chemical Physics Letters*, 22(3):613–615, 1973.

[10] Huimin Chen, Steve P. Meisburger, Suzette A. Pabit, Julie L. Sutton, Watt W. Webb, and Lois Pollack. Ionic strength-dependent persistence lengths of single-stranded RNA and DNA. *Proceedings of the National Academy of Sciences*, 109(3):799–804, 2012.

[11] Nicholas T. Crump and Thomas A. Milne. Is Enhancer Function Driven by Protein–Protein Interactions? From Bacteria to Leukemia. *BioEssays*, 47(6):e70006, 2025.

[12] Iain F. Davidson and Jan-Michael Peters. Genome folding through loop extrusion by SMC complexes. *Nature Reviews Molecular Cell Biology*, 22(7):445–464, 2021.

[13] Pierre Gilles de Gennes. *Scaling Concepts in Polymer Physics*. Cornell University Press, 1985.

[14] Job Dekker and Leonid A. Mirny. The chromosome folding problem and how cells solve it. *Cell*, 187(23):6424–6450, 2024.

[15] Sungho Do, Chanseok Lee, Taehyun Lee, Do-Nyun Kim, and Yongdae Shin. Engineering DNA-based synthetic condensates with programmable material properties, compositions, and functionalities. *Science Advances*, 8(41):eabj1771, 2022.

[16] M. Doi and S.F. Edwards. *The Theory of Polymer Dynamics*, chapter 8, pages

295–296. Oxford University Press, 1986.

[17] M. Doi and S.F. Edwards. *The Theory of Polymer Dynamics*. Oxford University Press, 1986.

[18] M. Doi and S.F. Edwards. *The Theory of Polymer Dynamics*, chapter 3, pages 56–58. Oxford University Press, 1986.

[19] Molly Gasperini, Andrew J. Hill, José L. McFaline-Figueroa, Beth Martin, Seungsoo Kim, Melissa D. Zhang, Dana Jackson, Anh Leith, Jacob Schreiber, William S. Noble, Cole Trapnell, Nadav Ahituv, and Jay Shendure. A Genome-wide Framework for Mapping Gene Regulation via Cellular Genetic Screens. *Cell*, 176(1-2):377–390.e19, 2019.

[20] Alexey A. Gavrilov, Ilya Shamovsky, Irina Zhegalova, Sergey Proshkin, Yosef Shamovsky, Grigory Evko, Vitaly Epshtein, Aviram Rasouly, Anna Blavatnik, Sudipta Lahiri, Eli Rothenberg, Sergey V. Razin, and Evgeny Nudler. Elementary 3D organization of active and silenced E. coli genome. *Nature*, pages 1–11, 2025.

[21] Bartosz A. Grzybowski and Wilhelm T. S. Huck. The nanotechnology of life-inspired systems. *Nature Nanotechnology*, 11(7):585–592, 2016.

[22] Sébastien Guilbaud, Laurence Salomé, Nicolas Destainville, Manoel Manghi, and Catherine Tardin. Dependence of DNA Persistence Length on Ionic Strength and Ion Type. *Physical Review Letters*, 122(2):028102, 2019.

[23] Tetsuya Hiraiwa and Takao Ohta. Linear viscoelasticity of a single semiflexible polymer with internal friction. *The Journal of chemical physics*, 133(4):044907, 2010.

[24] Tatsuya Hirano. Condensin-Based Chromosome Organization from Bacteria to Vertebrates. *Cell*, 164(5):847–857, 2016.

[25] Ryota Ibusuki, Tatsuya Morishita, Akane Furuta, Shintaro Nakayama, Maki Yoshio, Hiroaki Kojima, Kazuhiro Oiwa, and Ken'ya Furuta. Programmable molecular

transport achieved by engineering protein motors to move on DNA nanotubes. *Science*, 375(6585):1159–1164, 2022.

[26] Jiyoun Jeong and Harold D Kim. Determinants of cyclization-decyclization kinetics of short DNA with sticky ends. *Nucleic Acids Research*, 48(9):5147–5156, 2020.

[27] S. Jun, J. Bechhoefer, and B.-Y. Ha. Diffusion-limited loop formation of semiflexible polymers: Kramers theory and the intertwined time scales of chain relaxation and closing. *Europhysics Letters*, 64(3):420–426, 2003.

[28] Masaru Kawakami, Katherine Byrne, Bhavin S Khatri, Tom C B McLeish, and D. Alastair Smith. Viscoelastic properties of single poly(ethylene glycol) molecules. *Chemphyschem*, 7(8):1710–1716, Aug 2006.

[29] Bhavin S. Khatri and Tom Charles Buckland McLeish. Rouse model with internal friction: A coarse grained framework for single biopolymer dynamics. *Macromolecules*, 40:6770 –6777, 2007.

[30] Mary L. Kopka, Albert V. Fratini, Horace R. Drew, and Richard E. Dickerson. Ordered water structure around a B-DNA dodecamer A quantitative study. *Journal of Molecular Biology*, 163(1):129–146, 1983.

[31] W. Kuhn and H. Kuhn. *Helv. Chim. Acta*, 28:1533, 1945.

[32] Tung T. Le and Harold D. Kim. Measuring Shape-Dependent Looping Probability of DNA. *Biophysical Journal*, 104(9):2068–2076, 2013.

[33] Chi H. Mak. Hydration Waters Make Up for the Missing Third Hydrogen Bond in the A·T Base Pair. *ACS Physical Chemistry Au*, 4(2):180–190, 2024.

[34] Marshal Mandelkern, John G. Elias, Don Eden, and Donald M. Crothers. The dimensions of DNA in solution. *Journal of Molecular Biology*, 152(1):153–161, 1981.

[35] John F Marko and Eric D Siggia. Stretching DNA. *Macromolecules*, 28(26):8759–8770, 1995.

[36] Peter J. Mulligan, Yi-Ju Chen, Rob Phillips, and Andrew J. Spakowitz. Interplay

of Protein Binding Interactions, DNA Mechanics, and Entropy in DNA Looping Kinetics. *Biophysical Journal*, 109(3):618–629, 2015.

[37] Natalia Naumova, Maxim Imakaev, Geoffrey Fudenberg, Ye Zhan, Bryan R. Lajoie, Leonid A. Mirny, and Job Dekker. Organization of the Mitotic Chromosome. *Science*, 342(6161):948–953, 2013.

[38] Jacobus W M Noordermeer, John D Ferry, and Norio Nemoto. Viscoelastic Properties of Polymer Solutions in High-Viscosity Solvents and Limiting High-Frequency Behavior. III. Poly(2-substituted methyl acrylates). *Macromolecules*, 8(5):672–677, 1975.

[39] Kunihiro Osaki and John L Schrag. Viscoelastk Properties of Polymer Solutions in High-Viscosity Solvents and Limiting High-Frequency Behavior. I. Polystyrene and Poly(-methylstyreste). *Polymer Journal*, 2(4):541–549, 1971.

[40] Thomas E. Ouldridge, Ard A. Louis, and Jonathan P. K. Doye. DNA Nanotweezers Studied with a Coarse-Grained Model of DNA. *Physical Review Letters*, 104(17):178101, 2009.

[41] Anil Panigrahi and Bert W. O'Malley. Mechanisms of enhancer action: the known and the unknown. *Genome Biology*, 22(1):108, 2021.

[42] Georgii Pobegalov, Lee-Ya Chu, Jan-Michael Peters, and Maxim I. Molodtsov. Single cohesin molecules generate force by two distinct mechanisms. *Nature Communications*, 14(1):3946, 2023.

[43] Michael G Poirier and John F Marko. Effect of internal friction on biofilament dynamics. *Physical review letters*, 88(22):228103, 2002.

[44] Martina Richeldi, Georgii Pobegalov, Torahiko L. Higashi, Karolina Gmurczyk, Frank Uhlmann, and Maxim I. Molodtsov. Mechanical disengagement of the cohesin ring. *Nature Structural & Molecular Biology*, 31(1):23–31, 2024.

[45] Timothy J. Richmond and Curt A. Davey. The structure of DNA in the nucleosome core. *Nature*, 423(6936):145–150, 2003.

[46] Ahmed M. Rozza, Imre Bakó, and Julianna Oláh. Theoretical insights into water network of B-DNA duplex with Watson-Crick and Hoogsteen base pairing geometries. *Journal of Molecular Liquids*, 362:119728, 2022.

[47] Nadrian C. Seeman and Hanadi F. Sleiman. DNA nanotechnology. *Nature Reviews Materials*, 3(1):17068, 2017.

[48] Yeonee Seol, Jinyu Li, Philip C. Nelson, Thomas T. Perkins, and M.D. Betterton. Elasticity of Short DNA Molecules: Theory and Experiment for Contour Lengths of 0.6-7m. *Biophysical Journal*, 93(12):4360–4373, 2007.

[49] Friedrich C. Simmel, Bernard Yurke, and Hari R. Singh. Principles and Applications of Nucleic Acid Strand Displacement Reactions. *Chemical Reviews*, 119(10):6326–6369, 2019.

[50] Frank Uhlmann. SMC complexes: from DNA to chromosomes. *Nature Reviews Molecular Cell Biology*, 17(7):399–412, 2016.

[51] Reza Vafabakhsh and Taekjip Ha. Extreme Bendability of DNA Less than 100 Base Pairs Long Revealed by Single-Molecule Cyclization. *Science*, 337(6098):1097–1101, 2012.

[52] Jay R. Wenner, Mark C. Williams, Ioulia Rouzina, and Victor A. Bloomfield. Salt Dependence of the Elasticity and Overstretching Transition of Single DNA Molecules. *Biophysical Journal*, 82(6):3160–3169, 2002.

[53] Paul A. Wiggins, Thijn van der Heijden, Fernando Moreno-Herrero, Andrew Spakowitz, Rob Phillips, Jonathan Widom, Cees Dekker, and Philip C. Nelson. High flexibility of DNA on short length scales probed by atomic force microscopy. *Nature Nanotechnology*, 1(2):137–141, 2006.

[54] Hiromi Yamakawa and W H Stockmayer. Statistical Mechanics of Wormlike Chains. II. Excluded Volume Effects. *The Journal of Chemical Physics*, 57(7):2843–2854, 1972.

[55] Bernard Yurke, Andrew J. Turberfield, Allen P. Mills, Friedrich C. Simmel, and

635   Jennifer L. Neumann. A DNA-fuelled molecular machine made of DNA. *Nature*,
636   406(6796):605–608, 2000.