1 **Dicer accumulates in cytoplasmic foci upon alphavirus infection and**

2 **plays a proviral role in _Myotis myotis_ bat cells**

5 Léa Gaucherand*, Hugo Marie*, Julie Cremaschi, Sébastien Pfeffer[‡]

7 Université de Strasbourg, Architecture et Réactivité de l'ARN, Institut de Biologie

8 Moléculaire et Cellulaire du CNRS, 67000 Strasbourg, France

14 * These authors contributed equally and should be considered as co-first authors

16 [‡] To whom correspondence should be addressed: s.pfeffer@ibmc-cnrs.unistra.fr

Abstract

Bats are reservoirs for many viruses that frequently cause epidemics in humans and animals. It is thus critical to better understand their immune system and mechanisms of antiviral immunity. Despite an increasing number of studies, much is still unknown about the molecular mechanisms that govern bat-virus interactions, especially given the large diversity of bat species. Dicer is a conserved ribonuclease with multiple activities that can modulate antiviral immunity, including the detection of viral RNA as part of the RNA interference (RNAi) pathway, the maturation of micro RNAs, and the direct inhibition of innate immunity in mouse and human cells. In view of these complex activities of Dicer, we tested its antiviral activity in *Myotis myotis* nasal epithelial cells. Surprisingly, we did not see strong evidence of RNAi in these cells, but instead saw a proviral effect of Dicer for two alphaviruses, Sindbis and Semliki forest virus. We also observed a striking relocalization of Dicer to cytoplasmic foci upon infection with these viruses, which did not occur in the several human cell lines we tested. These foci contained dsRNA and viral plus strand RNA, suggesting that they are sites of viral replication. Finally, we found that factors specific to *M. myotis* cells are needed for Dicer relocalization. Overall, we propose that Dicer can play different roles in different bat species and/or cell types, and is being repurposed by alphaviruses to promote viral replication.

Introduction

Bats are unique mammals in many aspects, such as their ability to fly and their high longevity relative to their body size (Gonzalez & Banerjee, 2022). They belong to the order Chiroptera, which can be divided into two suborders that diverged over 50 million years ago (Lei & Dong, 2016). More than 1400 different species of bats have been identified so far, accounting for 20% of all mammals (Burgin *et al*, 2018). Importantly, bats are reservoirs for many viruses that can spillover into humans, either directly or through an intermediate host, such as Marburg virus, Hendra and Nipah virus, Middle East respiratory syndrome coronavirus (MERS-CoV), severe acute respiratory syndrome coronavirus (SARS-CoV) and likely SARS-CoV-2 (Halpin *et al*, 2000; Li *et al*, 2005; Memish *et al*, 2013; Moratelli & Calisher, 2015; Chua *et al*, 2002; Yob *et al*, 2001; Towner *et al*, 2007). Interestingly, while these viruses can be very pathogenic to humans and other animals, most of the studied bats so far show little to no clinical symptoms, whether they were naturally or experimentally infected (Gonzalez &

Banerjee, 2022; Pei *et al*, 2024). An increasing number of studies are highlighting the role of the bat innate immune system in this apparent tolerance to viruses.

The type I and III interferon (IFN) pathways are the main first lines of defense against viruses in mammals. In this pathway, sensing of viral nucleic acids by pattern recognition receptors (PRRs) leads to the expression and secretion of type I and III IFN cytokines. These cytokines are then sensed by cells in an autocrine and paracrine fashion, leading to the expression of interferon stimulated genes (ISGs) that can actively fight the infection (Pichlmair & Reis e Sousa, 2007). The IFN pathway is present and active in bats, but differences have been reported compared to humans, as well as between different bat species. These include a dampening of the response to DNA sensing, as well as differences in the baseline expression and temporal regulation of some IFNs and ISGs (Irving *et al*, 2021; Banerjee *et al*, 2020; Clayton & Munir, 2020). In contrast, invertebrates and plants mostly rely on the RNA interference (RNAi) pathway to fight viral infections. In antiviral RNAi, viral double-stranded (ds)RNAs are sensed and cleaved into small interfering (si)RNAs by a Dicer or Dicer-like ribonuclease, then loaded into a protein of the Argonaute family to target and degrade complementary viral RNAs (Guo *et al*, 2019). Interestingly, two recent studies have reported enhanced antiviral activity of Dicer through the RNAi pathway in the *Pteropus alecto* kidney (PaKi) cell line infected with SINV, an NS1 deletion mutant strain of infuenza A virus and a B2 deletion mutant strain of Nodamura virus, and in the *Tadarida brasiliensis* lung (TBLU) cell line infected with SARS-CoV-2 (Dai *et al*, 2024; Owolabi *et al*, 2025). The first study also proposed that viral dsRNA cleavage by Dicer limits the recognition of dsRNA by PRRs and thus the activation of the IFN response in *P. alecto*, thereby suggesting a role for Dicer in bat viral tolerance (Dai *et al*, 2024).

Dicer is well known for its essential role in the maturation of most micro (mi)RNAs, key regulators of gene expression (Bartel, 2018). In mammals, the same Dicer protein can generate both miRNAs and siRNAs of 22 nucleotides (nt) thanks to its regulatory cofactors transactivation response element RNA-binding protein (TRBP) and Protein ACTivator of the interferon-induced protein kinase (PACT) (Lee *et al*, 2013). However, the antiviral activity of Dicer through the RNAi pathway is controversial and appears to be cell-type and context-dependent in human and mouse cells (Cullen *et al*, 2013). Indeed, Dicer isoforms with increased RNAi activity have been identified in human and mouse stem cells (Poirier *et al*, 2021), and in mouse oocytes (Flemr *et al*, 2013). While

the former appears to have some antiviral properties against viruses such as Zika and
SARS-CoV-2 (Poirier *et al*, 2021), the mouse oocyte specific one does not confer an
antiviral phenotype when artificially expressed ubiquitously (Kulmann *et al*, 2025). One
important reason for the reduced antiviral activity of full-length Dicer is the existence
of a regulatory crosstalk between the IFN and RNAi pathways (Baldaccini & Pfeffer,
2021; Gaucherand *et al*, 2024; Maillard *et al*, 2019). Human Dicer is inhibited by the
PRR LGP2 (laboratory of genetics and physiology 2 / DexH box polypeptide 58) (van
der Veen *et al*, 2018). Moreover, inactivating the IFN pathway in differentiated mouse
cells unmasks long dsRNA processing by Dicer (Maillard *et al*, 2016). Conversely,
Dicer represses IFN and the ISG protein kinase R (PKR) activity in mouse embryonic
stem cells transfected with double stranded RNA (Gurung *et al*, 2021). In addition, we
recently showed that human Dicer interacts with PKR *via* its helicase domain and
regulates innate immune activation during Sindbis virus (SINV) infection (Montavon *et
al*, 2021; Baldaccini *et al*, 2024).

Overall, Dicer could modulate antiviral activity through its role in the RNAi pathway,
but also by maturing miRNAs that can affect viral infection, by directly regulating innate
immune activation, and potentially through other still unknown mechanisms. How
these multiple complex activities of Dicer could contribute to antiviral activity and viral
tolerance in different bat species is still poorly understood. This prompted us to
investigate the antiviral activity of Dicer in the greater mouse-eared bat (*Myotis
myotis*), a European microbat from the *Vespertilionidae* family. Unexpectedly,
expressing *M. myotis* Dicer in human Dicer knock out cells did not protect from viral
infection. Instead, we saw a proviral role of Dicer in *M. myotis* nasal epithelial cells
upon infection with two different alphaviruses, with no clear evidence of antiviral RNAi
activity. Interestingly, Dicer relocalizes to large cytoplasmic foci upon infection with
these viruses in *M. myotis* nasal epithelial cells but not in human cells. We find that
these foci are likely viral factories and that *M. myotis* cell-specific factors are required
for Dicer relocalization. Overall, we propose that the activity of bat Dicer is context-
dependent and could be used by some viruses like SINV in *M. myotis* cells to promote
viral replication.

Results
Expressing *M. myotis* Dicer in NoDice cells does not protect from SINV and VSV
infection

Since *P. alecto* and *T. brasiliensis* Dicer were found to be antiviral against a few viruses
(Dai *et al*, 2024; Owolabi *et al*, 2025), we tested the antiviral activity of *M. myotis* Dicer.
We cloned Dicer from *M. myotis* cells (hereafter referred to as mmDicer) and
expressed it with an HA tag in human embryonic kidney (HEK) 293T cells knock-out
for Dicer (NoDice cells (Bogerd *et al*, 2014)). As controls, we also expressed an HA-
tagged catalytically inactive mutant (amino acid changes E1562A and E1819A)
version of *M. myotis* Dicer (hereafter referred to as mmDicer CM), HA-tagged human
Dicer (hDicer), or an empty vector (Empty). Efficient expression of the different Dicer
constructs was checked by western blot (**Fig. 1A**). Expressing mmDicer but not
mmDicer CM also rescued maturation of the human miRNA miR-16 (**Fig. 1B**). These
results suggested that mmDicer is indeed expressed and active in NoDice cells. We
then tested its antiviral activity against a SINV that expresses GFP (SINV-GFP) (López
et al, 2020). Expressing mmDicer did not lead to a significant decrease in SINV-GFP
titers, RNA and protein levels compared to the empty vector control, hDicer or
mmDicer CM (**Fig. 1C-E**), suggesting that mmDicer is not particularly antiviral against
SINV in this context. We also tested a different virus, the rhabdovirus vesicular
stomatitis virus expressing GFP (VSV-GFP), and again saw no antiviral activity of
mmDicer (**Fig. 1F**). Overall, these results suggest that expressing mmDicer in a
human cell line does not protect against SINV and VSV infection.

<u>Dicer is proviral for two alphaviruses in *M. myotis* nasal epithelial cells</u>
To determine whether mmDicer displays antiviral activity in *M. myotis* cells, we used a
pool of multiple siRNAs at low concentrations to knock down Dicer in a SV40
transformed *M. myotis* nasal epithelial cell line (He *et al*, 2014b). This technique allows
for efficient knock down (**Fig. 2A**) while limiting off-target effects. We then infected
these cells with SINV-GFP and VSV-GFP. Interestingly, we saw that knocking down
Dicer actually reduced SINV-GFP replication, as seen by a decrease in GFP+ cells
over time (**Fig. 2B**). At 24h, we also saw a decrease in SINV-GFP titers, in GFP protein
levels and in SINV RNA levels (**Fig. 2C-E**). To test whether the proviral effect of Dicer
could be recapitulated with another alphavirus, we infected our *M. myotis* cell line with
Semliki forest virus (SFV). Similar to SINV infection, knocking down Dicer decreased
SFV titers and RNA levels at 24 hours post infection (**Fig. 2F-G**). In contrast, Dicer
knock down had an antiviral effect against VSV-GFP infection at the RNA and protein
levels, but no difference in titers was found (**Fig. 2H-J**). Overall, our results suggest

that Dicer plays a proviral role for SINV and SFV but not VSV infection in *M. myotis*
nasal epithelial cells.

M. myotis nasal epithelial cells only display a mild RNAi signature
Since *P. Alecto* and *T. brasiliensis* Dicer have enhanced antiviral RNAi activity, we
wondered whether there was any evidence of antiviral RNAi activity in our *M. myotis*
nasal epithelial cell line. We infected these cells with SINV-GFP and VSV-GFP, then
sequenced small RNAs isolated from these cells. As control, we followed the same
procedure with HEK293T cells, which we have previously showed display a weak
RNAi signature against SINV-GFP (Girardi *et al*, 2013). Consistent with the lack of
antiviral activity of mmDicer against SINV-GFP (**Fig. 1-2**), we did not observe a
stronger RNAi signature against SINV-GFP in *M. myotis* cells than in HEK293T cells
(**Fig. 3** and **S1**). This was also true for VSV-GFP (**Fig. 3** and **S1**), even though Dicer
seemed to interfere with VSV-GFP at the RNA and protein level in *M. myotis* cells (**Fig.**
2). In both HEK293T and *M. myotis* cells, only a very low percentage of reads aligned
to the SINV-GFP genome, and even lower to the VSV-GFP genome (**Fig. 3A**).
Focusing on the SINV-GFP reads, while they showed an enrichment at 22nt on both
strands in HEK293T cells, there was an enrichment of smaller size RNAs on the
positive strand in the *M. myotis* cells, suggesting the presence of viral RNA
degradation products (**Fig. 3B** and **S1A**). The same enrichment for smaller size RNAs
on the positive strand could be seen for VSV-GFP reads in *M. myotis* cells, while
almost no reads aligned to the VSV-GFP genome in HEK293T cells (**Fig. 3B** and **S1A**).
The few viral reads that were 22nt long did display typical RNAi characteristics in both
cell lines, as they aligned to both strands and throughout the viral genome (**Fig. 3C**
and **S1B**). In both cell lines there was also an enrichment for paired positive and
negative sense SINV-GFP reads that overlapped with a 2nt overhang, a signature of
Dicer cleavage as part of the RNAi pathway (**Fig. 3D** and **S1C**), although that was not
the case for VSV-GFP reads. However, the very low total number of viral reads of 22nt
in both cell lines argues against a strong antiviral RNAi activity of mmDicer in *M. myotis*
nasal epithelial cells against SINV and VSV. Altogether, our results indicate that
mmDicer does not appear to have increased RNAi activity in *M. myotis* nasal epithelial
cells.

Dicer relocalizes to cytoplasmic foci upon SINV infection in *M. myotis* nasal epithelial cells

To try to understand how mmDicer could play a proviral role in *M. myotis* nasal epithelial cells, we looked at the subcellular localization of Dicer upon SINV infection. In mock conditions, mmDicer exhibited a diffuse cytoplasmic localization, but interestingly upon WT SINV infection we saw a striking relocalization of Dicer into discrete cytoplasmic foci in *M. myotis* nasal epithelial cells. This relocalization could already be noticed at 8h (**Fig. 4A**) and was still apparent at 24h (**Fig. 4B**). There was no significant difference in both the number and size distribution of Dicer foci between 8h and 24h, although there was a small trend towards a decrease in the number of Dicer foci per cell (**Fig. 4C**). Importantly, we did not see any shift in Dicer localization upon SINV infection for HEK293T cells (**Fig. 4A-B**), as well as for the two other human cell lines we tested, human lung epithelial carcinoma A549 cells and human hepatocyte-derived carcinoma Huh7 cells (**Fig. S2A**). Overall, our results suggest a singular activity of Dicer during SINV infection in *M. myotis* nasal epithelial cells that leads to the relocalization of Dicer in large cytoplasmic foci, but that does not appear to occur in human cells.

Dicer likely relocalizes to alphavirus viral factories in *M. myotis* nasal epithelial cells

SINV infected cells are known to accumulate large quantities of double stranded (ds)RNA in the cytoplasm as a result of replication (Griffin, 2007). Since dsRNA is the primary substrate of Dicer, we wondered if Dicer clustered around dsRNA in SINV infected cells. We infected *M. myotis* nasal epithelial cells with SINV and measured the co-localization of Dicer and dsRNA by confocal microscopy, using J2 antibodies that recognize dsRNAs longer than 40bp (Richardson *et al*, 2010). We saw that many of the Dicer foci co-localized with dsRNA in SINV infected cells (**Fig. 5A**), with a high Pearson correlation coefficient across multiple experiments (**Fig. 5B**), suggesting that Dicer forms foci in close proximity to dsRNA in *M. myotis* cells. As control, we could also see dsRNA accumulating in SINV infected HEK293T, A549 and Huh7 cells but Dicer staining remained diffuse and did not co-localize with dsRNA (**Fig. 5A-B, Fig. S2A**). Importantly, infection with the other alphavirus SFV also led to Dicer foci that co-localized with dsRNA (**Fig. 5A-B**). In contrast, infection with WT VSV is known to not accumulate dsRNA by J2 staining (Weber *et al*, 2006). And indeed, we did not detect dsRNA and there was no relocalization of Dicer upon infection in *M. myotis* cells

(**Fig. S2B**). These results suggest that Dicer does not form large foci for every viral
infection but its relocalization may be specific to alphaviruses or linked to dsRNA
accessibility. dsRNA is often formed during viral replication, we thus wondered if Dicer
was present at sites of SINV replication. Since J2 antibodies cannot distinguish
between dsRNA of viral origin or originating from the host, we carried out fluorescence
in situ hybridization (FISH) to specifically stain for SINV plus strand RNA. As expected,
we saw discreet foci showing the accumulation of SINV RNA, likely at sites of viral
replication, in both HEK293T cells and *M. myotis* nasal epithelial cells (**Fig. 5C**). Again,
Dicer stayed diffuse upon infection in HEK293T cells. Importantly, the Dicer foci
matched the SINV plus strand RNA foci in *M. myotis* cells (**Fig. 5C**). Altogether, our
results argue for the recruitment of Dicer to sites of SINV viral replication.
232
<u>*M. myotis* Dicer alone is not sufficient for relocalization in NoDice cells</u>
Finally, we wondered if there was something inherently special about mmDicer that
triggered its relocalization upon SINV infection. To test this, we went back to our
human NoDice cells expressing mmDicer (**Fig. 1**). Immunoprecipitating mmDicer in
these cells showed that mmDicer still interacted with known hDicer interactors such
as PACT and TRBP in mock conditions, as well as PKR and ADAR1, which we
previously showed interact with hDicer during SINV infection (Montavon *et al*, 2021)
(**Fig. 6A**). As expected, while J2 staining showed discrete foci characteristic of SINV
infection, hDicer stayed diffuse throughout the cytoplasm (**Fig. 6B**). Importantly, there
was also no relocalization of mmDicer upon SINV infection (**Fig. 6B**). These results
suggest that mmDicer alone is not sufficient to form large foci, and that factors specific
to *M. myotis* nasal epithelial cells are required for relocalization.
245

Discussion
Recent studies have highlighted a role for Dicer in bat antiviral immunity and tolerance
to viruses through RNAi in two different bat species, *P. alecto* and *T. brasiliensis* (Dai
et al, 2024; Owolabi *et al*, 2025). In this study, we set out to characterize the antiviral
activity of Dicer in *M. myotis* nasal epithelial cells. Interestingly, mmDicer did not
protect from SINV and VSV infection when expressed in a human Dicer knock out cell
line (**Fig. 1**). Moreover, knocking down Dicer in *M. myotis* nasal epithelial cells led to
a decrease in SINV and SFV infection, while results were more mitigated with VSV
(**Fig. 2**). Consistent with these results, we did not find a strong antiviral RNAi signature

when analyzing the viral small RNA profiles of SINV and VSV infected *M. myotis* nasal epithelial cells (**Fig. 3** and **S1**). However, we did observe a striking relocalization of Dicer in large cytoplasmic foci upon SINV and SFV infection in *M. myotis* nasal epithelial cells that we did not witness in human cells (**Fig. 4** and **S2**). These Dicer foci are in close proximity with dsRNA and SINV plus strand RNA, suggesting that Dicer is recruited to sites of SINV replication (**Fig. 5**). Finally, mmDicer does not form these foci when expressed in human Dicer knock out cells, suggesting that its relocalization requires specific factors that are present in *M. myotis* nasal epithelial cells (**Fig. 6**).

Our results that mmDicer does not have increased antiviral RNAi activity is unexpected in view of the recently published results on *P. alecto* and *T. brasiliensis* Dicer, especially as SINV was also used in the *P. alecto* study (Dai *et al*, 2024; Owolabi *et al*, 2025). This discrepancy could come from differences in bat species, as *P. alecto* and *M. myotis* are thought to have diverged more than 60 million years ago, and *T. brasiliensis* and *M. myotis* at least 35 million years ago (Teeling *et al*, 2005; Agnarsson *et al*, 2011; Hao *et al*, 2024). Since in human and mouse cells the antiviral RNAi activity of Dicer can be cell-type dependent, it is also possible that the activity of bat Dicer is influenced by the cell type used in each study, especially if different bat species and cell types have differences in IFN activity (Geng *et al*, 2024; Bondet *et al*, 2021; Clayton & Munir, 2020). In mammalian cells, the same Dicer protein could play a role in antiviral immunity through its RNAi activity, but also through direct regulation of the innate immune reponse and through the maturation of miRNAs that regulate the innate immune response (Gaucherand *et al*, 2024; Maillard *et al*, 2019). These multiple interconnected activities of Dicer make the study of Dicer antiviral activity difficult. We employed a transient siRNA knockdown strategy instead of knocking out Dicer to limit the impact of removing Dicer on the expression of miRNAs, which have a notoriously long half-life (Marzi *et al*, 2016), but there could still be indirect effects of Dicer influencing its propensity to be anti- or proviral in specific cellular contexts. In general, it will be important to study the activity of Dicer in more bat species to resolve this discrepancy.

The case of VSV is interesting, as knocking down Dicer in *M. myotis* cells led to an increase in VSV-GFP RNA and proteins levels (**Fig. 2F-G**). Yet, there was almost no viral siRNAs (**Fig. 3, Fig S1**). The mechanism behind this apparent antiviral activity of Dicer is thus unclear. Nonetheless, it serves as a good control of a virus that does not trigger relocalization of Dicer in *M. myotis* cells, suggesting that this relocalization

is context-specific and not an artefact of infection. Since dsRNA is not detectable upon
VSV infection unlike infection with alphaviruses, it is tempting to hypothesize that the
relocalization of Dicer is linked to dsRNA accessibility. It will be interesting to test this
hypothesis.

Dicer relocalization to discreet cytoplasmic foci upon infection in *M. myotis* cells is
reminiscent of *Drosophila* D2 bodies (Nishida *et al*, 2013). These D2 bodies are
cytoplasmic foci containing Dicer-2 and its co-factor R2D2 that are important for the
processing of endogenous siRNAs in *Drosophila* cells. Yet, our small RNAseq results
did not identify such activity of Dicer. Moreover, we see a proviral effect instead of an
antiviral effect associated with Dicer relocalization, although it remains purely
correlative as we do not have concrete evidence so far that the two are directly linked.
We can nonetheless propose several hypotheses as to how the relocalization of Dicer
could play a proviral role. Potential consequences of this relocalization could be the
sequestration of Dicer by SINV, for example to prevent Dicer activity as part of the
antiviral RNAi pathway. Alternatively, Dicer could be recruited to sites of SINV
replication to hide dsRNA from PRRs or to directly tone down the local innate immune
response. One possible explanation for this recruitment, which remains to be tested,
could be that mmDicer has a higher affinity for dsRNA than its human counterpart in
M. myotis cells, maybe thanks to specific co-factors, but with no increased cleavage
activity. Dicer could also have a structural role to help viral factory formation and
alphavirus replication. More work will be needed to identify the purpose of Dicer
relocalization and the link between this relocalization and Dicer proviral activity.

It is intriguing that this relocalization does not appear to occur in human cells, or
at least in the human cell lines that we have tested. The fact that mmDicer alone is not
sufficient for relocalization is consistent with mmDicer and hDicer being very similar,
with 95% conservation between the two proteins at the amino acid level. Work to
identify the *M. myotis* cell specific factors responsible for Dicer relocalization is
ongoing. Host proteins that only interact with viral proteins in bat cells are good
candidates. On top of finding the mechanism of Dicer relocalization, it will also be
important to determine whether Dicer forms cytoplasmic foci during infection with other
viruses and in other stress conditions. It will especially be important to test viral strains
that are more biologically relevant to *M. myotis*. While many different alphaviruses
have been detected in several species of bats in the wild, including SINV and SFV, it
is still not clear whether bats constitute a reservoir for alphaviruses as this has not

been well studied (Fagre & Kading, 2019). Finally, it would be interesting to determine
if Dicer relocalization occurs in other bat species and cell types, and whether this
relocalization could explain the discrepancy in the observed antiviral RNAi activity of
Dicer.

Materials and methods

<u>Cell culture and viral infections</u>

SV40 transformed *M. myotis* nasal epithelial cells (He *et al*, 2014b) were a kind gift
from Bernd Köllner (FLI, Greifswald, Germany) through Lucie Etienne (CIRI, Lyon,
France). HEK293T/NoDice (2.20) cells (Bogerd *et al*, 2014) were a kind gift from Bryan
Cullen (Duke University, Durham NC, USA). All cells were maintened in Dulbecco's
modified Eagle medium (DMEM, Gibco, Life Technologies) supplemented with 10%
fetal bovine serum (FBS, bioSera FB1090/500, Batch# S00O0) in a humidified
atmosphere of 5% CO_2 at 37°C.
WT SINV (strain AR339) and SINV-GFP (kindly provided by M. C. Saleh, Institut
Pasteur, Paris, France, Saleh et al., 2009) viral stocks were produced and amplified
in BHK21 cells. SFV (strain UVE/SFV/UNK/XX1745; EVAg 001V-02468) was
propagated in Vero E6 cells from the initial stock. WT VSV (strain Indiana isolate PI10)
and VSV-GFP (Mueller *et al*, 2010) were propagated in BHK21 cells from the initial
stock. Stock titrations were performed by plaque assay on VERO E6 cells. Cells were
infected at MOIs ranging from 0.02 to 20, for 8h or 24h, as indicated in the figure
legends.

<u>Plasmids and cloning</u>

pLenti6 Flag-HA-V5 vector was modified from pLenti6-V5 gateway vector (Thermo
Fisher scientific V49610) by Gibson cloning (New England Biolabs) as previously
described (Baldaccini *et al*, 2024), then further modified to remove the V5 tag and
ccdB cassette. plenti6 Flag-HA hDicer was already described (Baldaccini *et al*, 2024).
mmDicer was cloned from *M. myotis* nasal epithelial cells mRNA using Superscript IV
one-step RT-PCR (Thermo Fisher Scientific) with primers Forward
(CAGTGTGGTGGAATTCTGCAGAAAA GCCCTGCTTTGCAACCC) and Reverse
(CCAAACTCATTACTAACCGGTTCAGCTGCT GGGAACCTGAG). plenti6 Flag-HA-
V5 was digested with AgeI and PstI restriction enzymes, then ligated with amplified
mmDicer with matching overhangs by Gibson cloning (New England Biolabs).

Catalytic mutations (E1562A and E1819A) were introduced by PCR inside primers
Forward (GACTGCGTGGCTGCCCTGCTGGGCTGC TATTTAA) + Reverse
(AGCAGGGCAGCCACGCAGTCGGCGATGCTTTTGT) and Forward
(GGGATATTTTTGCTTCACTTGCTGGTGCCATTTAC) + Reverse (GCAAGTGA
AGCAAAAATATCCCCCATGGCCTTG), then ligated to AgeI and PstI digested plenti6
Flag-HA-V5 by Gibson cloning (New England Biolabs).

<u>Lentivirus production and generation of stable cell lines</u>
HEK23T cells were plated in 6 well plates and transfected with 0.33 µg pVSV envelope
plasmid (Addgene #8454) and 1.33 µg pSPAX2 packaging plasmid (Addgene
#12260), and 1.7 µg of either plenti6 Flag-HA Empty, plenti6 Flag-HA hDicer, plenti6
Flag-HA mmDicer or plenti6 Flag-HA mmDicer CM, using Lipofectamine 2000 reagent
(Invitrogen, Fisher Scientific) according to the manufacturer's protocol. Media was
replaced 6h later, then the supernatent containing viral particles was collected at 48h
and filtered through a 0.45 µm PES filter. HEK293T NoDice (Bogerd *et al*, 2014) were
then transduced with the filtered lentiviruses mixed with 4µg/ml polybrene. Efficiently
transduced cells were selected for several days with blasticidine at 15 µg/mL and
polyclonal populations were used for experiments.

<u>Dicer knock down</u>
M. myotis Dicer specific siRNAs were designed by siTOOLs Biotech to form a siPOOL
of 30 siRNAs. A siPOOL of 30 non-targeting siRNAs was used as control (siTOOLs
Biotech). *M. myotis* nasal epithelial cells were reverse transfected with siNeg or siDicer
at 3nM final siRNA concentration using Lipofectamine RNAiMax (ThermoFisher)
following siTOOLs Biotech instructions. Cells were transfected again the following day
with siNeg or siDicer at 3nM final siRNA concentration. The next day, cells were
collected to check for efficient knock down or infected with different viruses.

<u>Live cell imaging</u>
M. myotis nasal epithelial cells were treated with two rounds of siRNAs then infected
with SINV-GFP at MOI 0.2. Cells were then placed into a CellCyte X live cell imaging
system (Cytena) where GFP fluorescence and phase contrast pictures (10X objective)
of every well were taken every 3h for 72h. The percentage of GFP+ area over the total

cell area was quantified with the CellCyte Studio software and averaged from six
photos of 2 individual wells per condition.
392

<u>Protein extraction and western blot</u>
Proteins were harvested and homogenized in appropriate quantity of ice cold lysis
buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 5 mM EDTA, 1% Triton X-100, 0.5%
SDS protease free inhibitor tablet (complete Mini; Sigma Aldrich)). Protein amounts
were quantified using Bradford reagent (Bio-Rad), diluted in lysis buffer + Laemmli and
boiled for 5 min before loading. Proteins were loaded on 10% acrylamide-bis-
acrylamide gels, or 4–20% Mini-PROTEAN TGX Precast Gels (Bio-Rad) for the Dicer
immunoprecipitation experiment. Proteins were separated by migration at 140 V in 1X
Tris-Glycine-SDS buffer (Euromedex), then electro-transferred on a nitrocellulose
membrane (Amersham 0.45 µm) in 1X Tris-Glycine buffer supplemented with 10%
ethanol. Appropriate loading and transfer efficiency were verified by Ponceau S
staining (Merck). Membranes were blocked in 5% milk diluted in PBS-Tween 0.2%.
Membranes were stained with primary antibodies overnight at 4 °C at the following
dilutions: anti-DICER (1:2000, A301-937A, Euromedex, Bethyl), anti-HA-HRP
(1:1000, 12013819001 Merck, Sigma Aldrich), anti-GFP (1:1000, 11814460001
Merck, Sigma Aldrich), anti-SINV nsp2 (1:5000, kind gift from Diane Griffin, Johns
Hopkins University School of Medicine, Baltimore, MD), anti-PKR (1:1000, ab32506
Abcam), anti-PACT (1:500, ab75749 Abcam), anti-TRBP (1:500, sc-514124,
Cliniscience, Santa Cruz), anti-ADAR1 (1:1000, E6X9R #81284 Cell Signaling), anti-
α-Tubulin-HRP (1:10000, Sigma, T6557). After several washes with PBS-Tween 0.2%,
the membranes were stained with the following secondary antibodies coupled to
horseradish peroxidase (HRP): anti-mouse-HRP (1:4000, A4416 Merck, Sigma-
Aldrich) or anti-rabbit-HRP (1:10,000, NA9340V Cytiva, Sigma-Aldrich). Detection was
carried out using SuperSignal West Femto maximum sensitivity substrate (Pierce,
Fisher Scientific) and visualized with a Fusion FX imaging system (Vilber).
418

<u>Plaque assay</u>
For SINV-GFP infections, Vero E6 cells were seeded in 96-well plates and infected
with 10-fold serial dilutions of infection supernatants. After 1 hour, the inoculum was
removed and cells were covered with 2.5% carboxymethyl cellulose and cultured for
72 hours at 37°C in a humidified atmosphere of 5% CO_2. Plaques were counted

manually under the microscope. For SFV and VSV-GFP infections, Vero E6 cells were

seeded in 12-well plates and infected with 10-fold serial dilutions of infection

supernatants. After 1 hour, the inoculum was removed, cells were covered with 1.2%

Avicel overlay (vivapur mcg, JRS Pharma) and cultured for 48 hours at 37°C in a

humidified atmosphere with 5% CO_2. Cells were then washed and fixed with 4%

formaldehyde (Merck, Sigma Aldrich) in PBS (phosphate buffered saline, Gibco) for

20 minutes, then stained with crystal violet for 5 minutes (Sigma-Aldrich). Plates were

washed with water and plaques were counted manually. Viral titers were calculated

according to the formula: PFU/mL = #plaques/ (Dilution*Volume of inoculum).

RNA and small RNA extractions

Total RNA was extracted from cell lysates using Trizol Reagent (15586018, InVitrogen)

to which 200 µl of chloroform was added before centrifugation for 15 minutes at 12,000

xg at 4°C. For regular RNA extraction, the aqueous layer was collected and nucleic

acids were precipitated in an equal volume of isopropanol for 10 min at room

temperature. For small RNA extraction, the aqueous layer was collected and nucleic

acids were precipitated in 3 volumes of 100% ethanol + 10% 1M sodium acetate for

30 minutes at -80°C. In both cases, precipitated RNA was then pelleted for 10 min at

12,000 xg at 4°C then washed with 70% ethanol and reconstituted in RNase free

water.

Northern blot analysis

3 µg of total RNA was loaded onto a 17% urea acrylamide gel and electro separated

in a 1X Tris-Borate-EDTA solution. The RNA was then electro transferred to a Hybond-

NX membrane (GE Healthcare), and chemically cross linked to the membrane with 1-

ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC) (Sigma Aldrich) for

90 min at 65°C. Membranes were then pre-hybridized for 30 min with PerfectHyb plus

(Sigma Aldrich) at 50°C. Probes consisting of an oligodeoxyribonucleotide sequence

for which the 5′- was labelled with 25 µCi of [γ-32P]dATP using T4 polynucleotide

kinase (Thermo Fisher Scientific) were incubated overnight with the membranes at

50°C. The membranes were then washed twice at 50°C for 10 min with 5X SSC/0.1%

SDS, then for 5 min with 1X SSC/0.1% SDS. Finally, membranes were placed in

contact with a phosphorimager plate overnight and imaged with a Bioimager FLA-7000

scanner (Fuji). Probe sequences: miR-16 (CGCCAATATTTACGTGCTGCTA), U6 (GCAGGGGCCATGCTAATCTTCTCTGT ATCG).

RT-qPCR analysis

After RNA extraction, samples were incubated with DNase I (Invitrogen, Fisher Scientific) for 20 min at 37°C. The RNA was then reverse-transcribed into cDNA using SuperScript IV (SSIV) reverse transcriptase (Invitrogen, Fisher Scientific) or AMV reverse transcriptase (Promega) with a random nonameric primer, following the manufacturer's protocol. Quantitative real-time PCR was performed using SYBR Green PCR Master Mix (Fisher Scientific) and the following primers: 18S F (CTCTTAGCTGAGTGTCCCGC), 18S R (CTTAATCATGGCCTCAGTTCCGA), SINV F (CCACTACGCAAGCAGAGACG), SINV R (AGTGCCCAGGGCCTGTGTCCG), VSV F (TGTACAATCAGGCGAGAGTC), VSV R (GGAACTCATGATGAATGGATTGGG).

Small RNA sequencing and bioinformatic analysis

After small RNA extraction, samples were ran on a Bioanalyzer chip to confirm RNA integrity. Samples were then shipped to BGI for library preparation with unique molecular identifiers, and sequenced as single-end 50 base reads on their DNBSEQ sequencing technology platform with at least 20 million reads per sample.

Raw reads were pre-cleaned by the sequencing platform, including adapter, contaminant and low-quality removal (Phred+33). Before alignment, redundant sequences were collapsed to one copy using fastq_collapser from FASTX-Toolkit 0.0.14 (http://hannonlab.cshl.edu/fastx_toolkit/) and a read-size filter was applied (18-35 bp). HEK293T and *M. myotis* short reads were then aligned to the human genome assembly GRCh38 (GCF_000001405.40) and the *M. myotis* genome mMyoMyo1.p (GCF_014108235.1), respectively. Mapping was performed using Bowtie 1.3.1 (Langmead *et al*, 2009) with parameters '-v 1 -a --best --strata'. Unmapped reads were next separately aligned against virus genomes SINV-GFP (private sequence derived from NC_001547.1 – RefSeq database) and VSV-GFP (FJ478454.1) using the following parameters '-v 2 -y -q -m 30 -a --best --strata'. Finally, siRNA signature was examined on 22nt mapped reads by the python tool signature by (Antoniewski, 2014) with options '22 22 1 22'. Sequencing data have been deposited on NCBI's Gene

Expression Omnibus (Edgar *et al*, 2002) and are accessible through GEO Series
accession number GSE301457.

<u>Immunostaining</u>
Cells were seeded onto Labtek eight-chamber slides (Nunc Lab Tek Chamber Slide
System) at a density of 8,000 cells per well. The next day, cells were infected for 8h or
24h with WT SINV, WT VSV or SFV. Cells were then fixed for 10 minutes at room
temperature with 4% formaldehyde (Sigma-Aldrich) diluted in 1X PBS. Blocking was
carried out with 5% normal goat serum in 0.1% Triton X-100 in 1X PBS for 1 hour at
room temperature. Primary antibody staining was carried out for 3h at room
temperature using the following antibodies diluted in blocking buffer: anti-Dicer
(1:1000, A301-937A, polyclonal, Bethyl) and J2 anti-dsRNA (1:1000, RNT-SCI-
10010200, Jena Bioscience). The cells were then incubated for 1 hour at room
temperature with secondary antibodies goat anti-mouse Alexa 594 (A11032,
Invitrogen) or goat anti-rabbit Alexa 488 (A11008, Invitrogen), both diluted at 1:1000
in 1X PBS with 0.1% Triton X-100. Between each step, cells were washed thoroughly
with 1X PBS containing 0.1% Triton X-100. Nuclei were stained with DAPI (4',6-
diamidino-2-phenylindole, dichlorhydrate, D1306, Thermo Fisher) at 1:5000 in 1X PBS
for 5 minutes. Finally, the slides were mounted with coverslips using Fluoromount-G
mounting media (Southern Biotech) and observed under a confocal microscope
(LSM780, Zeiss).

<u>Sequential immunostaining and FISH</u>
Mock- or WT SINV-infected cells were cultured on 18 mm round coverslips placed in
12-well plates. Cells were fixed with 3,7% formaldehyde (Sigma-Aldrich) diluted in 1X
PBS for 10 minutes at room temperature. Following fixation, cells were permeabilized
with 0.1% Triton X-100 in 1X PBS for 5 minutes at room temperature. Immunostaining
was performed as described in the previous section but with 1h incubation steps for
primary and secondary antibody staining. Subsequently, cells were fixed again with
3.7% formaldehyde (Sigma-Aldrich) diluted in 1X PBS for 10 minutes at room
temperature. For RNA detection, cells were incubated overnight at room temperature
with SINV genome-specific Stellaris Cal Fluor Red 610 RNA FISH Probes (LGC
Biosearch Technologies, custom made by Erika Girardi (Messmer *et al*, 2024)), kindly
provided by Erika Girardi (IBMC, Strasbourg, France). Probes were diluted in RNA

FISH hybridization buffer (Stellaris, Biosearch Technologies). Nuclei were stained with DAPI (D1306, Invitrogen, Thermo Fisher Scientific) for 30 minutes. Finally, coverslips were mounted using Fluoromount-G mounting medium (Invitrogen, Thermo Fisher Scientific), and the samples were imaged by fluorescence microscopy (Olympus BX51).

Image analysis

All image analyses were performed on confocal images and using the Fiji software (Schindelin *et al*, 2012). Detection thresholds and all parameters were standardized between the different images and cell lines to ensure consistency in the analysis. Each analysis was performed on two (SFV) or three (SINV) biological replicates, with each field containing approximately the same number of cells. The "Plot profile" tool on Fiji was used to visualize fluorescent intensity as a function of distance for each channel. The selection of the line or field of observation was made arbitrarily to best represent the whole slide. Threshold parameters were adjusted accordingly to measure the size of various objects, and the minimum size accepted was set to 0.1 µm² to limit background detection. The "Analyse particle" tool was used to extract numerical data, which was then systematically analyzed. The colocalization analysis between two fluorescent signals was performed using the "Colocalization Finder" tool in Fiji. This tool analyzes colocalization on a pixel-by-pixel basis by comparing the intensity of each pixel in one channel to the corresponding pixel in the second channel of a bi-color image. Importantly, the thresholds or brightness levels in the image are not changed during this process. The analysis generates a scatter plot, from which a Pearson correlation coefficient is calculated. A value of +1 indicates perfect correlation, 0 indicates no correlation, and -1 signifies perfect anti-correlation.

Dicer immunoprecipitation

NoDice cells expressing different HA-tagged Dicer constructs were plated and mock-infected or infected with SINV-GFP at MOI 0.02. 24 hours later, cells were harvested, washed twice with ice-cold 1X PBS (Gibco, Life Technologies), and resuspended in lysis buffer (50 mM Tris-HCl pH 7.5, 140 mM NaCl, 6 mM MgCl2, 0.1% NP-40), supplemented with a Complete-EDTA-free Protease Inhibitor Cocktail tablet (complete Mini; Sigma Aldrich). Cells were lysed on ice for 10 min and debris were removed by 10 min centrifugation at 12,000 *xg* at 4°C. An aliquot of the cleared lysates was kept

aside as protein input. The rest of the samples were divided in two equal volumes and
incubated with 40 µL magnetic microparticles coated with monoclonal anti-HA or anti-
MYC antibodies (MACS purification system, Miltenyi Biotech) at 4°C for 1 hour with
constant rotation. Next, samples were passed through µ Columns (MACS purification
system, Miltenyi Biotech) and washed four times with 200 µL lysis buffer. Finally,
proteins were eluted with 2X Laemmli buffer (10% glycerol, 4% SDS, 62.5 mM TrisHCl
pH 6.8, 5% (v/v) 2-β-mercaptoethanol, Bromophenol Blue) that was pre-warmed to
95℃.

Acknowledgements
We would like to thank members of the laboratory for insightful discussions and
suggestions. This work of the Interdisciplinary Thematic Institute IMCbio+, as part of
the ITI 2021-2028 program of the University of Strasbourg, CNRS and Inserm, was
supported by IdEx Unistra (ANR-10-IDEX-0002), by SFRI-STRAT'US project (ANR-
20-SFRI-0012), EUR IMCBio (IMCBio ANR-17-EURE-0023) under the framework of
the French Investments for the Future Program. It was also supported by an Agence
Nationale pour la Recherche PRCI grant to SP (ANR-21-CE35-0018-01). LG was
funded by a postdoctoral fellowship from the Fondation pour la Recherche Médicale
(funding number SPF202209015746).

References
Agnarsson I, Zambrana-Torrelio CM, Flores-Saldana NP & May-Collado LJ (2011) A
time-calibrated species-level phylogeny of bats (Chiroptera, Mammalia). *PLoS*
Currents 3: RRN1212
Antoniewski C (2014) Computing siRNA and piRNA Overlap Signatures. In *Animal*
Endo-SiRNAs: Methods and Protocols, Werner A (ed) pp 135–146. New York, NY:
Springer
Baldaccini M, Gaucherand L, Chane-Woon-Ming B, Messmer M, Gucciardi F & Pfeffer
S (2024) The helicase domain of human Dicer prevents RNAi-independent
activation of antiviral and inflammatory pathways. *EMBO J* 43: 806–835
Baldaccini M & Pfeffer S (2021) Untangling the roles of RNA helicases in antiviral
innate immunity. *PLOS Pathogens* 17: e1010072
Banerjee A, Baker ML, Kulcsar K, Misra V, Plowright R & Mossman K (2020) Novel
Insights Into Immune Systems of Bats. *Frontiers in Immunology* 11

Bartel DP (2018) Metazoan MicroRNAs. *Cell* 173: 20–51

Bogerd HP, Whisnant AW, Kennedy EM, Flores O & Cullen BR (2014) Derivation and characterization of Dicer- and microRNA-deficient human cells. *RNA* 20: 923–937

Bondet V, Le Baut M, Le Poder S, Lécu A, Petit T, Wedlarski R, Duffy D & Le Roux D (2021) Constitutive IFNα Protein Production in Bats. *Front Immunol* 12: 735866

Burgin CJ, Colella JP, Kahn PL & Upham NS (2018) How many species of mammals are there? *Journal of Mammalogy* 99: 1–14

Chua KB, Lek Koh C, Hooi PS, Wee KF, Khong JH, Chua BH, Chan YP, Lim ME & Lam SK (2002) Isolation of Nipah virus from Malaysian Island flying-foxes. *Microbes and Infection* 4: 145–151

Clayton E & Munir M (2020) Fundamental Characteristics of Bat Interferon Systems. *Front Cell Infect Microbiol* 10: 527921

Cullen BR, Cherry S & tenOever BR (2013) Is RNA Interference a Physiologically Relevant Innate Antiviral Immune Response in Mammals? *Cell Host & Microbe* 14: 374–378

Dai Y, Wang B, Wang J, Wei X, Liu X, Che X, Li J, Lun Ng W, Wang L-F & Li Y (2024) Increased viral tolerance mediates by antiviral RNA interference in bat cells. *Cell Reports* 43: 114581

Edgar R, Domrachev M & Lash AE (2002) Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res* 30: 207–210

Fagre AC & Kading RC (2019) Can Bats Serve as Reservoirs for Arboviruses? *Viruses* 11: 215

Flemr M, Malik R, Franke V, Nejepinska J, Sedlacek R, Vlahovicek K & Svoboda P (2013) A retrotransposon-driven dicer isoform directs endogenous small interfering RNA production in mouse oocytes. *Cell* 155: 807–816

Gaucherand L, Baldaccini M & Pfeffer S (2024) Beyond RNAi: How the Dicer protein modulates the antiviral innate immune response in mammalian cells. *Bioessays*: e2400173

Geng R, Wang Q, Yao Y-L, Shen X-R, Jia J-K, Wang X, Zhu Y, Li Q, Shi Z-L & Zhou P (2024) Unconventional IFNω-like Genes Dominate the Type I IFN Locus and the Constitutive Antiviral Responses in Bats. *The Journal of Immunology*: ji2300301

Girardi E, Chane-Woon-Ming B, Messmer M, Kaukinen P & Pfeffer S (2013)
Identification of RNase L-dependent, 3'-end-modified, viral small RNAs in Sindbis
virus-infected mammalian cells. *MBio* 4: e00698-00613

Gonzalez V & Banerjee A (2022) Molecular, ecological, and behavioral drivers of the
bat-virus relationship. *iScience* 25: 104779

Griffin D (2007) Alphaviruses, p 1023–1067. In *Fields virology, 5th ed. Lippincott*
Williams & Wilkins, Philadelphia, PA

Guo Z, Li Y & Ding S-W (2019) Small RNA-based antimicrobial immunity. *Nat Rev*
Immunol 19: 31–44

Gurung C, Fendereski M, Sapkota K, Guo J, Huang F & Guo Y-L (2021) Dicer
represses the interferon response and the double-stranded RNA-activated protein
kinase pathway in mouse embryonic stem cells. *Journal of Biological Chemistry*
296: 100264

Halpin K, Young PL, Field HE & Mackenzie JS (2000) Isolation of Hendra virus from
pteropid bats: a natural reservoir of Hendra virus. *Journal of General Virology* 81:
1927–1932

Hao X, Lu Q & Zhao H (2024) A molecular phylogeny for all 21 families within
Chiroptera (bats). *Integrative Zoology* 19: 989–998

He X, Korytář T, Schatz J, Freuling CM, Müller T & Köllner B (2014a) Anti-Lyssaviral
Activity of Interferons κ and ω from the Serotine Bat, Eptesicus serotinus. *Journal*
of Virology 88: 5444

He X, Korytář T, Zhu Y, Pikula J, Bandouchova H, Zukal J & Köllner B (2014b)
Establishment of Myotis myotis cell lines--model for investigation of host-pathogen
interaction in a natural host for emerging viruses. *PLoS One* 9: e109795

Irving AT, Ahn M, Goh G, Anderson DE & Wang L-F (2021) Lessons from the host
defences of bats, a unique viral reservoir. *Nature* 589: 363–370

Kulmann MIR, Taborska E, Benköova B, Palus M, Drobek A, Horvat F, Pasulka J, Malik
R, Salyova E, Hönig V, *et al* (2025) Enhanced RNAi does not provide efficient
innate antiviral immunity in mice. *Nucleic Acids Res* 53: gkae1288

Langmead B, Trapnell C, Pop M & Salzberg SL (2009) Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol* 10: R25

Lee HY, Zhou K, Smith AM, Noland CL & Doudna JA (2013) Differential roles of human
Dicer-binding proteins TRBP and PACT in small RNA processing. *Nucleic Acids*
Res 41: 6568–6576

Lei M & Dong D (2016) Phylogenomic analyses of bat subordinal relationships based
on transcriptome data. *Sci Rep* 6: 27726

Li W, Shi Z, Yu M, Ren W, Smith C, Epstein JH, Wang H, Crameri G, Hu Z, Zhang H,
et al (2005) Bats Are Natural Reservoirs of SARS-Like Coronaviruses. *Science*
310: 676–679

López P, Girardi E, Mounce BC, Weiss A, Chane-Woon-Ming B, Messmer M, Kaukinen
P, Kopp A, Bortolamiol-Becet D, Fendri A, *et al* (2020) High-Throughput
Fluorescence-Based Screen Identifies the Neuronal MicroRNA miR-124 as a
Positive Regulator of Alphavirus Infection. *J Virol* 94

Maillard PV, Van der Veen AG, Deddouche-Grass S, Rogers NC, Merits A & Reis e
Sousa C (2016) Inactivation of the type I interferon pathway reveals long double-
stranded RNA-mediated RNA interference in mammalian cells. *EMBO J* 35: 2505–
2518

Maillard PV, van der Veen AG, Poirier EZ & Reis e Sousa C (2019) Slicing and dicing
viruses: antiviral RNA interference in mammals. *EMBO J* 38

Marzi MJ, Ghini F, Cerruti B, Pretis S de, Bonetti P, Giacomelli C, Gorski MG, Kress T,
Pelizzola M, Muller H, *et al* (2016) Degradation dynamics of microRNAs revealed
by a novel pulse-chase approach. *Genome Res*: gr.198788.115

Memish ZA, Mishra N, Olival KJ, Fagbo SF, Kapoor V, Epstein JH, AlHakeem R,
Durosinloun A, Al Asmari M, Islam A, *et al* (2013) Middle East Respiratory
Syndrome Coronavirus in Bats, Saudi Arabia. *Emerg Infect Dis* 19: 1819–1823

Messmer M, Pierson L, Pasquier C, Djordjevic N, Chicher J, Hammann P, Pfeffer S &
Girardi E (2024) DEAD box RNA helicase 5 is a new pro-viral host factor for
Sindbis virus infection. *Virol J* 21: 76

Montavon TC, Baldaccini M, Lefèvre M, Girardi E, Chane-Woon-Ming B, Messmer M,
Hammann P, Chicher J & Pfeffer S (2021) Human DICER helicase domain recruits
PKR and modulates its antiviral activity. *PLoS Pathog* 17: e1009549

Moratelli R & Calisher CH (2015) Bats and zoonotic viruses: can we confidently link
bats with emerging deadly viruses? *Mem Inst Oswaldo Cruz* 110: 1–22

Mueller S, Gausson V, Vodovar N, Deddouche S, Troxler L, Perot J, Pfeffer S,
Hoffmann JA, Saleh M-C & Imler J-L (2010) RNAi-mediated immunity provides
strong protection against the negative-strand RNA vesicular stomatitis virus in
Drosophila. *Proc Natl Acad Sci USA* 107: 19390–19395

Nishida KM, Miyoshi K, Ogino A, Miyoshi T, Siomi H & Siomi MC (2013) Roles of R2D2, a cytoplasmic D2 body component, in the endogenous siRNA pathway in Drosophila. *Mol Cell* 49: 680–691

Owolabi IJ, Karim S-U, Khanal S, Valdivia S, Frenzel C, Bai F & Flynt AS (2025) Processing of genomic RNAs by Dicer in bat cells limits SARS-CoV-2 replication. *Virology Journal* 22: 86

Pei G, Balkema-Buschmann A & Dorhoi A (2024) Disease tolerance as immune defense strategy in bats: One size fits all? *PLOS Pathogens* 20: e1012471

Pichlmair A & Reis e Sousa C (2007) Innate recognition of viruses. *Immunity* 27: 370–383

Poirier EZ, Buck MD, Chakravarty P, Carvalho J, Frederico B, Cardoso A, Healy L, Ulferts R, Beale R & Reis e Sousa C (2021) An isoform of Dicer protects mammalian stem cells against multiple RNA viruses. *Science* 373: 231–236

Richardson SJ, Willcox A, Hilton DA, Tauriainen S, Hyoty H, Bone AJ, Foulis AK & Morgan NG (2010) Use of antisera directed against dsRNA to detect viral infections in formalin-fixed paraffin-embedded tissue. *J Clin Virol* 49: 180–185

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, *et al* (2012) Fiji: an open-source platform for biological-image analysis. *Nat Methods* 9: 676–682

Teeling EC, Springer MS, Madsen O, Bates P, O'Brien SJ & Murphy WJ (2005) A Molecular Phylogeny for Bats Illuminates Biogeography and the Fossil Record. *Science* 307: 580–584

Towner JS, Pourrut X, Albariño CG, Nkogue CN, Bird BH, Grard G, Ksiazek TG, Gonzalez J-P, Nichol ST & Leroy EM (2007) Marburg Virus Infection Detected in a Common African Bat. *PLOS ONE* 2: e764

van der Veen AG, Maillard PV, Schmidt JM, Lee SA, Deddouche-Grass S, Borg A, Kjær S, Snijders AP & Reis e Sousa C (2018) The RIG-I-like receptor LGP2 inhibits Dicer-dependent processing of long double-stranded RNA and blocks RNA interference in mammalian cells. *EMBO J* 37: e97479

Weber F, Wagner V, Rasmussen SB, Hartmann R & Paludan SR (2006) Double-Stranded RNA Is Produced by Positive-Strand RNA Viruses and DNA Viruses but Not in Detectable Amounts by Negative-Strand RNA Viruses. *J Virol* 80: 5059–5064

724   Yob JM, Field H, Rashdi AM, Morrissy C, van der Heide B, Rota P, bin Adzhar A, White

725   J, Daniels P, Jamaluddin A, *et al* (2001) Nipah virus infection in bats (order

726   Chiroptera) in peninsular Malaysia. *Emerg Infect Dis* 7: 439–441

 Figure legends

 Figure 1: Expressing *M. myotis* Dicer in human NoDice cells does not protect

from SINV and VSV infection. (A) Expression of HA-tagged Dicer was analyzed by western blot for NoDice cells expressing HA-tagged hDicer, mmDicer, mmDicer CM or an empty vector control. α-Tubulin was used as loading control. Images are representative of 3 independent experiments **(B)** Expression of miR-16 was analyzed by northern blot for NoDice cells expressing HA-tagged hDicer, mmDicer, mmDicer CM or an empty vector control. A probe for U6 snRNP was used as loading control. Images are representative of 2 independent experiments. **(C-F)** NoDice cells expressing HA-tagged hDicer, mmDicer, mmDicer CM or an empty vector control were infected with SINV-GFP **(C-E)** or VSV-GFP **(F)** at MOI 0.02 for 24h. **(C, F)** Supernatants were collected and viral titers were quantified by plaque assay. Means +/- standard deviation are plotted for 4 **(C)** or 3 **(F)** independent experiments. **(D)** Protein lysates were collected and analyzed by western blot using antibodies for SINV nsp2 and GFP. α-Tubulin was used as loading control. n = 3 independent experiments. Quantification of GFP or nsp2 band intensities relative to Tubulin band intensities were quantified with ImageJ and the mean +/- standard deviation was plotted relative to empty vector control for all three independent experiments. **(E)** RNAs were purified and levels of SINV RNA were quantified by qRT-PCR. Means +/- standard deviation normalized to 18S are plotted relative to empty vector control for 4 independent experiments. For all graphs, p values were calculated using standard one-way ANOVA with Dunnett's multiple comparison test. ns: not significant.

Figure 2: Dicer is proviral for SINV and SFV in *M. myotis* nasal epithelial cells.
(A) Expression of Dicer was analyzed by western blot in *M. myotis* nasal epithelial cells after two rounds of siRNA targeting Dicer (siDicer) or a non-targeting control (siNeg). α-Tubulin was used as loading control. Images are representative of 2 independent experiments. **(B-J)** *M. myotis* nasal epithelial cells treated with siRNAs targeting Dicer (siDicer) or a non-targeting control (siNeg) were infected with SINV-GFP for 24h at MOI 0.2 **(B-E)**, SFV for 24h at MOI 20 **(F-G)** or VSV-GFP for 8h at MOI 1 **(H-J)**. **(B)** The number of GFP+ cells as a proxy for infection were monitored over time by live microscopy imaging. Percentages of GFP+ area were plotted for each condition as the average of two wells, six photos per well, for one experiment. **(C, F, H)** Supernatants were collected and viral titers were quantified by plaque assay.

Means +/- standard deviation are plotted for 3 independent experiments. **(D, I)** Protein

lysates were collected and analyzed by western blot using antibodies for GFP. α-

Tubulin was used as loading control. n = 3 independent experiments. Quantification of

GFP band intensities relative to Tubulin band intensities were quantified with ImageJ

and the mean +/- standard deviation was plotted relative to siNeg mock samples for

all three independent experiments. **(E, G, J)** RNAs were purified and levels of SINV

RNA **(E)**, SFV RNA **(G)** or VSV RNA **(J)** were quantified by qRT-PCR. Means +/-

standard deviation are plotted relative to siNeg infected samples for 3 independent

experiments. For all graphs, p values were calculated using unpaired t test. * p<0.5 ;

ns: not significant.

Figure 3: No strong RNAi signature in *M. myotis* nasal epithelial cells. HEK293T

or *M. myotis* nasal epithelial cells were mock infected or infected with SINV-GFP or

VSV-GFP, then small RNAs were extracted and sequenced. n = 2 independent

experiments, see figure S1 for data from other replicate. **(A)** Percent of total

sequencing reads that align to the human or *M. myotis* genome (black), the SINV-GFP

genome (blue) or the VSV-GFP genome (red) for each sample in HEK293T cells (left)

or *M. myotis* nasal epithelial cells (right). "rep" indicates the replicate number. **(B)**

Number of reads that align to SINV-GFP or VSV-GFP for each cell type based on the

read size. The total number of reads is further broken down into colors based on the

identity of the first nucleotide of the read: yellow for U, blue for A, purple for G and

black for C. **(C)** Location of the 22 nucleotide (nt) reads along the SINV-GFP or VSV-

GFP genome for each sample. The number of reads falling on the same nucleotide is

represented in red if the reads align to the genome (+ strand) and in blue if they align

to the anti-genome (- strand). **(D)** For each sample, 22 nt small RNA pairs that overlap

on + and - strands were analyzed and associated z-scores (Antoniewski, 2014) were

plotted for the indicated nucleotide overlaps.

Figure 4: Dicer relocalizes to distinct cytoplasmic puncta upon infection in *M.*

***myotis* nasal epithelial cells. (A-C)** HEK293T were mock infected or infected with

WT SINV for 8h at MOI 0.2 **(A)** or 24h at MOI 0.02 **(B)**. *M. myotis* nasal epithelial cells

were mock infected or infected with WT SINV for 8h at MOI 5 **(A)** or 24h at MOI 2 **(B)**.

Localization of Dicer (yellow) was imaged by immunofluorescence and confocal

microscopy. DAPI staining (blue) indicates cell nuclei. Images are representative of at

least 3 independent experiments. Scale bars 10µm. **(C)** For each condition, the

number and size of Dicer foci in SINV infected *M. myotis* nasal epithelial cells were

plotted for 3 or 4 independent experiments with each 1 or 2 fields containing

approximately the same number of cells. The black horizontal bars represent the

median. The p value was calculated using an unpaired t test.

Figure 5: Dicer likely relocalizes to SINV viral factories in *M. myotis* nasal

epithelial cells. (A-B) HEK293T were mock infected or infected with WT SINV or SFV

for 24h at MOI 0.02. *M. myotis* nasal epithelial cells were mock infected or infected

with WT SINV or SFV for 24h at MOI 2 or 5 respectively. Localization of Dicer (yellow)

and dsRNA via J2 antibodies (magenta) was imaged by immunofluorescence and

confocal microscopy. DAPI staining (blue) indicates cell nuclei. Images are

representative of 3 (SINV) or 2 (SFV) independent experiments. Scale bars 10µm. Co-

localization between Dicer and dsRNA signals was quantified and Pearson correlation

R coefficients were calculated and plotted for 3 (SINV) or 2 (SFV) independent

experiments – mean (numerical value indicated in gray font above each bar) +/-

standard deviation **(B)**. A zoomed-in image of the boxed area in some merge images

was added. The graphs show the signals for each channel along the line drawn in the

zoomed in images. **(C)** HEK293T or *M. myotis* nasal epithelial cells were mock infected

or infected with WT SINV for 24h at MOI 0.02 or 4, respectively. Localization of Dicer

(yellow) by immunofluorescence and of SINV plus strand (+) RNA (magenta) by FISH

was imaged with a fluorescent microscope. DAPI staining (blue) indicates cell nuclei.

Images are representative of 2 independent experiments. Scale bars 10µm.

Figure 6: *M. myotis* Dicer alone is not sufficient for relocalization. HEK293T

NoDice cells expressing HA-tagged hDicer or mmDicer were mock infected or infected

with SINV-GFP for 24h at MOI 0.02. **(A)** Cells were then lysed, a small aliquot was

kept as input, and Dicer was immunoprecipitated with anti-HA coated magnetic beads

(IP Dicer) or anti-myc coated beads (IP ctrl). Eluted proteins were analyzed by western

blot using antibodies for the known hDicer interactors ADAR1, PKR, TRBP and PACT.

Anti-HA antibodies show efficient immunoprecipitation of hDicer and mmDicer. Images

representative of 2 independent experiments. **(B)** Localization of Dicer (yellow) and

dsRNA via J2 antibodies (magenta) was imaged by immunofluorescence and confocal

microscopy. DAPI staining (blue) indicates cell nuclei. Images are representative of 2
independent experiments. Scale bars 10μm.

Figure S1: Small RNAseq results for the rest of HEK293T and *M. myotis* nasal
epithelial samples. HEK293T or *M. myotis* nasal epithelial cells were mock infected
or infected with SINV-GFP or VSV-GFP, then small RNAs were extracted and
sequenced. n = 2 independent experiments, see figure 3 for data from other replicate.
(A) Number of reads that align to SINV-GFP or VSV-GFP for each cell type based on
the read size. The total number of reads is further broken down into colors based on
the identity of the first nucleotide of the read: yellow for U, blue for A, purple for G and
black for C. **(B)** Location of the 22 nucleotide (nt) reads along the SINV-GFP or VSV-
GFP genome for each sample. The number of reads falling on the same nucleotide is
represented in red if the reads align to the genome (+ strand) and in blue if they align
to the anti-genome (- strand). **(C)** For each sample, 22 nt small RNA pairs that overlap
on + and - strands were analyzed and associated z-scores (Antoniewski, 2014) were
plotted for the indicated nucleotide overlaps.

Figure S2: No Dicer foci are found in other human cell lines and in VSV infected
cells. (A) Human A549 and Huh7 cells were mock infected or infected with WT SINV
at 24h at MOI 0.2 and 0.02, respectively. Localization of Dicer (yellow) and dsRNA via
J2 antibodies (magenta) were imaged by immunofluorescence and confocal
microscopy. DAPI staining (blue) indicates cell nuclei. Images are representative of 2
independent experiments. Scale bars 10μm. **(B)** HEK293T or *M. myotis* nasal
epithelial cells were infected with WT VSV for 8h at MOI 1. Localization of Dicer
(yellow) and dsRNA via J2 antibodies (magenta) were imaged by immunofluorescence
and confocal microscopy. DAPI staining (blue) indicates cell nuclei. Images are
representative of 2 independent experiments.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

Figure S2