The ORB2 RNA-binding protein represses translation of its target transcripts during the

Drosophila maternal-to-zygotic transition via its functionally conserved Zinc-binding 'ZZ'

domain

Timothy C.H. Low[1], Hua Luo[1], Chalini Weerasooriya[2], Sichun Lin[3], Stephane Angers[2,3], Craig

A. Smibert[1,2], Howard D. Lipshitz[1]*

[1]Department of Molecular Genetics, University of Toronto, 661 University Avenue, Toronto

Ontario, Canada M5G 1M1

[2]Department of Biochemistry, University of Toronto, 661 University Avenue, Toronto Ontario,

Canada M5G 1M1

[3]Department of Pharmaceutical Sciences & Donnelly Centre, University of Toronto, 160 College

Street, Toronto, Ontario, Canada M5S 3E1

*Author for correspondence: howard.lipshitz@utoronto.ca

Running head: ORB2 represses mRNAs during the MZT

Keywords: ORB2, Cytoplasmic Polyadenylation Element Binding Protein (CPEB), RNA-

binding protein (RBP), translational regulation, post-transcriptional regulation, pre-initiation

complex (PIC), embryo, maternal-to-zygotic transition (MZT), S2 cell, Cup.

ARTICLE SUMMARY

We show that Drosophila ORB2, the ortholog of the human CPEB2 RNA-binding protein, represses translation of its target mRNAs during the maternal-to-zygotic transition via its C-terminal Zinc-binding ('ZZ') domain.

ABSTRACT

RNA-binding proteins (RBPs) are key components of the post-transcriptional regulatory machinery. We show that the ORB2 RBP, the Drosophila ortholog of human Cytoplasmic Polyadenylation Element Binding Protein (hCPEB) 2-4 protein family, binds to hundreds of maternally provided mRNAs in early embryos, identify a U-rich motif enriched in ORB2's targets, and show that this motif confers ORB2 binding and repression to a luciferase reporter mRNA in S2 tissue culture cells. ORB2's target transcripts are translationally repressed and unstable during the maternal-to-zygotic transition (MZT), a developmental phase during which a large proportion of maternally provided mRNAs are repressed and cleared. We show that, when tethered to a luciferase reporter, ORB2 and hCPEB2 (but not ORB and hCPEB1) repress translation and that the C-terminal Zinc-binding ('ZZ') domain of ORB2 is necessary and sufficient for repression. ORB2 interacts with a suite of post-transcriptional regulators in early embryos; a subset of these interactions is lost upon deletion of the ZZ domain, notably with the Cup repressive complex. Analysis of the early embryo's translatome in the presence or absence of the endogenous ZZ domain, shows that ORB2's targets move onto polysomes upon ZZ domain deletion, indicating that this domain mediates translational repression of ORB2's targets during the MZT. Together, our results assign a function to the ZZ domain and support a significant role for ORB2 in post-transcriptional regulation of maternal mRNAs during the Drosophila MZT.

INTRODUCTION

Proteins belonging to the Cytoplasmic Polyadenylation Element Binding (CPEB) family are highly conserved RNA-binding proteins (RBPs) that canonically interact with U-rich motifs, known as cytoplasmic polyadenylation elements (CPEs), which are typically found in the 3′UTR of target transcripts (HAKE AND RICHTER 1994). Vertebrates have four CPEBs (CPEB1-4), while flies have two (ORB and ORB2). The CPEBs can be divided into two groups based on sequence

similarity: the CPEB1-like subfamily (containing vertebrate CPEB1 and its homologs, such as Drosophila ORB) and the CPEB2-like subfamily (containing vertebrate CPEB2-4 and their homologs, such as Drosophila ORB2) (WANG AND COOPER 2009; FERNANDEZ-MIRANDA AND MENDEZ 2012; IVSHINA *et al.* 2014; DURAN-ARQUÉ *et al.* 2022).

CPEB proteins have been shown to function in the regulation of cytoplasmic mRNAs involved in egg maturation, embryonic stem cell division, memory formation, synaptic plasticity, and gametogenesis (HAKE AND RICHTER 1994; GROISMAN *et al.* 2001; KELEMAN *et al.* 2007; STANDART AND MINSHALL 2008; KRÜTTNER *et al.* 2012; XU *et al.* 2012). Xenopus CPEB1, for example, is a major specificity factor of an mRNA masking complex that represses translation in oocytes and mediates translational activation of its target mRNAs upon egg activation (HAKE AND RICHTER 1994; KIM AND RICHTER 2006).

Drosophila has a single ortholog of CPEB2-4, which is encoded by the *orb2* gene and has been shown to have a role in neural development, long-term memory formation, and spermatogenesis (KELEMAN *et al.* 2007; HAFER *et al.* 2011; MAJUMDAR *et al.* 2012; XU *et al.* 2012; KHAN *et al.* 2015; HERVÁS *et al.* 2016). The *orb2* gene encodes two protein isoforms, ORB2A and ORB2B, which have 542 amino acids in common (from N-to-C): a poly-glutamine (polyQ) tract, a predicted intrinsically disordered region (IDR2) that spans approximately 200 amino acids, the RNA-binding domain (RBD), which consists of tandem RNA recognition motif (RRM) domains; and a ZZ-class Zinc-binding domain (ORB2B is diagrammed in Figure 1a). ORB2A and ORB2B differ only in IDR1 at their N-termini. ORB2A-ORB2B hetero-oligomers serve as translational activators whereas, in the absence of ORB2A, ORB2B does not oligomerize and has been shown to repress translation of reporter transcripts *in vitro* and in S2 tissue culture cells (MASTUSHITA-SAKAI *et al.* 2010; KHAN *et al.* 2015; STEPIEN *et al.* 2016; HERVAS *et al.* 2020).

All animal embryos undergo an MZT that encompasses two global processes: First, a subset of maternal gene products loaded into the egg by the female parent is cleared; second, the zygotic genome becomes transcriptionally active (TADROS AND LIPSHITZ 2009; VASTENHOUW *et al.* 2019). In Drosophila embryos, the MZT is very rapid, reaching completion about 3 hours after fertilization. Several RBPs have been shown to function during the first phase of the Drosophila MZT (TADROS *et al.* 2007; CHEN *et al.* 2014; LAVER *et al.* 2015; HAUGEN *et al.*

2024) together with their co-regulatory complexes involved in translational repression (NELSON

et al. 2007; CAO *et al.* 2020) and/or mRNA degradation (SEMOTOK *et al.* 2005).

Here we elucidate the role of ORB2 during the Drosophila MZT and show that it

translationally represses hundreds of bound mRNAs through its ZZ domain. The early

Drosophila embryo is unusual in that only one of the two protein isoforms, ORB2B, is expressed,

affording us the opportunity to assess ORB2B's mechanism of action and function *in vivo* in the

absence of possible confounding by the presence of ORB2A. To identify protein and RNA

components of ORB2B-containing ribonucleoprotein complexes (RNPs) during the MZT, we

used a synthetic antibody to immunoprecipitate (IP) endogenously expressed ORB2-containing

RNPs followed by either next-generation sequencing of co-immunoprecipitated transcripts (RIP-

seq) or mass spectrometry of associated proteins (IP-MS). We identified over 450 transcripts

bound by ORB2; these are enriched for mRNAs that are translationally repressed and degraded

during the MZT. We also identified over 200 protein binding partners of ORB2, including

components of the 43S translation pre-initiation complex (PIC), the Cup-ME31B-TRAL-BEL

repressive complex (hereafter referred to as the Cup repressive complex). We leveraged cell

culture assays to map ORB2's repressive function to the C-terminal ZZ domain; showed that the

repressive function is conserved in its subfamily ortholog, human CPEB2 (hCPEB2) but not in

members of the other subfamily member, ORB or hCPEB1; and mapped the repressive function

to regions within the ZZ domain. We show that CRISPR deletion of the ZZ domain of

endogenous ORB2 results in loss of interaction with the PIC and the Cup repressive complex,

and derepression of ORB2's targets in early embryos, thus uncovering ORB2's post-

transcriptional role during the MZT.

MATERIALS AND METHODS

Drosophila stocks

The following *Drosophila melanogaster* stocks were used: w^{1118} and $orb^{\Delta ZZ}$. The latter

stock was generated for use in this study (see below). Flies were cultivated at 25°C under

standard laboratory conditions unless otherwise indicated.

Generation of the CRISPR deletion of the ZZ domain at the ORB2 endogenous locus

Generation of this CRISPR line was performed by Well Genetics (13F.-12, No. 93, Sec. 1,

Xintai 5th Rd., Xizhi Dist., New Taipei City 221416, Taiwan (R.O.C.)). The scarless PBac-

dsRed method was used to produce the *orb2*$^{\Delta ZZ}$ line (GRATZ *et al.* 2013; GRATZ *et al.* 2014). The

ZZ domain, defined as from D638 to C704 (using the numbering from the ORB2-PB isoform)

was targeted for deletion such that the endogenous 3'UTR was retained in its entirety. Two guide

RNAs (gRNAs) were used to direct CRISPR to make double-stranded cuts in the DNA

surrounding the mutation of interest; in our case, one gRNA was directed to the genome to a cut

site just after the beginning of the ZZ domain (-179 nt from STOP Codon of *orb2-RA/B/C/D/H*),

and the second gRNA was directed to cut +3 nt from STOP Codon. Cas9 mediates breaks in the

double-stranded DNA, which triggers homologous repair between the genomic locus and a donor

plasmid, which was co-injected with the gRNA plasmid.

The donor plasmid which contained the mutation of interest (a deletion of the ZZ domain,

which removes the genomic portion of the coding sequence after the end of the second RRM,

advancing the STOP codon to this point, such that this mutant protein would stop at residue

L637 in the PB isoform) followed by a dsRed marker flanked by Piggy Bac 5′ terminal repeats

(Figure S7a). The Piggy Bac transposon terminal repeats enabled this marker to be excised after

it had been used to screen for successful CRISPR incorporation into the genome, leaving behind

no trace of the marker afterwards (i.e. it is "scarless" on the genome). Flanking this dsRed

marker is genomic sequence that is homologous to the base genome, 1kb upstream and

downstream for the cut site; this provided the homology needed to facilitate homology-

dependent repair (HDR) between the donor plasmid and the genomic site.

After injections of the host *w*^{1118} strain (LWG182), balanced fly lines were established

from the resulting progeny and screened for successful integration of the mutation-containing

region of the donor plasmid. Successful recombination resulted in an ORB2 gene locus harboring

a precise deletion of the ZZ domain at the endogenous locus, marked by the expression of dsRed,

which was later removed from the genome using the Piggy Bac transposon machinery to excise

the marker by crossing them to a Piggy Bac expressing fly (BDSC #8285). This was done by

using primers which flanked the upstream and downstream insertion sites, where a successful

PCR product would only be observed when incorporation of the donor plasmid occurred at the

intended location in the correct orientation (Figure S7b,c).

Synthetic antibody (Fab) expression and harvesting

Phage displayed synthetic antibodies against ORB2 protein have been described previously (LAVER et al. 2012; NA et al. 2016). To summarize, antigens containing amino acids 163-446 (numbered according to ORB2-PA isoform) were expressed and purified from *E. coli* as GST fusion proteins. Synthetic antibody library F (LAVER et al. 2012; NA et al. 2016) was used in five rounds of binding selection, which included negative selection to remove antibodies that bound to GST, to obtain synthetic antibodies that bound to the ORB2 antigen. The two synthetic antibodies used in this study were anti-ORB2 E8 and anti-hEGFR C1 (control).

Synthetic antibodies ('Fabs') were expressed and harvested from *E. coli* as previously described (LAVER et al. 2012). In summary, DH5α *E. coli* was transformed with plasmid containing the FLAG- and 6xHis-tagged Fab open reading frame under the control of the *lac* operator. Transformed cells were used to inoculate bacterial culture using 2xYT media; the inoculated culture was incubated at 37°C on rotation at 225rpm until OD595 reached 0.8. The culture was then induced with isopropyl β- d-1-thiogalactopyranoside (IPTG) to drive expression of the Fab and allowed to grow at 18°C overnight on rotation at 225rpm. Bacteria were pelleted by centrifugation at 4°C, 4000 rpm for 30 minutes, then bacterial pellet was resuspended with B-PER bacterial protein extraction reagent liquid solution (Thermo Scientific) supplemented with protease inhibitors (150mM NaCl, 1mM AEBSF, 2mM Benzamidine, 2μg/mL Leupeptin, 2μg/mL Pepstatin) and Benzonase nuclease (2500 units/mL). Cells were allowed to lyse for 30 minutes at 4°C, then the lysate was cleared by centrifugation at 4°C, 15 000 rpm for 30 minutes. Supernatant was collected and synthetic antibodies were purified by passing lysates through a column filled with Protein A affinity gel matrix beads; the captured Fab was eluted with an acid buffer containing 50mM NaH_2PO_4, 100mM H_3PO_4, and 140mM NaCl, pH to 2.8 in eight fractions of 200μL each. Fab-containing fractions were pooled and dialyzed with 1L PBS at 4°C for eight hours, twice. Dialyzed Fab was diluted with an equal volume of 95% glycerol 1xPBS and stored at -20°C.

Collection of embryo lysates

Lysate was collected from embryos laid and aged at 25°C. Extract was prepared from embryos collected 0 to 3 hours post egg-lay on apple juice agar plates, dechorionated with 100% bleach for 2 minutes, washed with 0.1% Triton X-100, and disrupted by crushing in minimal

volume of lysis buffer consisting of 150 mM KCl, 20 mM HEPES-KOH (pH 7.4), 1 mM MgCl$_2$,

1 mM DTT, and protease inhibitors (1 mM AEBSF, 2 mM benzamidine, 2 μg/mL pepstatin, 2

μg/mL leupeptin). The lysate was cleared by centrifugation at 4°C, 13 000 RPM for 15 minutes,

and the supernatant was collected and either used immediately or stored at -80°C.

RIP-seq

For each replicate of RIP-seq, 1mL of 0-3 hour embryo extract (prepared as described

above) was supplemented with Triton X-100 to 0.1% final concentration, then diluted in an equal

volume of lysis buffer with 0.1% Triton X-100 and cleared again by centrifugation at 4°C, 13

000 RPM for 15 minutes. The supernatant was incubated for 3 to 4 hours at 4°C with end-over-

end rotation with 100μL of anti-FLAG M2 affinity gel beads (Sigma) which had been pre-

incubated at 4°C with 50μg of purified anti-ORB2 E8 Fab blocked with 5μg/μL BSA. After

incubation, the lysates were decanted and beads were washed four times with lysis buffer

supplemented with 0.1% Triton X-100 and then three times with equilibration buffer: 100 mM

NaCl, 100 mM HEPES-NaOH (pH 7.4), 1 mM MgCl$_2$. Fabs and associated ribonuceoprotein

(RNPs) were eluted from the beads with 200 μL of 200μg/μL FLAG-peptide (Sigma) suspended

in equilibration buffer at 4°C for 20 minutes. RNA was isolated from the eluate using TRI

Reagent (Sigma) according to the manufacturer's instructions.

RNA was prepared for sequencing using Ribo-Zero Gold rRNA depletion kit and TruSeq

Stranded Total RNA Library Prep Kit (Illumina) according to manufacturer's instructions.

Libraries were sequenced using the HiSeq 2500 System, V4 Chemistry (Illumina), with paired-

end reads 125nt in length. The raw data re available on GEO (accession number GSE299661).

Computational analysis of RIP-seq data

Paired-end reads obtained from the sequencing were mapped to the Drosophila

melanogaster transcriptome r6.23, FB2018_04 obtained from flybase.org (OZTURK-COLAK et al.

2024) using the Salmon read mapping program (PATRO et al. 2017). The reference transcriptome

was supplied as a FASTA file. The resulting quantification was then analyzed for enrichment in

the ORB2 IP compared to control IP using the DESeq2 software package for R (LOVE et al.

2014). Prior to formal analysis, the dataset was filtered for transcripts having >1TPM in at least

three of the six samples, and rlog transformation was performed to stabilize the variance. The

similarity between samples was calculated by determining the Poisson Distance between pairs of
samples using the *PoiClaClu* package in R (see Figure S1) (WHITTEN 2011).The differential
expression analysis was performed with the *dds* function using alpha = 0.01 cut-off for the False
Discovery Rate. This returned a fold enrichment for each transcript, accompanied by an adjusted
P-value. We defined an ORB2-bound RNA target as satisfying all three of the following criteria:
(1) it was enriched in the ORB2 IP >1.5-fold relative to Input; (2) it had an ORB2 IP/Control IP
ratio >1.5; (3) it had an adjusted *P*-value < 0.05. The results are presented in Figure 1 and File
S1.
For comparisons between our dataset and previously published datasets, each pairwise
comparison was done independently. For comparisons that required a Wilcoxon Rank Sum test,
for each given pair of datasets, a common background list of genes was first compiled by taking
the intersection between the input/background of both datasets; genes which only appeared in
one dataset and not the other were discarded. Then, the measurements of the remaining identified
genes were placed into a dataframe in which each row corresponded to one unique gene, the first
column contained the measurements taken from one dataset, and the second column contained
measurements from the other dataset. This dataframe was then used to perform a Wilcoxon Rank
Sum test using the wilcox.test(x, y, exact = FALSE) function in R, where x is the data in the first
column, y is the data in the second column. The results are shown in Figure 2.
For comparisons that required a Fisher's Exact Test, for each given pair of datasets a
common background list of genes was first compiled by taking the intersection between the
input/background of both datasets; genes that only appeared in one dataset and not the other were
discarded. Then, the remaining identified genes of interest within in each dataset were compared
to look for common elements; genes that were identified in both lists of interest were annotated
as "Overlap", whereas genes that only appeared in one list were annotated as "X Only". A
Fisher's Exact Test was performed by using a 2x2 contingency table, where first cell (top left)
contained the number of genes that did not appear in either list (equal to the total number of
genes in the common background minus the number of genes described in the other three cells),
the second cell (top right) contained number of genes that appeared only on the ORB2 list
(ORB2 Only), the third cell (bottom left) contained the number of genes that appeared only on
the other list of interest (Other Only), and the last cell (bottom right) contained the number of

genes appear on both lists of interest (Overlap). This contingency table was constructed and

evaluated using the fisher.test() function in R. The results are presented in Table 1.

The Gene Ontology (GO) term enrichment analysis was performed using the Functional

Profiling function of the web version of gProfiler2 (accessed at https://biit.cs.ut.ee/gprofiler/gost)

(KOLBERG *et al.* 2023). The list of ORB2 bound genes was provided to the program in the Query

field, and the Run Query option was selected in order to begin analysis. The results are presented

in Tables 2-4.

Number of Accessible Target Sites (#ATS) *de novo* motif discovery

The ORB2 motifs were identified using the #ATS model, as described (LI *et al.* 2010).

Briefly, the #ATS program is a *de novo* motif discovery algorithm which takes as input nucleic

acid sequences belonging to mRNA transcripts in two lists: a positive (in this case, the 3′UTR of

ORB2-bound) and negative (in this case, the 3′UTR of co-expressed and unbound) list, and

outputs *k*-mers that best differentiate the positive transcripts from the negative transcripts. These

k-mers are assessed based on enrichment in the positive list compared to the negative list and for

accessibility as predicted by computational modeling of RNA secondary structures, and assigned

a *P*-value (see File S2). The top five seed *k*-mers from the #ATS were combined into a consensus

sequence by aligning the G in the motif and converting the resulting table into a position weight

matrix (PWM) shown in Figure 3.

Cell culture and transient transfection

The Drosophila S2 cell line was maintained at 25°C with ExpressFive SFM (Fisher

Scientific) medium supplemented with 16mM L-glutamine, 100units/mL penicillin, and

100µg/mL streptomycin. 400µL of S2 cells at a density of 1.75×10^6 were plated and transfected

with a mixture of 1.5ng Firefly luciferase-6x TAR plasmid, 1.5ng Renilla luciferase plasmid,

20ng Tat-FLAG-effector plasmid (containing the open reading frame for either ORB2 whole

protein, fragments of ORB2 protein, Protein A, or GFP), 4ng of FLAG-GST, and 178.5ng pSP72

using 0.4µL of TransIT-Insect transection reagent (Mirius Bio) according to the manufacturer's

instructions. Both luciferase reporters were under the control of the metal-inducible

metallothionein promotor; all FLAG-tagged constructs were derived from pAc5.1/V5-His

plasmid (Thermo Fisher Scientific), which carries the Actin5C promotor. Luciferase expression

was induced 24 hours post-transfection by the addition of copper sulphate to final concentration

of 0.5mM. 24 hours post-induction (48 hours post-transfection), cells were harvested for

downstream assays.

Luciferase Assay

Cell lysate was prepared from S2 cell transient transfections by dislodging cells from the

surfaces of the cell culture container and transferred into conical tubes. Cells were pelleted by

centrifugation at 13,000 rpm at 4°C for 1 minute; the supernatant was removed, and the cell

pellet was resuspended in by adding 20 µL of lysis buffer (containing 1xPassive Lysis Buffer

[Promega], 10 mg/mL BSA, and 1 mM AEBSF). Lysis was facilitated by pipetting up and down

and allowing the cells to lyse at 4°C for 10 minutes. Firefly and Renilla luciferase activities were

measured using the Dual-Luciferase Reporter Assay system (Promega) according to

manufacturer's instructions.

Reverse Transcription – Quantitative Real-time Polymerase Chain Reaction (RT-qPCR)

RNA isolated from ORB2 co-immunoprecipitations was reverse transcribed into single-

stranded cDNA with the Superscript IV reverse transcriptase kit (Invitrogen) according to

manufacturer's instructions. 500ng of total RNA per sample were used to synthesize cDNA,

which was primed using random hexamer primers. The resulting single-stranded cDNA was

diluted 1:25 using RNase-free water and used to perform quantitative real-time PCR (qPCR)

with primers specific to the various transcripts assayed. Primers were designed using NCBU

Primer-BLAST to cover all isoforms produced by a gene and were required to span an exon-

exon junction. qPCR was performed with SensiFAST SYBR PCR mix (Bioline) following the

manufacturer's protocol and using 5 µL of diluted cDNA per reaction. The CFX384 Real-Time

System (Bio-Rad) was used to carry out the PCR reaction.

The results of the qPCR were analyzed using CFX Manager software (Bio-Rad). Each

biological replicate represented a measurement obtained for a separate embryo collection; values

from three technical replicates were averaged and relative gene expression was normalized to the

ribosomal protein-coding transcript *RpL32*. Normalized expression was presented as the average

of three biological replicates with error bars representing standard deviation calculated by one-

way ANOVA.

Western Blots

The following primary antibodies were used: mouse anti-ORB2 4GS (1:50) obtained from the Developmental Studies Hybridoma Bank (https://dshb.biology.uiowa.edu/orb2-4G8) and described in (HAFER *et al.* 2011), mouse M2 anti-FLAG (Sigma, 1:5,000), and mouse anti-Actin (Sigma Cat#A4700, 1:10,000) which was used as a loading control.

Dechorionated embryos were counted, then lysed in Laemmli buffer at a concentration of 1 embryo/µL and boiled for 2 minutes. Proteins were resolved by SDS-PAGE and transferred to PVDF membrane in the absence of SDS. The blot was blocked at room temperature with 2% non-fat milk in PBST for 30 minutes and incubated with primary antibodies diluted in 2% non-fat milk in PBST at 4°C overnight. After this incubation, the blot was washed 3 x 10 minutes with PBST at room temperature and incubated with the appropriate HRP-conjugated secondary antibodies (1:5000, Jackson ImmunoResearch) in 2% non-fat milk in PBST at room temperature for 1 hour. The blot was washed 3 x 10 minutes with PBST and developed using Immobilon Luminata Crescendo Western HRP substrate (Millipore), imaged using ImageLab (BioRad) and band intensities were quantified using ImageLab.

Structure Prediction with AlphaFold

For structural predictions the full amino acid sequence of the ZZ domain constructs used in the tethering assays (see above) was predicted using AlphaFold version 3 (JUMPER *et al.* 2021; ABRAMSON *et al.* 2024). Predictions were visualized and truncated to show regions A-to-C in ChimeraX version 1.9 (MENG *et al.* 2023). Structures were color-coded to depict their predicted local distance difference test (pLDDT) score using the AlphaFold Error Plot tool. The results are presented in Figures S4 and 5.

Mass Spectrometry

Immunoprecipitation of ORB2 from embryo lysates with synthetic antibodies

For IP-MS, 500 µL of 0-3 hour embryo extract (prepared as described above) was supplemented with Triton X-100 to 0.1% final concentration, then diluted in an equal volume of lysis buffer with 0.1% Triton X-100 and cleared again by centrifugation at 4°C, 13 000 RPM for 15 minutes. The supernatant was incubated for 3 to 4 hours at 4°C with end-over-end rotation

with 100μL of anti-FLAG M2 affinity gel beads (Sigma) which had been pre-incubated at 4°C

with 50μg of purified anti-ORB2 E8 Fab blocked with 5μg/μL BSA. After incubation, the lysates

were decanted and beads were washed twice with lysis buffer supplemented with 0.1% Triton X-

100, then another twice with lysis buffer without Triton X-100; beads were transferred into a

new tube and washed another twice with lysis buffer without Triton X-100. Fabs and associated

proteins were eluted by tryptic digest: Beads were resuspended in 200 μL of 50mM ammonium

bicarbonate pH 8 supplemented with 2 μg of MS-grade trypsin and incubated at room

temperature overnight with end-over-end rotation. The digested peptides were collected and a

second elution with 200 μL of 50mM ammonium bicarbonate was performed at room

temperature for 30 minutes. The two fractions were pooled, flash frozen with dry ice, dehydrated

in a rotary evaporator for 8 hours, then stored at -20°C.

Liquid chromatography and mass spectrometry

For liquid chromatography (LC) the following solvents were used: Solvent A (0.1%

(vol/vol) formic acid) and Solvent B (0.1% (vol/vol) formic acid/90% (vol/vol) acetonitrile). The

LC parameters were set using an Easy-nLC 1200 instrument. Peptides were loaded and separated

on a nanoViper trap column (75μm x 2 cm), Acclaim PepMap 100 (C18, 3μm, 100 Å, Thermo

Scientific, cat. no. 164946), and EASY-Spray analytical column (75μm x 50cm) Acclaim

PepMap RSLC (C18, 2μm, 100Å, Thermo Scientific) using a flow rate of 225nL/minute.

Tandem MS was performed using the Q Exactive HF-Orbitrap mass spectrometer (Thermo

Scientific), as previously described (LIU *et al.* 2014; JIANG *et al.* 2015; CHIU *et al.* 2016). The

parameters for acquisition were 1 MS scan (50 ms; mass range, 390 to 1800) at a resolution of

60,000K, followed by up to 20 MS/MS scans (50 ms each) at a resolution of 15,000 and an AGC

target of 1x105. Candidate ions with charge states +2 to +7 were isolated using a window of 1.4

amu with a 5s dynamic exclusion window.

Data processing and analysis

Four biological replicates were performed, both in the presence or absence of RNase A.

The ProHits software package was used to perform Significance Analysis of INTeractome

(SAINT) in order to identify ORB2-interacting proteins (LIU *et al.* 2010; CHOI *et al.* 2012; LIU *et*

al. 2012). Proteins with associated peptide counts were filtered through the Trans Proteomic

Pipeline (TPP) for iProphet probability > 0.95 and number of unique peptides ≥2. SAINT

express analysis was performed on the ProHits interface with only Drosophila detected proteins

(selected options: number of compressed controls = 3, burn-in period, nburn = 2000; iterations,

niter = 5000, lowMode = 1; minFold = 1; normalize = 1; nCompressBaits = 3). This analysis

determined the probability of interaction for each identified prey protein. Proteins with AvgP

score ≥ 0.95 and BFDR ≤ 0.01 were defined as ORB2-ineracting proteins. Raw data are available

at MassIVE MSV000098275. The results are presented in File S3.

To perform network analysis on the list of identified protein interactors, we used the

GeneMANIA application (v3.5.2) (WARDE-FARLEY *et al.* 2010; FRANZ *et al.* 2018) with the

Cytoscape network visualization program (v3.9.1) (SHANNON *et al.* 2003; MONTOJO *et al.* 2010).

Data resulting from the IP-MS was imported to Cytoscape in table form. The interaction network

was produced by performing a Network Search on the list of proteins using the *Drosophila*

melanogaster database of GeneMANIA. The nodes of this network were then annotated

according to the number of spectral counts detected in our IP-MS and the edges were annotated

with known physical and genetic interactions reported in the GeneMANIA database. The results

are presented in Figure 5.

Embryo Sample Collection for *orb*$^{\Delta ZZ}$ Time-course and Polysome Gradient RNA-seq

Embryos were collected from *orb2*$^{\Delta ZZ}$ and *w*^{1118} (pre-injected control, LWG182) strains at

1-hour intervals post-egglay (0-1 hour, 1-2 hour, 2-3 hour), dechorionated with ice cold 100%

bleach for 2 minutes, washed with 0.1% Triton X-100, and disrupted by crushing using 0.5 g

embryos per mL of either lysis buffer (150 mM KCl, 20 mM HEPES-KOH (pH 7.4), 1 mM

MgCl2, 1 mM DTT, 1 mM AEBSF, 2 mM benzamidine, 2 μg/mL pepstatin, 2 μg/mL leupeptin)

or polysome lysis buffer (7.5 mM MgCl2, 500 mM NaCl, 25 mM Tris (pH 7.5), 1 mM DTT, 1

mM AEBSF, 2 μg/mL leupeptin, 2 mM benzamidine, 2 μg/mL pepstatin A, 2 mg/mL heparin,

0.5 mg/mL cycloheximide). The lysate with lysis buffer was cleared by centrifugation at 4°C, 13

000 RPM for 15 minutes, TRIzol (Invitrogen) was added to the supernatant which used

immediately for RNA extraction or stored at -80°C. The lysate with polysome lysis buffer was

either used immediately for polysome gradient processing or stored at -80°C.

Polysome Gradients

The polysome gradient protocol was adapted from our published method with
modifications (CHEN *et al.* 2014). Lysed samples were diluted 1 in 10 in polysome lysis buffer
and 20% Triton X-100 was added to a final concentration of 1%, centrifuged at 6,000xg for 10
minutes and the resulting supernatant was diluted in polysome lysis buffer supplemented with
1% Triton X-100 to an A_{260} of 12.5. A 5 ml 15% to 45% linear sucrose gradient in 7.5 mM
$MgCl_2$, 500 mM NaCl, 50 mM Tris pH 7.5 was produced using a BioComp Model 117 Gradient
Mate gradient maker (Biocomp, Fredericton, New Brunswick, Canada) using a rotation angle of
86 degrees and a rotation speed of 25 for 51 seconds. After chilling the polysome gradient at 4ºC
for 1 hour, 200 µL of diluted embryo extract was loaded onto the top of the gradient, which was
then spun at 36,000 rpm in a Beckman SW50.1 rotor for 2.5 hours. 400 µL fractions (hand
fractionated) were collected, and A_{260} absorbance was measured using a Nanodrop (Thermo
Fisher Scientific). We added 20% SDS, 0.5 M EDTA and 20 mg/ml proteinase K to final
concentrations of 0.8%, 0.01 M and 0.128 mg/ml, respectively, and then incubated the fractions
for 30 minutes at 37ºC. 2.5 µL GlycoBlue (15 mg/mL, ThermoFisher) following which 2.5x
volume ethanol was added to each fraction and precipitated overnight. The resulting pellet was
washed with 75% ethanol and resuspended in phenol-saturated water.

The fractions were then separated into three pools: Pool 1 contained the (top) fractions 1-3,
pool 2 contained fractions 4-6, pool 3 contained fractions 7-13. Following two phenol-
chloroform extractions, the aqueous phase of each pool was collected with Invitrogen
Phasemaker Tubes (Fisher A33248) following the manufacturer's protocol. The samples were
precipitated by the addition of 7.5 M LiCl to a final concentration of 3.75 M followed by an
overnight incubation at -20°C. The pellets were washed with 70% ethanol and resuspended in
water.

RNA purification was performed using 2.2× volume NEBNext RNA Sample Purification
Beads (NEBNext Ultra II DNA Library Prep with Sample Purification Beads, E7767L)
following the manufacturer's protocol. RNA concentration was assessed with the Qubit RNA HS
Assay Kit (Thermo Fisher) and Agilent bioanalyzer at The Centre for Applied Genomics
(TCAG), The Hospital for Sick Children, Toronto, ON, Canada.

**rRNA Depletion, Library Preparation and RNA-seq for Time-course and Polysome
Gradient RNA-seq**

For the RNA-seq time-course rRNA depletion was performed as previously described

(HAUGEN *et al.* 2024). For polysome gradient RNA-seq, rRNA depletion was performed using

the QIAseq FastSelect rRNA Fly Kit (QIAGEN, Cat# 33326). Library preparation for RNA-seq

was conducted using the NEBNext Ultra II Directional RNA Library Prep Kit for Illumina, and

sequencing was performed on an Illumina NovaSeq 6000 (PE150) at the Next Generation

Sequencing Facility, The Centre for Applied Genomics (TCAG), The Hospital for Sick Children,

Toronto, ON, Canada.

Computational Analysis of Time-course and Polysome Gradient RNA-seq Data

RNA-seq data processing pipeline:

Adapter trimming and quality control used Trim Galore (Version 0.6.5)

[https://github.com/FelixKrueger/TrimGalore]. rRNA and 'unwanted' RNA removal used

SortMeRNA (version 2.1) (KOPYLOVA *et al.* 2012), filtering against Drosophila melanogaster

7SLRNA, rRNA, pseudogene-rRNA, tRNA, mtRNA and other species' rRNA databases.

Transcriptome alignment was carried out with Salmon (Version 1.9.1) (PATRO *et al.* 2017) in

mapping-based mode using FlyBase release 6.23 transcriptome RNA (FB2018_04) (OZTURK-

COLAK *et al.* 2024). Full decoy indexing was applied following the salmon website instructions

[https://salmon.readthedocs.io/en/latest/salmon.html].

Differential gene expression analysis:

Differential expression analysis used the R package tximeta (version 1.20.3) (LOVE *et al.*

2020) and Deseq2 (version 1.42.1) (LOVE *et al.* 2014) in R (version 4.3.2). For the $orb2^{\Delta ZZ}$ and

wild type time-course total RNA-seq, reads from each time point were normalized using the

DESeq2 package. For the $orb2^{\Delta ZZ}$ and wild type polysome gradient RNA-seq time-courses, reads

of each time point of $orb2^{\Delta ZZ}$ and wild type pool 1 and pool 3 RNA were normalized using the

DESeq2 package. For each gene, the mean of normalized counts per pool was computed at each

time point and genotype. The 'Translation Index' of each mRNA was calculated as the ratio =

mean normalized counts of pool 3 /mean normalized counts of pool 1. Normalized counts were

used for statistical analysis by the Wilcoxon Rank Sum Test and data were visualized with

ggplot2 in R. Raw data are available at GEO for the RNA-seq timecourse and polysome gradient

timecourse with accession numbers GSE299140 and GSE299019, respectively. The results are

presented in Figure 6 and Files S5 and 6.

RESULTS

ORB2 binds to over 450 targets in the early Drosophila embryo

To identify RNA transcripts that are bound by ORB2B (hereafter referred to as ORB2), we

used our previously produced Fab against an antigen N-terminal to the RRMs (Figure 1a)

(LAVER *et al.* 2012; NA *et al.* 2016). We used this antibody to perform RNA-

coimmunoprecipitation (RIP) of endogenously expressed ORB2 from embryo extracts prepared

from 0-3 hour old embryos produced by w^{1118} females. We enriched for mRNA in the sample by

rRNA depletion, generated randomly primed cDNA libraries from the resulting samples, and we

identified co-purifying RNAs by next-generation RNA sequencing (see Materials and Methods).

As a negative control, we performed parallel RIPs using a control synthetic antibody, C1, as

described previously (LAVER *et al.* 2015; LAVER *et al.* 2020; HAUGEN *et al.* 2024). Input samples

from the same extracts were also sequenced. We performed three biological replicates for each

of these three conditions, for a total of nine cDNA libraries.

To identify ORB2-bound target genes, we mapped the sequence reads to the Drosophila

transcriptome using Salmon read alignment software, and analyzed the results with the DESeq2

software package as described in Materials and Methods (LOVE *et al.* 2014; PATRO *et al.* 2017).

We identified 462 genes whose transcripts were bound by ORB2 (Figure 1b and File S1). ORB2-

bound transcripts were significantly less abundant than co-expressed unbound transcripts (Figure

1b). RIP-RT-qPCR verified several of the bound and unbound targets (Figure 1c).

While ours is the first identification of ORB2 targets in embryos, there are previously

published target lists from other tissues or cells. For example, ORB2-bound transcripts were

identified in a CLIP experiment derived from ORB2 overexpression in S2 cells (STEPIEN *et al.*

2016). Of the 462 ORB2 targets on our list, 298 were identified in their CLIP-seq background set

of which they identified 113 as targets (Fisher's exact $P = 6.52 \times 10^{-7}$, Odds Ratio = 1.88). We

also examined the overlap of our targets with a list of ORB2-associated transcripts obtained by

performing RIP-seq on Drosophila head extracts (KOZLOV *et al.* 2023). 336 genes in our target

gene list overlapped with theirs (Fisher's exact $P < 10^{-15}$, Odds Ratio = 5.00).

The strong overlap between our ORB2 targets and previously reported targets from other

cell types/tissues verifies that our list is enriched for true ORB2 interactors.

ORB2 targets are translationally repressed and degraded during the MZT

To assess the post-transcriptional fate of ORB2-bound transcripts in the early Drosophila

embryo we utilized three previously published datasets that scored each transcript's translational

status, poly(A) tail length, and abundance during the MZT.

Two studies measured Translation Efficiency (TE), calculated from ribosome profiling,

during the MZT (DUNN *et al.* 2013; EICHHORN *et al.* 2016). At all three timepoints during the

MZT that were studied, ORB2-bound transcripts had a significantly lower TE compared to co-

expressed, unbound transcripts (Figure 2a-c). A third study measured Translation Index (TI),

calculated as the ratio of that transcript's abundance in polysome fractions of a sucrose gradient

compared to its abundance in non-polysomal fractions (CHEN *et al.* 2014). ORB2-bound

transcripts had a significantly lower TI than co-expressed unbound transcripts (Figure 2d).

We next assessed changes in poly(A) tail length and abundance for ORB2 targets between

the 0-1 hour timepoint and the 2-3 hour timepoint, which had been measured in parallel to TE

(EICHHORN *et al.* 2016). ORB2 targets underwent a significantly greater decrease in poly(A) tail

length and abundance than co-expressed, unbound transcripts (Figure 2e,f). We also examined

previously published lists of transcripts that are degraded during the MZT (Table 1). ORB2

targets overlapped significantly with transcripts identified to be unstable in activated eggs

(TADROS *et al.* 2007) (Fisher's Exact P = 2.2 x 10^{-16}, odds ratio 3.85). In addition, ORB2 targets

were significantly enriched for genes that are degraded during the MZT (THOMSEN *et al.* 2010):

(Class III and V; respectively, Fisher's Exact P = 0.02, odds ratio 1.54 and Fisher's Exact P =

2.2 x 10^{-16}, odds ratio 2.54) and significantly depleted for transcripts that are stable (Class I;

Fisher's Exact P = 2.2 x 10^{-16}, odds ratio 0.23).

We conclude that ORB2's target mRNAs are enriched for transcripts that are

translationally repressed, deadenylated and degraded during the MZT.

ORB2 targets are enriched for roles in gamete generation, gene regulation, and

development

Having assessed the post-transcriptional fate of ORB2's targets, we next asked whether

these targets were enriched for particular molecular, cellular or biological functions by

performing Gene Ontology (GO) term analysis using the gProfiler2 tool (GENE ONTOLOGY *et al.*

2023; KOLBERG *et al.* 2023) (Tables 2-4).

The top GO terms under the Molecular Function category (Table 2) were dominated by

molecular binding functions; 'mRNA 3′UTR binding', and 'mRNA regulatory element binding'

were among the top ten results. Genes belonging to these terms included *orb2* itself as well as

patronin, piwi, and *bru1*. Other top terms included 'translation regulator activity' and 'translation

repressor activity'; genes belonging to these terms included *orb2* itself, *orb, pum, fmr1, sxl, heph,*

and *eIF4E-HP*. In the Biological Process category (Table 3), ORB2 targets were enriched for

terms related to regulation and development, such as 'anatomical structure development', 'cell

differentiation', and 'regulation of metabolic process'. ORB2 targets were also enriched for

terms related to gametogenesis, including 'gamete generation', 'oogenesis', and 'germ cell

development'. Genes related to these terms included *pum, rbfox1, piwi, heph, bru1, sxl,* and *mei-*

P26. Enrichment for GO terms related to gametogenesis in ORB2 targets is consistent with

previously reported functions for ORB2 in, for example, regulation of spermatogenesis (XU *et al.*

2012; GILMUTDINOV *et al.* 2021). The Cellular Component category (Table 4) was dominated by

terms that included 'ribonucleoprotein granule', 'P granule', 'pole plasm', and 'germ plasm'.

Genes related to these terms included *larp, tud, osk, tapas, ago1,* and *dcr-2*.

Many of the genes listed under enriched GO terms are previously identified ORB2 targets

(e.g., *orb2* itself, *ago1*, *bru1*, *dcr-2*, *eIF4E-HP*, *fmr1*, *heph*, *orb*, *pum*, *rbfox1*, *unr*)

(MASTUSHITA-SAKAI *et al.* 2010; XU *et al.* 2012; STEPIEN *et al.* 2016; KOZLOV *et al.* 2021) while

others are previously unidentified players in the same enriched function or process (e.g., *imp and*

piwi). Enrichment for mRNAs encoding known post-transcriptional regulators suggests that

ORB2 may play a key role in the cross-regulatory network that controls the MZT, a function that

is supported by our subsequent analyses reported below. The fact that ORB2 binds to and

potentially regulates mRNAs that encode other post-transcriptional regulators is also consistent

with a previous study in S2 cells showing that RNP complexes tend to be enriched for mRNAs

encoding post-transcriptional regulators (STOIBER *et al.* 2015).

ORB2-bound transcripts are enriched for a U-rich motif

CPEB-family proteins are known to bind to U-rich elements, typically located in the

3′UTR of their mRNA targets (HAKE AND RICHTER 1994; DE MOOR AND RICHTER 1999; STEPIEN

et al. 2016). To predict *cis*-elements recognized by ORB2 from our *in vivo* RIP-seq results, we

used our algorithm, Number of Accessible Target Sites (#ATS), which performs *de novo* motif

discovery given a target set and a co-expressed non-target set of transcripts (LI *et al.* 2010).

Using our input RNA-seq data, we identified the most abundant transcript isoform from each

expressed gene in the early embryo and then compared the 3′UTR sequences of ORB2 targets to

those of non-targets. The #ATS program identified a list of 6-mers enriched in ORB2 target

sequences relative to non-targets; these 6-mers were then combined into a consensus motif

(Figure 3a, File S2), which will henceforth be referred to as an ORB2-recognition element

(ORE). The ORE, NUUGUWU (where W = A or U) has similarities to published binding motifs:

KUUUKKK (where K = G or U) identified by an *in vitro* binding assay (RAY *et al.* 2013);

UUUU(A)$_{1-3}$U and UUUUGU identified in S2 cells (STEPIEN *et al.* 2016); UUUURU (where R =

A or G) in brain extracts (MASTUSHITA-SAKAI *et al.* 2010).

The *de novo* discovery of a U-rich ORE enriched in ORB2-bound RNAs in the early

embryo and its similarity to ORB2 motifs previously identified using a variety of methods

supports the conclusion that our list of targets is enriched for transcripts directly bound by

ORB2.

The ORE directs ORB2 binding to and repression of a luciferase reporter mRNA in S2

cells

To validate the biological activity of the ORE identified by #ATS, we used a luciferase

reporter assay in Drosophila S2 cells to measure ORB2 binding and ORB2-mediated regulation.

In the assay, firefly (FLuc) reporters harboured six copies in the 3′UTR of either a representative

ORE (UUUUGUU) or a mutated ORE (UUAUCUC) (the reporter is schematized in Figure 3b).

Each of these reporters was individually expressed in S2 cells together with either a control

protein (GFP), full-length ORB2 protein, or an ORB2 protein lacking its RBD (ORB2ΔRRM).

An unregulated Renilla luciferase (Rluc) reporter was co-transfected with all samples, allowing

us to normalize Fluc mRNA and protein levels.

ORB2 RIP followed by RT-qPCR confirmed that the reporters carrying the wild-type

OREs, but not the mutant OREs, were enriched in ORB2 IPs, compared to IPs of the

ORB2ΔRRM protein (Figure 3c). Furthermore, significant ORB2-mediated repression of the

reporter was seen when ORB2 protein was expressed together with a reporter construct carrying

wild-type OREs (Figure 3d). When either the OREs were mutated or the RRM was deleted, no

repression was observed (Figure 3d) despite the fact that ORB2ΔRRM protein was expressed at

higher levels than full-length ORB2 (Figure S2).

We conclude that ORB2 binds to OREs and that this binding confers repression on a

reporter RNA. The extent of repression we observed is similar to that previously reported for

ORB2B in similar reporter assays in S2 cells (STEPIEN *et al.* 2016).

The C-terminal Zinc-binding ('ZZ') domain of ORB2/hCPEB2 but not ORB/hCPEB1

represses a luciferase reporter mRNA in S2 cells

To identify domains of ORB2 necessary and sufficient for repression we used a luciferase

reporter tethering assay in S2 cells. We transfected a plasmid construct that would express a

protein containing the BIV-Tat RNA-binding domain fused to all or part of the ORB2 protein.

This fusion protein is recruited to six tandem repeats of the trans-activating response (TAR)

element (to which BIV-Tat binds) located in the 3′UTR of a co-transfected firefly luciferase

reporter (schematised in Figure 4a). Tethering of full-length ORB2B protein produced significant

repression relative to a tethered GFP control (Figure 4b); the extent of repression was consistent

with previously published data for ORB2 in S2 cells (STEPIEN *et al.* 2016). Tethering of ORB2

protein did not affect steady-state reporter RNA levels (Figure 4c).

To map regions of ORB2 that mediate repression, we fused the BIV-tat domain to

fragments of the ORB2 protein and assessed whether any individual region was capable of

repressing luciferase expression. We divided the ORB2 protein into five fragments: IDR1, the

polyQ region, IDR2, the RBD, and the ZZ domain (Figures 1a and 4b,c). There was no

significant effect on the expression of luciferase after tethering of IDR1, polyQ, or IDR2 (Figure

4b and see Figure S3A for western blot of expression of each fragment). The RBD and the ZZ

domain conferred repression (Figure 4b). We note that the RBD was over-expressed relative to

the other fragments; however, the ZZ domain produced very strong (~three-fold) repression

despite being expressed at a comparatively low level (Figure S3a). To determine whether the ZZ

domain was also necessary for repression, we tethered an ORB2 protein construct lacking the ZZ

domain (ORB2ΔZZ) to the luciferase reporter and found that it was unable to repress luciferase

expression, despite ORB2ΔZZ retaining the RBD and being expressed at higher levels than full-

length ORB2 (Figure 4b, Figure S3b). Because luciferase mRNA levels were not significantly

different from controls in either the ZZ sufficiency or the ZZ necessity experiments (Figure 4c,

ZZ and ORB2ΔZZ), ZZ domain mediated repression can be attributed largely to effects on

translation rather than RNA stability. We conclude that the ZZ domain of ORB2 is both

necessary and sufficient to confer translational repression when tethered to a reporter mRNA.

Finally, to assess possible conservation of function, we also tethered the ZZ domains of

ORB, hCPEB1 and hCPEB2. The hCPEB2 ZZ domain conferred significant repression, whereas

the ORB ZZ and hCPEB1 ZZ domain were unable to repress (Figure 4e; western blot is shown

in Figure S3c). We conclude that the repressive function of the ZZ domain is conserved between

ORB2 and its human ortholog, hCPEB2, while it is not conserved in ORB or hCPEB1. We note

that the ZZ domains of hCPEB3 and hCPEB4 are $\geq$ 90% identical to the hCPEB2 ZZ domain.

This suggests that, although not tested here, the hCPEB3 and hCPEB4 ZZ domains likely also

function in translational repression.

Regions within the ORB2 ZZ domain are sufficient to confer repression

To map regions of the ORB2 ZZ domain sufficient for repression, we first compared the

sequences of the ZZ domains of ORB, ORB2, hCPEB1 and hCPEB2. There is high level of

sequence identity between the ORB2 ZZ domain and that of hCPEB2 (81.3% amino acid

identity, Figure 4d). In contrast, the ZZ domain of ORB2 and ORB, exhibit only 35.6% sequence

identity and hCPEB1 has only 28.5% amino acid identity to ORB2 (Figure 4d). We then

considered the predicted structures of these domains (Figure S4). An NMR structure of the

hCPEB1 ZZ domain showed it consists of a rubredoxin (Rd) turn, followed by a beta hairpin,

and a short alpha helix (MERKEL *et al.* 2013). Predictions of the structures of the ORB, ORB2

and hCPEB2 ZZ domains using Alphafold 3 (ABRAMSON *et al.* 2024) showed a similar

arrangement. Indeed, Alphafold3 prediction of the ORB ZZ structure was virtually identical to

that of the NMR structure of the hCPEB1 ZZ domain (Figure S4a). Similarly, the predictions of

the ORB2 and hCPEB2 ZZ structures were very similar to one another (Figure S4a). Finally, an

overlap of the ORB and ORB2 ZZ domain structures showed that the only difference was the

length of the Rd turn (Figure S4b).

To map regions within the ORB2 ZZ domain sufficient for repression we focused on the 46

amino acids beginning two amino acids before the first zinc-coordinating residue (at Q640) and

ending at the C-terminal zinc-coordinating residue (H685). We asked whether we could

transform the non-repressive ORB ZZ domain into a repressive one by substituting the Rd turn

('A', Q640 to P656), the beta hairpin ('B', F657-Y668), and/or the alpha helix ('C', C669 to

H685) regions of the ORB2 ZZ domain into the corresponding portion of the ORB ZZ domain

and assessing their function using the tethering assay. Importantly, AlphaFold3 did not predict

any disruptions in the folding of the chimeric proteins when the ORB ZZ domain carried one of

these ORB2 regions (Figure S5).

When the entire 46 amino acid region was substituted from the ORB2 ZZ domain into the

ORB ZZ domain (ORB ZZ-$A^2B^2C^2$, where the superscript '2' refers to ORB2 and superscript '1'

refers to ORB), the chimeric construct produced significant repression relative to the GFP

control (Figure 4f). Region A (ORB ZZ-$A^2B^1C^1$) and region C (ORB ZZ-$A^1B^1C^2$) produced

significant repression whereas region B (ORB ZZ-$A^1B^2C^1$) did not (Figure 4f; western blots are

shown in Figure S6).

We conclude that regions A and C of the ORB2 ZZ domain are each sufficient to convert

the ORB ZZ domain into a repressor.

ORB2 interacts with translational repressors and the 43S translation preinitiation complex

in early embryos

We next turned to proteomic strategies to identify protein partners of ORB2 in early

embryos. We immunoprecipitated endogenous ORB2 from extracts prepared from embryos

produced by w^{1118} females, both in the presence and absence of RNase A, in biological

quadruplicates, using the same synthetic antibody used above to identify ORB2 target RNAs.

Co-purifying proteins were identified by mass spectrometry followed by Significance Analysis

of INTeractome (SAINT), which assigns confidence scores to protein-protein interactions based

on proteomic data (CHOI *et al.* 2012). We defined a protein interactor of ORB2 as any protein

that was significantly enriched in ORB2 immunoprecipitates compared to parallel purifications

using our negative control Fab, C1, with a SAINT (AvgP) score ≥ 0.95 and a BFDR < 0.01 (File

S3).

We identified 212 interactors of ORB2 in the absence of RNase A and 106 interactors in

the presence of RNase A, the latter representing RNA-independent interactions; 85 were in

common (File S3). The interactors largely fell into two distinct groups: The first consisted of

RBPs known to negatively regulate target stability and/or translation, including AGO1, AGO2,

PUM, and RBFOX1, as well as the Cup repressive complex (Cup, ME31B, TRAL, BEL)

(GOTZE *et al.* 2017) (Figure 5a, File S3). All components of the Cup repressive complex

copurified in the presence of RNase, consistent with an RNA-independent interaction with

ORB2. The second group consisted of translation-initiation factors, notably the components of

the 43S preinitiation complex (PIC): all eIF3 and eIF2 subunits, eIF1 and -1A, and eIF5A; most

(13/20) were RNA-independent interactors, including 11/14 detected eIF3 subunits (Figure 5b

and File S3).

We note that a subset of ORB2's protein interactors were also identified as mRNA targets

in our RIP-seq experiments (File S4). This result is consistent with an earlier study of RBP

mRNA targets and protein interactors carried out in S2 cells, which showed that RNP complexes

are enriched for both mRNAs and their encoded proteins involved in RNA binding and post-

transcriptional regulation (STOIBER *et al.* 2015).

Co-purification of ORB2 with negative regulators of mRNA stability and translation is

consistent with our analyses of the post-transcriptional fate of ORB2 targets during the MZT

(repressed and unstable; Figure 2) and our reporter experiments in S2 cells (repressed, Figures 3

and 4). In the Discussion, we consider possible reasons for the lack of an effect on stability in S2

cells as well as possible implications of the PIC interaction.

The ORB2 ZZ domain is required for interaction with the 43S PIC and the Cup repressive

complex

The experiments described above showed that the ZZ domain of ORB2 is necessary and

sufficient for translational repression in our S2-cell tethering assay; therefore, we assessed

whether this domain is important for association with ORB2's identified binding partners in

early embryos. To generate an endogenous deletion of the ZZ domain, we used a scarless dsRed

CRISPR method to remove all amino acids from ORB2 D638 to the C-terminus at C704,

inclusive, leaving the endogenous 3′UTR intact (see Figure 4d, Materials and Methods, Figure

S7) (GRATZ *et al.* 2013; GRATZ *et al.* 2014). The mutant fly lines resulting from the excision will

henceforth be referred to as *orb2*$^{\Delta ZZ}$. Western blots on whole-fly extract showed that ORB2ΔZZ

was expressed at higher levels than full-length ORB2 in the pre-CRISPRed w^{1118} wild-type

control flies (Figure S7d) and a similar result was obtained for 0-3 hour embryos (Figure S8a).

We confirmed by RIP-RT-qPCR that deletion of the ZZ domain did not significantly affect

ORB2ΔZZ's ability to bind to several of its RNA targets in embryos (Figure S8b).

Immunopurification of ORB2ΔZZ protein from 0-3 hour embryos produced by *orb2*$^{\Delta ZZ}$

homozygous females followed by mass spectrometry identified 91 high-confidence protein

interactors in the absence and 48 in the presence of RNase A, of which 43 overlapped (File S3).

When the union of these interactors (96 proteins) was compared to the union of full-length

ORB2 IP-MS interactors (233 proteins), we found that 65 overlapped, including several

translational repressors such as PUM and RBFOX1 (File S3). Strikingly, interactions with all

components of the PIC and the Cup repressive complex were lost upon deletion of the ZZ

domain.

We conclude that the ORB2 ZZ domain is required for interaction between ORB2 and the

43S PIC as well as the Cup repressive complex, but that additional repressors retain interaction

with ORB2 in the absence of its ZZ domain.

The ZZ domain is necessary for translational repression of ORB2 targets in the early

embryo

Since our experiments in S2 cells showed that the ZZ domain is necessary for ORB2-

mediated translational repression, and our IP-MS results indicated that ORB2 loses interactions

with the Cup repressive complex when the ZZ domain is deleted, we next asked whether deletion

of the ZZ domain had consequences for post-transcriptional regulation of ORB2's target mRNAs

during the MZT.

First, we assessed whether ORB2's targets, which are enriched for maternal transcripts that

are deadenylated and degraded during the MZT (see Figure 2e,f), were stabilized in ORB2ΔZZ

mutant embryos. We collected embryo extracts from embryos laid by control or *orb2*$^{\Delta ZZ}$ females

at 0-1, 1-2 and 2-3 hours after egg lay and carried out RNA-seq followed by analysis using

DEseq2. We compared the expression level of targets and co-expressed non-targets across this

time course and found that, at all three stages, there was no significant difference in expression

level between control versus ORB2ΔZZ embryos for either target or non-target transcripts

(Figure 6a and File S5). These results suggest that, while ORB2 binds to unstable mRNAs, it

does not induce their degradation or, alternatively, that ORB2 may direct destabilization of its

targets during the MZT in a ZZ domain-independent mechanism.

Second, we assessed whether ORB2's targets, which are enriched for transcripts with

significantly lower TE and TI than co-expressed non-targets (see Figure 2a-d), underwent an

increase in polysome association in ORB2ΔZZ embryos. We collected embryo extracts from

embryos laid by control or *orb2^{ΔZZ}* females at 0-1, 1-2 and 2-3 hours after egg lay, then ran

polysome gradients followed by RNA-seq of pooled polysome fractions and pooled 'free' (non-

polysome-associated) fractions. For each gene, we calculated the ratio of RNA in the polysome

fractions to that in the free fractions (see Materials and Methods); if translational repression were

relieved upon deletion of the ZZ domain then this ratio would increase (i.e., transcripts would

move onto polysomes).We found that, in ORB2ΔZZ embryos, ORB2 targets shifted onto

polysomes at the first timepoint (0-1 hour) whereas non-targets did not (Figure 6b and File S6).

We conclude that the ZZ domain of ORB2 mediates translational repression of ORB2

targets at the beginning of the MZT. At subsequent timepoints (1-2 and 2-3 hours), there was no

significant difference in polysome association of ORB2 targets in ORB2ΔZZ embryos versus

controls; possible reasons will be considered in the Discussion.

DISCUSSION

In this study, we identified hundreds of RNA targets and protein partners of the Drosophila

ORB2 protein in the early embryo. ORB2's targets are enriched for translationally repressed

mRNAs that are degraded during the MZT. These targets are enriched for a U-rich motif that

directs binding and repression of a reporter in S2 cells. Repression by ORB2 is mediated by the

ZZ domain, a function that is conserved in hCPEB2. Deletion of the ZZ domain from the

endogenous ORB2 gene results in loss of physical interaction with the Cup repressive complex

and derepression of ORB2's mRNA targets in the early embryo.

A conserved role for the ORB2/hCPEB2 ZZ domain in mediating translational repression

Structural analysis of the ZZ domain of hCPEB1 has implicated it in protein-protein

interaction (MERKEL *et al.* 2013; STEPIEN *et al.* 2016), consistent with earlier studies on, for

example, the ZZ domain of CREB-binding protein (LEGGE *et al.* 2004). Our identification of

multiple proteins that depend *in vivo* on the ZZ domain for interaction with ORB2 are consistent

with a major role in recruiting co-regulators to target transcripts. Our data showing that deletion

of the ZZ domain does not affect RNA binding are consistent with a previous study (STEPIEN *et*

al. 2016). The fact that, in our S2 cell tethering assay, the ZZ domain is necessary and sufficient

for repression permitted us to assess evolutionary conservation of this role as well as to map

regions of the ZZ domain that implement this function.

Interestingly, we found that the repressive function of the entire ZZ domain is conserved

between ORB2 and hCPEB2 but not with ORB and hCPEB1. Swapping regions from ORB2 into

ORB identified which of these are sufficient for repression. Notably, the Rd turn at the N-

terminus ('A', Q640 to P656) and the α-helix at the C-terminus ('C', C669 to H685) were

sufficient to transform the ORB ZZ domain into a repressor. The ZZ domains of hCPEB3 and 4

show ≥ 90% amino acid identity to hCPEB2, with 16 of the 17 residues of region A and 12 of

the 13 residues of region C identical between hCPEB2-4. As such, it is quite likely that the ZZ

domains of hCPEB3 and 4 are also capable of repressing target mRNA translation via regions A

and C.

That fact that region A or C of ORB2's ZZ domain are each sufficient to convert the ORB

ZZ domain into a translational repressor indicates that they can function independent of one

another, perhaps by binding to different co-repressors. Future work aimed at identifying the

ORB2 binding partners that interact with regions A and/or C should provide insights into the

molecular mechanisms that underlie ZZ domain function.

Regulation of post-transcriptional processes during the MZT

Our analyses of the post-transcriptional fate of ORB2's targets during the MZT indicate

that they are enriched for transcripts that are translationally repressed and degraded. Consistent

with this outcome, multiple negative regulators of mRNA translation and stability copurify with

ORB2 in early embryos; most of these copurify in the presence of RNase, consistent with an

RNA-independent interaction with ORB2. Several of the regulators that interact in an RNA-

independent manner require the ZZ domain for interaction (notably, the Cup complex

components – Cup, TRAL, ME31B and BEL) while others interact both in the presence and

absence of the ZZ domain (e.g., PUM, FMR1, RBFOX1).

The S2 cell reporter assay showed that, while full-length ORB2 and its isolated ZZ domain

are both able to repress translation, they did not affect steady-state RNA levels. The simplest

explanation for this difference between S2 cells and embryos is that key factors that direct

degradation in embryos may not be present or may be expressed at low levels in S2 cells. For

example, RBFOX1, which we identified as an ORB2-interactor by IP-MS in early embryos, is

expressed at low levels in S2 cells according to RNA-seq data (CHERBAS *et al.* 2011). At present

our results indicating a role for ORB2 in degrading mRNAs in the embryo are correlative;

experiments to assess whether ORB2 directly triggers degradation of target mRNAs during the

MZT and the role of ORB2-interactors in this regulation awaits further investigation. What is

clear is that if ORB2 does play such a role, it does not require the ZZ domain.

We have shown that the ZZ domain is necessary and sufficient for translational repression

of reporters in S2 cells and is required to repress ORB2's *in vivo* targets in 0-1 hour embryos.

That the Cup complex is expressed in both S2 cells and embryos, that ZZ domain deletion results

in loss of Cup complex interaction with ORB2 in embryos, and that ORB2's targets in embryos

lose repression when the ZZ domain is deleted from endogenous ORB2, are all consistent with

the hypothesis that ORB2-dependent repression is mediated by the Cup complex. The Cup

complex prevents interaction of eIF4G with eIF4E, thus abrogating translation initiation

(WILHELM *et al.* 2003; NAKAMURA *et al.* 2004; NELSON *et al.* 2004); consistent with this role, we

note that, while eIF4E was present in our ORB2 IP-MS data, eIF4G was absent. Previous

studies have shown that the Cup complex is brought to mRNAs by BRU1 in oocytes and by

SMG in embryos (WILHELM *et al.* 2003; NAKAMURA *et al.* 2004; NELSON *et al.* 2004). ORB2

would represent a third RBP that recruits this complex for repression, in this case early in the

MZT.

The Cup complex has been shown to be cleared by the CTLH E3 ligase part-way through

the MZT (CAO *et al.* 2020; ZAVORTINK *et al.* 2020), consistent with the ZZ domain's repressive

role in the early but not late MZT. However, ORB2's targets are repressed at both 0-1 hour

(when the Cup complex is present) and at 2-3 hours (after the Cup complex has been cleared);

furthermore, ORB2's targets are deadenylated and degraded between 0-1 and 2-3 hours, but

deletion of the ZZ domain has no effect on their clearance. These data suggest that additional

regulators may function either independent of or together with ORB2 to direct these processes; if

the latter, they would interact with domains of ORB2 distinct from the ZZ domain. PUM, FMR1

and RBFOX1 are present throughout the MZT (CASAS-VILA *et al.* 2017; CAO *et al.* 2020) and

we have shown here that their interaction with ORB2 is both RNA-independent and ZZ-domain

independent, making them candidates for implementing these roles.

The mystery of PIC interaction

Unexpectedly, we identified the 43S PIC as a major interactor of ORB2 via its ZZ domain.

An essential component of the PIC is the eIF3 complex (its subunits are encoded by 14 genes in

Drosophila), which is a highly regulated component of translation initiation (HARRIS *et al.* 2006;

CHIU *et al.* 2010; CATE 2017). Several of the mammalian eIF3 subunits have been shown to have

RNA-binding ability (LEE *et al.* 2015; LEE *et al.* 2016) and to mediate distinct regulatory

outcomes (PESTOVA *et al.* 1998; PESTOVA AND HELLEN 1999; HUI *et al.* 2003; HUI *et al.* 2005;

MARTINEAU *et al.* 2008; LEE *et al.* 2010; MARTINEAU *et al.* 2014).

At this time we do not have data that provide insights into the functional consequences of

ORB2's interaction with the PIC. ORB2's interaction with the PIC's eIF3 subunits is RNA-

independent, consistent with direct interaction with one or more of these subunits. One

possibility that we considered is that ORB2 may sequester the PIC, thus reducing translation

overall. However, in such a model, one would expect a global increase in translation upon loss of

PIC interaction rather than what we observed in ORB2ΔZZ embryos, which is an increase in

polysome association of ORB2's targets but no change in non-targets. An alternative hypothesis

is that ORB2-dependent binding to the PIC on target transcripts has a negative effect on

translation either in addition to or in concert with the Cup repressive complex. Indeed, a

speculative model might be that ORB2 binding to the 3′UTR and interaction with PIC at the

5′UTR might repress translation initiation by interfering with PIC function. The mechanistic and

functional consequences of ORB2-PIC interaction will be a fruitful area for future study.

Data Availability

Drosophila strains are available upon request. Raw data for RIP-seq, RNA-seq and IP-MS

are available at GEO (GSE299661 for RIP-seq, GSE299140 for RNA-seq time-course,

GSE299019 for polysome gradient seq) and at MassIVE (MSV000098275 for IP-MS). The

authors affirm that all of the other data necessary for confirming the conclusions of the article are

present within the article, figures, tables and supplementary files.

Acknowledgements

858
859   We thank Kun Nie for assisting with the *de novo* motif discovery using #ATS and Dr.

860   Dorothy Lerit for feedback on a draft of this manuscript.

Funding

This research was supported by grants from the Canadian Institutes of Health Research

PJT-159702 (HDL), PJT-190124 (HDL) and PJT-159494 (CAS). TCHL was supported in part

by a Canada Graduate Scholarship (CGS-M) and a University of Toronto Open Fellowship; CW

was supported in part by a University of Toronto Open Fellowship. Molecular graphics and

analyses were performed with UCSF ChimeraX, developed by the Resource for Biocomputing,

Visualization, and Informatics at the University of California, San Francisco, with support from

National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and

Computational Biology, National Institute of Allergy and Infectious Diseases.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

LITERATURE CITED

Abramson, J., J. Adler, J. Dunger, R. Evans, T. Green *et al.*, 2024 Accurate structure prediction
of biomolecular interactions with AlphaFold 3. Nature 630**:** 493-500.
Cao, W. X., S. Kabelitz, M. Gupta, E. Yeung, S. Lin *et al.*, 2020 Precise Temporal Regulation of
Post-transcriptional Repressors Is Required for an Orderly Drosophila Maternal-to-
Zygotic Transition. Cell Rep 31**:** 107783.
Casas-Vila, N., A. Bluhm, S. Sayols, N. Dinges, M. Dejung *et al.*, 2017 The developmental
proteome of Drosophila melanogaster. Genome Res 27**:** 1273-1285.
Cate, J. H., 2017 Human eIF3: from 'blobology' to biological insight. Philos Trans R Soc Lond B
Biol Sci 372.
Chen, L., J. G. Dumelie, X. Li, M. H. Cheng, Z. Yang *et al.*, 2014 Global regulation of mRNA
translation and stability in the early Drosophila embryo by the Smaug RNA-binding
protein. Genome Biol 15**:** R4.
Cherbas, L., A. Willingham, D. Zhang, L. Yang, Y. Zou *et al.*, 2011 The transcriptional diversity
of 25 Drosophila cell lines. Genome Res 21**:** 301-314.
Chiu, C. W. N., C. Monat, M. Robitaille, M. Lacomme, A. M. Daulat *et al.*, 2016 SAPCD2
Controls Spindle Orientation and Asymmetric Divisions by Negatively Regulating the
Galphai-LGN-NuMA Ternary Complex. Dev Cell 36**:** 50-62.
Chiu, W. L., S. Wagner, A. Herrmannová, L. Burela, F. Zhang *et al.*, 2010 The C-terminal
region of eukaryotic translation initiation factor 3a (eIF3a) promotes mRNA recruitment,
scanning, and, together with eIF3j and the eIF3b RNA recognition motif, selection of
AUG start codons. Mol Cell Biol 30**:** 4415-4434.
Choi, H., G. Liu, D. Mellacheruvu, M. Tyers, A. C. Gingras *et al.*, 2012 Analyzing protein-
protein interactions from affinity purification-mass spectrometry data with SAINT. Curr
Protoc Bioinformatics Chapter 8**:** Unit8.15.
de Moor, C. H., and J. D. Richter, 1999 Cytoplasmic polyadenylation elements mediate masking
and unmasking of cyclin B1 mRNA. EMBO J 18**:** 2294-2303.
Dunn, J. G., C. K. Foo, N. G. Belletier, E. R. Gavis and J. S. Weissman, 2013 Ribosome
profiling reveals pervasive and regulated stop codon readthrough in Drosophila
melanogaster. Elife 2**:** e01179.
Duran-Arqué, B., M. Cañete, C. L. Castellazzi, A. Bartomeu, A. Ferrer-Caelles *et al.*, 2022
Comparative analyses of vertebrate CPEB proteins define two subfamilies with
coordinated yet distinct functions in post-transcriptional gene regulation. Genome Biol
23**:** 192.
Eichhorn, S. W., A. O. Subtelny, I. Kronja, J. C. Kwasnieski, T. L. Orr-Weaver *et al.*, 2016
mRNA poly(A)-tail changes specified by deadenylation broadly reshape translation in
Drosophila oocytes and early embryos. Elife 5.
Fernandez-Miranda, G., and R. Mendez, 2012 The CPEB-family of proteins, translational
control in senescence and cancer. Ageing Res Rev 11**:** 460-472.
Franz, M., H. Rodriguez, C. Lopes, K. Zuberi, J. Montojo *et al.*, 2018 GeneMANIA update
2018. Nucleic Acids Res 46**:** W60-W64.
Gene Ontology, C., S. A. Aleksander, J. Balhoff, S. Carbon, J. M. Cherry *et al.*, 2023 The Gene
Ontology knowledgebase in 2023. Genetics 224.

Gilmutdinov, R., E. N. Kozlov, K. V. Yakovlev, L. V. Olenina, A. A. Kotov *et al.*, 2021 The
920   3'UTR of the Drosophila CPEB translation factor gene orb2 plays a crucial role in
921   spermatogenesis. Development 148.
922   Gotze, M., J. Dufourt, C. Ihling, C. Rammelt, S. Pierson *et al.*, 2017 Translational repression of
923   the Drosophila nanos mRNA involves the RNA helicase Belle and RNA coating by
924   Me31B and Trailer hitch. RNA 23**:** 1552-1568.
925   Gratz, S. J., A. M. Cummings, J. N. Nguyen, D. C. Hamm, L. K. Donohue *et al.*, 2013 Genome
926   engineering of Drosophila with the CRISPR RNA-guided Cas9 nuclease. Genetics 194**:**
927   1029-1035.
928   Gratz, S. J., F. P. Ukken, C. D. Rubinstein, G. Thiede, L. K. Donohue *et al.*, 2014 Highly
929   specific and efficient CRISPR/Cas9-catalyzed homology-directed repair in Drosophila.
930   Genetics 196**:** 961-971.
931   Groisman, I., Y. S. Huang, R. Mendez, Q. Cao and J. D. Richter, 2001 Translational control of
932   embryonic cell division by CPEB and maskin. Cold Spring Harb Symp Quant Biol 66**:**
933   345-351.
934   Hafer, N., S. Xu, K. M. Bhat and P. Schedl, 2011 The Drosophila CPEB protein Orb2 has a
935   novel expression pattern and is important for asymmetric cell division and nervous
936   system function. Genetics 189**:** 907-921.
937   Hake, L. E., and J. D. Richter, 1994 CPEB is a specificity factor that mediates cytoplasmic
938   polyadenylation during Xenopus oocyte maturation. Cell 79**:** 617-627.
939   Harris, T. E., A. Chi, J. Shabanowitz, D. F. Hunt, R. E. Rhoads *et al.*, 2006 mTOR-dependent
940   stimulation of the association of eIF4G and eIF3 by insulin. EMBO J 25**:** 1659-1668.
941   Haugen, R. J., C. Barnier, N. D. Elrod, H. Luo, M. K. Jensen *et al.*, 2024 Regulation of the
942   Drosophila transcriptome by Pumilio and the CCR4-NOT deadenylase complex. RNA
943   30**:** 866-890.
944   Hervás, R., L. Li, A. Majumdar, M. e. C. Fernández-Ramírez, J. R. Unruh *et al.*, 2016 Molecular
945   Basis of Orb2 Amyloidogenesis and Blockade of Memory Consolidation. PLoS Biol 14**:**
946   e1002361.
947   Hervas, R., M. J. Rau, Y. Park, W. Zhang, A. G. Murzin *et al.*, 2020 Cryo-EM structure of a
948   neuronal functional amyloid implicated in memory persistence in *Drosophila*. Science
949   367**:** 1230-1234.
950   Hui, D. J., C. R. Bhasker, W. C. Merrick and G. C. Sen, 2003 Viral stress-inducible protein p56
951   inhibits translation by blocking the interaction of eIF3 with the ternary complex
952   eIF2.GTP.Met-tRNAi. J Biol Chem 278**:** 39477-39482.
953   Hui, D. J., F. Terenzi, W. C. Merrick and G. C. Sen, 2005 Mouse p56 blocks a distinct function
954   of eukaryotic initiation factor 3 in translation initiation. J Biol Chem 280**:** 3433-3440.
955   Ivshina, M., P. Lasko and J. D. Richter, 2014 Cytoplasmic polyadenylation element binding
956   proteins in development, health, and disease. Annu Rev Cell Dev Biol 30**:** 393-415.
957   Jiang, T., R. F. McKinley, M. A. McGill, S. Angers and T. J. Harris, 2015 A Par-1-Par-3-
958   Centrosome Cell Polarity Pathway and Its Tuning for Isotropic Cell Adhesion. Curr Biol
959   25**:** 2701-2708.
960   Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov *et al.*, 2021 Applying and improving
961   AlphaFold at CASP14. Proteins 89**:** 1711-1721.
962   Keleman, K., S. Krüttner, M. Alenius and B. J. Dickson, 2007 Function of the Drosophila CPEB
963   protein Orb2 in long-term courtship memory. Nat Neurosci 10**:** 1587-1593.

Khan, M. R., L. Li, C. Pérez-Sánchez, A. Saraf, L. Florens *et al.*, 2015 Amyloidogenic
Oligomerization Transforms Drosophila Orb2 from a Translation Repressor to an
Activator. Cell 163**:** 1468-1483.
Kim, J. H., and J. D. Richter, 2006 Opposing polymerase-deadenylase activities regulate
cytoplasmic polyadenylation. Mol Cell 24**:** 173-183.
Kolberg, L., U. Raudvere, I. Kuzmin, P. Adler, J. Vilo *et al.*, 2023 g:Profiler-interoperable web
service for functional enrichment analysis and gene identifier mapping (2023 update).
Nucleic Acids Res 51**:** W207-W212.
Kopylova, E., L. Noe and H. Touzet, 2012 SortMeRNA: fast and accurate filtering of ribosomal
RNAs in metatranscriptomic data. Bioinformatics 28**:** 3211-3217.
Kozlov, E., Y. V. Shidlovskii, R. Gilmutdinov, P. Schedl and M. Zhukova, 2021 The role of
CPEB family proteins in the nervous system function in the norm and pathology. Cell
Biosci 11**:** 64.
Kozlov, E. N., E. V. Tokmatcheva, A. M. Khrustaleva, E. S. Grebenshchikov, R. V. Deev *et al.*,
2023 Long-Term Memory Formation in Drosophila Depends on the 3'UTR of CPEB
Gene orb2. Cells 12.
Krüttner, S., B. Stepien, J. N. Noordermeer, M. A. Mommaas, K. Mechtler *et al.*, 2012
Drosophila CPEB Orb2A mediates memory independent of Its RNA-binding domain.
Neuron 76**:** 383-395.
Laver, J. D., K. Ancevicius, P. Sollazzo, J. T. Westwood, S. S. Sidhu *et al.*, 2012 Synthetic
antibodies as tools to probe RNA-binding protein function. Mol Biosyst 8**:** 1650-1657.
Laver, J. D., X. Li, D. Ray, K. B. Cook, N. A. Hahn *et al.*, 2015 Brain tumor is a sequence-
specific RNA-binding protein that directs maternal mRNA clearance during the
Drosophila maternal-to-zygotic transition. Genome Biol 16**:** 94.
Laver, J. D., J. Ly, A. K. Winn, A. Karaiskakis, S. Lin *et al.*, 2020 The RNA-Binding Protein
Rasputin/G3BP Enhances the Stability and Translation of Its Target mRNAs. Cell Rep
30**:** 3353-3367 e3357.
Lee, A. S., P. J. Kranzusch and J. H. Cate, 2015 eIF3 targets cell-proliferation messenger RNAs
for translational activation or repression. Nature 522**:** 111-114.
Lee, A. S., P. J. Kranzusch, J. A. Doudna and J. H. Cate, 2016 eIF3d is an mRNA cap-binding
protein that is required for specialized translation initiation. Nature 536**:** 96-99.
Lee, J. P., A. Brauweiler, M. Rudolph, J. E. Hooper, H. A. Drabkin *et al.*, 2010 The TRC8
ubiquitin ligase is sterol regulated and interacts with lipid and protein biosynthetic
pathways. Mol Cancer Res 8**:** 93-106.
Legge, G. B., M. A. Martinez-Yamout, D. M. Hambly, T. Trinh, B. M. Lee *et al.*, 2004 ZZ
domain of CBP: an unusual zinc finger fold in a protein interaction module. J Mol Biol
343**:** 1081-1093.
Li, X., G. Quon, H. D. Lipshitz and Q. Morris, 2010 Predicting in vivo binding sites of RNA-
binding proteins using mRNA secondary structure. RNA 16**:** 1096-1107.
Liu, G., J. Zhang, H. Choi, J. P. Lambert, T. Srikumar *et al.*, 2012 Using ProHits to store,
annotate, and analyze affinity purification-mass spectrometry (AP-MS) data. Curr Protoc
Bioinformatics Chapter 8**:** 8.16.11-18.16.32.
Liu, G., J. Zhang, B. Larsen, C. Stark, A. Breitkreutz *et al.*, 2010 ProHits: integrated software for
mass spectrometry-based interaction proteomics. Nat Biotechnol 28**:** 1015-1017.

Liu, Y. C., A. L. Couzens, A. R. Deshwar, B. M.-C. LD, X. Zhang *et al.*, 2014 The PPFIA1-
PP2A protein complex promotes trafficking of Kif7 to the ciliary tip and Hedgehog
signaling. Sci Signal 7: ra117.
Love, M. I., W. Huber and S. Anders, 2014 Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol 15: 550.
Love, M. I., C. Soneson, P. F. Hickey, L. K. Johnson, N. T. Pierce *et al.*, 2020 Tximeta:
Reference sequence checksums for provenance identification in RNA-seq. PLoS Comput
Biol 16: e1007664.
Majumdar, A., W. C. Cesario, E. White-Grindley, H. Jiang, F. Ren *et al.*, 2012 Critical role of
amyloid-like oligomers of Drosophila Orb2 in the persistence of memory. Cell 148: 515-
529.
Martineau, Y., M. C. Derry, X. Wang, A. Yanagiya, J. J. Berlanga *et al.*, 2008 Poly(A)-binding
protein-interacting protein 1 binds to eukaryotic translation initiation factor 3 to stimulate
translation. Mol Cell Biol 28: 6658-6667.
Martineau, Y., X. Wang, T. Alain, E. Petroulakis, D. Shahbazian *et al.*, 2014 Control of Paip1-
eukayrotic translation initiation factor 3 interaction by amino acids through S6 kinase.
Mol Cell Biol 34: 1046-1053.
Mastushita-Sakai, T., E. White-Grindley, J. Samuelson, C. Seidel and K. Si, 2010 Drosophila
Orb2 targets genes involved in neuronal growth, synapse formation, and protein turnover.
Proc Natl Acad Sci U S A 107: 11987-11992.
Meng, E. C., T. D. Goddard, E. F. Pettersen, G. S. Couch, Z. J. Pearson *et al.*, 2023 UCSF
ChimeraX: Tools for structure building and analysis. Protein Sci 32: e4792.
Merkel, D. J., S. B. Wells, B. C. Hilburn, F. Elazzouzi, G. C. Pérez-Alvarado *et al.*, 2013 The C-
terminal region of cytoplasmic polyadenylation element binding protein is a ZZ domain
with potential for protein-protein interactions. J Mol Biol 425: 2015-2026.
Montojo, J., K. Zuberi, H. Rodriguez, F. Kazi, G. Wright *et al.*, 2010 GeneMANIA Cytoscape
plugin: fast gene function predictions on the desktop. Bioinformatics 26: 2927-2928.
Na, H., J. D. Laver, J. Jeon, F. Singh, K. Ancevicius *et al.*, 2016 A high-throughput pipeline for
the production of synthetic antibodies for analysis of ribonucleoprotein complexes. RNA
22: 636-655.
Nakamura, A., K. Sato and K. Hanyu-Nakamura, 2004 Drosophila cup is an eIF4E binding
protein that associates with Bruno and regulates oskar mRNA translation in oogenesis.
Dev Cell 6: 69-78.
Nelson, M. R., A. M. Leidal and C. A. Smibert, 2004 Drosophila Cup is an eIF4E-binding
protein that functions in Smaug-mediated translational repression. The EMBO journal
23: 150-159.
Nelson, M. R., H. Luo, H. K. Vari, B. J. Cox, A. J. Simmonds *et al.*, 2007 A multiprotein
complex that mediates translational enhancement in Drosophila. J Biol Chem 282:
34031-34038.
Ozturk-Colak, A., S. J. Marygold, G. Antonazzo, H. Attrill, D. Goutte-Gattat *et al.*, 2024
FlyBase: updates to the Drosophila genes and genomes database. Genetics 227.
Patro, R., G. Duggal, M. I. Love, R. A. Irizarry and C. Kingsford, 2017 Salmon provides fast and
bias-aware quantification of transcript expression. Nat Methods 14: 417-419.
Pestova, T. V., and C. U. Hellen, 1999 Internal initiation of translation of bovine viral diarrhea
virus RNA. Virology 258: 249-256.

Pestova, T. V., I. N. Shatsky, S. P. Fletcher, R. J. Jackson and C. U. Hellen, 1998 A prokaryotic-
like mode of cytoplasmic eukaryotic ribosome binding to the initiation codon during
internal translation initiation of hepatitis C and classical swine fever virus RNAs. Genes
Dev 12**:** 67-83.
Ray, D., H. Kazan, K. B. Cook, M. T. Weirauch, H. S. Najafabadi *et al.*, 2013 A compendium of
RNA-binding motifs for decoding gene regulation. Nature 499**:** 172-177.
Semotok, J. L., R. L. Cooperstock, B. D. Pinder, H. K. Vari, H. D. Lipshitz *et al.*, 2005 Smaug
recruits the CCR4/POP2/NOT deadenylase complex to trigger maternal transcript
localization in the early Drosophila embryo. Curr Biol 15**:** 284-294.
Shannon, P., A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang *et al.*, 2003 Cytoscape: a software
environment for integrated models of biomolecular interaction networks. Genome Res
13**:** 2498-2504.
Standart, N., and N. Minshall, 2008 Translational control in early development: CPEB, P-bodies
and germinal granules. Biochem Soc Trans 36**:** 671-676.
Stepien, B. K., C. Oppitz, D. Gerlach, U. Dag, M. Novatchkova *et al.*, 2016 RNA-binding
profiles of Drosophila CPEB proteins Orb and Orb2. Proc Natl Acad Sci U S A 113**:**
E7030-E7038.
Stoiber, M. H., S. Olson, G. E. May, M. O. Duff, J. Manent *et al.*, 2015 Extensive cross-
regulation of post-transcriptional regulatory networks in Drosophila. Genome Res 25**:**
1692-1702.
Tadros, W., A. L. Goldman, T. Babak, F. Menzies, L. Vardy *et al.*, 2007 SMAUG is a major
regulator of maternal mRNA destabilization in Drosophila and its translation is activated
by the PAN GU kinase. Dev Cell 12**:** 143-155.
Tadros, W., and H. D. Lipshitz, 2009 The maternal-to-zygotic transition: a play in two acts.
Development 136**:** 3033-3042.
Thomsen, S., S. Anders, S. C. Janga, W. Huber and C. R. Alonso, 2010 Genome-wide analysis
of mRNA decay patterns during early Drosophila development. Genome Biol 11**:** R93.
Vastenhouw, N. L., W. X. Cao and H. D. Lipshitz, 2019 The maternal-to-zygotic transition
revisited. Development 146.
Wang, X. P., and N. G. Cooper, 2009 Characterization of the transcripts and protein isoforms for
cytoplasmic polyadenylation element binding protein-3 (CPEB3) in the mouse retina.
BMC Mol Biol 10**:** 109.
Warde-Farley, D., S. L. Donaldson, O. Comes, K. Zuberi, R. Badrawi *et al.*, 2010 The
GeneMANIA prediction server: biological network integration for gene prioritization and
predicting gene function. Nucleic Acids Res 38**:** W214-220.
Whitten, D., 2011 Classification and clustering of sequencing data using a Poisson model, pp.
2493-2518. Annals of Applied Statistics.
Wilhelm, J. E., M. Hilton, Q. Amos and W. J. Henzel, 2003 Cup is an eIF4E binding protein
required for both the translational repression of oskar and the recruitment of Barentsz. J
Cell Biol 163**:** 1197-1204.
Xu, S., N. Hafer, B. Agunwamba and P. Schedl, 2012 The CPEB protein Orb2 has multiple
functions during spermatogenesis in Drosophila melanogaster. PLoS Genet 8**:** e1003079.
Zavortink, M., L. N. Rutt, S. Dzitoyeva, J. C. Henriksen, C. Barrington *et al.*, 2020 The E2
Marie Kondo and the CTLH E3 ligase clear deposited RNA binding proteins during the
maternal-to-zygotic transition. Elife 9.

1098

Figure Legends

Figure 1. RIP-seq identifies over 450 ORB2 targets in 0-3 hr embryos. (a) Domain structure of ORB2B and location of the antigen used to screen for phage-displayed synthetic antibodies. (b) Scatterplot depicting the $\log_2$(Fold Enrichment)/Input Abundance (TPM) for all genes identified in the RIP-seq analysis; each point represents a single gene. Values represent the average enrichment from three independent biological replicates. ORB2 targets (red) are defined as having fold enrichment >1.5 in the anti-ORB2 RIP-seq compared to input and a ratio of ORB2/Control > 1.5, with an FDR adjusted P-value < 0.05; genes enriched in both ORB2 and control RIP-seq (green) are defined as having fold enrichment >1.5 in both anti-ORB2 and control RIP-seq, with a ratio of ORB2/Control < 1.5 and an FDR adjusted P-value < 0.05; Control enriched transcripts (blue) are defined as having fold enrichment <-1.5 in the control RIP-seq, fold enrichment < -1.5 in the anti-ORB2 RIP-seq, and an FDR adjusted P-value < 0.05. Dotted lines indicate ±1.5 fold-change thresholds. (c) Violin and boxplots depicting the distribution of transcript abundance as measured by average TPM from three biological replicates based on bound status as described in (A). Wilcoxon Rank Sum Test with Bonferroni multiple-test correction: ** = $P < 10^{-10}$; *** = $P < 10^{-15}$. (d) Validation of RIP-seq targets by RT-qPCR; bar graphs depict the $\log_2$(fold enrichment) in an anti-ORB2 RIP-RT-qPCR compared to control RIP-RT-qPCR; P-values given are derived from Wilcoxon Signed Rank Test comparing ORB2 RIP-RT-qPCR to control RIP-RT-qPCR.

Figure 2. ORB2-bound transcripts are enriched in translationally repressed, unstable transcripts during the MZT. Violin and boxplots depicting the relationship between ORB2-association and translational status, change in poly(A) tail length, and change in transcript abundance. Unbound genes include all expressed genes that were not defined as ORB2 targets in the RIP-seq analysis. (a) Translation efficiency, calculated from ribosome footprinting performed on 0-1 hour embryo extracts (EICHHORN *et al.* 2016). (b) Translation efficiency, calculated from ribosome footprinting performed on 2-3 hour embryo extracts (EICHHORN *et al.* 2016). (c) Translation efficiency as measured by ribosome profiling performed on 0-2 hour embryo extracts (DUNN *et al.* 2013). (d) Translation index, calculated from polysome gradients performed on 0-3 hour embryo extracts (CHEN *et al.* 2014). (e) Change in transcript poly(A) tail length, calculated by comparing 0-1 hour and 2-3 hour embryo extracts, as measured by PAL-Seq (EICHHORN *et al.* 2016). (f) Change in abundance, calculated by comparing 0-1 hour and 2-3 hour embryo extracts, as measured by RNA-Seq (EICHHORN *et al.* 2016). Wilcoxon Rank Sum Test: ** = $P < 10^{-10}$; *** = $P < 10^{-15}$.

Figure 3. ORB2 binding motifs predicted from RIP-seq confer translational repression on a reporter mRNA in S2 cells. (a) ORB2 recognition element (ORE) enriched in ORB2 targets identified by *de novo* motif discovery using #ATS (LI *et al.* 2010) on results from the anti-ORB2 RIP-seq in this study. (b) Diagram depicting the reporter Firefly Luciferase mRNA with six copies of different ORB2-binding motifs in the 3′UTR. (c) Bar graph depicting the enrichment of the luciferase reporter mRNA in an anti-ORB2 RIP-qPCR compared to Renilla Luciferase mRNA, normalized to a reporter without ORE motifs. Mutation of either the OREs or deletion of the RNA-binding domain (RRM) abrogate binding to the mRNA. (d) Bar graph depicting the ratio of luminescence between Firefly luciferase and Renilla luciferase. Mutation of either the OREs or deletion of the RNA-binding domain (RRM) abrogate repression. One-way ANOVA with Tukey multiple test correction: * = $P < 0.05$.

Figure 4. The ORB2 and hCPEB2 ZZ domain but not the ORB and hCPEB1 ZZ domain
repress a luciferase reporter. (a) Diagram depicting the tethering assay in which a polypeptide
of interest fused to the BIV Tat domain is tethered to a Firefly luciferase reporter mRNA
carrying six tandem repeat TAR elements in the 3′UTR. (b) Bar graph depicting the expression
of the reporter firefly luciferase (normalized to co-transfected Renilla luciferase and normalized
to steady state RNA levels) when tethering whole ORB2 or ORB2 fragments compared to GFP
control. The ORB2 ZZ domain is necessary and sufficient for repression. (c) Bar graph depicting
the expression of the reporter firefly luciferase mRNA measured by RT-qPCR (normalized to co-
transfected Renilla luciferase mRNA) when tethering whole ORB2 or ORB2 fragments
compared to GFP control. Reporter mRNA levels are unaffected by ORB2 ZZ or ORB2ΔZZ. (d)
amino acid sequence alignment of the ZZ domains of ORB2, ORB, and their human orthologs,
hCPEB1 and hCPEB2. The arrowhead denotes the junction between the RBD and ZZ domains.
(e) Bar graph depicting the expression of the reporter firefly luciferase (normalized to co-
transfected Renilla luciferase) when tethering different CPEB ZZ domains. Only ORB2 and
hPEB2 repress. (f) Bar graph depicting the expression of the reporter firefly luciferase when
ORB ZZ chimeric domains harboring substitutions from the ORB2 ZZ domain, are tethered. The
ORB2 'A' and 'C' regions are each sufficient to convert the ORB ZZ domain into a repressor.
One-way ANOVA with Hold-Sidak multiple test correction: $* = P < 0.05$; $** = P < 0.001$; $*** =$
$P < 10^{-5}$.

Figure 5. Anti-ORB2 IP-MS from early embryo extracts identifies known translation
repressors and the PIC as binding partners. (a) Interaction network depicting the connections
between components of the Cup repressive complex and other translational repressors identified
as ORB2 interactors by SAINT analysis. (b) Interaction network depicting the connections
between components of eukaryotic PIC identified as ORB2 interactors by SAINT analysis.
Interaction networks were produced by the GeneMANIA app for Cytoscape (FRANZ *et al.* 2018);
size of node indicates average number of spectral counts obtained in the IP-MS; thickness of
edges indicates strength of interaction assigned by GeneMANIA.

Figure 6. Deletion of the ORB2 ZZ domain results in derepression of target RNAs in the
early embryo. (a) Violin and boxplots comparing target and non-target abundance as determined
by RNA-Seq in 0-1, 1-2 and 2-3 hour old embryos laid by w^{1118} and *orb2*$^{\Delta ZZ}$ females. No
significant difference was detected for either genotype at any of the time-points. (b) Violin and
boxplots comparing target and non-target translation index as determined by polysome gradients
followed by RNA-Seq in 0-1, 1-2 and 2-3 hour old embryos laid by w^{1118} and *orb2*$^{\Delta ZZ}$ females.
ORB2 targets were significantly derepressed in 0-1 hour old *orb2*$^{\Delta ZZ}$ embryos relative to w^{1118}.
Log2(Ratio of mean normalized counts of pool 3 /mean normalized counts of pool 1 mean) was
calculated for each transcript. Wilcoxon Rank Sum Test: $** = P < 0.006$.

Table Legends

Table 1. Overlap between the list of ORB2 targets and lists of genes associated with mRNA
stability or decay in the early embryo.

Table 2. Molecular Function GO-terms enriched in ORB2 targets. GO-term analysis was
performed using the gProfiler2 tool (KOLBERG *et al.* 2023).

Table 3. Biological Process GO-terms enriched in ORB2 targets. GO-term analysis was performed using the gProfiler2 tool (KOLBERG *et al.* 2023).

Table 4. Cellular Component GO-terms enriched in ORB2 targets. GO-term analysis was performed using the gProfiler2 tool (KOLBERG *et al.* 2023).

Supplemental Figure Legends

Figure S1. Replicate-to-replicate comparisons of transcript TPMs from RIP-seq experiments show high correlation between replicates. (a)-(c) Pairwise comparisons of three biological replicates performed with the anti-ORB2 E8 Fab. The Pearson correlation coefficient and its corresponding *P*-value were calculated and used to determine the degree of similarity between each pair. (d) Heatmap of the Poisson distance between three biological RIP-seq replicates performed with anti-ORB2 E8 Fab and three biological RIP-seq replicates performed with control C1 Fab. The Poisson distance is calculated by the difference in Poisson distribution between pairs of samples; a larger distance indicates a greater level of dissimilarity between samples.

Figure S2. Expression levels of ORB2, ORB2ΔRRM, and GFP in S2 cells. Each set of three lanes corresponds to the ORE validation experiments in which expression vectors for tagged proteins and reporter mRNA constructs containing various recognition motifs were transfected into S2 cells, described in Figure 3.

Figure S3. Expression of ORB2 full-length protein, portions of ORB2, and the ZZ domains of ORB, ORB2, hPCPEB1 and 2 in S2 cells. (a) Western blot showing the expression of ORB2 full-length protein and protein portions corresponding to the experiments described in Figure 4b and c. (b) Western blot showing the expression of ORB2B and ORB2BΔZZ to the experiments described in Figure 4b and c. (c) Western blot showing the expression of ORB2 ZZ, hCPEB2 ZZ, ORB ZZ and hCPEB1 ZZ corresponding to the experiments described in Figure 4e.

Figure S4. Structure of ZZ domains of CPEB family members. (a) The NMR structure of the ZZ domain of human CPEB1 Protein Data Bank accession 2M13 (MERKEL *et al.* 2013) is shown, along with the Alphafold 3 (ABRAMSON *et al.* 2024) predicted structures of the ZZ domains of ORB, hCPEB2 and ORB2. The colors in the Alphafold 3 predictions represent the predicted local distance difference test (pLDDT) scores, where higher numbers indicate greater confidence in the prediction. (b) On the left is an overlap of the ORB and ORB2 predicted structures with Region A (the Rd turn), Region B (the beta hairpin), and Region C (the alpha helix) of each protein indicated. The boxes on the right highlight each region individually in two different orientations.

Figure S5. The predicted structures of chimeric ZZ domains. Alphafold 3 (ABRAMSON *et al.* 2024) predictions for chimeric ZZ domains where one of Regions A, B or C of ORB was replaced with the corresponding region from ORB2. The top row of images indicates the identity each construct as described in Figure 4f, while the bottom row shows the pLDDT scores for each prediction.

Figure S6. Expression of chimeric ORB-ORB2 ZZ domain proteins in S2 cells. Western blot showing the expression of ORB-ORB2 chimeric proteins corresponding to the substitution

experiments which assessed transformation of the non-repressing ORB ZZ domain into a
translational repressor, as shown in Figure 4f.

Figure S7. CRISPR Cas9-mediated endogenous deletion of the ORB2 ZZ domain was used
to generate *orb2*$^{\Delta ZZ}$ flies. (a) Diagram depicting the *orb2* endogenous locus, located at 66E4-
66E5 on 3L, the corresponding mRNAs that are encoded, and the targeted deletion. (b) Agarose
gel of the PCR amplification of the region surrounding the CRISPR targeted deletion of the F1
generation post-injection. Genotypes indicate lines generated from individual F1 flies. Up PCR
indicates PCR amplification using primers designed against the 5′ insertion junction; Down PCR
indicates PCR amplification using primers designed against the 3′ insertion junction. (c) Agarose
gel of the PCR amplification of the region surrounding the PiggyBac transposon-mediated
excision of the dsRed marker. WT indicates the band representing PCR amplification of the
endogenous ORB2 locus containing an intact ZZ domain; Excised indicates the band
representing amplification of the locus with the ZZ domain deletion without the dsRed marker.
Genotypes indicate lines generated from individual flies resulting from the CRISPR injection. (d)
Western blot showing expression of endogenous ORB2 and CRISPRed ORB2ΔZZ in
heterozygous and homozygous whole flies. ORB2ΔZZ is expressed at higher levels than full-
length ORB2 (see heterozygote for comparison). Generation of CRISPR mutants was performed
by WellGenetics as described in the Materials and Methods.

Figure S8. ORB2ΔZZ retains RNA-binding to its targets in embryos. (a) Western blot
showing the expression of ORB2 and ORB2ΔZZ protein over the first four hours of embryonic
development. (b) Bar graph depicting the enrichment of five ORB2 target mRNAs in RIP-RT-
qPCR from 0-3 hour old embryos laid by w^{1118} and *orb2*$^{\Delta ZZ}$ females. The ORB2 RIPs used the
ORB2 E8 Fab; the control RIP used the C1 Fab. RNA levels were normalized to *RpL32* mRNA.
n = 3; error bars are standard deviation. Student's *t*-test *P*-values were not significant (n.s.; *P* >
0.05)

Supplemental File Legends

File S1. Results of RIP-seq of ORB2 in 0-3 hour embryos. See Materials and Methods for
details.

File S2. Results of *de novo* motif discovery on ORB2 targets in 0-3 hour embryos. Motif
discovery was carried out using #ATS as described (LI *et al.* 2010). See Materials and Methods
for details.

File S3. Results of IP-MS of ORB2 and ORB2ΔZZ in 0-3 hour embryos in the presence or
absence of RNase A. Interactors were identified using SAINT as described (CHOI *et al.* 2012).
See Materials and Methods for details.

File S4. Lists of mRNAs bound by ORB2 (RIP-seq) whose protein products were also
identified as interactors in ORB2 IP-MS.

File S5. Results of RNA-seq time-course in wild-type and ORB2ΔZZ embryos. See Materials
and Methods for details.

File S6. Results of polysome gradient RNA-seq time-course in wild-type and ORB2ΔZZ

embryos. See Materials and Methods for details.

FIGURE 1

a

b

c

d

FIGURE 2

FIGURE 3

a ORE derived from RIP-Seq

b

Six tandem copies of ORB2 motif in 3'UTR

c RNA enrichment in ORB2 IP
☐ GFP ☐ ORB2 ☐ ORB2ΔRRM

d Luminescence of Luciferase reporter
☐ GFP ☐ ORB2 ☐ ORB2ΔRRM

FIGURE 4

A

B

Control (GFP)
ORB2B ***
IDR1 n.s.
PolyQ n.s.
IDR1+PolyQ n.s.
IDR2 n.s.
RBD *
ZZ ***
ORB2ΔZZ n.s.

FLuc/RenLuc Luminescence

C

Control (GFP)
ORB2B n.s.
IDR1 *
PolyQ n.s.
IDR1+PolyQ n.s.
IDR2 n.s.
RBD n.s.
ZZ n.s.
ORB2ΔZZ n.s.

FLuc/RenLuc RNA

D

```
                    RRM end  ▼        Rd Turn      β1       β2       α1
ORB2 ZZ     KRVEVKPYVLDDQMCDECEGQRCGGKFAPFFCANVTCLQYYCEHCWAVIHSRPGREYHKPLVKEGADRPRAVPFRWC
hCPEB2 ZZ   KRVEVKPYVLDDQMCDECGARCGGKFAPFFCANVTCLQYYCEPCWANIHSRAGREFHKPLVKEGADRPRQIHFRWN
ORB ZZ      KKVQVDPY-LEDALCSICGVQHG-----PYYCRELSCFRYFCRSCWQWQHSCDIVKNHKPLRNSKSQSLVGIGPSS
hCPEB1 ZZ   KKVQHDPY-LEDSLCHICSSQPG-----PFFCRDQVCFKYFCRSCWHWRHSMEGLRHHSPLVRNQKNRDSS------
```

■ Zinc cooridnating residue
■ Different from ORB2
■ Conserved between ORB and hCPEB1 but different from ORB2

Region A   Region B   Region C

E

Control (GFP)
ORB ZZ n.s.
ORB2 ZZ ***
hCPEB1 ZZ n.s.
hCPEB2 ZZ ***

FLuc/RenLuc Luminescence

F

Control (GFP)
ORB ZZ n.s.
ORB2 ZZ ***
ORB ZZ-A²B²C² **
ORB ZZ-A²B¹C¹ ***
ORB ZZ-A¹B²C¹ n.s.
ORB ZZ-A¹B¹C² **

Region A Region B Region C

FLuc/RenLuc Luminescence

a

b

FIGURE 6

TABLE 1

Comparison	Backgound (Total)	Orb2 Only	Other Only	Overlap	Fisher's Exact p-value	Odds Ratio
Maternal Unstable (Unfertilized eggs, Tadros et al 2007)	3030	172	744	166	2.20E-16	3.849738
Thomsen et al 2010						
Class I (Stable transcripts)	4186	377	1335	35	2.20E-16	0.2334165
Class II (Maternal Decay only)	5108	395	394	36	0.3359	1.181531
Class III (Maternal Decay + Transcription)	5177	392	326	38	0.02109	1.539281
Class IV (Zygotic Decay only)	4332	322	1171	108	0.06763	1.240746
Class V (Both Maternal + Zygotic Decay)	4186	239	1317	191	2.20E-16	2.53962

TABLE 2. Molecular Function

GO Term Name	Adjusted p-value	GO Term Size	ORB2 targets	Overlap
protein binding	1.13E-21	3760	416	229
mRNA 3'-UTR binding	5.78E-11	48	416	17
mRNA binding	1.15E-10	238	416	35
mRNA regulatory element binding translation repressor activity	2.51E-08	18	416	10
translation repressor activity	7.18E-08	25	416	11
translation regulator activity	2.28E-04	106	416	16
catalytic activity, acting on DNA	2.49E-04	147	416	19
protein serine/threonine kinase activity	5.25E-04	184	416	21
DNA binding	6.42E-04	1127	416	69
poly-pyrimidine tract binding	1.28E-03	14	416	6
adenyl ribonucleotide binding	1.42E-03	846	416	55
catalytic activity, acting on a nucleic acid	1.62E-03	406	416	33
ATP binding	2.44E-03	840	416	54
purine ribonucleotide binding	2.99E-03	1021	416	62
purine nucleotide binding	3.63E-03	1072	416	64
adenyl nucleotide binding	3.73E-03	896	416	56
ribonucleotide binding	4.31E-03	1033	416	62

TABLE 3. Biological Process

GO Term Name	adjusted p-value	GO Term Size	ORB2 targets	Overlap
purine ribonucleoside triphosphate binding	4.48E-03	1012	416	61
anatomical structure development	5.77E-28	2775	435	212
developmental process	2.30E-27	2954	435	219
negative regulation of cellular process	1.73E-25	1520	435	144
cell development	6.92E-25	1450	435	139
multicellular organism development	8.26E-25	2276	435	182
cell differentiation	3.46E-24	1768	435	155
cellular component organization	1.07E-21	3178	435	217
regulation of metabolic process	1.81E-21	2253	435	174
gamete generation	1.19E-20	917	435	100
germ cell development	7.67E-20	721	435	86
anatomical structure morphogenesis	1.34E-19	1580	435	136
oogenesis	1.45E-19	582	435	76
system development	5.09E-19	1527	435	132
multicellular organismal reproductive process	1.60E-18	1096	435	107
regulation of primary metabolic process	1.69E-17	1860	435	146
nervous system development	4.18E-17	1196	435	110
animal organ development	5.43E-17	1200	435	110
regulation of macromolecule biosynthetic process	1.09E-15	1742	435	136

TABLE 4. Cellular Component

GO Term Name	adjusted p-value	GO Term Size	ORB2 targets	Overlap
nucleus	1.99E-14	3234	438	196
ribonucleoprotein granule	2.57E-12	119	438	27
cytoplasmic ribonucleoprotein granule	7.11E-12	114	438	26
cellular anatomical structure	6.48E-10	10708	438	426
cytoplasm	6.02E-09	5187	438	255
P granule	1.14E-07	46	438	14
cytosol	1.14E-07	1131	438	82
germ plasm	2.15E-07	48	438	14
nucleoplasm	1.40E-06	573	438	50
pole plasm	2.54E-06	57	438	14
nuclear lumen	3.31E-06	910	438	67
cytoplasmic stress granule	5.07E-06	21	438	9
protein-containing complex	2.53E-05	2861	438	151
chromosome	3.50E-05	787	438	58
PcG protein complex	8.68E-04	27	438	8
transferase complex	2.13E-03	550	438	41
intracellular protein-containing complex	4.15E-03	625	438	44
cell junction	4.67E-03	608	438	43
intracellular organelle lumen	4.94E-03	1158	438	69
membrane-enclosed lumen	5.08E-03	1159	438	69
organelle lumen	5.08E-03	1159	438	69
cell cortex	5.36E-03	208	438	21