The ORB2 RNA-binding protein negatively regulates its target transcripts during the Drosophila maternal-to-zygotic transition via its functionally conserved Zinc-binding 'ZZ' domain

Timothy C.H. Low[1], Hua Luo[1], Chalini Weerasooriya[2], Zhuyi Wang[1], Sichun Lin[3], Angelo Karaiskakis[1], Stephane Angers[2,3], Craig A. Smibert[1,2], Howard D. Lipshitz[1]*

[1]Department of Molecular Genetics, University of Toronto, 661 University Avenue, Toronto Ontario, Canada M5G 1M1

[2]Department of Biochemistry, University of Toronto, 661 University Avenue, Toronto Ontario, Canada M5G 1M1

[3]Department of Pharmaceutical Sciences & Donnelly Centre, University of Toronto, 160 College Street, Toronto, Ontario, Canada M5S 3E1

*Author for correspondence: howard.lipshitz@utoronto.ca

Running head: ORB2 negatively regulates mRNAs during the MZT

Keywords: ORB2, Cytoplasmic Polyadenylation Element Binding Protein (CPEB), RNA-binding protein (RBP), translational regulation, post-transcriptional regulation, pre-initiation complex (PIC), embryo, maternal-to-zygotic transition (MZT), S2 cell, Cup.

ARTICLE SUMMARY

We show that Drosophila ORB2, the ortholog of the human CPEB2 RNA-binding protein, negatively regulates its target mRNAs during the maternal-to-zygotic transition via its C-terminal Zinc-binding ('ZZ') domain.

ABSTRACT

Post-transcriptional regulation is particularly prominent during the maternal-to-zygotic transition (MZT), a developmental phase during which a large proportion of maternally provided mRNAs is repressed and cleared from metazoan embryos. RNA-binding proteins (RBPs) are key components of the post-transcriptional regulatory machinery. We show that the ORB2 RBP, the Drosophila ortholog of human Cytoplasmic Polyadenylation Element Binding Protein (hCPEB) 2-4 protein subfamily, binds to hundreds of maternally provided, rare-codon-enriched mRNAs in early embryos; that ORB2 targets are translationally repressed and unstable during the MZT; identify a U-rich motif enriched in ORB2 targets' 3′UTRs; and show that this motif confers ORB2 binding and repression to a luciferase reporter mRNA in S2 tissue culture cells. When tethered to a luciferase reporter, ORB2 and hCPEB2 (but not ORB and hCPEB1) repress translation; the C-terminal Zinc-binding ('ZZ') domain of ORB2 is necessary and sufficient for repression. ORB2 interacts with a suite of post-transcriptional regulators in early embryos; a subset of these interactions is lost upon deletion of the ZZ domain, notably with the Cup repressive complex. ORB2-targets significantly overlap with those previously identified for the repressive RBP, Smaug (SMG). Analysis of the early embryo's translatome in the presence or absence of the endogenous ZZ domain shows that mRNAs bound by ORB2 but not by SMG move onto polysomes upon ZZ domain deletion whereas co-bound transcripts do not, consistent with co-regulation of the latter set of transcripts by both RBPs. Our results assign a function to the ZZ domain and position ORB2 in the post-transcriptional network that regulates maternal transcripts during the Drosophila MZT.

INTRODUCTION

All animal embryos undergo a maternal-to-zygotic transition (MZT) that encompasses two global processes: First, a large subset of maternal gene products loaded into the egg by the female parent is cleared; second, the zygotic genome becomes transcriptionally active (TADROS

AND LIPSHITZ 2009; VASTENHOUW *et al.* 2019; HARRISON *et al.* 2023). In Drosophila embryos,

the MZT is both very rapid – reaching completion about 3 hours after fertilization – and

extensive: Thousands of maternally loaded transcripts are cleared (DE RENZIS *et al.* 2007;

LÉCUYER *et al.* 2007; TADROS *et al.* 2007; THOMSEN *et al.* 2010); hundreds of these transcripts

are translationally repressed (KRONJA *et al.* 2014); and thousands of genes are transcribed from

the embryo's own genome in a process known as zygotic genome activation (ZGA) (HARRISON

et al. 2023). Several RBPs have been shown to function in maternal transcript repression and

degradation in *Drosophila*: Smaug (SMG) directs repression and clearance of a subset of

maternal transcripts prior to ZGA (TADROS *et al.* 2007; CHEN *et al.* 2014); Brain tumor (BRAT)

binds to and represses a distinct subset of maternal mRNAs both before and after ZGA (LAVER

et al. 2015); Pumilio (PUM) co-binds a subset of the BRAT targets but primarily directs

transcript clearance after rather than before ZGA (THOMSEN *et al.* 2010; LAVER *et al.* 2015;

HAUGEN *et al.* 2024). In contrast to these 'repressive' RBPs, Rasputin (RIN), the Drosophila

ortholog of vertebrate G3BP, upregulates the stability and translation of its targets during the

MZT (LAVER *et al.* 2020a).

Here we focus on the ORB2 RBP, which is of particular interest because it and its

vertebrate homologs have been reported to mediate both negative and positive post-

transcriptional regulatory functions in a variety of organisms and cell types. ORB and ORB2 are

the fly homologs of the vertebrate Cytoplasmic Polyadenylation Element Binding (CPEB) family

of RBPs, which canonically interact with U-rich motifs known as cytoplasmic polyadenylation

elements (CPEs) in the 3′UTR of target transcripts (HAKE AND RICHTER 1994). Vertebrates have

four CPEBs (CPEB1-4), which can be divided into two groups based on sequence similarity: the

CPEB1-like subfamily (containing vertebrate CPEB1 and its homologs, such as Drosophila

ORB) and the CPEB2-like subfamily (containing vertebrate CPEB2-4 and their homologs, such

as Drosophila ORB2) (WANG AND COOPER 2009; FERNÁNDEZ-MIRANDA AND MÉNDEZ 2012;

IVSHINA *et al.* 2014; DURAN-ARQUÉ *et al.* 2022).

CPEB proteins are known to function in the regulation of cytoplasmic mRNAs involved in

egg maturation, embryonic stem cell division, memory formation, synaptic plasticity, and

gametogenesis (HAKE AND RICHTER 1994; GROISMAN *et al.* 2001; KELEMAN *et al.* 2007;

STANDART AND MINSHALL 2008; KRÜTTNER *et al.* 2012; XU *et al.* 2012). Xenopus CPEB1, for

example, is a major specificity factor of an mRNA masking complex that represses translation in

oocytes and mediates translational activation of its target mRNAs upon egg activation (HAKE

AND RICHTER 1994; KIM AND RICHTER 2006).

ORB2 has been shown to have a role in neural development, long-term memory formation,

and spermatogenesis (KELEMAN *et al.* 2007; HAFER *et al.* 2011; MAJUMDAR *et al.* 2012; XU *et al.*

2012; KHAN *et al.* 2015; HERVÁS *et al.* 2016). The *orb2* gene encodes two protein isoforms,

ORB2A and ORB2B, which have 542 amino acids in common (from N-to-C): a poly-glutamine

(polyQ) tract, a predicted intrinsically disordered region (IDR2) that spans approximately 200

amino acids, the RNA-binding domain (RBD) comprised of tandem RNA recognition motif

(RRM) domains, and a ZZ-class Zinc-binding domain. ORB2A and ORB2B differ only in IDR1

at their N-termini (ORB2B is diagrammed in Figure 1a). ORB2A-ORB2B hetero-oligomers

serve as translational activators whereas, in the absence of ORB2A, ORB2B does not readily

oligomerize and has been shown to repress translation of reporter transcripts *in vitro* and in S2

tissue culture cells (MASTUSHITA-SAKAI *et al.* 2010; KHAN *et al.* 2015; STEPIEN *et al.* 2016;

HERVAS *et al.* 2020). The early Drosophila embryo is unusual in that only one of the two protein

isoforms, ORB2B, is expressed, affording us the opportunity to assess ORB2B's mechanism of

action and function *in vivo* in the absence of ORB2A.

To identify protein and RNA components of ORB2B-containing ribonucleoprotein

complexes (RNPs) during the MZT, we used a synthetic antibody to immunoprecipitate (IP)

endogenously expressed ORB2-containing RNPs followed by either next-generation sequencing

of co-immunoprecipitated transcripts (RIP-seq) or mass spectrometry of associated proteins (IP-

MS). We show that ORB2 binds hundreds of mRNAs that are translationally repressed and

degraded during the MZT. These transcripts are enriched for rare codons in their open reading

frames and for a U-rich motif in their 3′UTRs. We identify over 200 protein binding partners of

ORB2, including components of the 43S translation pre-initiation complex (PIC) and the Cup-

ME31B-TRAL-BEL repressive complex (hereafter referred to as the Cup repressive complex).

We leverage cell culture assays to verify that ORB2 binding to the U-rich motif results in

translational repression of a reporter RNA; to map ORB2's repressive function to the C-terminal

ZZ domain; and to show that the repressive function of the ZZ domain is conserved in its

subfamily ortholog, human CPEB2 (hCPEB2), but not in members of the other subfamily, ORB

or hCPEB1. We produce a CRISPR deletion of the ZZ domain of endogenous ORB2 and show

that deletion of the ZZ domain results in loss of interaction with the PIC and the Cup repressive

complex and a shift of ORB2's targets onto polysomes in early embryos. We show that a subset

of ORB2's targets is co-bound by the SMG RBP. Loss of the ZZ domain results in 'ORB2 only'

targets (i.e., those not co-bound by SMG) moving onto polysomes; in contrast, ORB2-SMG co-

bound targets do not. Together our results provide a function for the ZZ domain and position

ORB2 within the RBP regulatory network that controls the post-transcriptional fate of maternal

mRNAs. We report elsewhere that the endogenous deletion of the ZZ domain leads to defects in

spermatogenesis (LOW *et al.* 2025) and brain development (HAILSTOCK *et al.* 2025).

MATERIALS AND METHODS

Drosophila stocks

The following *Drosophila melanogaster* stocks were used: w^{1118} and $orb^{\Delta ZZ}$. The latter

stock was generated for use in this study (see below). Flies were cultivated at 25°C under

standard laboratory conditions unless otherwise indicated.

Generation of the CRISPR deletion of the ZZ domain at the ORB2 endogenous locus

Generation of this CRISPR line was performed by Well Genetics (13F.-12, No. 93, Sec. 1,

Xintai 5th Rd., Xizhi Dist., New Taipei City 221416, Taiwan (R.O.C.)). The scarless PBac-

dsRed method was used to produce the $orb2^{\Delta ZZ}$ line (GRATZ *et al.* 2013; GRATZ *et al.* 2014). The

ZZ domain, defined as from D638 to C704 (using the numbering from the ORB2-PB isoform)

was targeted for deletion such that the endogenous 3'UTR was retained in its entirety. Two guide

RNAs (gRNAs) were used to direct CRISPR to make double-stranded cuts in the DNA

surrounding the mutation of interest; in our case, one gRNA was directed to the genome to a cut

site just after the beginning of the ZZ domain (-179 nt from STOP Codon of *orb2-RA/B/C/D/H*),

and the second gRNA was directed to cut +3 nt from STOP Codon. Cas9 mediates breaks in the

double-stranded DNA, which triggers homologous repair between the genomic locus and a donor

plasmid, which was co-injected with the gRNA plasmid.

The donor plasmid containing the mutation of interest (a deletion of the ZZ domain, which

removes the genomic portion of the coding sequence after the end of the second RRM,

advancing the STOP codon to this point, such that this mutant protein would stop at residue

L637 in the PB isoform) followed by a dsRed marker flanked by Piggy Bac 5′ terminal repeats

(Figure S1a). The Piggy Bac transposon terminal repeats enabled this marker to be excised after

it had been used to screen for successful CRISPR incorporation into the genome, leaving behind

no trace of the marker afterwards (i.e. it is "scarless" on the genome). Flanking this dsRed

marker is genomic sequence that is homologous to the base genome, 1kb upstream and

downstream for the cut site; this provided the homology needed to facilitate homology-

dependent repair (HDR) between the donor plasmid and the genomic site.

After injection of the host w^{1118} strain (LWG182), balanced fly lines were established from

the resulting progeny and screened for successful integration of the mutation-containing region

of the donor plasmid. Successful recombination resulted in an ORB2 gene locus harboring a

precise deletion of the ZZ domain at the endogenous locus, marked by the expression of dsRed,

which was later removed from the genome using the Piggy Bac transposon machinery to excise

the marker by crossing them to a Piggy Bac expressing fly (BDSC #8285). PCR was carried out

to confirm donor plasmid insertion at the intended location in the correct orientation (Figure S1b)

followed by deletion of the dsRed marker (Figure S1c), which was then verified by sequencing

(Figure S2). PCR was used to confirm that the $orb2^{\Delta ZZ}$ line was not infected with Wolbachia

(since BDSC #8285 is known to be infected) (data not shown).

Synthetic antibody (Fab) expression and harvesting

Phage displayed synthetic antibodies against ORB2 protein have been described previously

(LAVER et al. 2012; NA et al. 2016). To summarize, antigens containing amino acids 163-446

(numbered according to ORB2-PA isoform) were expressed and purified from *E. coli* as GST

fusion proteins. Synthetic antibody library F (LAVER et al. 2012; NA et al. 2016) was used in five

rounds of binding selection, which included negative selection to remove antibodies that bound

to GST, to obtain synthetic antibodies that bound to the ORB2 antigen. The two synthetic

antibodies used in this study were anti-ORB2 E8 and anti-hEGFR C1 (control).

Synthetic antibodies ('Fabs') were expressed and harvested from *E. coli* as previously

described (LAVER et al. 2012). In summary, DH5α *E. coli* was transformed with plasmid

containing the FLAG- and 6xHis-tagged Fab open reading frame under the control of the *lac*

operator. Transformed cells were used to inoculate bacterial culture using 2xYT media; the

inoculated culture was incubated at 37°C on rotation at 225rpm until OD595 reached 0.8. The

culture was then induced with isopropyl β- d-1-thiogalactopyranoside (IPTG) to drive expression

of the Fab and allowed to grow at 18°C overnight on rotation at 225rpm. Bacteria were pelleted

by centrifugation at 4°C, 4000 rpm for 30 minutes, then the bacterial pellet was resuspended with

B-PER bacterial protein extraction reagent liquid solution (Thermo Scientific) supplemented

with protease inhibitors (150mM NaCl, 1mM AEBSF, 2mM Benzamidine, 2μg/mL Leupeptin,

2μg/mL Pepstatin) and Benzonase nuclease (2500 units/mL). Cells were allowed to lyse for 30

minutes at 4°C, then the lysate was cleared by centrifugation at 4°C at 21,380 RCF for 30

minutes. Supernatant was collected and synthetic antibodies were purified by passing lysates

through a column filled with Protein A affinity gel matrix beads; the captured Fab was eluted

with an acid buffer containing 50mM NaH_2PO_4, 100mM H_3PO_4, and 140mM NaCl, pH to 2.8 in

eight fractions of 200μL each. Fab-containing fractions were pooled and dialyzed with 1L PBS at

4°C for eight hours, twice. Dialyzed Fab was diluted with an equal volume of 95% glycerol

1xPBS and stored at -20°C.

Collection of embryo lysates

Lysate was collected from embryos laid and aged at 25°C. Extract was prepared from

embryos collected 0 to 3 hours post egg-lay on apple juice agar plates, dechorionated with 100%

bleach for 2 minutes, washed with 0.1% Triton X-100, and disrupted by crushing in minimal

volume of lysis buffer consisting of 150 mM KCl, 20 mM HEPES-KOH (pH 7.4), 1 mM $MgCl_2$,

1 mM DTT, and protease inhibitors (1 mM AEBSF, 2 mM benzamidine, 2 μg/mL pepstatin, 2

μg/mL leupeptin). The lysate was cleared by centrifugation at 4°C at 21,380 RCF for 15 minutes,

and the supernatant was collected and either used immediately or stored at -80°C.

RIP-seq

For each replicate of RIP-seq, 1mL of 0-3 hour embryo extract (prepared as described

above) was supplemented with Triton X-100 to 0.1% final concentration, then diluted in an equal

volume of lysis buffer with 0.1% Triton X-100 and cleared again by centrifugation at 4°C, 13

000 RPM for 15 minutes. The supernatant was incubated for 3 to 4 hours at 4°C with end-over-

end rotation with 100μL of anti-FLAG M2 affinity gel beads (Sigma) which had been pre-

incubated at 4°C with 50μg of purified anti-ORB2 E8 Fab blocked with 5μg/μL BSA. After

incubation, the lysates were decanted and beads were washed four times with lysis buffer

supplemented with 0.1% Triton X-100 and then three times with equilibration buffer: 100 mM

NaCl, 100 mM HEPES-NaOH (pH 7.4), 1 mM $MgCl_2$. Fabs and associated ribonucleoprotein

(RNPs) were eluted from the beads with 200 μL of 200μg/μL FLAG-peptide (Sigma) suspended

in equilibration buffer at 4°C for 20 minutes. RNA was isolated from the eluate using TRI

Reagent (Sigma) according to the manufacturer's instructions.

RNA was prepared for sequencing using Ribo-Zero Gold rRNA depletion kit and TruSeq

Stranded Total RNA Library Prep Kit (Illumina) according to manufacturer's instructions.

Libraries were sequenced using the HiSeq 2500 System, V4 Chemistry (Illumina), with paired-

end reads 125nt in length. The raw data are available on GEO (accession number GSE299661).

Computational analysis of RIP-seq data

Paired-end reads obtained from the sequencing were mapped to the *Drosophila*

melanogaster transcriptome r6.23, FB2018_04 obtained from flybase.org (OZTURK-COLAK *et al.*

2024) using the Salmon read mapping program (PATRO *et al.* 2017). The reference transcriptome

was supplied as a FASTA file. The resulting quantification was then analyzed for enrichment in

the ORB2 IP compared to Input and control IP compared to Input using the DESeq2 software

package for R (LOVE *et al.* 2014). Prior to formal analysis, the dataset was filtered for transcripts

having > 1TPM in at least three of the six samples, and rlog transformation was performed to

stabilize the variance. The similarity between samples was calculated by determining the Poisson

Distance between pairs of samples using the *PoiClaClu* package in R (see Figure S3) (WHITTEN

2011). The differential expression analysis was performed with the *dds* function using alpha =

0.01 cut-off for the False Discovery Rate. This returned a fold enrichment for each transcript,

accompanied by an adjusted *P*-value. We defined an ORB2-bound RNA target as satisfying all

three of the following criteria: (1) it was enriched in the ORB2 IP >1.5-fold relative to Input; (2)

it had an ORB2 IP/Control IP ratio >1.5; (3) it had an adjusted *P*-value < 0.05. The results are

presented in Figure 1 and File S1.

For comparisons between our dataset and previously published datasets, each pairwise

comparison was done independently. For comparisons that required a Wilcoxon Rank Sum test,

for each given pair of datasets, a common background list of genes was first compiled by taking

the intersection between the input/background of both datasets; genes which only appeared in

one dataset and not the other were discarded. Then, the measurements of the remaining identified

genes were placed into a dataframe in which each row corresponded to one unique gene, the first

column contained the measurements taken from one dataset, and the second column contained

measurements from the other dataset. This dataframe was then used to perform a Wilcoxon Rank

Sum test using the wilcox.test(x, y, exact = FALSE) function in R, where x is the data in the first

column, y is the data in the second column. The results are shown in Figure 2.

For comparisons that required a Fisher's Exact Test, for each given pair of datasets a

common background list of genes was first compiled by taking the intersection between the

input/background of both datasets; genes that only appeared in one dataset and not the other were

discarded. Then, the remaining identified genes of interest within in each dataset were compared

to look for common elements; genes that were identified in both lists of interest were annotated

as "Overlap", whereas genes that only appeared in one list were annotated as "X Only". A

Fisher's Exact Test was performed by using a 2x2 contingency table, where first cell (top left)

contained the number of genes that did not appear in either list (equal to the total number of

genes in the common background minus the number of genes described in the other three cells),

the second cell (top right) contained number of genes that appeared only on the ORB2 list

(ORB2 Only), the third cell (bottom left) contained the number of genes that appeared only on

the other list of interest (Other Only), and the last cell (bottom right) contained the number of

genes appear on both lists of interest (Overlap). This contingency table was constructed and

evaluated using the fisher.test() function in R. The results are presented in Table 1.

The Gene Ontology (GO) term enrichment analysis was performed using the Functional

Profiling function of the web version of gProfiler2 (accessed at https://biit.cs.ut.ee/gprofiler/gost)

(KOLBERG *et al.* 2023). The list of ORB2-bound target genes was provided to the program in the

Query field, and the Run Query option was selected in order to begin analysis. The results are

presented in Tables 2-4.

Rare-codon enrichment analysis

MELP score (SUPEK AND VLAHOVICEK 2005): Assuming ribosomal proteins are the best

translated and the most codon optimized genes in a given genome, a chi-square based statistic

called Measure Independent of Length and Composition (MILC) was calculated to evaluate the

difference of codon usage frequencies between ribosomal proteins and all annotated CDS that

are longer than 80 codons from Flybase 6.58. Then, MELP (MILC-based Expression Level

Predictor) was calculated based on MILC to predict expression level for each mRNA, and the

isoform with the minimum score (the worst predicted expression) was chosen to represent each

expressed gene (File S2). The Wilcoxon Rank Sum Test was performed for protein coding genes

that were either ORB2 target or non-target.

CAI (PUIGBO *et al.* 2008): Codon Adaptation Index (CAI) is another metric that evaluates

the similarity of codon usage to a reference set. This score was calculated for all annotated CDS

longer than 80 codons from Flybase 6.58 with the local version of CAIcal and the *Drosophila*

melanogaster codon usage table from the Kazusa codon usage database (NAKAMURA *et al.*

2000). The isoform with the minimum score (the most enriched for rare codons) was chosen to

represent each expressed gene (File S2), and the Wilcoxon Rank Sum Test was performed for

protein coding genes that were either ORB2 target or non-target.

Number of Accessible Target Sites (#ATS) *de novo* motif discovery

The ORB2 motifs were identified using the #ATS model, as described (LI *et al.* 2010).

Briefly, the #ATS program is a *de novo* motif discovery algorithm which takes as input nucleic

acid sequences belonging to mRNA transcripts in two lists: a positive (in this case, the 3′UTR of

ORB2 targets) and negative (in this case, the 3′UTR of co-expressed and unbound) list, and

outputs *k*-mers that best differentiate the positive transcripts from the negative transcripts. To

begin the analysis, 6-nt long seed sequences are first assessed based on their enrichment in the

positive list compared to the negative list, and then the top seed sequences are used as the

lauching point upon which the program iterates in an attempt to improve enrichment scores.This

is assessed based on enrichment in the positive list compared to the negative list and accounts for

RNA accessibility, as predicted by computational modeling of RNA secondary structures, and

assigned a *P*-value (see File S3). Iterative enrichment yielded degenerate motifs, so the top five

seed 6-mers from the #ATS were combined into a consensus sequence by aligning the G in the

motif and converting the resulting table into a position weight matrix (PWM) shown in Figure 3.

Evaluation of ORB2 binding motif enrichment in RNA sequences

The enrichment of ORB2 binding motifs was computationally evaluated using a multi-step

pipeline modified from our previous method, which assessed Smaug recognition element (SRE)

enrichment (CHEN *et al.* 2014). Briefly, the computation program scans each RNA using

RNAplfold (ViennaRNA package version 2.3.1) using parameters W=170 -L=120 -T=25

(BERNHART *et al.* 2006; LORENZ *et al.* 2011). Next, the nucleotide sequence is scanned for the

motif of interest (either the ORB2 motif uncovered by our analysis, NUUGUWU, or the ORB2

motif determined by RNA Compete, KUUUKKK) (RAY *et al.* 2013). At candidate motif sites,

the program then assesses the accessibility of the RNA at that position using the results from the

RNAplfold analysis. A score is then assigned to each motif site based on the probability that the

motif sequence is unpaired with any other nucleotide in the sequence. The total score for a

transcript is represented by the sum of motif scores for all motifs found in the sequence.

Cell culture and transient transfection

The Drosophila S2 cell line was maintained at 25°C with ExpressFive SFM (Fisher

Scientific) medium supplemented with 16mM L-glutamine, 100units/mL penicillin, and

100µg/mL streptomycin. 400µL of S2 cells at a density of 1.75×10^6 were plated into one well of

a 24-well tissue culture tray and transfected with a plasmid mixture containing 1.5ng firefly

luciferase plasmid, 1.5ng Renilla luciferase plasmid, 20ng FLAG-effector plasmid (containing

the open reading frame for either ORB2 whole protein, fragments of ORB2 protein, Protein A, or

GFP), 4ng of FLAG-GST plasmid, and 178.5ng pSP72 plasmid using 0.4µL of TransIT-Insect

transection reagent (Mirius Bio) suspended in 20µl of tissue culture medium according to the

manufacturer's instructions. For tethering assays, the luciferase plasmid was modified such that

the luciferase 3′UTR contained six copies of the TAR element and the effector protein construct

was modified to include the BIV-Tat ORF N-terminal to the FLAG tag. For experiments which

measure ORB2 binding to ORE elements, the luciferase plasmid was modified such that the

luciferase 3′UTR contained six copies of either intact ORE elements (UUUUGUU) or mutant

ORE elements (UUAUCUC). Both luciferase reporters were under the control of the metal-

inducible metallothionein promoter; all FLAG-tagged effector constructs were derived from

pAc5.1/V5-His plasmid (Thermo Fisher Scientific), which carries the Actin5C promoter.

Luciferase expression was induced 24 hours post-transfection by the addition of copper sulphate

to final concentration of 0.5mM. 24 hours post-induction and at 48 hours post-transfection cells

were harvested for downstream assays.

Luciferase Assay

Cell lysate was prepared from S2 cell transient transfections by dislodging cells from the

surfaces of the cell culture container and transferred into conical tubes. Cells were pelleted by

centrifugation at 21,380 RCF at 4°C for 1 minute; the supernatant was removed, and the cell

pellet was resuspended in by adding 20 µL of lysis buffer (containing 1xPassive Lysis Buffer

[Promega], 10 mg/mL BSA, and 1 mM AEBSF). Lysis was facilitated by pipetting up and down

and allowing the cells to lyse at 4°C for 10 minutes. Firefly and Renilla luciferase activities were

measured using the Dual-Luciferase Reporter Assay system (Promega) according to

manufacturer's instructions using a Berthold Detection Systems SIRIUS Luminometer.

Normalized luminescence values were calculated by first dividing the Firefly luminescence by

the Renilla luminesence readings to obtain a Firefly/Renilla ratio; ratios for each biological

replicate were obtained from the average of two technical replicates. Then, this ratio was

normalized to the corresponding GFP control sample (transfected in parallel to the experimental

conditions). Normalized expression was presented as the average of at least three biological

replicates with error bars representing standard deviation.

Reverse Transcription – Quantitative Real-time Polymerase Chain Reaction (RT-qPCR)

RNA isolated from ORB2 co-immunoprecipitations was reverse transcribed into single-

stranded cDNA with the Superscript IV reverse transcriptase kit (Invitrogen) according to

manufacturer's instructions. 500ng of total RNA per sample were used to synthesize cDNA,

which was primed using random hexamer primers. The resulting single-stranded cDNA was

diluted 1:25 using RNase-free water and used to perform quantitative real-time PCR (qPCR)

with primers specific to the various transcripts assayed. Primers were designed using NCBI

Primer-BLAST to cover all isoforms produced by a gene and were required to span an exon-

exon junction. qPCR was performed with SensiFAST SYBR PCR mix (Bioline) following the

manufacturer's protocol and using 5 µL of diluted cDNA per reaction. The CFX384 Real-Time

System (Bio-Rad) was used to carry out the PCR reaction.

The results of the qPCR were analyzed using CFX Manager software (Bio-Rad). Each

biological replicate represented a measurement obtained for a separate embryo collection; values

from three technical replicates were averaged and relative gene expression was normalized to the

ribosomal protein-coding transcript *RpL32*. Normalized expression was presented as the average

of three biological replicates with error bars representing standard deviation.

Western Blots

The following primary antibodies were used: mouse anti-ORB2 4GS (1:50) obtained from the Developmental Studies Hybridoma Bank (https://dshb.biology.uiowa.edu/orb2-4G8) and described in (HAFER *et al.* 2011), mouse M2 anti-FLAG (Sigma, 1:5,000), and mouse anti-Actin (Sigma Cat#A4700, 1:10,000) which was used as a loading control.

Dechorionated embryos were counted, then lysed in Laemmli buffer at a concentration of 1 embryo/μL and boiled for 2 minutes. Proteins were resolved by loading 5μL of lysed embryos in Laemmli buffer per lane into an SDS-PAGE and transferred to PVDF membrane in the absence of SDS. The blot was blocked at room temperature with 2% non-fat milk in PBST for 30 minutes and incubated with primary antibodies diluted in 2% non-fat milk in PBST at 4°C overnight. After this incubation, the blot was washed 3 x 10 minutes with PBST at room temperature and incubated with the appropriate HRP-conjugated secondary antibodies (1:5000, Jackson ImmunoResearch) in 2% non-fat milk in PBST at room temperature for 1 hour. The blot was washed 3 x 10 minutes with PBST and developed using Immobilon Luminata Crescendo Western HRP substrate (Millipore), imaged using a BioRad Chemidoc MP Universal Hood III imager and band intensities were quantified using ImageLab.

Structure Prediction with AlphaFold

For structural predictions the full amino acid sequence of the ZZ domain constructs used in the tethering assays (see above) was predicted using AlphaFold version 3 (JUMPER *et al.* 2021; ABRAMSON *et al.* 2024). Predictions were visualized and truncated to show regions A-to-D in ChimeraX version 1.9 (MENG *et al.* 2023). Structures were color-coded to depict their predicted local distance difference test (pLDDT) score using the AlphaFold Error Plot tool. The results are presented in Figure S8.

Mass spectrometry

Immunoprecipitation of ORB2 from embryo lysates with synthetic antibodies

For IP-MS, 500 μL of 0-3 hour embryo extract (prepared as described above) was supplemented with Triton X-100 to 0.1% final concentration, then diluted in an equal volume of lysis buffer with 0.1% Triton X-100 and cleared again by centrifugation at 4°C at 21,380 RCF

for 15 minutes. The supernatant was incubated for 3 to 4 hours at 4°C with end-over-end rotation

with 100µL of anti-FLAG M2 affinity gel beads (Sigma) which had been pre-incubated at 4°C

with 50µg of purified anti-ORB2 E8 Fab blocked with 5µg/µL BSA. After incubation, the lysates

were decanted and beads were washed twice with lysis buffer supplemented with 0.1% Triton X-

100, then another twice with lysis buffer without Triton X-100; beads were transferred into a

new tube and washed another twice with lysis buffer without Triton X-100. Fabs and associated

proteins were eluted by tryptic digest: Beads were resuspended in 200 µL of 50mM ammonium

bicarbonate pH 8 supplemented with 2 µg of MS-grade trypsin and incubated at room

temperature overnight with end-over-end rotation. The digested peptides were collected and a

second elution with 200 µL of 50mM ammonium bicarbonate was performed at room

temperature for 30 minutes. The two fractions were pooled, flash frozen with dry ice, dehydrated

in a rotary evaporator for 8 hours, then stored at -20°C.

Liquid chromatography and mass spectrometry

For liquid chromatography (LC) the following solvents were used: Solvent A (0.1%

(vol/vol) formic acid) and Solvent B (0.1% (vol/vol) formic acid/90% (vol/vol) acetonitrile). The

LC parameters were set using an Easy-nLC 1200 instrument. Peptides were loaded and separated

on a nanoViper trap column (75µm x 2 cm), Acclaim PepMap 100 (C18, 3µm, 100 Å, Thermo

Scientific, cat. no. 164946), and EASY-Spray analytical column (75µm x 50cm) Acclaim

PepMap RSLC (C18, 2µm, 100Å, Thermo Scientific) using a flow rate of 225nL/minute.

Tandem MS was performed using the Q Exactive HF-Orbitrap mass spectrometer (Thermo

Scientific), as previously described (LIU *et al.* 2014; JIANG *et al.* 2015; CHIU *et al.* 2016). The

parameters for acquisition were 1 MS scan (50 ms; mass range, 390 to 1800) at a resolution of

60,000K, followed by up to 20 MS/MS scans (50 ms each) at a resolution of 15,000 and an AGC

target of $1x10^5$. Candidate ions with charge states +2 to +7 were isolated using a window of 1.4

amu with a 5s dynamic exclusion window.

Data processing and analysis

Four biological replicates were performed, both in the presence or absence of RNase A.

The ProHits software package was used to perform Significance Analysis of INTeractome

(SAINT) in order to identify ORB2-interacting proteins (LIU *et al.* 2010; CHOI *et al.* 2012; LIU *et*

al. 2012). Proteins with associated peptide counts were filtered through the Trans Proteomic

Pipeline (TPP) for iProphet probability > 0.95 and number of unique peptides ≥ 2. SAINT

express analysis was performed on the ProHits interface with only Drosophila detected proteins

(selected options: number of compressed controls = 3, burn-in period, nburn = 2000; iterations,

niter = 5000, lowMode = 1; minFold = 1; normalize = 1; nCompressBaits = 3). This analysis

determined the probability of interaction for each identified prey protein. Proteins with AvgP

score ('SAINT score') ≥ 0.95 and BFDR ≤ 0.01 were defined as ORB2-ineracting proteins. Raw

data are available at MassIVE MSV000098275. The results are presented in File S4.

To perform network analysis on the list of identified protein interactors, we used the

GeneMANIA application (v3.5.2) (WARDE-FARLEY *et al.* 2010; FRANZ *et al.* 2018) with the

Cytoscape network visualization program (v3.9.1) (SHANNON *et al.* 2003; MONTOJO *et al.* 2010).

Data resulting from the IP-MS was imported to Cytoscape in table form. GeneMANIA maintains

a database of functional association data, which includes protein interactions, genetic

interactions, pathways, co-expression, and co-localization data that is obtained from a wide

variety of published sources. The details regarding the GeneMANIA database can be found

online (https://genemania.org/data/). Two interaction networks were produced by performing a

Network Search on a subset of identified protein interactors using GeneMANIA's *Drosophila*

melanogaster functional association database (see

https://genemania.org/data/current/Drosophila_melanogaster/ for details). Then, the nodes of

each network were annotated according to the association type reported in the GeneMANIA

database (i.e. known physical interaction, known genetic interaction, or predicted interaction).

The results are presented in Figure 5c,d.

Embryo sample collection for *orb*$^{\Delta ZZ}$ time-course and polysome gradient RNA-seq

experiments

Embryos were collected from *orb2*$^{\Delta ZZ}$ and *w*^{1118} (pre-injected control, LWG182) strains at

1-hour intervals post-egglay (0-1 hour, 1-2 hour, 2-3 hour), dechorionated with ice cold 100%

bleach for 2 minutes, washed with 0.1% Triton X-100, and disrupted by crushing using 0.5 g

embryos per mL of either lysis buffer (150 mM KCl, 20 mM HEPES-KOH (pH 7.4), 1 mM

MgCl2, 1 mM DTT, 1 mM AEBSF, 2 mM benzamidine, 2 µg/mL pepstatin, 2 µg/mL leupeptin)

or polysome lysis buffer (7.5 mM MgCl2, 500 mM NaCl, 25 mM Tris (pH 7.5), 1 mM DTT, 1

mM AEBSF, 2 μg/mL leupeptin, 2 mM benzamidine, 2 μg/mL pepstatin A, 2 mg/mL heparin,

0.5 mg/mL cycloheximide). The lysate with lysis buffer was cleared by centrifugation at 4°C at

21,380 RCF for 15 minutes, TRIzol (Invitrogen) was added to the supernatant which used

immediately for RNA extraction or stored at -80°C. The lysate with polysome lysis buffer was

either used immediately for polysome gradient processing or stored at -80°C.

Polysome gradients

The polysome gradient protocol was adapted from our published method with

modifications (CHEN *et al.* 2014). Lysed samples were diluted 1 in 10 in polysome lysis buffer

and 20% Triton X-100 was added to a final concentration of 1%, centrifuged at 6,000xg for 10

minutes and the resulting supernatant was diluted in polysome lysis buffer supplemented with

1% Triton X-100 to an A_{260} of 12.5. A 5 ml 15% to 45% linear sucrose gradient in 7.5 mM

$MgCl_2$, 500 mM NaCl, 50 mM Tris pH 7.5 was produced using a BioComp Model 117 Gradient

Mate gradient maker (Biocomp, Fredericton, New Brunswick, Canada) using a rotation angle of

86 degrees and a rotation speed of 25 for 51 seconds. After chilling the polysome gradient at 4ºC

for 1 hour, 200 μL of diluted embryo extract was loaded onto the top of the gradient, which was

then spun at 36,000 rpm in a Beckman SW50.1 rotor for 2.5 hours. 400 μL fractions (hand

fractionated) were collected, and A_{260} absorbance was measured using a Nanodrop (Thermo

Fisher Scientific). We added 20% SDS, 0.5 M EDTA and 20 mg/ml proteinase K to final

concentrations of 0.8%, 0.01 M and 0.128 mg/ml, respectively, and then incubated the fractions

for 30 minutes at 37ºC. 2.5 μL GlycoBlue (15 mg/mL, ThermoFisher) following which 2.5x

volume ethanol was added to each fraction and precipitated overnight. The resulting pellet was

washed with 75% ethanol and resuspended in phenol-saturated water.

The fractions were then separated into three pools: Pool 1 contained the (top) fractions 1-3,

pool 2 contained fractions 4-6, pool 3 contained fractions 7-13. Following two phenol-

chloroform extractions, the aqueous phase of each pool was collected with Invitrogen

Phasemaker Tubes (Fisher A33248) following the manufacturer's protocol. The samples were

precipitated by the addition of 7.5 M LiCl to a final concentration of 3.75 M followed by an

overnight incubation at -20°C. The pellets were washed with 70% ethanol and resuspended in

water.

RNA purification was performed using 2.2× volume NEBNext RNA Sample Purification

Beads (NEBNext Ultra II DNA Library Prep with Sample Purification Beads, E7767L)

following the manufacturer's protocol. RNA concentration was assessed with the Qubit RNA HS

Assay Kit (Thermo Fisher) and Agilent bioanalyzer at The Centre for Applied Genomics

(TCAG), The Hospital for Sick Children, Toronto, ON, Canada.

rRNA depletion, library preparation and RNA-seq for time-course and polysome gradient

RNA-seq

For the RNA-seq time-course rRNA depletion was performed as previously described

(HAUGEN *et al.* 2024). For polysome gradient RNA-seq, rRNA depletion was performed using

the QIAseq FastSelect rRNA Fly Kit (QIAGEN, Cat# 33326). Library preparation for RNA-seq

was conducted using the NEBNext Ultra II Directional RNA Library Prep Kit for Illumina, and

sequencing was performed on an Illumina NovaSeq 6000 (PE150) at the Next Generation

Sequencing Facility, The Centre for Applied Genomics (TCAG), The Hospital for Sick Children,

Toronto, ON, Canada.

Computational analysis of time-course and polysome gradient RNA-seq data

RNA-seq data processing pipeline

Adapter trimming and quality control used Trim Galore (Version 0.6.5)

[https://github.com/FelixKrueger/TrimGalore]. rRNA and 'unwanted' RNA removal used

SortMeRNA (version 2.1) (KOPYLOVA *et al.* 2012), filtering against *Drosophila melanogaster*

7SLRNA, rRNA, pseudogene-rRNA, tRNA, mtRNA and other species' rRNA databases.

Transcriptome alignment was carried out with Salmon (Version 1.9.1) (PATRO *et al.* 2017) in

mapping-based mode using FlyBase release 6.23 transcriptome RNA (FB2018_04) (OZTURK-

COLAK *et al.* 2024). Full decoy indexing was applied following the Salmon website instructions

[https://salmon.readthedocs.io/en/latest/salmon.html].

Differential gene expression analysis

Differential expression analysis used the R package tximeta (version 1.20.3) (LOVE *et al.*

2020) and Deseq2 (version 1.42.1) (LOVE *et al.* 2014) in R (version 4.3.2). For the *orb2*$^{\Delta ZZ}$ and

wild type time-course total RNA-seq, reads from each time point were normalized using the

DESeq2 package. For the *orb2^{ΔZZ}* and wild type polysome gradient RNA-seq time-courses, reads

of each time point of *orb2^{ΔZZ}* and wild type pool 1 and pool 3 RNA were normalized using the

DESeq2 package. For each gene, the mean of normalized counts per pool was computed at each

time point and genotype. The 'Translation Index' of each mRNA was calculated as the ratio =

mean normalized counts of pool 3 /mean normalized counts of pool 1. Normalized counts were

used for statistical analysis by the Wilcoxon Rank Sum Test and data were visualized with

ggplot2 in R. Raw data are available at GEO for the RNA-seq timecourse and polysome gradient

timecourse with accession numbers GSE299140 and GSE299019, respectively. The results are

presented in Figure 6 and Files S6 and S7.

Comparisons between the ORB2 target list and published SMG, PUM, BRAT and RIN

target lists:

The total list of genes detected in this ORB2 RIP-Seq experiment was first compared with

the total list of genes present in the published RIP-Chip experiments for SMG, PUM and BRAT

(CHEN *et al.* 2014; LAVER *et al.* 2015; LAVER *et al.* 2020a). The overlap between these lists was

retained, and the non-overlapping genes (i.e. genes detected in only one of the two experiments)

were discarded. From the remaining genes, ORB2 targets were compared with the targets of

SMG, PUM, and BRAT. The translation level, as defined by the polysome RIP-Seq ratio

measured in 0-1h embryos (see above), was compared for the overlap between the ORB target

list and the other list of interest (i.e. ORB2 + SMG co-targets, ORB2 + PUM co-targets, and

ORB2 + BRAT co-targets), and separately for the ORB2-only targets, the SMG- PUM-, and

BRAT-only targets, and non targets.

RESULTS

ORB2 binds to over 450 targets in the early Drosophila embryo

To identify transcripts that are bound by ORB2B (hereafter referred to as ORB2), we used

our previously produced Fab against an antigen N-terminal to the RRMs (Figure 1a) (LAVER *et*

al. 2012; NA *et al.* 2016) to perform RNA-coimmunoprecipitation (RIP) of endogenously

expressed ORB2 from embryo extracts prepared from 0-3 hour old embryos produced by *w^{1118}*

females. We enriched for mRNA in the sample by rRNA depletion, generated randomly primed

cDNA libraries from the resulting samples, and identified co-purifying RNAs by next-generation

RNA sequencing (see Materials and Methods). As a negative control, we performed parallel

RIPs using a control synthetic antibody, C1, as described previously (LAVER *et al.* 2015; LAVER

et al. 2020b; HAUGEN *et al.* 2024). Input samples from the same extracts were also sequenced.

We performed three biological replicates for each of these three conditions, for a total of nine

cDNA libraries.

To identify ORB2-bound target genes, we mapped the sequence reads to the Drosophila

transcriptome using Salmon read alignment software, and analyzed the results with the DESeq2

software package as described in Materials and Methods (LOVE *et al.* 2014; PATRO *et al.* 2017).

We defined an ORB2 target as a gene whose transcripts had fold enrichment >1.5 in the anti-

ORB2 RIP-seq compared to input and a ratio of ORB2/Control > 1.5, with an FDR adjusted *P*-

value < 0.05; using these criteria, we identified 462 genes whose transcripts were bound by

ORB2 (Figure 1b and File S1). ORB2's target transcripts were significantly less abundant than

co-expressed non-target transcripts (Figure 1c). RIP-RT-qPCR verified several of the target and

non-target RNAs (Figure 1d).

While ours is the first identification of ORB2 targets in embryos, there are previously

published target lists from other tissues or cells. For example, ORB2-bound transcripts were

identified in a CLIP experiment derived from ORB2 overexpression in S2 cells (STEPIEN *et al.*

2016). Of the 462 ORB2 targets on our list, 298 were identified in their CLIP-seq background set

of which they identified 113 as targets (Fisher's exact $P = 6.52 \times 10^{-7}$, Odds Ratio = 1.88). We

also examined the overlap of our targets with a list of ORB2-associated transcripts obtained by

performing RIP-seq on Drosophila head extracts (KOZLOV *et al.* 2023). 336 genes in our target

gene list overlapped with theirs (Fisher's exact $P < 10^{-15}$, Odds Ratio = 5.00).

The strong overlap between our ORB2 targets and previously reported targets from other

cell types/tissues verifies that our list is enriched for true ORB2 interactors.

ORB2 targets are translationally repressed, deadenylated and degraded during the MZT

To assess the post-transcriptional fate of ORB2 target transcripts in the early Drosophila

embryo, we utilized three previously published datasets that scored each transcript's translational

status, poly(A) tail length, and abundance during the MZT.

Two studies measured Translation Efficiency (TE), calculated from ribosome profiling,

during the MZT (DUNN *et al.* 2013; EICHHORN *et al.* 2016). At all three timepoints during the

MZT that were studied, ORB2 targets had a significantly lower TE compared to co-expressed,

non-target transcripts (Figure 2a-c). A third study measured Translation Index (TI), calculated as

the ratio of that transcript's abundance in polysome fractions of a sucrose gradient compared to

its abundance in non-polysomal fractions (CHEN *et al.* 2014). ORB2 targets had a significantly

lower TI than co-expressed non-target transcripts (Figure 2d).

We next assessed changes in poly(A) tail length and abundance for ORB2 targets between

the 0-1 hour timepoint and the 2-3 hour timepoint, which had been measured in parallel to TE

(EICHHORN *et al.* 2016). Decrease in abundance and poly(A) tail length were significantly greater

for ORB2 targets than for co-expressed, non-target transcripts (Figure 2e,f). We also examined

previously published lists of transcripts that are degraded during the MZT (Table 1). ORB2

targets overlapped significantly with transcripts identified to be unstable in activated eggs

(TADROS *et al.* 2007) (Fisher's Exact $P = 2.2 \times 10^{-16}$, odds ratio 3.85). In addition, ORB2 targets

were significantly enriched for genes that are degraded during the MZT (THOMSEN *et al.* 2010):

(Class III and V; respectively, Fisher's Exact $P = 0.02$, odds ratio 1.54 and Fisher's Exact $P =$

2.2×10^{-16}, odds ratio 2.54) and significantly depleted for transcripts that are stable (Class I;

Fisher's Exact $P = 2.2 \times 10^{-16}$, odds ratio 0.23).

We conclude that ORB2's target mRNAs are enriched for transcripts that are

translationally repressed, deadenylated and degraded during the MZT.

ORB2 targets are enriched for rare codons

It has previously been reported that ORB2 binds to and upregulates the stability and

translation of rare-codon-enriched mRNAs in neurons in *Drosophila* (STEWART *et al.* 2024). We,

therefore, asked whether ORB2's embryonic targets are enriched for rare codons. Two different

algorithms – MELP (SUPEK AND VLAHOVICEK 2005) and CAI (PUIGBO *et al.* 2008) – found

significant enrichment of rare codons in ORB2 targets relative to non-target, co expressed

mRNAs (Figure 2g,h and File S2). Why the post-transcriptional outcome in embryos is the

opposite of that in neurons will be considered in the Discussion.

ORB2 targets are enriched for roles in gamete generation, gene regulation, and development

We next asked whether ORB2's targets were enriched for particular molecular, cellular or biological functions by performing Gene Ontology (GO) term analysis using the gProfiler2 tool (GENE ONTOLOGY *et al.* 2023; KOLBERG *et al.* 2023) (Tables 2-4).

The top GO terms under the Molecular Function category (Table 2) were dominated by molecular binding functions; 'mRNA 3′UTR binding', and 'mRNA regulatory element binding' were among the top ten results. Genes belonging to these terms included *orb2* itself as well as *patronin, piwi,* and *bru1*. Other top terms included 'translation regulator activity' and 'translation repressor activity'; genes belonging to these terms included *orb2* itself, *orb, pum, fmr1, sxl, heph,* and *eIF4E-HP*. In the Biological Process category (Table 3), ORB2 targets were enriched for terms related to regulation and development, such as 'anatomical structure development', 'cell differentiation', and 'regulation of metabolic process'. ORB2 targets were also enriched for terms related to gametogenesis, including 'gamete generation', 'oogenesis', and 'germ cell development'. Genes related to these terms included *pum, rbfox1, piwi, heph, bru1, sxl,* and *mei-P26*. Enrichment for GO terms related to gametogenesis in ORB2 targets is consistent with previously reported functions for ORB2 in, for example, regulation of spermatogenesis (XU *et al.* 2012; GILMUTDINOV *et al.* 2021). The Cellular Component category (Table 4) was dominated by terms that included 'ribonucleoprotein granule', 'P granule', 'pole plasm', and 'germ plasm'. Genes related to these terms included *larp, tud, osk, tapas, ago1,* and *dcr-2*.

Many of the genes listed under enriched GO terms are previously identified ORB2 targets (e.g., *orb2* itself, *ago1*, *bru1*, *dcr-2*, *eIF4E-HP*, *fmr1*, *heph*, *orb, pum, rbfox1, unr*) (MASTUSHITA-SAKAI *et al.* 2010; XU *et al.* 2012; STEPIEN *et al.* 2016; KOZLOV *et al.* 2021) while others are previously unidentified players in the same enriched function or process (e.g., *imp* and *piwi*). Enrichment for mRNAs encoding known post-transcriptional regulators suggests that ORB2 may play a key role in the cross-regulatory network that controls the MZT, a function that is supported by our subsequent analyses reported below. The fact that ORB2 binds to and potentially regulates mRNAs that encode other post-transcriptional regulators is also consistent with a previous study in S2 cells showing that RNP complexes tend to be enriched for mRNAs encoding post-transcriptional regulators (STOIBER *et al.* 2015).

ORB2 target transcripts are enriched for a U-rich motif

CPEB-family proteins are known to bind to U-rich elements, typically located in the 3′UTR of their mRNA targets (HAKE AND RICHTER 1994; DE MOOR AND RICHTER 1999; STEPIEN *et al.* 2016). To predict *cis*-elements recognized by ORB2 from our *in vivo* RIP-seq results, we used our algorithm, Number of Accessible Target Sites (#ATS), which performs *de novo* motif discovery given a target set and a co-expressed non-target set of transcripts (LI *et al.* 2010) as described in the Material and Methods. Using our input RNA-seq data, we identified the most abundant transcript isoform from each expressed gene in the early embryo and then ran #ATS analysis on the full-length transcripts or the 3′UTR sequences of ORB2 targets against non-targets. Whereas no motif was produced from the analysis of full-length transcripts, the #ATS program identified a list of 6-mers enriched in the 3′UTR of ORB2 target sequences relative to non-targets; these 6-mers were then combined into a consensus motif (Figure 3a, File S3), which will henceforth be referred to as an ORB2-recognition element (ORE: NUUGUWU (where W = A or U). We also calculated an ORE 'score' for each transcript by summing the accessibilities of each predicted ORE in that transcript (see Materials and Methods). ORB2 targets had significantly higher ORE scores than co-expressed non-targets (Figure 3b, File S3).

This ORE is similar to published binding motifs: UUUU(A)$_{1-3}$U and UUUUGU identified in S2 cells (STEPIEN *et al.* 2016); UUUURU (where R = A or G) in brain extracts (MASTUSHITA-SAKAI *et al.* 2010); and KUUUKKK (where K = G or U) identified by an *in vitro* binding assay (RAY *et al.* 2013). We calculated 'KUUUKKK' scores for each transcript and found that the ORB2 targets we identified in the embryo have significantly higher scores than co-expressed non-targets (Figure S3b, File S3).

The *de novo* discovery of the U-rich ORE enriched in ORB2 target RNAs in the early embryo and its similarity to ORB2 motifs previously identified using a variety of methods supports the conclusion that our list of targets is enriched for transcripts directly bound by ORB2.

The ORE directs ORB2 binding to and repression of a luciferase reporter mRNA in S2 cells

To validate the biological activity of the ORE identified by #ATS, we used a luciferase reporter assay in Drosophila S2 cells to measure ORB2 binding and ORB2-mediated regulation.

In the assay, firefly luciferase (FLuc) reporters harboured six copies in their 3′UTR of either a

representative ORE (UUUUGUU) or a mutated ORE (UUAUCUC) (the reporter is schematized

in Figure 3c). Each of these reporters was individually expressed in S2 cells together with either

a control protein (GFP), full-length ORB2 protein, or an ORB2 protein lacking its RBD

(ORB2ΔRRM). An unregulated Renilla luciferase (RLuc) reporter was co-transfected with all

samples, allowing us to normalize FLuc mRNA and protein levels.

ORB2 RIP followed by RT-qPCR confirmed that the reporters carrying the wild-type

OREs, but not the mutant OREs, were enriched in ORB2 IPs, compared to IPs of the

ORB2ΔRRM protein (Figure 3d). Furthermore, significant ORB2-mediated repression of the

reporter was seen when ORB2 protein was expressed together with a reporter construct carrying

wild-type OREs (Figure 3e). When either the OREs were mutated or the RRMs were deleted, no

repression was observed (Figure 3e) despite the fact that ORB2ΔRRM protein was expressed at

higher levels than full-length ORB2 (Figure S4).

We also assessed whether the firefly luciferase open reading frame used in the reporters

might be enriched for *Drosophila melanogaster* rare codons and found that this is indeed the

case (Figure S5). Our data do not address whether rare codon-enrichment per se contributes to

binding and/or repression; this would require the use of a codon-optimized version of luciferase,

which is beyond the scope of this study.

Our data do, however, clearly show that ORB2 binds to OREs in an RBD-dependent

manner and that this binding is sufficient to confer repression on the reporter RNA. The

magnitude of repression we observed is similar to that previously reported for ORB2B in similar

reporter assays in S2 cells (STEPIEN *et al.* 2016).

The C-terminal Zinc-binding ('ZZ') domain of ORB2/hCPEB2 but not ORB/hCPEB1

represses a luciferase reporter mRNA in S2 cells

To identify domains of ORB2 necessary and sufficient for repression we used a luciferase

reporter tethering assay in S2 cells. We transfected a plasmid construct that would express a

protein containing the BIV-Tat RNA-binding domain fused to all or part of the ORB2 protein.

This fusion protein is recruited to six tandem repeats of the trans-activating response (TAR)

element (to which BIV-Tat binds) located in the 3′UTR of a co-transfected firefly luciferase

reporter (schematised in Figure 4a) (WAKIYAMA *et al.* 2012; LAVER *et al.* 2020a). Tethering of

full-length ORB2B protein produced a significant, two-fold repression relative to a tethered GFP

control (Figure 4b); the extent of repression was consistent with previously published data for

ORB2 in S2 cells (STEPIEN *et al.* 2016). Tethering of ORB2 protein did not affect steady-state

reporter RNA levels (Figure 4c).

To map regions of ORB2 that mediate repression, we fused the BIV-Tat domain to

fragments of the ORB2 protein and assessed whether any individual region was capable of

repressing luciferase expression. We divided the ORB2 protein into five regions: IDR1, the

polyQ region, IDR2, the RBD, and the ZZ domain (Figures 1a and 4b,c). There was no

significant effect on the expression of luciferase after tethering of IDR1, polyQ, IDR1+polyQ,

IDR2 or RBD (Figure 4b and see Figure S6A for western blot of expression of each fragment).

Only the ZZ domain conferred strong (~three-fold) repression (Figure 4b) despite being

expressed at a comparatively low level (Figure S6a). To determine whether the ZZ domain was

also necessary for repression, we tethered an ORB2 protein construct lacking the ZZ domain

(ORB2ΔZZ) to the luciferase reporter and found that it was unable to repress luciferase

expression, despite ORB2ΔZZ retaining the RBD and being expressed at higher levels than full-

length ORB2 (Figure 4b, Figure S6b). Because luciferase mRNA levels were not significantly

different from controls in either the ZZ sufficiency or the ZZ necessity experiments (Figure 4c,

ZZ and ORB2ΔZZ), ZZ domain mediated repression can be attributed largely to effects on

translation rather than RNA stability. We conclude that the ZZ domain of ORB2 is both

necessary and sufficient to confer translational repression when tethered to a reporter mRNA.

Finally, to assess possible conservation of function, we also tethered the ZZ domains of

ORB, hCPEB1 and hCPEB2. The hCPEB2 ZZ domain conferred significant repression, whereas

the ORB ZZ and hCPEB1 ZZ domain were unable to repress (Figure 4e; western blot is shown

in Figure S6c). We conclude that the repressive function of the ZZ domain is conserved between

ORB2 and its human ortholog, hCPEB2, while it is not conserved in the more distantly related

proteins, ORB or hCPEB1. We note that the ZZ domains of hCPEB3 and hCPEB4 are ≥ 90%

identical to the hCPEB2 ZZ domain (Figure S7a). This suggests that, although not tested here,

the hCPEB3 and hCPEB4 ZZ domains likely also function in translational repression.

Regions within the ORB2 ZZ domain function in combination to confer repression

To map regions of the ORB2 ZZ domain sufficient for repression, we first compared the

sequences of the ZZ domains of ORB, ORB2, hCPEB1 and hCPEB2 (Figures 4d and S7a).

There is high level of sequence identity between the ORB2 ZZ domain and that of hCPEB2

(81.3% amino acid identity). In contrast, the ZZ domain of ORB2 and ORB, exhibit only 35.6%

sequence identity and hCPEB1 has only 28.5% amino acid identity to ORB2. We then

considered the predicted structures of these domains (Figure S8). An NMR structure of the

hCPEB1 ZZ domain showed it consists of a rubredoxin (Rd) turn, followed by a beta hairpin,

and a short alpha helix (MERKEL *et al.* 2013). Predictions of the structures of the ORB, ORB2

and hCPEB2 ZZ domains using Alphafold 3 (ABRAMSON *et al.* 2024) showed a similar

arrangement. Indeed, Alphafold3 prediction of the ORB ZZ structure was virtually identical to

that of the NMR structure of the hCPEB1 ZZ domain (Figure S8a). Similarly, the predictions of

the ORB2 and hCPEB2 ZZ structures were very similar to one another (Figure S8a). Finally, an

overlap of the ORB and ORB2 ZZ domain structures showed that the only difference was the

length of the Rd turn (Figure S8b).

To map regions within the ORB2 ZZ domain sufficient for repression we assessed whether

or not chimeric versions that included one or more of the Rd turn ('A', Q640 to P656), the beta

hairpin ('B', F657-Y668), the alpha helix ('C', C669 to Y684) and/or the C-terminal region ('D',

H685 to C-terminus) regions of the ORB or ORB2 ZZ domain functioned to repress firefly

luciferase expression using the tethering assay.

As expected, the full-length ORB2 ZZ domain repressed whereas the ORB ZZ domain did

not. Of the 14 chimeras tested, three resulted in significant repression relative to the GFP control

(Figure 4f, western blots are shown in Figure S7b,c): ORB2 ZZ-$A^1B^2C^2D^2$, ORB ZZ-$A^2B^1C^2D^1$

and ORB ZZ-$A^1B^1C^2D^2$ (where the superscript '2' refers to ORB2-derived sequences and

superscript '1' refers to ORB-derived sequences). The region shared by all of these mosaics is

ORB2 region C (denoted C^2), suggesting that it is a major contributor to the repressive function

of the ORB2 ZZ domain. However, ORB2 region C alone is not sufficient to confer repression

but needs to be combined with another ORB2 region (either A^2, B^2 or D^2) for repression.

ORB2 interacts with translational repressors and the 43S translation preinitiation complex

in early embryos

We next turned to proteomic strategies to identify protein partners of ORB2 in early

embryos. We immunoprecipitated endogenous ORB2 from extracts prepared from embryos

produced by w^{1118} females, both in the presence and absence of RNase A, in biological

quadruplicates, using the same synthetic antibody used above to identify ORB2 target RNAs.

Co-purifying proteins were identified by mass spectrometry followed by Significance Analysis

of INTeractome (SAINT), which assigns confidence scores to protein-protein interactions based

on comparisons between the Gaussian curves produced by the control IP data and the bait IP data

(CHOI et al. 2012). SAINT was performed with stringent parameters, where for each protein, the

lowest control and the highest bait were dropped prior to statistical analysis, which helped ensure

that only high-confidence interactors would be identified. We defined a protein interactor of

ORB2 as any protein that was significantly enriched in ORB2 immunoprecipitates compared to

parallel purifications using our negative control antibody, C1, with a SAINT (AvgP) score $\geq$

0.95 and a BFDR < 0.01 (File S4).

We identified 106 interactors in the presence of RNase A (Figures 5a and S10a),

representing RNA-independent interactions, and 212 interactors of ORB2 in the absence of

RNase A (Figure S10c,d); 85 were in common (File S4). The interactors largely fell into two

distinct groups: The first consisted of RBPs known to negatively regulate target stability and/or

translation, including AGO1, AGO2, PUM, and RBFOX1, as well as the Cup repressive

complex (Cup, ME31B, TRAL, BEL) (GOTZE et al. 2017) (Figure 5a,c File S4). All components

of the Cup repressive complex copurified in the presence of RNase (Figure 5a,c), consistent with

an RNA-independent interaction with ORB2. The second group consisted of translation-

initiation factors, notably the components of the 43S preinitiation complex (PIC): all eIF3 and

eIF2 subunits, eIF1 and -1A, and eIF5A; most (13/20) were RNA-independent interactors,

including 11/14 detected eIF3 subunits (Figure 5a,d and File S4).

We note that several of ORB2's protein interactors were also identified as mRNA targets

in our RIP-seq experiments (File S5). This result is consistent with an earlier study of RBP

mRNA targets and protein interactors carried out in S2 cells, which showed that RNP complexes

are enriched for both mRNAs and their encoded proteins involved in RNA binding and post-

transcriptional regulation (STOIBER et al. 2015).

Co-purification of ORB2 with post-transcriptional negative regulators is consistent with

our analyses of the post-transcriptional fate of ORB2 targets during the MZT and our reporter

experiments in S2 cells.

The ORB2 ZZ domain is required for interaction with the 43S PIC and the Cup repressive

complex

Our S2-cell tethering assay, described above, showed that the ZZ domain of ORB2 is

necessary and sufficient for translational repression; therefore, we assessed whether this domain

is important for association with ORB2's identified binding partners in early embryos. To

generate an endogenous deletion of the ZZ domain, we used a scarless dsRed CRISPR method to

remove all amino acids from ORB2 between D638 and the C-terminus at C704, inclusive,

leaving the endogenous 3′UTR intact (see Figure 4d, Materials and Methods, Figures S1, S2)

(GRATZ *et al.* 2013; GRATZ *et al.* 2014). The mutant fly lines resulting from the excision will

henceforth be referred to as *orb2^{ΔZZ}*. Western blots on whole-fly extract showed that ORB2ΔZZ

was expressed at higher levels than full-length ORB2 in the pre-CRISPRed *w^{1118}* wild-type

control flies (Figure S1d) and a similar result was obtained for 0-3 hour embryos (Figure S9a).

We confirmed by RIP-RT-qPCR that deletion of the ZZ domain did not significantly affect

ORB2ΔZZ's ability to bind to several of its RNA targets in embryos (Figure S9b). As assayed by

DAPI staining, *orb2^{ΔZZ}* females showed no overt oogenesis defects; females outcrossed to the

control, host strain used for CRISPR produced embryos that underwent normal syncytial nuclear

cycles, cellularization and gastrulation, and exhibited similar hatch rates into first instar larvae as

the control (control: 92.5%. n = 200; *orb2^{ΔZZ}*: 90.7%, n = 300). We report elsewhere that *orb2^{ΔZZ}*

mutants have defects in brain development (HAILSTOCK *et al.* 2025) and spermatogenesis (LOW

et al. 2025).

Immunopurification of ORB2ΔZZ protein from 0-3-hour embryos produced by *orb2^{ΔZZ}*

homozygous females (henceforth referred to as *orb2^{ΔZZ}* embryos) followed by mass spectrometry

identified 48 high-confidence protein interactors in the presence (Figure 5b, S10b) and 91 in the

absence of RNase A (Figure S10f), of which 43 overlapped (File S4). When the union of these

interactors (96 proteins) was compared to the union of full-length ORB2 IP-MS interactors (233

proteins), we found that 65 overlapped, including several translational repressors such as PUM

and RBFOX1 (Figure 5b, File S4). Strikingly, interactions with all components of the PIC and

the Cup repressive complex were lost upon deletion of the ZZ domain (compare Figure 5a and

5b).

We conclude that the ORB2 ZZ domain is required for interaction between ORB2 and the

43S PIC as well as the Cup repressive complex, but that additional repressors retain interaction

with ORB2 in the absence of its ZZ domain.

Deletion of the ZZ domain results in a shift of ORB2 targets onto polysomes in the early

embryo

Since our experiments in S2 cells showed that the ZZ domain is necessary for ORB2-

mediated translational repression, and our IP-MS results indicated that ORB2 loses interactions

with the Cup repressive complex when the ZZ domain is deleted, we next asked whether deletion

of the ZZ domain had consequences for post-transcriptional regulation of ORB2's target mRNAs

during the MZT. Notably, since embryos from *orb2$^{\Delta ZZ}$* females undergo normal development and

binding to targets is unaffected by loss of the ZZ domain, any alterations in the post-

transcriptional fate of targets are likely to be a direct effect of the domain deletion rather than a

secondary consequence of abnormal development.

First, we assessed whether ORB2's targets, which are enriched for maternal transcripts that

are deadenylated and degraded during the MZT (see Figure 2e,f), were stabilized in *orb2$^{\Delta ZZ}$*

mutant embryos. We collected extracts from embryos laid by control or *orb2$^{\Delta ZZ}$* females at 0-1,

1-2 and 2-3 hours after egg lay and carried out RNA-seq followed by analysis using DEseq2. We

compared the expression level of targets and co-expressed non-targets across this time course

and found that, at all three stages, there was no significant difference in expression level between

control versus ORB2ΔZZ embryos for either target or non-target transcripts (Figure 6a and File

S7). These results suggest that, while ORB2 binds to unstable mRNAs, it does not induce their

degradation or, alternatively, that ORB2 may direct destabilization of its targets during the MZT

in a ZZ domain-independent mechanism.

Second, we assessed whether ORB2's targets, which are enriched for transcripts with

significantly lower TE and TI than co-expressed non-targets (see Figure 2a-d), underwent an

increase in polysome association in ORB2ΔZZ embryos. We collected embryo extracts from

embryos laid by control or *orb2$^{\Delta ZZ}$* females at 0-1, 1-2 and 2-3 hours after egg lay, then ran

polysome gradients followed by RNA-seq of pooled polysome fractions and pooled 'free' (non-

polysome-associated) fractions. For each gene, we calculated the ratio of RNA in the polysome
fractions to that in the free fractions (see Materials and Methods); if translational repression were
relieved upon deletion of the ZZ domain then this ratio would increase (i.e., transcripts would
move onto polysomes). We found that, in ORB2ΔZZ embryos, ORB2 targets shifted onto
polysomes at the first timepoint (0-1 hour) whereas non-targets did not (Figure 6b and File S7).

These data are consistent with the hypothesis that the ZZ domain of ORB2 mediates
translational repression of ORB2 targets early in the MZT, prior to ZGA. We note that we have
not directly demonstrated an increase in protein synthesis from these target transcripts in
embryos; however our S2 cell experiments showed that the ZZ domain is both necessary and
sufficient to repress luciferase protein expression without a change in reporter RNA levels,
consistent with the ZZ domain acting in translational repression.

Co-regulation of ORB2 target transcripts by additional RBPs
Although the shift onto polysomes of ORB2 targets at 0-1 hours in ORB2ΔZZ embryos
versus wild type is statistically significant ($P < 0.006$), it is subtle. We, therefore, decided to
assess whether ORB2 might function together with other post-transcriptional regulators known
to act during the MZT.

First, we asked whether there was a significant overlap of ORB2's targets with the targets
of several RBPs that we had analyzed previously. We found that ORB2 targets were depleted for
targets of Rasputin (RIN), an RBP that positively regulates transcript stability and translation
during the MZT (Table 5) (LAVER *et al.* 2020a), consistent with a role for ORB2 in negative
rather than positive regulation of its targets. In contrast, ORB2 targets significantly overlapped
those of PUM, BRAT and SMG (Table 5), RBPs that direct mRNA repression/degradation
during the MZT (TADROS *et al.* 2007; CHEN *et al.* 2014; LAVER *et al.* 2015).

For each of PUM, BRAT and SMG, we subdivided ORB2's targets into ones bound only
by ORB2, ones bound by both ORB2 and that RBP, ones bound only by that RBP, and co-
expressed unbound transcripts (see Materials and Methods, File S8). We found that, upon loss of
the ZZ domain, ORB2 targets bound by ORB2 but not SMG exhibited a significant shift onto
polysomes ($P = 0.009$; Figure 7b 'ORB2 only') whereas those co-bound by SMG (Figure 7a),
bound by SMG but not ORB2 (Figure 7c) or bound by neither ORB2 nor SMG (Figure 7d) did
not significantly shift onto polysomes. This result is consistent with the hypothesis that co-

regulation of a subset of ORB2's targets by SMG maintains repression (i.e., they do not shift

onto polysomes when the ZZ domain is deleted) and explains at least in part the subtle albeit

significant shift seen when ORB2 targets are aggregated (Figure 6b). Consistent with the SMG

analyses, ORB2 targets co-regulated (or not) by BRAT and PUM, exhibited a similar trend but

with the ORB2-only targets just missing significance ($P = 0.062$ and 0.073 for the PUM and

BRAT comparisons, respectively; Figures S11 and S12).

These data are consistent with the hypothesis that a significant subset of ORB2's targets

are coregulated by additional repressive RBPs, which attenuate the effect of loss of ORB2

repression upon deletion of the ZZ domain.

DISCUSSION

In this study, we identified hundreds of RNA targets and protein partners of the Drosophila

ORB2 protein in the early embryo. ORB2's targets are enriched for translationally repressed

mRNAs that are degraded during the MZT. ORB2-target open reading frames are enriched for

rare codons and a U-rich motif in their 3′UTR, the latter of which directs binding and repression

of a reporter in S2 cells. Repression by ORB2 is mediated by the ZZ domain, a function that is

conserved in hCPEB2. Deletion of the ZZ domain from the endogenous ORB2 gene results in

loss of physical interaction with the Cup repressive complex and movement of ORB2's mRNA

targets onto polysomes in the early embryo. Subsets of ORB2's targets are co-bound and likely

co-regulated by SMG.

A conserved role for the ORB2/hCPEB2 ZZ domain in mediating translational repression

Structural analysis of the ZZ domain of hCPEB1 has implicated it in protein-protein

interaction (MERKEL et al. 2013; STEPIEN et al. 2016), consistent with earlier studies on, for

example, the ZZ domain of CREB-binding protein (LEGGE et al. 2004). Our identification of

multiple proteins that depend in vivo on the ZZ domain for interaction with ORB2 are consistent

with a major role in recruiting co-regulators to target transcripts. Our data showing that deletion

of the ZZ domain does not affect RNA binding are consistent with a previous study (STEPIEN et

al. 2016). The fact that, in our S2 cell tethering assay, the ZZ domain is necessary and sufficient

for repression permitted us to assess evolutionary conservation of this role as well as to map

regions of the ZZ domain that implement this function.

Interestingly, we found that the repressive function of the entire ZZ domain is conserved
between ORB2 and hCPEB2 but not with ORB and hCPEB1. Chimeras containing regions from
the ORB2 and ORB ZZ domain showed that the α-helix (region C, C669 to H685) of ORB2 can
convert the ORB ZZ domain into a repressor when paired with one of the following: ORB2's Rd
turn at the N-terminus (Region A, Q640 to P656), the beta hairpin (Region B, F657-Y668), or
the C-terminal beta-sheet region (Region D, H685 to C-terminus). The ZZ domains of hCPEB3
and 4 show ≥ 90% amino acid identity to hCPEB2, with 16/17 residues of region A, 14/15 of
region B, 12/13 of region C, and 15/20 of D identical between hCPEB2-4. As such, it is quite
likely that the ZZ domains of hCPEB3 and 4 are also capable of repressing target mRNA
translation via region C.

That fact that region C of ORB2's ZZ domain needs one of A, B or D to convert the ORB
ZZ domain into a translational repressor is consistent with a model in which A, B or D bind
different co-repressors, any of which cooperate with C to mediate repression, or a model in
which A, B or D establish a structural context for C to interact with co-repressor(s). Future work
aimed at identifying the ORB2 binding partners that interact with each of regions A-D should
provide insights into the molecular mechanisms that underlie ZZ domain function.
934

Regulation of post-transcriptional processes by ORB2 during the MZT: features of ORB2's
targets and their implications
Three features in ORB2's target transcripts have implications for post-transcriptional
regulation during the MZT: enrichment for OREs, enrichment for rare codons, and enrichment
for binding sites for additional RBPs (notably SMG, BRAT and PUM).
First, enrichment for OREs and our evidence from S2 cell reporter assays that ORB2 binds
to OREs, which then confers repression on the reporter, are consistent with previous studies of
the binding specificity of ORB2 and ORB2B's role as a repressor (KHAN *et al.* 2015; STEPIEN *et*
al. 2016). Notably, since ORB2B is the only isoform expressed during the MZT, we conclude
that the *orb2* gene's function during the MZT is exclusively negative, which contrasts with its
role in other tissues. For example, in the nervous system, ORB2A and ORB2B are co-expressed,
permitting hetero-oligomer formation and translational upregulation of ORB2 targets
(MAJUMDAR *et al.* 2012; KHAN *et al.* 2015). Thus, expression specifically of the ORB2B isoform
during the MZT is a mechanism to negatively regulate the post-transcriptional fate of its targets.

Second, enrichment of ORB2's targets for rare codons is consistent with previous

observations in neurons, where it has been shown that ORB2 binds rare-codon enriched

transcripts (STEWART *et al.* 2024). In contrast to neurons, where the outcome is upregulation of

transcript stability and translation, embryonic targets of ORB2 are repressed and degraded. As

discussed above, this may be in part due to an absence of ORB2A and a consequence of only

having ORB2B present, but it may also be because of additional specific, negative-regulatory co-

factors that interact with ORB2 and/or ORB2 targets in embryos, as discussed in the next

section. Either way, the observation that ORB2 targets are enriched for rare codons and are

downregulated is consistent with the larger literature on the outcome of non-optimal codon

usage, including during the MZT of flies and vertebrates (PRESNYAK *et al.* 2015; BAZZINI *et al.*

2016; IRIARTE *et al.* 2021).

Third, we have shown that significant subsets of ORB2's targets are co-bound during the

MZT by SMG, BRAT and PUM. Our analyses of the ORB2 targets that move onto polysomes

when the ZZ domain is deleted are consistent with the hypothesis that co-regulated targets (e.g.,

those bound by both ORB2 and SMG) continue to be repressed, likely because SMG directs

ongoing repression, whereas ORB2-targets not bound by SMG shift onto polysomes. Whether

the latter are co-bound by additional unidentified RBPs is not yet known, but the fact that they

move onto polysomes suggests that, even if co-bound by such RBPs, those RBPs are not

sufficient for repression, at least not at the level of translation initiation.

Regulation of post-transcriptional processes by ORB2 during the MZT: the role of

repressive co-factors

Our analyses of the post-transcriptional fate of ORB2's targets during the MZT indicate

that they are enriched for transcripts that are translationally repressed and degraded. Consistent

with this outcome, multiple co-factors that are known to mediate repression and degradation of

mRNAs copurify with ORB2 in early embryos; most of these copurify in the presence of RNase,

consistent with an RNA-independent interaction with ORB2. Several of the regulators that

interact in an RNA-independent manner require the ZZ domain for interaction, notably the Cup

complex components Cup, TRAL, ME31B and BEL.

We have shown that the ZZ domain is necessary and sufficient for translational repression

of reporters in S2 cells and deletion of the ZZ domain results in a shift of ORB2's *in vivo* targets

onto polysomes in 0-1-hour embryos. That the Cup complex is expressed in both S2 cells and

embryos, that ZZ domain deletion results in loss of Cup complex interaction with ORB2 in

embryos, and that ORB2's targets in embryos move onto polysomes when the ZZ domain is

deleted from endogenous ORB2, are all consistent with the hypothesis that ORB2-dependent

repression is mediated by the Cup complex. The Cup complex prevents interaction of eIF4G

with eIF4E, thus abrogating translation initiation (WILHELM *et al.* 2003; NAKAMURA *et al.* 2004;

NELSON *et al.* 2004a); consistent with this role, we note that, while eIF4E was present in our

ORB2 IP-MS data, eIF4G was absent. Previous studies have shown that the Cup complex is

brought to mRNAs by BRU or Squid (SQD) in oocytes and by SMG in embryos (WILHELM *et al.*

2003; NAKAMURA *et al.* 2004; NELSON *et al.* 2004a; CLOUSE *et al.* 2008). ORB2, thus, represents

a second RBP that acts in embryos to recruit this complex for repression, in this case early in the

MZT. A caveat is that we have not demonstrated that the targets that shift onto polysomes are

translated; post-initiation mechanisms may act to prevent protein synthesis.

The Cup complex has been shown to be cleared by the CTLH E3 ligase part-way through

the MZT (CAO *et al.* 2020; ZAVORTINK *et al.* 2020), consistent with the ZZ domain's repressive

role in the early but not late MZT. However, ORB2's targets are repressed at both 0-1 hours

(when the Cup complex is present) and at 2-3 hours (after the Cup complex has been cleared);

furthermore, ORB2's targets are deadenylated and degraded between 0-1 and 2-3 hours, but

deletion of the ZZ domain has no effect on their clearance. These data suggest that additional

regulators may function either independent of or together with ORB2 to direct these processes.

SMG is an example of a co-regulator present throughout the MZT that acts through the Cup

complex for repression and through the CCR4-NOT deadenylase to direct degradation (NELSON

et al. 2004b; SEMOTOK *et al.* 2005). SMG's co-purification with ORB2 in our IP-MS

experiments is RNA-dependent; thus, SMG's binding and regulation of a subset of ORB2-targets

may be independent of physical interaction with ORB2. PUM, FMR1 and RBFOX1 are also

present throughout the MZT (CASAS-VILA *et al.* 2017; CAO *et al.* 2020) and we have shown here

that their interaction with ORB2 is both RNA-independent and ZZ-domain independent, making

them candidates for implementing these roles together with ORB2 but independent of its ZZ

domain.

1009

1010   **The mystery of PIC interaction**

Unexpectedly, we identified the 43S PIC as a major interactor of ORB2 via its ZZ domain. An essential component of the PIC is the eIF3 complex (its subunits are encoded by 14 genes in *Drosophila*), which is a highly regulated component of translation initiation (HARRIS *et al.* 2006; CHIU *et al.* 2010; CATE 2017). Several of the mammalian eIF3 subunits have been shown to have RNA-binding ability (LEE *et al.* 2015; LEE *et al.* 2016) and to mediate distinct regulatory outcomes (PESTOVA *et al.* 1998; PESTOVA AND HELLEN 1999; HUI *et al.* 2003; HUI *et al.* 2005; MARTINEAU *et al.* 2008; LEE *et al.* 2010; MARTINEAU *et al.* 2014).

At this time we do not have data that provide insights into the functional consequences of ORB2's interaction with the PIC. ORB2's interaction with the PIC's eIF3 subunits is RNA-independent, consistent with direct interaction with one or more of these subunits. We considered one possibile mechanism where ORB2 may sequester the PIC, thus limiting its availability and reducing translation overall. However, in such a model, one would expect a global increase in translation upon loss of PIC interaction rather than what we observed in *orb2$^{\Delta ZZ}$* embryos, which is an increase in polysome association of ORB2's targets but no change in non-targets. An alternative hypothesis is that ORB2-dependent binding to the PIC on target transcripts has a negative effect on translation, either in addition to or in concert with the Cup repressive complex. Indeed, a speculative model might be that ORB2 binding to the 3′UTR and its interaction with PIC at the 5′UTR might repress translation initiation by interfering with PIC function or enhancing the function of repressive elements of the eIF3 complex. The mechanistic and functional consequences of ORB2-PIC interaction will be a fruitful area for future study.

Data Availability

Drosophila strains are available upon request. Raw data for RIP-seq, RNA-seq and IP-MS are available at GEO (GSE299661 for RIP-seq, GSE299140 for RNA-seq time-course, GSE299019 for polysome gradient seq) and at MassIVE (MSV000098275 for IP-MS). The authors affirm that all of the other data necessary for confirming the conclusions of the article are present within the article, figures, tables and supplementary files.

Acknowledgements

1040   We thank Kun Nie for assisting with the *de novo* motif discovery using #ATS, Dr. Dorothy

1041   Lerit and Dr. Donald Fox for feedback on draft and preprint versions of this manuscript,

1042   respectively.

1043

Funding

This research was supported by grants from the Canadian Institutes of Health Research

PJT-159702 (HDL), PJT-190124 (HDL) and PJT-159494 (CAS). TCHL was supported in part

by a Canada Graduate Scholarship (CGS-M) and a University of Toronto Open Fellowship; CW

and ZW were supported in part by University of Toronto Open Fellowships. Molecular graphics

and analyses were performed with UCSF ChimeraX, developed by the Resource for

Biocomputing, Visualization, and Informatics at the University of California, San Francisco,

with support from National Institutes of Health R01-GM129325 and the Office of Cyber

Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases.

1053

1054   **Conflicts of Interest**

1055   The authors declare that they have no conflicts of interest.

1056   **LITERATURE CITED**

1057

Abramson, J., J. Adler, J. Dunger, R. Evans, T. Green *et al.*, 2024 Accurate structure prediction
of biomolecular interactions with AlphaFold 3. Nature 630**:** 493-500.
Bazzini, A. A., F. Del Viso, M. A. Moreno-Mateos, T. G. Johnstone, C. E. Vejnar *et al.*, 2016
Codon identity regulates mRNA stability and translation efficiency during the maternal-
to-zygotic transition. EMBO J 35**:** 2087-2103.
Bernhart, S. H., I. L. Hofacker and P. F. Stadler, 2006 Local RNA base pairing probabilities in
large sequences. Bioinformatics 22**:** 614-615.
Cao, W. X., S. Kabelitz, M. Gupta, E. Yeung, S. Lin *et al.*, 2020 Precise Temporal Regulation of
Post-transcriptional Repressors Is Required for an Orderly Drosophila Maternal-to-
Zygotic Transition. Cell Rep 31**:** 107783.
Casas-Vila, N., A. Bluhm, S. Sayols, N. Dinges, M. Dejung *et al.*, 2017 The developmental
proteome of Drosophila melanogaster. Genome Res 27**:** 1273-1285.
Cate, J. H., 2017 Human eIF3: from 'blobology' to biological insight. Philos Trans R Soc Lond B
Biol Sci 372.
Chen, L., J. G. Dumelie, X. Li, M. H. Cheng, Z. Yang *et al.*, 2014 Global regulation of mRNA
translation and stability in the early Drosophila embryo by the Smaug RNA-binding
protein. Genome Biol 15**:** R4.
Chiu, C. W. N., C. Monat, M. Robitaille, M. Lacomme, A. M. Daulat *et al.*, 2016 SAPCD2
Controls Spindle Orientation and Asymmetric Divisions by Negatively Regulating the
Galphai-LGN-NuMA Ternary Complex. Dev Cell 36**:** 50-62.
Chiu, W. L., S. Wagner, A. Herrmannová, L. Burela, F. Zhang *et al.*, 2010 The C-terminal
region of eukaryotic translation initiation factor 3a (eIF3a) promotes mRNA recruitment,
scanning, and, together with eIF3j and the eIF3b RNA recognition motif, selection of
AUG start codons. Mol Cell Biol 30**:** 4415-4434.
Choi, H., G. Liu, D. Mellacheruvu, M. Tyers, A. C. Gingras *et al.*, 2012 Analyzing protein-
protein interactions from affinity purification-mass spectrometry data with SAINT. Curr
Protoc Bioinformatics Chapter 8**:** Unit8.15.
Clouse, K. N., S. B. Ferguson and T. Schupbach, 2008 Squid, Cup, and PABP55B function
together to regulate gurken translation in Drosophila. Dev Biol 313**:** 713-724.
de Moor, C. H., and J. D. Richter, 1999 Cytoplasmic polyadenylation elements mediate masking
and unmasking of cyclin B1 mRNA. EMBO J 18**:** 2294-2303.
De Renzis, S., O. Elemento, S. Tavazoie and E. F. Wieschaus, 2007 Unmasking activation of the
zygotic genome using chromosomal deletions in the Drosophila embryo. PLoS Biol 5**:**
e117.
Dunn, J. G., C. K. Foo, N. G. Belletier, E. R. Gavis and J. S. Weissman, 2013 Ribosome
profiling reveals pervasive and regulated stop codon readthrough in Drosophila
melanogaster. Elife 2**:** e01179.
Duran-Arqué, B., M. Cañete, C. L. Castellazzi, A. Bartomeu, A. Ferrer-Caelles *et al.*, 2022
Comparative analyses of vertebrate CPEB proteins define two subfamilies with
coordinated yet distinct functions in post-transcriptional gene regulation. Genome Biol
23**:** 192.

Eichhorn, S. W., A. O. Subtelny, I. Kronja, J. C. Kwasnieski, T. L. Orr-Weaver *et al.*, 2016
mRNA poly(A)-tail changes specified by deadenylation broadly reshape translation in
Drosophila oocytes and early embryos. Elife 5.

Fernández-Miranda, G., and R. Méndez, 2012 The CPEB-family of proteins, translational
control in senescence and cancer. Ageing Res Rev 11**:** 460-472.

Franz, M., H. Rodriguez, C. Lopes, K. Zuberi, J. Montojo *et al.*, 2018 GeneMANIA update
2018. Nucleic Acids Res 46**:** W60-W64.

Gene Ontology, C., S. A. Aleksander, J. Balhoff, S. Carbon, J. M. Cherry *et al.*, 2023 The Gene
Ontology knowledgebase in 2023. Genetics 224.

Gilmutdinov, R., E. N. Kozlov, K. V. Yakovlev, L. V. Olenina, A. A. Kotov *et al.*, 2021 The
3'UTR of the Drosophila CPEB translation factor gene orb2 plays a crucial role in
spermatogenesis. Development 148.

Gotze, M., J. Dufourt, C. Ihling, C. Rammelt, S. Pierson *et al.*, 2017 Translational repression of
the Drosophila nanos mRNA involves the RNA helicase Belle and RNA coating by
Me31B and Trailer hitch. RNA 23**:** 1552-1568.

Gratz, S. J., A. M. Cummings, J. N. Nguyen, D. C. Hamm, L. K. Donohue *et al.*, 2013 Genome
engineering of Drosophila with the CRISPR RNA-guided Cas9 nuclease. Genetics 194**:**
1029-1035.

Gratz, S. J., F. P. Ukken, C. D. Rubinstein, G. Thiede, L. K. Donohue *et al.*, 2014 Highly
specific and efficient CRISPR/Cas9-catalyzed homology-directed repair in Drosophila.
Genetics 196**:** 961-971.

Groisman, I., Y. S. Huang, R. Mendez, Q. Cao and J. D. Richter, 2001 Translational control of
embryonic cell division by CPEB and maskin. Cold Spring Harb Symp Quant Biol 66**:**
345-351.

Hafer, N., S. Xu, K. M. Bhat and P. Schedl, 2011 The Drosophila CPEB protein Orb2 has a
novel expression pattern and is important for asymmetric cell division and nervous
system function. Genetics 189**:** 907-921.

Hailstock, T., J. Buehler, B. V. Robinson, T. C. H. Low, Y. Hua *et al.*, 2025 The CPEB ortholog
Orb2 regulates brain size through the TRIM-NHL RNA-binding protein, Brain tumor.
bioRxiv.

Hake, L. E., and J. D. Richter, 1994 CPEB is a specificity factor that mediates cytoplasmic
polyadenylation during Xenopus oocyte maturation. Cell 79**:** 617-627.

Harris, T. E., A. Chi, J. Shabanowitz, D. F. Hunt, R. E. Rhoads *et al.*, 2006 mTOR-dependent
stimulation of the association of eIF4G and eIF3 by insulin. EMBO J 25**:** 1659-1668.

Harrison, M. M., A. J. Marsh and C. A. Rushlow, 2023 Setting the stage for development: the
maternal-to-zygotic transition in Drosophila. Genetics 225.

Haugen, R. J., C. Barnier, N. D. Elrod, H. Luo, M. K. Jensen *et al.*, 2024 Regulation of the
Drosophila transcriptome by Pumilio and the CCR4-NOT deadenylase complex. RNA
30**:** 866-890.

Hervás, R., L. Li, A. Majumdar, M. e. C. Fernández-Ramírez, J. R. Unruh *et al.*, 2016 Molecular
Basis of Orb2 Amyloidogenesis and Blockade of Memory Consolidation. PLoS Biol 14**:**
e1002361.

Hervas, R., M. J. Rau, Y. Park, W. Zhang, A. G. Murzin *et al.*, 2020 Cryo-EM structure of a
neuronal functional amyloid implicated in memory persistence in *Drosophila*. Science
367**:** 1230-1234.

Hui, D. J., C. R. Bhasker, W. C. Merrick and G. C. Sen, 2003 Viral stress-inducible protein p56 inhibits translation by blocking the interaction of eIF3 with the ternary complex eIF2.GTP.Met-tRNAi. J Biol Chem 278: 39477-39482.

Hui, D. J., F. Terenzi, W. C. Merrick and G. C. Sen, 2005 Mouse p56 blocks a distinct function of eukaryotic initiation factor 3 in translation initiation. J Biol Chem 280: 3433-3440.

Iriarte, A., G. Lamolle and H. Musto, 2021 Codon Usage Bias: An Endless Tale. J Mol Evol 89: 589-593.

Ivshina, M., P. Lasko and J. D. Richter, 2014 Cytoplasmic polyadenylation element binding proteins in development, health, and disease. Annu Rev Cell Dev Biol 30: 393-415.

Jiang, T., R. F. McKinley, M. A. McGill, S. Angers and T. J. Harris, 2015 A Par-1-Par-3-Centrosome Cell Polarity Pathway and Its Tuning for Isotropic Cell Adhesion. Curr Biol 25: 2701-2708.

Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov *et al.*, 2021 Applying and improving AlphaFold at CASP14. Proteins 89: 1711-1721.

Keleman, K., S. Krüttner, M. Alenius and B. J. Dickson, 2007 Function of the Drosophila CPEB protein Orb2 in long-term courtship memory. Nat Neurosci 10: 1587-1593.

Khan, M. R., L. Li, C. Pérez-Sánchez, A. Saraf, L. Florens *et al.*, 2015 Amyloidogenic Oligomerization Transforms Drosophila Orb2 from a Translation Repressor to an Activator. Cell 163: 1468-1483.

Kim, J. H., and J. D. Richter, 2006 Opposing polymerase-deadenylase activities regulate cytoplasmic polyadenylation. Mol Cell 24: 173-183.

Kolberg, L., U. Raudvere, I. Kuzmin, P. Adler, J. Vilo *et al.*, 2023 g:Profiler-interoperable web service for functional enrichment analysis and gene identifier mapping (2023 update). Nucleic Acids Res 51: W207-W212.

Kopylova, E., L. Noe and H. Touzet, 2012 SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. Bioinformatics 28: 3211-3217.

Kozlov, E., Y. V. Shidlovskii, R. Gilmutdinov, P. Schedl and M. Zhukova, 2021 The role of CPEB family proteins in the nervous system function in the norm and pathology. Cell Biosci 11: 64.

Kozlov, E. N., E. V. Tokmatcheva, A. M. Khrustaleva, E. S. Grebenshchikov, R. V. Deev *et al.*, 2023 Long-Term Memory Formation in Drosophila Depends on the 3'UTR of CPEB Gene orb2. Cells 12.

Kronja, I., B. Yuan, S. W. Eichhorn, K. Dzeyk, J. Krijgsveld *et al.*, 2014 Widespread changes in the posttranscriptional landscape at the Drosophila oocyte-to-embryo transition. Cell Rep 7: 1495-1508.

Krüttner, S., B. Stepien, J. N. Noordermeer, M. A. Mommaas, K. Mechtler *et al.*, 2012 Drosophila CPEB Orb2A mediates memory independent of Its RNA-binding domain. Neuron 76: 383-395.

Laver, J. D., K. Ancevicius, P. Sollazzo, J. T. Westwood, S. S. Sidhu *et al.*, 2012 Synthetic antibodies as tools to probe RNA-binding protein function. Mol Biosyst 8: 1650-1657.

Laver, J. D., X. Li, D. Ray, K. B. Cook, N. A. Hahn *et al.*, 2015 Brain tumor is a sequence-specific RNA-binding protein that directs maternal mRNA clearance during the Drosophila maternal-to-zygotic transition. Genome Biol 16: 94.

Laver, J. D., J. Ly, A. K. Winn, A. Karaiskakis, S. Lin *et al.*, 2020a The RNA-Binding Protein Rasputin/G3BP Enhances the Stability and Translation of Its Target mRNAs. Cell Rep 30: 3353-3367.e3357.

Laver, J. D., J. Ly, A. K. Winn, A. Karaiskakis, S. Lin *et al.*, 2020b The RNA-Binding Protein Rasputin/G3BP Enhances the Stability and Translation of Its Target mRNAs. Cell Rep 30**:** 3353-3367 e3357.

Lécuyer, E., H. Yoshida, N. Parthasarathy, C. Alm, T. Babak *et al.*, 2007 Global analysis of mRNA localization reveals a prominent role in organizing cellular architecture and function. Cell 131**:** 174-187.

Lee, A. S., P. J. Kranzusch and J. H. Cate, 2015 eIF3 targets cell-proliferation messenger RNAs for translational activation or repression. Nature 522**:** 111-114.

Lee, A. S., P. J. Kranzusch, J. A. Doudna and J. H. Cate, 2016 eIF3d is an mRNA cap-binding protein that is required for specialized translation initiation. Nature 536**:** 96-99.

Lee, J. P., A. Brauweiler, M. Rudolph, J. E. Hooper, H. A. Drabkin *et al.*, 2010 The TRC8 ubiquitin ligase is sterol regulated and interacts with lipid and protein biosynthetic pathways. Mol Cancer Res 8**:** 93-106.

Legge, G. B., M. A. Martinez-Yamout, D. M. Hambly, T. Trinh, B. M. Lee *et al.*, 2004 ZZ domain of CBP: an unusual zinc finger fold in a protein interaction module. J Mol Biol 343**:** 1081-1093.

Li, X., G. Quon, H. D. Lipshitz and Q. Morris, 2010 Predicting in vivo binding sites of RNA-binding proteins using mRNA secondary structure. RNA 16**:** 1096-1107.

Liu, G., J. Zhang, H. Choi, J. P. Lambert, T. Srikumar *et al.*, 2012 Using ProHits to store, annotate, and analyze affinity purification-mass spectrometry (AP-MS) data. Curr Protoc Bioinformatics Chapter 8**:** 8.16.11-18.16.32.

Liu, G., J. Zhang, B. Larsen, C. Stark, A. Breitkreutz *et al.*, 2010 ProHits: integrated software for mass spectrometry-based interaction proteomics. Nat Biotechnol 28**:** 1015-1017.

Liu, Y. C., A. L. Couzens, A. R. Deshwar, B. M.-C. LD, X. Zhang *et al.*, 2014 The PPFIA1-PP2A protein complex promotes trafficking of Kif7 to the ciliary tip and Hedgehog signaling. Sci Signal 7**:** ra117.

Lorenz, R., S. H. Bernhart, C. Honer Zu Siederdissen, H. Tafer, C. Flamm *et al.*, 2011 ViennaRNA Package 2.0. Algorithms Mol Biol 6**:** 26.

Love, M. I., W. Huber and S. Anders, 2014 Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol 15**:** 550.

Love, M. I., C. Soneson, P. F. Hickey, L. K. Johnson, N. T. Pierce *et al.*, 2020 Tximeta: Reference sequence checksums for provenance identification in RNA-seq. PLoS Comput Biol 16**:** e1007664.

Low, T. C. H., B. L. Falk, J. A. Brill and H. D. Lipshitz, 2025 The C-terminal ZZ domain of the Drosophila ORB2 RNA-binding protein is required for spermatid individualization. bioRxiv**:** 2025.2008.2022.671863.

Majumdar, A., W. C. Cesario, E. White-Grindley, H. Jiang, F. Ren *et al.*, 2012 Critical role of amyloid-like oligomers of Drosophila Orb2 in the persistence of memory. Cell 148**:** 515-529.

Martineau, Y., M. C. Derry, X. Wang, A. Yanagiya, J. J. Berlanga *et al.*, 2008 Poly(A)-binding protein-interacting protein 1 binds to eukaryotic translation initiation factor 3 to stimulate translation. Mol Cell Biol 28**:** 6658-6667.

Martineau, Y., X. Wang, T. Alain, E. Petroulakis, D. Shahbazian *et al.*, 2014 Control of Paip1-eukayrotic translation initiation factor 3 interaction by amino acids through S6 kinase. Mol Cell Biol 34**:** 1046-1053.

Mastushita-Sakai, T., E. White-Grindley, J. Samuelson, C. Seidel and K. Si, 2010 Drosophila Orb2 targets genes involved in neuronal growth, synapse formation, and protein turnover. Proc Natl Acad Sci U S A 107**:** 11987-11992.

Meng, E. C., T. D. Goddard, E. F. Pettersen, G. S. Couch, Z. J. Pearson *et al.*, 2023 UCSF ChimeraX: Tools for structure building and analysis. Protein Sci 32**:** e4792.

Merkel, D. J., S. B. Wells, B. C. Hilburn, F. Elazzouzi, G. C. Pérez-Alvarado *et al.*, 2013 The C-terminal region of cytoplasmic polyadenylation element binding protein is a ZZ domain with potential for protein-protein interactions. J Mol Biol 425**:** 2015-2026.

Montojo, J., K. Zuberi, H. Rodriguez, F. Kazi, G. Wright *et al.*, 2010 GeneMANIA Cytoscape plugin: fast gene function predictions on the desktop. Bioinformatics 26**:** 2927-2928.

Na, H., J. D. Laver, J. Jeon, F. Singh, K. Ancevicius *et al.*, 2016 A high-throughput pipeline for the production of synthetic antibodies for analysis of ribonucleoprotein complexes. RNA 22**:** 636-655.

Nakamura, A., K. Sato and K. Hanyu-Nakamura, 2004 Drosophila cup is an eIF4E binding protein that associates with Bruno and regulates oskar mRNA translation in oogenesis. Dev Cell 6**:** 69-78.

Nakamura, Y., T. Gojobori and T. Ikemura, 2000 Codon usage tabulated from international DNA sequence databases: status for the year 2000. Nucleic Acids Res 28**:** 292.

Nelson, M. R., A. M. Leidal and C. A. Smibert, 2004a Drosophila Cup is an eIF4E-binding protein that functions in Smaug-mediated translational repression. The EMBO journal 23**:** 150-159.

Nelson, M. R., A. M. Leidal and C. A. Smibert, 2004b Drosophila Cup is an eIF4E-binding protein that functions in Smaug-mediated translational repression. EMBO J 23**:** 150-159.

Ozturk-Colak, A., S. J. Marygold, G. Antonazzo, H. Attrill, D. Goutte-Gattat *et al.*, 2024 FlyBase: updates to the Drosophila genes and genomes database. Genetics 227.

Patro, R., G. Duggal, M. I. Love, R. A. Irizarry and C. Kingsford, 2017 Salmon provides fast and bias-aware quantification of transcript expression. Nat Methods 14**:** 417-419.

Pestova, T. V., and C. U. Hellen, 1999 Internal initiation of translation of bovine viral diarrhea virus RNA. Virology 258**:** 249-256.

Pestova, T. V., I. N. Shatsky, S. P. Fletcher, R. J. Jackson and C. U. Hellen, 1998 A prokaryotic-like mode of cytoplasmic eukaryotic ribosome binding to the initiation codon during internal translation initiation of hepatitis C and classical swine fever virus RNAs. Genes Dev 12**:** 67-83.

Presnyak, V., N. Alhusaini, Y. H. Chen, S. Martin, N. Morris *et al.*, 2015 Codon optimality is a major determinant of mRNA stability. Cell 160**:** 1111-1124.

Puigbo, P., I. G. Bravo and S. Garcia-Vallve, 2008 CAIcal: a combined set of tools to assess codon usage adaptation. Biol Direct 3**:** 38.

Ray, D., H. Kazan, K. B. Cook, M. T. Weirauch, H. S. Najafabadi *et al.*, 2013 A compendium of RNA-binding motifs for decoding gene regulation. Nature 499**:** 172-177.

Semotok, J. L., R. L. Cooperstock, B. D. Pinder, H. K. Vari, H. D. Lipshitz *et al.*, 2005 Smaug recruits the CCR4/POP2/NOT deadenylase complex to trigger maternal transcript localization in the early Drosophila embryo. Curr Biol 15**:** 284-294.

Shannon, P., A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang *et al.*, 2003 Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res 13**:** 2498-2504.

1280   Standart, N., and N. Minshall, 2008 Translational control in early development: CPEB, P-bodies
1281   and germinal granules. Biochem Soc Trans 36: 671-676.
1282   Stepien, B. K., C. Oppitz, D. Gerlach, U. Dag, M. Novatchkova *et al.*, 2016 RNA-binding
1283   profiles of Drosophila CPEB proteins Orb and Orb2. Proc Natl Acad Sci U S A 113:
1284   E7030-E7038.
1285   Stewart, R. K., P. Nguyen, A. Laederach, P. C. Volkan, J. K. Sawyer *et al.*, 2024 Orb2 enables
1286   rare-codon-enriched mRNA expression during Drosophila neuron differentiation. Nat
1287   Commun 15: 5270.
1288   Stoiber, M. H., S. Olson, G. E. May, M. O. Duff, J. Manent *et al.*, 2015 Extensive cross-
1289   regulation of post-transcriptional regulatory networks in Drosophila. Genome Res 25:
1290   1692-1702.
1291   Supek, F., and K. Vlahovicek, 2005 Comparison of codon usage measures and their applicability
1292   in prediction of microbial gene expressivity. BMC Bioinformatics 6: 182.
1293   Tadros, W., A. L. Goldman, T. Babak, F. Menzies, L. Vardy *et al.*, 2007 SMAUG is a major
1294   regulator of maternal mRNA destabilization in Drosophila and its translation is activated
1295   by the PAN GU kinase. Dev Cell 12: 143-155.
1296   Tadros, W., and H. D. Lipshitz, 2009 The maternal-to-zygotic transition: a play in two acts.
1297   Development 136: 3033-3042.
1298   Thomsen, S., S. Anders, S. C. Janga, W. Huber and C. R. Alonso, 2010 Genome-wide analysis
1299   of mRNA decay patterns during early Drosophila development. Genome Biol 11: R93.
1300   Vastenhouw, N. L., W. X. Cao and H. D. Lipshitz, 2019 The maternal-to-zygotic transition
1301   revisited. Development 146.
1302   Wakiyama, M., Y. Kaitsu, R. Muramatsu, K. Takimoto and S. Yokoyama, 2012 Tethering of
1303   proteins to RNAs using the bovine immunodeficiency virus-Tat peptide and BIV-TAR
1304   RNA. Anal Biochem 427: 130-132.
1305   Wang, X. P., and N. G. Cooper, 2009 Characterization of the transcripts and protein isoforms for
1306   cytoplasmic polyadenylation element binding protein-3 (CPEB3) in the mouse retina.
1307   BMC Mol Biol 10: 109.
1308   Warde-Farley, D., S. L. Donaldson, O. Comes, K. Zuberi, R. Badrawi *et al.*, 2010 The
1309   GeneMANIA prediction server: biological network integration for gene prioritization and
1310   predicting gene function. Nucleic Acids Res 38: W214-220.
1311   Whitten, D., 2011 Classification and clustering of sequencing data using a Poisson model, pp.
1312   2493-2518. Annals of Applied Statistics.
1313   Wilhelm, J. E., M. Hilton, Q. Amos and W. J. Henzel, 2003 Cup is an eIF4E binding protein
1314   required for both the translational repression of oskar and the recruitment of Barentsz. J
1315   Cell Biol 163: 1197-1204.
1316   Xu, S., N. Hafer, B. Agunwamba and P. Schedl, 2012 The CPEB protein Orb2 has multiple
1317   functions during spermatogenesis in Drosophila melanogaster. PLoS Genet 8: e1003079.
1318   Zavortink, M., L. N. Rutt, S. Dzitoyeva, J. C. Henriksen, C. Barrington *et al.*, 2020 The E2
1319   Marie Kondo and the CTLH E3 ligase clear deposited RNA binding proteins during the
1320   maternal-to-zygotic transition. Elife 9.

1321

Figure Legends

Figure 1. RIP-seq identifies over 450 ORB2 targets in 0-3 hr embryos. (a) Domain structure of ORB2B and location of the antigen used to screen for phage-displayed synthetic antibodies. (b) Scatterplot depicting the Log_2(Fold Enrichment compared to Input) for all genes identified in the RIP-seq analysis; each point represents a single gene. Values represent the average enrichment from three independent biological replicates. ORB2 targets (red) are defined as having fold enrichment > 1.5 in the anti-ORB2 RIP-seq compared to input and a ratio of ORB2/Control > 1.5, with an FDR adjusted *P*-value < 0.05; genes enriched in both ORB2 and control RIP-seq (green) are defined as having fold enrichment > 1.5 in both anti-ORB2 and control RIP-seq, with a ratio of ORB2/Control < 1.5 and an FDR adjusted *P*-value < 0.05; Control enriched transcripts (blue) are defined as having fold enrichment < -1.5 in the control RIP-seq, fold enrichment < -1.5 in the anti-ORB2 RIP-seq, and an FDR adjusted *P*-value < 0.05. Dotted lines indicate ±1.5 fold-change thresholds. (c) Violin and boxplots depicting the distribution of transcript abundance as measured by average TPM from three biological replicates based on target status as described in (A). Wilcoxon Rank Sum Test with Bonferroni multiple-test correction: ** = $P < 10^{-10}$; *** = $P < 10^{-15}$. (d) Validation of RIP-seq targets by RT-qPCR; bar graphs depict the log_2(fold enrichment) in an anti-ORB2 RIP-RT-qPCR compared to control RIP-RT-qPCR; *P*-values given are derived from Wilcoxon Signed Rank Test comparing ORB2 RIP-RT-qPCR to control RIP-RT-qPCR.

Figure 2. ORB2 target transcripts are enriched in translationally repressed, unstable, rare-codon containing transcripts during the MZT. Violin plots and boxplots depicting the relationship between ORB2-association and translational status, change in poly(A) tail length, and change in transcript abundance. Non-target genes include all expressed genes that were not defined as ORB2 targets in the RIP-seq analysis. (a) Translation efficiency, calculated from ribosome footprinting performed on 0-1 hour embryo extracts (EICHHORN *et al.* 2016). (b) Translation efficiency, calculated from ribosome footprinting performed on 2-3 hour embryo extracts (EICHHORN *et al.* 2016). (c) Translation efficiency as measured by ribosome profiling performed on 0-2 hour embryo extracts (DUNN *et al.* 2013). (d) Translation index, calculated from polysome gradients performed on 0-3 hour embryo extracts (CHEN *et al.* 2014). (e) Change in transcript poly(A) tail length, calculated by comparing 0-1 hour and 2-3 hour embryo extracts, as measured by PAL-Seq (EICHHORN *et al.* 2016). (f) Change in abundance, calculated by comparing 0-1 hour and 2-3 hour embryo extracts, as measured by RNA-Seq (EICHHORN *et al.* 2016). (g) MELP score, calculated for all annotated CDS longer than 80 codons on FlyBase 6.58 by coRdon with ribosomal proteins as the references for codon optimization, and the isoform with minimum score (the worst expressivity) is chosen to represent each expressed gene. (h) CAI score, calculated for all annotated CDS longer than 80 codons on FlyBase 6.58 by local version of CAIcal with the *Drosophila melanogaster* codon usage table from the Kazusa codon usage database, and the isoform with minimum score (the most enriched for rare codons) is chosen to represent each expressed gene. Wilcoxon Rank Sum Test: ** = $P < 10^{-10}$; *** = $P < 10^{-15}$.

Figure 3. ORB2 binding motifs predicted from RIP-seq confer translational repression on a reporter mRNA in S2 cells. (a) The #ATS *de novo* motif discovery program (LI *et al.* 2010) was performed on the results from the anti-ORB2 RIP-seq in this study. #ATS identified NUUGUWU as the ORB2 recognition element (ORE) enriched in ORB2 targets. (b) The transcriptome was evaluated for accessible enrichment of the NUUGUWU motif and each

transcript was scored based on the number and accessibility of these motif sites. ORB2 targets
had significantly higher scores compared to non-targets, as assessed by Wilcoxon rank sum test.
*** = $P < 10^{-15}$ (c) Diagram depicting the reporter firefly luciferase mRNA with six copies of
different ORB2-binding motifs in the 3′UTR. (d) Bar graph depicting the enrichment of the
luciferase reporter mRNA in an anti-ORB2 RIP-qPCR compared to Renilla luciferase mRNA,
normalized to a reporter without ORE motifs. Mutation of the OREs or deletion of the RNA-
binding domain (RRM) abrogate binding to the mRNA. (e) Bar graph depicting the ratio of
luminescence between firefly luciferase and Renilla luciferase. Mutation of the OREs or deletion
of the RNA-binding domain (RRM) abrogate repression. One-way Kruskal-Wallis non-
parametric test with Dunn multiple test correction: * = $P < 0.05$.

Figure 4. The ORB2 and hCPEB2 ZZ domain but not the ORB and hCPEB1 ZZ domain
repress a luciferase reporter. (a) Diagram depicting the tethering assay in which a polypeptide
of interest fused to the BIV-Tat domain is tethered to a firefly luciferase reporter mRNA carrying
six tandem repeat TAR elements in the 3′UTR. (b) Bar graph depicting the expression of the
reporter firefly luciferase (normalized to co-transfected Renilla luciferase and normalized to
steady state RNA levels) when tethering whole ORB2 or ORB2 fragments compared to GFP
control. (c) Bar graph depicting the expression of the reporter firefly luciferase mRNA measured
by RT-qPCR (normalized to co-transfected Renilla luciferase mRNA) when tethering whole
ORB2 or ORB2 fragments compared to GFP control. Reporter mRNA levels are unaffected by
ORB2 ZZ or ORB2ΔZZ. (d) amino acid sequence alignment of the ZZ domains of ORB2, ORB,
and their human orthologs, hCPEB1 and hCPEB2. The arrowhead denotes the junction between
the RBD and ZZ domains. (e) Bar graph depicting the expression of the reporter firefly luciferase
(normalized to co-transfected Renilla luciferase) when tethering different CPEB ZZ domains.
Only ORB2 and hCPEB2 repress. (f) Bar graph depicting the expression of the reporter firefly
luciferase when ORB ZZ chimeric domains harboring substitutions from the ORB2 ZZ domain,
are tethered. One-way Kruskal-Wallis non-parametric test with Dunn multiple test correction: *
= $P < 0.05$.

Figure 5. Anti-ORB2 IP-MS from early embryo extracts identifies known translation
repressors and the PIC as binding partners. (a-b) Scatterplots depicting all proteins identified
in the IP-MS performed in the presence of RNase. Enrichment is expressed as the Log_2 ratio of
the average spectral counts detected in the ORB2 IP compared to the average spectral counts
detected in the control IP. Protein interactors were identified by SAINT (with a SAINT score ≥
0.95 and BFDR < 0.01), shown in red. (a) IP-MS performed on embryos laid by w^{1118} females,
(b) IP-MS performed on embryos laid by $orb2^{ΔZZ}$ females. (c) Interaction network depicting the
connections between components of the Cup repressive complex and other translational
repressors identified as ORB2 interactors by SAINT analysis. (d) Interaction network depicting
the connections between components of eukaryotic PIC identified as ORB2 interactors by
SAINT analysis. Interaction networks were produced by the GeneMANIA app for Cytoscape
(FRANZ *et al.* 2018); size of node indicates average number of spectral counts obtained in the
ORB2 IP-MS; thickness of edges indicates strength of interaction assigned by GeneMANIA.

Figure 6. Deletion of the ORB2 ZZ domain results in derepression of target RNAs in the
early embryo. (a) Violin and boxplots comparing target and non-target abundance as determined
by RNA-Seq in 0-1, 1-2 and 2-3 hour old embryos laid by w^{1118} and $orb2^{ΔZZ}$ females. No
significant difference was detected for either genotype at any of the time-points. (b) Violin plots

and boxplots comparing target and non-target translation index as determined by polysome
gradients followed by RNA-Seq in 0-1, 1-2 and 2-3 hour old embryos laid by w^{1118} and $orb2^{\Delta ZZ}$
females. ORB2 targets were significantly derepressed in 0-1 hour old $orb2^{\Delta ZZ}$ embryos relative to
w^{1118}. Log2(Ratio of mean normalized counts of pool 3/mean normalized counts of pool 1) was
calculated for each transcript. Wilcoxon Rank Sum Test: ** = $P < 0.01$.

Figure 7. ORB2 targets which are co-bound by SMG are not repressed in the early embryo.
(a-d) Violin and boxplots comparing the polysome association of ORB2 and SMG co-targets (a),
ORB2 targets (b), SMG targets (c), and non-targets (d), as determined by polysome gradients
followed by RNA-Seq in 0-1 hour old embryos laid by w^{1118} control (WT) and $orb2^{\Delta ZZ}$ females.
Wilcoxon Rank Sum Test ** = $P < 0.01$.

1420

Table Legends

Table 1. Overlap between the list of ORB2 targets and lists of genes associated with mRNA
stability or decay in the early embryo.

Table 2. Molecular Function GO-terms enriched in ORB2 targets. GO-term analysis was
performed using the gProfiler2 tool (KOLBERG *et al.* 2023).

Table 3. Biological Process GO-terms enriched in ORB2 targets. GO-term analysis was
performed using the gProfiler2 tool (KOLBERG *et al.* 2023).

Table 4. Cellular Component GO-terms enriched in ORB2 targets. GO-term analysis was
performed using the gProfiler2 tool (KOLBERG *et al.* 2023).

Table 5. Overlap of ORB2 targets with those bound by SMG, BRAT, PUM and RIN.

1431

Supplemental Figure Legends

Figure S1. CRISPR Cas9-mediated endogenous deletion of the ORB2 ZZ domain was used
to generate $orb2^{\Delta ZZ}$ flies. (a) Diagram depicting the *orb2* endogenous locus, located at 66E4-
66E5 on 3L, the corresponding mRNAs that are encoded, and the targeted deletion. (b) Agarose
gel of the PCR amplification of the region surrounding the CRISPR targeted deletion of the F1
generation post-injection. Genotypes indicate lines generated from individual F1 flies. Up PCR
indicates PCR amplification using primers designed against the 5′ insertion junction; Down PCR
indicates PCR amplification using primers designed against the 3′ insertion junction. (c) Agarose
gel of the PCR amplification of the region surrounding the PiggyBac transposon-mediated
excision of the dsRed marker. WT indicates the band representing PCR amplification of the
endogenous ORB2 locus containing an intact ZZ domain; Excised indicates the band
representing amplification of the locus with the ZZ domain deletion without the dsRed marker.
Genotypes indicate lines generated from individual flies resulting from the CRISPR injection. (d)
Western blot showing expression of endogenous ORB2 and CRISPRed ORB2ΔZZ in
heterozygous and homozygous whole flies. ORB2ΔZZ is expressed at higher levels than full-

length ORB2 (see heterozygote for comparison). Generation of CRISPR mutants was performed
by WellGenetics as described in the Materials and Methods.

Figure S2. Sequence of the CRISPR Cas9-mediated endogenous deletion of the ORB2 ZZ
domain.

Figure S3. Replicate-to-replicate comparisons of transcript TPMs from RIP-seq
experiments show high correlation between replicates. (a)-(c) Pairwise comparisons of the
three biological replicates performed with the anti-ORB2 E8 Fab. The Pearson correlation
coefficient and its corresponding *P*-value were calculated and used to determine the degree of
similarity between each pair. (d) Heatmap of the Poisson distance between three biological RIP-
seq replicates performed with anti-ORB2 E8 Fab and three biological RIP-seq replicates
performed with control C1 Fab. The Poisson distance is calculated by the difference in Poisson
distribution between pairs of samples; a larger distance indicates a greater level of dissimilarity
between samples.

Figure S4. Evaluation of RNA Compete motif enrichment and expression levels of ORB2,
ORB2ΔRRM, and GFP in S2 cells. (a) The transcriptome was evaluated for accessible
enrichment of the KUUUKKK motif and each transcript was scored based on the number and
accessibility of these motif sites, as in Figure 3b. ORB2 targets had significantly higher scores
compared to non-targets, as assessed by Wilcoxon rank sum test. *** = $P < 10^{-15}$. (b) Each set of
three lanes corresponds to the ORE validation experiments in which expression vectors for
tagged proteins and reporter mRNA constructs containing various recognition motifs were
transfected into S2 cells, described in Figure 3d and 3e.

Figure S5. The firefly luciferase open reading frame used in the FLuc reporter is enriched
for rare codons. (a) Distribution of CAI scores, calculated for all annotated CDS longer than 80
codons on FlyBase 6.58 by local version of CAIcal with the *Drosophila melanogaster* codon
usage table from the Kazusa codon usage database, and the isoform with minimum score (the
most enriched for rare codons) is chosen to represent each expressed gene. (b) Distribution of
MELP scores, calculated for all annotated CDS longer than 80 codons on FlyBase 6.58 by
coRdon with ribosomal proteins as the references for codon optimization, and the isoform with
minimum score (the worst expressivity) is chosen to represent each expressed gene. In (a) and
(b) the scores for the firefly luciferase open reading frame are indicated by the dashed blue lines.

Figure S6. Expression of ORB2 full-length protein, portions of ORB2, and the ZZ domains
of ORB, ORB2, hPCPEB1 and 2 in S2 cells. (a) Western blot showing the expression of ORB2
full-length protein and protein portions corresponding to the experiments described in Figure 4b
and c. (b) Western blot showing the expression of ORB2B and ORB2BΔZZ to the experiments
described in Figure 4b and c. (c) Western blot showing the expression of ORB2 ZZ, hCPEB2
ZZ, ORB ZZ and hCPEB1 ZZ corresponding to the experiments described in Figure 4e.

Figure S7. Expression of chimeric ORB-ORB2 ZZ domain proteins in S2 cells. (a)
Schematic depicting the amino acid sequence alignment of the ZZ domains of ORB2, ORB, and
their human orthologs, hCPEB1, hCPEB2, hCPEB3, and hCPEB4. The arrowhead denotes the
junction between the RBD and ZZ domains. (b) Bar graph depicting the expression of the
reporter firefly luciferase when ORB ZZ chimeric domains harboring substitutions from the

ORB2 ZZ domain, are tethered. The ORB2 'A' and 'C' regions are each sufficient to convert the
ORB ZZ domain into a repressor. One-way Kruskal-Wallis non-parametric test with Dunn
multiple test correction was used to assess significance. (c) Western blot showing the expression
of ORB-ORB2 chimeric proteins corresponding to the substitution experiments which assessed
transformation of the non-repressing ORB ZZ domain into a translational repressor, as shown in
S7b above.

Figure S8. Structure of ZZ domains of CPEB family members. (a) The NMR structure of the
ZZ domain of human CPEB1 Protein Data Bank accession 2M13 (MERKEL *et al.* 2013) is
shown, along with the Alphafold 3 (ABRAMSON *et al.* 2024) predicted structures of the ZZ
domains of ORB, hCPEB2 and ORB2. The colors in the Alphafold 3 predictions represent the
predicted local distance difference test (pLDDT) scores, where higher numbers indicate greater
confidence in the prediction. (b) On the left is an overlap of the ORB and ORB2 predicted
structures with Region A (the Rd turn), Region B (the beta hairpin), Region C (the alpha helix),
and Region D (the beta sheet) of each protein indicated. The boxes on the right depict the most
confident structure within each region in two different orientations.

Figure S9. ORB2ΔZZ retains RNA-binding to its targets in embryos. (a) Western blot
showing the expression of ORB2 and ORB2ΔZZ protein over the first four hours of embryonic
development. (b) Bar graph depicting the enrichment of five ORB2 target mRNAs in RIP-RT-
qPCR from 0-3 hour old embryos laid by w^{1118} and *orb2*^{ΔZZ} females. The ORB2 RIPs used the
ORB2 E8 Fab; the control RIP used the C1 Fab. RNA levels were normalized to *RpL32* mRNA.
n = 3; error bars are standard deviation. Student's *t*-test *P*-values were not significant (n.s.; *P* >
0.05)

Figure S10. IP-MS identifies ORB2 protein interactors. (a,b,d,e) Scatterplots depicting all
proteins identified in the IP-MS. Log_2(average spectral counts) from the ORB2 IP is plotted
against the Log_2(average spectral counts) from the control IP. (a) IP-MS performed on embryos
laid by w^{1118} females in the presence of RNase (corresponding to Figure 5a), (b) IP-MS
performed on embryos laid by *orb2*^{ΔZZ} females in the presence of RNase (corresponding to
Figure 5b), (d) IP-MS performed on embryos laid by w^{1118} females in the absence of RNase, and
(e) IP-MS performed on embryos laid by *orb2*^{ΔZZ} females in the absence of RNase. (c, d)
Scatterplots depicting all proteins identified in the IP-MS performed in the absence of RNase.
Enrichment is expressed as the Log_2 ratio of the average spectral counts detected in the ORB2 IP
compared to the average spectral counts detected in the control IP. Protein interactors were
identified by SAINT (with a SAINT score ≥ 0.95 and BFDR < 0.01), shown in red.

Figure S11. ORB2 targets which are co-bound by PUM are not de-repressed in the early
embryo. (a-d) Violin plots and boxplots comparing the translation index of ORB2 and PUM co-
targets (a), ORB2 targets (b), PUM targets (c), and non-targets (d), as determined by polysome
gradients followed by RNA-Seq in 0-1 hour old embryos laid by w^{1118} and *orb2*^{ΔZZ} females.
Wilcoxon Rank Sum Test *P*-values are shown.
1526
1527 **Figure S12. ORB2 targets which are co-bound by BRAT are not de-repressed in the early**
1528 **embryo.** (a-d) Violin plots and boxplots comparing the translation index of ORB2 and BRAT
1529 co-targets (a), ORB2 targets (b), BRAT targets (c), and non-targets (d), as determined by

1530 polysome gradients followed by RNA-Seq in 0-1 hour old embryos laid by w^{1118} and $orb^{2\Delta ZZ}$
1531 females. Wilcoxon Rank Sum Test *P*-values are shown.
1532

Supplemental File Legends

File S1. Results of RIP-seq of ORB2 in 0-3 hour embryos. See Materials and Methods for
details.

File S2. Results of rare codon enrichment analyses on ORB2 targets in 0-3 hour embryos.
The MELP and CAI were calculated as described in Materials and Methods.

File S3. Results of *de novo* motif discovery on ORB2 targets in 0-3 hour embryos. Motif
discovery was carried out using #ATS as described (Lɪ *et al.* 2010). ORE scores and
'KUUUKKK' scores are listed. See Materials and Methods for details.

File S4. Results of IP-MS of ORB2 and ORB2ΔZZ in 0-3 hour embryos in the presence or
absence of RNase A. Interactors were identified using SAINT as described (Cʜᴏɪ *et al.* 2012).
See Materials and Methods for details.

File S5. Lists of mRNAs bound by ORB2 (RIP-seq) whose protein products were also
identified as interactors in ORB2 IP-MS.

File S6. Results of RNA-seq time-course in wild-type and ORB2ΔZZ embryos. See Materials
and Methods for details.

File S7. Results of polysome gradient RNA-seq time-course in wild-type and ORB2ΔZZ
embryos. See Materials and Methods for details.

File S8. Overlaps of ORB2 targets and previously identified SMG, BRAT and PUM targets.
See Materials and Methods for details.

FIGURE 1

a

b

c

d

FIGURE 2

FIGURE 3

a ORE derived from RIP-Seq

b ***

c m⁷G — Luciferase — □□□□□□ — AAA...
ORB2
Six tandem copies of ORB2 motif in 3'UTR

d RNA enrichment in ORB2 IP
☐ GFP ☐ ORB2 ☐ ORB2ΔRRM

e Luminescence of Luciferase reporter
☐ GFP ☐ ORB2 ☐ ORB2ΔRRM

FIGURE 4

FIGURE 5

a Proteins identified in IP-MS from *w¹¹¹⁸*

b Proteins identified in IP-MS from *orb2ᐃZZ*

FIGURE 6

FIGURE 7

TABLE 1

Comparison	Backgound (Total)	Orb2 Only	Other Only	Overlap	Fisher's Exact p-value	Odds Ratio
Maternal Unstable (Unfertilized eggs, Tadros et al 2007)	3030	172	744	166	2.20E-16	3.849738
Thomsen et al 2010						
Class I (Stable transcripts)	4186	377	1335	35	2.20E-16	0.2334165
Class II (Maternal Decay only)	5108	395	394	36	0.3359	1.181531
Class III (Maternal Decay + Transcription)	5177	392	326	38	0.02109	1.539281
Class IV (Zygotic Decay only)	4332	322	1171	108	0.06763	1.240746
Class V (Both Maternal + Zygotic Decay)	4186	239	1317	191	2.20E-16	2.53962

TABLE 2. Molecular Function

GO Term Name	Adjusted p-value	GO Term Size	ORB2 targets	Overlap
protein binding	1.13E-21	3760	416	229
mRNA 3'-UTR binding	5.78E-11	48	416	17
mRNA binding	1.15E-10	238	416	35
mRNA regulatory element binding translation repressor activity	2.51E-08	18	416	10
translation repressor activity	7.18E-08	25	416	11
translation regulator activity	2.28E-04	106	416	16
catalytic activity, acting on DNA	2.49E-04	147	416	19
protein serine/threonine kinase activity	5.25E-04	184	416	21
DNA binding	6.42E-04	1127	416	69
poly-pyrimidine tract binding	1.28E-03	14	416	6
adenyl ribonucleotide binding	1.42E-03	846	416	55
catalytic activity, acting on a nucleic acid	1.62E-03	406	416	33
ATP binding	2.44E-03	840	416	54
purine ribonucleotide binding	2.99E-03	1021	416	62
purine nucleotide binding	3.63E-03	1072	416	64
adenyl nucleotide binding	3.73E-03	896	416	56
ribonucleotide binding	4.31E-03	1033	416	62

TABLE 3. Biological Process

GO Term Name	adjusted p-value	GO Term Size	ORB2 targets	Overlap
purine ribonucleoside triphosphate binding	4.48E-03	1012	416	61
anatomical structure development	5.77E-28	2775	435	212
developmental process	2.30E-27	2954	435	219
negative regulation of cellular process	1.73E-25	1520	435	144
cell development	6.92E-25	1450	435	139
multicellular organism development	8.26E-25	2276	435	182
cell differentiation	3.46E-24	1768	435	155
cellular component organization	1.07E-21	3178	435	217
regulation of metabolic process	1.81E-21	2253	435	174
gamete generation	1.19E-20	917	435	100
germ cell development	7.67E-20	721	435	86
anatomical structure morphogenesis	1.34E-19	1580	435	136
oogenesis	1.45E-19	582	435	76
system development	5.09E-19	1527	435	132
multicellular organismal reproductive process	1.60E-18	1096	435	107
regulation of primary metabolic process	1.69E-17	1860	435	146
nervous system development	4.18E-17	1196	435	110
animal organ development	5.43E-17	1200	435	110
regulation of macromolecule biosynthetic process	1.09E-15	1742	435	136

TABLE 4. Cellular Component

GO Term Name	adjusted p-value	GO Term Size	ORB2 targets	Overlap
nucleus	1.99E-14	3234	438	196
ribonucleoprotein granule	2.57E-12	119	438	27
cytoplasmic ribonucleoprotein granule	7.11E-12	114	438	26
cellular anatomical structure	6.48E-10	10708	438	426
cytoplasm	6.02E-09	5187	438	255
P granule	1.14E-07	46	438	14
cytosol	1.14E-07	1131	438	82
germ plasm	2.15E-07	48	438	14
nucleoplasm	1.40E-06	573	438	50
pole plasm	2.54E-06	57	438	14
nuclear lumen	3.31E-06	910	438	67
cytoplasmic stress granule	5.07E-06	21	438	9
protein-containing complex	2.53E-05	2861	438	151
chromosome	3.50E-05	787	438	58
PcG protein complex	8.68E-04	27	438	8
transferase complex	2.13E-03	550	438	41
intracellular protein-containing complex	4.15E-03	625	438	44
cell junction	4.67E-03	608	438	43
intracellular organelle lumen	4.94E-03	1158	438	69
membrane-enclosed lumen	5.08E-03	1159	438	69
organelle lumen	5.08E-03	1159	438	69
cell cortex	5.36E-03	208	438	21

TABLE 5

Comparison	Backgound	Orb2 Only	Other Only	Overlap	Fisher's Exact p-value	Odds Ratio
Smaug (Chen et al 2012)	3758	254	290	41	0.0001416	2.09148
Brat (Laver et al 2015)	3347	228	701	69	0.02181	1.40294
Pum (Laver et al 2015)	3696	238	352	57	6.197E-08	2.51398
Rin (Laver et al 2020)	4429	365	507	6	2.15E-10	0.14363