Allosteric effects of the coupling cation in melibiose transporter MelB

Parameswaran Hariharan[1], Yuqi Shi[2], Rosa Viner[2], and Lan Guan[1§]

[1]Department of Cell Physiology and Molecular Biophysics, Center for Membrane Protein

Research, School of Medicine, Texas Tech University Health Sciences Center, Lubbock, TX,

USA

[2]Thermo Fisher Scientific, San Jose, CA, USA

Running title:

Allostery of Na^+ in MelB

[§]To whom correspondence should be addressed:

Lan Guan, E-mail: Lan.Guan@ttuhsc.edu

Keywords:

Allostery, symport, structural dynamics, sugar recognition, x-ray crystallography, HDX-MS

Abbreviations:

MelB$_{St}$, *Salmonella enterica* serovar Typhimurium melibiose permease

MFS, major facilitator superfamily

ITC, isothermal titration calorimetry

α-NPG, 4-nitrophenyl-α-D-galactopyranoside

Mel, melibiose

α-MG, α-methyl galactoside

Raff, raffinose

K_d, dissociation constant

HDX-MS, hydrogen-deuterium exchange coupled to mass spectrometry

D, deuterium

Abstract

The major facilitator superfamily (MFS) transporters play significant roles in human health and disease. *Salmonella enterica* serovar Typhimurium melibiose permease (MelB$_{St}$), which catalyzes the symport of galactosides with Na$^+$, H$^+$, or Li$^+$, is a prototype of this important transporter superfamilies. We have published the structures of the inward- and outward-facing conformations of MelB$_{St}$ with galactoside or Na$^+$ bound, and determined the binding thermodynamic cycle. We have proposed that positive cooperativity between the two co-transported solutes plays a key role in the symport mechanism of MelB$_{St}$; however, the molecular basis for this core mechanism remains unclear. In this study, we investigated the structural dynamics induced by melibiose, Na$^+$, or both on MelB$_{St}$ using hydrogen-deuterium exchange mass spectrometry (HDX-MS). We also refined the specific determinants for the sugar recognition in both protein and galactoside molecules by solving the crystal structures of D59C MelB$_{St}$ bound to melibiose and two other sugars that contain different numbers of sugar units, and identified a critical water molecule as part of the specific determinants from a α-NPG-bound structure. Our integrated structural and HDX-MS analyses support the notion that the binding of the coupling cation at a remote site stabilizes those dynamic sidechains in the sugar-binding pocket, leading to a high-affinity state. This study provides the molecular basis for the essential symport mechanism through positive cooperativity, which may serve as a general mechanism for most cation-coupled symporters.

Introduction

The Solute Carrier (SLC) family of transporters encompasses diverse superfamilies of membrane proteins with various protein folds and employing different transport mechanisms to facilitate the translocation of a wide range of solutes across cell membranes (1). The largest superfamily of SLC transporters is the major facilitator superfamily (MFS)(2). MFS transporters comprise a significant number of cation-coupled secondary active transporters and are responsible for the uptake of a broad spectrum of solutes across cell membranes, playing crucial roles in physiology, pathology, and pharmacokinetics, and are emerging as drug targets (3, 4). Recent rapid advancements in membrane protein research have greatly enhanced our understanding of protein conformation and mechanisms(5-8); however, critical details remain lacking. For example, in cation-coupled symport, it is still unclear how the coupling cations help in the binding, translocating, and accumulating the primary substrate.

For MFS secondary active transporters, most members use H^+ as the coupling cation, and a few members use Na^+, such as the Na^+-coupled lipid transporter (MFSD2A)(9, 10) that is expressed in the major organ barriers, including the blood-brain barrier or blood-retina barrier. MFSD2A plays a critical role in the uptake of essential lipids into these neural tissues (9-11). The Na^+-coupled melibiose transporter of *Salmonella enterica* serovar Typhimurium (MelB$_{St}$), which is a well-characterized representative for the Na^+-coupled MFS transporters, catalyzes the symport of a galactopyranoside with Na^+, H^+, or Li^+, and is a useful model system for studying cation-coupled transport mechanisms (12-22). Two major conformations of MelB$_{St}$ have been determined: one is an outward-facing conformation at apo or galactoside-bound states(20, 22), and the other is an inward-facing conformation at a Na^+-bound state(21). The primary substrate-specificity determinant pocket and the cation-specificity determinant pocket have been

structurally and functionally characterized (19-23). Binding at both pockets exhibits a positive

cooperativity; that is, the binding of one increases the affinity for the other (19). All three

coupling cations compete for the same binding pocket, and the transport stoichiometry is 1

galactoside: 1 cation (Na^+, H^+, or Li^+). The sugar affinity depends on the cation identity, with the

cooperativity numbers (fold of increase in affinity) being 8, 5, or 2 for Na^+, Li^+, and H^+,

respectively (19, 20). Notably, the sugar-binding affinity is also dependent on MelB$_{St}$

conformation (21), in addition to being sensitive to the binding of the coupling cation and its

identity. By trapping MelB$_{St}$ in an inward-facing state using the inward-facing conformation-

specific binder nanobody-725 (Nb725), both experimental sugar-binding assays and cryoEM

structural analysis support a low-affinity state of the sugar-binding pocket (21, 23). Remarkably,

Na^+ binding to the inward-facing conformation remains unchanged (21, 23). These results

provide experimental evidence supporting the previously proposed stepped-binding kinetic

model for melibiose/Na^+ symport, in which Na^+ binds first and is released after the sugar release

on the opposite surface (15, 18, 24, 25).

Positive cooperativity is proposed to be the key symport mechanism in MelB$_{St}$, but the

molecular basis for this critical mechanism remains unclear. The structures indicate that the

bound sugar and Na^+ have no direct contact, while the two binding pockets are in close

proximity (20, 21). The minimum free-energy landscape for sugar translocation has been

simulated based on the structures of the outward- and inward-facing conformations as two

starting points (26). The results suggest that the Na^+ contribution to the binding free energy of

sugar by direct contact is negligible. The cooperativity between melibiose and Na^+ occurs

through allosteric coupling, primarily via electrostatic effects. To experimentally evaluate this

concept, in this study, we enhanced the resolution of the structure with α-NPG bound, which

revealed a critical water molecule, and also determined crystal structures with three other α-

sugars (melibiose, raffinose, or α-MG) that contain different numbers of sugar units to verify the

sugar-recognition specific determinants. We also analyzed the structural dynamics with HDX-

MS induced by melibiose, Na^+, or both on MelB$_{St}$. The extensive integrated analyses of the

structure and structural dynamics by MD simulations and HDX-MS support the conclusion that

the binding of the coupling cation at a remote location stabilizes the sugar-binding residues to

switch to a higher-affinity state. Therefore, the coupling cation in this symporter is an allosteric

activator.

Results

Outward-facing structures of MelB complexed with varied sugar substrates

We have published the crystal structures of MelB$_{St}$ with bound sugar analogues α-NPG or

melibiose-derived detergent (DDMB), from a MelB$_{St}$ uniport mutant D59C (20), or a solely H$^+$-

coupled symporter mutant D55C(22). Here we report four crystal structures of D59C MelB$_{St}$

with the bound endogenous sugar substrate melibiose, and two other α-galactosides, α-methyl

galactoside (α-MG) with a single sugar unit or raffinose with three sugar units, as well as α-NPG

at an improved resolution. The statistics of the four structures are presented in **Table S1**. All

outward-facing structures are virtually identical with RMSD values of less than 0.4 (**Fig. S1**).

This typical MFS-fold transporter has 12 transmembrane helices and is organized into two six-

helix domains linked by the middle loop between helices VI and VII (Loop$_{6-7}$). There are two

cytoplasmic helices on the middle loop or the C-terminal tail or lid (**Fig. 1**). In all structures, one

sugar molecule is bound in the middle of the protein and sandwiched by the helices from both

the N- and C-terminal domains. As indicated by the sliced surface, the binding residues located

within the cytoplasmic leaflet of both domains shape the edge of the inner barrier that prevents

the sugar from passing across the transporter into the cytoplasm. On the periplasmic side, there is

an open vestibule connecting the solvent to the binding pocket.

α-NPG binding refined to a resolution of 2.7 Å. The improved resolution from 3.01 Å to 2.7

Å provides a better-resolved density map for the bound α-NPG molecule, which further supports

the originally assigned pose (**Fig. 2**) (20). As described, the binding pocket is formed by 14

residues on five helices, including N-terminal helices I (Lys18, Asp19, Ile22, and Tyr26), IV

(Tyr128, Asp124, Tyr120), and V (Arg149 and Ala152), and the C-terminal helices X (Trp342)

and XI (Gln372, Thr373, Val376, and Lys377).

A water molecule was modeled to a positive density, which located at a good hydrogen-

bonding distances from the C4-OH on the galactopyranosyl ring and Thr733 on helix XI, and

also surrounded by C6-OH, Gln372 on helix XI and Asp124 on helix IV (**Fig. 2**). Overall, the

galactosyl moiety is primarily stabilized by the N-terminal charged residues Asp19, Asp124, and

Arg149. Especially, strong and continuous density between Asp19 and the C2-OH and C3-OH

and Asp124 with C4-OH were observed in the previous maps (20) and current maps, indicating

that both negatively charged Asp resides and the water molecule play critical roles in the binding.

Five more water molecules were modeled, and three interact with the previously identified

charge network involved in 9 charged residues between the N- and C-terminal domains (**Fig. S2a,**

b). The water 2 and 3 interact with the charged pair Arg295 (helix IX) and Glu351 (helix XI)

and the water 4 interacts with Arg138 on helix IV, which also form a salt bridge with Glu142 to

stabilize the hydrogen-bonding interaction between Glu142 and Arg295 (**Fig. S2b**).

Melibiose binding. The endogenous substrate melibiose is formed from a galactose unit and a

glucose unit linked by an α-1,6 galactosyl bond (D-Gal-(α1→6)-D-Glc). The melibiose-bound

D59C MelB$_{St}$ structure was refined to a resolution of 3.05 Å, and the density map displayed a

clear two-unit blob in the binding pocket, fitting well with one molecule of melibiose. While the

water molecule was not modeled due to limited resolution, a positive density is also present at

W1 position. Two waters at W2 and W3 positions were also modeled in this structure.

Interestingly, the binding affinity between melibiose and α-NPG differs by a factor of 100; the

only difference in the binding sites is the pose and chemistry of the non-galactosyl moiety (**Fig.**

3a, e). In α-NPG, the hydrophobic phenyl group, positioned at 4 Å-distance from the phenyl

group of Tyr26, forms a strong stacking interaction. While melibiose has four additional OH

groups, there is no further polar interaction with MelB. This glucosyl ring rotates away and

becomes nearly perpendicular to the phenyl ring of Tyr26 **(Fig. S3)**. The absence of aromatic

stacking may contribute to the relatively poor affinity of melibiose.

α-Methyl galactoside binding. This sugar molecule has a methyl substituent at the anomeric

C1 position in an α-linkage. The α-MG-bound structure of the D59C MelB$_{St}$ mutant was refined

to a resolution of 3.68 Å, and the density map still clearly displayed a one-unit blob in the

binding pocket, where one α-MG was modeled similarly to the galactosyl moiety of melibiose

(Fig. 3b).

Raffinose binding. Raffinose is a trisaccharide formed from melibiose and fructose. The

raffinose-bound D59C MelB$_{St}$ mutant structure was refined to a resolution of 3.40 Å, and the

density map displayed a clear three-unit blob in the binding pocket, fitting well with this

trisaccharide. As expected, the raffinose-binding pocket is significantly larger than that of

melibiose and α-NPG **(Fig. 3c)**, involving two additional helices. Consequently, a set of polar

side chains on helix VII (Asn244, Ser247, Asn248, and Asn251), helix VIII (Asn279), and helix

V (Ser153) were found to be within 4-5 Å of either the glucosyl or fructosyl moieties.

Interestingly, the polar interactions at hydrogen-bonding distances between raffinose and

MelB$_{St}$ are nearly identical to those in α-MG, melibiose, or α-NPG, with most

occurring within the galactosyl moiety, whiles the Asp19 is at a hydrogen-bonding distance from

the OH-4 on the glucosyl moiety.

All four galactosyl moieties align well in the specificity-determinant pocket, regardless of the

number of monosaccharide units **(Fig. 3d, e; Fig. S1)**. This alignment strongly supports the

previous conclusion that the galactosyl moiety determines the specificity of the primary

substrates; however, while the non-galactosyl moiety contributes to the binding affinity. The

galactosyl-binding pocket is well formed with little room for flexibility in the sugar pose;

however, the spacious cavity can accommodate at least one more fructose unit, as shown in the

raffinose.

Isothermal titration calorimetry (ITC) measurement of the binding affinity

Binding affinity K_d values of the WT and D59C MelB$_{St}$ for melibiose of 1.25 mM ± 0.05

mM and 4.96 ± 0.11 mM, respectively, and for α-NPG of 16.46 ± 0.21 μM or 11.97 ± 0.09 μM,

respectively, have been determined by ITC (19, 20, 25, 27). To determine the binding affinity of

MelB$_{St}$ for α-MG and raffinose, ITC measurements were also utilized to determine the K_d values

in the presence of Na$^+$ (**Fig. 4**). By injecting the raffinose or α-MG in a buffer-matched solution

into the sample cell containing the WT MelB$_{St}$ in the presence of Na$^+$, the isotherm curve fitted

well, yielding K_d values of 0.88 ± 0.04 mM or 3.79 ± 0.53 mM or for α-MG or raffinose,

respectively. It is surprising that raffinose, which has more sugar units, exhibits a poor binding

affinity to MelB$_{St}$.

Structural dynamics measured by HDX-MS

To determine the structural dynamics of MelB$_{St}$ in response to the substrate and/or Na$^+$, the

bottom-up HDX-MS method was employed to measure the exchange rates of amide hydrogen

with deuterium by analyzing time-dependent mass spectra at the peptide level as described(28-

30). Three pairs of datasets were collected, including Apo vs. melibiose, Na$^+$, or both, as outlined

in the Methods section. The analyzed region of MelB$_{St}$ spans positions 2-470 out of 476,

achieving a coverage of over 87.47% and 86.62% from 149 or 152 overlapping deuterium-

labeled peptides (**Table S2; Fig. S4**). There are 59, 63, or 59 non-covered residues for the

datasets of Apo vs. melibiose-bound, Na$^+$-bound, and melibiose- and Na$^+$-bound states,

respectively. Most of these positions are localized within the transmembrane regions, particularly

on the C-terminal half (**Table S4**). Interestingly, the two longest helices, IV and X, display a

significantly higher number of non-covered positions compared to the other helices. While a

significant number of such positions are present on the cytoplasmic loops, all periplasmic loops

are covered by labeled peptides.

The HDX rates are varying in different regions (**Table S4**). When averaging the related

deuterium uptake (D%) from all peptides at all time points at a given region, the N-terminal

helices I and V, as well as the C-terminal helices VIII and IX exhibited higher values, indicating

they are more dynamic regions of Apo MelB$_{St}$.

The differential deuterium labeling (ΔD) was calculated based on the deuterium uptake in the

absence (Apo state) or presence of specific ligand(s) (Holo state), and is presented as the residual

plot of each peptide between the two states ($_{Holo - Apo}$) (**Fig. 5a**) at three labeling time points and

the sum of all. The dashed lines indicate the global thresholds (ΔD) for each dataset. The C-

terminal His tag and tail residues at positions 470-475 are highly dynamic and fully labeled with

greater variations and are excluded from the analysis. The residual plots include data from all

peptides, and the ΔD values greater than the global threshold and P values <0.05 are

considered statistically significant. Melibiose binding induced a wide range of structural effects

on MelB$_{St}$ throughout the entire MelB$_{St}$ polypeptide, and the effects include protection (less

deuterium uptake in the Holo state) and deprotection (greater deuterium uptake in the Holo state).

In contrast, Na$^+$ alone or in combination with melibiose primarily caused deprotection. As shown

in **Table S3**, more than 50% of residues (237, 264, or 257 positions) exhibited either

insignificant ΔD values ($D_{Mel - Apo} < |0.186|$, $\Delta D_{(Na+)}$ - Apo $< |0.224|$, or $\Delta D_{Na(+)Mel}$ - Apo $< |0.175|$)

or greater P values (> 0.05). A peptide with at least one significant ΔD or overlapping peptides

extending from the peptides with at least one significant ΔD were mapped onto the melibiose-

bound structure of each dataset individually (**Fig. S5**). The uncovered positions were indicated in

gray spheres, and the positions with insignificant ΔD values are shown in backbone

representation. Peptides with significant ΔD from all three datasets are presented together in

Figure 5b.

Consensus effects are observed from all three conditions. Interestingly, peptides with

significant ΔD values are clustered in several regions. Nearly full-length transmembrane helices I

and V, as well as the tail helix, demonstrated protections from melibiose, Na^+, or

Na^+ with melibiose. In particular, the peptides containing sugar-binding critical residues (Lys18,

Asp19, and Arg149 on helices I and V) showed statistically high confidence ($P < 0.01$, **Fig. 5b,**

Blue). The peptide 213-226 in the middle loop across the three conditions was also protected.

Additionally, the peptide 364-374 on helix XI, which contains sugar-binding residues Gln372

and Thr373, was protected by Na^+ alone ($P = 0.029$) or Na^+ with melibiose ($P = 0.011$) and was

weakly protected by melibiose but at a P value of 0.056.

In the presence of melibiose and Na^+, the deprotections of overlapping peptides 36-47

(loop1-2) and 284-293 (loop8-9) with high confidence ($P < 0.01$) were obtained. Melibiose alone

caused the deprotections of peptide 36-47 with medium confidence ($P < 0.05$) (**Fig. 5b, red**).

Five deprotected regions induced solely by melibiose are found at the ends of transmembrane

helices with extended loops (**Fig. 5b, pink**). Overlapping peptide 292-298 (**Fig. 5, Cyan**)

showed opposite effects when comparing melibiose alone to melibiose with Na^+.

Allosteric effects of Na^+ binding on sugar-binding residues.

The helices II and IV, which host the coupling cations, were shown to have significantly

lower deuterium uptake (**Fig. 5b; Fig. S5**) compared to helices I and V, which are specific for

sugar binding and do not participate in cation binding. The effects of deuterium uptake by

melibiose, Na^+, or melibiose plus Na^+ on the residues in the sugar- or Na^+-binding pockets are

summarized in **Table S5**. Seven peptides covering all N-terminal residues for sugar and cation

binding were presented with the time courses of deuterium uptake. Notably, peptides 53-62, 120-

123, and 121-130, which contain all cation-binding positions (Asp55, Asn58, Asp59, and

Thr121), exhibited poor deuterium uptake rates, and their uptake rates were not significantly

affected by the binding of melibiose, Na^+, or both (**Fig. 6, labeled in black (II) and blue (IV),**

respectively).

On the contrary, the two peptides (16-20 and 21-27) covering the sugar-binding residues 18,

19, 22, and 26 on helix I exhibited strong protection by melibiose, both in the absence and

presence of Na^+ (**Fig. 6, magenta label)**. Remarkably, both peptides on helix I are also protected

by Na^+, which binds at helices II and IV, remotely. Additionally, the sugar-binding residue

Arg149 on helix V was covered by several overlapping peptides that consistently

demonstrated strong protections by melibiose, Na^+, or melibiose plus Na^+, as illustrated

by peptide 140-151 (**Fig. 6, red label)**. The residue 152, located in the binding pocket but not in

direct contact with sugar, showed no detectable protection by melibiose or Na^+ alone; however,

significant protection was observed in the presence of both melibiose and Na^+. The protective

effects on the C-terminal sugar-binding residues Gln372 and Thr373 have been described; the

peptide 369-374 is protected by Na^+ alone and stronger protection was observed in the presence

of melibiose plus Na^+ (**Fig. 5b**).

Discussion

The dehydration of sugar molecules is required for sugar binding, yet little is understood

about this process in MelB. In the 2.7 Å-resolution α-NPG-bound structure, a water molecule

was modeled to a positive density, and surrounded by the OH-4 and OH-6 groups on the

galactopyranosyl ring, as well as by Gln372 and Thr373 on helix XI. In the published MelB$_{St}$

structures with α-NPG or DDMB bound (20, 22) and the current structures with native melibiose,

α-MG, or raffinose bound, due to the limited resolutions, the water molecule was not modeled;

however, the positive density at the same position always appeared in the mFo-DFc maps, which

indicate that this OH-4 water molecule is part of the sugar binding coordination in MelB.

The structures complexed with galactosides containing different numbers of sugar units,

ranging from one to three, clearly demonstrate that the specificity is directed toward the

galactosyl moiety (**Figs. 1-3**). The sugar-protein polar interactions primarily occur at positions 2,

3, 4, and 6. Notably, the orientation of the hydroxyl group at the C4 position is the crucial feature

for distinguishing between galactose and glucose. The OH-4 on α-NPG is within hydrogen-

bonding distances from the carboxyl group of Asp124 and the indole group of Trp128, and it

indirectly connects with Gln372 and Thr373 on helix XI through this water molecule bound to

OH-4 (**Fig. 2**). These interactions define the specificity of galactosyl molecules, as the glucose or

glucoside OH-4 points to the opposite side of the galactosyl plane; thus, it does not form the

hydrogen-bonding network around OH-4 and therefore would not be selected by MelB$_{St}$. Clearly,

the water molecule bound to OH-4 contributes to the sugar recognition.

Another important piece of structural information is that the OH-4 group and the bound water

are in close proximity to the Na$^+$-binding residue Thr121 and the cation-binding important

residue Lys377, indicating that the OH-4 water molecule appears on the pathway between the

two specificity-determining pockets. The OH-3 and OH-2 groups on the opposite edge of the

galactopyranosyl ring create multiple hydrogen-bonding interactions with the charged residues

Asp19 and Arg149. These interactions likely help to stabilize recognition at OH-4 and enhance

the binding affinity.

All available structures of MelB$_{St}$ (19-21) and MD simulations on the coupled translocation

of melibiose and Na$^+$ (26) indicate that there is no direct interaction between the bound Na$^+$ and

the bound galactosides. Binding affinity determinations of the primary substrate and the coupling

cations demonstrate their positive cooperativity (19, 20, 27). The MD simulations suggested that

Na$^+$ binding allosterically affects the local electrostatic interactions in the sugar binding pocket.

HDX-MS was used to obtain experimental evidence for the structural dynamics induced by

melibiose, Na$^+$, or melibiose combined with Na$^+$.

HDX determines the structural motion or dynamics by measuring exchange rate over

time(28). The resolution of this method is limited to the peptide level, and the deuterium uptake

rate is affected by the interplay of structural dynamics and solvent accessibility. Previously, we

used this technique to study the conformational dynamics between inward- and outward-facing

states of MelB$_{St}$ with bound Na$^+$. In this study, we tested the Apo state and Holo states with

different ligands. At Apo state, the peptides with faster deuterium uptake rates are distributed in

both N- and C-terminal domains, and helices I and V are the three helices with the fastest uptake

rates, indicating those regions at Apo state are more dynamic. The previous thermal denaturation

study detected by circular dichroism spectroscopy showed that the stability of the Apo state is

lower than that of any holo states, either with melibiose or Na$^+$; the most stable state is the

ternary form with both melibiose and Na$^+$ bound(27). This HDX study identified the dynamic

regions that contribute to the poor stability of Apo MelB$_{St}$.

There is a significant difference in the dynamics of the sugar-binding helices (I, IV, V, X, and XI) and the cation-binding helices (II and IV). The cation-binding helix II (containing Asp55, Asn58 and Asp59) exhibits poor deuterium uptake and helix IV (containing Thr121) contains more non-covered positions either not covered by labelling or not covered by peptides. In either condition, their solvent accessibility and H/D exchange should be expected to be low. Therefore, the areas hosting the cation binding are relatively rigid (**Figs. 5-6**). In contrary, most sugar-binding helices, e.g., helices I (containing Lys18, Asp19, Ile22 and Try126), V (Arg149 and Ala152) and XI (containing Gln372 and Thr373), are dynamics; however, helix IV (containing Tyr120, Asp124, and Try128) that also carries the Na^+-binding residue Thr121 is rigid. Notably, Asp19 and Arg149, which contribute to binding affinity, are located in the most dynamic helices I and V. Stabilization of both helices is expected to increase the affinity for sugar.

Melibiose-induced structural changes include dynamic restraints and enhancements (**Fig. S5a**). The structural restraints or protections were observed on the binding residue-hosting helices (I and V) and two loop helices (the middle loop and the C-terminal tail), while the deprotections affect a wide range of areas, which are mapped on the helices-loop regions and most are at the periplasmic side. Interestingly, most regions protected by melibiose are also protected by Na^+ alone; in addition, the water-mediated sugar-binding residues Gln372 and Thr373 are also protected (**Fig. S5b**). There is no significant deprotection detected by Na^+, consisting with the conclusion that the Na^+-bound MelB is more stable than the Apo state.

When both melibiose and Na^+ are presented during the H/D exchange reaction, most of the protected regions by melibiose or Na^+ are also protected (**Fig. S5c**). Two deprotected regions were identified. One is also deprotected by melibiose, which is the peptide 36-47 located in $Loop_{1-2}$ and the beginning of the helix II; another deprotected region 292-298 is on the extended

helix VIII. Both regions become more dynamic in the ternary state with both melibiose and Na^+.

Notably, the deprotected region 292-298 contains Arg295, and two water molecules are modeled

in the 2.70 Å map which stabilize the salt-bridge Arg295 and Asp351 on helix X, which is part

of the large comprehensive charged network to stabilize the packing between the N- and C-

halves at the cytoplasmic leaflet region for the inward-facing state. The binding of both prepared

the transporter to initiate the conformational transition from the outward to the inward-facing

states. Arg295 might respond to this binding signal and become more dynamics to destabilize the

charged network and facilitate the opening at the cytoplasmic surface.

The MD simulation and free energy calculations and distance distribution indicated that the

Asp19 is highly dynamic when Na^+ unbound, and the binding of Na^+ shifted the weak interaction

of melibiose to stronger single interaction(26). HDX-MS data showed that the binding of Na^+

alone causes the protections of the dynamic regions that carry Asp19 and Arg149, so the sugar-

binding residues near OH-3 and OH-2 groups are more rigid. Therefore, it can be concluded that

Na^+ binding rigidifies the sugar-binding pocket through an allosteric effect on the local

electrostatic interactions as proposed from the MD simulations.

The symporter H^+-coupled lactose permease LacY and MelB share several common features

(5, 31, 32). Both transporters share a similar substrate spectrum, and all are selective for

galactosides. In addition, the cation-binding affinity in both LacY and MelB does not appear to

be influenced by conformational transition(21, 32, 33). Furthermore, their cation sites do not

have direct contact with their primary substrate galactosides. Glu325 of LacY is the sole

protonation residue (34) and located in a hydrophobic milieu with no direct contact with

galactosides(31, 33, 35). It has also been demonstrated that the sugar-binding affinity depends on

the protonation status of LacY (36), which could be postulated to be an allosteric effect. The

role of the coupling cation as an allosteric activator in cation-coupled symporters might be a

common mechanism. Understanding this could provide broader insights into how these transport

systems operate.

Experimental Procedures

Reagents. Melibiose was purchased from *Acros Organics* (*ThermoFisher Scientific*). Raffinose (RPI Chemicals), α-methyl galactoside andα-nitrophenyl galactoside are purchased from Sigma-Aldrich. Detergents undecyl-β-D-maltopyranoside (UDM) and DDM (dodecyl-β-D-maltopyranoside (DDM) were purchased from *Anatrace*. *E. coli* lipids (Extract Polar, 100600) were purchased from *Avanti Polar Lipids, Inc*. All other materials were reagent grade and obtained from commercial sources.

Strains and plasmids. *E. coli* DW2 cells (*mel*A$^+$, *melB$^-$, lacZ'Y*) (37) were used for protein expression and functional studies. The expression plasmids pK95/ΔAH/WT MelB$_{St}$/CHis$_{10}$(15) and pK95/ΔAH/D59CMelB$_{St}$/CHis$_{10}$ (18) were used for constitutive expression.

MelB$_{St}$ protein expression and purification. Cell growth for the large-scale production of WT MelB$_{St}$ or D59C MelB$_{St}$ were carried out in *E. coli* DW2 cells (18, 38). Briefly, MelB$_{St}$ purification by cobalt-affinity chromatography (Talon Superflow Metal Affinity Resin, Takara) after extraction by 1.5% UDM. MelB$_{St}$ protein was eluted with 250 mM imidazole in a buffer containing 50 mM NaPi, pH 7.5, 200 mM NaCl, 0.035% UDM, and 10% glycerol, and further dialyzed to change the buffer conditions accordingly.

Protein concentration assay. The Micro BCA Protein Assay (Pierce Biotechnology, Inc.) was used to determine the protein concentration.

Isothermal titration calorimetry. All ITC ligand-binding assays were performed with the TA Instruments (Nano-ITC device) as described (19), which yields the exothermic binding as a positive peak. The MelB$_{St}$ was dialyzed overnight with assay buffer containing 20 mM Tris-HCl (pH7.5), 100 mM NaCl, 10% glycerol and 0.035% UDM. The ligands are prepared by dissolving in the same batch of dialysis buffer for buffer matching. In a typical experiment, the titrand

(MelB$_{St}$) placed in the ITC Sample Cell was titrated with the specified titrant raffinose, α-methyl

galactoside, melibiose, or α-NPG (placed in the Syringe) in the assay buffer by an incremental

injection of 2.5-μL aliquots at an interval of 2500 sec at a constant stirring rate of 250 rpm

(nano-ITC). MelB$_{St}$ protein samples were buffer-matched to the assay buffer by dialysis. The

normalized heat changes were subtracted from the heat of dilution elicited by the last few

injections, where no further binding occurred, and the corrected heat changes were plotted

against the mole ratio of titrant versus titrand. The values for the binding association constant (K_a)

were obtained by fitting the data using the one-site independent-binding model included in the

NanoAnalyze software (version 3.7.5). The dissociation constant (K_d) = $1/K_a$.

Crystallization, native diffraction data collection, and processing. The D59C MelB$_{St}$ dialyzed

overnight against the sugar-free dialysis buffer (20 mM Tris-HCl, pH 7.5, 100 mM NaCl, 0.035%

UDM, and 10% glycerol), concentrated with Vivaspin column at 50 kDa cutoff, and stored at -

80 °C. A phospholipid stock solution of 20 mM was prepared by dissolving the *E. coli* Extract

Polar (Avanti, 100600) with a dialysis buffer containing 0.01% DDM. The protein sample was

diluted to a final concentration of 10 mg/ml with the same sugar-free dialysis buffer,

supplemented with phospholipids at 3.6 mM and 30 mM of melibiose or α-MG, 40 mM raffinose,

or 6 mM of α-NPG in DMSO solution. Crystallization trials were carried out by the hanging-

drop vapor-diffusion method at 23 °C by mixing 2-μL protein with 2-μL reservoir. Crystals from

D59C MelBSt protein with the melibiose, α-MG, or α-NPG appeared against a reservoir

consisting of 100 mM Tris-HCl, pH 8.5, 100 mM NaCl$_2$, 50 mM CaCl$_2$, and 32-35% PEG 400.

For the raffinose-containing sample, the crystals were collected from 100 mM Tris-HCl, pH 8.5,

50 mM CaCl$_2$, 50 mM BaCl$_2$, and 32.5% PEG 400. All crystals were frozen with liquid nitrogen

in 2 weeks, and tested for X-ray diffraction at the Lawrence Berkeley National Laboratory ALS

BL 5.0.1 (dataset for melibiose, α-MG, and α-NPG-containing complex) or 5.0.2 (raffinose-

containing datasets) via remote data collection method.

ALS auto-processing XDS or DIALS programs output files were further reduction by

AIMLESS in the ccp4i2 program for the structure solution (39). The statistics in data collection

are described in **Table S1**.

Structure determination. The structure determination was performed by the Molecular

Replacement method using the α-NPG-bound D59C MelB$_{St}$ mutant structure [PDB ID 7L17] as

the search template, followed by rounds of manual building and refinement to resolutions of 2.70

Å, 3.05, 3.45, or 3.68 Å for structure with α-NPG, melibiose, raffinose, or α-MG, respectively, in

Phenix (40). The structures were modeled from positions 2 to 453 or 456, respectively, without a

gap, and the missing side chains due to the density disorder, and Ramachandran assessment were

listed in the **Tables S6**.

Sugar docking and modeling. A strong positive density, with varying sizes and shapes, was

observed in the difference maps of each of the four structures. The size and shape matched the

sugars that co-crystallized, and the docked sugar molecules fitted well with the densities. The

sugar refinement restrains were generated from SMILES using the ELBOW program in Phenix.

To the 2.7 Å α-NPG-bound map, six water molecules were modeled. In addition, a PEG

molecule (ligand ID 1PE) was also model to a strong positive density with a sausage shape

aligning with the helix IX.

Hydrogen-deuterium exchange coupled to mass spectrometry (HDX-MS). In-solution HDX-

MS experiment was performed to study the substrate-induced structural dynamics of MelB$_{St}$. As

described (21), the labeling, quenching, lipids removal, and online digestion were achieved using

a fully automated manner using a HDx3 extended parallel system (LEAP Technologies,

Morrisville, NC)(41, 42). MelB$_{St}$ was prepared at 50.0 μM in a Na$^+$-free buffer (25 mM Tris-

HCl, pH 7.5, 150 mM choline chloride, 10% glycerol, 0.035% UDM in H$_2$O), either in the

absence of a substrate (Apo) or in the presence of 100 mM melibiose, 100 mM NaCl, or both.

The hydrogen/deuterium exchange reaction as described previously (21). Briefly, aliquots of 4 μl

of each sample were diluted10-fold into the labeling buffer (25 mM Tris-HCl, pD 7.5, 50 mM

choline chloride, 10% glycerol, 0.035% UDM in D$_2$O), without or with 100 mM of melibiose,

Na$^+$, or both. Labeled samples were incubated in D$_2$O buffer at 20 °C for multiple time points

(30 sec, 300 sec, and 3000 sec) in triplicates and non-deuterated controls were prepared in a

similar manner except H$_2$O buffer was used in the labeling step.

At each designated time point, reaction was quenched by adding an equal volume of ice-cold

quench buffer (6 M urea, 100 mM citric acid, pH, 2.3 in H$_2$O) for 180 seconds at 0 °C and

immediately subjected to a lipid filtration module integrated on the LEAP PAL system. After

incubation of a 60 sec with ZrO2 particles, the LEAP X-Press then compressed the filter

assembly to separate proteins from the ZrO2 particles-bound phospholipids and detergents. The

filtered protein sample was injected into a cool box for online digestion and separation.

LC/MS bottom-up HDX was performed using a Thermo Scientific™ Ultimate™ 3000

UHPLC system and Thermo Scientific^{TM} Orbitrap Eclipse^{TM} Tribrid^{TM} mass spectrometer.

Samples were digested with a Nepenthesin-2 (Affipro, Czech Republic) column at 8 °C and then

trapped in a 1.0 mm x 5.0 mm, 5.0 μm trap cartridge for desalting over 180 sec. The resulting

peptides were then separated on a Thermo Scientific™ Hypersil Gold^{TM}, 50 x1 mm, 1.9 μm, C18

column with a gradient of 10 % to 40 % gradient (A: water, 0.1 % formic acid; B: acetonitrile,

0.1 % formic acid) for 15 minutes at a flow rate of 40 μL/min. A pepsin wash was added in

between runs to minimize the carryover.

A nonspecific digested peptide database has been created for MelB$_{St}$ with a separate MS/MS

measurement of non-deuterated samples as described (21). Digested peptides from undeuterated

MelB$_{St}$ protein were identified on the orbitrap mass spectrometer using the same LC gradient as

the HDX-MS experiment. Using the Thermo BioPharma Finder software (v 5.1), MS2 spectra

were matched to the MelB$_{St}$ sequence with fixed modifications.

A number of 146 or 150 peptide assignments (confident HDX data cross all labeling times)

were confirmed for MelB$_{St}$ samples giving 86-87% sequence coverage. The MS data were

processed using the Sierra Analytics HDExaminer software with the MelB$_{St}$ peptide database.

Following the automated HDX-MS analysis, manual curation was performed. Upon the

completion of the data review, a single charge state with high-quality spectra for all replicates

across all HDX labeling times was chosen to represent HDX for each peptide. Differential HDX

data were tested for statistical significance using the hybrid significance testing criteria method

with an in-house MATLAB script, where the HDX differences at different protein states were

calculated ($\Delta D = D_{Holo} - D_{Apo}$). Mean HDX differences from the three replicates were assigned

as significant according to the hybrid criteria based on the pooled standard deviation and

Welch's t-test with $P < 0.05$. The statistically significant differences observed at each residue

($D_{Mel - Apo} < |0.186|$, $\Delta D_{(Na+)}$ - Apo $< |0.224|$, or $\Delta D_{Na(+)Mel}$ - Apo $< |0.175|$) were used to map

HDX consensus effects based on overlapping peptides onto the structure models.

Statistics and reproducibility. All experiments were performed 2-4 times. The average values

were presented in the table with standard errors. An unpaired t-test was used for statistical

analysis.

Graphs. Pymol (3.1.5.1) (43) and UCSF ChimeraX (X) were used to generate all graphs. The

program Origin 2024 was used to plot the ITC curves.

Data availability statement

The x-ray diffraction datasets and models have been deposited to wwPDB under the accession

codes 9OLD for the α-nitrophenyl galactoside-bound, 9OLI for the melibiose-bound, 9OLR for

the α-methyl galactoside, as well as 9OLP for the raffinose-bound D59C MelB$_{St}$.

Competing Interests

Y. S. and R. V. are employees of *Thermo Fisher Scientific*.

Acknowledgments

The x-ray diffraction datasets were collected at ALS BL 5.01 or 5.0.2. This work was

supported by the National Institutes of Health Grant R35GM153222 to L.G.

References

1. Ferrada, E., and Superti-Furga, G. (**2022**) A structure and evolutionary-based classification of solute carriers. *iScience* 25, 105096

2. Pao, S. S., Paulsen, I. T., and Saier, M. H., Jr. (**1998**) Major facilitator superfamily. *Microbiol Mol Biol Rev* 62, 1-32

3. Cesar-Razquin, A., Snijder, B., Frappier-Brinton, T., Isserlin, R., Gyimesi, G., Bai, X. *et al.* (**2015**) A Call for Systematic Research on Solute Carriers. *Cell* 162, 478-487

4. Lin, L., Yee, S. W., Kim, R. B., and Giacomini, K. M. (**2015**) SLC transporters as therapeutic targets: emerging opportunities. *Nat Rev Drug Discov* 14, 543-560

5. Guan, L., and Kaback, H. R. (**2006**) Lessons from lactose permease. *Annu Rev Biophys Biomol Struct* 35, 67-91

6. Yan, N. (**2015**) Structural Biology of the Major Facilitator Superfamily Transporters. *Annu Rev Biophys* 44, 257-283

7. Drew, D., North, R. A., Nagarathinam, K., and Tanabe, M. (**2021**) Structures and General Transport Mechanisms by the Major Facilitator Superfamily (MFS). *Chem Rev* 121, 5289-5335

8. Guan, L. (**2023**) The rapid developments of membrane protein structure biology over the last two decades. *BMC Biol* 21, 300

9. Nguyen, L. N., Ma, D., Shui, G., Wong, P., Cazenave-Gassiot, A., Zhang, X. *et al.* (**2014**) Mfsd2a is a transporter for the essential omega-3 fatty acid docosahexaenoic acid. *Nature* 509, 503-506

10. Chua, G. L., Tan, B. C., Loke, R. Y. J., He, M., Chin, C. F., Wong, B. H. *et al.* (**2023**) Mfsd2a utilizes a flippase mechanism to mediate omega-3 fatty acid lysolipid transport. *Proc Natl Acad Sci U S A* 120, e2215290120

11. Cater, R. J., Chua, G. L., Erramilli, S. K., Keener, J. E., Choy, B. C., Tokarz, P. *et al.* (**2021**) Structural basis of omega-3 fatty acid transport across the blood-brain barrier. *Nature* 595, 315-319

12. Wilson, T. H., and Ding, P. Z. (**2001**) Sodium-substrate cotransport in bacteria. *Biochim Biophys Acta* 1505, 121-130

13. Maehrel, C., Cordat, E., Mus-Veteau, I., and Leblanc, G. (**1998**) Structural studies of the melibiose permease of *Escherichia coli* by fluorescence resonance energy transfer. I. Evidence for ion-induced conformational change. *J Biol Chem* 273, 33192-33197

14. Meyer-Lipp, K., Sery, N., Ganea, C., Basquin, C., Fendler, K., and Leblanc, G. (**2006**) The inner interhelix loop 4-5 of the melibiose permease from *Escherichia coli* takes part in conformational changes after sugar binding. *J Biol Chem* 281, 25882-25892

15. Guan, L., Nurva, S., and Ankeshwarapu, S. P. (**2011**) Mechanism of melibiose/cation symport of the melibiose permease of *Salmonella typhimurium*. *J Biol Chem* 286, 6367-6374

16. Granell, M., Leon, X., Leblanc, G., Padros, E., and Lorenz-Fonfria, V. A. (**2010**) Structural insights into the activation mechanism of melibiose permease by sodium binding. *Proc Natl Acad Sci U S A* 107, 22078-22083

17. Guan, L. (2018) Na(+)/Melibiose Membrane Transport Protein, MelB In In: Roberts G., Watts A., European Biophysical Societies (eds) Encyclopedia of Biophysics, 2nd Ed. Roberts G, Watts A, eds. Springer, Berlin, Heidelberg, 10.1007/978-3-642-35943-9_10082-1

18.   Ethayathulla, A. S., Yousef, M. S., Amin, A., Leblanc, G., Kaback, H. R., and Guan, L.
(**2014**) Structure-based mechanism for Na(+)/melibiose symport by MelB. *Nat Commun*
5, 3009
19.   Hariharan, P., and Guan, L. (**2017**) Thermodynamic cooperativity of cosubstrate binding
and cation selectivity of Salmonella typhimurium MelB. *J Gen Physiol* 149, 1029-1039
20.   Guan, L., and Hariharan, P. (**2021**) X-ray crystallography reveals molecular recognition
mechanism for sugar binding in a melibiose transporter MelB. *Commun Biol* 4, 931
21.   Hariharan, P., Shi, Y., Katsube, S., Willibal, K., Burrows, N. D., Mitchell, P. *et al.* (**2024**)
Mobile barrier mechanisms for Na(+)-coupled symport in an MFS sugar transporter. *Elife*
12,
22.   Hariharan, P., Bakhtiiari, A., Liang, R., and Guan, L. (**2024**) Distinct roles of the major
binding residues in the cation-binding pocket of the melibiose transporter MelB. *J Biol*
Chem 107427
23.   Katsube, S., Liang, R., Amin, A., Hariharan, P., and Guan, L. (**2022**) Molecular Basis for
the Cation Selectivity of Salmonella typhimurium Melibiose Permease. *J Mol Biol* 434,
167598
24.   Guan, L., Jakkula, S. V., Hodkoff, A. A., and Su, Y. (**2012**) Role of Gly117 in the
cation/melibiose symport of MelB of *Salmonella typhimurium*. *Biochemistry* 51, 2950-
2957
25.   Hariharan, P., and Guan, L. (**2014**) Insights into the inhibitory mechanisms of the
regulatory protein IIA(Glc) on melibiose permease activity. *J Biol Chem* 289, 33012-
33019
26.   Liang, R., and Guan, L. (**2024**) Atomic-Level Free Energy Landscape Reveals
Cooperative Symport Mechanism of Melibiose Transporter. *bioRxiv*
27.   Hariharan, P., and Guan, L. (**2021**) Cooperative binding ensures the obligatory
melibiose/Na+ cotransport in MelB. *J Gen Physiol* 153,
28.   Masson, G. R., Burke, J. E., Ahn, N. G., Anand, G. S., Borchers, C., Brier, S. *et al.*
(**2019**) Recommendations for performing, interpreting and reporting hydrogen deuterium
exchange mass spectrometry (HDX-MS) experiments. *Nat Methods* 16, 595-602
29.   Jia, R., Martens, C., Shekhar, M., Pant, S., Pellowe, G. A., Lau, A. M. *et al.* (**2020**)
Hydrogen-deuterium exchange mass spectrometry captures distinct dynamics upon
substrate and inhibitor binding to a transporter. *Nat Commun* 11, 6162
30.   Zmyslowski, A. M., Baxa, M. C., Gagnon, I. A., and Sosnick, T. R. (**2022**) HDX-MS
performed on BtuB in E. coli outer membranes delineates the luminal domain's allostery
and unfolding upon B12 and TonB binding. *Proc Natl Acad Sci U S A* 119, e2119436119
31.   Guan, L., Mirza, O., Verner, G., Iwata, S., and Kaback, H. (**2007**) Structural
determination of wild-type lactose permease. *Proceedings of the National Academy of*
Sciences of the United States of America 104, 15294-15298
32.   Kaback, H. R., and Guan, L. (**2019**) It takes two to tango: The dance of the permease. *J*
Gen Physiol 151, 878-886
33.   Kumar, H., Kasho, V., Smirnova, I., Finer-Moore, J. S., Kaback, H. R., and Stroud, R. M.
(**2014**) Structure of sugar-bound LacY. *Proc Natl Acad Sci U S A* 111, 1784-1788
34.   Grytsyk, N., Sugihara, J., Kaback, H. R., and Hellwig, P. (**2017**) pKa of Glu325 in LacY.
Proc Natl Acad Sci U S A 114, 1530-1535
35.   Mirza, O., Guan, L., Verner, G., Iwata, S., and Kaback, H. R. (**2006**) Structural evidence
for induced fit and a mechanism for sugar/H+ symport in LacY. *EMBO J* 25, 1177-1183

36.   Smirnova, I. N., Kasho, V., and Kaback, H. R. (**2008**) Protonation and sugar binding to
LacY. *Proc Natl Acad Sci U S A* 105, 8896-8901

37.   Botfield, M. C., and Wilson, T. H. (**1988**) Mutations that simultaneously alter both sugar
and cation specificity in the melibiose carrier of Escherichia coli. *J Biol Chem* 263,
12909-12915

38.   Pourcher, T., Leclercq, S., Brandolin, G., and Leblanc, G. (**1995**) Melibiose permease of
Escherichia coli: large scale purification and evidence that H(+), Na(+), and Li(+) sugar
symport is catalyzed by a single polypeptide. *Biochemistry* 34, 4412-4420

39.   Potterton, L., Agirre, J., Ballard, C., Cowtan, K., Dodson, E., Evans, P. R. *et al.* (**2018**)
CCP4i2: the new graphical user interface to the CCP4 program suite. *Acta Crystallogr D*
Struct Biol 74, 68-84

40.   Adams, P. D., Afonine, P. V., Bunkoczi, G., Chen, V. B., Davis, I. W., Echols, N. *et al.*
(**2010**) PHENIX: a comprehensive Python-based system for macromolecular structure
solution. *Acta Crystallogr D Biol Crystallogr* 66, 213-221

41.   Hamuro, Y., Coales, S. J., Southern, M. R., Nemeth-Cawley, J. F., Stranz, D. D., and
Griffin, P. R. (**2003**) Rapid analysis of protein structure and dynamics by
hydrogen/deuterium exchange mass spectrometry. *J Biomol Tech* 14, 171-182

42.   Hamuro, Y., and Coales, S. J. (**2018**) Optimization of Feasibility Stage for
Hydrogen/Deuterium Exchange Mass Spectrometry. *J Am Soc Mass Spectrom* 29, 623-
629

43.   Schrodinger, L. (**2013**) The PyMOL Molecular Graphics System, Version 1.5.

Figure Legends

Fig. 1. Crystal structure of D59C MelBSt in complex with α-sugars or sugar analogs. The

identities of the bound sugars are indicated at the top. Upper row: Cartoon representation of the

structures of D59C MelB$_{St}$ bound with α-nitrophenyl galactoside (α-NPG), melibiose (α-

disaccharide), and raffinose (α-trisaccharide), respectively, along with a surface presentation of

D59C MelB$_{St}$ complexed with α-methyl galactoside (α-MG). All structures were oriented with

the cytoplasmic side on top and the N-terminal domain (colored green) on the left. Each sugar

molecule is colored yellow. The blue sticks and surfaces indicate residues within 5 Å of the

sugar molecules. Lower row: the sugar-binding pockets. Residues from the N-terminal and C-

terminal domains were shown in surface representation, and colored in green and blue,

respectively.

Fig. 2. α-NPG binding. The cross-eye stereo view of α-NPG binding. **(a)** Isomesh map of the

α-NPG-binding site, contoured at a level of σ =1.5. Dashed lines indicate distances within

hydrogen-bonding or salt-bridge interactions (Å). W1, water molecule 1 in red. The helices are

labeled with Roman numerals.

Fig. 3. Binding of melibiose, α-MG, and raffinose. All residues within a 5-Å distance to the

bound substrates were shown in sticks. Isomesh maps for each sugar and Asp19 were contoured

at levels of σ = 1.5 for melibiose and raffinose or σ = 1.0 for α-MG. Dashed lines, the distances

within hydrogen-bonding and salt-bridge interactions (Å). **(a)** Melibiose binding. **(b)** α-MG

binding. **(c)** Raffinose binding. **(d)** Residues in the sugar-binding pockets from the alignment of

all four structures. **(e)** Substrates from the alignment of all four structures. Carbon positions on

the galactosyl (C1-6) and glucosyl moiety (C1-6′) are labeled on the melibiose molecule. Trp342

was removed for clarity in panels a, b, and d.

Fig. 4. Binding affinity determination via ITC. The binding measurement of α-MG or

raffinose to the WT MelB$_{St}$ was performed with a NanoITC calorimeters (TA Instruments) at

25 °C as described in Methods. The thermograms were plotted as baseline-corrected heat rate

(μJ/sec; left axis) vs. time (bottom axis) for the titrant to MelB$_{St}$ (red for α-MG and blue for

raffinose) or to buffer (light blue). The heat change ΔQ (μJ; filled black symbol) was plotted

against the mole ratio of the sugar to WT MelB$_{St}$ (top/right axes). **(a)** α-MG. (b) Raffinose.

Fig. 5. HDX-MS. HDX experiments on WT MelB$_{St}$ in Apo or Holo states (with melibiose, Na$^+$,

or Na$^+$ plus melibiose) were conducted as described in Methods. **(a)** Residual plots (D$_{Holo - Apo}$).

Differential deuterium uptakes ΔDs, calculated from paired conditions for position 2-470, were

plotted against overlapping peptides for each time point and the total uptake. Black, cyan, and

purple bars, the deuterium uptake at 30, 300, 3000 sec, respectively; dark gray curve, total uptake

from all three time points. Deprotection, $\Delta D_{Holo - Apo} > 0$; protection, $\Delta D_{Holo - Apo} < 0$. Each

sample was analyzed in triplicates. Regions not covered by labeled peptides were excluded.

Dashed lines, the global threshold values calculated from each dataset. **(b)** Mapping onto the 3D

structure of the melibiose-bound state. Peptides with ΔD value greater than the absolute value of

the threshold and P < 0.05 at any time point were colored according to the legend on the figure.

The membrane region of MelB$_{St}$ is indicated by a gray bar. Each peptide or overlapping peptide

was indicated by arrows pointing to the start of the peptide. The helices I, V, VIII, and IX, as

well as the C-terminal helix (lid), are labeled individually. Mel, melibiose.

Fig. 6. Allosteric effects on deuterium uptake by remote binding of Na$^+$. The time course of

percentage deuterium uptake was plotted against the labeling time at 0, 30, 300, and 3000 sec for

the representative peptides that cover the residues responsible for the specificity of sugar or

cation binding in MelB$_{St}$. The peptide sequences are shown for each panel. *, statistically

significant; **, statistically highly significant. Helices I and V (sugar-binding only), II (cation

binding only), and IV (both), were shown in tube representation. Residues involved in sugar

binding are represented by sticks and highlighted in magenta (helix I), blue (helix IV), and red

(helix V). Residues involved in cation binding are highlighted in black. Labels in the time course

and on the model are color-coded, except for the stick representing Thr121, which is black.

Helices are labeled with Roman numerals and positioned at the beginning of each helix.

Fig. 1

α-NPG   Melibiose   Raffinose   α-MG

C-

-N

Cytoplsmic side

Inner Barrier

N-terminal domain   C-terminal domain

Periplasmic side

Fig. 2

Fig. 3

Fig. 4

Fig. 5

(a) Residual plot

$D_{Melibiose} - D_{Apo}$

Deprotection

Peptide 209-218
(Middle loop)

+ 0.186
- 0.186

Protection

$D_{Na(+)} - D_{Apo}$

+ 0.224
- 0.224

Peptide 213-226
(Middle loop)

$D_{[Na(+)Melibiose]} - D_{Apo}$

+ 0.175
- 0.175

Peptide 213-226
(Middle loop)

Peptide ID number

▬ Uptake at 30 sec ── Total deuterium uptake
▬ Uptake at 300 sec --- Global threshold
▬ Uptake at 3000 sec

(b) Structural Mapping

C-   Cytoplasmic side

Lid or tail helix

213-226 (*)
209-218 (*)
2-27 (**)
137-159 (**)

N

441-453 (**)
364-374 (**)
292-298 (**)
284-293 (**)#

Inner

Na+-binding pocket

Membrane

Melibiose

394-405 (*)
317-326 (**)

Outer

99-109 (**)
36-47 (**)

264-271 (**)

Blue, protected by Mel, Na(+), or Na(+)Mel
Red, deprotected by Mel or Na(+)Mel
Magenta, deprotected by Mel only
Cyan, deprotected by Mel but protected by Na(+)Mel
#, deprotected by Na(+)Mel (P<0.001), and deprotectd by Mel (P=0.15)
(*), ΔD values are statistically significant, P<0.05
(**), ΔD values are statistically highly significant, P<0.0

Fig. 6

H-V

Mel	Na⁺	Mel + Na⁺

140-151
EREQLVPF
PREF

151-156
FASLAG

H-I

Mel	Na⁺	Mel + Na⁺

16-20
FGKDF

21-27
AIGLVYM

21-26

21-27

H-IV

Mel	Na⁺	Mel + Na⁺

121-130
TIMDIPFWSL

120-123
YTIM

H-II

Mel	Na⁺	Mel + Na⁺

53-62
IWDAINDPIM