Chromosome Size as a Robust Predictor of Recombination Rate: Insights from
Holocentric and Monocentric Systems

Authors: František Zedek[1], Petr Bureš[1], Tammy L. Elliott[1,2], Marcial Escudero[3], Kay Lucek[4],
André Marques[5]

[1]Department of Botany and Zoology, Faculty of Science, Masaryk University, Kotlářská 2,
61137 Brno, Czech Republic

[2]Bolus Herbarium, Department of Biological Sciences, University of Cape Town, Private Bag
X3, Rondebosch, 7701, Cape Town, South Africa

[3]Department of Plant Biology and Ecology, Faculty of Biology, University of Seville, Reina
Mercedes 6, 41012 Seville, Spain

[4]Biodiversity Genomics Laboratory, Institute of Biology, University of Neuchâtel, 11 rue
Emile-Argand, 2000 Neuchâtel, Switzerland

[5]Department of Chromosome Biology, Max Planck Institute for Plant Breeding Research,
Cologne, Germany

Corresponding author: František Zedek, email: fzedek@gmail.com

Abstract

Recombination is a fundamental evolutionary process essential for generating genetic
diversity, facilitating adaptation, and driving speciation. However, direct measurement of
recombination rate remains challenging, as standard methods—such as chiasma counts or
genetic linkage maps—are labor-intensive and often infeasible for non-model species. In this
study, we identify chromosome number and mean chromosome size as practical proxies for
genome-wide recombination rate by analyzing genetic map data from 73 insect species and
supplementary analyses of 157 monocentric flowering plants. We confirm the long-standing
hypothesis that monocentric species have nearly twice as many crossovers per
chromosome as holocentric species, reflecting structural constraints imposed by diffuse
centromeres. Using both ordinary and phylogenetically informed Bayesian regression
models, we show that recombination rate increases with chromosome number and
decreases with mean chromosome size. Crucially, mean chromosome size is a significantly
better predictor, particularly in holocentric species. This insight enables recombination rate
estimation in thousands of species with known genome and chromosome sizes, thereby
allowing hypothesis testing at scales previously unattainable. Building on these results, we
present predictive models applicable to poorly studied holocentric plants. Overall, our study
highlights the pivotal role of chromosome architecture in recombination evolution and
provides an accessible framework for evolutionary genomic research across diverse
lineages.

Keywords: Recombination rate, Chromosome size, Chromosome number, Crossover
frequency, Holocentric chromosomes, Monocentric chromosomes, Evolutionary genomics,
Meiotic recombination, Phylogenetic comparative methods, Genome architecture

Article summary: Why do some species have more genetic shuffling than others? We
investigated 73 insect species with detailed genetic maps and found that mean chromosome
size is the best predictor of genome-wide recombination rate, not the number of
chromosomes. This means we can now estimate recombination rates even in species where
genetic maps are unavailable, using only genome size and chromosome number. We also
show that chromosome structure matters—species with a single centromere per
chromosome (monocentric) have nearly double the recombination compared to those with
diffuse centromeres (holocentric). These insights open new doors for studying evolution
across many organisms.

Introduction

The key evolutionary implications of recombination are twofold: recombination can facilitate adaptation across environments by generating and maintaining genetic variation, or it can break up co-adapted allele combinations, potentially disrupting current optimal gene combinations (Ritz et al., 2017; Stapley et al., 2017a). Recombination also plays a crucial role in speciation as the formation of new species often involves a progressive reduction in genetic exchange in specific genomic regions that act as barriers to gene flow (Butlin, 2005). Structural changes along the chromosomes may suppress recombination locally, promoting the evolution of speciation genes (Zhang et al., 2021). On a genome-wide level, a lot of attention has been given to the evolutionary advantage of sex, which promotes recombination. However, the reason why average genome-wide recombination also varies between sexually reproducing organisms is less understood (Stapley et al., 2017a). In sexual eukaryotes, genome-wide recombination rate is determined by the number of crossovers (COs; i.e., chiasmata) per meiosis and by random meiotic segregation of homologs (Stapley et al., 2017b; Otto & Payseur, 2019; Hofstatter et al., 2021). Because estimating recombination rates with standard approaches, such as counting chiasmata or establishing linkage maps (Stapley et al. 2017b), demands time-consuming analysis and is typically impractical for non-model species, it remains challenging to investigate recombination rate-related hypotheses at a macroevolutionary scale across species and environments.

Since at least one CO per chromosome is usually required for proper segregation during meiosis (a phenomenon known as CO assurance; Hunter, 2015) and the formation of additional COs is tightly regulated by CO interference (a mechanism that prevents COs from occurring too closely together; Hunter, 2015), the number of COs per chromosome is maintained within a narrow range (up to 3 COs in 80% of cases; Lian et al., 2023). Therefore, chromosome number is expected to predict genome-wide recombination rate across species (Stapley et al., 2017b). Having more chromosomes also increases the potential for diverse combinations during the assortment of chromosomes in gametes. However, variation in genome size can obscure this relationship.

All else being equal, the necessity of at least one CO per chromosome results in proportionally higher recombination rates in shorter chromosomes (Lynch et al., 2011; Tigano et al., 2022). Heterochromatin is moreover associated with reduced recombination, for instance, in pericentromeric regions, which often accumulate transposable elements and display suppressed CO activity (Kent et al., 2017). Consequently, larger chromosomes, which tend to contain proportionally larger heterochromatic regions (Leitch et al., 2023), may experience a lower rate of recombination per unit length compared to smaller chromosomes. This dynamic could explain the commonly observed negative association between recombination rate and chromosome size of individual chromosomes inside a single karyotype (Kaback et al., 1992; Jensen-Seaman et al., 2004; Backström et al., 2010; Martin et al., 2019; Brazier & Glémin, 2022; Shipilina et al., 2022; Näsvall et al., 2023; Palahí I Torres et al., 2023), which appears to translate to a negative relationship between mean chromosome size and average genome-wide recombination rate (Lynch et al., 2011; Brazier & Glémin, 2022). Thus, chromosome counts and mean chromosome size can potentially be used as proxies for genome-wide recombination rate comparison across a large number of species.

In eukaryotes, two predominant types of chromosomes have evolved that differ regarding their centromere organization, i.e. monocentric and holocentric chromosomes. Importantly, the accuracy with which chromosome number and mean chromosome size predict genome-wide recombination rate is expected to differ between these two types of chromosome organization. For species with monocentric chromosomes, where kinetochores and chromatid cohesion are restricted to a single centromeric region, the number of COs per chromosome mostly varies between one to four (Crismani & Mercier, 2012; Mercier et al., 2015; Blary & Jenczewski, 2019; Otto & Payseur, 2019; Brazier & Glémin, 2022; Lian et al., 2023), but higher numbers have also been reported (Jones & Franklin, 2006; Mancera et al., 2008; Brazier & Glémin, 2022; Lian et al., 2023). In contrast, species with holocentric chromosomes, where kinetochores are maintained along the entire chromosome length (Bureš et al., 2013; Mandrioli & Manicardi, 2020; Hofstatter et al., 2021) cannot rely on the two-step loss of cohesion during meiosis and restrict the CO frequency per homolog to one or two chiasmata at maximum (Schvarzstein et al., 2010; Metlers et al., 2012; Cuacos et al., 2015; Marques & Pedrosa-Harand, 2016; Mandrioli & Manicardi, 2020; Castellani et al., 2024). This upper limit in holocentric species is primarily supported by cytological observations showing that additional chiasmata tend to disrupt proper chromosome segregation (Nokkala et al., 2024; Castellani et al., 2024). The exact mechanism underlying this constraint remains unknown, but it likely relates to the absence of a localized centromere and to the distinct mode of cohesion release along holocentric chromosomes. Such decreased recombination has been shown for both holocentric plants (e.g., Nordenskiöld, 1962; Cabral et al., 2014; Heckmann et al., 2014; Hofstatter et al., 2021; Castellani et al., 2024) and animals (e.g., Barnes et al., 1995; Nokkala et al., 2004; Viera et al., 2009; Lukhtanov et al., 2018). Another difference between holocentric and monocentric species is the composition of chromatin. In holocentric species, chromatin is often not compartmentalized and shows no obvious euchromatic or heterochromatic clusters as in monocentric species (Mandrioli & Manicardi, 2012; Heckmann et al., 2013; Hofstatter et al., 2022; Mata-Sucre et al., 2024) with homogeneous TE distributions along chromosomes (Hofstatter et al. 2022; Shipilina et al., 2022; Mata-Sucre et al., 2024; Castellani et al., 2024). This could potentially result in a more uniform – yet still telomere-biased – distribution of recombination rates along holocentric chromosomes compared to monocentric ones (Haenel et al., 2018; Palahí i Torres et al., 2022; Hofstatter et al., 2022; Shipilina et al., 2022; Castellani et al., 2024). Furthermore, due to their extended kinetochore and the attachment of spindle microtubules along their entire length, holocentric species are more likely to tolerate chromosomal fissions and fusions that have negligible fitness costs (Bureš et al., 2013; Jankowska et al., 2015; Zedek & Bureš, 2019; Zedek et al., 2020; Mandrioli & Manicardi, 2020; Lucek et al., 2022). Therefore, holocentric species are expected to more readily change chromosome numbers and size and, as a consequence, recombination rates in response to selection compared to monocentric species (Bell, 1982; Escudero et al., 2012; Bureš et al., 2013; Escudero et al. 2013; Bureš & Zedek, 2014). Thus, a higher recombination rate in monocentric species is usually expected to be reached by increasing the number of COs per chromosome (Burt, 2000; Stapley et al., 2017b), whereas in holocentric species, a higher recombination rate is predicted to primarily result from increasing chromosome numbers (and thereby decreasing chromosome size).

Although chromosome numbers and mean chromosome size have been used as proxies for genome-wide recombination rate inference in both monocentric (Pessia et al., 2012; Carta et al., 2018) and holocentric species (Bell, 1982; Escudero et al., 2012; Escudero et al., 2013;

Spalink et al., 2018; Mackintosh et al., 2019; Márquez-Corro et al., 2021; Elliott et al., 2022),
the assumption that these two genomic traits accurately represent recombination rates
should be critically assessed. A system encompassing multiple holocentric and monocentric
lineages, for which chromosome numbers, genome sizes, and linkage maps or properly
counted chiasmata are available to estimate genome-wide recombination rates is required to
address these questions. Insects may constitute such a group (Figure 1), where holocentric
chromosomes have independently evolved in the orders Lepidoptera, Trichoptera,
Hemiptera, Thysanoptera, Psocodea, Zoraptera, Dermaptera, and Odonata, whereas
monocentric chromosomes are more broadly found, for example in Diptera, Coleoptera,
Hymenoptera, Orthoptera, or Blattodea (Melters et al., 2012; Zedek & Bureš, 2019; Mandrioli
& Manicardi, 2020; Marques & Drinnenberg, 2025). Moreover, chromosome number and
size, genome size information, and linkage maps are available for a reasonable amount of
species of both holocentric and monocentric insect orders.

By analyzing recombination data from insect species with well-characterized genetic maps
(or counted chiasmata), we aimed to assess the suitability of chromosome numbers and
mean chromosome size as predictors of genome-wide recombination rate and test the
hypothesis that these predictors are more accurate in holocentric than in monocentric
species. Specifically, we expect that (1) the average number of COs per chromosome is
higher in monocentric species than in holocentric ones, which is the key assumption
stemming from the structural differences between holocentric and monocentric
chromosomes (see above), and (2) chromosome number and size predict genome-wide
recombination rate better in holocentric than monocentric species. To assess the broader
applicability of our results beyond insects, we evaluated the accuracy of chromosome
number and mean chromosome size as predictors of recombination rate in flowering plants
with monocentric chromosomes. Holocentric species could not be meaningfully included, as
only two holocentric plant species currently have established genetic maps (Escudero et al.,
2018; Castellani et al., 2024), but we made a prediction of holocentric plants based on the
results we obtained from insects and monocentric plants.
189
190

Materials and Methods

Data Collection

We compiled genetic map length data for 70 species and chiasma count data for three
additional species, spanning both holocentric and monocentric insects (Supplementary
Table S1). Genome size and chromosome number information were obtained from the NCBI
Genome database (Sayers et al., 2022), the Animal Genome Size Database (Gregory,
2025), and additional publications (Supplementary Table S1). We defined the recombination
rate as the haploid genetic map length divided by the 1C genome size (Stapley et al.,
2017b).

For all but three species, recombination rates were derived directly from available genetic
maps. For *Philaenus spumarius* (Hemiptera), *Aeshna subarctica* (Odonata), and *Forficula*
auricularia (Dermaptera), where linkage maps were not available, we used published
chiasma counts as a proxy for crossover frequency. We converted the mean number of

chiasmata per meiosis into a genetic map length by multiplying by 50, assuming that one
crossover corresponds to 50 centiMorgans (cM) (Anderson et al., 2004; Chuang & Smith,
2023). We then estimated the mean number of crossovers per chromosome by dividing this
value by the haploid chromosome number.

To assess potential artefacts from variation in marker spacing, we tested for associations
between map length and average marker distance (cM per marker). No significant
correlation was detected (Spearman $\rho \approx 0$, $p > 0.05$) and a linear model controlling for
chromosome type likewise showed no effect ($\beta \approx 0$, $p > 0.05$; Supplementary Figure S1).
This confirms that estimates of map length are not biased by marker density.

For all final analyses we used exclusively autosomal map lengths, along with autosomal
mean chromosome sizes and autosomal chromosome numbers. This standardization
ensures consistency across species, as some published linkage maps report only
autosomes. Focusing on autosomes also avoids biases introduced by heteromorphic sex
chromosomes, which vary in size among species and are often missing or inconsistently
represented in linkage maps. In cases where the size of sex chromosomes (N=14) or the
length of sex chromosome maps (N=6) was not reported, we subtracted the average
chromosome size or average map length per chromosome from the total genome size or
total map length, respectively, to approximate the autosomal component. Genome size
estimates were preferentially taken from whole-genome assemblies; this was possible for all
but two species, where flow cytometry remained the only source of information.
Chromosome counts correspond to the haploid number excluding sex chromosomes, unless
explicitly stated otherwise.

Phylogenetic tree

The phylogenetic tree (Figure 1) used in the following analyses was based on the TimeTree
(Kumar et al., 2022), which amalgamates the phylogenetic information of several thousand
studies to build an overarching phylogeny of the Eukaryotic Tree of Life. If a species was not
available from TimeTree (N=10), we used the tip of its most closely related relative, which
was renamed, or added manually based on published phylogenies.

Recombination rate predicted by chromosome number and mean chromosome size

We analyzed autosomal recombination rate (cM/Mb) and the average number of crossovers
(COs) per chromosome in a Bayesian framework using the *brms* R package (Bürkner 2017,
2018, 2021). Because recombination rates are strictly positive and right-skewed, we
modeled them on their original scale using a Gamma regression with a log link in all
analyses.
240
We log-transformed the continuous chromosomal predictors and standardized them prior to
analysis. Specifically, we considered (i) the haploid number of autosomes, (ii) the mean
autosome size (Mb), and (iii) both predictors jointly. Chromosome type (holocentric vs.
monocentric) was included as a categorical factor. Models were first specified with

interaction terms; if the 95% credible interval for an interaction overlapped with zero, we also
analyzed the corresponding additive formulation.
247
248 Phylogenetically informed versions of all models were implemented by including a random
249 intercept of the form (1 | gr(tip, cov = A)), where A is the phylogenetic covariance matrix
250 derived from the species tree using the vcv function in the R package *ape* (Paradis & Schliep
251 2019).
252
For non-phylogenetic models, we used mildly informative priors: normal(0, 2) on regression
coefficients and gamma(2, 0.1) on the Gamma shape parameters. For phylogenetic models,
the same priors were applied to coefficients and shape, with an additional exponential(1)
prior on the phylogenetic random-effect standard deviation. All models were estimated with
four Markov chains and 7–8,000 total iterations per chain, including 2,000 warmup iterations
and stringent sampler settings (adapt_delta = 0.99–0.999, max_treedepth = 15).
259
260 *Evaluation of the Models and Predictive Accuracy*
261
Model fits were evaluated using approximate leave-one-out cross-validation (LOOIC) and
Bayesian R^2. To further assess predictive accuracy specifically in the comparison between
holocentric and monocentric species, we also calculated error metrics based on observed
versus fitted recombination rates. We report root mean squared error (RMSE) and mean
absolute error (MAE) as absolute measures, and their standardized versions relative to the
observed variation in recombination rates (rel_RMSE and rel_MAE). These relative metrics
facilitate interpretation by accounting for differences in the magnitude of recombination rates
between chromosome types.
270

Assessing the relation of crossover number per chromosome to chromosome type

To test whether the average number of COs per chromosome differed between holocentric
and monocentric species, we pursued two complementary approaches. Given that the
number of COs per chromosome is strictly positive, we first modeled it using a Gamma
regression with a log link and used chromosome type as the main predictor. The same priors
as in the recombination rate analyses (normal(0,2) for regression coefficients and
gamma(2,0.1) for the shape parameter) were applied. Phylogenetic non-independence was
incorporated by including a random intercept specified as (1│gr(phylo, cov = A)).

In addition, we employed Ornstein–Uhlenbeck (OU) phylogenetic modeling to evaluate the
evolutionary dynamics of CO number variation between holocentric and monocentric
lineages. OU models extend Brownian motion by allowing traits to evolve towards
evolutionary optima (θ), thereby testing whether different chromosome types are associated
with distinct selective regimes. Analyses were implemented in the R package *OUwie*
(Beaulieu et al., 2012). To assign chromosome type states across the phylogeny, we used
stochastic character mapping (SIMMAP) under an equal-rates (ER) model, which generates
probabilistic reconstructions of ancestral states and thus incorporates uncertainty in the
evolutionary history of chromosome type.

We compared seven standard models: two Brownian motion models (BM1 and BMS), which
assume a constant variance of the trait through time (BM1 with a single global rate, BMS
with regime-specific rates), and five Ornstein–Uhlenbeck models (OU1, OUM, OUMV,
OUMA, and OUMVA). OU1 assumes a single optimum θ with constant strength of selection
(α) and variance ($σ^2$) across the tree, while OUM allows θ to vary between regimes
(holocentric vs. monocentric). OUMV allows both θ and $σ^2$ to vary, OUMA allows θ and α to
vary, and OUMVA allows all three parameters (θ, α, and $σ^2$) to vary among regimes. Model
selection was based on weighted AICs, and the best-fitting model(s) were further evaluated
using 500 bootstrap replicates in to obtain empirical estimates and 95% confidence intervals
for θ.
298
299 *Evaluation of chromosome number and mean chromosome size predictors in angiosperms*
300
To assess the broader applicability of our results beyond insects, we evaluated the accuracy
of chromosome number and mean chromosome size as predictors of recombination rate in
flowering plants with monocentric chromosomes. Holocentric species could not be
meaningfully included, as only two holocentric plant species currently have genetic maps
(Escudero et al., 2018; Castellani et al., 2024). To this end, we obtained data on
recombination rates, genome sizes, and chromosome numbers from Stapley et al. (2017b)
and randomly selected one phylogenetic tree from a set of 100 angiosperm phylogenies
provided by Forest (2023). The dataset and phylogenetic tree overlapped with 157
angiosperm species (Supplementary Table S2), which we used in the regression models
and their evaluation as described above for insects. To facilitate convergence and
comparability of parameter estimates, predictors in phylogenetic models were z-transformed
(centered and scaled to unit variance).

Results

We obtained recombination rate, chromosome number, chromosome size, and the average
number of COs per chromosome for 33 holocentric (1 Dermaptera, 5 Hemiptera, 25
Lepidoptera, and 2 Odonata) and 40 monocentric (2 Blattodea, 7 Coleoptera, 9 Diptera, 19
Hymenoptera, and 3 Orthoptera) insect species (Figure 1; Supplementary Table S1).

Differences in Crossover Number between Chromosome Types

We first assessed whether the average number of COs per chromosome differs between
holocentric and monocentric insect species. Using a phylogenetically informed Bayesian
Gamma regression, we modeled COs per chromosome as a function of chromosome type
with species relatedness included as a random effect (Figure 2A; Supplementary Table S3).
The intercept for holocentric species was 0.15 (95% credible interval [CI]: –0.32 to 0.61),
corresponding to an expected mean of ≈ 1.16 COs per chromosome. The effect of
monocentric species was positive and significant (Estimate = 0.55; 95% CI: 0.02–1.05),
implying nearly double the average crossover number relative to holocentric species [≈ 1.82
COs per chromosome].

To further evaluate the evolutionary dynamics underlying this difference, we applied
Ornstein–Uhlenbeck phylogenetic modeling. Across the seven models compared, the OUMV
model (allowing optima θ and trait variance σ² to vary between regimes) received
overwhelming support (AIC weight ≈ 1.0; Supplementary Table S3). This model estimated
distinct adaptive optima for holocentric and monocentric lineages (Figure 2B): θ₁ = 1.16
COs per chromosome (95% CI: 1.06–1.23) for holocentric species and θ₂ = 2.52 (95% CI:
2.26–2.79) for monocentric species. These results are consistent with the regression
analysis and demonstrate that monocentric insects maintain a significantly higher average
number of crossovers per chromosome than holocentric insects, in line with expectations
from their contrasting meiotic architectures.

Chromosome Number and Size as Predictors of Recombination Rate

Across the 73 analyzed insect species, recombination rate decreased with increasing mean
chromosome size but increased with chromosome number (Figure 3; Supplementary Table
S4). Models including mean chromosome size consistently outperformed those including
chromosome number in predicting genome-wide recombination rate (Supplementary Table
S5). In non-phylogenetic analyses, models with log(mean chromosome size) explained
substantially more variance (Bayes R^2 ≈ 0.61–0.64) than models with chromosome number
(Bayes R^2 ≈ 0.44–0.45). Phylogenetic models confirmed this pattern: those including
chromosome size achieved high explanatory power (Bayes R^2 ≈ 0.85–0.86) with marginal R^2
values around 0.22–0.29, indicating that chromosome size contributed meaningfully beyond
phylogenetic structure (Supplementary Table S5). By contrast, phylogenetic models
including chromosome number, despite similar conditional R^2 values (~0.92), had marginal
R^2 values near zero (0.02–0.03).

Model comparison using leave-one-out cross-validation (LOOIC/ELPD) further highlighted
this contrast (Supplementary Table S5). The best-fitting models were consistently those that
included chromosome size, while models containing chromosome number performed
substantially worse (ΔELPD ≈ –47 to –54). This indicates that the apparent explanatory
power of chromosome-number models was almost entirely attributable to phylogenetic
signal, whereas chromosome size remained an independent and robust predictor of
recombination rate even after accounting for shared ancestry.

For practical use, the best-fitting model (Gamma regression with a log link) can be
expressed as:

ln(expected recombination rate) = 4.19 – 1.04 × ln(mean chromosome size [Mbp]) + 0.80 ×
I(monocentric),

where *I(monocentric)* = 1 for monocentric species and 0 for holocentric species. Here, ln
denotes the natural logarithm.

Assessing Predictive Accuracy in Holocentric and Monocentric Species

In the best-supported non-phylogenetic model (recombination rate ~ log(mean chromosome
size) + chromosome type, see Supplementary Table S5), prediction accuracy differed
between chromosome types. Root mean square error (RMSE) and mean absolute error
(MAE) quantify the absolute size of prediction errors, while their relative versions (rel_RMSE
and rel_MAE) scale these errors by the observed variation in recombination rates. For
holocentric species, prediction errors were relatively low (RMSE = 1.13 [95% CI: 1.10–1.36];
MAE = 0.74 [0.67–1.04]; rel_RMSE = 0.52 [0.51–0.63]), indicating a close match between
observed and predicted values. In contrast, monocentric species exhibited larger errors
(RMSE = 3.89 [3.68–4.79]; MAE = 2.48 [2.38–2.91]; rel_RMSE = 0.73 [0.69–0.89]),
suggesting greater unexplained variance. These results demonstrate that recombination rate
is more accurately predicted in holocentric species, both in absolute and relative terms.
Importantly, the relative metrics show that the poorer fit in monocentrics cannot be explained
solely by their higher recombination rates; instead, recombination in monocentric lineages
appears intrinsically less predictable from chromosome size.

Evaluation of Chromosome Number and Mean Chromosome Size as Predictors in
Angiosperms

Across 157 angiosperm species with monocentric chromosomes, recombination rate
decreased with increasing mean chromosome size but increased with chromosome number
(Figure 4; Supplementary Table S6). As in the case of insects, chromosome size was by far
the stronger predictor (Supplementary Table S7).

In non-phylogenetic analyses, models including log(mean chromosome size) explained
substantially more variance (Bayes $R^2 \approx 0.49$–0.50) than those including chromosome
number (Bayes $R^2 \approx 0.07$). The estimated effect of size was strongly negative ($\beta \approx -0.89 \pm$
0.04), whereas the positive effect of chromosome number was small ($\beta \approx 0.14 \pm 0.06$).
Phylogenetic models confirmed this pattern: size-based models achieved higher explanatory
power (conditional $R^2 \approx 0.54$–0.55; marginal $R^2 \approx 0.48$), while number-based models had
much lower values (conditional $R^2 \approx 0.43$; marginal $R^2 \approx 0.05$), indicating that their apparent
fit largely reflected phylogenetic structure.

Model comparison using leave-one-out cross-validation further supported these results
(Supplementary Table S7). The best-fitting models were consistently those including mean
chromosome size, whereas models based on chromosome number performed substantially
worse (ΔELPD ≈ -90 to -120). Adding chromosome number or interaction terms did not
improve model performance.

Overall, these analyses show that mean chromosome size is a robust, independent predictor
of recombination rate in monocentric angiosperms, with smaller chromosomes associated
with higher recombination intensity, while the positive relationship with chromosome number
is weak and largely phylogenetically driven (Figure 4; Supplementary Tables S6 and S7).

405
406 *Predicting recombination in holocentric plants*
407
Building our results from flowering plants (Figure 4), we generated a prediction for
holocentric plants by taking the best monocentric plant model (with mean chromosome size
as a sole predictor) and shifting its intercept by the monocentric–holocentric offset estimated

in insects (Figure 3), while keeping the size slope identical (Figure 5). When we compared
this prediction with the two holocentric plants with maps, *Rhynchospora breviuscula* closely
matched the predicted curve, whereas *Carex scoparia* lay above it (Figure 5).

Discussion

The evolution of recombination rates plays a pivotal role in speciation, balancing the benefits
of adaptability with the risks of disrupting established genetic combinations (Ritz et al., 2017;
Stapley et al., 2017a). High recombination rates, while fostering evolutionary potential by
generating novel allele combinations for adaptation to diverse environments (Wang et al.,
2019) or for highly competitive communities (Bell, 1992), can also dismantle favorable co-
adapted gene complexes (Ritz et al., 2017). Conversely, lower recombination rates may
provide greater reproductive assurance by preserving existing adaptive combinations but at
the cost of reduced evolutionary flexibility. This trade-off is particularly evident when
considering the factors influencing recombination.

Here, we show that the number of COs per chromosome is predominantly higher and more
variable in monocentric insects compared to holocentric ones (Figure 2) which provides
support for a long-standing hypothesis based on karyological observations suggesting that
holocentric chromosomes typically exhibit more stringent constraints on chiasma formation
(White, 1973; Nokkala et al., 2004; Gokhman & Kuznetsova, 2006; Schvarzstein et al., 2010;
Metlers et al., 2012; Bureš et al., 2013; Cabral et al., 2014). Mechanistically, these
differences likely stem from the centromere organization in holocentric species, which
prevents the two-step loss of cohesion typical of monocentric meiosis and often results in
only one, or at most two, chiasmata per homologous pair (Schvarzstein et al., 2010;
Marques & Pedrosa-Harand, 2016; Hofstatter et al., 2021). However, the exact mechanism
underlying this limitation remains unknown. More than two chiasmata per bivalent appear to
be either absent or non-viable in holocentric species studied so far (Nokkala et al., 2004;
Hollis et al., 2020; Castellani et al., 2024), although the existence of holocentric lineages
capable of more than two viable crossovers per chromosome cannot be ruled out. In
contrast, monocentric chromosomes can achieve more COs per homolog pair (even over
10) and apparently do so across a range of species (Jones & Franklin, 2006; Fernandes et
al., 2018; Otto & Payseur, 2019; Lian et al., 2023).

In line with the tight regulation of CO number per chromosome through CO assurance and
interference (Hunter, 2015), both chromosome number and mean chromosome size initially
appeared to predict genome-wide recombination rate in insects (Figures 3A,C). However,
chromosome number consistently showed lower explanatory power, and across both
phylogenetic and non-phylogenetic models, mean chromosome size emerged as the
stronger predictor (Figure 3; Supplementary Table S5). Including both variables in additive or
interaction models did not improve fit (Supplementary Table S5), indicating that chromosome
number contributes little independent information beyond mean chromosome size. This likely
reflects that chromosome number captures coarse differences among higher taxonomic
groups, while within genera or families, where chromosome numbers vary little (as in
Lepidoptera; de Vos et al., 2020), variation in chromosome size more effectively explains
subtle differences in recombination rate. Moreover, polyploidization multiplies chromosome
number and likely has little effect on genome‑wide recombination rates (Ross-Ibarra, 2007;
Tiley & Burleigh, 2015; Stapley et al., 2017b), although autopolyploids may reduce the CO

number per chromosome to avoid multivalent formation (Bomblies et al., 2016). This may
further undermine the predictive accuracy of chromosome numbers. In contrast, mean
chromosome size effectively controls the multiplicative effects of (neo)polyploidy on
chromosome number (Bureš et al., 2024), although there is still unexplained variation left
(Figure 3A; Supplementary Table S5). The superior model fit observed for holocentric
species across both predictors further suggests that the strict limitations on CO numbers
(Figure 2)—imposed by the constraints of their kinetochore architecture—result in more
predictable recombination patterns than those seen in monocentric species. Together, these
results indicate that mean chromosome size is generally a better proxy for recombination
rate and that predictions are more accurate in holocentric than in monocentric systems.

While our findings support that chromosome number and especially mean chromosome size
are valid proxies for recombination rate, particularly in holocentric insects, it is important to
acknowledge the limitations of our study. Recombination varies along a genome and is
undoubtedly influenced by many factors beyond chromosome number and mean
chromosome size, such as structural variation, centromere position (metacentric vs.
acrocentric in monocentrics), transposable element activity, chromatin structure, and specific
meiotic genes (Johnston, 2024). Both sex chromosomes and sex determination systems
further differ across our studied insects, which is likely adding additional variation. To
estimate the impact of sex determination system and sex chromosomes, data for more
species is, however, required with better assemblies for sex chromosomes unique to the
heterogametic sex. In addition, the reliability of our approach inevitably depends on the
accuracy of genome size and chromosome number estimates. Variation in assembly
completeness or sequencing technology can affect genome size estimates and thus the
calculated mean chromosome size. These effects are likely minor compared to biological
variation but should be considered when applying the model across heterogeneous
datasets. Although the broad taxonomic sampling across insects is valuable, extrapolating
these findings to other monocentric and holocentric animals — and especially to plants —
should be done with caution.

On the other hand, our analyses encompassing 157 monocentric species of flowering plants
showed very similar results (Figure 4; Supplementary Tables S6, S7) to our findings in
insects (Figure 3, Supplementary Tables S4, S5), suggesting their general applicability.
Building on results from insects and monocentric plants, we predicted how genome-wide
recombination rate should relate to mean chromosome size in holocentric plants (Figure 5).
Notably, *Rhynchospora breviuscula* (Castellani et al., 2024) closely follows the predicted
relationship, whereas *Carex scoparia* (Escudero et al., 2018) deviates from it (Figure 5). This
deviation may reflect random variation, lineage-specific biological factors, lack of a reference
genome assembly, or systematic differences in recombination dynamics between holocentric
and monocentric plants. An additional possibility is that *Carex scoparia* lies outside the range
of chromosome sizes represented in our monocentric plant dataset; the extrapolated
prediction curve may therefore not accurately capture recombination behavior at such small
chromosome sizes. Moreover, while our insect analyses suggest a shared slope between
chromosome types (Figure 3; Supplementary Table S4), the relationship between
chromosome size and recombination rate could differ more fundamentally between
monocentric and holocentric plants. However, with data currently limited to just two
holocentric plant species, we cannot determine whether the observed deviation is

meaningful or incidental. Additional data from diverse holocentric plant lineages will be
essential to assess the robustness and generality of our predictions.

Our demonstration of the links among CO number, mean chromosome size, chromosome
type, and recombination rate provides a valuable framework for future large-scale
evolutionary studies aimed at unraveling the evolution of recombination and its role in
adaptation and speciation across the Tree of Life. In light of our findings, it is possible that
previous studies on the holocentric plant genus *Carex* (Bell, 1982; Escudero et al., 2012;
Escudero et al., 2013; Spalink et al., 2018; Márquez-Corro et al., 2021) and monocentric
angiosperms (Carta et al., 2018) may have suffered from reduced power by using
chromosome number rather than mean chromosome size as a proxy for recombination rate.
However, chromosome numbers are available for many more species because the mean
chromosome size requires knowledge of both chromosome number and genome size. For
instance, in flowering plants, chromosome numbers are available for ca. 60,000 species
(Rice et al., 2015) while mean chromosome size is available for only about 14,000 species
(Bureš et al., 2024). Therefore, continuing measurements of genome size (e.g., using flow
cytometry) and more initiatives like the Darwin Tree of Life (Darwin Tree of Life Project
Consortium, 2022) that generate chromosome-resolved reference genomes for thousands of
species will allow us to obtain mean chromosome size data and chromosome numbers at an
unprecedented scale.

While holocentric chromosomes appear to offer certain evolutionary advantages, such as the
ability to more readily tolerate increased rates of chromosomal fissions and fusions without
necessarily reducing fitness (Zedek & Bureš, 2019; Lucek et al., 2022; Zedek et al., 2022)—
allowing for rapid karyotypic evolution and potentially increased genetic diversity through
altered recombination rates and linkage groups—the prevalence of monocentric
chromosomes in the vast majority (80-85%) of eukaryotes (Márquez-Corro et al., 2018) and
comparable rates of chromosome number evolution in monocentric and holocentric insects
(Ruckman et al., 2020) suggest a different story. This apparent discrepancy–advantages
ascribed to holocentrics versus the predominance and comparable dynamics of
monocentrics–may be explained by the spatial separation of chromosome segregation and
recombination in monocentric chromosomes. In monocentric species, these two crucial
meiotic processes are functionally and physically distinct. Holocentric chromosomes, by
contrast, must manage both segregation and recombination across their diffuse centromere
structure. This constraint likely necessitates a reduction in the number of chiasmata per
chromosome to ensure proper segregation, as multiple chiasmata in holocentric species can
lead to meiotic errors (Nokkala et al., 2004; Hollis et al., 2020). However, some holocentric
insects and plants have evolved non-canonical meioses that decouple where crossovers
form from where spindle forces act (Melters et al. 2012; Heckmann et al. 2016; Pedrosa-
Harand & Marques 2016; Lukhtanov et al. 2018): inverted (post-reductional) meiosis, in
which sister chromatids bi-orient and separate in meiosis I and homologous chromatids
segregate in meiosis II, and telokinetic meiosis, which restricts kinetochore activity to one
chromosome end so homologs segregate reductionally in meiosis I; additionally, several
lineages limit crossovers and remodel meiotic chromosome domains to keep cohesion loss
spatially separated from kinetochore function (Pedrosa-Harand & Marques 2016). Our
findings, demonstrating a significantly higher number of chiasmata per homolog in
monocentric species compared to holocentric species (Figure 2), directly support this
explanation. The increased chiasma frequency in monocentrics likely contributes to their

greater evolutionary success by allowing for higher levels of genetic recombination. Thus,
while holocentric species may offer flexibility in karyotypic evolution, the inherent trade-off
between recombination rate specifically for chiasma formation and proper segregation, might
ultimately limit their overall evolutionary success compared to the more specialized and
efficient mechanisms found in monocentric chromosomes. In summary, our findings reveal a
fundamental difference in how insect recombination is regulated. While monocentric species
rely on increased CO rates, holocentric species appear to compensate through changes in
chromosome number. This stark contrast underscores centromere architecture's critical role
in shaping insect genomes' evolutionary trajectory.

Data availability statement

557 All the data used in this article are available in supplementary materials online on the
558 publisher's website.

Funding

561 This work was supported by the Czech Science Foundation, grant no. 24-11400S to FZ. KL
562 was supported by the Swiss National Science Foundation grants 202869 and 220868 as well
563 as the Fondation Pierre Mercier pour la science. ME was supported by the Spanish AEI-
564 MICINN grant DiversiChrom *PID2021-122715NB-I00.* AM was financially supported by the
565 Max Planck Society.

Author contributions
568 FZ conceived the study and performed analyses. FZ, PB, TLE, ME, KL, and AM wrote the
569 manuscript.

Conflicts of interests

572 None declared.

References

575 **Anderson LK, Salameh N, Bass HW, Harper LC, Cande WZ, Weber G, Stack SM. 2004.**
576 Integrating genetic linkage maps with pachytene chromosome structure in maize. *Genetics*
577 **166**: 1923–1933. doi: 10.1534/genetics.166.4.1923.
578
Backström N, Forstmeier W, Schielzeth H, Mellenius H, Nam K, Bolund E, Webster MT,
Ost T, Schneider M, Kempenaers B, Ellegren H. 2010. The recombination landscape of

the zebra finch *Taeniopygia guttata* genome. *Genome Research* **20**: 485–495. doi:
10.1101/gr.101410.109.

Barnes TM, Kohara Y, Coulson A, Hekimi S. 1995. Meiotic recombination, noncoding DNA
and genomic organization in *Caenorhabditis elegans. Genetics* **141**: 159–179. doi:
10.1093/genetics/141.1.159.

Beaulieu JM, Jhwueng DC, Boettiger C, O'Meara BC. 2012. Modeling stabilizing
selection: expanding the Ornstein-Uhlenbeck model of adaptive evolution. *Evolution* **66**:
2369–2383. doi: 10.1111/j.1558-5646.2012.01619.x.

Bell G. 1982. The masterpiece of nature: the evolution and genetics of sexuality. University
of California Press.

Blary A, Jenczewski E. 2019. Manipulation of crossover frequency and distribution for plant
breeding. *Theoretical and Applied Genetics* **132**: 575–592.

Bomblies K, Jones G, Franklin C, Zickler D, Kleckner N. 2016. The challenge of evolving
stable polyploidy: could an increase in "crossover interference distance" play a central role?
Chromosoma **125**: 287–300. doi: 10.1007/s00412-015-0571-4.

Brazier T, Glémin S. 2022. Diversity and determinants of recombination landscapes in
flowering plants. *PLoS Genetics* **18**: e1010141. doi: 10.1371/journal.pgen.1010141.

Bureš P, Elliott TL, Veselý P, Šmarda P, Forest F, Leitch IJ, Nic Lughadha E, Soto
Gomez M, Pironon S, Brown MJM, Šmerda J, Zedek F. 2024. The global distribution of
angiosperm genome size is shaped by climate. *New Phytologist* **242**: 744–759. doi:
10.1111/nph.19544.

Bureš P, Zedek F. 2014. Holokinetic drive: centromere drive in chromosomes without
centromeres. *Evolution* **68**: 2412–2420. doi: 10.1111/evo.12437.

Bureš P, Zedek F, Marková M. 2013. Holocentric chromosomes. In: *Plant genome*
diversity. Vol. 2., Vienna: Springer, 187-208.

Burt A. 2000. Perspective: sex, recombination, and the efficacy of selection--was Weismann
right? *Evolution* **54**: 337–351. doi: 10.1111/j.0014-3820.2000.tb00038.x.

Butlin RK. 2005. Recombination and speciation. *Molecular Ecology* **14**: 2621–2635. doi:
10.1111/j.1365-294X.2005.02617.x.

Bürkner P. 2017. brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal*
of Statistical Software **80**: 1–28. doi:10.18637/jss.v080.i01.

Bürkner P. 2018. Advanced Bayesian Multilevel Modeling with the R Package brms. *The R*
Journal **10**: 395–411. doi:10.32614/RJ-2018-017.

Bürkner P. 2021. Bayesian Item Response Modeling in R with brms and Stan. *Journal of*
Statistical Software **100**: 1–54. doi:10.18637/jss.v100.i05.
627
628 **Cabral G, Marques A, Schubert V, Pedrosa-Harand A, Schlögelhofer P. 2014.**
629 Chiasmatic and achiasmatic inverted meiosis of plants with holocentric chromosomes.
630 *Nature Communications* **5**: 5070. doi: 10.1038/ncomms6070.
631
632 **Carta A, Bedini G, Peruzzi L.** 2018. Unscrambling phylogenetic effects and ecological
633 determinants of chromosome number in major angiosperm clades. *Scientific Reports* **8**:
634 14258.
635
636 **Castellani M, Zhang M, Thangavel G, Mata-Sucre Y, Lux T, Campoy JA, Marek M,**
637 **Huettel B, Sun H, Mayer KFX, Schneeberger K, Marques A. 2024.** Meiotic recombination
638 dynamics in plants with repeat-based holocentromeres shed light on the primary drivers of
639 crossover patterning. *Nature Plants* **10**: 423–438. doi: 10.1038/s41477-024-01625-y.
640
641 **Chuang YC, Smith GR. 2023.** Meiotic crossover interference: Methods of analysis and
642 mechanisms of action. *Current Topics in Developmental Biology* **151**: 217–244. doi:
643 10.1016/bs.ctdb.2022.04.006.
644
645 **Crismani W, Mercier R. 2012.** Meiotic recombination: mechanisms and implications.
646 *Genetics* **190**: 1325–1334.
647
Cuacos MH, Franklin FC, Heckmann S. 2015. Atypical centromeres in plants—what they
can tell us. *Frontiers in Plant Science* **6**: 913.

Darwin Tree of Life Project Consortium. 2022. Sequence locally, think globally: The
Darwin Tree of Life Project. *Proc Natl Acad Sci U S A* **119**: e2115642118. doi:
10.1073/pnas.2115642118.

de Vos JM, Augustijnen H, Bätscher L, Lucek K. 2020. Speciation through chromosomal
fusion and fission in Lepidoptera. *Philosophical Transactions of the Royal Society B:*
Biological Sciences **375**: 20190539. doi: 10.1098/rstb.2019.0539.

Elliott TL, Zedek F, Barrett RL, Bruhl JJ, Escudero M, Hroudová Z, Joly S, Larridon I,
Luceño M, Márquez-Corro JI, Martín-Bravo S, Muasya AM, Šmarda P, Thomas WW,
Wilson KL, Bureš P. 2022. Chromosome size matters: genome evolution in the cyperid
clade. *Annals of Botany* **130**: 999–1014. doi: 10.1093/aob/mcac136.

Escudero M, Hipp AL, Hansen TF, Voje KL, Luceño M. 2012. Selection and inertia in the
evolution of holocentric chromosomes in sedges (*Carex*, Cyperaceae). *New Phytologist* **195**:
237–247. doi: 10.1111/j.1469-8137.2012.04137.x.

Escudero M, Maguilla E, Modesto L. 2013. Selection by climatic regime and neutral
evolutionary processes in holocentric chromosomes (*Carex* gr. *laevigata*: Cyperaceae): A
microevolutionary approach. *Perspectives in Plant Ecology, Evolution and Systematics* **15**:
118–129.

Escudero M, Hahn M, Hipp AL. 2018. RAD-seq linkage mapping and patterns of
segregation distortion in sedges: meiosis as a driver of karyotypic evolution in organisms
with holocentric chromosomes. *Journal of Evolutionary Biology* **31**: 833–843. doi:
10.1111/jeb.13267.

Fernandes JB, Séguéla-Arnaud M, Larchevêque C, Lloyd AH, Mercier R. 2018.
Unleashing meiotic crossovers in hybrid plants. *Proc Natl Acad Sci U S A* **115**: 2431–2436.
doi: 10.1073/pnas.1713078114.
674
675   **Forest F. 2023.** Species-level phylogenetic trees of all angiosperm species (100 trees).
676   https://zenodo.org/record/7600341
677
678   **Gokhman VE, Kuznetsova VG. 2006.** Comparative insect karyology: current state and
679   applications. *Entomological Review* **86**: 352–368.
680
681   **Gregory TR. 2025.** Animal Genome Size Database. http://www.genomesize.com.
682
683   **Haenel Q, Laurentino TG, Roesti M, Berner D. 2018.** Meta-analysis of chromosome-scale
684   crossover rate variation in eukaryotes and its significance to evolutionary genomics.
685   *Molecular Ecology* **27**: 2477–2497. doi: 10.1111/mec.14699.
686
687   **Heckmann S, Jankowska M, Schubert V, Kumke K, Ma W, Houben A. 2014.** Alternative
688   meiotic chromatid segregation in the holocentric plant *Luzula elegans*. *Nature*
689   *Communications* **5**: 4979.
690
691   **Heckmann S, Macas J, Kumke K, Fuchs J, Schubert V, Ma L, Novák P, Neumann P,**
692   **Taudien S, Platzer M, Houben A. 2013.** The holocentric species *Luzula elegans* shows
693   interplay between centromere and large-scale genome organization. *The Plant Journal* **73**:
694   555–565. doi: 10.1111/tpj.12054.
695
696   **Hofstatter PG, Thangavel G, Castellani M, Marques A. 2021.** Meiosis Progression and
697   Recombination in Holocentric Plants: What Is Known? *Frontiers in Plant Science* **12**:
698   658296. doi: 10.3389/fpls.2021.658296.
699
Hofstatter PG, Thangavel G, Lux T, Neumann P, Vondrak T, Novak P, Zhang M, Costa
L, Castellani M, Scott A, Toegelová H, Fuchs J, Mata-Sucre Y, Dias Y, Vanzela ALL,
Huettel B, Almeida CCS, Šimková H, Souza G, Pedrosa-Harand A, Macas J, Mayer
KFX, Houben A, Marques A. 2022. Repeat-based holocentromeres influence genome
architecture and karyotype evolution. *Cell* **185**: 3153–3168.e18. doi:
10.1016/j.cell.2022.06.045.
706
707   **Hollis JA, Glover ML, Schlientz AJ, Cahoon CK, Bowerman B, Wignall SM, Libuda DE.**
708   **2020.** Excess crossovers impede faithful meiotic chromosome segregation in *C. elegans*.
709   *PLoS Genetics* **16**: e1009001. doi: 10.1371/journal.pgen.1009001.
710
711   **Hunter N. 2015.** Meiotic Recombination: The Essence of Heredity. *Cold Spring Harbor*
712   *Perspectives in Biology* **7**: a016618. doi: 10.1101/cshperspect.a016618.
713

714 **Jankowska M, Fuchs J, Klocke E, Fojtová M, Polanská P, Fajkus J, Schubert V,**
715 **Houben A. 2015.** Holokinetic centromeres and efficient telomere healing enable rapid
716 karyotype evolution. *Chromosoma* **124**: 519–528. doi: 10.1007/s00412-015-0524-y.
717
718 **Jones GH, Franklin FCH. 2006.** Meiotic crossing-over: obligation and interference. *Cell*
719 **126**: 246–248.
720
721 **Jensen-Seaman MI, Furey TS, Payseur BA, Lu Y, Roskin KM, Chen CF, Thomas MA,**
722 **Haussler D, Jacob HJ. 2004.** Comparative recombination rates in the rat, mouse, and
723 human genomes. *Genome Research* **14**: 528–538. doi: 10.1101/gr.1970304.
724
725 **Johnston SE. 2024.** Understanding the Genetic Basis of Variation in Meiotic
726 Recombination: Past, Present, and Future. *Molecular Biology and Evolution* **41**: msae112.
727 doi: 10.1093/molbev/msae112.
728
729 **Kaback DB, Guacci V, Barber D, Mahon JW. 1992.** Chromosome size-dependent control
730 of meiotic recombination. *Science* **256**: 228–232. doi: 10.1126/science.1566070.
731
732 **Kent TV, Uzunović J, Wright SI. 2017.** Coevolution between transposable elements and
733 recombination. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**:
734 20160458. doi: 10.1098/rstb.2016.0458.
735
736 **Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, Stecher G, Hedges**
737 **SB. 2022.** TimeTree 5: An Expanded Resource for Species Divergence Times. Molecular
738 Biology and Evolution **39**: msac174. doi: 10.1093/molbev/msac174.
739
740 **Leitch AR, Ma L, Dodsworth S, Fuchs J, Houben A, Leitch IJ. 2023.** The Role of
741 Chromatin Modifications in the Evolution of Giant Plant Genomes. *Plants (Basel)* **12**: 2159.
742 doi: 10.3390/plants12112159.
743
744 **Lian Q, Maestroni L, Gaudin M, Llorente B, Mercier R. 2023.** Meiotic recombination is
745 confirmed to be unusually high in the fission yeast *Schizosaccharomyces pombe*. *iScience*
746 **26**: 107614. doi: 10.1016/j.isci.2023.107614.
747
748 **Lucek K, Augustijnen H, Escudero M. 2022.** A holocentric twist to chromosomal
749 speciation? *Trends in Ecology and Evolution* **37**: 655–662. doi: 10.1016/j.tree.2022.04.002.
750
751 **Lukhtanov VA, Dincă V, Friberg M, Šíchová J, Olofsson M, Vila R, Marec F, Wiklund C.**
752 **2018.** Versatility of multivalent orientation, inverted meiosis, and rescued fitness in
753 holocentric chromosomal hybrids. *Proc Natl Acad Sci U S A* **115**: E9610–E9619. doi:
754 10.1073/pnas.1802610115.
755
756 **Lynch M, Bobay LM, Catania F, Gout JF, Rho M. 2011.** The repatterning of eukaryotic
757 genomes by random genetic drift. *Annual Review of Genomics and Human Genetics* **12**:
758 347–366. doi: 10.1146/annurev-genom-082410-101412.
759

Mackintosh A, Laetsch DR, Hayward A, Charlesworth B, Waterfall M, Vila R, Lohse K. 2019. The determinants of genetic diversity in butterflies. *Nature Communications* **10**: 3466. doi: 10.1038/s41467-019-11308-4.

Mancera E, Bourgon R, Brozzi A, Huber W, Steinmetz LM. 2008. High-resolution mapping of meiotic crossovers and non-crossovers in yeast. *Nature* **454**: 479–485. doi: 10.1038/nature07135.

Mandrioli M, Manicardi GC. 2012. Unlocking holocentric chromosomes: new perspectives from comparative and functional genomics? *Current Genomics* **13**: 343–349. doi: 10.2174/138920212801619250.

Mandrioli M, Manicardi GC. 2020. Holocentric chromosomes. *PLoS Genetics* **16**: e1008918.

Marques A, Drinnenberg IA. 2025. Same but different: Centromere regulations in holocentric insects and plants. *Current Opinion in Cell Biology* **93**: 102484. doi: 10.1016/j.ceb.2025.102484.

Marques A, Pedrosa-Harand A. 2016. Holocentromere identity: from the typical mitotic linear structure to the great plasticity of meiotic holocentromeres. *Chromosoma* **125**: 669–681.

Márquez-Corro JI, Escudero M & Luceño M. 2018. Do holocentric chromosomes represent an evolutionary advantage? A study of paired analyses of diversification rates of lineages with holocentric chromosomes and their monocentric closest relatives. *Chromosome Research* **26:** 139-152.

Márquez-Corro JI, Martín-Bravo S, Jiménez-Mejías P, Hipp AL, Spalink D, Naczi RF, Roalson EH, Luceño M, Escudero M. 2021. Macroevolutionary insights into sedges (*Carex*: Cyperaceae): The effects of rapid chromosome number evolution on lineage diversification. *Journal of Systematics and Evolution* **59**: 776–790. doi: 10.1111/jse.12730.

Martin SH, Davey JW, Salazar C, Jiggins CD. 2019. Recombination rate variation shapes barriers to introgression across butterfly genomes. *PLoS Biology* **17**: e2006288. doi: 10.1371/journal.pbio.2006288.

Mata-Sucre Y, Krátká M, Oliveira L, Neumann P, Macas J, Schubert V, Huettel B, Kejnovský E, Houben A, Pedrosa-Harand A, Souza G, Marques A. 2024. Repeat-based holocentromeres of the woodrush Luzula sylvatica reveal insights into the evolutionary transition to holocentricity. *Nature Communications* **15**: 9565. doi: 10.1038/s41467-024-53944-5.

Melters DP, Paliulis LV, Korf IF, Chan SW. 2012. Holocentric chromosomes: convergent evolution, meiotic adaptations, and genomic analysis. *Chromosome Research* **20**: 579–593.

Mercier R, Mézard C, Jenczewski E, Macaisne N, Grelon M. 2015. The molecular biology of meiosis in plants. *Annual Review of Plant Biology* **66**: 297–327.

Näsvall K, Boman J, Höök L, Vila R, Wiklund C, Backström N. 2023. Nascent evolution
of recombination rate differences as a consequence of chromosomal rearrangements. *PLoS*
Genetics **19**: e1010717. doi: 10.1371/journal.pgen.1010717.

Nokkala S, Kuznetsova VG, Maryanska-Nadachowska A, Nokkala C. 2004. Holocentric
chromosomes in meiosis. I. Restriction of the number of chiasmata in bivalents.
Chromosome Research **12**: 733–739.

Nordenskiöld H. 1963. A study of meiosis in the progeny of X-irradiated *Luzula purpurea*.
Hereditas **49**: 33–47.

Otto SP, Payseur BA. 2019. Crossover interference: shedding light on the evolution of
recombination. *Annual Review of Genetics* **53**: 19–44.

Palahí I Torres A, Höök L, Näsvall K, Shipilina D, Wiklund C, Vila R, Pruisscher P,
Backström N. 2023. The fine-scale recombination rate variation and associations with
genomic features in a butterfly. *Genome Research* **33**: 810–823. doi:
10.1101/gr.277414.122.

Paradis E, Schliep K .2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**: 526–528. doi:10.1093/bioinformatics/bty633.

Pessia E, Popa A, Mousset S, Rezvoy C, Duret L, Marais GA. 2012. Evidence for
widespread GC-biased gene conversion in eukaryotes. *Genome Biology and Evolution* **4**:
675–682. doi: 10.1093/gbe/evs052.

Rice A, Glick G, Abadi S, Einhorn M, Kopelman NM, Salman-Minkov A, Mayzel J, Chay
O, Mayrose I. 2015. The chromosome counts database (CCDB) – a community resource of
plant chromosome numbers. *New Phytologist* **206**: 16–26. doi: 10.1111/nph.13191.

Ritz KR, Noor MA, Singh ND. 2017. Variation in recombination rate: adaptive or not?.
Trends in Genetics **33**: 364–374.

Ross-Ibarra J. 2007. Genome size and recombination in angiosperms: a second look.
Journal of Evolutionary Biology **20**: 800–806. doi: 10.1111/j.1420-9101.2006.01275.x.

Ruckman SN, Jonika MM, Casola C, Blackmon H. 2020. Chromosome number evolves at
equal rates in holocentric and monocentric clades. *PLoS Genetics* **16**: e1009076. doi:
10.1371/journal.pgen.1009076.

Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, Connor R, Funk K,
Kelly C, Kim S, Madej T, Marchler-Bauer A, Lanczycki C, Lathrop S, Lu Z, Thibaud-
Nissen F, Murphy T, Phan L, Skripchenko Y, Tse T, Wang J, Williams R, Trawick BW,
Pruitt KD, Sherry ST. 2022. Database resources of the National Center for Biotechnology
Information. *Nucleic Acids Research* **50**: D20–D26. doi: 10.1093/nar/gkab1112.

Shipilina D, Näsvall K, Höök L, Vila R, Talavera G, Backström N. 2022. Linkage mapping
and genome annotation give novel insights into gene family expansions and regional
recombination rate variation in the painted lady (Vanessa cardui) butterfly. *Genomics* **114**:
110481. doi: 10.1016/j.ygeno.2022.110481.

Spalink D, Pender J, Escudero M, Hipp AL, Roalson EH, Starr JR, Waterway MJ, Bohs
L, Sytsma KJ. 2018. The spatial structure of phylogenetic and functional diversity in the
United States and Canada: An example using the sedge family (Cyperaceae). *Journal of*
Systematics and Evolution **56**: 449–465. doi: 10.1111/jse.12423.

Stapley J, Feulner PG, Johnston SE, Santure AW, Smadja CM. 2017a. Recombination:
the good, the bad and the variable. *Philosophical Transactions of the Royal Society B:*
Biological Sciences **372**: 20170279.

Stapley J, Feulner PG, Johnston SE, Santure AW, Smadja CM. 2017b. Variation in
recombination frequency and distribution across eukaryotes: patterns and processes.
Philosophical Transactions of the Royal Society B: Biological Sciences **372**: 20160455.

Schvarzstein M, Wignall SM, Villeneuve AM. 2010. Coordinating cohesion, co-orientation,
and congression during meiosis: lessons from holocentric chromosomes. *Genes and*
Development **24**: 219–228. doi: 10.1101/gad.1863610.

Tigano A, Khan R, Omer AD, Weisz D, Dudchenko O, Multani AS, Pathak S, Behringer
RR, Aiden EL, Fisher H, MacManes MD. 2022. Chromosome size affects sequence
divergence between species through the interplay of recombination and selection. *Evolution*
76: 782–798. doi: 10.1111/evo.14467.

Tiley GP, Burleigh JG. 2015. The relationship of recombination rate, genome structure, and
patterns of molecular evolution across angiosperms. *BMC Evolutionary Biology* **15**: 194. doi:
10.1186/s12862-015-0473-3.

Viera A, Page J, Rufas JS. 2009. Inverted meiosis: the true bugs as a model to study.
Meiosis **5**: 137–156.

Wang S, Veller C, Sun F, Ruiz-Herrera A, Shang Y, Liu H, Zickler D, Chen Z, Kleckner
N, Zhang L. 2019. Per-Nucleus Crossover Covariation and Implications for Evolution. *Cell*
177: 326–338.e16. doi: 10.1016/j.cell.2019.02.021.

White MJD. 1973. *Animal cytology and evolution.* Cambridge University Press, Cambridge,
UK
865
866 **Zedek F, Bureš P. 2019.** Pest Arthropods with Holocentric Chromosomes are More
867 Resistant to Sterilizing Ionizing Radiation. *Radiation Research* **191**: 255–261. doi:
868 10.1667/RR15208.1.
869
Zedek F, Plačková K, Veselý P, Šmerda J, Šmarda P, Horová L, Bureš P. 2020.
Endopolyploidy is a common response to UV-B stress in natural plant populations, but its
magnitude may be affected by chromosome type. *Annals of Botany* **126**: 883–889. doi:
10.1093/aob/mcaa109.

Zedek F, Veselý P, Tichý L, Elliott TL, Garbolino E, de Ruffray P, Bureš P. 2022.
Holocentric plants are more competitive under higher UV-B doses. *New Phytologist* **233**: 15–
21. doi: 10.1111/nph.17750.

Zhang L, Reifová R, Halenková Z, Gompert Z. 2021. How Important Are Structural
Variants for Speciation? *Genes (Basel)* **12**:1084. doi: 10.3390/genes12071084.

 Figure legends

Figure 1. Phylogenetic tree of 33 holocentric (blue) and 40 monocentric (red) species
included in our study. Holocentric species include butterflies (Lepidoptera), true bugs
(Hemiptera), earwigs (Dermaptera), and dragonflies and damselflies (Odonata). Monocentric
species include flies (Diptera), beetles (Coleoptera), sawflies, wasps, bees, and ants
(Hymenoptera), termites (Blattodea), and crickets (Orthoptera). The panels with barplots
show from right to left haploid autosome numbers (n, log10-transformed), Mean autosome
size (Mbp, log10-transformed), autosomal recombination rate (cM/Mb), and number of
crossovers per autosome per meiosis. The arrows display putative transitions to holocentric
chromosomes (Melters et al., 2021; Marques and Drinnenberg, 2025). The phylogenetic tree
is based on the TimeTree resource (Kumar et al., 2022). If a species was not available from
TimeTree, we used the tip of its most closely related relative, which was renamed, or added
manually based on published phylogenies.

Figure 2. Number of crossovers (COs) in holocentric and monocentric insects. A - Observed
COs per chromosome in holocentric (blue) and monocentric (red) species. Individual data
points are jittered to display variation, and the overlaid symbols with error bars represent the
group mean and 95% prediction credible intervals estimated from the model. B - Bootstrap
density distributions (based on 100 replicates) of the evolutionary optimum (θ) for the
number of COs estimated under the OUMVA model. The density for holocentric lineages
(blue) and monocentric lineages (red) illustrate the uncertainty and differences in the
adaptive optima between chromosome types.

Figure 3. (A) Recombination rate (cM/Mbp) vs. mean chromosome size without accounting
for phylogeny. (B) The same relationship when phylogenetic relatedness is considered. (C)
Recombination rate vs. haploid autosome number (n) without accounting for phylogeny. (D)
Recombination rate vs. haploid autosome number when phylogeny is considered. Solid lines
indicate model predictions and shaded ribbons represent 95% prediction credible intervals.
Holocentric species are shown in blue, monocentric in red.

Figure 4. Supplementary Figure S1. Relationships between chromosome number, mean
chromosome size, and recombination rate in flowering plants with monocentric
chromosomes. (A) Relationship between recombination rate (cM/Mbp) and mean
chromosome size. (B) Relationship between recombination rate (cM/Mbp) and mean
chromosome size when phylogenetic relatedness is considered. (C) Relationship between
recombination rate (cM/Mbp) and haploid chromosome number. (D) Relationship between
recombination rate (cM/Mbp) and haploid chromosome number when phylogenetic
relatedness is considered. Solid lines indicate model predictions and shaded ribbons
represent 95% prediction credible intervals.

Figure 5. Relationship between mean chromosome size (Mbp) and recombination rate
(cM/Mbp) in plants. Light green points represent empirical data for monocentric plants. The
light green solid line and shaded area indicate the regression fit and 95% confidence
intervals, respectively, for monocentric plants. The dark green dashed line and shaded area
represent the predicted regression and 95% confidence intervals for holocentric plants,
derived by adjusting the monocentric plant intercept using the intercept difference between
monocentric and holocentric insect species. Black triangles show the only two plant species

with holocentric chromosomes for which both recombination rate and mean chromosome
size are known: *Carex scoparia* (Escudero et al., 2018) and *Rhynchospora breviuscula*
(Castellani et al., 2024).

Supplementary Figure S1. Associations between genetic map length and average marker
distance (cM per marker). No significant correlation was detected (Spearman $\rho \approx 0$, p >
0.05) and a linear model controlling for chromosome type likewise showed no effect ($\beta \approx 0$, p
> 0.05). This confirms that estimates of map length are not biased by marker density.

Recombination rate vs mean chromosome size in monocentric plants with a predicted holocentric shift