Chromosome Size as a Universal Predictor of Recombination Rate: Insights from
Holocentric and Monocentric Systems

Authors: František Zedek[1], Petr Bureš[1], Tammy L. Elliott[1,2], Marcial Escudero[3], Kay Lucek[4],
André Marques[5]

[1]Department of Botany and Zoology, Faculty of Science, Masaryk University, Kotlářská 2,
61137 Brno, Czech Republic

[2]Bolus Herbarium, Department of Biological Sciences, University of Cape Town, Private Bag
X3, Rondebosch, 7701, Cape Town, South Africa

[3]Department of Plant Biology and Ecology, Faculty of Biology, University of Seville, Reina
Mercedes 6, 41012 Seville, Spain

[4]Biodiversity Genomics Laboratory, Institute of Biology, University of Neuchâtel, 11 rue
Emile-Argand, 2000 Neuchâtel, Switzerland

[5]Department of Chromosome Biology, Max Planck Institute for Plant Breeding Research,
Cologne, Germany

Corresponding author: František Zedek, email: fzedek@gmail.com

Abstract

Recombination is a fundamental evolutionary process essential for generating genetic diversity, facilitating adaptation, and driving speciation. However, direct measurement of recombination rate remains challenging, as standard methods—such as chiasma counts or genetic linkage maps—are labor-intensive and often infeasible for non-model species. In this study, we identify chromosome number and mean chromosome size as practical proxies for genome-wide recombination rate by analyzing genetic map data from 69 insect species and supplementary analyses of 157 monocentric flowering plants. Using both ordinary and phylogenetically informed Bayesian regression models, we show that recombination rate increases with chromosome number and decreases with mean chromosome size. Crucially, mean chromosome size is a significantly better predictor, particularly in holocentric species. This insight enables recombination rate estimation in thousands of species with known genome and chromosome sizes, thereby allowing hypothesis testing at scales previously unattainable. We also confirm the long-standing hypothesis that monocentric species have nearly twice as many crossovers per chromosome as holocentric species, reflecting structural constraints imposed by diffuse centromeres. Building on these results, we present predictive models applicable to poorly studied holocentric plants. Overall, our study highlights the pivotal role of chromosome architecture in recombination evolution and provides an accessible framework for evolutionary genomic research across diverse lineages.

Keywords: Recombination rate, Chromosome size, Chromosome number, Crossover frequency, Holocentric chromosomes, Monocentric chromosomes, Evolutionary genomics, Meiotic recombination, Phylogenetic comparative methods, Genome architecture

Article summary: Why do some species have more genetic shuffling than others? We investigated 69 insect species with detailed genetic maps and found that mean chromosome size is the best predictor of genome-wide recombination rate, not the number of chromosomes. This means we can now estimate recombination rates even in species where genetic maps are unavailable, using only genome size and chromosome number. We also show that chromosome structure matters—species with a single centromere per chromosome (monocentric) have nearly double the recombination compared to those with diffuse centromeres (holocentric). These insights open new doors for studying evolution across many organisms.

Introduction

The key evolutionary implications of recombination are twofold: recombination can facilitate adaptation across environments by generating and maintaining genetic variation, or it can break up co-adapted allele combinations, potentially disrupting current optimal gene combinations (Ritz et al., 2017; Stapley et al., 2017a). Recombination also plays a crucial role in speciation as the formation of new species often involves a progressive reduction in genetic exchange in specific genomic regions that act as barriers to gene flow (Butlin, 2005). In this context, a lot of attention has been given to the evolutionary advantage of sex, which promotes recombination. However, the reason why average genome-wide recombination also varies between sexually reproducing organisms is less understood (Stapley et al., 2017a). In sexual eukaryotes, genome-wide recombination rate is determined by the number of crossovers (COs; i.e., chiasmata) per meiosis and by random meiotic segregation of homologs (Stapley et al., 2017b; Otto & Payseur, 2019; Hofstatter et al., 2021). Because estimating recombination rates with standard approaches, such as counting chiasmata or establishing linkage maps (Stapley et al. 2017b), demands time-consuming analysis and is typically impractical for non-model species, it remains challenging to investigate recombination rate-related hypotheses at a macroevolutionary scale across species and environments.

Since at least one CO per chromosome is usually required for proper segregation during meiosis (a phenomenon known as CO assurance; Hunter, 2015) and the formation of additional COss is tightly regulated by CO interference (a mechanism that prevents COs from occurring too closely together; Hunter, 2015), the number of COs per chromosome is maintained within a narrow range (Lian et al., 2023). Therefore, chromosome number is expected to predict genome-wide recombination rate across species (Stapley et al., 2017b). Having more chromosomes also increases the potential for diverse combinations during the assortment of chromosomes in gametes. However, variation in genome size can obscure this relationship.

All else being equal, the necessity of at least one CO per chromosome results in proportionally higher recombination rates in shorter chromosomes (Lynch et al., 2011; Tigano et al., 2022). Heterochromatin is moreover associated with reduced recombination, for instance, in pericentromeric regions, which often accumulate transposable elements and display suppressed CO activity (Kent et al., 2017). Consequently, larger chromosomes, which tend to contain proportionally larger heterochromatic regions (Leitch et al., 2023), may experience a lower rate of recombination per unit length compared to smaller chromosomes. This dynamic could explain the commonly observed negative association between recombination rate and chromosome size of individual chromosomes inside a single karyotype (Kaback et al., 1992; Jensen-Seaman et al., 2004; Backström et al., 2010; Martin et al., 2019; Brazier & Glémin, 2022; Shipilina et al., 2022; Näsvall et al., 2023; Palahí I Torres et al., 2023), which appears to translate to a negative relationship between mean chromosome size and average genome-wide recombination rate (Lynch et al., 2011; Brazier & Glémin, 2022). Thus, chromosome counts and mean chromosome size can potentially be used as proxies for genome-wide recombination rate comparison across a large number of species.

In eukaryotes, two predominant types of chromosomes have evolved that differ regarding their centromere organization, i.e. monocentric and holocentric chromosomes. Importantly, the accuracy with which chromosome number and mean chromosome size predict genome-wide recombination rate is expected to differ between these two types of chromosome organization. For species with monocentric chromosomes, where kinetochores and chromatid cohesion are restricted to a single centromeric region, the number of COs per chromosome mostly varies between one to four (Crismani & Mercier, 2012; Mercier et al., 2015; Blary & Jenczewski, 2019; Otto & Payseur, 2019; Brazier & Glémin, 2022; Lian et al., 2023), but higher numbers have also been reported (Jones & Franklin, 2006; Mancera et al., 2008; Brazier & Glémin, 2022; Lian et al., 2023). In contrast, species with holocentric chromosomes, where kinetochores are maintained along the entire chromosome length (Bureš et al., 2013; Mandrioli & Manicardi, 2020; Hofstatter et al., 2021) cannot rely on the two-step loss of cohesion during meiosis and restrict the CO frequency per homolog to one or two chiasmata at maximum (Schvarzstein et al., 2010; Metlers et al., 2012; Cuacos et al., 2015; Marques & Pedrosa-Harand, 2016; Mandrioli & Manicardi, 2020; Castellani et al., 2024). Such decreased recombination has been shown for both holocentric plants (e.g., Nordenskiöld, 1962; Cabral et al., 2014; Heckmann et al., 2014; Hofstatter et al., 2021; Castellani et al., 2024) and animals (e.g., Barnes et al., 1995; Nokkala et al., 2004; Viera et al., 2009; Lukhtanov et al., 2018). Another difference between holocentric and monocentric species is the composition of chromatin. In holocentric species, chromatin is often not compartmentalized and shows no obvious euchromatic or heterochromatic clusters as in monocentric species (Mandrioli & Manicardi, 2012; Heckmann et al., 2013; Hofstatter et al., 2022; Mata-Sucre et al., 2024) with homogeneous TEs distributions along chromosomes (Hofstatter et al. 2022; Shipilina et al., 2022; Mata-Sucre et al., 2024; Castellani et al., 2024). This could potentially result in a more uniform – yet still telomere-biased – distribution of recombination rates along holocentric chromosomes compared to monocentric (Haenel et al., 2018; Palahí i Torres et al., 2022; Hofstatter et al., 2022; Shipilina et al., 2022; Castellani et al., 2024). Furthermore, due to their extended kinetochore and the attachment of spindle microtubules along their entire length, holocentric species are more likely to tolerate chromosomal fissions and fusions that have negligible fitness costs (Bureš et al., 2013; Jankowska et al., 2015; Zedek & Bureš, 2019; Mandrioli & Manicardi, 2020; Lucek et al., 2022). Therefore, holocentric species are expected to more readily change chromosome numbers and size and, as a consequence, recombination rates in response to selection compared to monocentric species (Bell, 1982; Escudero et al., 2012; Bureš et al., 2013; Escudero et al. 2013; Bureš & Zedek, 2014). Thus, a higher recombination rate in monocentric species is usually expected to be reached by increasing the number of COs per chromosome (Burt, 2000; Stapley et al., 2017b), whereas in holocentric species, a higher recombination rate is predicted to primarily result from increasing chromosome numbers (and thereby decreasing their size).

Although chromosome numbers and mean chromosome size have been used as proxies for genome-wide recombination rate inference in both monocentric (Pessia et al., 2012; Carta et al., 2018) and holocentric species (Bell, 1982; Escudero et al., 2012; Escudero et al., 2013; Spalink et al., 2018; Mackintosh et al., 2019; Márquez- Corro et al., 2021; Elliott et al., 2022), the assumption that these two genomic traits accurately represent recombination rates should be critically assessed. A system encompassing multiple holocentric and monocentric lineages, for which chromosome numbers, genome sizes, and linkage maps or properly counted chiasmata are available to estimate genome-wide recombination rates is

required to address these questions. Insects may constitute such a group (Figure 1), where
holocentric chromosomes have independently evolved in the orders Lepidoptera,
Trichoptera, Hemiptera, Thysanoptera, Psocodea, Zoraptera, Dermaptera, and Odonata,
whereas monocentric chromosomes are more broadly found, for example in Diptera,
Coleoptera, Hymenoptera, Orthoptera, or Blattodea (Melters et al., 2012; Zedek & Bureš,
2019; Mandrioli & Manicardi, 2020; Marques & Drinnenberg, 2025). Moreover, chromosome
number and size, genome size information, and linkage maps are available for a reasonable
amount of species of both holocentric and monocentric insect orders.

By analyzing recombination data from insect species with well-characterized genetic maps
(or counted chiasmata), we aimed to assess the suitability of chromosome numbers and
mean chromosome size as predictors of genome-wide recombination rate and test the
hypothesis that these predictors are more accurate in holocentric than in monocentric
species. Specifically, we expect that (1) the average number of COs per chromosome is
higher in monocentric species than in holocentric ones, which is the key assumption
stemming from the structural differences between holocentric and monocentric
chromosomes (see above), and (2) chromosome number and size predict genome-wide
recombination rate better in holocentric than monocentric species. To assess the broader
applicability of our results beyond insects, we evaluated the accuracy of chromosome
number and mean chromosome size as predictors of recombination rate in flowering plants
with monocentric chromosomes. Holocentric species could not be meaningfully included, as
only two holocentric plant species currently have established genetic maps (Escudero et al.,
2018; Castellani et al., 2024), but we made a prediction of holocentric plants based on the
results we obtained from insects and monocentric plants.
182

Figure 1. Phylogenetic tree of 30 holocentric and 39 monocentric species included in our study. Holocentric species include
butterflies (Lepidoptera), true bugs (Hemiptera), earwigs (Dermaptera), and dragonflies and damselflies (Odonata).
Monocentric species include flies (Diptera), beetles (Coleoptera), sawflies, wasps, bees, and ants (Hymenoptera), termites
(Blattodea), and crickets (Orthoptera).

Materials and Methods

Data Collection

We compiled genetic map length (67 species) and chiasmata count (2 species) data from
the literature for a range of insect species, spanning both holocentric and monocentric
species (Supplementary Table S1). Genome size and chromosome count information were
obtained from the NCBI Genome database (Sayers et al., 2022), the Animal Genome Size
Database (Gregory, 2025), and additional publications (Supplementary Table S1). We
defined the recombination rate as the haploid genetic map length divided by the 1C genome
size (Stapley et al., 2017b). In the case of the holocentric odonata *Aeshna subarctica* and
the earwig *Forficula auricularia*, we calculated the genetic map length from the number of
chiasmata as $50 \times$ average number of chiasmata per meiosis because 50 centiMorgans
(cM) correspond to an average of one CO per meiosis (Anderson et al., 2004; Chuang &
Smith, 2023). We estimated the mean number of COs per chromosome by dividing the
genetic map length by 50 divided by the chromosome count. The final dataset included the
haploid chromosome number (n), mean chromosome size (Mbp), the recombination rate
(cM/Mbp), the mean number of COs per chromosome, and the chromosome type

(holocentric or monocentric) (Supplementary Table S1). The phylogenetic tree (Figure 1)
used in the following analyses was based on the TimeTree (Kumar et al., 2022). If a species
was not available from TimeTree, we used the tip of its most closely related relative, which
was renamed, or added manually based on published phylogenies.

Recombination rate predicted by chromosome number and mean chromosome size

Recombination rate (cM/Mb) and the average number of COs per chromosome (COperChr)
were analyzed within a Bayesian framework and through comparative phylogenetic methods
to account for the data's distributional properties and the non-independence among species
employing the *brms* R package (Bürkner, 2017; 2018; 2021).

Because recombination rates are strictly positive and right- skewed, different modeling
strategies were employed depending on the focal chromosomal predictor. For chromosome
number as a predictor, we observed that heteroscedasticity persisted even after log-
transforming recombination rate. Therefore, recombination rate was modeled on its original
scale using a Gamma regression with a log link. This approach accommodates the positive
nature of the data and the residual heteroscedasticity, permitting a multiplicative
interpretation of regression coefficients. A model with interaction effects was initially
specified as *recombination rate~chromosome number×chromosome type*, where
chromosome type distinguishes holocentric from monocentric species. If the 95% credible
interval for the interaction term overlapped zero, a simpler additive formulation was adopted.

For analyses involving chromosome size as a predictor, both recombination rate and
chromosome size were log-transformed. This transformation effectively stabilized the
variance and linearized the multiplicative relationship, allowing us to apply a Gaussian model
to the log–log data: *log(recombination rate)~log(chromosome size)×chromosome type*. As
before, evidence for interaction was evaluated via the 95% credible intervals, with additive
models used if the interaction term was not supported.

For both sets of analyses, models were estimated using four Markov chains with total

iterations ranging from 4,000 to 100,000 (including 100 to 5,000 warmup iterations) as

needed. In Gamma models, mildly informative priors (normal(0,2)) were placed on

regression coefficients, and a gamma(2,0.1) prior was applied to the shape parameter;

analogous priors were used for Gaussian models on log-transformed data. Sampling

convergence and model diagnostics were evaluated with R-hat statistics, trace plots, and

effective sample sizes. To account for shared evolutionary history, phylogenetically informed

versions of these models were implemented by including a random intercept of the form

(1|gr(phylo, cov = A)), where the covariance matrix A was derived from the species tree

using the vcv function in the R package ape (Paradis & Schliep 2019).

Residual Analysis for Predictive Accuracy

To further evaluate model performance, we conducted a residual analysis comparing the
predictive accuracy of recombination rate from chromosome number and mean chromosome
size. For the Gamma regression model using chromosome number as the predictor,
posterior expected predictions were obtained on the original scale and residuals were
calculated as the difference between the observed recombination rate and the predicted
value. For the Gaussian model based on log- transformed recombination rate and mean
chromosome size, predictions were first obtained on the log scale and then
back- transformed using the exponential function to yield estimates on the original scale;
residuals were computed analogously. We quantified model performance using the root
mean squared error (RMSE) and mean absolute error (MAE). Furthermore, absolute
residuals of holocentric versus monocentric species were compared using Wilcoxon
rank- sum tests to determine whether the models predicted recombination rate more
accurately in one group than the other. This analysis allowed us to compare both the overall
predictive precision of the two chromosomal predictors and the relative fit across
chromosome types.

Assessing the relation of crossover number per chromosome to chromosome type

To test whether the average number of COs per chromosome differed between holocentric

and monocentric species, we pursued two complementary approaches. Given that COs per

chromosome is strictly positive, we first modeled it using a Gamma regression with a log link

and used chromosome type as the main predictor. The same priors as in the recombination

rate analyses (normal(0,2) for regression coefficients and gamma(2,0.1) for the shape

parameter) were applied. Phylogenetic non-independence was incorporated by including a

random intercept specified as (1|gr(phylo, cov = A)).

To further evaluate the evolutionary dynamics underlying variation in COs per chromosome
between holocentric and monocentric species, we employed an OUwie-based approach
(Beaulieu et al., 2012). A SIMMAP tree was constructed by mapping the discrete character
(chromosome type) onto the species' phylogeny using an equal rates (ER) model. To
account for uncertainty in ancestral state reconstructions, we generated a SIMMAP tree
based on Drinnenberg & Marques (2025). A suite of OUwie models (BM1, BMS, OU1, OUM,
OUMV, OUMA, and OUMVA) was fitted to COs per chromosome. Model selection was
based on weighted AIC computed across replicates, thereby identifying the best-fitting
evolutionary model while explicitly incorporating uncertainty in the character history. Once

the best model(s) was selected, we ran 100 bootstrap replicates using the OUwie.boot
function in OUwie to obtain empirical estimates and 95% confidence intervals for the
evolutionary optimum (θ).

Evaluation of chromosome number and mean chromosome size predictors in angiosperms

To assess the broader applicability of our results beyond insects, we evaluated the accuracy
of chromosome number and mean chromosome size as predictors of recombination rate in
flowering plants with monocentric chromosomes. Holocentric species could not be
meaningfully included, as only two holocentric plant species currently have genetic maps
(Escudero et al., 2018; Castellani et al., 2024). To this end, we obtained data on
recombination rates, genome sizes, and chromosome numbers from Stapley et al. (2017b)
and randomly selected one phylogenetic tree from a set of 100 angiosperm phylogenies
provided by Forest (2023). The dataset and phylogenetic tree overlapped with 157
angiosperm species (Supplementary Table S2), which we used in the regression models
and residual analyses as described above for insects.

Results

We obtained recombination rate, chromosome number, chromosome size, and the average
number of COs per chromosome for 30 holocentric (1 Dermaptera, 4 Hemiptera, 23
Lepidoptera, and 2 Odonata) and 39 monocentric (2 Blattodea, 7 Coleoptera, 9 Diptera, 19
Hymenoptera, and 2 Orthoptera) insect species (Figure 1; Supplementary Table S1).

Differences in Crossover Number between Chromosome Types

We first assessed whether the average number of COs per chromosome (COperChr) differs
between holocentric and monocentric insect species. Using a phylogenetically informed
Bayesian Gamma regression model, we modeled COperChr as a function of chromosome
type, incorporating species' relatedness via a random intercept (Figure 2A; Supplementary
Table S3). In this analysis, the intercept for holocentric species was 0.01 (95% Credible
Interval [CI]: –0.54 to 0.53), corresponding to an expected mean of exp(0.01) ≈ 1.01 COs per
chromosome. The effect of chromosome type (with monocentric species coded as the focal
level) was positive and significant (Estimate = 0.65; 95% CI: 0.10–1.21). This implies that,
on average, monocentric species exhibit nearly double the number of crossovers per
chromosome relative to holocentric species [exp(0.01 + 0.65) ≈ 1.93 COs per
chromosome], supporting our prediction based on chromosomal architecture.

Figure 2. Number of crossovers in holocentric and monocentric insects. **A -** Observed COs per chromosome in holocentric
(blue) and monocentric (red) species. Individual data points are jittered to display variation, and the overlaid symbols with error
bars represent the group mean and 95% prediction credible intervals estimated from the model. **B -** Bootstrap density
distributions (based on 100 replicates) of the evolutionary optimum (θ) for the number of COs estimated under the OUMVA
model. The density for holocentric lineages (blue) and monocentric lineages (red) illustrate the uncertainty and differences in
the adaptive optima between chromosome types.

To further explore the evolutionary dynamics underlying variation in CO numbers, we
conducted an OUwie-based analysis. This approach allowed us to compare models that
assume different evolutionary optima for holocentric versus monocentric lineages. The two
best-fitting models were OUMVA and OUMV (Supplementary Table S3), which yielded
almost identical results and estimated distinct adaptive optima (θ) for each chromosome type
(Figure 2B; Supplementary Table S3). Therefore, here we report only the results from
OUMVA model. Consistent with the regression analysis, the OUwie results indicated a lower
optimum for holocentric species than monocentric ones. The optimum CO number for
holocentric species was estimated at θ_h = 1.00 (95% CI: 0.59–1.27), whereas for
monocentric species it was θ_m = 2.49 (95% CI: 2.05–3.08). Together, these results confirm
that monocentric species maintain a significantly higher average number of COs per
chromosome than holocentric species in insect orders, in line with the expectations derived
from their distinct meiotic architectures.

Chromosome Number as a Predictor of Recombination Rate

Across the 69 analyzed insect species, genome- wide recombination rate was positively
associated with haploid chromosome number (Figure 3; Supplementary Table S4). In a
Bayesian Gamma regression that modeled recombination rate as a function of chromosome
number and chromosome type (holocentric vs. monocentric), the estimated effect of
chromosome number was positive and significant (β = 0.09; 95% Credible Interval [CI]: 0.07–
0.12). Monocentric species exhibited higher baseline recombination rates than holocentric
species (β = 1.65 95% CI: 1.16–2.13; Figure 3A). The interaction between chromosome
number and chromosome type was not supported (β = 0.00; 95% CI: –0.05 to 0.06;
Supplementary Table S4), indicating that the positive relationship between chromosome
number and recombination rate does not differ appreciably between the two chromosomal
architectures. Consistent results were obtained when accounting for phylogenetic non-
independence, although the magnitude of the effect of chromosome number variation was
somewhat reduced (β = 0.05; 95% CI: 0.01–0.08; Figure 3B; Supplementary Table S4). Our

results indicate that genome- wide recombination rates increase with haploid chromosome
number, with monocentric species consistently exhibiting higher baseline rates than
holocentric species, and this relationship remains robust even after accounting for
phylogenetic relatedness.

Figure 3. Relationships between chromosome number, recombination rate, and crossovers per chromosome in insects. **(A)**
Relationship between recombination rate (cM/Mbp) and haploid chromosome number, showing holocentric (blue) and
monocentric (red) species. Solid lines indicate model predictions and shaded ribbons represent 95% prediction credible
intervals. **(B)** Relationship between recombination rate (cM/Mbp) and haploid chromosome number when phylogenetic
relatedness is considered. Dashed lines indicate model predictions and shaded ribbons represent 95% prediction credible
intervals.

Chromosome Size as a Predictor of Recombination Rate

A parallel set of analyses examined the effect of mean chromosome size on recombination
rate by modeling log- transformed recombination rate against log- transformed chromosome
size (Figure 4; Supplementary Table S5). In the non- phylogenetic additive model, mean
chromosome size was a strong negative predictor of recombination rate ($\beta = -1.08$; 95% CI:
-1.21 to -0.95; Figure 4A), indicating that species with smaller chromosomes tend to
experience higher recombination rates per megabase. Although the interaction between
log(chromosome size) and chromosome type was not significant ($\beta = 0.10$ 95% CI: -0.16 to
0.36; Supplementary Table S5), monocentric species consistently maintained higher
recombination rates than holocentric species at a given chromosome size ($\beta = 0.74$; 95% CI:
0.48–0.99; Figure 4A; Supplementary Table S5). Phylogenetic corrections did not alter these
qualitative patterns, with log(chromosome size) remaining negatively associated with
recombination rate ($\beta = -0.93$; 95% CI: -1.15 to -0.70) and the effect of chromosome type
persisting ($\beta = 0.64$; 95% CI: 0.08–1.19) (Figure 4B; Supplementary Table S5). Our analyses
indicate that smaller chromosome sizes predict higher recombination rates per megabase,

and that monocentric species consistently exhibit elevated rates compared to holocentric
species—even after accounting for phylogenetic relationships.

Figure 4. Relationships between mean chromosome size and recombination rate in insects. **(A)** Relationship between
recombination rate (cM/Mbp) and chromosome number without considering phylogeny, showing holocentric (blue) and
monocentric (red) species. Solid lines indicate model predictions and shaded ribbons represent 95% prediction credible
intervals. **(B)** Relationship between recombination rate (cM/Mbp) and mean chromosome size when phylogenetic relatedness is
considered. Dashed lines indicate model predictions and shaded ribbons represent 95% prediction credible intervals.

Residual Analyses: Assessing Predictive Accuracy

The Gamma regression model using chromosome number as the predictor yielded an
RMSE of 3.73 and an MAE of 2.51, whereas the Gaussian model based on log- transformed
recombination rate and mean chromosome size produced lower error metrics (RMSE = 2.75;
MAE = 1.54). Following this, the model with mean chromosome size as a predictor shows
lower absolute residuals than the model with chromosome number as a predictor (Figure 5).
These results highlight that mean chromosome size is a more precise predictor of
recombination rate than chromosome number when considering the overall dataset types.

When considering chromosome type, for the model using chromosome number as the
predictor, holocentric species had significantly lower residuals (mean = 1.35, median = 1.24)
than monocentric species (mean = 3.40, median = 2.73; Wilcoxon test: W = 331, p = 0.0018;
Figure 5), indicating more accurate predictions. Similarly, when predicting log- transformed
recombination rate from log- transformed chromosome size, holocentric species again
showed lower residuals (mean = 0.60, median = 0.31) compared to monocentric species
(mean = 2.27, median = 1.14; W = 304, p = 0.0005; Figure 5). These analyses demonstrate

that the predictive accuracy of recombination rate from both chromosome number and size
is higher in holocentric than in monocentric species.

Figure 5. Absolute residuals from the non-phylogenetic recombination rate models, grouped by predictor (chromosome number
or mean chromosome size) and chromosome type (holocentric or monocentric). The residuals from the model with
chromosome size as the predictor were back-transformed from the log scale to make them comparable with residuals from the
model with chromosome number as the predictor (see Methods for details).

*Evaluation of Chromosome Number and Mean Chromosome Size as Predictors in
Angiosperms*

Regression analyses of 157 angiosperm species with monocentric chromosomes revealed
that recombination rate is positively associated with chromosome number (Supplementary
Figure S1A,B; Supplementary Table S6). In the non- phylogenetic Gamma model, the
estimated coefficient for chromosome number was 0.03 (95% CI: 0.01–0.05). When
accounting for phylogenetic non- independence, the chromosome number coefficient was
0.02 (95% CI: 0.00–0.04). The Gaussian model predicting log- transformed recombination
rate from log- transformed chromosome size (Supplementary Figure S1C,D; Supplementary
Table S7) produced a coefficient for log mean chromosome size of –0.90 (95% CI: –0.96 to
–0.83). With phylogenetic corrections, the coefficient remained similar at –0.91 (95% CI: –
0.98 to –0.83).

Residual analyses further demonstrated that the model based on mean chromosome size
yielded lower error metrics than the model based on chromosome number (Supplementary
Figure S2). Specifically, the Gamma model (chromosome number predictor) had an RMSE
of 1.46 and an MAE of 1.12, with absolute residuals averaging 1.12 (median = 0.98). In
contrast, the log- transformed model (mean chromosome size predictor) produced an RMSE

of 0.92 and an MAE of 0.63, with absolute residuals averaging 0.63 (median = 0.44). These
results indicate that mean chromosome size is a more precise predictor of recombination
rate than chromosome number in angiosperms.

Discussion

The evolution of recombination rates plays a pivotal role in speciation, balancing the benefits
of adaptability with the risks of disrupting established genetic combinations (Ritz et al., 2017;
Stapley et al., 2017a). High recombination rates, while fostering evolutionary potential by
generating novel allele combinations for adaptation to diverse environments (Wang et al.,
2019) or for highly competitive communities (Bell, 1992), can also dismantle favorable co-
adapted gene complexes (Ritz et al., 2017). Conversely, lower recombination rates may
provide greater reproductive assurance by preserving existing adaptive combinations but at
the cost of reduced evolutionary flexibility. This trade-off is particularly evident when
considering the factors influencing recombination.

Here, we show that the number of COs per chromosome is predominantly higher and more
variable in monocentric insects compared to holocentric ones (Figure 2) which provides
support for a long-standing hypothesis based on karyological observations suggesting that
holocentric chromosomes typically exhibit more stringent constraints on chiasma formation
(White, 1973; Nokkala et al., 2004; Gokhman & Kuznetsova, 2006; Schvarzstein et al., 2010;
Metlers et al., 2012; Bureš et al., 2013; Cabral et al., 2014). Mechanistically, these
differences can be attributed to the centromere structure in holocentric species, which
complicates the two-step loss of cohesion that is typical for monocentric species and often
leads to one or at most two chiasmata per homolog pair (Schvarzstein et al., 2010; Marques
& Pedrosa-Harand, 2016; Hofstatter et al., 2021). More than two chiasmata appear to be
either absent or non-viable in holocentric species studied so far (Nokkala et al., 2004; Hollis
et al., 2020; Castellani et al., 2024). In contrast, monocentric chromosomes can achieve
more COs per homolog pair (even over 10) and apparently do so across a range of species
(Jones & Franklin, 2006; Fernandes et al., 2018; Otto & Payseur, 2019; Lian et al., 2023).

In accordance with the tight regulation of the number of COs per chromosome via CO
assurance and CO interference (Hunter, 2015), we identified both chromosome number and
mean chromosome size as good predictors of genome-wide recombination rate in insects
when phylogenetic relatedness is not taken into account (Figures 3A and 4A). However,
once phylogeny is considered, the chromosome number does not perform as well as the
mean chromosome size (Figures 3B and 4B). Our residual analyses further indicate that
mean chromosome size is a more precise predictor of genome- wide recombination rate
than chromosome number, even in the non-phylogenetic approach (Figure 5). It is likely that
the number of chromosomes captures larger differences in recombination rates among
higher taxonomic units (e.g., orders or families), which also differ in chromosome numbers.
In contrast, within genera or families—where closely related species often differ little or not
at all in chromosome number, as e.g. in Lepidoptera (de Vos et al., 2020)—more subtle
differences in recombination rate related to chromosome size are not effectively captured by
chromosome number. Moreover, polyploidization multiplies chromosome number and likely
has little effect on genome- wide recombination rates (Ross- Ibarra, 2007; Tiley & Burleigh,

2015; Stapley et al., 2017b), although autopolyploids may reduce the CO number per
chromosome to avoid multivalent formation (Bomblies et al., 2016). This may further
undermine the predictive accuracy of chromosome numbers. In contrast, mean chromosome
size effectively controls the multiplicative effects of (neo)polyploidy on chromosome number
(Bureš et al., 2024). The superior model fit observed for holocentric species across both
predictors (Figure 5) further suggests that the strict limitations on CO numbers—imposed by
the constraints of their kinetochore architecture—result in more predictable recombination
patterns than those seen in monocentric species. Together, these results indicate that mean
chromosome size is generally a better proxy for recombination rate and that predictions are
more accurate in holocentric than in monocentric systems.

While our findings provide significant support for chromosome number and mean
chromosome size being valid proxies for recombination rate, particularly in holocentric
insects, it is important to acknowledge the limitations of our study. Recombination varies
along a genome and is undoubtedly influenced by many factors beyond chromosome
number and mean chromosome size, such as structural variation, transposable element
activity, chromatin structure, and specific meiotic genes (Johnston, 2024). Although the
broad taxonomic sampling across insects is valuable, extrapolating these findings to other
monocentric and holocentric animals — and especially to plants — should be done with
caution. On the other hand, our analyses encompassing 157 monocentric species of
flowering plants showed very similar results (Supplementary Figures S1, S2) to our findings
in insects (Figures 3, 4, and 5), suggesting their general applicability. Building on results
from insects and monocentric plants, we predicted how genome-wide recombination rate
should relate to mean chromosome size in holocentric plants (Figure 6). Notably,
Rhynchospora breviuscula (Castellani et al., 2024) closely follows the predicted relationship,
whereas *Carex scoparia* (Escudero et al., 2018) deviates from it (Figure 6). This deviation
may reflect random variation, lineage-specific biological factors, lack of a reference genome
assembly, or systematic differences in recombination dynamics between holocentric and
monocentric plants. Notably, while our insect analyses support a shared slope between
chromosome types (Figure 4; Supplementary Table S5), it remains possible that the
relationship between chromosome size and recombination rate differs more fundamentally
between monocentric and holocentric plants. However, with data currently limited to just two
holocentric plant species, we cannot determine whether the observed deviation is
meaningful or incidental. Additional data from diverse holocentric plant lineages will be
essential to assess the robustness and generality of our predictions.

Figure 6. Relationship between mean chromosome size (Mbp) and recombination rate (cM/Mbp) in plants. Black points represent empirical data for monocentric plants. The dark green solid line and shaded area indicate the regression fit and 95% confidence intervals, respectively, for monocentric plants. The blue dotted line and shaded area represent the predicted regression and 95% confidence intervals for holocentric plants, derived by adjusting the monocentric plant intercept using the intercept difference between monocentric and holocentric insect species. Specifically, the intercept for holocentric plants was calculated by subtracting the insect-derived intercept difference (0.80 ± 0.14 SE) from the monocentric plant intercept (4.23 ± 0.15 SE), yielding a holocentric intercept of 3.43 ± 0.205 SE. The slope for holocentric plants was assumed identical to monocentric plants (-0.90 ± 0.03 SE). Blue triangles show the only two plant species with holocentric chromosomes for which both recombination rate and mean chromosome size are known: *Carex scoparia* (Escudero et al., 2018) and *Rhynchospora breviuscula* (Castellani et al., 2024).

Our demonstration of the links among CO number, mean chromosome size, chromosome
type, and recombination rate provides a valuable framework for future large-scale
evolutionary studies aimed at unraveling the evolution of recombination and its role in
adaptation and speciation across the Tree of Life. In light of our findings, it is possible that
previous studies on the holocentric plant genus *Carex* (Bell, 1982; Escudero et al., 2012;
Escudero et al., 2013; Spalink et al., 2018; Márquez- Corro et al., 2021) and monocentric
angiosperms (Carta et al., 2018) may have suffered from reduced power by using
chromosome number rather than mean chromosome size as a proxy for recombination rate.
However, chromosome numbers are available for many more species because the mean
chromosome size requires knowledge of both chromosome number and genome size. For
instance, in flowering plants, chromosome numbers are available for ca. 60,000 species
(Rice et al., 2015) while mean chromosome size is for only about 14,000 species (Bureš et
al., 2024). Therefore, continuing measurements of genome size (e.g., using flow cytometry)
and more initiatives like the Darwin Tree of Life (Darwin Tree of Life Project Consortium,
2022) that generate chromosome-resolved reference genomes for thousands of species will
allow us to obtain mean chromosome size data and chromosome numbers at an
unprecedented scale.

While holocentric chromosomes appear to offer certain evolutionary advantages, such as the
ability to more readily tolerate increased rates of chromosomal fissions and fusions without
necessarily reducing fitness (Zedek & Bureš, 2019; Lucek et al., 2022; Zedek et al., 2022)—
allowing for rapid karyotypic evolution and potentially increased genetic diversity through
altered recombination rates and linkage groups—the prevalence of monocentric

chromosomes in the vast majority (80-85%) of eukaryotes (Márquez-Corro et al., 2018) and comparable rates of chromosome number evolution in monocentric and holocentric insects (Ruckman et al., 2020) suggest a different story. This apparent paradox may be explained by the spatial separation of chromosome segregation and recombination in monocentric chromosomes. In monocentric species, these two crucial meiotic processes are functionally and physically distinct. Holocentric chromosomes, by contrast, must manage both segregation and recombination across their diffuse centromere structure. This constraint likely necessitates a reduction in the number of chiasmata per chromosome to ensure proper segregation, as multiple chiasmata in holocentric species can lead to meiotic errors (Nokkala et al., 2004; Hollis et al., 2020). However, some holocentric insects and plants have also evolved non-canonical meiosis to solve these problems (Melters et al., 2012; Lukhtanov et al., 2018). Our findings, demonstrating a significantly higher number of chiasmata per homolog in monocentric species compared to holocentric species, directly support this explanation and help to resolve the paradox. The increased chiasma frequency in monocentrics likely contributes to their greater evolutionary success by allowing for higher levels of genetic recombination. Thus, while holocentric species may offer flexibility in karyotypic evolution, the inherent trade-off between recombination rate specifically for chiasma formation and proper segregation, might ultimately limit their overall evolutionary success compared to the more specialized and efficient mechanisms found in monocentric chromosomes. In summary, our findings reveal a fundamental difference in how insect recombination is regulated. While monocentric species rely on increased CO rates, holocentric species appear to compensate through changes in chromosome number. This stark contrast underscores centromere architecture's critical role in shaping insect genomes' evolutionary trajectory.

Data availability statement

All the data used in this article are available in supplementary materials online on the publisher's website.

Funding

This work was supported by the Czech Science Foundation, grant no. 24-11400S to FZ. KL was supported by the Swiss National Science Foundation grants 202869 and 220868 as well as the Fondation Pierre Mercier pour la science. ME was supported by the Spanish AEI-MICINN grant DiversiChrom *PID2021-122715NB-I00.* AM was financially supported by the Max Planck Society.

Author contributions
FZ conceived the study and performed analyses. FZ, PB, TLE, ME, KL, and AM wrote the manuscript.

Conflicts of interests

None declared.

References

Anderson LK, Salameh N, Bass HW, Harper LC, Cande WZ, Weber G, Stack SM. 2004.
Integrating genetic linkage maps with pachytene chromosome structure in maize. *Genetics*
166: 1923–1933. doi: 10.1534/genetics.166.4.1923.
578

579 **Backström N, Forstmeier W, Schielzeth H, Mellenius H, Nam K, Bolund E, Webster MT,**
580 **Ost T, Schneider M, Kempenaers B, Ellegren H. 2010.** The recombination landscape of
581 the zebra finch *Taeniopygia guttata* genome. *Genome Research* **20**: 485–495. doi:
582 10.1101/gr.101410.109.
583

Barnes TM, Kohara Y, Coulson A, Hekimi S. 1995. Meiotic recombination, noncoding DNA
and genomic organization in *Caenorhabditis elegans*. *Genetics* **141**: 159–179. doi:
10.1093/genetics/141.1.159.

Beaulieu JM, Jhwueng DC, Boettiger C, O'Meara BC. 2012. Modeling stabilizing
selection: expanding the Ornstein-Uhlenbeck model of adaptive evolution. *Evolution* **66**:
2369–2383. doi: 10.1111/j.1558-5646.2012.01619.x.

Bell G. 1982. The masterpiece of nature: the evolution and genetics of sexuality. University
of California Press.

Blary A, Jenczewski E. 2019. Manipulation of crossover frequency and distribution for plant
breeding. *Theoretical and Applied Genetics* **132**: 575–592.
594

595 **Bomblies K, Jones G, Franklin C, Zickler D, Kleckner N. 2016.** The challenge of evolving
596 stable polyploidy: could an increase in "crossover interference distance" play a central role?
597 *Chromosoma* **125**: 287–300. doi: 10.1007/s00412-015-0571-4.
598

599 **Brazier T, Glémin S. 2022.** Diversity and determinants of recombination landscapes in
600 flowering plants. *PLoS Genetics* **18**: e1010141. doi: 10.1371/journal.pgen.1010141.
601

602 **Bureš P, Elliott TL, Veselý P, Šmarda P, Forest F, Leitch IJ, Nic Lughadha E, Soto**
603 **Gomez M, Pironon S, Brown MJM, Šmerda J, Zedek F. 2024.** The global distribution of
604 angiosperm genome size is shaped by climate. *New Phytologist* **242**: 744–759. doi:
605 10.1111/nph.19544.
606

607 **Bureš P, Zedek F. 2014.** Holokinetic drive: centromere drive in chromosomes without
608 centromeres. *Evolution* **68**: 2412–2420. doi: 10.1111/evo.12437.
609

610 **Bureš P, Zedek F, Marková M. 2013.** Holocentric chromosomes. In: *Plant genome*
611 *diversity. Vol. 2.*, Vienna: Springer, 187-208.
612

Burt A. 2000. Perspective: sex, recombination, and the efficacy of selection--was Weismann
right? *Evolution* **54**: 337–351. doi: 10.1111/j.0014-3820.2000.tb00038.x.

Butlin RK. 2005. Recombination and speciation. *Molecular Ecology* **14**: 2621–2635. doi:
10.1111/j.1365-294X.2005.02617.x.

Bürkner P. 2017. brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal*
of Statistical Software **80**: 1–28. doi:10.18637/jss.v080.i01.

Bürkner P. 2018. Advanced Bayesian Multilevel Modeling with the R Package brms. *The R*
Journal **10**: 395–411. doi:10.32614/RJ-2018-017.

Bürkner P. 2021. Bayesian Item Response Modeling in R with brms and Stan. *Journal of*
Statistical Software **100**: 1–54. doi:10.18637/jss.v100.i05.

Cabral G, Marques A, Schubert V, Pedrosa-Harand A, Schlögelhofer P. 2014.
Chiasmatic and achiasmatic inverted meiosis of plants with holocentric chromosomes.
Nature Communications **5**: 5070. doi: 10.1038/ncomms6070.

Carta A, Bedini G, Peruzzi L. 2018. Unscrambling phylogenetic effects and ecological
determinants of chromosome number in major angiosperm clades. *Scientific Reports* **8**:
14258.

Castellani M, Zhang M, Thangavel G, Mata-Sucre Y, Lux T, Campoy JA, Marek M,
Huettel B, Sun H, Mayer KFX, Schneeberger K, Marques A. 2024. Meiotic recombination
dynamics in plants with repeat-based holocentromeres shed light on the primary drivers of
crossover patterning. *Nature Plants* **10**: 423–438. doi: 10.1038/s41477-024-01625-y.

Chuang YC, Smith GR. 2023. Meiotic crossover interference: Methods of analysis and
mechanisms of action. *Current Topics in Developmental Biology* **151**: 217–244. doi:
10.1016/bs.ctdb.2022.04.006.

Crismani W, Mercier R. 2012. Meiotic recombination: mechanisms and implications.
Genetics **190**: 1325–1334.

Cuacos MH, Franklin FC, Heckmann S. 2015. Atypical centromeres in plants—what they
can tell us. *Frontiers in Plant Science* **6**: 913.

Darwin Tree of Life Project Consortium. 2022. Sequence locally, think globally: The
Darwin Tree of Life Project. *Proc Natl Acad Sci U S A* **119**: e2115642118. doi:
10.1073/pnas.2115642118.

de Vos JM, Augustijnen H, Bätscher L, Lucek K. 2020. Speciation through chromosomal
fusion and fission in Lepidoptera. *Philosophical Transactions of the Royal Society B:*
Biological Sciences **375**: 20190539. doi: 10.1098/rstb.2019.0539.

Elliott TL, Zedek F, Barrett RL, Bruhl JJ, Escudero M, Hroudová Z, Joly S, Larridon I,
Luceño M, Márquez-Corro JI, Martín-Bravo S, Muasya AM, Šmarda P, Thomas WW,

Wilson KL, Bureš P. 2022. Chromosome size matters: genome evolution in the cyperid clade. *Annals of Botany* **130**: 999–1014. doi: 10.1093/aob/mcac136.

Escudero M, Hipp AL, Hansen TF, Voje KL, Luceño M. 2012. Selection and inertia in the evolution of holocentric chromosomes in sedges (*Carex*, Cyperaceae). *New Phytologist* **195**: 237–247. doi: 10.1111/j.1469-8137.2012.04137.x.

Escudero M, Maguilla E, Modesto L. 2013. Selection by climatic regime and neutral evolutionary processes in holocentric chromosomes (*Carex* gr. *laevigata*: Cyperaceae): A microevolutionary approach. *Perspectives in Plant Ecology, Evolution and Systematics* **15**: 118–129.

Escudero M, Hahn M, Hipp AL. 2018. RAD-seq linkage mapping and patterns of segregation distortion in sedges: meiosis as a driver of karyotypic evolution in organisms with holocentric chromosomes. *Journal of Evolutionary Biology* **31**: 833–843. doi: 10.1111/jeb.13267.

Fernandes JB, Séguéla-Arnaud M, Larchevêque C, Lloyd AH, Mercier R. 2018. Unleashing meiotic crossovers in hybrid plants. *Proc Natl Acad Sci U S A* **115**: 2431–2436. doi: 10.1073/pnas.1713078114.

Forest F. 2023. Species-level phylogenetic trees of all angiosperm species (100 trees). https://zenodo.org/record/7600341

Gokhman VE, Kuznetsova VG. 2006. Comparative insect karyology: current state and applications. *Entomological Review* **86**: 352–368.

Gregory TR. 2025. Animal Genome Size Database. http://www.genomesize.com.

Haenel Q, Laurentino TG, Roesti M, Berner D. 2018. Meta-analysis of chromosome-scale crossover rate variation in eukaryotes and its significance to evolutionary genomics. *Molecular Ecology* **27**: 2477–2497. doi: 10.1111/mec.14699.

Heckmann S, Jankowska M, Schubert V, Kumke K, Ma W, Houben A. 2014. Alternative meiotic chromatid segregation in the holocentric plant *Luzula elegans*. *Nature Communications* **5**: 4979.

Heckmann S, Macas J, Kumke K, Fuchs J, Schubert V, Ma L, Novák P, Neumann P, Taudien S, Platzer M, Houben A. 2013. The holocentric species *Luzula elegans* shows interplay between centromere and large-scale genome organization. *The Plant Journal* **73**: 555–565. doi: 10.1111/tpj.12054.

Hofstatter PG, Thangavel G, Castellani M, Marques A. 2021. Meiosis Progression and Recombination in Holocentric Plants: What Is Known? *Frontiers in Plant Science* **12**: 658296. doi: 10.3389/fpls.2021.658296.

Hofstatter PG, Thangavel G, Lux T, Neumann P, Vondrak T, Novak P, Zhang M, Costa L, Castellani M, Scott A, Toegelová H, Fuchs J, Mata-Sucre Y, Dias Y, Vanzela ALL,

Huettel B, Almeida CCS, Šimková H, Souza G, Pedrosa-Harand A, Macas J, Mayer KFX, Houben A, Marques A. 2022. Repeat-based holocentromeres influence genome architecture and karyotype evolution. *Cell* **185**: 3153–3168.e18. doi: 10.1016/j.cell.2022.06.045.

Hollis JA, Glover ML, Schlientz AJ, Cahoon CK, Bowerman B, Wignall SM, Libuda DE. 2020. Excess crossovers impede faithful meiotic chromosome segregation in *C. elegans*. *PLoS Genetics* **16**: e1009001. doi: 10.1371/journal.pgen.1009001.

Hunter N. 2015. Meiotic Recombination: The Essence of Heredity. *Cold Spring Harbor Perspectives in Biology* **7**: a016618. doi: 10.1101/cshperspect.a016618.

Jankowska M, Fuchs J, Klocke E, Fojtová M, Polanská P, Fajkus J, Schubert V, Houben A. 2015. Holokinetic centromeres and efficient telomere healing enable rapid karyotype evolution. *Chromosoma* **124**: 519–528. doi: 10.1007/s00412-015-0524-y.

Jones GH, Franklin FCH. 2006. Meiotic crossing-over: obligation and interference. *Cell* **126**: 246–248.

Jensen-Seaman MI, Furey TS, Payseur BA, Lu Y, Roskin KM, Chen CF, Thomas MA, Haussler D, Jacob HJ. 2004. Comparative recombination rates in the rat, mouse, and human genomes. *Genome Research* **14**: 528–538. doi: 10.1101/gr.1970304.

Johnston SE. 2024. Understanding the Genetic Basis of Variation in Meiotic Recombination: Past, Present, and Future. *Molecular Biology and Evolution* **41**: msae112. doi: 10.1093/molbev/msae112.

Kaback DB, Guacci V, Barber D, Mahon JW. 1992. Chromosome size-dependent control of meiotic recombination. *Science* **256**: 228–232. doi: 10.1126/science.1566070.

Kent TV, Uzunović J, Wright SI. 2017. Coevolution between transposable elements and recombination. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**: 20160458. doi: 10.1098/rstb.2016.0458.

Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, Stecher G, Hedges SB. 2022. TimeTree 5: An Expanded Resource for Species Divergence Times. Molecular Biology and Evolution **39**: msac174. doi: 10.1093/molbev/msac174.

Leitch AR, Ma L, Dodsworth S, Fuchs J, Houben A, Leitch IJ. 2023. The Role of Chromatin Modifications in the Evolution of Giant Plant Genomes. *Plants (Basel)* **12**: 2159. doi: 10.3390/plants12112159.

Lian Q, Maestroni L, Gaudin M, Llorente B, Mercier R. 2023. Meiotic recombination is confirmed to be unusually high in the fission yeast *Schizosaccharomyces pombe*. *iScience* **26**: 107614. doi: 10.1016/j.isci.2023.107614.

Lucek K, Augustijnen H, Escudero M. 2022. A holocentric twist to chromosomal speciation? *Trends in Ecology and Evolution* **37**: 655–662. doi: 10.1016/j.tree.2022.04.002.

Lukhtanov VA, Dincă V, Friberg M, Šíchová J, Olofsson M, Vila R, Marec F, Wiklund C. 2018. Versatility of multivalent orientation, inverted meiosis, and rescued fitness in holocentric chromosomal hybrids. *Proc Natl Acad Sci U S A* **115**: E9610–E9619. doi: 10.1073/pnas.1802610115.

Lynch M, Bobay LM, Catania F, Gout JF, Rho M. 2011. The repatterning of eukaryotic genomes by random genetic drift. *Annual Review of Genomics and Human Genetics* **12**: 347–366. doi: 10.1146/annurev-genom-082410-101412.

Mackintosh A, Laetsch DR, Hayward A, Charlesworth B, Waterfall M, Vila R, Lohse K. 2019. The determinants of genetic diversity in butterflies. *Nature Communications* **10**: 3466. doi: 10.1038/s41467-019-11308-4.

Mancera E, Bourgon R, Brozzi A, Huber W, Steinmetz LM. 2008. High-resolution mapping of meiotic crossovers and non-crossovers in yeast. *Nature* **454**: 479–485. doi: 10.1038/nature07135.

Mandrioli M, Manicardi GC. 2012. Unlocking holocentric chromosomes: new perspectives from comparative and functional genomics? *Current Genomics* **13**: 343–349. doi: 10.2174/138920212801619250.

Mandrioli M, Manicardi GC. 2020. Holocentric chromosomes. *PLoS Genetics* **16**: e1008918.

Marques A, Drinnenberg IA. 2025. Same but different: Centromere regulations in holocentric insects and plants. *Current Opinion in Cell Biology* **93**: 102484. doi: 10.1016/j.ceb.2025.102484.

Marques A, Pedrosa-Harand A. 2016. Holocentromere identity: from the typical mitotic linear structure to the great plasticity of meiotic holocentromeres. *Chromosoma* **125**: 669–681.

Márquez-Corro JI, Escudero M & Luceño M. 2018. Do holocentric chromosomes represent an evolutionary advantage? A study of paired analyses of diversification rates of lineages with holocentric chromosomes and their monocentric closest relatives. *Chromosome Research* **26:** 139-152.

Márquez- Corro JI, Martín- Bravo S, Jiménez- Mejías P, Hipp AL, Spalink D, Naczi RF, Roalson EH, Luceño M, Escudero M. 2021. Macroevolutionary insights into sedges (*Carex*: Cyperaceae): The effects of rapid chromosome number evolution on lineage diversification. *Journal of Systematics and Evolution* **59**: 776–790. doi: 10.1111/jse.12730.

Martin SH, Davey JW, Salazar C, Jiggins CD. 2019. Recombination rate variation shapes barriers to introgression across butterfly genomes. *PLoS Biology* **17**: e2006288. doi: 10.1371/journal.pbio.2006288.

Mata-Sucre Y, Krátká M, Oliveira L, Neumann P, Macas J, Schubert V, Huettel B, Kejnovský E, Houben A, Pedrosa-Harand A, Souza G, Marques A. 2024. Repeat-based

holocentromeres of the woodrush Luzula sylvatica reveal insights into the evolutionary
transition to holocentricity. *Nature Communications* **15**: 9565. doi: 10.1038/s41467-024-
53944-5.

Melters DP, Paliulis LV, Korf IF, Chan SW. 2012. Holocentric chromosomes: convergent
evolution, meiotic adaptations, and genomic analysis. *Chromosome Research* **20**: 579–593.

Mercier R, Mézard C, Jenczewski E, Macaisne N, Grelon M. 2015. The molecular biology
of meiosis in plants. *Annual Review of Plant Biology* **66**: 297–327.

Näsvall K, Boman J, Höök L, Vila R, Wiklund C, Backström N. 2023. Nascent evolution
of recombination rate differences as a consequence of chromosomal rearrangements. *PLoS*
Genetics **19**: e1010717. doi: 10.1371/journal.pgen.1010717.

Nokkala S, Kuznetsova VG, Maryanska-Nadachowska A, Nokkala C. 2004. Holocentric
chromosomes in meiosis. I. Restriction of the number of chiasmata in bivalents.
Chromosome Research **12**: 733–739.

Nordenskiöld H. 1963. A study of meiosis in the progeny of X-irradiated *Luzula purpurea*.
Hereditas **49**: 33–47.

Otto SP, Payseur BA. 2019. Crossover interference: shedding light on the evolution of
recombination. *Annual Review of Genetics* **53**: 19–44.

Palahí I Torres A, Höök L, Näsvall K, Shipilina D, Wiklund C, Vila R, Pruisscher P,
Backström N. 2023. The fine-scale recombination rate variation and associations with
genomic features in a butterfly. *Genome Research* **33**: 810–823. doi:
10.1101/gr.277414.122.

Paradis E, Schliep K .2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**: 526–528. doi:10.1093/bioinformatics/bty633.

Pessia E, Popa A, Mousset S, Rezvoy C, Duret L, Marais GA. 2012. Evidence for
widespread GC-biased gene conversion in eukaryotes. *Genome Biology and Evolution* **4**:
675–682. doi: 10.1093/gbe/evs052.

Rice A, Glick G, Abadi S, Einhorn M, Kopelman NM, Salman-Minkov A, Mayzel J, Chay
O, Mayrose I. 2015. The chromosome counts database (CCDB) – a community resource of
plant chromosome numbers. *New Phytologist* **206**: 16–26. doi: 10.1111/nph.13191.

Ritz KR, Noor MA, Singh ND. 2017. Variation in recombination rate: adaptive or not?.
Trends in Genetics **33**: 364–374.

Ross-Ibarra J. 2007. Genome size and recombination in angiosperms: a second look.
Journal of Evolutionary Biology **20**: 800–806. doi: 10.1111/j.1420-9101.2006.01275.x.

Ruckman SN, Jonika MM, Casola C, Blackmon H. 2020. Chromosome number evolves at
equal rates in holocentric and monocentric clades. *PLoS Genetics* **16**: e1009076. doi:
10.1371/journal.pgen.1009076.

Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, Connor R, Funk K,
Kelly C, Kim S, Madej T, Marchler-Bauer A, Lanczycki C, Lathrop S, Lu Z, Thibaud-
Nissen F, Murphy T, Phan L, Skripchenko Y, Tse T, Wang J, Williams R, Trawick BW,
Pruitt KD, Sherry ST. 2022. Database resources of the National Center for Biotechnology
Information. *Nucleic Acids Research* **50**: D20–D26. doi: 10.1093/nar/gkab1112.

Shipilina D, Näsvall K, Höök L, Vila R, Talavera G, Backström N. 2022. Linkage mapping
and genome annotation give novel insights into gene family expansions and regional
recombination rate variation in the painted lady (Vanessa cardui) butterfly. *Genomics* **114**:
110481. doi: 10.1016/j.ygeno.2022.110481.

Spalink D, Pender J, Escudero M, Hipp AL, Roalson EH, Starr JR, Waterway MJ, Bohs
L, Sytsma KJ. 2018. The spatial structure of phylogenetic and functional diversity in the
United States and Canada: An example using the sedge family (Cyperaceae). *Journal of*
Systematics and Evolution **56**: 449–465. doi: 10.1111/jse.12423.

Stapley J, Feulner PG, Johnston SE, Santure AW, Smadja CM. 2017a. Recombination:
the good, the bad and the variable. *Philosophical Transactions of the Royal Society B:*
Biological Sciences **372**: 20170279.

Stapley J, Feulner PG, Johnston SE, Santure AW, Smadja CM. 2017b. Variation in
recombination frequency and distribution across eukaryotes: patterns and processes.
Philosophical Transactions of the Royal Society B: Biological Sciences **372**: 20160455.

Ruckman SN, Jonika MM, Casola C, Blackmon H. 2020. Chromosome number evolves at
equal rates in holocentric and monocentric clades. *PLoS Genetics* **16**: e1009076.

Schvarzstein M, Melters DP. 2010. Recombination dynamics in holocentric chromosomes.
Chromosome Research **18**: 213–224.

Tigano A, Khan R, Omer AD, Weisz D, Dudchenko O, Multani AS, Pathak S, Behringer
RR, Aiden EL, Fisher H, MacManes MD. 2022. Chromosome size affects sequence
divergence between species through the interplay of recombination and selection. *Evolution*
76: 782–798. doi: 10.1111/evo.14467.

Tiley GP, Burleigh JG. 2015. The relationship of recombination rate, genome structure, and
patterns of molecular evolution across angiosperms. *BMC Evolutionary Biology* **15**: 194. doi:
10.1186/s12862-015-0473-3.

Viera A, Page J, Rufas JS. 2009. Inverted meiosis: the true bugs as a model to study.
Meiosis **5**: 137–156.

Wang S, Veller C, Sun F, Ruiz-Herrera A, Shang Y, Liu H, Zickler D, Chen Z, Kleckner
N, Zhang L. 2019. Per-Nucleus Crossover Covariation and Implications for Evolution. *Cell*
177: 326–338.e16. doi: 10.1016/j.cell.2019.02.021.

White MJD. 1973. *Animal cytology and evolution.* Cambridge University Press, Cambridge,
UK
866

Zedek F, Bureš P. 2019. Pest Arthropods with Holocentric Chromosomes are More
Resistant to Sterilizing Ionizing Radiation. *Radiation Research* **191**: 255–261. doi:
10.1667/RR15208.1.

Zedek F, Veselý P, Tichý L, Elliott TL, Garbolino E, de Ruffray P, Bureš P. 2022.
Holocentric plants are more competitive under higher UV-B doses. *New Phytologist* **233**: 15–
21. doi: 10.1111/nph.17750.

Supplementary Figures

Supplementary Figure S1. Relationships between chromosome number, mean chromosome size, and recombination rate in
flowering plants with monocentric chromosomes. (A) Relationship between recombination rate (cM/Mbp) and haploid
chromosome number. (B) Relationship between recombination rate (cM/Mbp) and haploid chromosome number when
phylogenetic relatedness is considered. (C) Relationship between recombination rate (cM/Mbp) and mean chromosome size
(both variables are on a logarithmic scale). (D) Relationship between recombination rate (cM/Mbp) and mean chromosome size
when phylogenetic relatedness is considered (both variables are on a logarithmic scale). Solid and dashed lines indicate model
predictions and shaded ribbons represent 95% prediction credible intervals.

Supplementary Figure S2. Absolute residuals from the non-phylogenetic recombination rate models using chromosome
number and mean chromosome size as predictors in flowering plants with monocentric chromosomes. Residuals from the
model with chromosome size were back-transformed from the log scale to enable direct comparison with those from the
chromosome number model.