Systematic and quantitative analyses of pre-clinical *Mycobacterium avium* Lung Disease

Tools for Drug Development and Transition from Animal Models to New Approach

Methodologies

Moti Chapagain,[1,2] Devyani Deshpande,[3] Shashikant Srivastava,[4,5] Tawanda Gumbo[,1,2,6]

[1]Mathematical Modeling and AI, Praedicare inc., Dallas, Texas

[2]Hollow Fiber System & Experimental Therapeutics Laboratories, Praedicare Inc., Dallas, Texas,

USA

[3]Baylor University Medical Center, Dallas, Texas, USA.

[4]Division of Infectious Diseases, Department of Medicine, The University of Texas at Tyler

School of Medicine, Tyler, Texas, USA.

[5]Department of Cellular and Molecular Biology, Center for Biomedical Research, University of

Texas Health Science Center at Tyler, Tyler, Texas, USA.

[6]IMPI Biotech Consortium Inc, Tateguru, Zimbabwe

ABSTRACT

Animal and new approach methodologies such as the hollow fiber system model [HFS] are used

for *Mycobacterium avium-complex* [MAC] lung disease [LD] preclinical drug development. Our

objective was to perform a systematic review to benchmark these pre-clinical tools. We

performed a literature search to identify preclinical pharmacokinetics/pharmacodynamics

[PK/PD] studies for MAC-LD. Preferred Reporting Items for Systematic Reviews and Meta-

analyses was used for bias minimization. Twenty HFS-MAC and 3 mouse studies met PK/PD

inclusion criteria. We created a novel quality score tool based on predictors of clinical response,

design optimization, and information theory. The quality score was judged high in 10%, good in

50%, adequate in 30%, and poor in 10% of studies. Monte Carlo experiments for PK/PD target

attainment were reported in 61% of studies. On repetitive sampling, the PK/PD target exposure

estimate varied significantly between sampling days in 76% of studies. The solution was

ordinary differential equations with parameter outputs such as γ [nonlinear kill-slope] and time-

to-extinction applied to both HFS-MAC and patients' sputa CFU/mL output. Next, we ranked the

antibiotics by extent of microbial kill as fold-improvement over guideline-based therapy. The

three top ranked drugs for microbial kill were omadacycline [69-fold], tedizolid [19-fold], and

ceftriaxone [8-fold]. We recommend the HFS-MAC as tractable for exposure-effect and dose-

fractionation studies, ranking antibiotics effect, and for translation to clinical doses. The

analyses inform us of HFS-MAC recommendations for regulatory authorities and drug

developers, including quality scores for optimal PK/PD design for target identification, resistance

suppression, and choice of the best novel regimen.

Keywords: hollow fiber system model of MAC lung disease; quality score tool; efficacy ranking;

PK/PD design criteria; factorial design; quality control

Mycobacterium avium complex [MAC] is responsible for nearly 80% of all non-tuberculous

mycobacteria [NTM] lung disease [LD] (1). Guideline-based therapy [GBT] includes a

combination of macrolide [azithromycin or clarithromycin], ethambutol, and a rifamycin [rifampin

or rifabutin] (2). Meta-analyses of clinical trials have demonstrated that GBT achieves 6-month

sustained sputum culture conversion [SSCC] in only 43-53% of patients; the number needed to

be treated is five to benefit one patient (3, 4). The predictors of patient clinical response to GBT

are [1] pretreatment bacterial burden (B_0), [2] drugs with high intracellular penetration such as

macrolides and macrolide-based regimens, [3] presence of cavitary disease, and [4] cavities

of >2cm diameter (2-5). Thus, therapy failure is due to pharmacokinetics [PK]-

pharmacodynamics [PK] and the pathobiology of the disease. MAC-LD clinicopathological

features include nodules and/or cavities, intracellular MAC in monocyte-lineage cells in

granulomatous and necrotic lesions, with MAC B_0 median [range] of 5.17 [4.23-6.2] $\log_{10}$

CFU/mL in cavities and 3.0 [0.48-3.85] $\log_{10}$ CFU/mL in nodular/bronchiectatic lesions (6-9).

The recent US Food and Drug Administration [FDA] and European Medicines Agencies [EMA]

roadmaps state that new approach methodologies [NAMs] "offer the tools to assess safety,

efficacy, and pharmacology of drugs and therapeutics *without* traditional animal models" and

that such tools must be of "high relevance to human biology" (10, 11). Consistent with these

recent regulatory agencies roadmaps, and MAC-LD clinicopathological factors associated with

therapy failure, the hollow fiber system model [HFS] of MAC-LD [HFS-MAC] could be relevant to

human biology as regards lung PKs, B_0, intracellular nature of MAC, and drug diffusion

properties (10-12). Here we performed a literature search for ALL preclinical PK/PD models,

animal models and the HFS-MAC, to systematically characterize these models' success and

failures for benchmarking.

Antimicrobial PK/PD is a quantitative science of the relationship between drug PKs' and

antimicrobial effect such as bacterial [1] growth or [2] kill or (3) antimicrobial resistance [AMR]

(13). PKs are captured by concentration-time profiles of drug and their shapes in lesions, as well

as indices such as peak concentration [C_{max}], 0-24h area under the concentration-time curve

[AUC_{0-24}], or % time concentration persists above MIC [$\%T_{MIC}$]. These are used to identify the

PK/PD target exposure, which is the lowest exposure that kills the microbe and/or suppresses

AMR, and the exposure that must be attained in the lung by a dose to cure MAC-LD. Ambrose

et al have shown a strong correlation between the probability of PK/PD target attainment [PTA]

of doses versus approval of new drug applications for antibiotic use in lung disease (14, 15).

This PK/PD approach also improves the chances of success in treating patients in the clinic,

while being cost and time effective, important for an orphan disease such as MAC-LD (15).

Starting four decades ago PK/PD science was applied to Gram-negative bacilli and Gram-

positive cocci, and then we applied to tuberculosis [TB] two decades ago, and more recently to

MAC-LD and *Mycobacterium abscessus* LD (16-24). Here, we performed a systematic analysis

for lessons learnt in MAC-LD preclinical PK/PD models.

METHODS

Rationale

Following work with the HFS model of TB [HFS-TB], which led to the qualification of the HFS-TB

as a drug development tool by the EMA, we propose to achieve the same for preclinical models

of MAC-LD (10, 11, 16-19, 25-27). Second, PK/PD work for MAC-LD has been ongoing for 15

years and has not been examined using meta-analytic approaches.

Objectives and study questions

Objective 1: To develop a quality score tool to judge the quality of pre-clinical models for MAC-

LD, for use by the academe, scientific journals, biopharma industry, and regulatory authorities.

Objective 2: To perform a systematic review of all pre-clinical models for MAC-LD that fulfil

definitions of full PK/PD studies, for regulatory submission to the FDA and EMA.

Objective 3: To perform a quantitative analysis of all pre-clinical models for MAC-LD to rank

new/repurposed drugs based on the extent of microbial kill, to design novel regimens.

Definitions used as criteria for inclusion and exclusion of studies

The reason to perform antimicrobial PK/PD studies is to identify target exposures and dosing

schedules. This is translated to optimal dosing regimens in the clinic, for maximal efficacy and

minimal AMR. *In-silico dose* finding is most often implemented in Monte Carlo experiments, to

account for PK variability in patients (28, 29). This approach integrates MIC distributions and

generates PK/PD susceptibility breakpoints. These foundational concepts were used to define

criteria for inclusion and exclusion of studies.

First, a PK/PD approach requires a PK system. This is because in patients both microbial kill

and AMR are a consequence of fluctuating concentrations, the shape of the concentration-time

curves, and periodicity [dosing schedule], not captured by static concentrations in time-kill

curves (13, 16-19, 30, 31). In fact, the exact PK profile determines outcomes. As an example, in

the HFS model of anthrax and in mice and macaques lung disease, the levofloxacin human half-

life of 7.5h was compared to the mouse and rhesus macaque half-life of 2h at the same AUC

(32, 33). The human PKs produced persistent reduction in bacterial burden while animal PKs

led to failure and AMR (32, 33). Thus, static concentrations studies such as time-kill studies and

stand-alone MIC results were defined as not fulfilling a PK/PD definition. Consequently, a

preclinical model that did not report PKs from concentrations *measured* in the same experiment,

did not fulfil the full PK/PD definition. An intended PK profile without measurements to back it up

did not fulfil PK/PD criteria: "Trust but verify".

Second, an exposure-response study must test several exposures to identify the PK/PD target

exposure. So, what is the minimum number of exposures that define a full PK/PD study? The

canonical model used for exposure-response analysis, the inhibitory sigmoid maximal effect

[E_{max}] or Hill equation, has the following formalism:

$$\text{Effect } [\log_{10} \text{CFU/mL}] = E_{con} - E_{max} * EC^H / [EC^H + EC_{50}{}^H] \qquad [1]$$

with the 4 parameters and [1] E_{con}, [2] E_{max}, (3) H, and (4) EC_{50} (24, 34, 35). EC_{50} or potency is

the drug exposure mediating 50% of E_{max}. If there are more model parameters [n] than data

points [i.e., exposures or doses], the model will not have a unique solution and will be

undetermined. Therefore, at a minimum n+1 or 5 exposure points are needed for optimal design

of an exposure-response study. Moreover, based on both Shannon entropy and optimal Fisher

information, exposures between those mediating 20% [EC_{20}] and 80% [EC_{80}] of E_{max}, at least

three exposures between of EC_{20} and EC_{80}, are a minimum requirement for dose-fractionation

studies (36). The definitions of at least five exposures for exposure-response study and three

between EC_{20} and EC_{80} for dose-fractionation studies were used as inclusion criteria for PK/PD

studies in the presented analysis.

Third, the same drug pair or triplets can have antagonism at some concentrations, additivity at

others, and antagonism at yet other concentrations, at different sampling times (37-40).

Therefore, for combination therapy, more than one dose needs to be examined in pre-clinical

models. The PK/PD definitions of synergy or antagonism or additivity are most often based on

two 2-parameter models and probability theory: Bliss independence and Loewe additivity. Bliss

Independence has the following formalism:

$$Y_{ab,P} = Y_{a.} + Y_b + Y_a * Y_b \qquad [2]$$

for interaction index, with Y_a effect [i.e., inhibition or microbial kill] of drug "*a*" at specified dose,

and Y_b effect of drug "*b*" at specified dose. Therefore, there is need for at least three dosing

regimens consisting of drug "*a*" alone, drug "*b*" alone, and "*a*b*" combined, plus non-treated

controls [four regimens]. Exposures of each of the drugs to be examined should be at least

three [EC_{20}, EC_{50}, and EC_{80}] (41). Thus, for two drugs there is need of 16 combination pairs

[including zero] at a minimum. Further, since interaction factor is defined based on the 95%

confidence interval crossing zero, replicates are required. We defined combination therapy

PK/PD study as being acceptable if there was minimum factorial design with at least the four

regimens: *0, a, b, a*b*. These must have at least 2 replicates.

Literature search strategy and study selection

We performed a literature search to identify PK/PD studies for MAC-LD, as defined by the

criteria above. Stand-alone Monte Carlo experiments which used the PK/PD targets for dose

selection were also included. The Medical Subject Heading [MeSH] terms and strategy used

included either [1] "pharmacokinetics-pharmacodynamics" or "hollow fiber" OR "hollow fibre" or

"mouse AND pharmacokinetics-pharmacodynamics" AND [2] either "*Mycobacterium avium"* or

"non-tuberculous mycobacteria". Bibliographies of original articles and guidelines were

examined for additional relevant studies. Two authors [SS and TG] searched PubMed, Google

Scholar, Inter-Science Conference on Antimicrobial Agents and Chemotherapy [ICAAC], ID

week [the annual conference of the Infectious Diseases Society of America] and the yearly

American Thoracic Society meeting, for studies published through March 31, 2025. When a

conference abstract was found we also sought out the posters or oral presentations. If a

conference presentation was published later in a scientific journal, the full publication was taken

as the main reference. The two authors independently extracted the data into the prespecified

table format. Consensus was reached for study inclusion after the discussion of each study.

There was no exclusion of articles by language. Bias minimization was according to Preferred

Reporting Items for Systematic Reviews and Meta-Analyses [PRISMA] (42).

Data and outcomes collected

Data collected included number of units [animals, in vitro experiments units], replicates used, B_0

achieved in CFU/mL or CFU/lung, number of doses tested, type of study design [exposure-

effect or dose fractionation or factorial design], number of bacterial isolates tested, and number

of Monte Carlo experiments doses. The outcomes recorded were CFU/mL or CFU/lung on each

sampling day, AMR population in CFU/mL, drug concentrations [mg/L or mg/g], PK/PD

exposure values, and PK/PD parameters reported or linked to efficacy and AMR suppression.

Calculated outcomes captured from the data or calculated by us were PK/PD target exposures,

microbial kill below B_0, EC_{80}, and *in silico* dose-ranging to identify probability of target attainment

[PTA] in published Monte Carlo experiments.

Quality criteria based on PK/PD science and predictors of clinical response to therapy

We proposed quality criteria based on experimental design optimization characteristics for the

laboratory work, and on modeling the important predictors of patient response identified by

clinicians, and "high relevance to human biology" (2-11). Since it is important to use multiple

MAC isolates in order not to misestimate PK/PD targets, and since regulatory agencies

guidelines state "that ~4-5 organisms of the major target species or organism groups should be

tested", we also scored based on number of non-ATCC clinical isolates used (43). The

proposed quality criteria were as detailed in **Supplementary Methods** and **Supplementary**

Table S1. This initial tool was refined based on statistical analyses, to a final scoring tool.

Data Synthesis and Analysis

The first level of analysis was qualitative. This was based on the main findings reported in each

study. This approach is descriptive.

The second type of analysis was quantitative. For each study we captured the following data,

and if data were not provided, we calculated the data:

1) Microbial kill below B_0 at E_{max} in CFU/mL and then ranked drugs by that metric. The

microbial kill was weighted by the number of isolates used in the study, and fold difference

with the three-drug GBT microbial kill among 5 isolates was calculated for normalization.

2) The target exposure, EC_{80}.

3) Dose that achieves EC_{80} in lungs and the probability of toxicity at that dose.

4) Interaction index for combinations using factorial design.

5) The reliability of each study was qualified by the final quality score tool.

RESULTS

Selected PK/PD studies

We found 16 animal studies [37% of all studies identified], shown in the PRISMA flow diagram

in **Figure 1** (44-59). Thirteen studies, shown in **Supplementary data Table S2** did not fulfill the

PK/PD criteria for inclusion: most had less than 5 exposures tested, and all combination therapy

studies excluded did not employ factorial design (44-55, 59). Three animal studies met the

criteria, De *et al* exposure-effect and combination therapy [considered two studies], and Remal

combination therapy study in BALB/c mice (56-58).

We found 27 HFS-MAC PK/PD studies [63% of all studies identified], with first publication in

2010, shown in **Figure 1** (12, 60-85). Twenty HFS-MAC PK/PD studies and Monte Carlo

experiments were met inclusion criteria (12, 60-78). Seven combination therapy studies were

excluded because they did not meet the minimum definition of factorial design and are shown in

Supplementary Table S3 (79-85).

Quality scores

The histogram of the quality scores using the initial tool is shown in **Figure 2A**. **Figure 2A**

shows the distribution of scores including the combination therapy in HFS-MAC not otherwise

discussed further because they did not meet PK/PD criteria and were included for tool

development purposes only. The mean score for the excluded studies was 8.43 (95%

Confidence interval (CI): 7.35-9.51) and would be categorized as poor quality or worse. **Figure**

2B shows the correlation coefficient between the different criteria was high [r>0.5 or R<-0.5]

between "potency" [i.e., reports of EC_{50} and PK/PD target (potency)] versus doses [i.e., number

of doses/exposures tested] criteria [Pearson r=0.91; p= 5.6 $\times 10^{-8}$]. Thus, we removed the

"potency" criteria from the tool. In addition, since all HFS-MAC studies basically scored the

same for intracellular MAC criteria, and all mice had intracellular disease, this criterion was

removed and enfolded into basic definition of an acceptable MAC-LD PK/PD study. The final

criteria scoring system that was developed was as shown in **Table 1**. The total possible score

was categorized as follows, [1] high score of >20, [2] good score of 15-20, [3] adequate score of

10-14, [4] poor quality score of 5-9, and [5] deficient and unreliable of less than 5.

The histogram of quality scores for this final tool is shown in **Figure 2C** for animal and HFS-

MAC studies**.** The mean score for studies that met inclusion criteria was 15.35 (95%CI: 13.33-

17.37). The quality score was judged high in 10%, good in 50%, adequate in 30%, and poor in

10% of studies. **Figure 2D** depicts a Bland-Altman agreement analysis between the observers

for the quality scores. **Figure 2D** shows the average score minus observer's score as an

agreement metric; the overall average for the score was 11.28 [95% CI: 10.36-13.43] and the

difference of scores between observers 0.03 [95% CI: -0.99 to 1.04]. The overall bias between

all observers was 1.98 (95% confidence interval: 0.36-3.9), which is close to zero.

Qualitative Systemic analysis findings

Table 2 shows a qualitative analysis of the different studies. The characteristics of the drugs

tested in these preclinical PK/PD models are shown in **Figure 3**. The most common

pharmacophores studied were β-lactams [23.5%], followed by tetracyclines and oxazolidinones

at 17.7% each, while macrolides were tested in only 11.8%. In the first three years, HFS-MAC

studies lasted only seven days, but switched to 28 days starting 2012 leading to better mapping

and PK/PD modeling of AMR sub-populations. All HFS-TB experiments underwent repetitive

sampling and measured the drug concentrations versus time points that were achieved [which

were all reported], for THP-1 cell count, as well as for both total bacterial burden and drug-

resistant bacteria. No study reported the % coefficient of variance [%CV] between HFS-MAC

replicates for drug concentrations or cell and CFU/mL counts. AMR universally arose to

monotherapy and was concentration dependent.

In the first 10 years, CFU/mL output from HFS-MAC versus exposure on each sampling day

was examined in the inhibitory sigmoid E_{max} and the antibiotic resistance arrow of time models,

with parameter estimates used being those on the day with best Akaike information criteria (12,

60-66). However, for HFS-MAC studies that were longer than 7 days repetitive sampling

revealed that in 13 out of 17 [76%] monotherapy studies, the EC_{50} and H varied between

sampling days by a factor of 2 or more outside 95% confidence intervals, as could the PK/PD

index [%T_{MIC} or AUC/MIC or C_{max}/MIC] linked to effect (12, 63-67, 71-77). This presents

problems as to which sampling day data to use to calculate the PK/PD target exposure.

Therefore, in 2019 sets of ordinary differential equations [ODEs] that gave parameter outputs

that are vectors such as γ [non-linear kill slope), mutation rates [*m*], growth rates [*r*], and time to

extinction [*τ*] were introduced for both HFS-MAC and patients sputa, with γ as new PD

parameter (67, 71, 74, 75). Second, several studies demonstrated that the PK/PD target

identification became more robust if up to five clinical MAC isolates were used in the PK/PD

studies (65, 73, 76, 77). Commonly the reference ATCC MAC isolate mis-estimated PK/PD

targets when compared to 5-isolate studies and to clinical studies (3, 4, 65, 73, 76, 77).

Monte Carlo experiments

Fourteen of the 23 studies [61%] reported results of Monte Carlo experiments, for dose

selection. These were reported mainly after monotherapy exposure-response studies.

Quantitative analyses and ranking of drugs by extent of microbial kill below B_0

Since the whole point of PK/PD studies is [1] identification of PK/PD targets in tandem with *in*

silico based dose versus PTA in Monte Carlo experiments and [2] to find drugs with better

efficacy [maximal microbial effect] than current GBT which performs poorly (3, 4), we performed

a quantitative analysis that ranked drugs by how much better they compared to GBT [i.e., as

fold of GBT]. **Table 3** ranks the microbial kill from the HFS-MAC studies and demonstrates that

the top three ranked drugs, at doses that can be tolerated by people, were [1] 300 mg/day oral

omadacycline or 35 mg/day inhaled tigecycline at 66-69-fold-GBT, [2] 200mg/day oral tedizolid

at 19-fold-GBT, and 2g/day intravenous ceftriaxone at 8-fold-GBT. Based on Monte Carlo

experiments to achieve PK/PD target exposure [PTA], the doses of each drug and the PK/PD

susceptibility breakpoints shown in **Table 3**.

Quality control HFS-MAC

For comparison and quality control purposes, we added a clarithromycin row in **Tables 2** and **3**,

that is part of our unpublished internal controls for HFS-MAC over the last 15 years, given that

clarithromycin is considered the primary drug driving outcome in MAC-LD. The data is from 32

HFS-MAC replicates [and 256 data points] of the American Type Culture Collection 700898

[ATCC] (12 HFS-MAC replicates) and five clinical isolates [20 HFS-MAC replicates]. The

clarithromycin time-kill curves for the ATCC isolate versus the 5 clinical isolates that are part of

our internal standards, are shown in **Figure 4A**, which demonstrates that the ATCC isolate

gives an overoptimistic picture. **Figure 4B** shows the %CV between replicates on each

sampling day; the higher %CV with ATCC is from work in earlier 10 years, while those in the

recent 5 years had %CV below 20% between replicates on each sampling day. Overall, when

all studies were considered, the %CV between replicates was 10.12% [95% CI: 7.01-13.24%].

Clarithromycin achieved 0.03-fold-GBT in ATCC isolate and 0.02-fold among 5 clinical isolates.

DISCUSSION AND RECOMMENDATIONS

Sixty eight percent of MAC-LD PK/PD studies were performed in the HFS-MAC while 32% were

in mice. Thus, the HFS-MAC has come a long way since the seven-day study with ethambutol

16 years ago (60). The HFS-MAC has become more sophisticated with repeated use, with

longer duration of therapy in the system, which is now routinely over 28 days. The %CV

between replicates has fallen from 24.24% in first 10 years to 4.53% recently. The HFS-MAC

allows repetitive sampling and tracking of AMR evolution and microbial kill and their

relationships to drug exposure. When the HFS-TB was qualified as a drug development tool, 22

HFS-TB studies had been published, a similar situation as with the HFS-MAC here (17-19). Our

systematic analyses of both the HFS-MAC and murine models led to multiple recommendations.

Quality scores [QS]

First, we developed a tool that scores pre-clinical laboratory models for quality and adequacy of

the models, and for adequacy of PK/PD design. The final tool has 6 scoring criteria **[Table 1]**.

We recommend this QS tool for HFS-MAC and mouse PK/PD work. The tool could also be used

by academia for standardization and to assess rigor and reproducibility by the academe and

granting institutes such as the National Institutes of Health. Indeed, the same concerns of rigor

and reproducibility are often raised by editors and reviewers of scientific journals. For regulatory

authorities [FDA or EMA], this tool could be used as minimal criteria for acceptance of PK/PD

data biopharma for medical product development for MAC-LD. It could also be used to

determine how much weight to give to a PK/PD result and dose choice for new drug

applications. For biopharma companies, this could also be used as criteria to judge the quality

of data from contract research organizations doing laboratory MAC-LD work.

PK/PD design criteria for monotherapy and combination therapy for MAC-LD (5-5-5-25)

Based on pathological features that predict therapeutic outcomes, we recommend that the

preclinical models of MAC-LD be intracellular and have the B_0 in the range observed in patient

lesions (2, 6-9). Exposure-response PK/PD require at least 5 doses, and greater weight should

be given to evidence from studies that have greater than 5 exposures. For combination therapy,

since concentration-dependent synergy and antagonism are a fact of life, and since PK

variability in patients will lead to situations in which each combination results *de facto* in a

spectrum of exposures, at least three exposures should be tested in factorial design. To identify

unbiased PK/PD targets, at least 5 MAC isolates should be examined in these studies (65, 73,

76, 77). Moreover, to account for biological variability, and to fulfil quality criteria scores,

replicates should be used. We propose that a %CV between replicates on any sampling above

25% be defined as a failure. We recommend documenting %CV between replicates for drug

concentrations or cell and CFU/mL counts at each sampling time for all reports. We propose

that the Monte Carlo experiments themselves examine at least 5 doses *in silico* for PTA to

achieve target exposure and for probability of toxicity. Data generated from these more robust

types of experimental design, and quality control, used to identify optimal exposures to be

translated to human doses using Monte Carlo experiments, are crucial for better precision.

The antibiotic resistance arrow of time

All PK/PD experiments should simultaneously identify and report the exposures associated with

optimal kill and those associated with AMR suppression. In preclinical PK/PD studies, drug-

resistant CFUs are captured by growing on agar supplemented with three times the MIC of a

drug, in essence a three-fold increase in MIC, chosen to capture low-level resistance from efflux

pump induction (12). Repetitive sampling in the HFS-MAC allows documentation of the

evolution of efflux-pump induction and high-level chromosomal-mutation-related AMR versus

drug exposure (12). On the other hand, the PK/PD susceptibility breakpoint identified in Monte

Carlo experiment PTA defines resistance as the MIC above which patients fail therapy and is

dependent on the dose used in the clinic and dosing route. Here in **Table 3**, we documented the

PK/PD breakpoints for MAC-LD for drugs beyond clarithromycin. Clinical validation is still

required. However, in the HFS-TB, such PK/PD breakpoints were identical to those picked by

machine learning when clinical data was later examined (86-91).

Recommendations of the best novel regimen for clinical trial testing

MAC-LD is an orphan disease. In addition, antibiotics have a negative net present value (92).

This means that resources for developing new regimens are limited, as are the number of

volunteers available for clinical trials. The most common approach has been to iteratively

replace one of the drugs in GBT such as macrolides and rifamycins with repurposed ones (79-

85). However, with this approach, it takes many years to develop a novel regimen and often can

only examine single doses of each drug in combination (79-85). Moreover, this perpetuates the

psychological "favorite child" based choice by academe: judging a drug's utility based on being

the discoverer or first adopter of the drug. We propose to select drugs for novel regimens-based

ranking of how well they better than the combination GBT (macrolide plus ethambutol plus

rifamycin). The top three ranked drugs in **Table 3** were omadacycline, tedizolid, and ceftriaxone:

each surpassed GBT. We propose first performing HFS-MAC studies using factorial design for

omadacycline, tedizolid, and ceftriaxone, with HFS-MAC output converted into ODEs, which can

then be translated to SSCC in patients. A virtual 10,000 patient clinical trial can then be

performed, followed by a clinical trial in patients that utilizes few patient volunteers.

Limitations

First, there were few well designed mouse and combination therapy PK/PD studies. The large

portion of studies excluded based on PK/PD quantitative criteria, which include three of our own,

also scored poorly on our new quality score tool. There was no bias in our search, which means

the poor scores are intrinsic to the studies. The most common reason what that the studies

were not designed to identify target exposures in monotherapy or for combinations and could

not inform on clinical doses or dosing schedules, which is the point of PK/PD studies. They

missed the "5-5-5-25" criteria. Second, our novel quality score tool is yet to be tested by other

investigators and thus remains a work in progress. Third, exposure-effect studies often did not

include replicates. While technically the inhibitory sigmoid E_{max} is a regression that can handle

lack of replicates, biological replicates are needed for better rigor and for quality control

purposes. One of the recommendations resulting from current systemic analysis is to always

have replicates. Fourth, use of multiple clinical isolates, at least 4, will be important for more

robust estimation of target exposures. Both HFS-MAC and murine studies often had only the

ATCC strains, with the limitation that results cannot be generalized. Fifth, while the role of

biofilm in MAC lung is yet to be shown from human histology specimens, it cannot just be

discounted. None of the models used explored that bacterial phenotype. Finally, there is always

the inherent bias within ourselves given that we came up with the HFS-MAC model which is our

own psychological "favorite child" over mice. That is the reason we introduced quantitative tools

and criteria that are agnostic of models to address that bias. Our conclusions that the HFS-MAC

model was more commonly used than mice in PK/PD studies, harmonizes with the FDA

Roadmap statement: "*Over 90% of drugs that appear safe and effective in animals do not go on

to receive FDA approval in humans predominantly due to safety and/or efficacy issues*" (10).

REGISTRATION OF REVIEW: Not registered.

ACKNOWLEDGEMENT. None.

CONFLICT OF INTEREST. TG Founded Praedicare Inc, which uses the hollow fiber

methodologies for drug development

FUNDING SOURCE. None.

DATA AVAILABILITY STATEMENT. The data for the results presented in the manuscript are

publicly available.

ETHICAL APPROVAL. Not applicable.

AUTHOR CONTRIBUTIONS. Conceptualization and design, TG; Data analysis, MC, DD, TG

and SS. TG wrote the first draft of the manuscript. All authors reviewed and approved the final

version of the manuscript.

REFERENCES

1. Prevots DR, Shaw PA, Strickland D, Jackson LA, Raebel MA, Blosky MA, Montes de Oca R, Shea YR, Seitz AE, Holland SM, Olivier KN. 2010. Nontuberculous mycobacterial lung disease prevalence at four integrated health care delivery systems. Am J Respir Crit Care Med 182:970-6.

2. Kang HR, Hwang EJ, Kim SA, Choi SM, Lee J, Lee CH, Yim JJ, Kwak N. 2021. Clinical Implications of Size of Cavities in Patients With Nontuberculous Mycobacterial Pulmonary Disease: A Single-Center Cohort Study. Open Forum Infect Dis 8:ofab087.

3. Pasipanodya JG, Ogbonna D, Deshpande D, Srivastava S, Gumbo T. 2017. Meta-analyses and the evidence base for microbial outcomes in the treatment of pulmonary Mycobacterium avium-intracellulare complex disease. J Antimicrob Chemother 72:i3-i19.

4. Kwak N, Park J, Kim E, Lee CH, Han SK, Yim JJ. 2017. Treatment Outcomes of Mycobacterium avium Complex Lung Disease: A Systematic Review and Meta-analysis. Clin Infect Dis 65:1077-1084.

5. Lam PK, Griffith DE, Aksamit TR, Ruoss SJ, Garay SM, Daley CL, Catanzaro A. 2006. Factors related to response to intermittent treatment of Mycobacterium avium complex lung disease. Am J Respir Crit Care Med 173:1283-9.

6. Ushiki A, Yamazaki Y, Koyama S, Tsushima K, Yamamoto H, Hanaoka M, Kubo K. 2011. Bronchoscopic microsampling for bacterial colony counting in relevant lesions in patients with pulmonary Mycobacterium avium complex infection. Intern Med 50:1287-92.

7. Hibiya K, Shigeto E, Iida K, Kaibai M, Higa F, Tateyama M, Fujita J. 2012. Distribution of mycobacterial antigen based on differences of histological characteristics in pulmonary

Mycobacterium avium infectious diseases--consideration of the extent of surgical

resection from the pathological standpoint. Pathol Res Pract 208:53-8.

8.   Fujita J, Ohtsuki Y, Shigeto E, Suemitsu I, Yamadori I, Shiode M, Nishimura K,

Hirayama T, Matsushima T, Ishida T. 2002. Pathological analysis of the cavitary wall in

Mycobacterium avium intracellulare complex pulmonary infection. Intern Med 41:617-21.

9.   Fujita J, Ohtsuki Y, Suemitsu I, Shigeto E, Yamadori I, Obayashi Y, Miyawaki H,

Dobashi N, Matsushima T, Takahara J. 1999. Pathological and radiological changes in

resected lung specimens in Mycobacterium avium intracellulare complex disease. Eur

Respir J 13:535-40.

10.   US Food and Drug Administration. 2025. FDA Roadmap to Reducing Animal Testing in

Preclinical Safety Studies.

https://www.fda.gov/files/newsroom/published/roadmap_to_reducing_animal_testing_in_

preclinical_safety_studies.pdf. 186092. FDA, Bethesda, Maryland.

11.   European Medicines Agencies. 2025. Regulatory acceptance of new approach

methodologies (NAMs) to reduce animal use testing.

https://www.ema.europa.eu/en/human-regulatory-overview/research-

development/ethical-use-animals-medicine-testing/regulatory-acceptance-new-

approach-methodologies-nams-reduce-animal-use-testing.

12.   Schmalstieg AM, Srivastava S, Belkaya S, Deshpande D, Meek C, Leff R, van Oers NS,

Gumbo T. 2012. The antibiotic resistance arrow of time: efflux pump induction is a

general first step in the evolution of mycobacterial drug resistance. Antimicrob Agents

Chemother 56:10.

13.   Drusano GL. 2004. Antimicrobial pharmacodynamics: Critical interactions of 'bug and

455   drug'. Nature Reviews Microbiology 2:289-300.

456   14.   Bulik CC, Bhavnani SM, Hammel J, Forrest A, Dudley MN, Ellis-Grosse E, Drusano G,

457   Ambrose PG. Relationship between regulatory approval and pharmacokinetic-

458   pharmacodynamic target attainment: Focus on community- and hospital-acquired

459   pneumonia, p Abstract A-295. *In* (ed),

460   15.   Palmer ME, Andrews LJ, Abbey TC, Dahlquist AE, Wenzler E. 2022. The importance of

461   pharmacokinetics and pharmacodynamics in antimicrobial drug development and their

462   influence on the success of agents developed to combat resistant gram negative

463   pathogens: A review. Front Pharmacol 13:888079.

464   16.   Ambrose PG, Bhavnani SM, Rubino CM, Louie A, Gumbo T, Forrest A, Drusano GL.

465   2007. Pharmacokinetics-pharmacodynamics of antimicrobial therapy: it's not just for

466   mice anymore. Clin Infect Dis 44:79-86.

467   17.   Gumbo T, Pasipanodya JG, Nuermberger E, Romero K, Hanna D. 2015. Correlations

468   Between the Hollow Fiber Model of Tuberculosis and Therapeutic Events in

469   Tuberculosis Patients: Learn and Confirm. Clin Infect Dis 61 Suppl 1:S18-24.

470   18.   Pasipanodya JG, Nuermberger E, Romero K, Hanna D, Gumbo T. 2015. Systematic

471   Analysis of Hollow Fiber Model of Tuberculosis Experiments. Clin Infect Dis 61 Suppl

472   1:S10-7.

473   19.   Gumbo T, Pasipanodya JG, Romero K, Hanna D, Nuermberger E. 2015. Forecasting

474   Accuracy of the Hollow Fiber Model of Tuberculosis for Clinical Therapeutic Outcomes.

475   Clin Infect Dis 61 Suppl 1:S25-31.

476   20.   Ferro BE, Srivastava S, Deshpande D, Sherman CM, Pasipanodya JG, van Soolingen D,

477   Mouton JW, van Ingen J, Gumbo T. 2015. Amikacin

478   pharmacokinetics/pharmacodynamics in a novel hollow-fiber *Mycobacterium abscessus*

479   disease model. Antimicrob Agents Chemother 60:1242-8.

480   21.   Ferro BE, Srivastava S, Deshpande D, Pasipanodya JG, van Soolingen D, Mouton JW,

481   van Ingen J, Gumbo T. 2016. Tigecycline Is highly efficacious against *Mycobacterium*

482   *abscessus* pulmonary disease. Antimicrob Agents Chemother 60:2895-900.

483   22.   Ferro BE, Srivastava S, Deshpande D, Pasipanodya JG, van Soolingen D, Mouton JW,

484   van Ingen J, Gumbo T. 2016. Failure of the amikacin, cefoxitin, and clarithromycin

485   combination regimen for treating pulmonary *Mycobacterium abscessus* infection.

486   Antimicrob Agents Chemother 60:6374-6.

487   23.   Gumbo T. 2018. Chemotherapy of tuberculosis, *Mycobacterium avium* complex

488   disease, and leprosy. *In* Brunton LL, Chabner B, Knollmann B (ed), Goodman &

489   Gilman's The Pharmacological Basis of Therapeutics, vol 13. McGraw Hill Medical, New

490   York, NY.

491   24.   Gumbo T. 2018. General principles of chemotherapy of infectious diseases. *In* Brunton

492   LL, Chabner B, Knollmann B (ed), Goodman & Gilman's The Pharmacological Basis of

493   Therapeutics, vol 13. McGraw Hill Medical, New York, NY.

494   25.   European Medicines Agencies. 2014. Dossier submitted for Qualification Opinion.

495   Hollow Fiber System Model of Tuberculosis (HFS-TB) as an in vitro preclinical tool for

496   optimization of dose selection and drug regimen in anti-TB drug development.

497   Procedure number: EMEA/H/SAB/049/1/QO/2014/SME.

498   https://www.ema.europa.eu/en/documents/other/dossier-submitted-qualification-opinion-

vitro-hollow-fibre-system-model-tuberculosis-hfs-tb_en.pdf. EMA, London, UK.

https://www.ema.europa.eu/en/documents/other/dossier-submitted-qualification-opinion-

vitro-hollow-fibre-system-model-tuberculosis-hfs-tb_en.pdf.

26.   European-Medicines-Agencies. 2015. Qualification opinion on in-vitro hollow-fibre-

system model of tuberculosis (HFS-TB).

http://www.ema.europa.eu/ema/index.jsp?curl=pages/includes/document/document_det

ail.jsp?webContentId=WC500181899&mid=WC0b01ac058009a3dc. Accessed

01.31.2017.

27.   Cavaleri M, Manolis E. 2015. Hollow Fiber System Model for Tuberculosis: The

European Medicines Agency Experience. Clin Infect Dis 61 Suppl 1:S1-4.

28.   Hall RG, 2nd, Swancutt MA, Meek C, Leff RD, Gumbo T. 2012. Ethambutol

pharmacokinetic variability is linked to body mass in overweight, obese, and extremely

obese people. Antimicrob Agents Chemother 56:1502-7.

29.   Hall RG, 2nd, Swancutt MA, Meek C, Leff R, Gumbo T. 2013. Weight drives caspofungin

pharmacokinetic variability in overweight and obese people: fractal power signatures

beyond two-thirds or three-fourths. Antimicrob Agents Chemother 57:2259-64.

30.   Ishii K, Matsuda H, Iwasa Y, Sasaki A. 1989. Evolutionarily Stable Mutation Rate in a

Periodically Changing Environment. Genetics 121:163-174.

31.   Carja O, Liberman U, Feldman MW. 2014. Evolution in changing environments:

modifiers of mutation, recombination, and migration. Proc Natl Acad Sci U S A

111:17935-40.

32.   Deziel MR, Heine H, Louie A, Kao M, Byrne WR, Basset J, Miller L, Bush K, Kelly M,

521   Drusano GL. 2005. Effective antimicrobial regimens for use in humans for therapy of

522   Bacillus anthracis infections and postexposure prophylaxis. Antimicrobial Agents and

523   Chemotherapy 49:5099-5106.

524   33.   Kao LM, Bush K, Barnewall R, Estep J, Thalacker FW, Olson PH, Drusano GL, Minton N,

525   Chien S, Hemeryck A, Kelley MF. 2006. Pharmacokinetic considerations and efficacy of

526   levofloxacin in an inhalational anthrax (postexposure) rhesus monkey model. Antimicrob

527   Agents Chemother 50:3535-42.

528   34.   Hill AV. 1910. The possible effects of the aggregation of the molecules of hemoglobin on

529   its dissociation curves. J Physiol 40:iv-vii.

530   35.   Weiss JN. 1997. The Hill equation revisited: uses and misuses. FASEB J 11:835-841.

531   36.   Shannon CE. 1997. The mathematical theory of communication. 1963. MD Comput

532   14:306-17.

533   37.   Swaminathan S, Pasipanodya JG, Ramachandran G, Hemanth Kumar AK, Srivastava S,

534   Deshpande D, Nuermberger E, Gumbo T. 2016. Drug Concentration Thresholds

535   Predictive of Therapy Failure and Death in Children With Tuberculosis: Bread Crumb

536   Trails in Random Forests. Clin Infect Dis 63:S63-S74.

537   38.   Rockwood N, Pasipanodya JG, Denti P, Sirgel F, Lesosky M, Gumbo T, Meintjes G,

538   McIlleron H, Wilkinson RJ. 2017. Concentration-Dependent Antagonism and Culture

539   Conversion in Pulmonary Tuberculosis. Clin Infect Dis 64:1350-1359.

540   39.   Deshpande D, Srivastava S, Nuermberger E, Pasipanodya JG, Swaminathan S, Gumbo

541   T. 2016. Concentration-dependent synergy and antagonism of linezolid and moxifloxacin

542   in the treatment of childhood tuberculosis: the dynamic duo. Clin Infect Dis 63:S88-S94.

543   40.   Gumbo T, Chapagain M, Magombedze G, Srivastava S, Deshpande D, Pasipanodya JG,

544   Gumbo R, Dheda K, Diacon A, Hermann D, Hanna D. 2022. Novel tuberculosis

545   combination regimens of two and three-months therapy duration. bioRxiv

546   doi:10.1101/2022.03.13.484155:2022.03.13.484155.

547   41.   Greco WR, Bravo G, Parsons JC. 1995. The search for synergy: a critical review from a

548   response surface perspective. Pharmacol Rev 47:331-85.

549   42.   Moher D, Liberati A, Tetzlaff J, Altman DG. 2009. Preferred reporting items for

550   systematic reviews and meta-analyses: the PRISMA statement. BMJ 339:b2535.

551   43.   European Medicines Agencies. 2016. Guideline on the use of pharmacokinetics and

552   pharmacodynamics in the development of antimicrobial medicinal products.

553   EMA/CHMP/594085/2015. EMA/CHMP/594085/2015. Agency EM, London, United

554   Kingdom. https://www.ema.europa.eu/en/documents/scientific-guideline/guideline-use-

555   pharmacokinetics-pharmacodynamics-development-antimicrobial-medicinal-

556   products_en.pdf.

557   44.   Düzgüneş N, Ashtekar DR, Flasher DL, Ghori N, Debs RJ, Friend DS, Gangadharam PR.

558   1991. Treatment of Mycobacterium avium-intracellulare complex infection in beige mice

559   with free and liposome-encapsulated streptomycin: role of liposome type and duration of

560   treatment. J Infect Dis 164:143-51.

561   45.   Tomioka H, Saito H, Sato K, Yoneyama T. 1991. Therapeutic efficacy of liposome-

562   encapsulated kanamycin against Mycobacterium intracellulare infection induced in mice.

563   Am Rev Respir Dis 144:575-9.

564   46.   Klemens SP, Cynamon MH. 1991. In vivo activities of newer rifamycin analogs against

565   Mycobacterium avium infection. Antimicrob Agents Chemother 35:2026-30.

566   47.   Tomioka H, Saito H, Sato K, Yamane T, Yamashita K, Hosoe K, Fujii K, Hidaka T. 1992.

567   Chemotherapeutic efficacy of a newly synthesized benzoxazinorifamycin, KRM-1648,

568   against Mycobacterium avium complex infection induced in mice. Antimicrob Agents

569   Chemother 36:387-93.

570   48.   Klemens SP, Cynamon MH. 1992. Activity of rifapentine against Mycobacterium avium

571   infection in beige mice. J Antimicrob Chemother 29:555-61.

572   49.   Kailasam S, Wise DL, Gangadharam PR. 1994. Bioavailability and chemotherapeutic

573   activity of clofazimine against Mycobacterium avium complex infections in beige mice

574   following a single implant of a biodegradable polymer. J Antimicrob Chemother 33:273-9.

575   50.   Le Conte P, Le Gallou F, Potel G, Struillou L, Baron D, Drugeon HB. 1994.

576   Pharmacokinetics, toxicity, and efficacy of liposomal capreomycin in disseminated

577   Mycobacterium avium beige mouse model. Antimicrob Agents Chemother 38:2695-701.

578   51.   Jagannath C, Emanuele MR, Hunter RL. 1999. Activities of poloxamer CRL-1072

579   against Mycobacterium avium in macrophage culture and in mice. Antimicrob Agents

580   Chemother 43:2898-903.

581   52.   Peters K, Leitzke S, Diederichs JE, Borner K, Hahn H, Muller RH, Ehlers S. 2000.

582   Preparation of a clofazimine nanosuspension for intravenous use and evaluation of its

583   therapeutic efficacy in murine Mycobacterium avium infection. J Antimicrob Chemother

584   45:77-83.

585   53.   Ellis S, Kalinowski DS, Leotta L, Huang ML, Jelfs P, Sintchenko V, Richardson DR,

586   Triccas JA. 2014. Potent antimycobacterial activity of the pyridoxal isonicotinoyl

587   hydrazone analog 2-pyridylcarboxaldehyde isonicotinoyl hydrazone: a lipophilic transport

588   vehicle for isonicotinic acid hydrazide. Mol Pharmacol 85:269-78.

589   54.   Grewal TK, Majeed S, Sharma S. 2018. Therapeutic implications of nano-encapsulated

590   rifabutin, azithromycin & ethambutol against experimental Mycobacterium avium

591   infection in mice. Indian J Med Res 147:594-602.

592   55.   Banaschewski B, Verma D, Pennings LJ, Zimmerman M, Ye Q, Gadawa J, Dartois V,

593   Ordway D, van Ingen J, Ufer S, Stapleton K, Hofmann T. 2019. Clofazimine inhalation

594   suspension for the aerosol treatment of pulmonary nontuberculous mycobacterial

595   infections. J Cyst Fibros 18:714-720.

596   56.   De K, DeStefano MS, Shoen C, Cynamon M, Alley MR. 2022. Epetraborole, a Novel

597   Bacterial Leucyl-tRNA Synthetase Inhibitor, Demonstrates Potent Efficacy and Improves

598   Efficacy of Standard of Care Regimen Against Mycobacterium avium complex in a

599   Chronic Mouse Lung Infection Model. Open Forum Infectious Diseases 9:ofac492.1334.

600   57.   Bhavnani SM, Hammel JP, Ganesan H, Safir MC, Rubino CM, Ambrose PG, Krause KM.

601   2022. Pharmacokinetic-Pharmacodynamic (PK-PD) Target Attainment Analyses to

602   Support Epetraborole Dose Selection for the Treatment of Patients with Mycobacterium

603   avium Complex (MAC) Lung Disease Open Forum Infect Dis 9:ofac492.671.

604   58.   Rimal B, Howe RA, Panthi C, Lamichhane G. 2024. Regimen comprising clarithromycin,

605   clofazimine and bedaquiline is more efficacious than monotherapy in a mouse model of

606   chronic Mycobacterium avium lung infection. bioRxiv doi:10.1101/2024.12.11.627976.

607   59.   Cotroneo N, Stokes SS, Pucci MJ, Rubio A, Hamed KA, Critchley IA. 2024. Efficacy of

608   SPR720 in murine models of non-tuberculous mycobacterial pulmonary infection. J

609   Antimicrob Chemother 79:875-882.

610   60.   Deshpande D, Srivastava S, Meek C, Leff R, Gumbo T. 2010. Ethambutol optimal

611   clinical dose and susceptibility breakpoint identification by use of a novel

612   pharmacokinetic-pharmacodynamic model of disseminated intracellular *Mycobacterium*

613   *avium*. Antimicrob Agents Chemother 54:1728-33.

614   61.   Deshpande D, Srivastava S, Meek C, Leff R, Hall GS, Gumbo T. 2010. Moxifloxacin

615   pharmacokinetics/pharmacodynamics and optimal dose and susceptibility breakpoint

616   identification for treatment of disseminated *Mycobacterium avium* infection. Antimicrob

617   Agents Chemother 54:2534-9.

618   62.   Deshpande D, Pasipanodya JG, Gumbo T. 2016. Azithromycin Dose To Maximize

619   Efficacy and Suppress Acquired Drug Resistance in Pulmonary Mycobacterium avium

620   Disease. Antimicrob Agents Chemother 60:2157-63.

621   63.   Deshpande D, Srivastava S, Musuka S, Gumbo T. 2016. Thioridazine as Chemotherapy

622   for Mycobacterium avium Complex Diseases. Antimicrobial Agents and Chemotherapy

623   60:4652-4658.

624   64.   Deshpande D, Srivastava S, Pasipanodya JG, Gumbo T. 2017. Linezolid as treatment

625   for pulmonary Mycobacterium avium disease. J Antimicrob Chemother 72:i24-i29.

626   65.   Deshpande D, Srivastava S, Pasipanodya JG, Lee PS, Gumbo T. 2017. Tedizolid is

627   highly bactericidal in the treatment of pulmonary *Mycobacterium avium* complex disease.

628   Journal of Antimicrobial Chemotherapy 72:30-35.

629   66.   Deshpande D, Srivastava S, Chapagain ML, Lee PS, Cirrincione KN, Pasipanodya JG,

630   Gumbo T. 2017. The discovery of ceftazidime/avibactam as an anti-*Mycobacterium*

631   *avium* agent. Journal of Antimicrobial Chemotherapy 72:36-42.

632   67.   Ruth MM, Magombedze G, Gumbo T, Bendet P, Sangen JJN, Zweijpfenning S,

633   Hoefsloot W, Pennings L, Koeken V, Wertheim HFL, Lee PS, van Ingen J, Deshpande D.

634   2019. Minocycline treatment for pulmonary Mycobacterium avium complex disease

635   based on pharmacokinetics/pharmacodynamics and Bayesian framework mathematical

636   models. J Antimicrob Chemother doi:10.1093/jac/dkz143.

637   68.   Deshpande D, Kuret D, Cirrincione K, Cotroneo N, Melnick D, Lister T, Stokes S, Gumbo

638   T. 2020. 1659. Pharmacokinetics/pharmacodynamics of the Novel Gyrase Inhibitor

639   SPR719/SPR720 and Clinical Dose Selection to Treat Pulmonary Mycobacterium

640   avium-complex Disease. Open Forum Infectious Diseases 7:S817-S817.

641   69.   Athale S, Chapagain M, Pasipanodya J, Howe D, Alley MRK, Gumbo T. 2022. 1698.

642   Pharmacokinetics/pharmacodynamics of Epetraborole, a Novel Bacterial Leucyl-tRNA

643   Synthetase Inhibitor, and High Intracellular Penetration in the Intracellular Hollow Fiber

644   System Model of Mycobacterium avium Complex Lung Disease. Open Forum Infectious

645   Diseases 9.

646   70.   Chapagain M, Athale S, Pasipanodya J, Howe D, Alley MRK, Gumbo T. 2022. 1697.

647   Dose-response Studies of the Novel Bacterial Leucyl-tRNA Synthetase Inhibitor,

Epetraborole, in the Intracellular Hollow Fiber System Model of Mycobacterium avium

complex Lung Disease. Open Forum Infectious Diseases 9.

71.   Chapagain M, Pasipanodya JG, Athale S, Bernal C, Trammell R, Howe D, Gumbo T.

2022. Omadacycline efficacy in the hollow fibre system model of pulmonary

Mycobacterium avium complex and potency at clinically attainable doses. J Antimicrob

Chemother 77:1694-1705.

72.   Singh S, Boorgula GD, Aryal S, Philley JV, Gumbo T, Srivastava S. 2024. Sarecycline

pharmacokinetics/pharmacodynamics in the hollow-fibre model of Mycobacterium avium

complex: so near and yet so far. J Antimicrob Chemother 79:96-99.

73.   Deshpande D, Magombedze G, Boorgula GD, Chapagain M, Srivastava S, Gumbo T.

2023. Ceftriaxone efficacy for Mycobacterium avium complex lung disease in the hollow

fiber and translation to sustained sputum culture conversion in patients. J Infect Dis

doi:10.1093/infdis/jiad545.

74.   Deshpande D, Srivastava S, Gumbo T. 2024. Ertapenem's therapeutic potential for

Mycobacterium avium lung disease in the hollow fiber model. Int J Antimicrob Agents

doi:10.1016/j.ijantimicag.2024.107204:107204.

75.   Deshpande D, Magombedze G, Srivastava S, Gumbo T. 2024. Antibacterial action of

penicillin against Mycobacterium avium complex. IJTLD Open 1:362-368.

76.   Deshpande D, Srivastava S, Gumbo T, Martin K. Tiger Roars: Tigecycline

Pharmacokinetics/Pharmacodynamic [PK/PD] Based Treatment of Pulmonary

Mycobacterium Avium Complex [MAC] Disease, p A4349-A4349, B106 Breakthroughs in

NTM Diagnosis and Treatment. American Thoracic Society. doi:10.1164/ajrccm-

conference.2020.201.1_MeetingAbstracts.A4349.

77.   Deshpande D, Srivastava S, Gumbo T. 2025. Pharmacodynamic Evidence for Tedizolid

Use in Mycobacterium avium Lung Disease. IJTLD Submitted.

78.   Chapagain M, Pasipanodya JG, Athale S, Bernal C, Trammell R, Howe D, Gumbo T.

2022. Omadacycline efficacy in the hollow fibre system model of pulmonary

Mycobacterium avium complex and potency at clinically attainable doses. J Antimicrob

Chemother 77:1694-1705.

79.   Srivastava S, Deshpande D, Gumbo T. 2017. Failure of the azithromycin and ethambutol

combination regimen in the hollow-fibre system model of pulmonary Mycobacterium

avium infection is due to acquired resistance. J Antimicrob Chemother 72:i20-i23.

80.   Deshpande D, Srivastava S, Pasipanodya JG, Lee PS, Gumbo T. 2017. A novel

ceftazidime/avibactam, rifabutin, tedizolid and moxifloxacin (CARTM) regimen for

pulmonary *Mycobacterium avium* disease. J Antimicrob Chemother 72:i48-i53.

81.   Srivastava S, Deshpande D, Sherman CM, Gumbo T. 2017. A 'shock and awe'

thioridazine and moxifloxacin combination-based regimen for pulmonary Mycobacterium

avium-intracellulare complex disease. J Antimicrob Chemother 72:i43-i47.

82.   Ruth MM, Raaijmakers J, van den Hombergh E, Aarnoutse R, Svensson EM, Susanto

BO, Simonsson USH, Wertheim H, Hoefsloot W, van Ingen J. 2022. Standard therapy of

Mycobacterium avium complex pulmonary disease shows limited efficacy in an open

source hollow fibre system that simulates human plasma and epithelial lining fluid

pharmacokinetics. Clin Microbiol Infect 28:448.e1-448.e7.

83.   Schildkraut JA, Raaijmakers J, Aarnoutse R, Hoefsloot W, Wertheim HFL, van Ingen J.

2023. The role of rifampicin within the treatment of Mycobacterium avium pulmonary

disease. Antimicrob Agents Chemother 67:e0087423.

84.   Salillas S, Raaijmakers J, Aarnoutse RE, Svensson EM, Asouit K, van den Hombergh E,

695   Te Brake L, Stemkens R, Wertheim HFL, Hoefsloot W, van Ingen J. 2024. Clofazimine

696   as a substitute for rifampicin improves efficacy of Mycobacterium avium pulmonary

697   disease treatment in the hollow-fiber model. Antimicrob Agents Chemother 68:e0115723.

698   85.   Raaijmakers J, Ruth MM, Schildkraut JA, Hombergh EVD, Aarnoutse RE, Svensson EM,

699   Wertheim HFL, Hoefsloot W, van Ingen J. 2025. Replacing rifampicin with minocycline

700   increases the activity of the treatment regimen for Mycobacterium avium complex

701   pulmonary disease in a dynamic hollow-fibre system. Int J Antimicrob Agents 65:107423.

702   86.   Gumbo T. 2010. New susceptibility breakpoints for first-line antituberculosis drugs based

703   on antimicrobial pharmacokinetic/pharmacodynamic science and population

704   pharmacokinetic variability. Antimicrob Agents Chemother 54:1484-91.

705   87.   Gumbo T, Pasipanodya JG, Wash P, Burger A, McIlleron H. 2014. Redefining multidrug-

706   resistant tuberculosis based on clinical response to combination therapy. Antimicrob

707   Agents Chemother 58:6111-5.

708   88.   Gumbo T, Chigutsa E, Pasipanodya J, Visser M, van Helden PD, Sirgel FA, McIlleron H.

709   2014. The pyrazinamide susceptibility breakpoint above which combination therapy fails.

710   J Antimicrob Chemother 69:2420-5.

711   89.   Deshpande D, Pasipanodya JG, Srivastava S, Bendet P, Koeuth T, Bhavnani SM,

712   Ambrose PG, Smythe W, McIlleron H, Thwaites G, Gumusboga M, Van Deun A, Gumbo

713   T. 2018. Gatifloxacin Pharmacokinetics/Pharmacodynamics-based Optimal Dosing for

714   Pulmonary and Meningeal Multidrug-resistant Tuberculosis. Clin Infect Dis 67:S274-

715   S283.

90.   Deshpande D, Pasipanodya JG, Mpagama SG, Bendet P, Srivastava S, Koeuth T, Lee

PS, Bhavnani SM, Ambrose PG, Thwaites G, Heysell SK, Gumbo T. 2018. Levofloxacin

Pharmacokinetics/Pharmacodynamics, Dosing, Susceptibility Breakpoints, and Artificial

Intelligence in the Treatment of Multidrug-resistant Tuberculosis. Clin Infect Dis 67:S293-

S302.

91.   Deshpande D, Pasipanodya JG, Mpagama SG, Srivastava S, Bendet P, Koeuth T, Lee

PS, Heysell SK, Gumbo T. 2018. Ethionamide Pharmacokinetics/pharmacodynamics-

derived dose, the role of MICs in clinical outcome, and the resistance arrow of time in

multidrug-resistant tuberculosis. Clin Infect Dis 67(suppl_3):S317-S326.

92.   Årdal C, Balasegaram M, Laxminarayan R, McAdams D, Outterson K, Rex JH,

Sumpradit N. 2020. Antibiotic development - economic, regulatory and societal

challenges. Nat Rev Microbiol 18:267-274.

Table 1. Final Proposed Quality Scores for preclinical pharmacokinetics/pharmacodynamics criteria
729

Criterion	No	1	2	3	4	5	Total Possible Score
EXPOSURE EFFECT							
B_0: 0 score if not reported 1 if density is reported but is outside the range seen in patients 2 if reported and within range encountered in patients 3 if both density and total bacterial burden are reported	0	1	2	3			3
Number of exposures: <5 exposures =0, 5 exposures =1 6 exposures =2 7 exposures =3 8 exposures =4 At least one exposure of steep portion = 1 extra point	0	1	2	3	4	5	5
Dose fractionation study was performed: 0 if none, 2 if fractionation for at least 2 dose, 3 if at least 3 doses fractionated	0		2	3			3
Monte Carlo Experiments: score by number of doses tested	0	1	2	3	4	5	5
Number of non-lab-reference (ATCC) isolates for PK/PD target	0	1	2	3	4	5	5
Number of replicates: One =0 2 replicates=1 3 replicates in dose response=2 If more than 1 replicate for both reference lab strain and the clinical isolates = 3	0	1	2	3			3
Total Possible Score							**24**
COMBINATION (similar to exposure-effect but replace [1] number of exposures, [2] dose-fractionation, and [3] Monte Carlo Experiments with-							
Score: 0 if one dose/exposure of each drug was used, 3 if at least 3 non-zero exposures were used for each drug, 4 if 2 replicates were used, and 5 if at least 3 replicates were used.	0	1	2	3	4	5	5
Score: 0 if there is no synergy or antagonism interaction index 3 if there is an interaction index calculated, 4 if the optimal exposures of each component for microbial kill calculated, 5 if optimal exposures are calculated for both microbial kill and resistance suppression	0	1	2	3	4	5	5
Total Possible Score							**21**

732 **Table 2. Qualitative Systematic Analyses Findings and Quality of Studies by Year of Publication**

Ref.	Type of model	Year	Drug	Findings and conclusions	Quality Score	Study Quality
(60)	HFS-MAC	2010	Ethambutol	1. First HFS-MAC published, 7-day exposure-effect study 2. Studies of static concentrations of ethambutol with extracellular MAC over-estimate efficacy versus HFS-MAC 3. C_{max}/MIC linked efficacy and resistance	16	Good
(61)	HFS-MAC	2010	Moxifloxacin	1. 7-day exposure-effect study 2. Intracellular penetration overcomes protein binding 3. Proposed susceptibility breakpoint meant most clinical isolates resistant to moxifloxacin	11	Adequate
(12, 62)	HFS-MAC	2012	Azithromycin	1. First 28-day exposure-effect study and 28 day dose fractionation study 2. The Antibiotic resistance arrow of time model proposed 3. Microbial kill and resistance linked to AUC/MIC, however on day 7 linked to % time above MIC (%T_{MIC}) based on reading off the curves 4. Standard clinical doses sub-optimal	16	Good
	HFS-MAC	2016	Thioridazine	1. Massive intracellular concentrations due to lysosomal trapping 2. C_{max}/MIC linked efficacy and resistance based on day 7 results 3. Based on day 5 and day 14 results pattern was AUC/MIC driven microbial kill	11	Adequate
	HFS-MAC	2017	Linezolid	1. Linezolid bactericidal effect better than GBT components 2. In Monte Carlo experiments ~50% of clinical isolates resistant to linezolid	17	Good
(65)	HFS-MAC	2017	Tedizolid	Tedizolid killed more than 2.0 $\log_{10}$ cfu/mL below B_0, highest seen yet	10	Adequate
(66)	HFS-MAC	2017	Ceftazidime/ avibactam	1. First β-lactam shown effective against MAC, at clinically achievable exposures 2. Ceftazidime was enhanced by avibactam 3. Efficacy higher than azithromycin and ethambutol.	15	Adequate
	HFS-MAC	2019	Minocycline	1. Minocycline is expected to be bactericidal at 400 mg/day 2. Use of Bayesian framework bacterial dynamics mathematical model based on competition model for drug-susceptible and resistant isolates	13	Adequate
(68)	HFS-MAC	2020	SPR720	1. PK/PD parameter linked to microbial kill was AUC_{0-24}/MIC 2. High intracellular penetration	20	High
(69, 70)	HFS-MAC	2022	Epetraborole (EBO)	1. At the EC_{80}, EBO killed at least 2.0 $\log_{10}$ CFU/mL below B_0 2. EBO + GBT demonstrated resistance suppression and faster microbial kill 3. Both microbial kill and AMR were AUC_{0-24}/MIC linked	24	High
(56, 57)	C57BL/6 mice	2022	Epetraborole	1. Epetraborole demonstrated potent in vivo efficacy against <u>5 MAC isolates</u> and significantly improved efficacy when combined with GBT 2. Epetraborole monotherapy better than GBT 3. No resistance emergence	20	Good

(56, 57)	C57BL/6 mice	2022	Epetraborole + GBT	More extensive microbial kill for epetraborole and GBT compared to either alone	15	Good
(71)	HFS-MAC	2022	Omadacycline	1. Omadacycline concentration declines by 50% every 24h in MIC assay versus MAC doubling time of 16.3h, artefactually increasing MICs 2. γ-slope-based PD modeling more robust than day-to-day CFU/mL 3. Omadacycline more effective than GBT in same study	14	Adequate
(72)	HFS-MAC	2024	Sarecycline	1. Due to high MICs and poor lung penetration EC_{80} unachievable 2. "Paradoxical effect" observed with the AUC_{0-24}/MIC of 285.39 exposure	8	Poor
	HFS-MAC	2024	Ceftriaxone	1. Exposure-response with reference MAC strain 2. GBT (azithromycin, ethambutol, rifabutin) killed 2 of <u>5 MAC clinical isolates</u>, was partially successful in one, and failed in 2 3. Once-daily ceftriaxone monotherapy killed all for the 5 MAC below B_0 4. ODEs identified γ-slopes as PD parameter in 5 MAC strains 5. γ-slopes translated to SSCC in patients in 1,000 virtual patients, and ceftriaxone monotherapy predicted to shorten SSCC 2.35-fold	20	Good
	HFS-MAC	2024	Ertapenem	1. E_{max} and EC_{50} was better without avibactam than with avibactam 2. Ertapenem microbial kill below B_0 better than GBT drug monotherapies	15	Good
	HFS-MAC	2024	Benzylpenicillin	1. Better EC_{50} without avibactam than with avibactam 2. Use of ODEs and γ-slopes that include model for resistance 3. Benzylpenicillin $\gamma=0.98$ (0.95–0.99) >azithromycin $\gamma=0.74$(0.65-0.86) and = azithromycin+ ethambutol $\gamma=0.94$ (0.82-0.99)	16	Good
(76)	HFS-MAC	2025	Tigecycline	1. MICs of a drug not good indicators of potency or efficacy for MAC drug development because MAC is mostly intracellular 2. Static concentrations studies (e.g. time-kill) provide limited to no information as regards to PK/PD target exposures or efficacy 3. PK/PD target identified with ATCC reference strain was biased and imprecise 4. Inclusion of 5 clinical isolates gives a more robust target selection and estimate of E_{max} 5. Inhalation therapy was studied	18	Good
	HFS-MAC	2025	Tedizolid	1. Inclusion of 5 clinical isolates gives a more robust target selection that was 7 times higher than reference lab isolate (65) 2. 5 strains gave a more robust estimate of kill below B_0 that was very different from that derived with laboratory reference isolate (65)	20	Good
	BALB/c mice	2024	Bedaquiline + clarithromycin	Bedaquiline plus clarithromycin combination therapy demonstrated enhanced efficacy compared to monotherapies	8	Poor
This study	HFS-MAC	2025	Clarithromycin	A total of 32 replicates, used to propose quality control criteria	NA	NA

734
735

736 **Table 3. Quantitative Systemic Analysis Findings and Ranking of Efficacy of Drugs for Monotherapy**

Drug [reference]	Weighted E_{max} (log_{10} CFU/mL kill below B_0)	Fold of 3-Drug GBT Efficacy	PK/PD Target & Dosing Schedule	Monte Carlo Experiments-generated Optimal Dose (mg per day)	Route	PK/PD breakpoint (MIC; mg/L)	Predicted Tolerability of Dose
MONOTHERAPY							
HFS-MAC							
Thioridazine (63)	4.68	227	C_{max}/MIC=0.88	None	Oral	None	Poor
Omadacycline (71)	4.16	68.63	AUC_{0-24}/MIC=9.9	300	Oral	16	Good
Tigecycline (76)	4.14	65.576	AUC_{0-24}/MIC=33.65	35	Inhaled		Good
Tedizolid (77)	3.61	19.31	AUC_{0-24}/MIC=155.5	200	Oral	8	Good
Ceftriaxone (73)	3.22	7.92	%T_{MIC}=100%	2,000	IV		Good
Minocycline (67)	2.9	3.77	AUC_{0-24}/MIC=63.1	200	Oral	8	Good
Moxifloxacin (61)	2.30	0.95	AUC_{0-24}/MIC=163	1,600	Oral	0.25	Poor
Ceftazidime/avibactam (66)	1.74	0.26	%T_{MIC}=52%	NA	NA	-	Good
SPR720 (68)	1.54	0.17	AUC_{0-24}/MIC=4.09	1,000	Oral	4	Good
Ertapenem (74)	1.47	0.14	%T_{MIC}=100	2,000 twice daily	IV	4	Good
Benzylpenicillin (75)	1.4	0.12	%T_{MIC}=84.6%	1,200	IV	64	Good
Sarecycline (72)	0.7	0.02	AUC_{0-24}/MIC=139.5	None		None	Poor
Clarithromycin (this study)	0.64	0.02	-	-	Oral		Poor
Linezolid (64)	0.38	0.01	AUC_{0-24}/MIC=43.7	1,800	Oral	64	Very poor
Azithromycin (12, 62)	0	<0.01	AUC_{0-24}/MIC=7.5	8,000	Oral	16	Poor
Ethambutol (60)	0		C_{max}/MIC=1.23	50 (mg/kg)	Oral	2	Poor
MOUSE							
Epetraborole					Oral		
COMBINATION THERAPY							
3-drug GBT in HFS-MAC (73)	211.0	1.0			Oral		Poor
Epetraborole in mice					Oral		
Bedaquiline + clarithromycin in mice					Oral		

IV= intravenous

Figure 1. PRISMA Flow Diagram for Studies

778 **Figure 2. Hollow fiber system model of MAC Quality Scores**

779
A. Quality scores for all 43 studies. **B.** Correlation coefficients (Pearson) between different

scoring criteria. **C.** Quality scoring with final tool, with only the studies that met inclusion criteria.

D. Bland-Altman agreement analysis between the observers for the quality scores. The average

score among all observers is shown on x-axis, while the difference between average and

observer are on the Y-axis.

Figure 3. Drugs studied in HFS-MAC and murine MAC-LD studies

787
788   **A.** Number of drugs in studies that fulfilled study acceptance criteria. **B.** Percentage of drugs

789   studied, by pharmacophore.

790   **Figure 4. Clarithromycin effect in ATCC isolate versus a panel of 5 clinical isolates**

791
792   Symbols depict mean values. Error bars are standard deviation. **A.** Effect of clarithromycin on

793   treatment with doses achieving EC_{80} in the lungs. **B.** Coefficient of variation of CFU/mL between

794   replicates.