From Concrete to Canopy: Illuminating Moth Biodiversity in New York City's Urban Jungle

Shira Linsk[1], Anna Thonis[1], & Kristin M. Winchell[1,2]

[1]New York University, Department of Biology

[2]Correspondence: kristin.winchell@nyu.edu

Author contact information: slinsk@uchicago.edu, at6689@nyu.edu, kristin.winchell@nyu.edu

Data Availability: All data and associated code are archived on Zenodo with doi:

10.5281/zenodo.14756963

Funding statement: This work was supported by New York University College of Arts and

Science's URA and DURF programs (Undergraduate Research Assistant, Dean's

Undergraduate Research Fund)

Conflict of interest disclosure: The authors declare no conflicts of interest.

RRH: URBAN MOTH BIODIVERSITY

Words: 4,473

0 tables, 5 figures

Keywords: Lepidoptera, moths, urbanization, ALAN, urban greenspace, invertebrates, urban ecology, biodiversity, path analysis

Abstract

Moths (Lepidoptera) are sensitive to anthropogenic threats and serve as valuable

bioindicators. Despite the remarkable diversity and abundance of Lepidoptera globally, there is a

lack of information on how moth species are impacted by urbanization. Notably, very little is

known about moths in the largest city of the United States, New York City, where pervasive

urban pollutants, artificial light at night, land cover change, and habitat fragmentation are severe.

We examined the effects of urbanization on moth biodiversity in New York City, with a focus on

green spaces. We used citizen science records from *iNaturalist* and complemented these data

with ground sampling at twelve locations across six parks at night. While the *iNaturalist* dataset

is comprehensive both spatially and temporally, it failed to detect rarer species we observed on

the ground. However, the scope of the field survey dataset is limited in geographical breadth and

seasonal coverage. Overall, we found a negative relationship between greater urbanization and

moth diversity, with community similarity correlated with environmental similarity. Our results

found greater biodiversity with less light at night and urban development, and more deciduous

tree cover and open land. Our structural equation model reveals additional insight: although we

detected a strong direct negative effect of developed land on moth diversity, urbanization also

negatively impacts diversity via indirect effects of reducing open space and deciduous tree cover.

Developed open space alone does not directly affect diversity but may positively impact

diversity through its covariance with vegetation cover. These findings support the importance of

mitigating artificial light at night in urban green spaces and maintaining urban vegetation to

ensure nocturnal Lepidoptera can persist in rapidly urbanizing landscapes.

Introduction

Urbanization is linked to declines in biodiversity through biotic homogenization and

species extirpation [1-2]. Although these impacts are seen across the tree of life, the effects of

urbanization on invertebrates have been particularly drastic, with steep declines across insect

orders [3-4]. Studies have demonstrated local extinctions of insect populations and reduced

insect diversity with increasing urbanization [5-6]. Much of this work has focused on bees and

butterflies, revealing decreased diversity and abundance with increased urbanization [5, 7-9].

Urbanization presents many challenges for insects, including the urban heat island effect, habitat

fragmentation, pollution (including artificial light), impervious surfaces, and exotic plants [10].

Habitat fragmentation and landscape maintenance are major contributors to pollinator decline,

including Lepidoptera [11]. However, alongside these challenges come opportunities in terms of

anthropogenic resources that may be exploited, as seen in common urban species like

cockroaches, flies, and carpet beetles [12]. Many species may thrive in cities despite the threats

of pest control and habitat modification. For example, urban plantings may generate habitat and

resources that support invertebrate diversity, including Hymenoptera and spiders [13-16].

Relatively little research on urban invertebrates has focused on moths, and there is a

notable bias toward butterflies and other diurnal Lepidoptera [7, 17-18]. This is perhaps due to

the challenges associated with observing moths at night (e.g., low visibility, general safety

considerations) and the minimal public engagement surrounding them compared to butterflies

[19]. Some recent studies on moths have found decreasing populations with increased artificial

light at night [20-22]. Additionally, anthropogenic environments generate intense selection

pressures and filtering processes that may favor the survival of only a few species, such as heat-

tolerant moths [23] or produce phenotypic shifts, such as increased body size in urban moths

[24]. Indeed, studies of industrial melanism in moths represent some of the first examples of

rapid evolutionary change in cities [25-27].

Understanding the effects of urbanization on moths is important because of the crucial

roles they play in ecosystem services (e.g., pollination, food, and nutrient cycling). Moths are

fundamental to ecosystem health and stability and serve as bioindicators [22, 28-29] since they

respond quickly to environmental change [20, 30-31]. There are approximately 150,000 known

species of moths globally [32], and hundreds of new species are described annually [33]. Yet, we

know very little about the species that persist in urban landscapes. For conservation initiatives to

be successful in these spaces, we must first take stock of the species that reside there and relate

patterns of diversity to urban stressors.

We investigate patterns of moth diversity in New York City, where artificial light at

night, pollution, and habitat fragmentation are extreme. We hypothesize that landscape-scale

features influence species diversity and predict that developed land use negatively impacts moth

diversity. In addition, we hypothesize that habitat-scale features such as pollution, light at night,

and microclimate correlate with moth diversity. By connecting moth diversity to urban habitat

features, we aim to improve our understanding of the ecological impacts of urbanization on

nocturnal Lepidoptera species in one of the largest urban centers globally.

Methods

Statistical analyses were conducted using R studio (ver. 2023.12.0.369) [34] and R (ver.

4.3.2) [35]. For all linear models, we evaluated multicollinearity with the 'VIF' function in the

package *car* [36] assessed statistical significance using t-tests for individual coefficients, and

evaluated model fit based on Akaike Information Criterion (AIC) and residual deviance.

iNaturalist data

We downloaded 31,793 Research-Grade observations of the Order Lepidoptera using a

polygon spatial filter around all five New York City boroughs. We manually removed butterfly

Families (i.e., Papilionidae, Nymphalidae, Pieridae, Lycaenidae, Hesperiidae), resulting in

12,956 moth observations. We imported records into QGIS (ver. .3.34.3-Prizren) [37] and used

the New York City administrative boundary layer [38] to reduce this dataset to 10,740 records

located within New York City boundaries. We used the plugin *Density Analysis* [39] to quantify

observation density using a grid of 1 km hexagonal cells across the region (Fig. 1). We excluded

hexagonal zones (33 zones) with undefined landscape types or >70% water cover (according to

the National Land Cover Database "NLCD" [40]), to ensure the zones had relevant habitat data

for terrestrial species. We filtered the *iNaturalist* records to include observations from 2000

onwards to better capture moth diversity over time, rather than at one specific point in time.

Additionally, we removed moth observations that we ourselves contributed. This resulted in a

total of 9,374 records.

Figure 1. Study sites and moth diversity patterns across New York City (a) On-the-ground

sampling locations for 12 locations in Brooklyn and Queens, New York City, USA, where we

sampled Lepidoptera, and National Land Cover (NLCD) data for New York City. (b) Species

diversity in 1-km hexagonal bins derived from iNaturalist data. Base maps by Open Street Map.

Field Site Selection

To complement our *iNaturalist* dataset and provide field-based comparisons with any

observed trends, we identified 12 on-the-ground sampling sites in six greenspaces across

Brooklyn and Queens, New York, that reflect variation in urban-associated land cover and

habitat features encountered across the city (Fig. 1). We visually assessed satellite imagery to

identify green spaces and then chose sampling locations within each space spanning a gradient in

light and air pollution based on on-the-ground measurements (rather than remote-sensed) to

capture variation in this variable at an organismally relevant spatial scale.

Field Sampling

We surveyed each location for one hour shortly after dusk (8-10 pm) between March 3-

April 29, 2024. The exact time at which sampling occurred varied throughout our survey

timeframe as sundown began later into the evening as the season progressed. We restricted the

temporal sampling period to minimize seasonal variation in species detected across sites. We

constructed a light trap using a suspended 5 m^2 white sheet directly illuminated with two flood

lights (6000 Kelvin, visible light spectrum) and one ultraviolet LED light (385-395 nm) (as in

[41]). Moths are easily sampled with light traps, allowing for robust estimates of species richness

and abundance [30]. Each survey consisted of two people photographing all moths observed

landing on or in the immediate vicinity (on illuminated ground within 2m) of the light trap. We

identified species from the photographs and counted the total number of individuals per species

encountered during each sampling event. In addition, for each sampling event, we measured the

following site-level variables: air pollution (PM 2.5; Temtop M2000C), light at night (lux;

EXTECH Light Meter LT300), temperature, humidity, and wind speed (Kestrel 3000).

Landscape analysis

We mapped *iNaturalist* records and on-the-ground sampling locations using QGIS. We

extracted zonal statistics of NLCD land cover classifications [40] within each 1 km hexagonal

zone for the *iNaturalist* dataset. For on-the-ground sampling locations, we used a 150 m buffer

around each sampling point to capture the immediate landscape characteristics of each site and at

a spatial scale relevant to Lepidoptera community structure (100-200 m) [42]. We calculated the

percentage of each land cover class for hexagonal zones and buffered areas. We summarized

percentage of land cover classifications (i.e., Deciduous, Mixed Forest, Grassland, Pasture/Hay,

Developed Open Space, Developed Low-Intensity, Developed Medium-Intensity, Developed

High-Intensity) with principal component analysis (PCA) using the function 'prcomp' in base R

stats [35]. We visualized results with packages *factoextra* [43] and *corrplot* [44].

Statistical analyses

127   We summarized species diversity with Simpson's Diversity Index (Simpson's D), which

considers both species richness and evenness [45]. Simpson's Diversity Index is robust in urban

settings as it is not sensitive to rare taxa that may be under-sampled or absent in disturbed

habitats where common generalist species might dominate [23]. Additionally, this index can

overcome issues of small sample size and spatial biases [46]. We calculated Simpson's D (1-D)

using species-level identification for each *iNaturalist* hex bin and each on-the-ground sampling

location in R using the function 'diversity' in the package *vegan* [47].

134   We conducted three analyses using the *iNaturalist* data. First, we used multiple matrix

regression (MRM) to analyze the similarity of community composition with respect to

environmental similarity across sites This entailed generating a distance matrix (Euclidean

Distance) of species composition similarity across hexagonal zones, as well as for the

environment (percentages of all NLCD land cover classes). We used the function 'dist' in the R

base *stats* for all distance matrix analyses [35] and performed matrix regression using the

function 'MRM' in R package *ecodist* [48]. Second, we used linear regression to analyze the

relationship between moth diversity (Simpson's D) and landscape-level (remotely sensed) habitat

features. Explanatory variables in the model included NLCD categories (as a proportion of hex-

bin area): deciduous tree cover, developed land cover (high-, medium-, low-intensity), pasture,

developed open land, shrub, and grassland. We simplified this model using stepwise model

simplification (backwards and forwards) based on AIC using the function 'step' in R base *stats*.

146   In addition, we developed a structural equation model (SEM) using *iNaturalist*

observations to examine relationships between moth diversity, vegetation (deciduous forest,

grassland, pasture, shrub), developed land (high-, medium-, low-intensity), and developed open

land cover. SEMs analyze multiple relationships at the same time while considering latent factors

(i.e., unobserved relationships, such as between deciduous land cover and grassland). We

specified the model to reflect the hypothesis that different components of the urban landscape

(i.e., vegetation, developed land, and open space) each exert direct effects on moth diversity, and

that these land cover types may covary due to the spatial structure of urban environments (e.g.,

lower vegetation cover in highly developed areas). Latent variables were used to group related

land cover types, reducing collinearity and capturing broader ecological gradients. We fit our

SEM using maximum likelihood with the 'sem' function in the *lavaan* R package (ver. 0.6-19)

[49]. Model fit was assessed using Chi-squared (χ^2), Comparative Fit Index (CFI), Tucker-Lewis

Index (TLI), and Root Mean Square Error of Approximation (RMSEA). To account for shared

variance between variables, we specified covariances between vegetation cover and development

intensity, development intensity and developed open space, and between several vegetation

variables, including between deciduous and grassland, and between grassland and shrub. We

used the z-statistic to assess the significance of each path coefficient in the SEM and visualized

observed relationships with the function 'semCors' in the R package *semPlot* [50].

164   We conducted two additional analyses using our on-the-ground field sampling data. We

again analyzed the similarity of community composition concerning environmental similarity

across sites using MRM. As with the *iNaturalist* MRM analysis, this entailed generating a

distance matrix of species composition similarity across sampling sites, and for the environment:

percentages of all NLCD land cover classes, as well as PM2.5, lux, temperature, humidity, and

wind speed measured on the ground at time of sampling. Second, we used linear regression to

analyze the relationship between moth diversity and site-level habitat features. We did not

include remotely sensed land cover data in this model because the resolution of these data and

our sample size resulted in low statistical power (overparameterization) and multicollinearity

when these variables were included. Explanatory variables used in our model included

temperature, pollution (PM 2.5), and light intensity (lux) measured during on-the-ground

surveys. We simplified this model using stepwise model simplification (backwards and

forwards) based on AIC using the function 'step' in R base *stats*. Lastly, we combined

iNaturalist data with our on-the-ground survey data to evaluate the relationship between

diversity and urbanization more broadly. Specifically, we analyzed land cover PCA, with no

interactions or model simplification.

Results

Urban environmental variation

183   Land cover analyses using remotely sensed data verified that our on-the-ground sampling

locations spanned a broad range of urbanization (Fig. 2a). Larger parks (Forest Park, Highland

Park, Prospect Park) exhibited greater habitat heterogeneity and had more deciduous forest

cover. Small parks (Maria Hernandez, Grover Cleveland, Irving Square), by contrast, had little to

no deciduous tree cover and were composed almost entirely of developed land. Grover Cleveland

Park stood out with only a small percentage of high-intensity developed land as it borders a

private cemetery with considerable greenspace. These differences are clearly visible in satellite

imagery: Forest Park is dominated by forest, Prospect Park has less tree cover, and Maria

Hernandez is substantially smaller and consists mostly of developed land (Fig. 2c).

Figure 2. Land cover composition, principal components analysis, and site imagery of sampling

locations (a) Stacked bar plot of NLCD land cover composition at on-the-ground sampling sites.

Colors represent the proportion of each land cover type in 150 m buffers around each sampling

location. (b) Principal components analysis of NLCD land cover for our on-the-ground sampling

locations and all hex-bins used with iNaturalist data. PC1 represents high-intensity developed

land cover. (c) Satellite images of sampling locations in Brooklyn and Queens, New York.

Our PCA incorporated land cover for both *iNaturalist* records and on-the-ground sites

(1115 *iNaturalist* hexagonal zones, 12 on-the-ground sampling locations). The PCA captured

95.31% of the variance in the first four principal components, with PC1 capturing 44.68% of the

variance. Visual inspection of the scree plot suggests that four principal components should be

retained to describe variation in the dataset. The first and second principal components (PC1 and

PC2) represent urbanization, with strong positive loadings for developed high-intensity land

cover on PC1 (Fig. 2b) and for medium-intensity developed land cover on PC2. The third

component (PC3) was negatively correlated with water and positively correlated with developed

open space and deciduous tree cover. The fourth principal component (PC4) was positively

correlated with developed open space and developed low-intensity land cover and negatively

associated with deciduous tree cover. Grassland, pasture, and shrub land cover classifications did

not load strongly on any of the first four principal components.

Diversity of Lepidoptera

214   From *iNaturalist*, we extracted data from 9,377 Research Grade observations,

representing 972 species, 535 genera, and 50 families (Fig. 3b). In our on-the-ground surveys,

we observed a total of 69 individuals, representing 11 families, 31 genera, and 33 species (Fig.

3a). Observations were identified to species in 100% of samples. When comparing *iNaturalist*

moth richness with surveyed richness, *iNaturalist* richness was markedly higher than that of our

surveys for Prospect Park (110 species on *iNaturalist* versus 12 in our sampling), but similar for

Grover Cleveland (two on *iNaturalist* versus three). By contrast, in two parks, we detected more

species in our on-the-ground sampling: Forest Park (29 on *iNaturalist* versus 35) and Highland

Park (nine on *iNaturalist* versus 19). In the two parks where no species were detected in our on-

the-ground sampling (Maria Hernandez and Irving Square), there were also no observations on

iNaturalist. Notably, we detected individuals from the family Eriocraniidae in our on-the-ground

sampling, but this family was not present in the *iNaturalist* dataset. Conversely, *iNaturalist*

observations documented 40 additional families not detected in our on-the-ground sampling.

Figure 3. Lepidoptera family diversity from field sampling and iNaturalist data (a) Stacked bar

plot of Lepidoptera Family diversity per site for on-the-ground sampling. (b) Number of

observations of each Lepidoptera Family across all iNaturalist hex bins. Bars are colored by

Families observed. (c) Images of moths taken during on-the-ground sampling.

Based on our on-the-ground sampling, large parks (Forest Park, Highland Park, Prospect

Park) displayed the highest levels of biodiversity (1-D: 0.44–0.92, mean±sd: 0.72±0.15). Among

them, Forest Park had a mean diversity index of 0.80±0.1, Highland Park had a mean diversity of

0.71±0.19, and Prospect Park had a mean diversity of 0.64±0.17. In contrast, the small parks

(Maria Hernandez, Grover Cleveland, Irving Square) exhibited lower diversity (1-D: 0–0.66,

mean±sd: 0.22±0.38). No species were detected at Maria Hernandez and Irving Square in our on-

the-ground sampling; since Simpson's Diversity Index is undefined in the absence of

observations, we recorded 1-D as 0 for these sites.

Community Composition

In both our *iNaturalist* data and on-the-ground sampling, locations with similar

environments had similar community composition at the species level (multiple-matrix

regression; on-the-ground: F=43.127, R^2=0.501, p=0.031; *iNaturalist*: F=3892.81, R^2=0.013,

p<0.001). Species diversity is strongly negatively correlated with PC1 and PC2 (representing

increased high- and medium-intensity development, respectively) across both datasets combined

(PC1 estimate=-0.200±0.041, t=-4.848, p<0.001; PC2 estimate=-0.449±0.052, t=-8.692,

p<0.001) (Fig. 4). Diversity was also positively correlated with PC3, representing less water and

more developed open space and deciduous tree cover (estimate=0.415±0.073, t=5.654, p<0.001).

Diversity was not significantly correlated with PC4 (t=0.046, p=0.963).

Figure 4. Correlations of Simpson's Diversity Index and NLCD land cover. Simpson's diversity

(D-1) as it relates to land cover summarized by principal components analysis: (a) PC1

representing high-intensity developed land, (b) PC2 representing medium-intensity developed

land, and (c) PC3 representing open developed land and deciduous tree cover.

Environmental Correlations

Our two linear models—one using *iNaturalist* observations with remotely sensed

environmental variation and the other using on-the-ground sampling with field-measured

environmental variation—revealed correlational relationships between environmental features

and biodiversity across both spatial scales. Our simplified linear regression of Simpson's

Diversity by land cover using *iNaturalist* and remotely sensed data retained five significant

variables: medium-intensity developed land, low-intensity developed land, developed open land,

shrubland, and deciduous tree cover. Diversity was negatively correlated with medium-intensity

developed land cover (estimate=-0.350±0.064, t=-5.493, p<0.001) and positively correlated with

deciduous tree cover (estimate=0.465±0.107, t=4.347, p<0.001) and developed open space

(estimate=0.375±0.118, t=3.169, p=0.002). There was also a near-significant positive trend

between diversity and low-intensity developed space (estimate=0.268±0.153, t=1.755, p=0.080),

and the correlation with shrubland was non-significant (t=1.598, p=0.110). In comparison, our

simplified regression of Simpson's Diversity for on-the-ground sampling retained two variables:

air pollution and light at night. Simpson's diversity was not significantly correlated with air

pollution (t=1.280, p=0.233) but was significantly negatively correlated with light at night

(estimate=-0.512±0.062 lux, t=-8.214, p<0.001).

Structural Equation Model

Our SEM exhibited an overall good fit to the data (CFI: 0.957, TLI: 0.918, RMSEA:

0.066). The chi-square statistic was significant (χ^2=80.896, df=19, p<0.001), although this is

expected with larger sample sizes, and the overall fit indices suggest the model effectively

captured the relationships in the data.

Our SEM detected a significant positive effect of vegetation cover on diversity

(estimate=1.905±0.398, z=4.791, p<0.001; Fig. 5) and a significant negative effect of developed

land on diversity (estimate=-0.301±0.101, z=-2.979, p=0.003). We did not find a significant

direct effect of developed open space (z=0.391, p=0.695) on diversity (Fig. 5). Significant

negative covariance was observed between vegetation cover and development intensity

(covariance=-0.012, z=-10.728, p<0.001), and significant positive covariance between vegetation

cover and open space (covariance=0.002, z=4.191, p<0.001). Similarly, we found a significant

negative covariance between development intensity and developed open space (covariance=-

0.014, z=-11.707, p<0.001). We also observed weak but significant positive covariance between

developed low-intensity and developed open space (covariance=0.005, z=9.382, p<0.001) and

grassland and shrub (covariance<0.001, z=10.954, p<0.001). We observed weak but significant

negative covariance between developed high-intensity and developed low-intensity

(covariance=0.015, z=-11.237, p<0.001), and deciduous and grassland (covariance<-0.001, z=-

2.650, p = 0.008). Further inspection of relationships among variables indicates that developed

medium-intensity land cover is driving the direct effect of development on diversity, whereas

high and low-intensity development impact diversity more indirectly via their relationships with

open land and deciduous tree cover. Further, among the vegetation variables, only deciduous tree

cover has a strong direct effect on diversity, whereas grassland, shrubland, and pasture all have

indirect effects related to their covariance with developed land.

Figure 5. Pathways linking urban land cover to Lepidoptera diversity revealed by structural

equation modeling. Relationships among predictor variables and Simpson's diversity generated

by the structural equation model. Nodes represent variables included in the model, while edges

indicate associations between them. Thicker lines represent stronger relationships, with positive

correlations shown in green and negative correlations in red. Edge weights are proportional to

the strength of standardized regression coefficients.

Discussion

Urbanization drastically alters landscapes, replacing naturally occurring habitats and

vegetation with impervious surfaces and buildings. This process not only impacts land cover but

also increases artificial light at night (ALAN), both of which can have significant consequences

for biodiversity. Lepidoptera (moths and butterflies) are highly sensitive to land cover changes

[31, 51], and are also influenced by habitat patch size, quality, and shape [52-53]. In this study,

we examined how urbanization relates to the diversity of nocturnal Lepidoptera (moths), which

perform vital ecosystem services and act as bioindicators [28]. Our findings suggest that

urbanization is associated with lower moth diversity, with specific elements of the urban

landscape shaping patterns of biodiversity in complex ways.

We found comparable moth community composition between locations with similar

environments in both *iNaturalist* and our on-the-ground sampling datasets, and a general trend of

decreasing diversity with increasing urbanization. The *iNaturalist* MRM had a low R^2 but was

highly significant, likely reflecting high spatial coverage but heterogeneous observer effort. The

on-the-ground MRM explained much more variation in community composition despite its

smaller sample size and its limited spatial and temporal scope, suggesting that controlled

sampling better captured ecological patterns among sites. These results suggest that moth

community composition across New York City is strongly influenced by the environmental

features that define different locations. Our linear models provide some insight into those

features: we found strong negative correlations between moth diversity and medium-intensity

urbanization, and strong positive correlations between diversity and deciduous tree cover and

open land. In addition, our on-the-ground sampling found a strong negative relationship between

moth diversity and light at night, with a high explained variance (R^2), suggesting light at night

accounts for a substantial portion of the observed variation in species diversity across sampled

locations. Although several studies have found no consistent effects of light at night on

biodiversity [11, 54], our results are in line with those that identify significant impacts on moth

community composition [55-57]. These negative effects on biodiversity are likely attributed to

how light at night disrupts moth feeding, reproduction, migration, and development [21].

Our PCAs of *iNaturalist* and on-the-ground data reinforce these findings: we found

strong negative correlations between diversity and the principal components primarily defined by

high- and medium-intensity development (i.e., PC1, PC2), and strong positive correlations with

the principal component representing deciduous tree cover and open space (i.e., PC3). The

repeated emergence of deciduous tree cover, open space, and high- and medium-intensity

developed land as key variables across multiple analyses highlights the importance of these

features. Our results are consistent with those of similar studies performed in metropolitan areas

where highly disturbed habitats have been shown to correlate with reduced moth species richness

[20, 23, 58].

While we did not perform a SEM using our on-the-ground data due to the larger sample

sizes required, our SEM using *iNaturalist* data and remotely sensed land cover found, similar to

our other analyses, that higher diversity is associated with greater vegetation cover and less

developed land. Further inspection of these relationships revealed that although vegetation and

development have direct impacts on diversity, multiple indirect effects exist. Our strongest

observed path within the SEM was the direct positive effect of vegetation on diversity while

development showed a weaker but significant negative direct effect. Indeed, the strong negative

covariance between vegetation and development and a strong positive covariance between

vegetation and developed open land, suggests that characteristics of more intense urbanization

may indirectly drive reduced moth diversity by altering land cover compositions. Taken together,

these results indicate that high- and medium-intensity development reduce moth diversity not

only directly, but also indirectly through their association with reduced vegetation cover.

Together, these results suggest that developed land conversion drives species declines

independently and through strong indirect effects. This finding points to potential avenues for

moth conservation through the maintenance of tree cover, vegetation, and open space even in

highly developed parts of the city. Therefore, urban planning efforts focused on restoring parks

and increasing vegetation across the city may make habitat more suitable for local moth

populations.

Surprisingly, we did not find a relationship between air pollution and moth diversity at

our on-the-ground sampling study sites. This finding contradicts our expectations, as researchers

have documented declines across various taxa in response to higher levels of air pollution [59-

60]. However, others suggest that the negative effects of air pollution on insects may have been

overestimated [61]. It is also possible that our findings were influenced by confounding variables

such as temperature and wind speed, as higher levels of particulate matter (PM2.5) coincided

with warmer temperatures and lower wind speeds during sampling. A longer temporal scale may

be required to detect a correlation between air pollution and moth biodiversity, given the high

variation and sensitive nature of this variable. These factors highlight the complexity of isolating

individual environmental drivers of diversity decline and underscore several limitations of our

study.

374   We also acknowledge the inherent limitations of using *iNaturalist* data, which can be

influenced by biases in socioeconomic factors and uneven sampling efforts. Research suggests

that areas with higher socioeconomic status often experience greater sampling efforts by

community scientists [62]. Taxonomic biases also exist; charismatic fauna, such as butterflies,

are reported more frequently despite the greater overall diversity of moth species [63]. For these

reasons we conducted on the ground surveys to reinforce our *iNaturalist* analyses. While our

field data overall reiterated the same findings, it also further highlighted the limitations of citizen

science data. For example, our field sampling documented high diversity in Forest and Highland

Parks, even though these parks have few moth records on *iNaturalist*. In fact, our on-the-ground

sampling documented more moth species over a single evening at three locations in each of these

parks than have ever been documented on *iNaturalist* for the entire park. Forest Park had 29

records on *iNaturalist* in over 20 years of observations in New York City, versus our 35

observations from our on-the-ground surveys; Highland Park is similar with 9 *iNaturalist*

observations versus our 19. Indeed, we detected members of a family (Eriocraniidae) that had

never been observed in New York City on *iNaturalist*, perhaps because of their particularly small

size (~10 mm wingspan) and early spring emergence. However, the *iNaturalist* data recorded 40

families not observed in person. Moreover, *iNaturalist* has many records of larval stage moths,

which would not be recorded during a light trap survey, and other adult stages of certain species

which are difficult to record via light trapping due to variation in phototaxis [41]. The

discrepancies between datasets highlight the importance of integrating *iNaturalist* data with on-

the-ground field surveys to develop a more comprehensive understanding of biodiversity in a

given area, and particularly in cities where habitat is highly heterogeneous and strong sampling

biases can arise.

In conclusion, our analyses revealed significant relationships between moth diversity and

urbanization at both landscape and local spatial scales. Our landscape-level analysis found that

higher vegetation cover and developed open space support greater moth diversity, while our on-

the-ground sampling revealed negative effects of light at night. Further, our findings highlight

the importance of using advanced modeling approaches, such as SEM, alongside linear

regression to reveal hidden or unobserved relationships in environmental data. Understanding

these relationships is essential for designing targeted conservation strategies, as singular actions

such as increasing vegetation alone, may be insufficient. We hope this study sparks further

research into urban moth diversity, particularly given the essential role these organisms serve in

the world's ecosystems.

Acknowledgements

We are thankful to Colin Ainsworth and Cassie Lustig who assisted in field-based data

collection. We would also like to recognize New York University's DURF and Undergraduate

Research Assistant Programs for making this research possible.

References

1. McKinney ML. Urbanization as a major cause of biotic homogenization. Biological Conservation. 2006 Jan 1;127(3):247-60.

2. Piano E, Souffreau C, Merckx T, Baardsen LF, Backeljau T, Bonte D, et al. Urbanization drives cross-taxon declines in abundance and diversity at multiple spatial scales. Global Change Biology. 2020 Mar;26(3):1196-211.

3. Hallmann CA, Sorg M, Jongejans E, Siepel H, Hofland N, Schwan H, et al. More than 75 percent decline over 27 years in total flying insect biomass in protected areas. PloS one. 2017 Oct 18;12(10):e0185809.

4. Vaz S, Manes S, Khattar G, Mendes M, Silveira L, Mendes E, et al. Global meta-analysis of urbanization stressors on insect abundance, richness, and traits. Science of the Total Environment. 2023 Dec 10;903:165967.

5. Bergerot B, Fontaine B, Julliard R, Baguette M. Landscape variables impact the structure and composition of butterfly assemblages along an urbanization gradient. Landscape Ecology. 2011 Jan;26(1):83-94.

6. Fattorini S. Insect extinction by urbanization: a long term study in Rome. Biological Conservation. 2011 Jan 1;144(1):370-5.

7. Giuliano WM, Accamando AK, McAdams EJ. Lepidoptera-habitat relationships in urban parks. Urban Ecosystems. 2004 Dec;7(4):361-70.

8. Lizee MH, Tatoni T, Deschamps-Cottin M. Nested patterns in urban butterfly species assemblages: respective roles of plot management, park layout and landscape features. Urban Ecosystems. 2016 Mar;19(1):205-24.

9. Ramírez-Restrepo L, MacGregor-Fors I. Butterflies in the city: a review of urban diurnal

Lepidoptera. Urban ecosystems. 2017 Feb;20(1):171-82.

10. Fenoglio MS, Calviño A, González E, Salvo A, Videla M. Urbanisation drivers and

underlying mechanisms of terrestrial insect diversity loss in cities. Ecological Entomology.

2021 Aug;46(4):757-71.

11. Sanetra D, Berger J, Hartlieb M, Simons NK, Walther G, Blüthgen N, Staab M.

Disentangling how urbanisation influences moth diversity in grasslands. Insect Conservation

and Diversity. 2024 Mar;17(2):229-42.

12. Frankie GW, Ehler LE. Ecology of insects in urban environments. Annual Review of

Entomology. 1978 Jan;23(1):367-87.

13. Sharmin M, Tjoelker MG, Esperon-Rodriguez M, Katlav A, Gilpin AM, Rymer PD, et al.

Urban greening with shrubs can supercharge invertebrate abundance and diversity. Scientific

Reports. 2024 Apr 16;14(1):8735.

14. Fenoglio MS, González E, Calviño A, Videla M. Urban insect communities in the

Neotropics: a systematic literature review and a green path to promote biodiversity

conservation. Insect Decline and Conservation in the Neotropics. 2024 Apr 22:163-88.

15. Theodorou P, Radzevičiūtė R, Lentendu G, Kahnt B, Husemann M, Bleidorn C, et al. Urban

areas as hotspots for bees and pollination but not a panacea for all insects. Nature

Communications. 2020 Jan 29;11(1):576.

16. Argañaraz CI, Rubio GD, Gleiser RM. Spider communities in urban green patches and their

relation to local and landscape traits. Biodiversity and Conservation. 2018 Mar;27(4):981-

1009.

17. Fox R. The decline of moths in Great Britain: a review of possible causes. Insect

Conservation and Diversity. 2013 Jan;6(1):5-19.

18. Anderson M, Rotheray EL, Mathews F. Marvellous moths! pollen deposition rate of bramble

(*Rubus futicosus L. agg.*) is greater at night than day. Plos one. 2023 Mar 29;18(3):e0281810.

19. Williams N. Raising moth awareness. Current biology. 2007 Jul 17;17(14):R527-8.

20. Belitz MW, Sawyer A, Hendrick LK, Kawahara AY, Guralnick RP. Substantial urbanization-

driven declines of larval and adult moths in a subtropical environment. Global Change

Biology. 2024 Mar;30(3):e17241.

21. Boyes DH, Evans DM, Fox R, Parsons MS, Pocock MJ. Is light pollution driving moth

population declines? A review of causal mechanisms across the life cycle. Insect

Conservation and Diversity. 2021 Mar;14(2):167-87.

22. Liang H, He YD, Theodorou P, Yang CF. The effects of urbanization on pollinators and

pollination: A meta-analysis. Ecology Letters. 2023 Sep;26(9):1629-42.

23. Franzén M, Betzholtz PE, Pettersson LB, Forsman A. Urban moth communities suggest that

life in the city favours thermophilic multi-dimensional generalists. Proceedings of the Royal

Society B. 2020 Jun 10;287(1928):20193014.

24. Merckx T, Kaiser A, Van Dyck H. Increased body size along urbanization gradients at both

community and intraspecific level in macro-moths. Global Change Biology. 2018

Aug;24(8):3837-48.

25. Kettlewell HB. Selection experiments on industrial melanism in the Lepidoptera. Heredity.

1955 Dec;9(3):323-42.

26. Kettlewell HB. Further selection experiments on industrial melanism in the Lepidoptera.

Heredity. 1956 Dec;10(3):287-301.

27. Hagen JB. Retelling experiments: HBD Kettlewell's studies of industrial melanism in

peppered moths. Biology and Philosophy. 1999 Jan;14(1):39-54.

28. Dar AA, Jamal K. Moths as ecological indicators: A review. Munis Entomol. Zool. J. 2021

Jun 29;16:830-6.

29. New TR. Moth ecology and conservation importance. In The Other Lepidoptera: Moth

Conservation in Australia 2023 Jul 11 (pp. 21-37). Cham: Springer International Publishing.

30. Choi SW. Diversity and composition of larger moths in three different forest types of

Southern Korea. Ecological Research. 2008 May;23(3):503-9.

31. Gaona FP, Iñiguez-Armijos C, Brehm G, Fiedler K, Espinosa CI. Drastic loss of insects

(Lepidoptera: Geometridae) in urban landscapes in a tropical biodiversity hotspot. Journal of

Insect Conservation. 2021 Jun;25(3):395-405.

32. Carter DJ. Butterflies and moths. DK Publishing (Dorling Kindersley); 1992.

33. Kristensen NP, Scoble MJ, Karsholt OL. Lepidoptera phylogeny and systematics: the state of

inventorying moth and butterfly diversity. Zootaxa. 2007 Dec 21;1668(1):699-747.

34. Posit Team. RStudio: Integrated Development Environment for R (ver. 2023.12.0.369). Posit

Software, PBC, Boston, MA; 2023. Available from: http://www.posit.co/

35. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing, Vienna, Austria; 2023. Available from: https://www.r-project.org/

36. Fox J, Weisberg S. An R Companion to Applied Regression, Third Ed. John-fox.ca. 2019.

Available from: https://www.john-fox.ca/Companion/

37. QGIS Development Team. QGIS Geographic Information System (ver. 3.34.3-Prizren).

QGIS Association; Available from: https://www.qgis.org/

38. GIS DCP. NYC Borough Boundary [shapefile] [Internet]. 2025. Available from:

https://hub.arcgis.com/datasets/DCP::nyc-borough-boundary/about

39. Hamilton C. QGIS Density Analysis Plugin (Version 2024.1.23) [Internet]. GitHub; 2024.

Available from: https://github.com/hamiltoncj/qgis-densityanalysis-plugin

40. Dewitz J. National Land Cover Database (NLCD) 2021 Products: U.S. Geological Survey

data release [Internet]. 2023. Available from: https://doi.org/10.5066/P9JZ7AO3

41. Merckx T, Slade EM. Macro-moth families differ in their attraction to light: implications for

light-trap monitoring programmes. Insect Conservation and Diversity. 2014 Sep;7(5):453-61.

42. Wang K, Qi Y, Hu Y, Han D. Distribution and Influences on Butterfly Diversity in Urban

Park Green Spaces: A Case Study of Harbin, China. Ecology and Evolution. 2025

Jun;15(6):e71554.

43. Kassambara A, Mundt F. factoextra: Extract and Visualize the Results of Multivariate Data

Analyses (ver. 1.0.7) [Internet]. 2020. Available from: https://cran.r

project.org/package=factoextra

44. Wei T, Simko V. R package "corrplot": Visualization of a Correlation Matrix (ver. 0.92)

[Internet]. 2024. Available from: https://github.com/taiyun/corrplot

45. Simpson EH. Measurement of diversity. nature. 1949 Apr 30;163(4148):688-688.

46. Qiao H, Orr MC, Hughes AC. Measuring metrics: what diversity indicators are most

appropriate for different forms of data bias?. Ecography. 2024 Sep;2024(9):e07042.

47. Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, et al. vegan:

Community Ecology Package (ver. 2.6-4) [Internet]. 2022. Available from: https://cran.r-

project.org/package=vegan

48. Goslee SC, Urban DL. The ecodist package for dissimilarity-based analysis of ecological

data. Journal of Statistical Software. 2007 Sep 30;22:1-9.

49. Rosseel Y. lavaan: An R package for structural equation modeling. Journal of statistical

software. 2012 May 24;48(1):1-36.

50. Epskamp S. semPlot: Path Diagrams and Visual Analysis of Various SEM Packages' Output

(ver. 1.1.6) [Internet]. 2022. Available from: https://cran.r-project.org/package=semPlot

51. Beck JA, Schulze CH, Linsenmair KE, Fiedler K. From forest to farmland: diversity of

geometrid moths along two habitat gradients on Borneo. Journal of tropical ecology. 2002

Jan;18(1):33-51.

52. Usher MB, Keiller SW. The macrolepidoptera of farm woodlands: determinants of diversity

and community structure. Biodiversity & Conservation. 1998 Jun;7(6):725-48.

53. Lintott PR, Bunnefeld N, Fuentes-Montemayor E, Minderman J, Blackmore LM, Goulson D,

Park KJ. Moth species richness, abundance and diversity in fragmented urban woodlands:

implications for conservation and management strategies. Biodiversity and Conservation.

2014 Oct;23(11):2875-901.

54. White PJ. An Aerial Approach to Investigating the Relationship between Macromoths and

Artificial Nighttime Lights Across an Urban Landscape1. Journal of Agricultural and Urban

Entomology. 2018 Jan;34(1):1-4.

55. van Langevelde F, Braamburg-Annegarn M, Huigens ME, Groendijk R, Poitevin O, van

Deijk JR, Ellis WN, van Grunsven RH, de Vos R, Vos RA, Franzén M. Declines in moth

populations stress the need for conserving dark nights. Global Change Biology. 2018

Mar;24(3):925-32.

56. Wilson JF, Baker D, Cheney J, Cook M, Ellis M, Freestone R, Gardner D, Geen G,

Hemming R, Hodgers D, Howarth S. A role for artificial night-time lighting in long-term

changes in populations of 100 widespread macro-moths in UK and Ireland: a citizen-science

study. Journal of Insect Conservation. 2018 Apr;22(2):189-96.

57. Grenis K, Nufio C, Wimp GM, Murphy SM. Does artificial light at night alter moth

community composition?. Philosophical Transactions of the Royal Society B. 2023 Dec

18;378(1892):20220365.

58. Rice AJ. Community patterns in urban moth assemblages. The Journal of the Lepidopterists'

Society. 2015 May;69(3):149-56.

59. Lovett GM, Tear TH, Evers DC, Findlay SE, Cosby BJ, Dunscomb JK, Driscoll CT,

Weathers KC. Effects of air pollution on ecosystems and biological diversity in the eastern

United States. Annals of the New York Academy of Sciences. 2009 Apr;1162(1):99-135.

60. Barker JR, Tingey DT, editors. Air pollution effects on biodiversity. Springer Science &

Business Media; 2012 Dec 6.

61. Kozlov MV. Population dynamics of herbivorous insects in polluted landscapes. Current

Opinion in Insect Science. 2022 Dec 1;54:100987.

62. Carlen EJ, Estien CO, Caspi T, Perkins D, Goldstein BR, Kreling SE, et al. A framework for

contextualizing social-ecological biases in contributory science data. People and Nature.

2024 Apr;6(2):377-90.

63. Shirey V, Belitz MW, Barve V, Guralnick R. A complete inventory of North American

butterfly occurrence data: narrowing data gaps, but increasing bias. Ecography. 2021

Apr;44(4):537-47.

A

Land Cover
- Deciduous
- Developed, High Intensity
- Developed, Med. Intensity
- Developed, Low Intensity
- Developed, Open Space
- Grassland
- Pasture
- Shrub
- Water

Proportion of Land Cover

Prospect Park (c)
Prospect Park (b)
Prospect Park (a)
Maria Hernandez
Irving Square
Highland Park (c)
Highland Park (b)
Highland Park (a)
Grover Cleveland
Forest Park (c)
Forest Park (b)
Forest Park (a)

B

Landcover PC1

On-the-ground   iNaturalist

- Forest Park (a)
- Forest Park (b)
- Forest Park (c)
- Grover Cleveland
- Highland Park (a)
- Highland Park (b)
- Highland Park (c)
- Irving Square
- Maria Hernandez
- Prospect Park (a)
- Prospect Park (b)
- Prospect Park (c)

C

Propsect Park, Brooklyn   Forest Park, Queens   Maria Hernandez, Brooklyn

200 m   200 m   200 m