From Concrete to Canopy: Illuminating Moth Biodiversity in New York City's Urban Jungle

Shira Linsk[1], Anna Thonis[1], & Kristin M. Winchell[1,2]

[1]New York University, Department of Biology
[2]Correspondence: kristin.winchell@nyu.edu

Author contact information: sl8825@nyu.edu, at6689@nyu.edu, kristin.winchell@nyu.edu

Data Availability: All data and associated code are archived on Zenodo

with doi: 10.5281/zenodo.14756963

Funding statement: This work was supported College of Arts and Science,

New York University, Dean's Undergraduate Research Fund

Conflict of interest disclosure: The authors declare no conflicts of

interest.

Words: 3,989

tables, 5 figures

Abstract

Moths (Lepidoptera) are sensitive to anthropogenic threats and serve

as valuable bioindicators. Despite the remarkable diversity and abundance

of Lepidoptera globally, there is a lack of information on how moth species

are impacted by urbanization. Notably, very little is known about moths in

the largest city of the United States, New York City, where pervasive urban

pollutants, artificial light at night, land cover change, and habitat

fragmentation are severe. We examined the effects of urbanization on moth

biodiversity in New York City, with a focus on green spaces. We sampled

twelve locations across six parks at night and complemented these surveys

with citizen science records of moths from *iNaturalist*. Overall, we found a

negative relationship between greater urbanization and moth biodiversity,

with community similarity correlated with environmental similarity. Our

results support greater biodiversity with less light at night, urban

development, and more deciduous tree cover and open land. Our structural

equation model reveals additional insight: although we detected a strong

negative direct effect of developed land, urbanization also impacts diversity

via indirect effects of reducing open space and deciduous tree cover.

Developed open space alone does not directly affect diversity, but may

positively impact diversity through its covariance with vegetation cover.

These findings support the importance of mitigating artificial light at night

in urban green spaces and maintaining urban vegetation to ensure

nocturnal Lepidoptera are able to persist in rapidly urbanizing landscapes.

Keywords: Lepidoptera, moths, urbanization, ALAN, urban greenspace,

invertebrates, urban ecology, biodiversity, path analysis

Introduction

Urbanization is linked to declines in biodiversity through biotic

homogenization and species extirpation (McKinney *et al.*, 2006; Piano *et al.*,

2020). Although these impacts are seen across the tree of life, the effects of

urbanization on invertebrates have been particularly drastic, with steep

declines across insect orders (Hallman *et al.,* 2017; Vaz *et al.*, 2023).

Studies have demonstrated local extinctions of insect populations and

reduced insect diversity with increasing urbanization (Bergerot *et al.*, 2010;

Fattorini, 2011). Much of this work has focused on bees and butterflies,

revealing decreased diversity and abundance with increased urbanization

(Giuliano, 2004; Bergerot *et al.*, 2010; Lizee *et al.*, 2015; Ramírez-Restrepo

& MacGregor-Fors, 2016).

Urbanization presents many challenges for insects, including the

urban heat island effect, habitat fragmentation, pollution, impervious

surfaces, and exotic plants (Fenoglio *et al.*, 2021). Habitat fragmentation

and landscape maintenance have been major contributors to pollinator

decline, including Lepidoptera (Sanetra *et al.*, 2024). However, alongside

these challenges come opportunities in terms of anthropogenic resources

that may be exploited, as seen in common urban species like cockroaches,

flies, and carpet beetles (Frankie & Ehler, 1978). Many species may thrive

in cities despite the threats of pest control and habitat modification (Frankie

& Ehler, 1978). For example, urban plantings may generate habitat and

resources that support invertebrate diversity (Sharmin *et al.*, 2024;

Fenoglio *et al.*, 2024), including Hymenoptera (Theodorou *et al.*, 2020) and

spiders (Argañaraz *et al.*, 2017).

Relatively little research on urban invertebrates has focused on

moths, and there is a notable bias toward diurnal Lepidoptera (Fox, 2012;

Anderson *et al.*, 2023; Giuliano, 2004). A few studies on moths have found

decreasing populations with increased artificial light at night (Belitz *et al.*,

2024; Boyes *et al.*, 2020; Liang *et al.,* 2023). Additionally, anthropogenic

environments generate intense selection pressures and filtering processes

that may favor the survival of only a few species, such as heat-tolerant

moths (Franzen *et al.*, 2020), or produce phenotypic shifts, such as

increased body size in urban moths (Merckx *et al.*, 2018). Indeed, studies of

industrial melanism in moths represent some of the first examples of rapid

evolutionary change in cities (Kettlewell, 1955; Kettlewell, 1956; Hagen,

1999).

Understanding the effects of urbanization on moths is important

because of the crucial roles they play in ecosystem services (e.g.,

pollination, food, and nutrient cycling). Moths are fundamental to ecosystem

health and stability and serve as bioindicators (Dar & Jamal, 2021; Liang *et*

al., 2023; New, 2023) since they respond quickly to environmental change

(Belitz *et al.*, 2024; Choi, 2008; Gaona *et al.*, 2021). There are approximately

170,000 known species of moths globally (Carter, 1992), and estimates

range as high as 500,000 (Kristensen *et al.,* 2007). Yet, we know very little

about the species that persist in urban landscapes. For conservation

initiatives to be successful in these spaces, we must first take stock of the

species that reside there and relate patterns of diversity to urban stressors.

We investigate patterns of moth diversity in New York City, where

artificial light at night, pollution, and habitat fragmentation are extreme.

We hypothesize that habitat-scale features such as pollution, light at night,

and microclimate correlate with moth diversity. In addition, we hypothesize

that landscape-scale features influence species diversity and predict

developed land use negatively impacts moth biodiversity. By connecting

moth biodiversity to specific urban features, our findings contribute to our

understanding of the ecological effects of urbanization and can aid the

development of urban conservation strategies.

Methods

Statistical analyses were conducted using R studio (Posit, 2023, ver.

2023.12.0.369) and R (v.4.3.2; R Core Team, 2023). For all linear models,

we evaluated multicollinearity with the 'VIF' function in the package *car*

(Fox & Weisberg, 2019), assessed statistical significance using t-tests for

individual coefficients, and evaluated model fit based on Akaike Information

Criterion (AIC) and residual deviance.

Site Selection: We identified 12 on-the-ground sampling sites in six

green spaces in Brooklyn and Queens, New York, that reflect variation in

urbanization (Fig. 1). We visually assessed satellite imagery to identify

green spaces, then chose general locations within each based on their

position along a continuum of urbanization quantified from remote-sensed

landcover (see below). We chose specific sampling sites spanning a gradient

of light pollution based on on-the-ground measurement of light (rather than

remote-sensed) to capture variation in this variable at an organismally

relevant spatial scale.

Field Sampling: We surveyed each location once shortly after dusk (8-

10 pm) between March 3-April 29, 2024. We restricted the temporal

sampling period to minimize seasonal variation in species detected across

sites. We constructed a light trap using a suspended 5 m^2 white sheet

directly illuminated with two flood lights (6000 Kelvin, visible light

spectrum) and one ultraviolet LED light (385-395 nm) (as in Merckx &

Slade, 2014). Moths are easily sampled with light traps, allowing for robust

estimates of species richness and abundance (Choi, 2008). Each survey

consisted of two people photographing all moths observed landing on or in

the immediate vicinity (on illuminated ground within 2m) of the light trap.

We identified species from the photographs and counted the total number of

individuals per species encountered during each sampling event. In

addition, for each sampling event, we measured the following site-level

variables: air pollution (PM 2.5; Temtop M2000C), light at night (lux;

EXTECH Light Meter LT300), temperature, humidity, and wind speed

(Kestrel 3000).

iNaturalist data: We downloaded 31,793 Research-Grade observations

of the Order Lepidoptera using a polygon spatial filter around all five

boroughs of New York City. We manually removed butterfly Families

(Papilionidae, Nymphalidae, Pieridae, Lycaenidae, Hesperiidae), resulting in

12,956 observations of moths. We imported records into QGIS (QGIS, 2024,

v.3.34.3-Prizren) and used the New York City administrative boundary layer

(GIS, 2025) to reduce this dataset to 10,740 records. We used the plugin

Density Analysis (Hamilton, 2024) to visualize observation density using a

grid of 1 km hexagonal cells across the region (Fig. 1). We excluded

hexagonal zones with undefined landscape types or >70% water cover to

help ensure the zones have relevant habitat data for terrestrial species,

resulting in 9,394 records. Three additional observations were excluded

because they were from our on-the-ground sampling, resulting in a final

dataset of 9,391 records.

Landscape analysis: We mapped on-the-ground sampling locations

and *iNaturalist* records using QGIS. We extracted zonal statistics of land

cover classification (National Land Cover Database "NLCD"; Dewitz, 2023)

within a 150-meter buffer around each on-the-ground sampling location and

within each hexagonal zone, and calculated the percentage of each land

cover class within each. We summarized land cover classifications

(Deciduous, Mixed Forest, Grassland, Pasture/Hay, Developed Open Space,

Developed Low-Intensity, Developed Medium-Intensity, Developed High-

Intensity) with principal component analysis (PCA) using the function

'prcomp' in base R *stats* (R Core Team, 2023). We visualized results with

packages *factoextra (*Kassambara & Mundt, 2020) and *corrplot* (Wei &

Simko, 2021).

Statistical analyses: We summarized species diversity with Simpson's

Diversity Index (Simpson's D; Simpson, 1949), which considers both species

richness and evenness. We calculated Simpson's D (1-D) using species-level

identification for each on-the-ground location and each *iNaturalist* hex bin

in R using the function 'diversity' in the package *vegan* (Oksanen *et al.*,

2022).

For on-the-ground sampling, we conducted two analyses. We first

analyzed the similarity of community composition with respect to

environmental similarity across sites using multiple matrix regression. This

entailed generating a distance matrix of species composition similarity

across sites, as well as for the environment (percentages of all NLCD land

cover classes, PM2.5, lux, temperature, humidity, and wind speed). We used

the function 'dist' in the R base *stats* for all distance matrix analyses (R

Core Team, 2023) and performed matrix regression using the function

'MRM' in R package *ecodist* (Goslee & Urban, 2007). Second, we analyzed

diversity using a linear regression of Simpson's Diversity and site-level

habitat features. We did not include remote-sensed land cover in this model

because the resolution of these data and our sample size resulted in low

statistical power (overparameterization) and multicollinearity when these

variables were included. Explanatory variables included temperature,

pollution (PM 2.5), and light intensity (lux) measured during on-the-ground

sampling. We simplified this model using stepwise model simplification

(backwards and forwards) based on AIC using the function 'step' in R base

stats.

For the *iNaturalist* data, we conducted three analyses. First, we

analyzed the similarity of community composition with respect to

environmental similarity across sites using multiple matrix regression. As

with the on-the-ground MRM analysis, this entailed generating a distance

matrix of species composition similarity across hex-bins, and for the

environment (percentages of all NLCD land cover classes only). Second, we

analyzed diversity using a linear regression of Simpson's Diversity and

landscape-level (remote-sensed) habitat features. Explanatory variables in

the model included NLCD categories (as a proportion of hex-bin area):

deciduous tree cover, developed land cover (high-, medium-, low-intensity),

pasture, developed open land, shrub, and grassland. We simplified this

model using stepwise model simplification (backwards and forwards) based

on AIC using the function 'step' in R base *stats*.

Third, we developed a structural equation model (SEM) using

iNaturalist observations to examine relationships between species diversity,

vegetation (deciduous forest, grassland, pasture, shrub), developed land

(high-, medium-, low-intensity), and developed open land cover. SEMs

analyze multiple relationships at the same time while considering latent

factors (i.e., unobserved relationships, such as between deciduous land

cover and grassland). We fit our SEM using maximum likelihood with the

'sem' function in the *lavaan* R package (v.0.6-19, Rosseel 2012). Model fit

was assessed using Chi-squared (χ^2), Comparative Fit Index (CFI), Tucker-

Lewis Index (TLI), and Root Mean Square Error of Approximation (RMSEA).

To account for shared variance between variables, we specified covariances

between vegetation cover and development intensity, development intensity

and developed open space, and between several vegetation variables,

including between deciduous and grassland, and between grassland and

shrub. We used the z-statistic to assess the significance of each path

coefficient in the SEM and visualized observed relationships with the

function 'semCors' in the R package *semPlot* (Epskamp, 2022*)*.

In addition, we evaluated relationships between diversity and

urbanization more generally across all on-the-ground sample sites combined

with *iNaturalist* data using a linear model of Simpson's Diversity Index and

the first four principal components of our PCA. We retained four principal

components in this analysis based on the percentage of variance captured

and visual inspection of the scree plot.

Results

Urban environmental variation: Land cover analyses using remote-

sensed data verified that our on-the-ground sampling locations spanned a

broad range of urbanization (Fig. 2a). Larger parks (Forest Park, Highland

Park, Prospect Park) exhibited greater habitat heterogeneity and had more

deciduous forest cover. Small parks (Maria Hernandez, Grover Cleveland,

Irving Square), by contrast, had little to no deciduous tree cover and were

composed almost entirely of developed land. Grover Cleveland Park stood

out with only a small percentage of high-intensity developed land as it

borders a private cemetery with green space. These differences are clearly

visible in satellite imagery: Forest Park is dominated by forest, Prospect

Park has less tree cover, and Maria Hernandez is substantially smaller and

consists mostly of developed land (Fig. 2c).

Our PCA incorporated land cover for both on-the-ground sites and

iNaturalist records (12 on-the-ground locations, 1115 *iNaturalist* hexagonal

zones). The PCA captured 95.22% of the variance in the first four principal

components, with PC1 capturing 44.65% of the variance. Visual inspection

of the scree plot suggests that 4 principal components should be retained to

describe variation in the dataset. The first and second principal components

(PC1 and PC2) represent urbanization, with strong positive loadings for

developed high-intensity land cover on PC1 (Fig. 2b) and for medium-

intensity developed land cover on PC2. The third component (PC3) was

negatively correlated with water and positively correlated with developed

open space and deciduous tree cover. The fourth principal component (PC4)

was positively correlated with developed open space and developed low-

intensity land cover and negatively associated with deciduous tree cover.

Grassland, pasture, and shrub land cover classifications did not load

strongly on any of the first four principal components.

Diversity of Lepidoptera: In our on-the-ground surveys, we observed a

total of 69 individuals, representing 11 families, 31 genera, and 33 species

(Fig. 3a). Observations were identified to species in 100% of samples. From

iNaturalist, we extracted data from 9391 "Research Grade" observations,

representing 955 species, 537 genera, and 50 families (Fig. 3b). The

number of species documented on *iNaturalist* from the locations where we

sampled on the ground were much higher for Prospect Park (110 species on

iNaturalist versus 12 in our sampling) and similar for Grover Cleveland (2

on *iNaturalist* versus 3). By contrast, in two parks, we detected more

species in our on-the-ground sampling: Forest Park (29 on *iNaturalist*

versus 35) and Highland Park (9 on *iNaturalist* versus 19). In the two parks

where no species were detected in our on-the-ground sampling (Maria

Hernandez and Irving Square), there were also no observations in

iNaturalist. Notably, we detected individuals from the family Eriocraniidae

in our on-the-ground sampling, but this family was not present in the

iNaturalist dataset. Conversely, *iNaturalist* observations documented 40

additional families not detected in our on-the-ground sampling.

Based on our on-the-ground sampling, large parks (Forest Park,

Highland Park, Prospect Park) displayed the highest levels of biodiversity

(1-D: 0.44–0.92, mean±sd: 0.72±0.15). Among them, Forest Park had a

mean diversity of 0.80±0.1, Highland Park had a mean diversity of

0.71±0.19, and Prospect Park had a mean diversity of 0.64±0.17. In

contrast, the small parks (Maria Hernandez, Grover Cleveland, Irving

Square) exhibited lower biodiversity (1-D: 0–0.66, mean±sd: 0.22±0.38). No

species were detected at Maria Hernandez and Irving Square in our on-the-

ground sampling; since Simpson's Diversity Index is undefined in the

absence of observations, we recorded 1-D as 0 for these sites.

Community Composition: In both our on-the-ground sampling and the

iNaturalist data, sites with similar environments had similar community

composition at the species level (multiple-matrix regression; on-the-ground:

F=43.127, R^2=0.501, p=0.031; *iNaturalist*: F=3943.21, R^2=0.014,

p<0.001). Species diversity is strongly negatively correlated with PC1 and

PC2 (representing increased high- and medium-intensity development,

respectively) across both datasets combined (PC1 estimate = -0.198±0.041,

t=-4.778, p<0.001; PC2 estimate=

-0.446±0.052, t=-8.637, p<0.001) (Fig. 4). Diversity was also positively

correlated with PC3, representing less water and more developed open

space and deciduous tree cover (estimate= 0.415±0.073, t=5.653, p<0.00).

Diversity was not significantly correlated with PC4 (t=0.095, p=0.924).

Environmental Correlations: Our two linear models—one using on-the-

ground sampling with field-measured environmental variation and the other

using *iNaturalist* sampling with remote-sensed environmental variation—

revealed correlational relationships between environmental features and

biodiversity across both spatial scales. Our simplified regression of

Simpson's Diversity for on-the-ground sampling retained two variables: air

pollution and light at night. Simpson's diversity was not significantly

correlated with air pollution (t=1.280, p=0.233) but was negatively

correlated with light at night (estimate=-0.512±0.062 lux, t=-8.214,

p<0.001). In comparison, our simplified linear regression of Simpson's

Diversity by land cover using *iNaturalist* and remote-sensed data retained

five significant variables: medium-intensity developed land, low-intensity

developed land, developed open land, shrubland, and deciduous tree cover.

Biodiversity was negatively correlated with medium-intensity developed

land cover (estimate=-0.348±0.064, t=-5.465, p<0.001) and positively

correlated with deciduous tree cover (estimate=0.468±0.107, t=4.364,

p<0.001) and developed open space (estimate=0.374±0.119, t=3.159,

p=0.002). There was also a near-significant positive trend between diversity

and low-intensity developed space (estimate=0.280±0.153, t=1.832,

p=0.068), and the correlation with shrubland was non-significant (t=1.457,

p=0.145).

Structural Equation Model: Our structural equation model (SEM)

exhibited an overall good fit to the data (CFI: 0.955, TLI: 0.915, RMSEA:

0.067). The chi-square statistic was significant (χ^2=83.670, df=19,

p<0.001), although this is expected with larger sample sizes, and the

overall fit indices suggest the model effectively captures the relationships in

the data. Our SEM detected a significant positive effect of vegetation cover

on diversity (estimate=1.856±0.394, z=4.715, p<0.001; Fig. 5) and a

significant negative effect of developed land on diversity (estimate=-

0.301±0.100, z=-3.011, p=0.003). We did not find a significant direct effect

of developed open space (z=0.435, p=0.664) on diversity (Fig. 5).

Significant negative covariance was observed between vegetation cover and

development intensity (covariance=-0.012, z=-10.725, p<0.001), and

significant positive covariance between vegetation cover and open space

(covariance=0.002, z=4.15, p<0.001). Similarly, we found a significant

negative covariance between development intensity and developed open

space

(covariance=-0.014, z=-11.702, p<0.001). We also observed weak but

significant positive covariance between developed low-intensity and

developed open space (covariance=0.005, z=9.396, p<0.001) and grassland

and shrub (covariance<0.001, z=11.033, p<0.001). We observed weak but

significant negative covariance between developed high-intensity and

developed low-intensity (covariance=0.015, z=-11.270, p<0.001), and

deciduous and grassland (covariance<-0.001, z=-2.655, p = 0.008). Further

inspection of relationships among variables indicates that developed

medium-intensity land cover is driving the direct effect of development on

diversity, whereas high and low-intensity development impact diversity

more indirectly via their relationships with open land and deciduous tree

cover. Further, among the vegetation variables, only deciduous tree cover

has a strong direct effect on diversity, whereas grassland, shrubland, and

pasture all have indirect effects related to their covariance with developed

land.

Discussion

Urbanization drastically alters landscapes, replacing naturally

occurring habitats and vegetation with impervious surfaces and buildings.

This process not only impacts land cover but also increases artificial light at

night (ALAN), which can both have significant consequences for

biodiversity. Lepidoptera (moths and butterflies) are highly sensitive to land

cover changes (Gaona et al., 2021; Beck et al., 2002), and are also

influenced by habitat patch size, quality, and shape (Usher & Keiller, 1998;

Lintott et al., 2014). In this study, we examined how urbanization relates to

biodiversity of nocturnal Lepidoptera (moths), which perform vital

ecosystem services and act as bioindicators (Dar & Jamal, 2021). Our

findings suggest that urbanization is associated with lower moth diversity,

with specific elements of the urban landscape shaping patterns of

biodiversity in complex ways.

We found comparable moth community composition between locations

with similar environments in both our on-the-ground and *iNaturalist*

datasets, and a general trend of decreasing diversity with increasing

urbanization. This result suggests that moth community composition across

New York City is strongly influenced by the environmental features that

define different locations. Our linear models provide some insight into those

features: we found strong negative correlations between diversity medium-

intensity urbanization and strong positive correlations between diversity

and deciduous tree cover and open land. In addition, our on-the-ground

sampling found a strong negative relationship between moth diversity and

light at night, with a high explained variance (R^2), suggesting light at night

accounts for a substantial portion of the observed variation in species

diversity across sampled locations. Although several studies have found no

consistent effects of light at night on biodiversity (White, 2018; Sanetra *et*

al., 2024), others have identified significant impacts on moth community

composition (Langevelde *et al.*, 2017; Wilson *et al.*, 2018; Grenis *et al.*,

2023).

Our PCA analysis across both on-the-ground and *iNaturalist* data

reinforces these findings: we found strong negative correlations between

diversity and principal components primarily defined by high- and medium-

intensity development, and strong positive correlations with the principal

component representing deciduous tree cover and open space. The

identification of deciduous tree cover, open space, and high- and medium-

intensity developed land across multiple analyses highlights the importance

of these features. Our results are consistent with those of similar studies

performed in metropolitan areas where highly disturbed habitats have been

shown to correlate with reduced moth species richness (Franzen *et al.*,

2020; Rice, 2015; Belitz *et al.,* 2024).

We used structural equation modeling (SEM) to gain deeper insight

into the correlations and interactions among environmental variables

shaping biodiversity. SEMs more holistically represent the complex effects

of urbanization and can be insightful for informing conservation (Yang *et*

al., 2022; Wang *et al.*, 2019; Santibañez-Andrade *et al.*, 2015). While we did

not perform a SEM using our on-the-ground data due to the larger sample

sizes required, our SEM using *iNaturalist* data and remote-sensed land

cover found, similar to our other analyses, that higher diversity is

associated with greater vegetation cover and less developed land. Further

inspection of these relationships revealed that although vegetation and

development have direct impacts on diversity, multiple indirect effects exist.

We found a strong negative covariance between vegetation and

development and a strong positive covariance between vegetation and

developed open land, suggesting that characteristics of more intense

urbanization may drive reduced moth diversity. The many indirect effects

detected in our SEM provide additional insight: both high- and medium-

intensity development negatively impact moth diversity through their

negative impact on tree cover and open space. Together, these results

suggest that developed land conversion itself does not necessarily drive

species declines but may lead to declines through strong indirect effects.

This finding points to potential avenues for moth conservation through the

maintenance of tree cover, vegetation, and open space in more developed

parts of the city.

Surprisingly, we did not find a relationship between air pollution and

moth diversity at our study sites. This finding contradicts our expectations,

as researchers have documented declines across various taxa in response to

higher levels of air pollution (Lovett *et al.,* 2009; Barker & Tingey, 2012).

However, Zvereva and Kozlov (2009) suggest that the negative effects of air

pollution on insects may have been overestimated. It is also possible that

our findings were influenced by confounding variables such as temperature

and wind speed, as higher levels of particulate matter (PM2.5) coincided

with warmer temperatures and lower wind speeds during sampling. A

longer temporal scale may be required to detect a correlation between air

pollution and moth biodiversity, given the high variation and covarying

nature of this variable. These factors highlight the complexity of isolating

individual environmental drivers and underscore some limitations of our

study.

Incorporating *iNaturalist* observations enabled us to expand beyond

the spatial and temporal scope of our on-the-ground sampling. This resulted

in detecting an additional 40 families not observed in person, most of which

were not active during the sampling timeframe. Moreover, *iNaturalist* has

many records of larval stage moths, which would not be recorded during a

light trap survey, and adult stages of some species are difficult to record via

light trapping due to variation in phototaxis (Merckx & Slade, 2014). We

also acknowledge the inherent limitations of using *iNaturalist* data, which

can be influenced by biases in socioeconomic factors and uneven sampling

efforts. Research suggests that areas with higher socioeconomic status

often experience greater sampling efforts by community scientists (Carlen

et al. 2024). Taxonomic biases also exist; charismatic fauna, such as

butterflies, are reported more frequently despite the greater overall

diversity of moth species (Shirey *et al.*, 2021). Although our sampling

documented high diversity in Forest and Highland Park, these parks have

few moth records on *iNaturalist*. In fact, our on-the-ground sampling

documented more moth species over a single evening at three locations in

each of these parks than have ever been documented on *iNaturalist* for the

entire park. Forest Park had 29 records on *iNaturalist* in over 20 years of

observations in New York City, versus our 35 observations from our on-the-

ground surveys; Highland Park is similar with 9 *iNaturalist* observations

versus our 19. Indeed, we detected members of a family (Eriocraniidae) that

had never been observed in New York City on *iNaturalist*, perhaps because

of their particularly small size (~10mm wingspan) and early spring

emergence. This discrepancy between datasets highlights the importance of

integrating *iNaturalist* data with on-the-ground field surveys to develop a

more comprehensive understanding of biodiversity in a given area, and

particularly in cities where habitat is highly heterogeneous and strong

sampling biases can arise.

In conclusion, our analyses revealed significant relationships between

moth biodiversity and urbanization at both local and landscape spatial

scales. Our on-the-ground sampling revealed negative effects of light at

night, while our landscape-level analyses found higher vegetation cover and

developed open space support greater moth diversity. Further, our findings

highlight the importance of using advanced modeling approaches, such as

SEM, alongside linear regression to reveal hidden or unobserved

relationships in environmental data. Understanding these relationships is

essential for designing targeted conservation strategies, as singular

actions—such as increasing vegetation alone—may be insufficient. We hope

this study sparks further research into urban moth biodiversity, as these

remarkable and essential creatures serve vital functions in our ecosystems.

References

Anderson, M., Rotheray, E. L., & Mathews, F. (2023). Marvellous moths!

pollen deposition rate of bramble (Rubus futicosus L. agg.) is greater

at night than day. *PLOS ONE*, *18*(3), e0281810.

https://doi.org/10.1371/journal.pone.0281810

Argañaraz, C. I., Rubio, G. D., & Gleiser, R. M. (2017). Spider communities

in urban green patches and their relation to local and landscape

traits. *Biodiversity and Conservation*, *27*(4), 981–1009.

https://doi.org/10.1007/s10531-017-1476-8

Barker, J. R., & Tingey, D. T. (2012). *Air Pollution Effects on Biodiversity*.

Springer Science & Business Media.

Beck, J., Schulze, C. H., Linsenmair, K. E., & Fielder, K. (2002). From forest

to farmland: diversity of geometrid moths along two habitat gradients

on Borneo. *Journal of Tropical Ecology*, *18*(1), 33–51.

https://doi.org/10.1017/s026646740200202x

Belitz, M. W., Sawyer, A., Hendrick, L. K., Kawahara, A. Y., & Guralnick, R.

P. (2024). Substantial urbanization-driven declines of larval and adult

moths in a subtropical environment. *Global Change Biology*, *30*(3).

https://doi.org/10.1111/gcb.17241

Bergerot, B., Fontaine, B., Julliard, R., & Baguette, M. (2010). Landscape

variables impact the structure and composition of butterfly

assemblages along an urbanization gradient. *Landscape Ecology*,

26(1), 83–94. https://doi.org/10.1007/s10980-010-9537-3

Boyes, D. H., Evans, D. M., Fox, R., Parsons, M. S., & Pocock, M. J. O.

(2020). Is light pollution driving moth population declines? A review of

causal mechanisms across the life cycle. *Insect Conservation and*

Diversity, *14*(2). https://doi.org/10.1111/icad.12447

Carlen, E. J., Estien, C. O., Caspi, T., Perkins, D., Goldstein, B. R., Kreling,

S., Hentati, Y., Williams, T. D., Stanton, L. A., Des Roches, S., Johnson,

R. F., Young, A. N., Cooper, C. B., & Schell, C. J. (2024). A framework

for contextualizing social-ecological biases in contributory science

data. *People and Nature*, *6*(2). https://doi.org/10.1002/pan3.10592

Carter, D. J. (1992). *Butterflies and Moths*. Dorling Kindersley.

Choi, S.-W. (2007). Diversity and composition of larger moths in three

different forest types of Southern Korea. *Ecological Research*, *23*(3),

503–509. https://doi.org/10.1007/s11284-007-0406-8

Dar, A. A., & Jamal, K. (2021). Moths As Ecological Indicators: A REVIEW.

Munis Entomology & Zoology.

https://www.munisentzool.org/yayin/Vol_16/Issue_2/20210520-

BX9WJO4X.pdf

Dewitz, J., 2023, National Land Cover Database (NLCD) 2021 Products: U.S.

Geological Survey data release, https://doi.org/10.5066/P9JZ7AO3.

Epskamp S (2022). _semPlot: Path Diagrams and Visual Analysis of Various

SEM Packages' Output_. R package version 1.1.6, https://CRAN.R-

project.org/package=semPlot

Fattorini, S. (2011). Insect extinction by urbanization: A long term study in

Rome. *Biological Conservation*, *144*(1), 370–375.

https://doi.org/10.1016/j.biocon.2010.09.014

Fenoglio, M. S., Calviño, A., González, E., Salvo, A., & Videla, M. (2021).

Urbanisation drivers and underlying mechanisms of terrestrial insect

diversity loss in cities. *Ecological Entomology*, *46*(4), 757–771.

https://doi.org/10.1111/een.13041

Fenoglio, M. S., González, E., Calviño, A., & Videla, M. (2024). Urban Insect

Communities in the Neotropics: A Systematic Literature Review and a

Green Path to Promote Biodiversity Conservation. *Springer EBooks*,

163–188. https://doi.org/10.1007/978-3-031-49255-6_8

Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression,*

Third Ed. Sage Publications. https://www.john-fox.ca/Companion/

Fox, R. (2012). The decline of moths in Great Britain: a review of possible

causes. *Insect Conservation and Diversity*, *6*(1), 5–19.

https://doi.org/10.1111/j.1752-4598.2012.00186.x

Frankie, G. W., & Ehler, L. E. (1978). Ecology of Insects in Urban

Environments. *Annual Review of Entomology*, *23*(1), 367–387.

https://doi.org/10.1146/annurev.en.23.010178.002055

Franzén, M., Betzholtz, P.-E., Pettersson, L. B., & Forsman, A. (2020).

Urban moth communities suggest that life in the city favours

thermophilic multi-dimensional generalists. *Proceedings of the Royal*

Society B: Biological Sciences, *287*(1928), 20193014.

https://doi.org/10.1098/rspb.2019.3014

Gaona, F. P., Iñiguez-Armijos, C., Brehm, G., Fiedler, K., & Espinosa, C. I.

(2021). Drastic loss of insects (Lepidoptera: Geometridae) in urban

landscapes in a tropical biodiversity hotspot. *Journal of Insect*

Conservation, *25*(3), 395–405. https://doi.org/10.1007/s10841-021-

00308-9

GIS DCP. (2025). *NYC Borough Boundary* [shapefile]. Arcgis.com.

https://hub.arcgis.com/datasets/DCP::nyc-borough-boundary/about

Giuliano, W. M., Accamando, A. K., & Mcadams, E. J. (2004). Lepidoptera-

habitat relationships in urban parks. *Urban Ecosystems*, *7*(4), 361–

370. https://doi.org/10.1007/s11252-005-6835-7

Golsee, S. C., & Urban, D. (2007). The ecodist package for dissimilarity-

based analysis of ecological data. *Journal of Statistical Software*,

22(7). 10.18637/jss.v022.i07

Grenis, K., Nufio, C. R., Wimp, G. M., & Murphy, J. M. (2023). Does artificial

light at night alter moth community composition? *Philosophical*

Transactions of the Royal Society B, *378*(1892).

https://doi.org/10.1098/rstb.2022.0365

Hagen, J. B. (1999). Retelling Experiments: H.B.D. Kettlewell's Studies of

Industrial Melanism in Peppered Moths. *Biology & Philosophy*, *14*(1),

39–54. https://doi.org/10.1023/a:1006576518561

Hallmann, C. A., Sorg, M., Jongejans, E., Siepel, H., Hofland, N., Schwan,

H., Stenmans, W., Müller, A., Sumser, H., Hörren, T., Goulson, D., &

de Kroon, H. (2017). More than 75 percent decline over 27 years in

total flying insect biomass in protected areas. *PLOS ONE*, *12*(10),

e0185809. https://doi.org/10.1371/journal.pone.0185809

Hamilton, C. (2024). *QGIS Density Analysis Plugin* (Version 2024.1.23)

[Computer software]. GitHub. https://github.com/hamiltoncj/qgis-

densityanalysis-plugin

Kassambara, A., & Mundt, F. (2020). _factoextra: Extract and Visualize the

Results of Multivariate Data Analyses_. R package version 1.0.7,

Kettlewell, H. (1955). Selection experiments on industrial melanism in the

Lepidoptera. *Heredity*, *9*(3), 323–342.

https://doi.org/10.1038/hdy.1955.36

Kettlewell, H. (1956). Further selection experiments on industrial melanism

in the Lepidoptera. *Heredity*, *10*(3), 287–301.

https://doi.org/10.1038/hdy.1956.28

Kristensen, N. P., Scoble, M. J., & Karsholt, O. (2007). "Lepidoptera

phylogeny and systematics: the state of inventorying moth and

butterfly diversity." Arthropod Systematics & Phylogeny, 65(1), 1-50.

Langevelde, F., Braamburg-Annegarn, M., Huigens, M. E., Groendijk, R.,

Poitevin, O., Deijk, J. R., Ellis, W. N., Grunsven, R. H. A., Vos, R., Vos,

R. A., Franzén, M., & WallisDeVries, M. F. (2018). Declines in moth

populations stress the need for conserving dark nights. *Global Change Biology*, *24*(3), 925–932. https://doi.org/10.1111/gcb.14008

Liang, H., He, Y.-D., Theodorou, P., & Yang, C.-F. (2023). The effects of urbanization on pollinators and pollination: A meta-analysis. *Ecology Letters*, *26*(9). https://doi.org/10.1111/ele.14277

Lintott, P. R., Bunnefeld, N., Fuentes-Montemayor, E., Minderman, J., Blackmore, L. M., Goulson, D., & Park, K. J. (2014). Moth species richness, abundance and diversity in fragmented urban woodlands: implications for conservation and management strategies. *Biodiversity and Conservation*, *23*(11), 2875–2901. https://doi.org/10.1007/s10531-014-0753-z

Lizee, M.-H., Tatoni, T., & Deschamps-Cottin, M. (2015). Nested patterns in urban butterfly species assemblages: respective roles of plot management, park layout and landscape features. *Urban Ecosystems*, *19*(1), 205–224. https://doi.org/10.1007/s11252-015-0501-5

Lovett, G. M., Tear, T. H., Evers, D. C., Findlay, S. E. G., Cosby, B. J., Dunscomb, J. K., Driscoll, C. T., & Weathers, K. C. (2009). Effects of Air Pollution on Ecosystems and Biological Diversity in the Eastern United States. *Annals of the New York Academy of Sciences*, *1162*(1), 99–135. https://doi.org/10.1111/j.1749-6632.2009.04153.x

McKinney, M. L. (2006). Urbanization as a major cause of biotic homogenization. *Biological Conservation*, *127*(3), 247–260. https://doi.org/10.1016/j.biocon.2005.09.005

Merckx, T., Kaiser, A., & Van Dyck, H. (2018). Increased body size along

urbanization gradients at both community and intraspecific level in

macro-moths. *Global Change Biology*, *24*(8), 3837–3848.

https://doi.org/10.1111/gcb.14151

Merckx, T., & Slade, E. M. (2014). Macro-moth families differ in their

attraction to light: implications for light-trap monitoring programmes.

Insect Conservation and Diversity, *7*(5), 453–461.

https://doi.org/10.1111/icad.12068

New, T. R. (2023). Moth Ecology and Conservation Importance. *Springer*

EBooks, 21–37. https://doi.org/10.1007/978-3-031-32103-0_2

Oksanen, J., Simpson, G., Blanchet, F., Kindt, R., Legendre, P., Minchin, P.,

O'Hara, R., Solymos, P., Stevens, M., Szoecs, E., Wagner, H.,

Barbour, M., Bedward, M., Bolker, B., Borcard, D., Carvalho, G.,

Chirico, M., De Caceres, M., Durand, S., Evangelista, H., FitzJohn, R.,

Friendly, M., Furneaux, B., Hannigan, G., Hill, M., Lahti, L., McGlinn,

D., Ouellette, M., Ribeiro, Cunha, E., Smith, T., Stier, A., Ter, Braak

C., Weedon, J. (2022). _vegan: Community Ecology Package_. R

package version 2.6-4,

Piano, E., Souffreau, C., Merckx, T., Baardsen, L. F., Backeljau, T., Bonte,

D., Brans, K. I., Cours, M., Dahirel, M., Debortoli, N., Decaestecker,

E., De Wolf, K., Engelen, J. M. T., Fontaneto, D., Gianuca, A. T.,

Govaert, L., Hanashiro, F. T. T., Higuti, J., Lens, L., & Martens, K.

(2020). Urbanization drives cross-taxon declines in abundance and

diversity at multiple spatial scales. *Global Change Biology*, *26*(3).

https://doi.org/10.1111/gcb.14934

Posit team (2023). RStudio: Integrated Development Environment for R

(ver. 2023.12.0.369). Posit Software, PBC, Boston, MA.

http://www.posit.co/

QGIS Development Team. (2024). QGIS Geographic Information System

(ver. 3.34.3-Prizren). QGIS Association. https://www.qgis.org/

Ramírez-Restrepo, L., & MacGregor-Fors, I. (2016). Butterflies in the city: a

review of urban diurnal Lepidoptera. *Urban Ecosystems*, *20*(1), 171–

182. https://doi.org/10.1007/s11252-016-0579-4

R Core Team (2023). _R: A Language and Environment for Statistical

Computing_. R Foundation for Statistical Computing, Vienna, Austria.

Rice, A. J. (2015). Community Patterns in Urban Moth Assemblages. *Journal*

of the Lepidopterists' Society, *69*(3), 149–156.

https://doi.org/10.18473/lepi.69i3.a2

Rosseel, Y. (2012). lavaan: An R Package for Structural Equation Modeling.

Journal of Statistical Software, 48(2), 1-36.

https://doi.org/10.18637/jss.v048.i02

Sanetra, D., Berger, J., Hartlieb, M., Simons, N. K., Walther, G., Nico

Blüthgen, & Staab, M. (2024). Disentangling how urbanisation

influences moth diversity in grasslands. *Insect Conservation and*

Diversity, *17*(2), 229–242. https://doi.org/10.1111/icad.12713

Santibañez-Andrade, G., Castillo-Argüero, S., Vega-Peña, E. V., Lindig-

Cisneros, R., & José Alejandro Zavala-Hurtado. (2015). Structural

equation modeling as a tool to develop conservation strategies using

environmental indicators: The case of the forests of the Magdalena

river basin in Mexico City. *Ecological Indicators*, *54*, 124–136.

https://doi.org/10.1016/j.ecolind.2015.02.022

Sharmin, M., Tjoelker, M. G., Esperon-Rodriguez, M., Katlav, A., Gilpin, A.-

M., Rymer, P. D., & Power, S. A. (2024). Urban greening with shrubs

can supercharge invertebrate abundance and diversity. *Scientific*

Reports, *14*(1), 8735. https://doi.org/10.1038/s41598-024-58909-8

Shirey, V., Belitz, M. W., Barve, V., & Guralnick, R. (2021). A complete

inventory of North American butterfly occurrence data: narrowing

data gaps, but increasing bias. *Ecography*, *44*(4), 537–547.

https://doi.org/10.1111/ecog.05396

Simpson, E. H. (1949). Measurement of Diversity. *Nature*, *163*(4148), 688–

688. https://doi.org/10.1038/163688a0

Theodorou, P., Radzevičiūtė, R., Lentendu, G., Kahnt, B., Husemann, M.,

Bleidorn, C., Settele, J., Schweiger, O., Grosse, I., Wubet, T., Murray,

T. E., & Paxton, R. J. (2020). Urban areas as hotspots for bees and

pollination but not a panacea for all insects. *Nature Communications*,

11(1), 1–13. https://doi.org/10.1038/s41467-020-14496-6

Usher, M. B., & Keiller, S. W. (1998). The macrolepidoptera of farm

woodlands: determinants of diversity and community

structure. *Biodiversity & Conservation*, *7*, 725-748.

https://doi.org/10.1023/a:1008836302193

Vaz, S., Manes, S., Khattar, G., Mendes, M., Lima, F., Teles, E., de, E.,

Gama-Maia, D., Maria Lucia Lorini, Valverde, M., & Paulo Cesar

Paiva. (2023). Global meta-analysis of urbanization stressors on insect

abundance, richness, and traits. *Science of the Total Environment*,

165967–165967. https://doi.org/10.1016/j.scitotenv.2023.165967

Wang, Y., Kotze, D. J., Vierikko, K., & Niemelä, J. (2019). What makes urban

greenspace unique – Relationships between citizens' perceptions on

unique urban nature, biodiversity and environmental factors. *Urban*

Forestry & Urban Greening, *42*, 1–9.

https://doi.org/10.1016/j.ufug.2019.04.005

Wei, T., & Simko, V. (2021). R package 'corrplot': Visualization of a

Correlation Matrix (Version 0.92). https://github.com/taiyun/corrplot

White, P. J. T. (2018). An Aerial Approach to Investigating the Relationship

between Macromoths and Artificial Nighttime Lights Across an Urban

Landscape. *Journal of Agricultural and Urban Entomology*, *34*(1), 1–

14. https://doi.org/10.3954/1523-5475-34.1.1

Wilson, J. F., Baker, D., Cheney, J., Cook, M., Ellis, M., Freestone, R.,

Gardner, D., Geen, G., Hemming, R., Hodgers, D., Howarth, S., Jupp,

A., Lowe, N., Orridge, S., Shaw, M., Smith, B., Turner, A., & Young, H.

(2018). A role for artificial night-time lighting in long-term changes in

populations of 100 widespread macro-moths in UK and Ireland: a

citizen-science study. *Journal of Insect Conservation*, *22*(2), 189–196.

https://doi.org/10.1007/s10841-018-0052-1

Yang, L., Zhao, S., & Liu, S. (2022). A global analysis of urbanization effects

on amphibian richness: Patterns and drivers. *Global Environmental*

Change, *73*, 102476. https://doi.org/10.1016/j.gloenvcha.2022.102476

Figure Captions

Figure 1. (a) On-the-ground sampling locations for 12 locations in Brooklyn

and Queens, New York City, USA, where we sampled Lepidoptera, and

National Land Cover (NLCD) data for New York City. (b) Species diversity

in 1-km hexagonal bins derived from iNaturalist data. Base maps by Open

Street Map.

Figure 2. (a) Stacked bar plot of NLCD land cover composition at on-the-

ground sampling sites. Colors represent the proportion of each land cover

type in 150m buffers around each sampling location. (b) Principal

components analysis of NLCD land cover for our on-the-ground sampling

locations and all hex-bins used with iNaturalist data. PC1 represents high-

intensity developed land cover. (c) Satellite images of sampling locations in

Brooklyn and Queens, New York.

Figure 3. (a) Stacked bar plot of Lepidoptera Family diversity per site for

on-the-ground sampling. (b) Number of observations of each Lepidoptera

Family across all iNaturalist sampling sites. Colored bars are Families

observed in iNaturalist and on-the-ground sampling. (c) Images of moths

taken during on-the-ground sampling.

Figure 4. Simpson's diversity (D-1) as it relates to land cover summarized

by principal components analysis: (a) PC1 representing high-intensity

developed land, (b) PC2 representing medium-intensity developed land, and

(c) PC3 representing open developed land and deciduous tree cover.

Figure 5. Relationships among predictor variables and Simpson's diversity

generated by the structural equation model. Nodes represent variables

included in the model, while edges indicate associations between them.

Thicker lines represent stronger relationships, with positive correlations

shown in green and negative correlations in red. Edge weights are

proportional to the strength of standardized regression coefficients.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Linsk et al. – NOCTURNAL URBAN MOTHS

731   *Figure 5*

MOTH DIVERSITY ACROSS NYC URBAN MOSAIC

Methods: Sample Moths Across Urban Gradient

-PC1
less urban

Land Cover PC1

+PC1
more urban

Late Night – Species Diversity
est = 0.812±0.062, p<0.001

Sampling

Light-trapping on the ground

Combined with
iNaturalist Observations

Higher Moth Diversity with:

Deciduous Trees – Species Diversity
at Night

est = -0.512±0.062
p<0.001

Med Night – Species Diversity

Intense-Urbanization p<0.001

est = 0.468±0.107
p<0.001

Results: Urbanization and Biodiversity

Diversity (1 – Simpson's D)

iNaturalist

on-the-ground

-0.25   0.0   0.25

Landcover PC1

Moth
Diversity

Deciduous
Trees

Open
Space

Grassland

Low-Intensity
Developed

Pasture

Mid-Intensity
Developed

High-Intensity
Developed

Shrub

↑ Deciduous Tree Cover

est = -0.348±0.064
p<0.001