The *Magnaporthe oryzae* MAX effector AVR-Pia binds a novel group of rice HMA domain-containing proteins

Josephine H.R. Maidment[1,2]*, Svenja C. Saile[1]*, Aurélien Bocquet[2], Céline Thivolle[1], Léo Bourcet[2], Lisa-Fatimatou Planel[1], Muriel Gelin[2], Thomas Kroj[1], André Padilla[2], Karine de Guillen[2]**, Stella Cesari[1]**

[1] PHIM Plant Health Institute, Université de Montpellier, INRAE, CIRAD, Institut Agro, IRD, Montpellier, France.
[2] Centre de Biologie Structurale, INSERM, CNRS, Université de Montpellier, 29 rue de Navacelles, 34090, Montpellier, France

*Contributed equally
**Corresponding authors

Stella Cesari

Tel: +33 4 99 62 48 62

Email: stella.cesari@inrae.fr

Karine de Guillen

Tel: +33 4 67 41 77 08

Email: karine.deguillen@cbs.cnrs.fr

ORCID

Josephine H.R. Maidment: 0000-0002-8229-2718
Svenja C. Saile: 0000-0003-2258-3166
Aurélien Bocquet: 0009-0007-0082-6098
Céline Thivolle: 0009-0000-5048-3619
Léo Bourcet: 0009-0004-4568-3543
Lisa-Fatimatou Planel: 0009-0007-5159-7109
Muriel Gelin 0000-0003-1320-8663
Thomas Kroj: 0000-0002-3752-1788
André Padilla: 0000-0001-9702-4612
Karine de Guillen: 0000-0002-5037-2115
Stella Cesari: 0000-0001-8558-0371

Keywords: pathogen effector, host virulence targets, rice blast fungus, rice, protein structure, plant defence, heavy metal associated domain, MAX effector

Abstract

Phytopathogenic fungi secrete effector proteins to promote virulence. MAX (*Magnaporthe* Avrs and ToxB-like) effectors form a sequence-diverse family sharing a conserved protein structure. AVR-Pia, a MAX effector from the rice blast fungus *Magnaporthe oryzae,* is recognised by the paired rice nucleotide-binding leucine-rich repeat immune receptors OsRGA4/OsRGA5 through direct binding to an integrated heavy metal-associated (HMA) domain in OsRGA5. Here, we identify previously unknown host targets of AVR-Pia: four HMA domain-containing rice proteins, belonging to the HMA Isoprenylated Plant Proteins (HIPPs) and HMA Plant Proteins (HPPs). AVR-Pia interacted with all four proteins, both in vitro and in planta, and bound their HMA domains with varying affinities. The crystal structure of AVR-Pia in complex with the HMA domain of OsHPP09 revealed the molecular details of the binding interface. Structure-guided mutagenesis of OsHPP09 identified a single point mutation which prevents AVR-Pia binding, providing a foundation for targeted engineering of HMA domains to evade effector binding.

Introduction

The fungal pathogen *Magnaporthe oryzae* (syn. *Pyricularia oryzae*) is the causative agent of rice blast, one of the most devastating diseases affecting cultivated rice worldwide[1,2]. *M. oryzae* also causes severe yield losses in other cereals, including millets, barley and wheat, threatening global food security[1].

To successfully colonise host plants, *M. oryzae* must evade plant defence responses and manipulate host cellular pathways to its advantage. Central to infection is the secretion of a large repertoire of effector proteins, many of which can be categorised into structural families[3]. In *M. oryzae,* the <u>Magnaporthe</u> <u>A</u>VRs and To<u>x</u>B-like (MAX) effectors form the largest structural family[4]. MAX effectors are defined by a conserved fold consisting of two β-sheets, each composed of three antiparallel β-strands and stabilised by one or more disulphide bonds. Despite this shared structure, there is little amino acid sequence similarity between MAX effectors. They differ in surface properties and some carry variable N- and C-terminal extensions, which can influence their folding pathway[5] and mediate interactions with host proteins[3,6–9].

AVR-Pia is a MAX effector present in approximately 20% of rice-infecting *M. oryzae* isolates. It is highly expressed during the biotrophic phase of host colonisation and contributes to virulence on rice[4,7,10]. AVR-Pia can be recognised by the intracellular immune receptor OsRGA5, a nucleotide-binding leucine-rich repeat (NLR) protein that functions together with the helper NLR OsRGA4 to activate immune responses and restrict the spread of *M. oryzae*[11–13]. Canonical plant NLRs consist of an N-terminal signalling domain, a central nucleotide-binding domain and a C-terminal leucine-rich repeat domain[14]. However, many NLRs also carry non-canonical integrated domains (IDs), thought to originate from host proteins targeted by pathogen effectors[15]. These IDs, found across diverse plant species, are believed to function as mimics or baits for effector recognition ("integrated decoy" model)[15–18]. Despite sharing limited sequence similarity, both AVR-Pia and a second MAX effector, AVR1-CO39, bind to the heavy metal-associated (HMA) ID of OsRGA5[11,12]. While this suggests that these effectors may target host proteins containing HMA domains to promote disease, no host targets of AVR-Pia or AVR1-CO39 have been identified, and their specific virulence functions remain unknown.

The OsRGA5 HMA ID shares sequence similarity with a family of HMA domain-containing metallochaperone-like proteins. The family can be subdivided into two groups: <u>H</u>MA domain-containing <u>I</u>soprenylated <u>P</u>lant <u>P</u>roteins (HIPPs) that carry a C-terminal CaaX isoprenylation

motif, and <u>H</u>MA domain-containing <u>P</u>lant <u>P</u>roteins (HPPs) that lack this motif[19,20]. The two

groups are collectively referred to hereafter as H(I)PPs. Isoprenylation, a post-translational

modification involving the covalent attachment of an isoprenoid lipid to the cysteine residue in

the CaaX motif, has been reported to facilitate membrane association[21]. Many HIPPs harbour

proline-rich regions between the HMA domain and the CaaX motif, potentially mediating

protein-protein interactions[19,22]. Typically, HMA domains feature a metal-binding MxCxxC

motif, implicating H(I)PPs in roles related to metal homeostasis[23,24]. However, approximately

a third of rice H(I)PPs lack one or both cysteines in their MxCxxC motif, suggesting functional

diversification beyond metal binding. H(I)PPs have undergone extensive expansion in

vascular plants, with approximately 100 members identified in rice[19,25–27]. Most remain

functionally uncharacterized, although several have been implicated in response to various

biotic and abiotic stresses[19,28–30]. Increasing evidence points to a broader role of H(I)PPs as

susceptibility factors in interactions with pathogens, pests, and viruses[27,31–41].

In cereals, H(I)PPs have been identified as susceptibility factors for *M. oryzae* infection[27,33,34].

A loss-of-function allele of *OsHIPP05* (*Pi21*) contributes to blast resistance in the field.

OsHIPP20 is also a susceptibility gene; a loss-of-function mutation reduces the susceptibility

of rice to *M. oryzae*[27]. Some H(I)PPs are targeted by *M. oryzae* MAX effectors AVR-Pik and

APikL2[8,27,42]. AVR-Pik specifically interacts with members of a phylogenetic subclade of

OsH(I)PPs, referred to as clade A, including OsHIPP19, OsHIPP20, OsHPP03, and

OsHPP04[8,27]. The paired rice NLR proteins OsPik-1 and OsPik-2 mediate recognition of AVR-

Pik, which requires direct binding of the effector to an HMA ID in OsPik-1[43–45]. The HMA

domain of OsPik-1 belongs to the same phylogenetic clade as the H(I)PPs targeted by AVR-

Pik, reinforcing the idea that effector targets may serve as templates for domain integration

into NLRs.

Structural analyses of complexes between AVR-PikF or APikL2 and the HMA domains of their

HIPP targets or cognate NLR receptors have revealed a conserved binding interface centring

on β3 of the MAX fold and β4 of the HMA domain[8,42,45,46]. By contrast, the crystal structure of

AVR1-CO39 bound to the HMA ID of OsRGA5 revealed a markedly different interface

involving β2 of both AVR1-CO39 and OsRGA5-HMA[12]. NMR titration experiments, structural

modelling, and mutagenesis assays indicated that AVR-Pia binds OsRGA5-HMA at a similar

interface[12,13,47]. Additionally, the crystal structure of AVR-Pia in complex with Os-Pikp-HMA

revealed a similar binding interface, indicating that AVR-Pia binds different HMA domains in a

structurally similar manner[48]. The crystal structure of the MAX effector Pwl2 bound to

OsHIPP43 also revealed an interface involving β2 of the Pwl2 MAX fold and β2 of OsHIPP43-

HMA[9]. However, unlike AVR-Pia and AVR1-CO39, Pwl2 has a C-terminal extension,

comprising an α-helix and an unstructured region, which contributes significantly to HMA

binding[9]. Collectively, these structural studies underscore the remarkable diversity with which

MAX effectors have evolved to bind various HMA domain-containing proteins. They support

the hypothesis that AVR-Pia and AVR1-CO39 target distinct H(I)PPs through interfaces that

differ from those employed by other MAX effectors.

Here, we identified four OsH(I)PPs as specific interactors of AVR-Pia, none of which are bound

by the MAX effectors AVR1-CO39, AVR-Pik and Pwl2. Using biochemical and biophysical

approaches, we characterised the interaction between AVR-Pia and one of the interactors,

OsHPP09. The crystal structure of AVR-Pia/OsHPP09-HMA complex showed global similarity

to the AVR1-CO39/OsRGA5-HMA complex but revealed more extensive intermolecular

hydrogen bonding. Guided by the AVR-Pia/OsHPP09-HMA structure, we introduced a point

mutation in OsHPP09 which disrupts its interaction with AVR-Pia without compromising the

HMA fold.

Results

AVR-Pia interacts with rice H(I)PPs in Y2H screening

As AVR-Pia is recognised by OsRGA5 through direct binding to its HMA ID[11,13], we

hypothesised that AVR-Pia may target HMA domain-containing proteins to promote host

susceptibility. We conducted pairwise yeast two-hybrid (Y2H) assays, using a signal peptide-

deleted version of AVR-Pia (dSP-AVR-Pia, 20-85 aa) fused to the GAL4 activation domain

(AD) against a rice HMA domain library[49] comprising 78 rice H(I)PP HMA domains fused to

the GAL4 DNA binding domain (BD). Five HMA domains interacted with AVR-Pia: OsHPP09

(LOC_Os03g02070), OsHIPP41 (LOC_Os03g06080), OsHPP11 (LOC_Os04g45130),

OsHIPP14 (LOC_Os04g39320) and OsHIPP39 (LOC_Os03g02860) (**Fig. 1a,**

Supplementary Fig. 1, Supplementary Table 1). When paired with AVR-Pia, OsHPP09-

HMA conferred strong yeast growth on interaction-selective media, indicating a robust

interaction, while yeast growth for the remaining candidates was weaker.

We additionally tested the HMA library against the well-characterised MAX effectors Pwl2 and

AVR-PikD. Consistent with earlier publications, we observed dSP-Pwl2 (22-145 aa) interacting

with the HMA domain of OsHIPP43, and dSP-AVR-PikD (22-113 aa) interacting with the HMA

domains of OsHIPP19, OsHIPP20, and other phylogenetically related HMA domains[8,9,27,34]

(**Supplementary Fig. 1**), confirming the sensitivity of our screen.

In parallel, we performed an independent Y2H screen using a rice cDNA library. This identified
OsHIPP21 (LOC_Os09g09830) as a putative interactor of AVR-Pia (**Supplementary Table**
1, Supplementary Table 2).

For each interacting HMA domain, we next tested whether full-length (FL) proteins also bind
AVR-Pia in pairwise Y2H assays. Interestingly, interaction between AVR-Pia and FL OsH(I)PP
proteins was only observed for OsHPP09, OsHPP11 and OsHIPP21 (**Fig. 1a and**
Supplementary Fig. 2), despite confirmed expression of all proteins (**Supplementary Fig.**
3a, 3b, 3c). To exclude the possibility that the isoprenylation (CaaX) motif of studied FL HIPPs
affects nuclear localisation in yeast[21], we mutated the cysteine to serine (**Supplementary Fig.**
3a, 3b). However, OsHIPP14^{C187S}, OsHIPP39^{C190S} and OsHIPP41^{C152S} did not interact with
AVR-Pia in Y2H assays (**Supplementary Fig. 2**).

The HMA domains of OsHPP09 and OsHPP11 share 91.5 % and 74.6 % amino acid sequence
identity with OsHPP10 (LOC_Os10g36200) (**Supplementary Fig. 4**). Surprisingly, AVR-Pia
did not interact with OsHPP10-HMA nor the FL protein in Y2H assays (**Fig. 1a**). By contrast,
OsHIPP21-HMA shares only 30-40% sequence identity with the HMA domains of the three
OsHPPs (**Supplementary Fig. 4**), and while the HMA domain of OsHIPP21 did not interact
with AVR-Pia, FL OsHIPP21 did. Taken together, these results suggest that AVR-Pia binds
and potentially targets OsHPP09, OsHPP11, and OsHIPP21.

Previous work classified MAX effectors into 20 structural groups[6]. MAX58 belongs to the same
group as AVR-Pia, but differs in surface properties[6]. dSP-MAX58 (20-86 aa) did not interact
with any of the tested OsH(I)PPs in Y2H assays (**Fig. 1a, Supplementary Fig. 1**), although
the effector protein was expressed (**Supplementary Fig. 3c**), suggesting that, despite their
shared structural fold, effectors with distinct surface properties target different host proteins.

AVR-Pia interacts with rice H(I)PPs in planta
To test whether AVR-Pia and the four OsH(I)PPs interact in planta, we performed split-
luciferase complementation assays in *Nicotiana benthamiana*. 3xFlag-CLuc-OsH(I)PP FL
proteins were co-expressed with 3xHA-NLuc-dSP-AVR-Pia or 3xHA-NLuc-dSP-MAX58. Co-
expression of FL OsHPP09, OsHPP11, and OsHIPP21 with AVR-Pia led to high
luminescence levels, exceeding those observed for the positive control OsRGA5$_{Cter}$ (883-1116
aa, including the HMA ID) (**Fig. 1b**). Strikingly, a strong luminescence signal was observed for
OsHPP10/AVR-Pia, within a similar range to OsHPP09 and OsHPP11 and exceeding
OsHIPP21 (**Fig. 1b**). Consistent with the Y2H results, no interaction was observed in planta
between AVR-Pia and other tested FL OsHIPPs, including OsHIPP14, OsHIPP39 and
OsHIPP41 (**Fig. 1b**). MAX58 showed no interaction with any of the tested OsH(I)PPs (**Fig.**
1b). Expression of all proteins was confirmed by immunoblot analyses (**Supplementary Fig.**

5). These data demonstrate that AVR-Pia interacts in planta with OsHPP09,

OsHPP10, OsHPP11 and OsHIPP21.

AVR-Pia interacts with the HMA domains of OsHPP09, OsHPP10, OsHPP11 and

OsHIPP21 in vitro with varying affinities

Recombinant AVR-Pia and HMA domains were produced and purified from *E. coli*

(**Supplementary Fig. 6**). We first tested for interaction between purified OsHPP09-HMA and

AVR-Pia using analytical size exclusion chromatography. When analysed alone, the peak

elution volumes for AVR-Pia and OsHPP09-HMA were 15.3 ml and 13.0 ml, respectively (the

latter is consistent with dimerisation, as reported for other HMA domains in solution[12,45]).

Following incubation with OsHPP09-HMA (1:1 effector:HMA ratio), the peak elution volume of

AVR-Pia shifted to 12.7 ml, consistent with formation of an AVR-Pia/OsHPP09-HMA complex

(**Supplementary Fig. 7**). The co-elution of the two proteins was supported by SDS-PAGE

analysis of elution fractions (**Supplementary Fig. 7**).

To determine the affinity with which AVR-Pia binds to the HMA domains of OsHPP09,

OsHPP10, OsHPP11 and OsHIPP21, we used isothermal titration calorimetry (ITC). Titration

of AVR-Pia into a solution containing any of the purified HMA domains resulted in negative

peaks in the titration curve (**Fig. 2, Supplementary Fig. 8**), indicating an exothermic binding

interaction. Dissociation constant (K_D) values were determined from binding isotherms (**Fig.**

2, Supplementary Fig. 8) fitted to a single site model.

AVR-Pia bound to OsHPP09-HMA with an apparent K_D of 115 nM (**Fig. 2**). Despite differing

in just six amino acids, AVR-Pia bound OsHPP10-HMA with significantly lower affinity, with

an apparent K_D of 5.26 µM. The binding affinities of AVR-Pia and OsHPP11-HMA or

OsHIPP21-HMA fell between these two extremes, with apparent K_Ds of 772 nM and 1.29 µM,

respectively (**Fig. 2, Supplementary Fig. 8, Supplementary Fig. 9, Supplementary Table**

3). The apparent K_Ds, determined by ITC, for the interactions between OsRGA5-HMA and

AVR-Pia or AVR1-CO39 were previously reported as 7.8 µM[13] and 5.1 µM[12], respectively.

AlphaFold Multimer models cannot differentiate between HMA domains which are

bound by AVR-Pia and those which are not

Several studies have used AlphaFold Multimer to distinguish interacting from non-interacting

pairs of effectors/host proteins[27,50,51]. We therefore tested whether structural models of AVR-

Pia in complex with experimentally verified interactors (OsHPP09-HMA, OsHPP10-HMA,

OsHPP11-HMA and OsHIPP21-HMA) showed differences, either in the models themselves

or in the associated confidence metrics, compared to structural models of AVR-Pia with non-

interacting HMA domains. We modelled AVR-Pia/HMA complexes using AlphaFold2 and

AlphaFold3. High confidence models were obtained for all AVR-Pia/HMA pairs except AVR-Pia/OsHIPP42-HMA using AlphaFold2 (**Supplementary Fig. 10**) and AVR-Pia/OsHIPP41-HMA using AlphaFold3 (**Supplementary Fig. 11**). For the AlphaFold2 models, interface analysis with qtPISA revealed a tendency towards increased interface area for AVR-Pia-interacting HMA domains compared to non-interacting HMA domains. (**Supplementary Fig. 10)**. By contrast, there were no significant differences for AlphaFold3 models (**Supplementary Fig. 11**). Overall, AlphaFold could not clearly differentiate between HMA domains bound or not bound by AVR-Pia.

AVR-Pia-H3 does not interact with OsHPP09, OsHPP10, OsHPP11 and OsHIPP21

Some rice-infecting *M. oryzae* isolates carry an allele of AVR-Pia, AVR-Pia-H3, which contains two non-synonymous polymorphisms (F24S and T46N)[11] (**Supplementary Fig. 12**). These polymorphisms interfere with OsRGA5-HMA binding and allow evasion of OsRGA4/OsRGA5-mediated immunity[11,13]. Pairwise Y2H assays revealed that AVR-Pia-H3 failed to interact with FL OsHPP09, OsHPP10, OsHPP11 and OsHIPP21 or their HMA domains (**Fig. 3a**). In planta split-luciferase complementation assays gave only weak luminescence signals for AVR-Pia-H3 with OsH(I)PPs, comparable to the negative control pairs OsRGA5$_{Cter}$/AVR-Pia-H3 and OsHPP09/MAX58, and markedly lower than the positive control OsHPP09/AVR-Pia (**Fig. 3b**). Protein expression in yeast and in planta was confirmed by immunoblot (**Supplementary Fig. 3c and 13**). Analysis of a 1:1 effector:HMA mixture of purified recombinant AVR-Pia-H3 and OsHPP09-HMA by analytical size exclusion chromatography showed no co-elution of the two proteins, indicating that they did not form a stable complex in vitro (**Fig. 3c**).

Together, these data indicate that the F24S and T46N polymorphisms that disrupt binding of AVR-Pia to the HMA ID of OsRGA5 also interfere with binding to its candidate host targets.

AVR1-CO39 does not interact with OsHPP09, OsHPP10, OsHPP11 and OsHIPP21

As the HMA ID of OsRGA5 also binds the MAX effector AVR1-CO39[11,12], we tested whether AVR1-CO39 binds to the same H(I)PPs as AVR-Pia. dSP-AVR1-CO39 (22-89 aa) did not interact with FL OsHPP09, OsHPP10, OsHPP11 or OsHIPP21, nor their HMA domains, in Y2H, whereas interaction with OsRGA5$_{Cter}$ was confirmed (**Fig. 4a**). Effector expression was confirmed by immunoblot (**Supplementary Fig. 3c**).

In split-luciferase complementation assays, we observed weak luminescence for AVR1-CO39 with FL OsHPP09, OsHPP10 and OsHPP11, similar to the luminescence observed with OsRGA5$_{Cter}$ but substantially lower than the level observed for AVR-Pia/OsHPP09 (**Fig. 4b**). Consistent with the Y2H results, no interaction was detected between FL OsHIPP21 and AVR1-CO39, similar to the negative control OsPi21/AVR1-CO39 (**Fig. 4b**). Protein expression was validated by immunoblot (**Supplementary Fig. 14**). Analytical size-exclusion

chromatography with purified proteins further supported the absence of complex formation (**Fig. 4c**).

Collectively, these results suggest that, while AVR1-CO39 and AVR-Pia are both recognised by the same HMA ID in OsRGA5, they do not bind the same H(I)PPs.

AVR-Pia interacts with a distinct group of H(I)PPs to AVR-Pik and Pwl2

Previous studies have shown that the MAX effectors AVR-PikC, AVR-PikD and Pwl2 target H(I)PPs[8,9,27,34]. However, dSP-AVR-PikC (22-113 aa), dSP-AVR-PikD (22-113 aa) and dSP-Pwl2 (22-145 aa) did not bind to the AVR-Pia interactors OsHPP09, OsHPP10, OsHPP11 and OsHIPP21 in Y2H assays (**Fig. 5a**), while we confirmed the known interactions of AVR-PikC/AVR-PikD with OsHIPP19/OsHIPP20, and Pwl2 with OsHIPP43 (**Fig. 5a**). Interestingly, AVR-Pia interacted with FL OsHIPP19 under low-stringency Y2H conditions (**Fig. 5a**).

The lack of interaction between AVR-PikC and AVR-PikD with FL OsHPP09, OsHPP10, OsHPP11 and OsHIPP21 was validated through split-luciferase complementation assays (**Fig. 5b**). AVR-Pia showed an interaction with OsHIPP19 similar to that observed for AVR-PikD, and a weaker interaction with OsHIPP20 (**Fig. 5b**). Co-expression of Pwl2 with OsHPP09, OsHPP10, OsHPP11 and OsHIPP21 resulted in very weak luminescence. Protein expression was confirmed by immunoblot (**Supplementary. Fig. 15**).

Together, our findings indicate that the HMA domains bound by AVR-Pia are not targeted by AVR-Pik and Pwl2, suggesting that despite sharing a conserved MAX effector fold, these effectors target different H(I)PPs.

The crystal structure of the AVR-Pia/OsHPP09-HMA complex reveals interface contacts underpinning high affinity binding

To understand the structural basis of the interaction between AVR-Pia and OsHPP09-HMA, we purified and crystallised the effector/HMA complex. Needle-like crystals were obtained in commercial screens and subsequent optimisation of conditions resulted in crystals suitable for diffraction studies. X-ray diffraction data were collected at the European Synchrotron Radiation Facility (ESRF) to a resolution of 1.65 Å. The structure was solved by molecular replacement, refined and validated as described in the Materials and Methods. Data collection, processing and refinement statistics are presented in **Supplementary Table 4**. The final refined model has been deposited in the Protein Data Bank (PDB accession code 9RSV).

As expected, OsHPP09-HMA adopts the well-characterised HMA domain fold, consisting of a four-stranded antiparallel β-sheet and two α-helices arranged with βαββαβ topology (**Fig. 6**). Structures of AVR-Pia have previously been determined by NMR spectroscopy[4,52] (PDB

accession codes 2MYW, 2N37) and X-ray crystallography[48] (PDB accession code 6Q76;

complex with Pikp-1 HMA ID). The structure of AVR-Pia determined here is highly similar

(RMSD of 0.357 Å to 6Q76, 0.951 Å to 2MYW, and 1.690 Å to 2N37) and adopts the six-

stranded β-sandwich fold characteristic of the MAX effectors (**Fig. 6**)[4,6].

Globally, the AVR-Pia/OsHPP09-HMA complex resembles the previously determined

structures of AVR-Pia/OsPikp-1-HMA (PDB 6Q76) and AVR1-CO39/OsRGA5-HMA (PDB

5ZNG) (**Supplementary Fig. 16**). The AVR-Pia/OsHPP09-HMA interface area is 521.7Å^2,

fractionally larger than AVR-Pia/OsPikp-HMA (460.7 Å^2) and AVR1-CO39/OsRGA5-HMA

(492.8 Å^2) (**Supplementary Table 5**). In all three complexes, β2 of the MAX effector is aligned

with β2 of the HMA domain, forming an antiparallel β-sheet comprising β6, β1, and β2 of the

MAX effector and the four-stranded β-sheet of the HMA domain (**Supplementary Fig. 16a**),

with hydrogen bonding between adjacent β-strands (**Supplementary Fig. 16b**).

In contrast to AVR-Pia/OsPikp-HMA and AVR1-CO39/OsRGA5-HMA, where the interface is

dominated by main chain hydrogen bonding, in OsHPP09-HMA/AVR-Pia, the side chains of

multiple residues in α1 of the HMA domain form intermolecular hydrogen bonds with the

effector. The side chains of OsHPP09^{E16} and OsHPP09^{K19} form hydrogen bonds with the

peptide backbone of AVR-Pia^{L38} (**Supplementary Fig. 16c**). OsPikp-1^{R203} is in the

corresponding position to OsHPP09^{K19}, however it is OsPikp-1^{R226}, in the β2-β3 loop, which

forms an analogous hydrogen bond with AVR-Pia^{L38} (**Supplementary Fig. 16c**). OsRGA5-

HMA has a basic residue (R1012) at the corresponding position to OsHPP09^{K19}, and the side

chain of OsRGA5-HMA^{R1012} is orientated towards the AVR1-CO39^{D35} side chain, however the

distance between them (4.1 Å) is greater than the 4 Å cutoff typically used when considering

salt bridge interactions (**Supplementary Fig. 16c**). At the start of β2, the OsHPP09^{D33} side

chain forms salt bridge interactions with the AVR-Pia^{R43} side chain (**Supplementary Fig. 16d**).

This aspartate residue is conserved in both OsPikp-HMA (D217), where it similarly forms a

salt bridge with AVR-Pia^{R43}, and OsRGA5-HMA (D1026), where it forms an indirect, water-

mediated contact with the backbone of AVR1-CO39^{T41} (**Supplementary Fig. 16d**). The

OsHPP09^{Q20} side chain forms hydrogen bonds with the side chain hydroxyl of AVR-Pia^{Y41}

(**Supplementary Fig. 16e**). The serine in the corresponding position in OsPikp-HMA (S204)

forms a water-mediated contact with the side chain hydroxyl of AVR-Pia^{Y41} (**Supplementary**

Fig. 16e). Finally, towards the C-terminal end of OsHPP09-α1, the side chain of OsHPP09^{D28}

forms a salt bridge with AVR-Pia^{R23} (**Supplementary Fig. 16f**). By contrast, the corresponding

amino acid in OsPikp-1, OsPikp-1^{S212}, forms a hydrogen bond with the side chain hydroxyl of

AVR-Pia^{Y85} (**Supplementary Fig. 16f**). Overall, the increased number of intermolecular

hydrogen bonds in the AVR-Pia/OsHPP09-HMA structure compared to the structure of AVR1-CO39/OsRGA5-HMA likely explains the difference in binding affinities.

To validate the interface observed in the crystal structure, NMR titration with ^{1}H-^{15}N-HSQC NMR 2D spectra of ^{15}N-labelled AVR-Pia alone and in the presence of unlabelled OsHPP09-HMA were performed. HSQC spectra for ^{15}N-labelled AVR-Pia had previously been collected, and the resonances of amide cross-peaks had been assigned[4]. Binding of AVR-Pia to OsHPP09-HMA alters the chemical environment of the amino acids located at the binding interface, resulting in a change in the chemical shift observed for those amino acids in the ^{15}N HSQC spectra. The ^{15}N HSQC spectra of the complex AVR-Pia with unlabelled OsHPP09-HMA were reassigned using ^{15}N,^{13}C-AVR-Pia and two- and three-dimensional NMR experiments **(Supplementary Fig. 17).** Depending on the exchange rate constant (kex) between the bound and unbound states and on the frequency difference ($\Delta\omega$) between the corresponding resonances in these two states, different NMR exchange regimes occur. NMR titration showed that the AVR-Pia:OsHPP09-HMA complex was in slow exchange with kex $<< \Delta\omega$ since separate resonances appeared for individual species **(Supplementary Fig. 18)**. An NMR slow exchange regime indicates a relatively high affinity (<0.1 µM) between the two proteins. The chemical shift perturbations of AVR-Pia in the absence and presence of OsHPP09-HMA were calculated, and significant changes in chemical shift ($> \overline{\Delta\delta} + 1\sigma$) were observed for multiple residues in β2 and β3 of AVR-Pia, consistent with the interface in the crystal structure **(Supplementary Fig. 19)**.

Replacement of OsHPP09^{E16} with alanine or arginine disrupts AVR-Pia binding

We aimed to identify point mutations in OsHPP09-HMA that interfere with AVR-Pia binding. First, we selected residues whose side chains contribute intermolecular hydrogen bonds or salt bridges and mutated these to alanine (E16A, K19A, Q20A, D28A, D33A) and, for acidic residues, to arginine (E16R, D28R, D33R). D33 is conserved in OsHPP09, OsHPP10, OsHPP11, and in the HMA IDs of Pikp-1 (D217) and RGA5 (D1026) (**Supplementary Fig. 16D**). By contrast, the corresponding residue in Pikm-HMA, which does not bind AVR-Pia[48], is histidine. We therefore also mutated D33 to histidine. Second, we used FoldX[53] to predict the effect of mutating each interface residue in OsHPP09-HMA to all other amino acids. Based on the calculated difference of interaction free energy (ΔΔG) upon mutation, we made three further mutations: K19W, I35K, and A37W.

Introduction of the point mutations E16A, E16R, D28A, D28R and I35K in OsHPP09 resulted in a complete loss of interaction with AVR-Pia in Y2H experiments (**Fig. 7a**). The remaining mutations (i.e., K19A, K19W, D33A, D33H, D33R, A37W) with the exception of Q20A, led to

reduced yeast growth on selective media suggesting a weakening of the interaction compared
to the wildtype OsHPP09/AVR-Pia pair (**Fig. 7a**). Immunoblot analysis revealed that the K19A,
K19W and I35K mutants exhibited lower protein expression levels, while all other mutant
constructs showed similar expression to wildtype OsHPP09 (**Supplementary Fig. 3d**).

To assess the effect of the mutations in the context of the full-length protein in planta, we
performed split-luciferase complementation assays. The E16A and E16R mutants showed
significantly reduced interaction with AVR-Pia, while the I35K mutant completely lost its ability
to interact with AVR-Pia, consistent with the Y2H results (**Fig. 7b**). In contrast, the D28A and
D28R mutations, which abolished interaction in Y2H assays, had only a minor effect on the
interaction with AVR-Pia in planta, similar to the other tested mutants (**Fig. 7b**). Immunoblot
analysis demonstrated that all proteins were expressed at comparable levels in planta
(**Supplementary Fig. 20**).

Next, we aimed to establish whether the E16A, E16R, D33H and I35K mutations affect the
structure of the HMA domain. Mutated HMA domains were produced in *E. coli*, purified to
homogeneity, and molecular weights were confirmed by intact mass spectrometry. Circular
dichroism (CD) spectra for OsHPP09^{E16A}-HMA, OsHPP09^{E16R}-HMA and OsHPP09^{D33H}-HMA
were comparable to the spectra for wildtype OsHPP09-HMA with features characteristic of α-
helices (negative peaks around 222 nm and 208 nm, positive peak around 193 nm) and β-
sheets (negative peak around 218 nm, positive peak around 195 nm) consistent with the HMA
domain fold (**Fig. 7c**). By contrast, OsHPP09^{I35K}-HMA showed features of disorder (low
ellipticity above 210 nm and a negative peak around 200 nm) (**Fig. 7c**), suggesting that the
lack of AVR-Pia binding to the I35K mutant is likely due to disruption of the HMA domain fold.

We obtained ^{1}H-^{15}N-HSQC NMR spectra of ^{15}N-AVR-Pia in the presence of (unlabelled) HMA
domains of OsHPP09^{E16A}, OsHPP09^{E16R}, and OsHPP09^{D33H}. The spectra of ^{15}N-AVR-Pia with
OsHPP09^{D33H}-HMA was highly similar to the spectra of ^{15}N-labelled AVR-Pia with WT
OsHPP09-HMA (**Fig. 7d**, **Supplementary Fig. 21**), although the exchange regime changes
at certain cross-peaks, suggesting lower binding affinity. By contrast, the spectra of ^{15}N-AVR-
Pia with OsHPP09^{E16A} or OsHPP09^{E16R} showed no chemical shift perturbation and largely
superimposed with the spectra of ^{15}N-AVR-Pia alone (**Fig. 7d**, **Supplementary Fig. 22**,
Supplementary Fig. 23). Taken together, these results demonstrate that E16A and E16R
mutations block AVR-Pia binding without disrupting the HMA fold, while the I35K mutation
interferes with correct folding of the HMA domain.

Discussion

In the present study, we found that the *M. oryzae* MAX effector AVR-Pia binds to the HMA domains of four rice H(I)PPs: OsHPP09, OsHPP10, OsHPP11, and OsHIPP21. Neither AVR-Pik nor Pwl2 interacts with these H(I)PPs, demonstrating that AVR-Pia binds a distinct group of H(I)PPs to those previously identified as MAX effector targets. Indeed, AVR-Pik interacts with multiple H(I)PPs within a large phylogenetic clade (referred to as "clade A")[27], while Pwl2 interacts specifically with OsHIPP43 and its homologs from other grass species[9].

Strikingly, the interactions between each of these three MAX effectors and HMA domains involve distinct interfaces. In the crystal structures of AVR-Pia/OsHPP09-HMA and Pwl2/OsHIPP43-HMA, β2 of the core MAX fold and β2 of the HMA domain are aligned in an antiparallel orientation[9], forming an antiparallel β-sheet comprising the four β-strands of the HMA domain and β2, β1, and β6 of the MAX fold. When the HMA domains are superposed, the relative positions of β2 of the MAX fold of Pwl2 and AVR-Pia are very similar (**Supplementary Fig. 24**). However, the other β-strands of Pwl2 are shifted compared to AVR-Pia, enabling residues in β3, β4, and the β3-β4 loop of the MAX fold to contribute to the HMA-binding interface. By contrast, in the OsHIPP19-HMA/AVR-PikF complex, an antiparallel β-sheet forms from the four β-strands of the HMA domain and β3, β4 and β5 of the MAX fold, mediated by hydrogen bonding between β3 of AVR-PikF and β4 of HIPP19-HMA[8]. Furthermore, while AVR-Pia consists solely of the core MAX fold, AVR-Pik has an unstructured extension at the N-terminus, while in Pwl2 the characteristic MAX fold is followed by an α-helix and a C-terminal unstructured region, and these extensions are involved in HMA binding (**Supplementary Fig. 24**). Pwl2^{E89}, in the α-helix, forms a hydrogen bond with OsHIPP43-HMA^{K51}, while the C-terminal unstructured segment of Pwl2 extends across the HMA domain, forming an extensive interface involving both hydrogen bonds and π-stacking interactions. Similarly, the N-terminal extension of AVR-Pik forms multiple hydrogen bonds with residues in β3, β2, and the β2-β3 loop of OsHIPP19-HMA. The interface in the AVR-Pia/OsHPP09-HMA complex is smaller than in AVR-PikF/OsHIPP19-HMA and Pwl2/OsHIPP43-HMA (521.6 Å^2 compared to 1044.3 Å^2 and 1965.2 Å^2, respectively), and is formed almost exclusively by residues in β2 of the core MAX fold.

It is highly interesting that at least three MAX effectors have evolved to bind HMA domains in different ways. This highlights how structural variation within this effector family can underpin different interactions with host targets. It is unknown whether the interactions of AVR-Pia, AVR-Pik, and Pwl2 with OsH(I)PPs lead to similar outcomes (redundancy) or if each binding event has a distinct and specific impact on plant susceptibility and/or fungal virulence.

While AVR-Pia, AVR-Pik, and Pwl2 all target OsH(I)PPs, the MAX effector AvrPiz-t has been

reported to interact with a range of host proteins[54,55]. From a structural perspective, it is unclear

how a single MAX effector binds to such diverse targets. However, this suggests that MAX

effectors may target many different host proteins and processes during infection.

AVR-Pia is recognised by the paired rice NLR immune receptors OsRGA5 and OsRGA4

through direct binding of the effector to the non-canonical HMA domain integrated into

OsRGA5[11]. Such non-canonical domains are hypothesised to have originated in effector host

targets and were incorporated into NLR proteins to function as effector "sensors"[15]. Consistent

with this, the crystal structure of AVR-Pia/OsHPP09 revealed that AVR-Pia binds OsHPP09-

HMA and OsRGA5-HMA through similar interfaces[13]. Interestingly, the F24S and T46N

polymorphisms of AVR-Pia-H3 interfered with binding to both OsRGA5-HMA and the

candidate OsH(I)PP targets, in contrast to naturally-occurring polymorphisms in AVR-Pik

which prevent binding to the HMA ID of Pikp-1 without disrupting interaction with the HMA

domains of OsH(I)PP19 and OsH(I)PP20[8,27].

AVR1-CO39 is also recognised by direct binding to the HMA ID of OsRGA5. However, AVR1-

CO39 did not interact with the same OsH(I)PPs as AVR-Pia. AVR1-CO39 also did not interact

with the "clade A" OsH(I)PPs bound by AVR-PikD[27]. AVR1-CO39 is broadly absent from rice-

infecting isolates of *M. oryzae*[7,56,57] and its loss has potentially contributed to a host shift from

foxtail millet (*Setaria* sp.) to rice. It is therefore possible that AVR1-CO39 has evolved to target

H(I)PPs from other grass species, and does not interact with those present in rice.

Interestingly, phylogenetic analysis based on HMA domains clusters OsRGA5-HMA with the

H(I)PPs targeted by AVR-Pik and the HMA ID of OsPik-1, and not with OsHPP09, OsHPP10,

OsHPP11, and OsHIPP21[27]. This suggests that the RGA5-ID did not arise through duplication

and integration of these AVR-Pia candidate targets, but instead from another H(I)PP. We

observed interaction between AVR-Pia and FL OsHIPP19, but not with the HMA domain

alone, in low stringency Y2H experiments and split-luciferase assays. This suggests that the

C-terminal part of OsHIPP19 contributes to AVR-Pia binding and merits further study.

A previous study reported that AVR-Pia binds OsRGA5-HMA with a K_D of 7.8 μM[13]. This is

comparable to the binding affinity of AVR-Pia to OsHPP10-HMA, but at least one order of

magnitude weaker than the AVR-Pia/OsHPP09-HMA interaction. This higher affinity can be

explained by the number of intermolecular hydrogen bonds and salt bridges at the AVR-

Pia/OsHPP09 interface; available structures of AVR1-CO39/OsRGA5-HMA and AVR-

Pia/OsPikp-HMA, combined with modelling approaches, suggest that the AVR-Pia/OsRGA5-

HMA interface likely involves fewer intermolecular contacts. A similar pattern was reported for

AVR-Pik/HMA interactions; surface plasmon resonance experiments showed that AVR-PikD interacts with the HMA domain of OsHIPP19 with higher affinity than the HMA ID of OsPik-1[8]. It is important to note that these binding affinities are determined in vitro using purified HMA domains. These experiments therefore lack the biological context of the FL protein or the plant cell environment.

H(I)PPs, which form extensive families in plants[19,25,26], have been reported to be involved in intracellular metal transport and homeostasis, responses to cold and drought stress, and heavy metal tolerance. In addition to the interactions between *M. oryzae* MAX effectors and OsH(I)PPs, there are multiple examples of plant H(I)PPs targeted by proteins from diverse pests and pathogens. For example, the putative effector RsMf8HN from the necrotrophic fungus *Rhizoctonia solani* interacts with OsHIPP28[38], while a HIPP from sugarcane (ScPi21) is a target for the *Fusarium sacchari* effector Fs00367[35]. CsHIPP03 was identified by Y2H screening as an interactor of SDE34, an effector secreted by the phloem-colonising bacteria *Candidatus* Liberibacter asiaticus[40]. A root-knot nematode (*Meloidogyne graminicola*) effector binds rice OsHPP04 and suppresses the flg22-induced ROS burst[36]. Finally, the potato mop-top virus movement protein interacts with NbHIPP26 to promote systemic spread of the virus[31]. Together, these results suggest that the manipulation of H(I)PPs is required for successful infection by diverse pests and pathogens. However, little is known about the molecular mechanisms through which H(I)PPs affect plant resistance and/or susceptibility.

HMA domains typically coordinate metal ions via two cysteine residues in an MxCxxC motif. Intriguingly, the metal-binding motif is degenerate in all four HMA domains bound by AVR-Pia, with both cysteines replaced by other amino acids (MT<u>D</u>EK<u>T</u> in OsHPP09, MT<u>D</u>DK<u>I</u> in OsHPP10, MS<u>D</u>TKM in OsHPP11, and MT<u>D</u>ER<u>K</u> in OsHIPP21), indicating the absence of metal-binding. Consistent with this, no electron density suggestive of bound metal ions was observed in the OsHPP09-HMA/AVR-Pia crystal structure. Interestingly, OsHPP09[E16], which is important for AVR-Pia binding, is located in the degenerate metal binding motif. Of the "clade A" H(I)PPs bound by AVR-Pik, some contain an intact MxCxxC motif, while others, such as OsHIPP19 and OsHIPP20, lack one or both cysteine residues. OsHIPP43 has an intact MxCxxC motif, though it is unknown whether it binds metal ions in planta. In cases where HMA domains lack a complete MxCxxC motif, and presumably also the capacity to bind metals, the functions of the H(I)PPs are particularly intriguing. Functional diversification, beyond metal transport and homeostasis, may have been facilitated by the stable core fold of the HMA domain supporting diversification of surface properties to deliver new functions. Protein prenylation can affect subcellular localisation, protein stability, and protein-protein interactions[58]. For some HIPPs, the C-terminal CaaX isoprenylation motif has been shown to

drive localisation to plasmodesmata. Further investigation is required to establish the

localisation and function of the AVR-Pia interactors in planta.

Pathogens require certain host factors to establish a successful infection, with the underlying

genes typically referred to as susceptibility (S) genes. Previous studies have shown that

OsHIPP05 (*Pi21*) and *OsHIPP20* are S genes for blast infection; *oshipp05* and *oshipp20*

CRISPR-Cas9 knockout plants exhibited increased blast resistance compared to WT plants

[27,33,34,59–61]. Further, a recent study reported that overexpression of barley (*Hordeum vulgare*)

HvHIPP43 increased susceptibility to *M. oryzae,* suggesting that *HvHIPP43* may also be an S

gene[34]. Other plant H(I)PPs are also S genes for different pests and pathogens; *Arabidopsis*

thaliana plants carrying loss-of-function mutations in *AtHIPP27* or *AtHMAD* (also known as

AtHIPP40[29]) exhibited reduced susceptibility to the beet cyst nematode *Heterodera schachtii*[32]

and the bacterial pathogen *Pseudomonas syringae*[41], respectively. Silencing of *TaHIPP1* in

wheat reduced sporulation of the wheat stripe rust fungus *Puccinia striiformis* f. sp. *tritici*[39].

OsHPP09, *OsHPP10*, *OsHPP11,* and/or *OsHIPP21* may also be S genes for rice blast

disease, and deletion of these genes may increase blast resistance.

Knocking out S genes has the potential to confer durable resistance, but can cause

undesirable pleiotropic effects on plant growth and development[62–65]. Base editing offers a

more precise alternative, allowing single amino acid changes that minimise disruption of

protein function[66]. While the functions of OsHPP09, OsHPP10, OsHPP11, and OsHIPP21,

and the consequences of AVR-Pia binding remain unknown, the virulence function of the MAX

effector Pwl2 depends on interaction with OsHIPP43[34]. Altering effector targets to block

effector binding could compromise the virulence function of effectors without compromising

the folding or endogenous function of the host protein. However, it is crucial to verify that such

mutations do not interfere with the folding and function of the host targets before commencing

time-consuming and costly base editing experiments in plants.

In summary, we have characterised the interactions between AVR-Pia and a novel subset of

HMA domain-containing rice proteins. Biophysical and structural analyses guided

mutagenesis of these effector targets and identified a mutation that blocks effector binding

without compromising structural integrity of the HMA domain. This provides a foundation for

targeted modification of these proteins for enhanced disease resistance.

Methods

Cloning

The coding sequences (CDS) of *OsH(I)PPs*, including the stop codon, were synthesised by Twist Bioscience. The CDS of *OsHIPP14*, *OsPi21*, *OsHIPP19*, *OsHIPP21,* and *OsHPP09* were codon-optimised to enable synthesis.

For Y2H and split luciferase experiments, synthesised *OsH(I)PP* genes were used as template for amplification with primers containing *attB1* and *attB2* recombination sites (**Supplementary Table S6**) and Phusion High-Fidelity DNA Polymerase (Thermo Fisher Scientific). Resulting PCR products were subsequently introduced into the pDONR207 vector (Invitrogen) via Gateway® BP recombination (Thermo Fisher Scientific). Entry constructs for the MAX effectors dSP-AVR-Pia[15], dSP-AVR1-CO39[12], dSP-AVR-PikD[47] and dSP-Pwl2[13] were previously published. The dSP-AVR-PikC fragment was amplified from pCambia1300_dSP-AVR-PikC[45] and dSP-MAX58 from a synthesised gene, both using primers containing *attB1* and *attB2* recombination sites (Supplementary Table S6) and Phusion High-Fidelity DNA Polymerase (Thermo Fisher Scientific). Resulting PCR products were cloned into the pDONR207 vector (Invitrogen) via Gateway® BP recombination (Thermo Fisher Scientific).

For recombinant protein production in *E. coli,* synthesised genes were used as templates for PCR amplification of HMA domains (**Supplementary Table S1**) with primers containing *Bpi*I restriction sites (**Supplementary Table S6**) using CloneAmp™ HiFi PCR Premix (Clontech Laboratories). Resulting PCR products were introduced into pICH41308 (Addgene #47998[67]) by Golden Gate assembly[68] to generate level 0 CDS modules. The CDS of OsHIPP21-HMA was codon-optimised for *E. coli* expression and supplied with appropriate overhangs and restriction sites for direct assembly into the vector pICH41308 without PCR amplification. Level 0 CDS modules were used in Golden Gate reactions with the pOPIN-GG vector pPGN-C (Addgene #174578[69]) and the level 0 module pICSL30015 (Addgene #174582[69]) to generate *E. coli* expression constructs with a cleavable N-terminal 6xHis-MBP tag.

Site-directed mutagenesis of OsHPP09-HMA for Y2H, split luciferase, and recombinant protein production was performed using a QuikChange Site-Directed Mutagenesis Kit (Agilent). Prenylation motif mutations (*OsHIPP14^{C187S}*, *OsHIPP21^{C133S}*, *OsHIPP39^{C190S}*, *OsHIPP41^{C152S}*) in the pDONR constructs*,* and mutations in AVR-Pia to give *AVR-Pia-H3*, were introduced by site-directed mutagenesis PCR using Phusion High-Fidelity DNA Polymerase (Thermo Fisher Scientific) and mutation-specific primers listed in **Supplementary Table S6**.

Gateway® LR recombination using LR Clonase® II mix (Thermo Fisher Scientific) was used

to transfer *OsH(I)PPs* and *RGA5$_{Cter}$*[11] (883-1116 aa) into Gateway®-compatible destination

vectors, including a modified Y2H vector pGBKT7-GW[70] and the plant expression vector pBIN-

35S-3Flag-Cluc-GWY (Deslandes et al., unpublished). Similarly, MAX effector

pDONR constructs were recombined into a modified pGADT7-GW[70] and pBIN-35S-3HA-Nluc-

GWY (Deslandes et al., unpublished) destination vectors. The pBIN-35S-3HA-Nluc-GWY and

pBIN-35S-3Flag-Cluc-GWY vectors were generated by ligation of a 3HA-Nluc-FrameB or

3Flag-Cluc-FrameB fragment, respectively, into a XhoI/XbaI digested pBIN-35S vector.

Previously published expression constructs for AVR1-CO39, AVR-Pia, and AVR-Pia-H3[4,12,13]

were used for protein production in the present study.

A complete list of plasmids used in this study is provided in **Supplementary Table S7.**

Yeast two-hybrid

The yeast strain Y2HGold (mating type a; Takara Bio) was transformed with the bait construct

expressed from the pGBKT7 vector, which contains the GAL4-BD and the *TRP1* marker gene.

These constructs included previously described HMA domains [49], as well as newly generated

constructs: OsHIPP21-HMA, OsH(I)PP^{WT} FL and OsH(I)PP$^{C/S}$ FL. The yeast strain Y187

(mating type α; Takara Bio) was transformed with the prey construct (MAX effectors)

expressed from the pGADT7 vector, which carries the GAL4-AD and the *LEU2* marker

gene. Yeast transformation was performed using a lithium acetate (LiAc)-based protocol

according to the Yeastmaker Yeast Transformation System (Takara Bio). Transformed yeast

cells were plated on synthetic defined (SD) media lacking either tryptophan (-W) or leucine (-

L) and incubated at 30 °C for 3 days.

Mating of the transformed yeast strains followed the Matchmaker Gold Yeast Two-Hybrid

System User Manual (Takara Bio). Diploid yeast cells were selected on SD/-L-W plates (SD

medium lacking leucine and tryptophan) after incubation at 30 °C for 2 days. For Y2H assays,

diploid yeast cells were inoculated in SD/-L-W medium, grown for 24 hours at 28°C with

shaking at 100 rpm, then pelleted and washed twice with sterile water. Serial 10-fold dilutions

(10^{-1}, 10^{-2}, and 10^{-3}) were spotted onto SD selection plates: SD/-L-W (growth control), SD/-

L-W-H (lacking leucine, tryptophan, and histidine), SD/-L-W-H supplemented with varying

concentrations of 3-amino-1,2,4-triazole (3AT). After 7 days incubation at 30 °C, plates were

photographed using the presentation station UF-130DX (Samsung).

Yeast two-hybrid cDNA library screening

The Y2H cDNA library derived from the rice cultivar CO39 and the BD:dSP-AVR-Pia construct, have been previously described[11,47]. Library screening followed the Matchmaker Gold Y2H system protocol (Takara Bio). Mating was performed between the Y2H Gold strain harbouring GAL4-BD:AVR-Pia and the Y187 strain carrying the CO39 cDNA library. Diploid yeasts were plated on SD/-L-W-H medium and supplemented with 20 mM 3AT. Colonies that grew were re-streaked onto fresh SD/-L-W-H medium containing 20 mM 3AT to confirm interaction-dependent growth. Yeast colony PCR was used to amplify cDNA inserts from positive clones, and PCR products were sequenced. Sequences were aligned to the cDNAs of two publicly available annotations of the *Oryza sativa* cv. Nipponbare reference genome (IRGSP-1.0[71] and MSU7.0)[72] using BLAST to identify the corresponding genes. Only genes for which the cDNA inserts (or fragments of cDNAs) were cloned in-frame with the GAL4-AD were retained as candidate AVR-Pia targets.

Agrobacterium-mediated transient expression in *N. benthamiana*

Agrobacterium tumefaciens strains AGL1 (carrying split-luciferase constructs) and GV3101-pMP90 (carrying 35S::P19) were grown overnight at 28°C in 5 mL LB medium supplemented with appropriate antibiotics. Overnight cultures were pelleted at 4,000 x *g* for 10 minutes, and pellets resuspended in 2 mL infiltration buffer (10 mM $MgCl_2$, 10 mM MES pH 5.6, 150 µM acetosyringone). Bacterial suspensions were incubated in the dark for 2 hours. The optical density at 600 nm (OD_{600}) was adjusted to 0.05 for 35S::P19 and 0.1 for OsH(I)PPs and MAX effectors, respectively. In all infiltration mixes, 35S::P19 was co-infiltrated to suppress gene silencing. Agrobacterial suspensions were mixed and infiltrated into leaves of 5–6-week-old *N. benthamiana* plants using a needleless syringe. Leaf discs were harvested 2 days post-infiltration for luminescence measurements or protein extraction. *N. benthamiana* plants were grown in a growth chamber on soil under a 16-hour light / 8-hour dark cycle, at 20 °C and 65 - 75% relative humidity.

Split luciferase assay

For luminescence measurements, 4 mm leaf discs (collected using a Miltex Biopsy Punch (Delta Microscopie)) were placed into wells of a white 96-well plate containing water. The water was removed and replaced with a 1 mM Xenolight D-luciferin-K$^+$ salt bioluminescent substrate solution (Perkin Elmer). Luminescence was quantified using a Spark® microplate reader (Tecan) over three cycles of 10 minutes each. For all experiments, data from the third cycle were used for analysis. Each interaction was tested with eight technical replicates (eight leaf discs per plate from four individual plants per biological replicate, unless otherwise indicated in the figure legend). The luminescence assay was performed in three independent

experiments. Statistical analyses from three independent replicates (24 leaf discs in total, unless otherwise indicated in the figure legend) and data visualisation were done using R (version 4.4.1). Multiple comparisons were performed using pairwise Wilcoxon tests followed by a Benjamini-Hochberg p-value adjustment with the stats package. Boxplots were generated using the ggplot2 package.

Protein extraction and Western blotting

Protein extraction from yeast was performed using a post-alkaline extraction protocol, as described by Kushnirov[73]. For BD-OsHIPP14/19/20/39/41 and AD-MAX, yeast crude extracts were resuspended in 50 μL of 1x LDS loading buffer (1x Bolt™ LDS sample buffer (Thermo Fisher Scientific), 1x NuPAGE™ sample reducing agent (Thermo Fisher Scientific)), boiled at 95°C for 10 minutes, and loaded onto a 10% NuPAGE™ Bis-Tris gels (Invitrogen). For BD-OsHPP09/10/11 and OsHIPP21, yeast crude extracts were resuspended in 50 μL of 1x SDS-Tricine buffer (1x SDS-Tricine buffer (Thermo Fisher Scientific), 1x NuPAGE™ sample reducing agent (Thermo Fisher Scientific)) and boiled at 85 °C for 2 minutes before loading onto a 10-20% Tricine gel (Novax). Proteins were transferred to iBlot® nitrocellulose membrane (Invitrogen) using the iBlot 2 Gel Transfer Device (program: 20 V 1 min, 23 V 4 min, 25 V, 2 min, Thermo Fisher Scientific), and analysed by immunoblotting.

For protein extraction from *N. benthamiana*, four leaf discs (0.8 mm diameter) were collected, immediately frozen in liquid nitrogen and ground using a tissue homogeniser (Qiagen). The ground plant material was solubilised in 200 μL extraction buffer (20 mM Tris-Hcl pH 7.5, 150 mM NaCl, 1 mM EDTA pH 8.0, 1% Triton X-100, 0.1% SDS, 5 mM DTT, 1x Protease inhibitor cocktail (Sigma-Aldrich), 1x EDTA-free cOmplete protease inhibitor (Roche)). Protein extracts were centrifuged at 18,000 x g for 15 minutes at 4 °C, and the resulting supernatants retained. 2x LDS buffer (Bolt™ LDS sample buffer (Thermo Fisher Scientific) and NuPAGE™ sample reducing agent (Thermo Fisher Scientific)) was added and samples were denatured at 70 °C for 10 minutes. Proteins were resolved by electrophoresis on NuPAGE™ 4-12% Bis-Tris gels (Invitrogen). Proteins were transferred to iBlot® nitrocellulose membrane (Invitrogen) as described above and analysed by immunoblotting.

For immunodetection of proteins, rabbit anti-GAL4-BD (sc-577, Santa Cruz Biotechnology, 1:2,000), goat anti-rabbit IgG-HRP (1:20,000, Sigma-Aldrich, A0545), mouse anti-Flag M2-HRP (1:60,000, Sigma-Aldrich) and rat anti-HA-HRP (clone 3F10, 1:1,1000, Roche) were used. Chemiluminescence was detected using the Immobilon western kit (Millipore) or the SuperSignal™ West Femto Maximum Sensitivity substrate (Thermo Fisher Scientific) and the Syngene 680X EF GBOX imaging system (Syngene).

Protein production and purification

HMA domains

HMA domains were produced with a cleavable N-terminal 6xHis-MBP tag from *E. coli* BL21 (DE3) or BL21 (DE3) pLysS. 700 mL cell cultures were grown in LB media at 37 °C from a starting OD_{600} of 0.05-0.07 to an OD_{600} of 0.5-0.8. Protein production was induced by addition of IPTG (final concentration 1 mM) and cultures were grown for a further 14-16 hours at 20 °C. Cells were pelleted by centrifugation and the pellets stored at -70 °C. Thawed pellets were resuspended in 50 mM Tris-HCl pH 7, 300 mM NaCl, 30 mM imidazole supplemented with cOmplete EDTA-free protease inhibitor cocktail (Roche). Cells were lysed by sonication and the lysate clarified by centrifugation at 40,000 x *g* at 6 °C for 30 minutes. Subsequent chromatography steps were carried out with an ÄKTA Pure system (Cytiva) at 4 °C. The filtered (0.45 μm) supernatant was applied to a 5 ml HisTrap[TM] FF column (Cytiva) and, after washing with 50 mM Tris-HCl pH 7, 300 mM NaCl, 30 mM imidazole, bound protein was step-eluted with 50 mM Tris-HCl pH 7, 300 mM NaCl, 500 mM imidazole. The eluate was injected onto a Superdex 75 26/60 size exclusion chromatography column (Cytiva) pre-equilibrated in 50 mM Tris-HCl pH 7, 150 mM NaCl. Fractions containing the fusion protein were incubated with 3C protease overnight at 4 °C. HMA domains were separated from the 6xHis-MBP tag and GST-tagged 3C protease by injecting the protease-treated sample onto a 5 ml HisTrap[TM] FF column (Cytiva), 5 ml MBPTrap[TM] HP column (Cytiva), and 5 ml GSTrap[TM] HP column (Cytiva) connected in tandem and equilibrated in 50 mM Tris-HCl pH 7, 300 mM NaCl, 30 mM imidazole. Pooled fractions containing HMA domains were concentrated to < 8 ml then injected onto a Superdex 75 26/60 size exclusion chromatography column (Cytiva) pre-equilibrated in 50 mM Tris-HCl pH 7, 150 mM NaCl. Pooled fractions containing the purified HMA domains were concentrated. At each stage, fractions containing the HMA domain were determined by SDS-PAGE. Samples were combined with 5X SDS loading buffer (60 mM Tris-HCl pH 6.8, 25% glycerol, 4% SDS, 0.1% bromophenol blue, 10 mM DTT), heated at 95 °C for 5 minutes, then loaded onto a Bolt™ 10 %, Bis-Tris gel (Invitrogen). After migration, gels were stained in InstantBlue® Coomassie protein stain (Abcam). Final protein concentration was determined using a Pierce™ BCA Protein Assay Kit (ThermoFisher Scientific), due to the extreme molar extinction coefficients of the proteins (0 $M^{-1}cm^{-1}$ for OsHIPP21-HMA, 1490 $M^{-1}cm^{-1}$ for OsHPP-HMA domains) at 280 nm. Proteins were stored at -80 °C.

MAX effectors

AVR-Pia, AVR-Pia-H3 and AVR1-CO39 were produced and purified as previously described[4,12,13]. Concentrations of MAX effectors were determined from molar extinction coefficients and absorbance at 280 nm.

Analytical size exclusion chromatography

Analytical size exclusion chromatography experiments were conducted using a Superdex 75 10/300 GL column (Cytiva) connected to an ÄKTA Pure system (Cytiva) at 4 °C. Running buffer was 50 mM Tris pH 7, 150 mM NaCl, 1 mM DTT for experiments with AVR1-CO39 and 20 mM sodium citrate pH 5.6, 150 mM NaCl, 1 mM DTT for experiments with AVR-Pia and AVR-Pia-H3. To investigate whether OsHPP09-HMA forms a complex with the effectors, the two proteins were combined in a 1:1 molar ratio and incubated on ice for 1 h prior to analysis. Each protein was also analysed alone, at a concentration equivalent to that present in the mixture. For each experiment, 100 µl protein was injected onto the column and eluted at a flow rate of 0.5 ml/min. 500 µl fractions were collected for analysis by SDS-PAGE (as described in the protein purification section above).

Circular dichroism spectroscopy

Circular dichroism (CD) spectra were obtained using a Chirascan™ CD Spectrometer (Applied Photophysics). Proteins were dialysed into 10 mM sodium phosphate, pH 7.2 using Slide-A-Lyzer™ MINI dialysis devices (Thermo Fisher Scientific). 40 µl protein was used to fill a 0.1 mm cuvette. The sample compartment temperature was 20 °C. Spectra were obtained for wavelengths from 180 nm to 260 nm with step size of 1 nm, bandwidth of 2 nm and time-per-point of 2 s. Ten spectra were acquired for buffer and each sample. The average trace for buffer was subtracted from the average trace for each protein. Units were converted from millidegrees to mean residue molar ellipticity (MRME) using the Chirascan™ software.

Nuclear magnetic resonance titration

Samples of ^{15}N-labelled AVR-Pia (40 µM) alone and complexed to unlabelled OsHPP09-HMA (1:0.5 or 1:1 molar ratio), or OsHPP09-HMA mutants (1:1 molar ratio), were prepared in 20 mM sodium citrate, pH 5.6, 150 mM NaCl, and 1mM DTT, with addition of 10% D_2O for the lock and 1 µM DSS as internal reference for the ^1H dimension and indirectly referenced for the ^{15}N dimension[74]. The backbone (Hn, Nh, Cα) and Cβ resonances of ^{15}N,^{13}C-labeled AVR-Pia complexed to OsHPP09-HMA (1:1 ratio) were assigned using three-dimensional (3D) HNCO, HNCA, HN(CO)CACB, HN(CA)CO and HNCACB experiments recorded at 305K on a Bruker Avance 800 MHz spectrometer equipped with a triple resonance (^1H, ^{15}N, ^{13}C) z-gradient cryo-probe, as described in detail for AVR-Pia alone in solution[13]. Spectra were processed using Topspin (v. 3.5pl6) and analysed with Cindy in-house software or CCPN[75] [analysis v 2.5.2].

720   Chemical shift perturbations ($\Delta\delta$) were calculated using the equation[76]:

$$\Delta\delta = \sqrt{0.5\left[(\Delta\delta_H)^2 + \left(\frac{\Delta\delta_N}{5}\right)^2\right]}$$

722    $\Delta\delta_H$ and $\Delta\delta_N$ are the chemical shift differences measured in the proton dimension and nitrogen
723   dimension, respectively, of 2D [^{1}H,^{15}N] HSQC spectra recorded for AVR-Pia alone or in the
724   presence of OsHPP09-HMA (1:1 ratio). $\Delta\delta$ greater than $\overline{\Delta\delta}$ + 1σ or $\overline{\Delta\delta}$ + 2σ were considered
725   as significant or very significant, respectively, σ being the standard deviation measured on the
726   distribution of chemical-shift perturbations.

Isothermal titration calorimetry (ITC)

ITC experiments were performed using a MicroCal PEAQ-ITC (Malvern Panalytical). Proteins
were dialysed overnight into 20 mM sodium citrate pH 6.0, 150 mM NaCl, 1 mM TCEP.
Experiments were carried out at 25 °C with a reference power of 5 µcal/s. The calorimetric cell
was filled with purified HMA domain at 18 µM (OsHPP09-HMA), 35 µM (OsHIPP21-HMA and
OsHPP11-HMA) or 60 µM (OsHPP10-HMA). This was titrated with AVR-Pia at a concentration
10X greater than that of the HMA domain (180 uM, 350 uM or 600 uM). Each experiment
consisted of a single injection of 0.5 µL followed by 18 injections of 2 µL each at 150 s intervals
with a stirring speed of 750 rpm. Data were processed with the MicroCal PEAQ-ITC analysis
software (Malvern Panalytical). Experiments were carried out in triplicate.

Crystallisation of OsHPP09-HMA and AVR-Pia

Sitting drop, vapour diffusion crystallisation trials were set up in 96-well Swissci 3 lens
crystallisation plates using dragonfly and mosquito dispensing robots (SPT Labtech). 100 nL
of protein at 8.6 mg/ml was combined with 100 nL reservoir solution. Needle-like crystals were
observed in multiple conditions in the commercial Structure Screen 1 + 2 (HT-96; Molecular
Dimensions) and PEGs Suite (NeXtal) screens. Optimisation around these conditions yielded
crystals suitable for diffraction studies. The X-ray diffraction data used to solve the structure
were obtained from a crystal from 0.1 M MES pH 6.5, 0.2 M ammonium sulphate, 20% (w/v)
PEG 4000. The crystal was mounted in a nylon loop and flash-frozen in liquid nitrogen prior to
shipping to the European Synchrotron Radiation Facility (ESRF).

X-ray diffraction data collection and processing, and model refinement

X-ray diffraction data were collected on beamline MASSIF-1 (ID30A-1) at the ESRF. Data were
processed with the EDNA autoprocessing pipeline[77]. The scaled, unmerged data file was
passed to AIMLESS for data reduction[78,79] (implemented in CCP4i2[80]). The AVR-
Pia/OsHPP09-HMA structure was solved by molecular replacement with PHASER[81], using a

monomer of AVR-Pia (PDB accession 6Q76, chain B) and a monomer of OsHPP09-HMA

(modelled with ColabFold[82]). Iterative cycles of manual adjustment, refinement, and validation

were carried out using COOT[83], REFMAC5[84,85] and MolProbity[86,87]. Analysis of the interface

was performed with qtPISA[88]. Interatomic distances were measured and structure

superposition/RMSD calculations were performed with PyMOL v2.5.0 Open-Source[89].

Structure figures were prepared using the CCP4 molecular graphics (CCP4mg) software[90].

Structural modelling of AVR-Pia/HMA complexes

Structural modelling of AVR-Pia/HMA complexes was carried out with ColabFold v1.5.5 [82]

(based on AlphaFold2 Multimer v3[91,92]) and the AlphaFold3 web server (alphafoldserver.com;

models generated between 2024-05-27 and 2024-07-02)[93]. For models generated with

ColabFold, no template information from the PDB was used (template_mode: none) and

number of recycles was set to 20 (num_recycles: auto). Five models were generated for each

complex and relaxed using AMBER (num_relax: 5). Default settings were otherwise used. For

models generated with AlphaFold3, default settings were used. At the time of use, AlphaFold3

did not support excluding template information from the PDB. Five models were generated for

each complex using five different seed values. Interface analysis was carried out with

qtPISA[88].

Acknowledgements

We thank Mark Youles and Mark Banfield for generating and providing the pOPIN-GG vector and tag modules, Laurent Deslandes for providing the split-luciferase vectors, and Florian Veillet for providing the pFV79 construct. We thank Thorsten Langner for providing the Y2H HMA library. We thank Julie Noell for assistance with protein purification, Corinne Lionne and Manon Mallet for helpful discussions about ITC methods, and Christian Roumestand for advice about NMR data. We thank Sebastien Ribeiro, Laura Mathieu and Vincent Chochois for valuable discussions and support with data analysis. This work benefited from access to the NMR and RX facilities of the Integrated Biophysics and Structural Biology Platform (PIBBS) of the CBS. PIBBS is a GIS-IBISA platform and belongs to the French Infrastructure for Integrated Structural Biology (FRISBI), supported by the National Research Agency (ANR-10-INBS-05). We acknowledge the European Synchrotron Radiation Facility (ESRF) for provision of synchrotron radiation facilities. We thank the staff of the ESRF and EMBL Grenoble for assistance in using beamline ID30-A1 under proposal number mx-2409. This work was funded by the HORIZON EUROPE ERC-2019-STG-852482-ii-MAX awarded to Stella Cesari.

Author contributions

Josephine H.R. Maidment: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing. **Svenja C. Saile**: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing. **Aurélien Bocquet**: Investigation; Methodology; Resources; Validation; Writing – review & editing. **Céline Thivolle**: Investigation; Methodology; Resources; Validation; Writing – review & editing. **Léo Bourcet**: Investigation; Methodology; Resources; Validation; Writing – review & editing. **Lisa-Fatimatou Planel**: Investigation; Methodology; Resources; Validation; Writing – review & editing. **Muriel Gelin**: Methodology; Resources; Writing – review & editing. **Thomas Kroj**: Conceptualization; Writing – review & editing. **André Padilla**: Conceptualization; Methodology; Writing – review & editing. **Karine de Guillen**: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing. **Stella Cesari**: Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing.

Competing interests

The authors declare that they have no conflicts of interest with the contents of this article.

Data availability

The protein structure of the complex between OsHPP09-HMA and AVR-Pia has been deposited in the Protein Data Bank with accession number 9RSV (pdb_00009RSV). X-ray diffraction data are publicly available via the European Synchrotron Radiation Facility [94].

Figures

Fig. 1. AVR-Pia interacts with OsH(I)PPs in yeast two-hybrid and in planta.

A Y2H assay of AVR-Pia and MAX58 (both without signal peptides, ΔSP) with either the HMA domain of OsH(I)PPs or full-length (FL) proteins. AVR-Pia/OsRGA5$_{Cter}$ (883-1116 aa) and AVR-Pia/OsHPP09 served as positive controls (upper and lower panels, respectively). Serial dilutions of diploid yeast clones were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW), or to assess protein-protein interactions (SD/-LWH and SD/-LWH supplemented with 0.5 mM 3-amino-1,2,4-triazole (3AT)). Photos were taken after 7 days of incubation. Ø indicates positions where no yeast was spotted. AD, GAL4 activation domain; BD, GAL4 DNA binding domain. **b** Split luciferase complementation assay of AVR-Pia and MAX58 (both ΔSP; fused to the N-terminal part of luciferase; 3xHA:NLuc:MAX) with FL OsH(I)PPs (fused to the C-terminal part of luciferase: 3xFlag:CLuc:OsH(I)PP). Indicated constructs were transiently co-expressed in *N. benthamiana* leaves and leaf discs were harvested 2 days post-infiltration for luminescence measurements. AVR-Pia/OsRGA5$_{Cter}$ (883-1116 aa) was used as a positive control, and leaves infiltrated with P19 alone as background (BG) control. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the

827   interquartile range, with outliers plotted individually. Three independent replicates were

828   performed (n = 8 per replicate) except for AVR-Pia/OsHPP09: n = 16 in replicates 1 and 2.

829   Different letters indicate statistically significant differences based on a pairwise Wilcoxon test

830   (α = 0.05).

Fig. 2. AVR-Pia binds the HMA domains of OsH(I)PPs with varying affinities.

a Representative ITC experiments for each of the AVR-Pia/HMA interactions tested. Top panel shows the raw thermogram obtained from titration of AVR-Pia into a solution containing the purified HMA domains. Central panel shows the integrated heats (coloured dots) and binding isotherm fitted to a single site model (black line) using the MicroCal PEAQ-ITC analysis software (Malvern Panalytical). Bottom panel shows the differences (coloured crosses) between the modelled and observed values (residuals). **b** Dot plot representation of the apparent K_D for each of the AVR-Pia/HMA interactions in three independent experiments. Coloured dots represent the K_D values determined in each replicate with coloured lines representing the error associated with the K_D value as determined by the MicroCal PEAQ-ITC analysis software (Malvern Panalytical).

Fig. 3. AVR-Pia-H3 does not interact with OsHPP09, OsHPP10, OsHPP11 and HIPP21.

a Y2H interaction analysis between AVR-Pia-H3 (ΔSP) and either the HMA domain of OsH(I)PPs or full-length (FL) proteins. OsHPP09/AVR-Pia served as positive control. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH). Photos were taken after 7 days of incubation. AD, GAL4 activation domain; BD, GAL4 DNA binding domain. **b** Split luciferase complementation assay of AVR-Pia-H3 (ΔSP; fused to the N-terminal part of luciferase; 3xHA:NLuc:MAX) with FL OsH(I)PPs (fused to the C-terminal part of luciferase: 3xFlag:CLuc:OsH(I)PP). Indicated constructs were transiently co-expressed in *N. benthamiana* leaves and leaf discs were harvested 2 days post-infiltration for luminescence measurements. AVR-Pia/OsHPP09 served as a positive control, MAX58/OsHPP09 and AVR-Pia-H3/OsRGA5$_{Cter}$ (883-1116 aa) as negative controls, and leaves infiltrated with P19 alone as background (BG) control. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range, with outliers plotted individually. Three independent replicates were performed, with n = 8 per combination per replicate. Due to experimental constraints, for AVR-Pia/OsHPP09, n = 16 in replicates 1 and 2. Data points are shown with shapes indicating the replicate. Different letters indicate statistically significant differences based on a pairwise Wilcoxon test (α = 0.05). **c** Analytical gel filtration traces obtained from injection of OsHPP09-

865   HMA alone (top panel), AVR-Pia-H3 alone (middle panel) and the proteins combined (1:1

866   molar ratio; bottom panel). Significant peaks are indicated by coloured dashed lines with

867   elution volume labelled. SDS-PAGE gel inserts show fractions from peak elution volumes

868   indicated by grey shaded regions. OsHPP09-HMA absorbs light at 280 nm poorly (molar

869   extinction coefficient: 1490 $M^{-1}cm^{-1}$) so the peak corresponding to OsHPP09-HMA is small.

Fig. 4. OsH(I)PPs interacting with AVR-Pia do not interact with AVR1-CO39.

a Y2H interaction analysis between AVR1-CO39 (ΔSP) and either the HMA domain of OsH(I)PPs or full-length (FL) proteins. AVR1-CO39/OsRGA5$_{Cter}$ (883-1116 aa) served as a positive control. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH). Photos were taken after 7 days of incubation. Ø indicates positions where no yeast was spotted. AD, GAL4 activation domain; BD, GAL4 DNA binding domain. **b** Split-luciferase complementation assay of ΔSP AVR1-CO39 with FL OsH(I)PPs and OsRGA5$_{Cter}$ (883-1116 aa). MAX effectors were fused to the N-terminal parts of luciferase (3xHA:NLuc:MAX), while OsH(I)PP FL proteins and OsRGA5$_{Cter}$ were fused to its C-terminal part (3xFlag:CLuc:OsH(I)PP/OsRGA5$_{Cter}$). Constructs were transiently co-expressed in *N. benthamiana* leaves and leaf discs were harvested 2 days post-infiltration for luminescence measurements. AVR-Pia or AVR1-CO39 with OsPi21 served as negative controls and AVR-Pia/OsHPP09 as a positive control. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range, with outliers plotted individually. Three independent replicates were performed, with n = 8 per combination per replicate. All data points are shown with shapes indicating the replicate. Different letters indicate statistically significant differences based on a pairwise Wilcoxon test (α = 0.05). **c** Analytical gel filtration traces obtained from injection of

OsHPP09-HMA alone (top panel), AVR1-CO39 alone (middle panel) and the proteins
combined (1:1 molar ratio; bottom panel). Significant peaks are indicated by coloured dashed
lines with elution volume labelled. SDS-PAGE gel inserts show fractions from peak elution
volumes indicated by grey shaded regions. The small peak at ~13.7 ml for AVR1-CO39 is
likely due to dimerisation of the AVR1-CO39 protein; the quantity of the dimer is insufficient to
observe a band on the SDS-PAGE gel. OsHPP09-HMA absorbs light at 280 nm poorly (molar
extinction coefficient: 1490 $M^{-1}cm^{-1}$) so the peak corresponding to OsHPP09-HMA is small.
898

Fig. 5. OsH(I)PPs interacting with AVR-Pia do not interact with AVR-Pik or Pwl2.

a Y2H interaction analysis between AVR-PikC, AVR-PikD, Pwl2 and AVR-Pia (all ΔSP) and either the HMA domain of OsH(I)PPs or full-length (FL) proteins. OsHIPP19/AVR-PikC, OsHIPP20/AVR-PikC, OsHIPP19/AVR-PikD, OsHIPP20/AVR-PikD and OsHIPP43/Pwl2 served as positive controls. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH). Photos were taken after 7 days of incubation. AD, activating domain; BD, binding domain. **b** Split luciferase complementation assay of MAX effectors (ΔSP; fused to the N-terminal part of luciferase: 3xHA:NLuc:MAX) with OsH(I)PP FL proteins (fused to the C-terminal part of luciferase: 3xFlag:CLuc:OsH(I)PP). AVR-PikC/AVR-PikD with OsHIPP19/20 and Pwl2 with OsHIPP43 served as positive controls, AVR-Pia with OsHIPP43 served as a negative control. Leaves infiltrated with P19 alone served as background (BG) control. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range, with outliers plotted individually Three independent replicates were performed, with n = 8 per combination per replicate, except for

AVR-Pia/OsHIPP19 and AVR-Pia/OsHIPP20 in AVR-PikC and AVR-PikD panels: n = 4 in

replicates 1 and 3. For BG control in AVR-PikC and AVR-PikD panels: n = 3 in replicates 1

and 3 and for BG control in Pwl2 panel: n = 2 in replicates 1 and 3. For AVR-PikD/OsHIPP19

n = 16 in replicates 1 and 3 and n = 24 in replicate 2. Data points are shown with shapes

indicating the replicate. Different letters indicate statistically significant differences based on a

pairwise Wilcoxon test (α = 0.05).

Fig. 6. Crystal structure of the HMA domain of OsHPP09 in complex with AVR-Pia.

The structures of OsHPP09-HMA and AVR-Pia are represented as teal and gold ribbons,
respectively. The amino acids comprising β2 of AVR-Pia and β2 of the HMA domain are
represented as cylinders. Side chains involved in intermolecular hydrogen bonds/salt bridges
are also represented as cylinders. Hydrogen bonds are represented by black dashed lines
labelled with the bond length. Left insert shows main chain hydrogen bonds between AVR-
Pia-β2 and OsHPP09-β2. Right insert shows intermolecular contacts between side chains of
residues in OsHPP09-α1 and AVR-Pia.

Fig. 7. Mutation of OsHPP09 E16 reduces AVR-Pia binding.

a Y2H interaction analysis between AVR-Pia (ΔSP) and full-length (FL) OsHPP09 carrying single point mutations. OsHPP09[WT]/AVR-Pia served as positive control. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH and SD/-LWH supplemented with 5 mM 3AT). Photos were taken after 7 days of incubation. AD, activating domain; BD, binding domain. **b** Split luciferase complementation assays of AVR-Pia (ΔSP; fused to the N-terminal part of luciferase; 3xHA:NLuc:AVR-Pia) with FL OsHPP09 carrying indicated mutations (fused to the C-terminal part of luciferase: 3xFlag:CLuc:OsHPP09). Co-expression of AVR-Pia with OsHPP09 served as a positive control, co-expression of OsHIPP43 with AVR-Pia as a negative control. Leaves infiltrated with P19 alone served as background (BG) control. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile rang, with outliers plotted individually. Three independent replicates were performed, with n = 8 per combination per replicate. Data points are shown with shapes corresponding to the replicate. Different letters

indicate statistically significant differences based on a pairwise Wilcoxon test (α = 0.05). **c**

Circular dichroism (CD) spectra obtained for WT OsHPP09-HMA (pale blue) and variants

carrying E16A, E16R, I35K or D33H mutations. Negative peaks at 222 nm/208 nm and a

positive peak at 193 nm indicate α-helices while a negative peak at 218 nm and positive peak

at 195 nm indicate β-sheets, consistent with the βαββαβ HMA domain topology. Low ellipticity

above 210 nm and a negative peak at 200 nm indicates disorder. **d** Overlays of cross-peaks

from HSQC spectra corresponding to residues G30, I44, E56 and Y85 of [15]N-labelled AVR-

Pia alone (orange) and in the presence of either WT OsHPP09-HMA or variants carrying

E16A, E16R or D33H mutations (blue/green/purple). Overlays of the full spectra are provided

in **Supplementary Figures 21-23**.

References

1.   Baudin, M. *et al.* Pyricularia oryzae: Lab star and field scourge. *Mol Plant Pathol* **25**,
(2024).

2.   Savary, S. *et al.* The global burden of pathogens and pests on major food crops. *Nat*
Ecol Evol **3**, 430–439 (2019).

3.   Oliveira-Garcia, E., Yan, X., Oses-Ruiz, M., de Paula, S. & Talbot, N. J. Effector-
triggered susceptibility by the rice blast fungus Magnaporthe oryzae. *New Phytologist*
241, 1007–1020 (2024).

4.   Le Naour--Vernet, M. *et al.* Structure-guided insights into the biology of fungal effectors.
New Phytologist **246**, 1460–1477 (2025).

5.   de Guillen, K. *et al.* Structure analysis uncovers a highly diverse but structurally
conserved effector family in phytopathogenic fungi. *PLoS Pathog* **11**, (2015).

6.   Lahfa, M. *et al.* Does a Similar 3D Structure Mean a Similar Folding Pathway? The
Presence of a C-Terminal α-Helical Extension in the 3D Structure of MAX60 Drastically
Changes the Folding Pathway Described for Other MAX-Effectors from Magnaporthe
oryzae. *Molecules* **28**, 6068 (2023).

7.   Lahfa, M. *et al.* The structural landscape and diversity of Pyricularia oryzae MAX
effectors revisited. *PLoS Pathog* **20**, e1012176 (2024).

8.   Le Naour--Vernet, M. *et al.* Adaptive evolution in virulence effectors of the rice blast
fungus Pyricularia oryzae. *PLoS Pathog* **19**, e1011294 (2023).

9.   Maidment, J. H. R. *et al.* Multiple variants of the fungal effector AVR-Pik bind the HMA
domain of the rice protein OsHIPP19, providing a foundation to engineer plant defense.
Journal of Biological Chemistry **296**, (2021).

10.   Zdrzałek, R. *et al.* Bioengineering a plant NLR immune receptor with a robust binding
interface toward a conserved fungal pathogen effector. *Proc Natl Acad Sci U S A* **121**,
e2402872121 (2024).

11.   Liao, J. *et al.* Pathogen effectors and plant immunity determine specialization of the
blast fungus to rice subspecies. *Elife* **5**, (2016).

12.   Cesari, S. *et al.* The rice resistance protein pair RGA4/RGA5 recognizes the
Magnaporthe oryzae effectors AVR-Pia and AVR1-CO39 by direct binding. *Plant Cell*
25, 1463–1481 (2013).

13.   Guo, L. *et al.* Specific recognition of two MAX effectors by integrated HMA domains in
plant immune receptors involves distinct binding surfaces. *Proc Natl Acad Sci U S A*
115, 11637–11642 (2018).

14.   Ortiz, D. *et al.* Recognition of the Magnaporthe oryzae effector AVR-Pia by the decoy
domain of the rice NLR immune receptor RGA5. *Plant Cell* **29**, 156–168 (2017).

15.   Cesari, S. Multiple strategies for pathogen perception by plant immune receptors. *New*
Phytologist **219**, 17–24 (2018).

16.   Cesari, S., Bernoux, M., Moncuquet, P., Kroj, T. & Dodds, P. N. A novel conserved
mechanism for plant NLR protein pairs: The 'integrated decoy' hypothesis. *Front Plant*
Sci **5**, (2014).

17.   Kroj, T., Chanclud, E., Michel-Romiti, C., Grand, X. & Morel, J. B. Integration of decoy
domains derived from protein targets of pathogen effectors into plant immune receptors
is widespread. *New Phytologist* **210**, 618–626 (2016).

18.   Sarris, P. F., Cevik, V., Dagdas, G., Jones, J. D. G. & Krasileva, K. V. Comparative
analysis of plant immune receptor architectures uncovers host proteins likely targeted
by pathogens. *BMC Biol* **14**, 1–19 (2016).

19.   Bailey, P. C. *et al.* Dominant integration locus drives continuous diversification of plant
immune receptors with exogenous domain fusions. *Genome Biol* **19**, 1–18 (2018).

20.   Oikawa, K. *et al.* The blast pathogen effector AVR-Pik binds and stabilizes rice heavy
metal-associated (HMA) proteins to co-opt their function in immunity. *PLoS Pathog* **20**,
e1012647 (2024).

21.   De Abreu-Neto, J. B., Turchetto-Zolet, A. C., De Oliveira, L. F. V., Bodanese Zanettini,
M. H. & Margis-Pinheiro, M. Heavy metal-associated isoprenylated plant protein (HIPP):
characterization of a family of proteins exclusive to plants. *FEBS J* **280**, 1604–1616
(2013).

22.   Tehseen, M., Cairns, N., Sherson, S. & Cobbett, C. S. Metallochaperone-like genes in
Arabidopsis thaliana. *Metallomics* **2**, 556–564 (2010).

23. Crowell, D. N. & Huizinga, D. H. Protein isoprenylation: the fat of the matter. *Trends*
Plant Sci **14**, 163–170 (2009).

24. Kay, B. K., Williamson, M. P. & Sudol, M. The importance of being proline: the
interaction of proline-rich motifs in signaling proteins with their cognate domains. *The*
FASEB Journal **14**, 231–241 (2000).

25. Dykema, P. E. *et al.* A new class of proteins capable of binding transition metals. *Plant*
Mol Biol **41**, 139–150 (1999).

26. Suzuki, N., Yamaguchi, Y., Koizumi, N. & Sano, H. Functional characterization of a
heavy metal binding protein Cdl19 from Arabidopsis. *Plant Journal* **32**, 165–173 (2002).

27. Khan, I. U. *et al.* Identification of novel rice (Oryza sativa) HPP and HIPP genes tolerant
to heavy metal toxicity. *Ecotoxicol Environ Saf* **175**, 8–18 (2019).

28. Li, J. *et al.* Heavy metal stress-associated proteins in rice and Arabidopsis: genome-
wide identification, phylogenetics, duplication, and expression profiles analysis. *Front*
Genet **11**, (2020).

29. Barr, Z. K., Werner, T. & Tilsner, J. Heavy Metal-Associated Isoprenylated Plant
Proteins (HIPPs) at plasmodesmata: Exploring the link between localization and
function. *Plants* **12**, (2023).

30. Barth, O., Vogt, S., Uhlemann, R., Zschiesche, W. & Humbeck, K. Stress induced and
nuclear localized HIPP26 from Arabidopsis thaliana interacts via its heavy metal
associated domain with the drought stress related zinc finger transcription factor
ATHB29. *Plant Mol Biol* **69**, 213–226 (2009).

31. Barth, O., Zschiesche, W., Siersleben, S. & Humbeck, K. Isolation of a novel barley
cDNA encoding a nuclear protein involved in stress response and leaf senescence.
Physiol Plant **121**, 282–293 (2004).

32. Cowan, G. H. *et al.* Potato mop-top virus co-opts the stress sensor HIPP26 for long-
distance movement. *Plant Physiol* **176**, 2052–2070 (2018).

33. Radakovic, Z. S. *et al.* Arabidopsis HIPP27 is a host susceptibility gene for the beet
cyst nematode Heterodera schachtii. *Mol Plant Pathol* **19**, 1917–1928 (2018).

34. Fukuoka, S. *et al.* Loss of function of a proline-containing protein confers durable
disease resistance in Rice. *Science (1979)* **325**, 998–1001 (2009).

35. Were, V. M. *et al.* The Magnaporthe oryzae effector Pwl2 alters HIPP43 localization to
suppress host immunity. *Plant Cell* (2025).

36. Huang, Z. *et al.* Small secreted effector protein from Fusarium sacchari suppresses
host immune response by inhibiting ScPi21-induced cell death. *Mol Plant Pathol* **25**,
e13414 (2024).

37. Song, H. *et al.* The Meloidogyne graminicola effector MgMO289 targets a novel copper
metallochaperone to suppress immunity in rice. *J Exp Bot* **72**, 5638–5655 (2021).

38. Huang, Q. *et al.* CRISPR/Cas9-mediated mutagenesis of the susceptibility gene
OsHPP04 in rice confers enhanced resistance to rice root-knot nematode. *Front Plant*
Sci **14**, (2023).

39. Niu, X. *et al.* A small secreted protein, RsMf8HN, in Rhizoctonia solani triggers plant
immune response, which interacts with rice OsHIPP28. *Microbiol Res* **266**, 127219
(2023).

40. Zhang, X. *et al.* Isolation and characterisation of cDNA encoding a wheat heavy metal-
associated isoprenylated protein involved in stress responses. *Plant Biol* **17**, 1176–
1186 (2015).

41. Huang, G. *et al.* Genome-wide characterization of heavy metal-associated
isoprenylated plant protein gene family from Citrus sinensis in response to
huanglongbing. *Front Plant Sci* **15**, (2024).

42. Imran, Q. M. *et al.* Nitric oxide responsive heavy metal-associated gene AtHMAD1
contributes to development and disease resistance in Arabidopsis thaliana. *Front Plant*
Sci **7**, 232346 (2016).

43. Bentham, A. R. *et al.* A single amino acid polymorphism in a conserved effector of the
multihost blast fungus pathogen expands host-target binding spectrum. *PLoS Pathog*
17, e1009957- (2021).

44. Ashikawa, I. *et al.* Two adjacent nucleotide-binding site-leucine-rich repeat class genes
are required to confer Pikm-specific rice blast resistance. *Genetics* **180**, 2267–2276
(2008).

45.   Zdrzałek, R., Kamoun, S., Terauchi, R., Saitoh, H. & Banfield, M. J. The rice NLR pair
1076   Pikp-1/Pikp-2 initiates cell death through receptor cooperation rather than negative
1077   regulation. *PLoS One* **15**, (2020).

1078   46.   Cesari, S. *et al.* New recognition specificity in a plant immune receptor by molecular
1079   engineering of its integrated domain. *Nat Commun* **13**, (2022).

1080   47.   Varden, F. A. *et al.* Cross-reactivity of a rice NLR immune receptor to distinct effectors
1081   from the rice blast pathogen Magnaporthe oryzae provides partial disease resistance.
1082   *J Biol Chem* **294**, 13006–13016 (2019).

1083   48.   Langner, T., Kellner, R., Harant, A., Terauchi, R. & Kamoun, S. A clone resource of
1084   cereal heavy-metal associated (HMA) domains— putative targets of plant pathogen
1085   effector proteins. Preprint at https://doi.org/10.5281/zenodo.5148558 (2021).

1086   49.   Maqbool, A. *et al.* Structural basis of pathogen recognition by an integrated HMA
1087   domain in a plant NLR immune receptor. *Elife* **4**, e08709 (2015).

1088   50.   Gómez De La Cruz, D., Zdrzałek, R., Banfield, M. J., Talbot, N. J. & Moscou, M. J.
1089   Molecular mimicry of a pathogen virulence target by a plant immune receptor. *bioRxiv*
1090   Preprint at https://doi.org/10.1101/2024.07.26.605320 (2024).

1091   51.   Homma, F., Huang, J. & van der Hoorn, R. A. L. AlphaFold-Multimer predicts cross-
1092   kingdom interactions at the plant-pathogen interface. *Nat Commun* **14**, 6040 (2023).

1093   52.   Ose, T. *et al.* Solution structure of an avirulence protein, AVR-Pia, from Magnaporthe
1094   oryzae. *J Biomol NMR* **63**, 229–235 (2015).

1095   53.   Schymkowitz, J. *et al.* The FoldX web server: an online force field. *Nucleic Acids Res*
1096   **33**, W382–W388 (2005).

1097   54.   Park, C. H. *et al.* The Magnaporthe oryzae effector AvrPiz-t targets the RING E3
1098   ubiquitin ligase APIP6 to suppress pathogen-associated molecular pattern–triggered
1099   immunity in rice. *Plant Cell* **24**, 4748–4762 (2012).

1100   55.   Park, C. H. *et al.* The E3 ligase APIP10 connects the effector AvrPiz-t to the NLR
1101   receptor Piz-t in rice. *PLoS Pathog* **12**, (2016).

1102   56.   Couch, B. C. *et al.* Origins of host-specific populations of the blast pathogen
1103   Magnaporthe oryzae in crop domestication with subsequent expansion of pandemic
1104   clones on rice and weeds of rice. *Genetics* **170**, 613–630 (2005).

57.   Tosa, Y. *et al.* Evolution of an avirulence gene, AVR1-CO39, concomitant with the
evolution and differentiation of Magnaporthe oryzae. *Molecular Plant-Microbe*
Interactions **18**, 1148–1160 (2005).

58.   Hála, M., Žárský, V., Swiezewska, E. & Surmacz, L. Protein prenylation in plant stress
responses. *Molecules* **24**, 3906 (2019).

59.   Nawaz, G. *et al.* Knockout of Pi21 by CRISPR/Cas9 and iTRAQ-based proteomic
analysis of mutants revealed new insights into M. oryzae resistance in elite rice line.
Genes (Basel) **11**, 1–24 (2020).

60.   Tao, H. *et al.* Engineering broad-spectrum disease-resistant rice by editing multiple
susceptibility genes. *J Integr Plant Biol* **63**, 1639–1648 (2021).

61.   Zhou, Y. *et al.* Engineering of rice varieties with enhanced resistances to both blast and
bacterial blight diseases via CRISPR/Cas9. *Plant Biotechnol J* **20**, 876–885 (2022).

62.   Hanika, K. *et al.* Impairment of tomato WAT1 enhances resistance to vascular wilt fungi
despite severe growth defects. *Front Plant Sci* **12**, 721674 (2021).

63.   Kieu, N. P., Lenman, M., Wang, E. S., Petersen, B. L. & Andreasson, E. Mutations
introduced in susceptibility genes through CRISPR/Cas9 genome editing confer
increased late blight resistance in potatoes. *Sci Rep* **11**, 1–12 (2021).

64.   Wolter, M., Hollricher, K., Salamini, F. & Schulze-Lefert, P. The mlo resistance alleles
to powdery mildew infection in barley trigger a developmentally controlled defence
mimic phenotype. *MGG Molecular & General Genetics* **239**, 122–128 (1993).

65.   Kusch, S. *et al.* Arabidopsis mlo3 mutant plants exhibit spontaneous callose deposition
and signs of early leaf senescence. *Plant Mol Biol* **101**, 21–40 (2019).

66.   Garcia-Ruiz, H., Szurek, B. & Van den Ackerveken, G. Stop helping pathogens:
engineering plant susceptibility genes for durable resistance. *Curr Opin Biotechnol* **70**,
187–195 (2021).

67.   Weber, E., Engler, C., Gruetzner, R., Werner, S. & Marillonnet, S. A modular cloning
system for standardized assembly of multigene constructs. *PLoS One* **6**, e16765
(2011).

68.   Engler, C., Kandzia, R. & Marillonnet, S. A one pot, one step, precision cloning method
with high throughput capability. *PLoS One* **3**, (2008).

69.   Bentham, A. R. *et al.* pOPIN-GG: A resource for modular assembly in protein
expression vectors. *bioRxiv* Preprint at https://doi.org/10.1101/2021.08.10.455798
(2021).

70.   Bernoux, M. *et al.* Structural and functional analysis of a plant resistance protein TIR
domain reveals interfaces for self-association, signaling, and autoregulation. *Cell Host*
Microbe **9**, 200–211 (2011).

71.   Matsumoto, T. *et al.* The map-based sequence of the rice genome. *Nature* **436**, (2005).

72.   Kawahara, Y. *et al.* Improvement of the Oryza sativa nipponbare reference genome
using next generation sequence and optical map data. *Rice* **6**, (2013).

73.   Kushnirov, V. V. Rapid and reliable protein extraction from yeast. *Yeast* **16**, 857–860
(2000).

74.   Wishart, D. S. *et al.* 1H, 13C and 15N chemical shift referencing in biomolecular NMR.
J Biomol NMR **6**, 135–140 (1995).

75.   Vranken, W. F. *et al.* The CCPN data model for NMR spectroscopy: Development of a
software pipeline. *Proteins: Structure, Function and Genetics* **59**, 687–696 (2005).

76.   Grzesiek, S. *et al.* The solution structure of HIV-1 Nef reveals an unexpected fold and
permits delineation of the binding surface for the SH3 domain of Hck tyrosine protein
kinase. *Nat Struct Biol* **3**, 340–345 (1996).

77.   Incardona, M. F. *et al.* EDNA: A framework for plugin-based applications applied to X-
ray experiment online data analysis. *J Synchrotron Radiat* **16**, 872–879 (2009).

78.   Evans, P. R. An introduction to data reduction: space-group determination, scaling and
intensity statistics. *Acta Crystallogr D Biol Crystallogr* **67**, 282 (2011).

79.   Evans, P. R. & Murshudov, G. N. How good are my data and what is the resolution?
Acta Crystallogr D Biol Crystallogr **69**, 1204–1214 (2013).

80.   Agirre, J. *et al.* The CCP4 suite: integrative software for macromolecular
crystallography. *Acta Crystallogr D Struct Biol* **79**, 449–461 (2023).

81.   McCoy, A. J. *et al.* Phaser crystallographic software. *J Appl Crystallogr* **40**, 658–674
(2007).

82. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679–682 (2022).

83. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486–501 (2010).

84. Murshudov, G. N., Vagin, A. A. & Dodson, E. J. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr D Biol Crystallogr* **53**, 240–255 (1997).

85. Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal structures. *Acta Crystallogr D Biol Crystallogr* **67**, 355–367 (2011).

86. Davis, I. W., Murray, L. W., Richardson, J. S. & Richardson, D. C. MolProbity: Structure validation and all-atom contact analysis for nucleic acids and their complexes. *Nucleic Acids Res* **32**, (2004).

87. Chen, V. B. *et al.* MolProbity: All-atom structure validation for macromolecular crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12–21 (2010).

88. Krissinel, E. & Henrick, K. Inference of macromolecular assemblies from crystalline state. *J Mol Biol* **372**, 774–797 (2007).

89. Schrödinger LLC. *The PyMOL Molecular Graphics System, Version 2.5.0 Open-Source*. (2015).

90. McNicholas, S., Potterton, E., Wilson, K. S. & Noble, M. E. M. Presenting your structures: the CCP4mg molecular-graphics software. *Acta Crystallogr D Biol Crystallogr* **67**, 386–394 (2011).

91. Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. *bioRxiv* Preprint at https://doi.org/10.1101/2021.10.04.463034 (2022).

92. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

93. Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with AlphaFold 3. *Nature* **630**, 493–500 (2024).

94. Maidment, J. H. R., Bocquet, A., Gelin, M., De Guillen, K. & Cesari, S. BAG Montpellier [Dataset]. (2025) doi:10.15151/ESRF-DC-2199982133.

1192