The *Magnaporthe oryzae* MAX effector AVR-Pia binds a novel group of rice HMA domain-containing proteins

Josephine H.R. Maidment[1,2]*, Svenja C. Saile[1]*, Aurélien Bocquet[2], Céline Thivolle[1], Léo Bourcet[2], Lisa-Fatimatou Planel[1], Muriel Gelin[2], Thomas Kroj[1], André Padilla[2], Karine de Guillen[2]**, Stella Cesari[1]**

[1] PHIM Plant Health Institute, Univ Montpellier, INRAE, CIRAD, Institut Agro, IRD, Montpellier, France.
[2] Centre de Biologie Structurale, INSERM, CNRS, Université de Montpellier, 29 rue de Navacelles, 34090, Montpellier, France

*Contributed equally
**Corresponding authors

Stella Cesari

Tel: +33 4 99 62 48 62

Email: stella.cesari@inrae.fr

Karine de Guillen

Tel: +33 4 67 41 77 08

Email: karine.deguillen@cbs.cnrs.fr

ORCID

Josephine H.R. Maidment: 0000-0002-8229-2718
Svenja C. Saile: 0000-0003-2258-3166
Aurélien Bocquet: 0009-0007-0082-6098
Céline Thivolle: 0009-0000-5048-3619
Léo Bourcet: 0009-0004-4568-3543
Lisa-Fatimatou Planel: 0009-0007-5159-7109
Muriel Gelin 0000-0003-1320-8663
Thomas Kroj: 0000-0002-3752-1788
André Padilla: 0000-0001-9702-4612
Karine de Guillen: 0000-0002-5037-2115
Stella Cesari: 0000-0001-8558-0371

Keywords: pathogen effector, host virulence targets, rice blast fungus, rice, protein structure, plant defence, heavy metal associated domain, MAX effector

Abstract

Phytopathogenic fungi secrete effector proteins to promote virulence. MAX (*Magnaporthe*

Avrs and ToxB-like) effectors form a sequence-diverse family sharing a conserved protein

structure. AVR-Pia, a MAX effector from the rice blast fungus *Magnaporthe oryzae,* is

recognised by the paired rice nucleotide-binding leucine-rich repeat immune receptors

OsRGA4/OsRGA5 through direct binding to an integrated heavy metal-associated (HMA)

domain in OsRGA5. Here, we identify previously unknown host targets of AVR-Pia: four HMA

domain-containing rice proteins, belonging to the HMA Isoprenylated Plant Protein (HIPP) and

HMA Plant Protein (HPP) families. AVR-Pia interacts with all four proteins both in vitro and in

planta and binds their HMA domains with varying affinities. The crystal structure of AVR-Pia

bound to the HMA domain of OsHPP09 revealed the molecular details of the binding interface.

Structure-guided mutagenesis of OsHPP09 identified a single point mutation which prevents

AVR-Pia binding, providing a foundation for targeted engineering of HMA domains to evade

effector binding.

Introduction

The fungal pathogen *Magnaporthe oryzae* (syn. *Pyricularia oryzae*) is the causative agent of rice blast, one of the most devastating diseases affecting cultivated rice worldwide [1,2]. Beyond rice, this pathogen also causes severe yield losses in other cereals, including millets, barley and wheat, and poses a threat to global food security [3].

To successfully colonize host plants, *M. oryzae* has evolved mechanisms to evade plant defense responses and manipulate host cellular pathways to its advantage. A key element of its pathogenic strategy is the secretion of a large collection of effector proteins. A subset of these belong to the Magnaporthe AVRs and ToxB-like (MAX) effector family which has undergone massive expansion in *M. oryzae* [4]. A characteristic feature of MAX effectors is their conserved structural fold consisting of two β-sheets, each composed of three antiparallel β-strands, stabilised by one or more disulphide bonds. Despite this structural conservation, MAX effectors lack sequence similarity, resulting in differences in surface properties. Some members also carry variable N- and C-terminal extensions, which may enable them to target a broad range of host proteins [5–7].

AVR-Pia is a MAX effector present in approximately 20% of the *M. oryzae* isolates that infect rice. It is highly expressed during the biotrophic phase of host colonization and contributes to fungal virulence on rice [4,6,8]. In certain rice varieties, AVR-Pia is directly recognised by the intracellular immune receptor OsRGA5, a nucleotide-binding leucine-rich repeat (NLR) protein that functions together with the helper NLR OsRGA4 to activate immune responses and restrict the spread of *M. oryzae* isolates expressing AVR-Pia [9–11]. Canonical plant NLRs consist of an N-terminal signaling domain, a central nucleotide-binding domain and a C-terminal leucine-rich repeat domain [12]. However, many NLRs also carry non-canonical integrated domains (IDs), which are thought to originate from host proteins targeted by pathogen effectors [13]. These IDs, found across diverse plant species, are believed to function as mimics or baits for effector recognition ("integrated decoy" model) [13–16]. Both AVR-Pia and a second MAX effector, AVR1-CO39, bind to the HMA ID of OsRGA5, despite sharing limited sequence similarity [9,10]. This suggests that these effectors may target host proteins containing HMA domains as part of their virulence strategy. However, to date, no host targets of AVR-Pia or AVR1-CO39 have been identified, and their specific virulence functions remain unknown.

Sequence analyses have shown that the OsRGA5-HMA domain shares sequence similarity with the HMA domains of two related families of HMA domain-containing proteins [17]: HMA

domain-containing Plant Proteins (HPPs) and HMA domain-containing Isoprenylated Plant Proteins (HIPPs). These two families are collectively referred to hereafter as H(I)PPs. H(I)PPs are small metallochaperone-like proteins characterized by the presence of one or two HMA domains, each adopting an α/β sandwich structure with a βαββαβ topology. Typically, the HMA domain features a metal-binding MxCxxC motif, implicating H(I)PPs in roles related to metal homeostasis [18,19]. However, approximately a third of rice H(I)PPs possess degenerate MxCxxC motifs, suggesting functional diversification beyond metal binding.

Unlike HPPs, HIPPs carry a C-terminal CaaX motif that undergoes isoprenylation [20,21], a post-translational modification involving the covalent attachment of an isoprenoid lipid to the cysteine residue in the motif. This modification promotes membrane association of proteins [22]. Additionally, many HIPPs harbour proline-rich regions between the HMA domain and the CaaX motif, which may be important in mediating protein-protein interactions [20,23].

Notably, H(I)PPs have undergone extensive expansion in vascular plants, with approximately 100 members identified in rice [17,20,24,25]. Most remain functionally uncharacterized, although several have been implicated in response to various biotic and abiotic stresses [20,26–28]. Increasing evidence points to a broader role of H(I)PPs as susceptibility factors in interactions with pathogens, pests, and viruses [17,29–39].

In cereals, H(I)PPs have been identified as susceptibility factors for *M. oryzae* infection [17,31,32]. Notably, some of these H(I)PPs are targeted by the MAX effectors AVR-Pik and APikL2[17,40,41] AVR-Pik specifically interacts with members of a defined phylogenetic subclade (referred to as clade A) of OsH(I)PPs, including among others OsHIPP19, OsHIPP20, OsHPP03, and OsHPP04 [17,41]. OsHIPP20 has been functionally validated as a susceptibility gene, as its loss-of-function mutation results in reduced sensitivity to *M. oryzae* [17].

Similar to AVR-Pia and AVR1-CO39, recognition of AVR-Pik in rice involves direct binding of the effector to an HMA ID in the NLR receptor OsPik-1, which operates together with another NLR protein, OsPik-2, to confer resistance [42,43]. The HMA domain of OsPik-1 belongs to the same phylogenetic clade as the H(I)PPs targeted by AVR-Pik, reinforcing the idea that effector targets may serve as templates for domain integration into NLRs.

Structural analyses of complexes between the MAX effectors AVR-PikF or APikL2 and the HMA domains of their respective H(I)PP targets have revealed a conserved binding interface [40,41]. In both cases, crystal structures show that β3 of the MAX fold and β4 of the HMA domain are aligned in an antiparallel orientation, resulting in the formation of a β-sheet comprising the four β-strands of the HMA domain and β-strands 3, 4 and 5 of the MAX fold. The interface is

globally similar to that observed in the crystal structures of AVR-Pik and the HMA ID of OsPik-

1, although the interface between AVR-PikF and OsHIPP19-HMA extends across a larger

area and involves additional hydrogen bonds [41].

By contrast, the crystal structure of AVR1-CO39 bound to the HMA ID of OsRGA5 revealed a

markedly different interface. Here, β2 of the core MAX fold of AVR1-CO39 and β2 of OsRGA5-

HMA are aligned, resulting in a β-sheet comprising the four β-strands of the HMA domain, and

β2, β1 and β6 of the MAX fold [10]. NMR titration experiments were used to map the OsRGA5-

interacting surface of AVR-Pia, and the results, combined with structural modelling and

mutagenesis, indicated that both effectors bind to OsRGA5-HMA at a similar interface [10,11,44].

Additionally, a crystal structure of AVR-Pia in complex with the HMA ID of OsPikp-1 revealed

a comparable binding interface, indicating that AVR-Pia binds different HMA domains in a

structurally similar manner [45].

More recently, the crystal structure of the MAX effector Pwl2 bound to OsHIPP43 showed that,

similar to the interface between AVR1-CO39 and AVR-Pia with OsRGA5-HMA, β2 of the core

MAX fold of Pwl2 and β2 of OsHIPP43-HMA are aligned [46]. However, unlike AVR-Pia and

AVR1-CO39, in addition to its core MAX fold, Pwl2 has a C-terminal extension comprising an

α-helix and unstructured region. This extension contributes significantly to the HMA binding

interface [46].

Collectively, these structural studies underscore the remarkable diversity with which MAX

effectors have evolved to target various HMA domain-containing proteins. They support the

hypothesis that AVR-Pia and AVR1-CO39 may engage distinct H(I)PP targets, different from

those bound by other MAX effectors, through unique interfaces.

MAX effectors constitute the largest arsenal of molecular weapons in *M. oryzae*. Identifying

their virulence targets, particularly for effectors that contribute directly to pathogenicity such

as AVR-Pia, and elucidating the molecular and structural basis of these interactions is of

critical importance. This knowledge will be instrumental in revealing the mechanisms exploited

by *M. oryzae* during the early stages of host infection. Furthermore, the identification of new

H(I)PP targets of *M. oryzae* effectors may provide opportunities for engineering blast

resistance, and possibly resistance to other pathogens that exploit H(I)PPs as susceptibility

factors. Such strategies could include bioengineering approaches aimed at disrupting effector-

target interactions in crop species. Finally, uncovering the targets of AVR-Pia would also bring

valuable evolutionary insights into the origin of its recognition by the NLR-ID receptor

OsRGA5.

Here, we identified four OsH(I)PPs as specific interactors of AVR-Pia. Our analyses revealed

that none of them are bound by other MAX effectors, including AVR1-CO39, AVR-Pik and

Pwl2. This finding suggests that although these MAX effectors commonly bind to HMA

domains, they have evolved to recognise different host targets. Using biochemical and

biophysical approaches, we validated and further characterized the interaction between AVR-

Pia and one of the interactors, OsHPP09. The crystal structure of AVR-Pia/OsHPP09-HMA

complex revealed extensive intermolecular hydrogen bonding and shows overall structural

similarity to the AVR1-CO39/OsRGA5-HMA complex. Guided by the AVR-Pia/OsHPP09-HMA

crystal structure, we identified a point mutation in OsHPP09 which disrupts its interaction with

AVR-Pia without compromising the HMA fold.

Results

AVR-Pia interacts with rice H(I)PPs in Y2H screening

As AVR-Pia is recognised by OsRGA5 through direct binding to its HMA ID [9,11], we hypothesized that AVR-Pia may target HMA domain-containing proteins to promote host susceptibility.

To identify potential H(I)PP host targets of AVR-Pia, we conducted extensive pairwise yeast two-hybrid (Y2H) assays. We screened a signal peptide-deleted version of AVR-Pia (dSP-AVR-Pia, 20 – 85 aa) fused to the Gal4 activation domain (AD) against a rice HMA domain library [47] comprising 80 rice HMA domains fused to the Gal4 DNA binding domain (BD). Through this analysis, we identified five HMA domains that interact with AVR-Pia: OsHPP09 (LOC_Os03g02070), OsHIPP41 (LOC_Os03g06080), OsHPP11 (LOC_Os04g45130), OsHIPP14 (LOC_Os04g39320) and OsHIPP39 (LOC_Os03g02860) (**Fig. 1a, Supplementary Fig. 1, Supplementary Table 1**). Among these, BD:OsHPP09 paired with AD:AVR-Pia conferred strong yeast growth on interaction-selective media, indicating a robust interaction. In contrast, the remaining candidates led to only weak yeast growth when co-expressed with AD:AVR-Pia.

We additionally tested the rice HMA library against the well-characterized MAX effectors Pwl2 and AVR-PikD. Consistent with earlier publications, we observed dSP-Pwl2 (22 – 145 aa) interacting with the HMA domain of OsHIPP43, and dSP-AVR-PikD (22 – 113 aa) interacting with the HMA domains of OsHIPP19, OsHIPP20, and other phylogenetically related HMA domains [17,32,41,46] (**Supplementary Fig. 1**), confirming the integrity and sensitivity of our screen.

In addition to this targeted pairwise analysis, an independent Y2H screen using a rice cDNA library identified another HMA domain-containing protein, OsHIPP21 (LOC_Os09g09830), as a putative interactor of AVR-Pia (**Supplementary Table 1, Supplementary Table 2**). Together, these findings suggest that AVR-Pia might target a specific set of OsH(I)PPs.

We next examined whether the corresponding full-length (FL) proteins of all identified putative interactors also associate directly with AVR-Pia, and tested them along with their isolated HMA domains in a pairwise Y2H assay. Interestingly, interaction between AVR-Pia and OsH(I)PP FL proteins was retained for only a subset of proteins: OsHPP09, OsHPP11 and OsHIPP21 (**Fig. 1a and Supplementary Fig. 2**), even though immunoblot analyses confirmed the expression of all tested proteins (**Supplementary Fig. 3a, 3b, 3c**). Given that FL HPPs and HIPPs differ by the presence of a C-terminal isoprenylation motif in HIPPs, and since isoprenylation is known to promote membrane association of proteins [22], we aimed to rule out

Fig. 1. AVR-Pia interacts with OsH(I)PPs.

a Y2H interaction analysis between AVR-Pia (upper panel) or MAX58 (lower panel) and either the heavy metal-associated (HMA) domain or full-length (FL) proteins of OsH(I)PPs. The OsH(I)PP candidates were previously identified in a comprehensive pairwise Y2H analysis using AVR-Pia as prey and a library of HMA domains from OsH(I)PPs as bait. OsRGA5$_{Cter}$ (883 – 1116 aa) was included as a positive control in the experiment with AVR-Pia/OsH(I)PPs (top panel). After an interaction was observed between OsHPP09 and AVR-Pia, this interaction was used as a positive control in the subsequent experiment with MAX58 (bottom panel).. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH and SD/-LWH supplemented with 0.5 mM 3AT). Photos were taken after 7 days of incubation. Ø indicates positions where no yeast was spotted. AD, activating domain; BD, binding domain. **b** Quantitative measurement of luciferase-derived signals shows that AVR-Pia, but not MAX58, interacts with OsHPP09, OsHPP10, OsHPP11 and OsHIPP21 in a split-luciferase complementation assay. Neither AVR-Pia nor MAX58 interact with any other tested OsHIPP in planta. AVR-Pia and MAX58 were fused to the N-terminal part of luciferase (3xHA:NLuc:AVR-Pia, 3xHA:NLuc:MAX58) and co-expressed with OsH(I)PP FL proteins fused at their N-termini to the C-terminal part of luciferase (3xFlag:CLuc:OsH(I)PP). Co-expression of AVR-Pia with OsRGA5$_{Cter}$ (883 – 1116 aa) was used as a positive control. Leaves infiltrated with P19 alone served as background (BG) control. Leaf discs were collected 2 days post infiltration. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range. Values outside this range are shown as individual data points. Three independent replicates were performed (n = 8 per replicate). Due to experimental constraints, the following sample sizes apply to specific conditions: AVR-Pia/OsHPP09: n = 16 in replicates 1 and 2. Data points are individually plotted and shaped according to the replicate. Different letters indicate statistically significant differences based on a pairwise Wilcoxon test (α = 0.05).

the possibility that the studied FL HIPPs might not localize efficiently to the yeast nucleus,
potentially causing false-negative results in Y2H assays. To test this, we generated mutant
versions of the respective FL HIPPs, replacing the cysteine residue within the isoprenylation
motif with a serine residue, thereby preventing isoprenylation (**Supplementary Fig. 3a, 3b**).
However, this mutation did not restore interactions of FL OsHIPPs (OsHIPP14, OsHIPP39,
OsHIPP41) with AVR-Pia in Y2H assays (**Supplementary Fig. 2**).

The HMA domains of OsHPP09 and OsHPP11 share high sequence similarity with OsHPP10
(LOC_Os10g36200) (91.5 % and 74.6 % identical, respectively; **Supplementary Fig. 4**).
Surprisingly, despite this high level of conservation, neither the HMA domain nor the FL
version of OsHPP10 interacted with AVR-Pia in Y2H assays (**Fig. 1a**). By contrast, OsHIPP21
shares only 30-40% sequence identity with the HMA domains of the three OsHPPs
(**Supplementary Fig. 4**). Similar to OsHPP10, the HMA domain of OsHIPP21 did not interact
with AVR-Pia. However, AVR-Pia interacted with FL OsHIPP21. Taken together, these results
suggest that AVR-Pia interacts with and potentially targets OsHPP09, OsHPP11, and
OsHIPP21 in rice.

A previous study classified MAX effectors into 20 groups based on structural features [5].
MAX58 belongs to the same group as AVR-Pia, but the two effectors differ in surface
properties [5]. dSP-MAX58 (20-86 aa) did not interact with any of the tested OsH(I)PPs (neither
their HMA domains nor the FL versions) in Y2H assays (**Fig. 1a, Supplementary Fig. 1**),
although the effector protein was expressed (**Supplementary Fig. 3c**). These findings
suggest that, despite their shared structural fold, effectors with distinct surface properties
target different host proteins.

AVR-Pia interacts with rice H(I)PPs in planta
To test whether AVR-Pia and the four OsH(I)PPs also interact in planta, we performed split-
luciferase complementation assays in *Nicotiana benthamiana*. 3xFlag-CLuc-OsH(I)PP FL
proteins were co-expressed with 3xHA-NLuc-dSP-AVR-Pia or 3xHA-NLuc-dSP-MAX58.
Luminescence was measured from leaf discs collected 2 days after infiltration. Co-expression
of FL OsHPP09, OsHPP11, and OsHIPP21 with AVR-Pia led to high luminescence levels,
exceeding even those observed for the positive control OsRGA5$_{Cter}$ (883 – 1116 aa), which
includes the HMA ID, in complex with AVR-Pia (**Fig. 1b**). Strikingly, we also detected a strong
luminescence signal for OsHPP10/AVR-Pia, although no interaction between OsHPP10 and
AVR-Pia was observed in Y2H assays (**Fig. 1b**). The luminescence level for OsHPP10/AVR-
Pia was slightly lower than those of OsHPP09 and OsHPP11 but within a similar range, and
exceeded that of OsHIPP21. Consistent with our Y2H results, no interaction was observed in
planta between AVR-Pia and other tested FL OsHIPPs, including OsHIPP14, OsHIPP39 and

OsHIPP41 (**Fig. 1b**). Additionally, MAX58 showed no detectable interaction with any of the

tested OsH(I)PPs in split-luciferase complementation experiments (**Fig. 1b**). Expression of all

proteins in *N. benthamiana* was confirmed by immunoblot analyses (**Supplementary Fig. 5**).

Collectively, these data demonstrate that AVR-Pia interacts with OsHPP09, OsHPP10,

OsHPP11 and OsHIPP21 in planta.

AlphaFold Multimer models cannot differentiate between HMA domains which are

bound by AVR-Pia and those which are not

Several studies have reported the use of AlphaFold Multimer to differentiate between pairs of

effectors/host proteins which interact and those which do not [17,48,49]. We therefore tested

whether structural models of AVR-Pia in complex with experimentally verified interactors

(OsHPP09-HMA, OsHPP10-HMA, OsHPP11-HMA and OsHIPP21-HMA) showed

differences, either in the models themselves or in the associated confidence metrics,

compared to structural models of AVR-Pia with non-interacting HMA domains. We modelled

AVR-Pia/HMA complexes using AlphaFold2 and AlphaFold3. High confidence models were

obtained for all AVR-Pia/HMA pairs except AVR-Pia/OsHIPP42-HMA using AlphaFold2

(**Supplementary Fig. 6**) and for all AVR-Pia/HMA pairs except AVR-Pia/OsHIPP41-HMA

using AlphaFold3 (**Supplementary Fig. 7**). For the AlphaFold2 models, interface analysis with

qtPISA revealed a tendency towards increased interface area for AVR-Pia-interacting HMA

domains compared to non-interacting HMA domains. (**Supplementary Fig. 6)**. By contrast,

interface analysis (qtPISA) did not indicate any significant differences in the interface area,

binding energy, or number of intermolecular contacts for AlphaFold3 models of AVR-Pia-

interacting HMA domains compared to non-interacting HMA domains (**Supplementary Fig.**

7). Overall, AlphaFold models do not show a clear differentiation between HMA domains which

are bound by AVR-Pia and those which are not.

AVR-Pia-H3 does not interact with OsHPP09, OsHPP10, OsHPP11 and OsHIPP21

Two non-synonymous polymorphisms in AVR-Pia (F24S and T46N) have emerged in a subset

of rice-infecting *M. oryzae* isolates, disrupting the interaction with OsRGA5 and thereby

enabling these isolates to evade recognition by rice varieties carrying the OsRGA4/OsRGA5

NLR pair [9,11] (**Supplementary Fig. 8**). This virulence-associated allele is referred to as AVR-

Pia-H3 [9]. We next investigated whether these polymorphisms also affect direct binding to the

putative host target H(I)PPs. Pairwise Y2H assays revealed that, unlike wild-type AVR-Pia,

the AVR-Pia-H3 variant failed to interact with the HMA domains and FL versions of OsHPP09,

OsHPP10, OsHPP11 and OsHIPP21 (**Fig. 2a**). Protein expression of AVR-Pia-H3 was

confirmed by immunoblot (**Supplementary Fig. 3c**).

Fig. 2. AVR-Pia-H3 does not interact with OsHPP09, OsHPP10, OsHPP11 and HIPP21.

a Y2H interaction analysis between AVR-Pia-H3 and either the heavy metal-associated (HMA) domain or full-length (FL) proteins of OsH(I)PPs. OsHPP09/AVR-Pia served as positive control. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH). Photos were taken after 7 days of incubation. AD, activating domain; BD, binding domain. **b** Quantitative measurement of luciferase-derived signals shows that AVR-Pia-H3 does not interact with OsHPP09, OsHPP10, OsHPP11 and OsHIPP21 in a split-luciferase complementation assay. MAX effectors were fused to the N-terminal part of luciferase (3xHA:NLuc:MAX) and co-expressed with OsH(I)PP FL proteins fused at their N-termini to the C-terminal part of luciferase (3xFlag:CLuc:OsH(I)PP). Co-expression of AVR-Pia with OsHPP09 served as positive control, co-expression of OsHPP09/MAX58 and OsRGA5$_{Cter}$ (883 – 1116 aa)/AVR-Pia-H3 served as negative controls. Leaves infiltrated with P19 alone served as background (BG) control. Leaf discs were collected 2 days post infiltration. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range. Values outside this range are shown as individual data points. Three independent replicates were performed, with n = 8 per combination per replicate. Due to experimental constraints, the following sample sizes apply to specific conditions: AVR-Pia/OsHPP09: n = 16 in replicates 1 and 2. Data points are individually plotted and shaped according to the replicate. Different letters indicate statistically significant differences based on a pairwise Wilcoxon test (α = 0.05). **c** Analytical gel filtration traces obtained from injection of OsHPP09-HMA alone (top panel), AVR-Pia-H3 alone (middle panel) and the two proteins in a 1:1 molar ratio (bottom panel). Significant peaks are indicated by coloured dashed lines with the elution volume labelled. SDS-PAGE gel inserts show fractions from peak elution volumes indicated by the grey shaded regions. OsHPP09-HMA absorbs light at 280 nm poorly (molar extinction coefficient of 1490 M^{-1}cm^{-1}) so the peak corresponding to OsHPP09-HMA is small. The elution volume observed for OsHPP09-HMA is consistent with dimerisation, as has been observed for other purified HMA domains [10,50]. There is no shift in the elution volume for AVR-Pia-H3 in the presence of OsHPP09-HMA indicating that the two proteins do not form a complex.

To validate these findings in planta, we performed split-luciferase complementation assays

following transient co-expression of OsH(I)PPs and AVR-Pia-H3 in *N. benthamiana*. Only

weak luminescence signals were detected for AVR-Pia-H3 with OsH(I)PPs, comparable to the

negative control pair OsRGA5$_{Cter}$/AVR-Pia-H3 and OsHPP09/MAX58, and markedly lower

than the positive control OsHPP09/AVR-Pia (**Fig. 2b**). These results suggest that the AVR-

Pia-H3 polymorphisms severely diminish effector-target binding. Protein expression was

confirmed by immunoblot analysis (**Supplementary Fig. 9**). We also tested for an interaction

between purified AVR-Pia-H3 and OsHPP09-HMA proteins in vitro using analytical size

exclusion chromatography. AVR-Pia-H3 had a peak elution volume of 14.7 ml when analysed

alone, while OsHPP09-HMA had a peak elution volume of 12.9 ml (consistent with

dimerisation, as has been reported for other HMA domains in solution) (**Fig. 2c**). The elution

volumes of the two proteins were unchanged when a 1:1 mixture of the proteins was analysed,

indicating that they did not form a stable complex in vitro (**Fig. 2c**). By contrast, the peak

elution volume of AVR-Pia shifted from 15.3 ml when analysed alone to 12.7 ml following

incubation with OsHPP09-HMA (1:1 effector:HMA ratio), consistent with the formation of an

AVR-Pia/OsHPP09-HMA complex (**Supplementary Fig. 10**).

Together, these data indicate that the same mutations that disrupt NLR recognition also

abolish the ability of AVR-Pia to bind to its candidate host targets, suggesting that AVR-Pia

might use a similar binding interface for interacting with both the host targets and the immune

receptor during recognition.

AVR1-CO39 does not interact with OsHPP09, OsHPP10, OsHPP11 and OsHIPP21

As the HMA ID of OsRGA5 also binds the MAX effector AVR1-CO39 [9,10] we tested whether

AVR-Pia and AVR1-CO39 bind to the same subset of H(I)PPs. To this end, we tested

interactions between dSP-AVR1-CO39 (22 – 89 aa) and the HMA domain as well as FL

version of OsHPP09, OsHPP10, OsHPP11 and OsHIPP21 in Y2H assays. AVR1-CO39 did

not interact with either the HMA domains or FL versions of any of these rice proteins, whereas

interaction with OsRGA5$_{Cter}$ including the HMA ID was confirmed (**Fig. 3a**). Effector

expression in yeast was confirmed by immunoblot analysis (**Supplementary Fig. 3c**).

To assess whether these interactions occur in planta, we performed split-luciferase

complementation assays in *N. benthamiana*. We observed only weak luminescence signals

indicating low-level interactions between AVR1-CO39 and FL OsHPP09, OsHPP10 and

OsHPP11. However, these signals were substantially lower than those observed for the AVR-

Pia/OsHPP09 interaction (**Fig. 3b**). Consistent with our Y2H results, no interaction was

detected between FL OsHIPP21 and AVR1-CO39, similar to the negative control combination

OsPi21/AVR1-CO39 (**Fig. 3b**). Protein expression was validated by immunoblot analysis

Fig. 3. OsH(I)PPs interacting with AVR-Pia do not interact with AVR1-CO39.

a Y2H interaction analysis between AVR1-CO39 and either the heavy metal-associated (HMA) domain or full-length (FL) proteins of OsH(I)PPs. $OsRGA5_{Cter}$ (883 – 1116 aa)/AVR1-CO39 served as positive control. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH). Photos were taken after 7 days of incubation. Ø indicates positions where no yeast was spotted. AD, activating domain; BD, binding domain. **b** Quantitative measurement of luciferase-derived signals shows that AVR1-CO39 does weakly interact with OsHPP09, OsHPP10 and OsHPP11 but not with OsHIPP21 in a split-luciferase complementation assay. MAX effectors were fused to the N-terminal part of luciferase (3xHA:NLuc:MAX) and co-expressed with OsH(I)PP FL proteins fused at their N-termini to the C-terminal part of luciferase (3xFlag:CLuc:OsH(I)PP). Co-expression of AVR-Pia or AVR1-CO39 with OsPi21 served as negative controls, while AVR-Pia/OsHPP09 served as a positive control. Leaf discs were collected 2 days post infiltration. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range. Values outside this range are shown as individual data points. Three independent replicates were performed, with n = 8 per combination per replicate. Data points are individually plotted and shaped according to the replicate. Different letters indicate statistically significant differences based on a pairwise Wilcoxon test ($\alpha = 0.05$). **c** Analytical gel filtration traces obtained from injection of OsHPP09-HMA alone (top panel), AVR1-CO39 alone (middle panel) and the two proteins in a 1:1 molar ratio (bottom panel). Significant peaks are indicated by coloured dashed lines with the elution volume labelled. SDS-PAGE gel inserts show fractions from peak elution volumes indicated by the grey shaded regions. OsHPP09-HMA absorbs light at 280 nm poorly (molar extinction coefficient of 1490 $M^{-1}cm^{-1}$) so the peak corresponding to OsHPP09-HMA is small. The small peak at ~13.7 ml in the trace for AVR1-CO39 is likely due to some dimerisation of the AVR1-CO39 protein; the quantity of protein is insufficient to observe a corresponding band on the SDS-PAGE gel. The elution volume observed for OsHPP09-HMA is consistent with

dimerisation, as has been observed for other purified HMA domains [10,50]. There is no shift in the elution volume

for AVR1-CO39 in the presence of OsHPP09-HMA indicating that the two proteins do not form a complex.

(**Supplementary Fig. 11**). Analytical size-exclusion chromatography with purified proteins
further supported the absence of complex formation (**Fig. 3c**).

Collectively, these results suggest that although AVR1-CO39 and AVR-Pia are both
recognised by the same HMA ID in OsRGA5, they do not target the same H(I)PPs in the host.

AVR-Pia interacts with a distinct group of H(I)PPs to AVR-Pik and Pwl2
Previous studies have shown that the MAX effectors AVR-PikC, AVR-PikD and Pwl2 target
H(I)PPs [17,32,41,46]. To investigate whether the OsH(I)PPs identified as AVR-Pia interactors
(OsHPP09, OsHPP10, OsHPP11 and OsHIPP21) also associate with AVR-Pik or Pwl2, we
tested their pairwise interactions using Y2H assays. Our results revealed that neither the HMA
domain nor the FL versions of these proteins bound directly to dSP-AVR-PikC (22 – 113 aa),
dSP-AVR-PikD (22 – 113 aa) or dSP-Pwl2 (22 – 145 aa) (**Fig. 4a**). As a control, we confirmed
the known interactions of AVR-PikC and AVR-PikD with OsHIPP19 and OsHIPP20, and Pwl2
with OsHIPP43 (**Fig. 4a**). Interestingly, we observed that AVR-Pia also directly interacted with
FL OsHIPP19 under low-stringency Y2H conditions (**Fig. 4a**).

The lack of interaction between AVR-PikC and AVR-PikD with the FL proteins OsHPP09,
OsHPP10, OsHPP11 and OsHIPP21 was further validated through split-luciferase
complementation assays using proteins transiently expressed in *N. benthamiana* (**Fig. 4b**).
AVR-Pia showed a strong interaction with OsHIPP19 and a weaker interaction with OsHIPP20
(**Fig. 4b**). Co-expression of Pwl2 with OsHPP09, OsHPP10, OsHPP11 and OsHIPP21
resulted in very weak luminescence. Protein expression was confirmed by immunoblot
analysis (**Supplementary. Fig. 12**).

Together, our findings indicate that the HMA domains bound by AVR-Pia are not targeted by
AVR-Pik and Pwl2. Despite sharing a conserved MAX effector fold, these effectors seem to
generally engage different target H(I)PPs, highlighting the specificity of effector-host target
interactions.

AVR-Pia interacts with the HMA domains of OsHPP09, OsHPP10, OsHPP11 and
OsHIPP21 with varying affinities
To determine the affinity with which AVR-Pia binds to the HMA domains of OsHPP09,
OsHPP10, OsHPP11 and OsHIPP21, we used isothermal titration calorimetry (ITC).
Recombinant AVR-Pia and HMA domains were produced and purified from *E. coli*
(**Supplementary Fig. 13**). Titration of AVR-Pia into a solution containing any of the purified
HMA domains resulted in negative peaks in the titration curve (**Fig. 5, Supplementary Fig.**
14), indicative of an exothermic binding interaction. Dissociation constant (K_D) values were

Fig. 4. OsH(I)PPs interacting with AVR-Pia do not interact with AVR-Pik or Pwl2.

a Y2H interaction analysis between AVR-PikC, AVR-PikD, Pwl2 and AVR-Pia and either the heavy metal-
associated (HMA) domain or full-length (FL) proteins of OsH(I)PPs. OsHIPP19/AVR-PikC, OsHIPP20/AVR-PikC,
OsHIPP19/AVR-PikD, OsHIPP20/AVR-PikD and OsHIPP43/Pwl2 served as positive controls. Serial dilutions of
diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-
protein interactions (SD/-LWH). Photos were taken after 7 days of incubation. AD, activating domain; BD, binding
domain. **b** Quantitative measurement of luciferase-derived signals shows that AVR-PikC and AVR-PikD do not
interact with OsHPP09, OsHPP10, OsHPP11 and OsHIPP21, while Pwl2 shows weak luminescence signals with
these proteins, suggesting weak interactions. AVR-Pia also interacts with OsHIPP19 and OsHIPP20 in planta.
MAX effectors were fused to the N-terminal part of luciferase (3xHA:NLuc:MAX) and co-expressed with OsH(I)PP
FL proteins fused at their N-termini to the C-terminal part of luciferase (3xFlag:CLuc:OsH(I)PP). Co-expression of
AVR-PikC/AVR-PikD with OsHIPP19/20 and Pwl2 with OsHIPP43 served as positive controls, AVR-Pia with
OsHIPP43 served as negative control. Leaves infiltrated with P19 alone served as background (BG) control. Leaf
discs were collected 2 days post infiltration. Box plots show the median (line), mean (cross) and upper/lower
quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range. Values
outside this range are shown as individual data points. Unless otherwise stated, three independent replicates were
performed, with n = 8 per combination per replicate. Due to experimental constraints, the following sample sizes
apply to specific conditions: AVR-Pia/OsHIPP19 and AVR-Pia/OsHIPP20 in AVR-PikC and AVR-PikD panels: n =
4 in replicates 1 and 3. BG control in AVR-PikC and AVR-PikD panels: n = 3 in replicates 1 and 3 and BG control

in Pwl2 panel: n = 2 in replicates 1 and 3. AVR-PikD/OsHIPP19 n = 16 in replicates 1 and 3 and n = 24 in replicate

2. Data points are individually plotted and shaped according to the replicate. Different letters indicate statistically

significant differences based on a pairwise Wilcoxon test (α = 0.05).

Fig. 5. AVR-Pia binds the HMA domains of OsH(I)PPs with varying affinities.

a Representative ITC experiments for each of the AVR-Pia/HMA interactions tested. Top panel shows the raw thermogram obtained from titration of AVR-Pia into a solution containing the purified HMA domains. Central panel shows the integrated heats (coloured dots) and binding isotherm fitted to a single site model (black line) using the MicroCal PEAQ-ITC analysis software (Malvern Panalytical). Bottom panel shows the differences (coloured crosses) between the modelled and observed values (residuals). **b** Dot plot representation of the apparent KD for each of the AVR-Pia/HMA interactions in three independent experiments. Coloured dots represent the KD vales determined in each replicate with coloured lines representing the error associated with the KD value as determined by the MicroCal PEAQ-ITC analysis software (Malvern Panalytical).

determined from binding isotherms (**Fig. 5, Supplementary Fig. 14**) fitted to a single site

model.

AVR-Pia bound to the HMA domain of OsHPP09 with an apparent K_D of 115 nM (**Fig. 5**).

Despite differing in just six amino acids, AVR-Pia bound OsHPP10-HMA with significantly

lower affinity, with an apparent K_D of 5.26 µM. The affinities of the interactions of AVR-Pia with

OsHPP11-HMA and OsHIPP21-HMA fall between these two extremes, with apparent K_Ds of

772 nM and 1.29 µM, respectively. (**Fig. 5, Supplementary Fig. 14, Supplementary Table**

3). Dissociation constants and thermodynamic parameters and profiles are provided in

Supplementary Table 3 and Supplementary Fig. 15. The apparent K_Ds, as determined by

ITC, for the interactions between OsRGA5-HMA and AVR-Pia or AVR1-CO39 have previously

been reported as 7.8 µM [11] and 5.1 µM[10], respectively.

The crystal structure of AVR-Pia in complex with the HMA domain of OsHPP09 reveals

interface contacts underpinning high affinity binding

To understand the structural basis of the interaction between AVR-Pia and OsHPP09-HMA,

we purified and crystallised the effector/HMA complex. Needle-like crystals were obtained in

commercial screens and subsequent optimisation of conditions (see Materials and Methods)

resulted in crystals suitable for diffractions studies. X-ray diffraction data were collected at the

European Synchrotron Radiation Facility (ESRF) to a resolution of 1.65 Å. The structure was

solved by molecular replacement and refined and validated as described in the Materials and

Methods. Data collection, processing and refinement statistics are presented in

Supplementary Table 4. The final refined model has been deposited in the Protein Data Bank

with accession code 9RSV.

As expected, OsHPP09-HMA adopts the well-characterised HMA domain fold, consisting of a

four-stranded antiparallel β-sheet and two α-helices arranged with βαββαβ topology (**Fig. 6**).

Solution structures of AVR-Pia alone have previously been determined by NMR spectroscopy

[4,51] (PDB accession codes 2MYW, 2N37) and the crystal structure of AVR-Pia bound to the

integrated HMA domain of the NLR protein OsPikp-1 has also been solved [45] (PDB accession

code 6Q76). The structure of AVR-Pia determined here is highly similar (RMSD of 0.357 Å to

6Q76, 0.951 Å to 2MYW, and 1.690 Å to 2N37) and adopts the six-stranded β-sandwich

structure characteristic of the MAX effector family (**Fig. 6**) [4,5].

The global structure of the complex between AVR-Pia and OsHPP09-HMA resembles the

previously determined structures of AVR-Pia bound to the HMA ID of OsPikp-1 (PDB 6Q76)

and AVR1-CO39 in complex with the HMA ID of OsRGA5 (PDB 5ZNG) (**Supplementary Fig.**

16). Analysis with qtPISA [52] (**Supplementary Table 5**) determined an interface area of 521.7

Fig. 6. Crystal structure of the HMA domain of OsHPP09 in complex with AVR-Pia.

The structures of OsHPP09-HMA and AVR-Pia are represented as teal and gold ribbons, respectively. The amino

acids comprising β2 of AVR-Pia and β2 of the HMA domain are represented as cylinders. Side chains involved in

intermolecular hydrogen bonds/salt bridges are also represented as cylinders. Hydrogen bonds are represented

by black dashed lines labelled with the bond length. Left insert shows main chain hydrogen bonds between AVR-

Pia-β2 and OsHPP09-β2. Right insert shows intermolecular contacts between side chains of residues in OsHPP09-

α1 and AVR-Pia.

Å^2, fractionally larger than the 460.7 Å^2 interface between AVR-Pia/OsPikp-HMA and the

492.8 Å^2 interface between AVR1-CO39/OsRGA5-HMA. In all three cases, β2 of the MAX

effector is aligned in an antiparallel orientation with β2 of the HMA domain, resulting in the

formation of an antiparallel β-sheet comprising β6, β1 and β2 of the MAX effector and the four-

stranded β-sheet of the HMA domain (**Supplementary Fig. 16a**), with hydrogen bonding

between the backbone amine and carbonyl groups of the peptide backbone of the adjacent β-

strands (**Supplementary Fig. 16b**).

In contrast to the structures of OsPikp-HMA with AVR-Pia and OsRGA5-HMA with AVR1-

CO39, where the interface is dominated by main chain hydrogen bonding, in the OsHPP09-

HMA/AVR-Pia structure, the side chains of multiple residues in α1 of the HMA domain form

intermolecular hydrogen bonds with the effector. The side chains of OsHPP09^{E16} and

OsHPP09^{K19} form hydrogen bonds with the peptide backbone of AVR-Pia^{L38} (**Supplementary

Fig. 16c**). The amino acid in the corresponding position to OsHPP09^{K19} in OsPikp-HMA is

OsPikp-1^{R203}, however it is OsPikp-1^{R226}, in the β2-β3 loop, which forms an analogous

hydrogen bond with AVR-Pia^{L38} (**Supplementary Fig. 16c**). OsRGA5-HMA has a basic

residue (R1012) at the corresponding position to OsHPP09^{K19}, and the side chain of OsRGA5-

HMA^{R1012} is orientated towards the side chain of AVR1-CO39^{D35}, however the distance

between them (4.1 Å) is greater than the 4 Å cutoff typically used when considering salt bridge

interactions (**Supplementary Fig. 16c**). At the start of β2, the side chain of OsHPP09^{D33} forms

salt bridge interactions with the side chain of AVR-Pia^{R43} (**Supplementary Fig. 16d**). This

aspartate residue is conserved in both OsPikp-HMA (D217), where it similarly forms a salt

bridge with AVR-Pia^{R43}, and in OsRGA5-HMA (D1026), where it forms an indirect, water-

mediated contact with the backbone of AVR1-CO39^{T41} (**Supplementary Fig. 16d**). The

OsHPP09^{Q20} side chain forms hydrogen bonds with the side chain hydroxyl of AVR-Pia^{Y41}

(**Supplementary Fig. 16e**). The amino acid in the corresponding position in OsPikp-HMA is

serine, and a water-mediated contact is formed between the smaller side chain of OsPikp-

HMA^{S204} and the side chain hydroxyl of AVR-Pia^{Y41} (**Supplementary Fig. 16e**). Finally,

towards the C-terminal end of OsHPP09-α1, the side chain of OsHPP09^{D28} forms a salt bridge

with AVR-Pia^{R23} (**Supplementary Fig. 16f**). By contrast, the corresponding amino acid in

OsPikp-1, OsPikp-1^{S212}, forms a hydrogen bond with the side chain hydroxyl of AVR-Pia^{Y85}

(**Supplementary Fig. 16f**). Overall, the increased number of intermolecular hydrogen bonds

in the AVR-Pia/OsHPP09-HMA structure compared to the structure of AVR1-CO39/OsRGA5-

HMA likely explain the reported difference in binding affinities.

To validate the interface observed in the crystal structure of OsHPP09-HMA in complex with

AVR-Pia, NMR titration with ^{1}H-^{15}N-HSQC NMR spectra of ^{15}N-labelled AVR-Pia alone and in

the presence of purified unlabelled OsHPP09-HMA were performed. HSQC spectra for [15]N-
labelled AVR-Pia had previously been collected and peaks assigned to the protein [4], and these
assignments were used in the present study. Binding of AVR-Pia to OsHPP09-HMA alters the
chemical environment of the amino acids located at the binding interface, resulting in a change
in the chemical shift observed for those amino acids in the HSQC spectra. Depending on the
rate of complex formation and dissociation, expressed by the exchange rate constant kex, and
the chemical shift difference $\Delta\omega$ between the unbound and bound states ($\Delta\omega$ = difference
between the resonance frequencies of the exchanging sites), different exchange regimes
occur. NMR titration showed that the AVR-Pia/OsHPP09-HMA complex was in slow exchange
with kex << $\Delta\omega$ since separate resonances appeared for individual species. For certain peaks
in the spectra of [15]N-labelled AVR-Pia with OsHPP09-HMA, it was not possible to confidently
identify the corresponding peak in the spectra for [15]N-labelled AVR-Pia alone. However,
significant changes in chemical shift ($\Delta\delta$ AV > 0.4 ppm) were observed for multiple residues
in β2 and β3 of AVR-Pia, consistent with the interface in the crystal structure **(Supplementary**
Fig. 17).

Replacement of OsHPP09[E16] with alanine or arginine disrupts AVR-Pia binding

Next, we aimed to identify point mutations in the HMA domain of OsHPP09 which interfere
with AVR-Pia binding. First, using the crystal structure, we selected residues where the side
chains contribute intermolecular hydrogen bonds or salt bridges and mutated each residue to
alanine (E16A, K19A, Q20A, D28A, D33A) and, for the acidic residues, also to arginine (E16R,
D28R, D33R). D33 is conserved in OsHPP09, OsHPP10 and OsHPP11, but also in the HMA
IDs of Pikp-1 (D217) and RGA5 (D1026) (**Supplementary Fig. 16C**). By contrast, the residue
in the corresponding position in Pikm-HMA, which does not bind AVR-Pia [45], is histidine. We
therefore also included the D33H mutation. Second, we used FoldX [53] to predict the effect of
mutating each interface residue in OsHPP09-HMA to all other amino acids. Based on the
calculated difference of interaction free energy ($\Delta\Delta$G) upon mutation, we selected three further
mutations (K19W, I35K, A37W) for experimental characterisation.

Mutations were introduced into expression constructs by site-directed mutagenesis. The effect
of the mutations on AVR-Pia binding was tested in Y2H experiments. Introduction of the point
mutations E16A, E16R, D28A, D28R and I35K in OsHPP09 resulted in a complete loss of
interaction with AVR-Pia (**Fig. 7a**). In contrast, the remaining mutations (K19A, K19W, Q20A,
D33A, D33H, D33R, A37W) did not completely abolish the interaction (**Fig. 7a**). However,
several including K19A, K19W, D33A, D33H, D33R and A37W led to reduced yeast growth
on interaction-selective media when co-expressed with AVR-Pia, suggesting a weakening of
the interaction compared to the wildtype OsHPP09/AVR-Pia pair. Immunoblot analysis

Fig. 7. Mutation of OsHPP09 E16 reduces AVR-Pia binding.

a Y2H interaction analysis between AVR-Pia and full-length (FL) OsHPP09 mutant versions carrying single mutations in the binding interface. OsHPP09^{WT}/AVR-Pia served as positive control. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH and SD/-LWH supplemented with 5 mM 3AT). Photos were taken after 7 days of incubation. AD, activating domain; BD, binding domain. **b** Quantitative measurement of luciferase-derived signals shows that the E16A, E16R and I35K mutations in OsHPP09 significantly reduce its interaction with AVR-Pia in a split-luciferase complementation assay. AVR-Pia was fused to the N-terminal part of luciferase (3xHA:NLuc:AVR-Pia) and co-expressed with OsH(I)PP FL proteins fused at their N-termini to the C-terminal part of luciferase (3xFlag:CLuc:OsH(I)PP). Co-expression of AVR-Pia with OsHPP09 served as positive control, co-expression of OsHIPP43 with AVR-Pia served as negative control. Leaves infiltrated with P19 alone served as background (BG) control. Leaf discs were collected 2 days post infiltration. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range. Values outside this range are shown as individual data points. Three independent replicates were performed, with n = 8 per combination per replicate. Data points are individually plotted and shaped according to the replicate. Different letters indicate statistically significant differences based on a pairwise Wilcoxon test (α = 0.05). **c** Circular dichroism (CD) spectra obtained for WT OsHPP09-HMA (pale blue) and variants carrying E16A, E16R, I35K or D33H mutations. Negative peaks at 222 nm and 208 nm and a positive peak at 193 nm are indicative of α-helices while a negative peak at 218 nm and positive peak at 195 nm are indicative of β-sheets, consistent

with the expected βαββαβ topology of an HMA domain. In the case of the I35K mutant, low ellipticity above 210

nm and a negative peak at 200 nm is indicative of disorder. **d** Overlays of cross-peaks from HSQC spectra

corresponding to residues L39, I44, E56 and Y85 of ^{15}N-labelled AVR-Pia alone (orange) and in the presence of

either WT OsHPP09-HMA or variants carrying E16A, E16R or D33H mutations (blue/green/purple). Overlays of

the full spectra are provided in Supp. Figures S19-S21. Chemical shift perturbation of ^{15}N-labelled AVR-Pia is

indicative of an interaction with the HMA domain.

revealed that the K19A, K19W and I35K mutants exhibited lower protein expression levels,
while all other mutant constructs showed similar expression to wildtype OsHPP09
(**Supplementary Fig. 3d**).

To assess the effect of the mutations in the context of the full-length protein in planta, we
performed split-luciferase complementation assays in *N. benthamiana*. The E16A and E16R
mutants showed significantly reduced interaction with AVR-Pia, while the I35K mutant
completely lost its ability to interact with AVR-Pia, consistent with the Y2H results (**Fig. 7b**). In
contrast, the D28A and D28R mutations, which abolished interaction in Y2H assays, had only
a minor effect on the interaction with AVR-Pia in planta, similar to the other tested mutants
(**Fig. 7b**). Immunoblot analysis demonstrated that all proteins were expressed at comparable
levels in planta (**Supplementary Fig. 18**).

Next, we aimed to establish whether the E16A, E16R and I35K mutations affect the structure
of the HMA domain. OsHPP09^{E16A}-HMA, OsHPP09^{E16R}-HMA and OsHPP09^{I35K}-HMA, as well
as OsHPP09^{D33H}-HMA (as a control), were produced in *E. coli* and purified to homogeneity.
The molecular weights of the modified HMA domains were confirmed by intact mass
spectrometry. To test whether the mutation affects the overall structure of the protein, we first
obtained circular dichroism (CD) spectra for the HMA domains. The spectra for native
OsHPP09-HMA showed features characteristic of α-helices (negative peaks around 222 nm
and 208 nm, positive peak around 193 nm) and β-sheets (negative peak around 218 nm,
positive peak around 195 nm) consistent with the HMA domain fold (**Fig. 7c**). While CD
spectra obtained for OsHPP09^{E16A}-HMA, OsHPP09^{E16R}-HMA and OsHPP09^{D33H}-HMA were
comparable to the spectra obtained for the native protein, OsHPP09^{I35K}-HMA showed features
of disorder (low ellipticity above 210 nm and a negative peak around 200 nm) (**Fig. 7c**). These
results suggest that the loss of AVR-Pia binding observed for the I35K mutant is likely due to
disruption of the HMA domain fold.

We obtained ^{1}H-^{15}N-HSQC NMR spectra of ^{15}N-AVR-Pia in the presence of purified unlabelled
HMA domains of OsHPP09^{E16A}, OsHPP09^{E16R}, OsHPP09^{D33H} and OsHPP09^{I35K}. The spectra of
^{15}N-AVR-Pia with OsHPP09^{D33H}-HMA was highly similar to the spectra of ^{15}N-labelled AVR-Pia
with WT OsHPP09-HMA (**Fig. 7d**, **Supplementary Fig. 19**). By contrast, the spectra of ^{15}N-
AVR-Pia with OsHPP09^{E16A} or OsHPP09^{E16R} showed no chemical shift perturbation and
superimposed with the spectra of ^{15}N-AVR-Pia alone (**Fig. 7d**, **Supplementary Fig. 20**,
Supplementary Fig. 21). Taken together, these results demonstrate that E16A and E16R
mutations block AVR-Pia binding without disrupting the HMA domain fold.

Discussion

The *Magnaporthe oryzae* effector AVR-Pia is a member of the MAX family of sequence-unrelated, structurally conserved effectors. Here, we identified four small HMA domain-containing proteins which are bound by AVR-Pia.

Structural variation between MAX effectors underpins interactions with different host targets

AVR-Pia is the third MAX effector which has been shown to interact with H(I)PPs. AVR-Pik was found to interact with the HMA domains of multiple H(I)PPs within a large phylogenetic clade, referred to as "clade A" [17]. By contrast, Pwl2 interacts specifically with OsHIPP43 and its homologs from other grass species [46]. In the present study, we found that AVR-Pia binds to the HMA domains of OsHPP09, OsHPP10, OsHPP11, and OsHIPP21. Neither AVR-Pik nor Pwl2 interacts with these H(I)PPs, demonstrating that AVR-Pia binds a distinct group of H(I)PPs to those previously identified as MAX effector targets.

Strikingly, the interactions between each of the three MAX effectors and HMA domains involve distinct interfaces. In the crystal structures of AVR-Pia/OsHPP09-HMA and Pwl2/OsHIPP43-HMA, β2 of the core MAX fold and β2 of the HMA domain are aligned in an antiparallel orientation [46]. Hydrogen bonds between the two β-strands lead to the formation of an antiparallel β-sheet comprising the four β-strands of the HMA domain and β2, β1 and β6 of the MAX fold. When the HMA domains are superposed, the relative positions of β2 of the MAX fold of Pwl2 and AVR-Pia are very similar (**Supplementary Fig. 22**). However, the other β-strands of Pwl2 are shifted compared to AVR-Pia, enabling residues in β3, β4, and the β3- β4 loop of the MAX fold to contribute to the HMA-binding interface. By contrast, in the complex of OsHIPP19-HMA/AVR-PikF, an antiparallel β-sheet forms from the four β-strands of the HMA domain and β3, β4 and β5 of the MAX fold, mediated by main chain hydrogen bonding between β3 of the MAX fold of AVR-PikF and β4 of HIPP19-HMA [41]. Furthermore, while AVR-Pia consists solely of the core MAX fold, AVR-Pik has an unstructured extension at the N-terminus, while in Pwl2 the characteristic MAX fold is proceeded by an α-helix and a C-terminal unstructured region. In both cases, these extensions are involved in HMA binding (**Supplementary Fig. 22**). Pwl2^{E89}, in the α-helix, forms a hydrogen bond with OsHIPP43-HMA^{K51}, while the C-terminal unstructured segment of Pwl2 extends across the HMA domain, forming an extensive interface involving both hydrogen bonds and π-stacking interactions. Similarly, the N-terminal extension of AVR-Pik binds OsHIPP19-HMA by forming multiple hydrogen bonds with residues in β3, β2 and the β2- β3 loop of the HMA domain. The interface in the AVR-Pia/OsHPP09-HMA complex is smaller than in AVR-PikF/OsHIPP19-HMA and

Pwl2/OsHIPP43-HMA (521.6 $Å^2$ compared to 1044.3 $Å^2$ and 1965.2 $Å^2$, respectively) and
formed almost exclusively by residues in β2 of the core MAX fold.

It is interesting that at least three MAX effectors have evolved to bind HMA domains in different
ways. This highlights how structural variation within this effector family can underpin different
interactions with host targets. At present, the impacts of effector binding to H(I)PPs are
unclear, and it remains to be seen whether binding to different interfaces leads to different
outcomes.

While AVR-Pia, AVR-Pik and Pwl2 all target OsH(I)PPs, AvrPiz-t is reported to interact with
diverse host proteins, including rice E3 ubiquitin ligases [54,55], a potassium channel [56], a bZIP-
type transcription factor [57] and a nucleoporin [58]. This suggests that MAX effectors may target
many different host proteins and processes during infection.

The HMA domains of the OsH(I)PPs bound by AVR-Pia share limited sequence
similarity with the HMA-ID of OsRGA5

AVR-Pia is recognised by the paired rice NLR immune receptors OsRGA5 and OsRGA4
through direct binding of the effector to a non-canonical HMA domain integrated into OsRGA5
[9]. The integrated domain model [13] posits that such non-canonical domains originated in the
host targets of effectors, and were incorporated into NLR proteins to function as effector
"sensors". Consistent with this, the crystal structure of AVR-Pia/OsHPP09 revealed that AVR-
Pia binds OsHPP09-HMA and OsRGA5-HMA through similar interfaces [11].

AVR1-CO39 is also recognised by direct binding to the HMA ID of OsRGA5. However, AVR1-
CO39 did not interact with the same OsH(I)PPs as AVR-Pia. AVR1-CO39 also did not interact
with the "clade A" OsH(I)PPs bound by AVR-PikD [17]. AVR1-CO39 is broadly absent from rice-
infecting isolates of *M. oryzae* [6,59,60] and loss of this effector has potentially contributed to a
host shift from foxtail millet (*Setaria* sp.) to rice. It is therefore possible that AVR1-CO39 has
evolved to target H(I)PPs from other grass species, and does not interact with those present
in rice.

Interestingly, phylogenetic analysis based on HMA domains clusters OsRGA5-HMA with the
H(I)PPs targeted by AVR-Pik and the HMA ID of OsPik-1, and not with OsHPP09, OsHPP10,
OsHPP11, and OsHIPP21 [17]. We observed interaction between AVR-Pia and FL OsHIPP19,
but not with the HMA domain alone, in low stringency Y2H experiments and split-luciferase
assays. This suggests that the C-terminal part of OsHIPP19 contributes to AVR-Pia binding
and merits further study.

Previously published ITC experiments showed that AVR-Pia binds OsRGA5-HMA with a K_D

of 7.8 μM [11]. This is comparable to the binding affinity of AVR-Pia to OsHPP10-HMA

determined in the present study, but at least one order of magnitude weaker than the

interaction between AVR-Pia and OsHPP09-HMA. This higher affinity binding can be

explained by the number of intermolecular hydrogen bonds and salt bridges at the interface

between AVR-Pia and OsHPP09; while no experimental structure is available for AVR-Pia

bound to OsRGA5-HMA, available structures of AVR1-CO39/OsRGA5-HMA and AVR-

Pia/OsPikp-HMA, combined with modelling approaches, suggest that the AVR-Pia/OsRGA5-

HMA interface likely involves fewer intermolecular contacts. A similar pattern was previously

reported for AVR-Pik/HMA interactions; surface plasmon resonance experiments showed that

AVR-PikD interacts with the HMA domain of OsHIPP19 with higher affinity than the HMA ID

of OsPik-1 [41].

It is important to note that these binding affinities are determined in vitro using the purified

HMA domain. These experiments therefore lack the biological context of the FL protein or the

plant cell environment. In the case of the AVR-Pia/OsRGA5 interaction, regions outside of the

OsRGA5 HMA domain contribute to AVR-Pia binding, and it is possible that AVR-Pia also

interacts with the C-terminal regions of the OsH(I)PPs. While the AVR-Pia/OsRGA5-HMA

interaction may be of lower affinity than the AVR-Pia/OsHPP09-HMA interaction, it is sufficient

for OsRGA5/OsRGA4 to activate immune signalling. A higher affinity interaction is unlikely to

confer any additional evolutionary benefit to the plant. If OsRGA5 recognises other pathogen

effectors, mutations which increase affinity for AVR-Pia may come at the cost of reducing

affinity/binding to other effectors.

H(I)PPs are targeted by diverse pests and pathogens, though their precise function in
plant susceptibility/resistance remains unclear

H(I)PPs, which form extensive families in plants [20,24,25], have been reported to be involved in

intracellular metal transport and homeostasis, responses to cold and drought stress and

tolerance to heavy metals. In addition to the interactions between *M. oryzae* MAX effectors

and OsH(I)PPs, there are a growing number of reports of plant H(I)PPs targeted by proteins

from other pests and pathogens. For example, the putative effector RsMf8HN from the

necrotrophic fungus *Rhizoctonia solani* interacts with OsHIPP28 [36], while a HIPP from

sugarcane (ScPi21) is a target for the *Fusarium sacchari* effector Fs00367 [33]. CsHIPP03 was

identified in a Y2H screen as a putative target of the core effector SDE34 secreted by

Candidatus Liberibacter asiaticus (CLas), a phloem-colonising bacteria which causes citrus

huanglongbing disease [38]. An effector secreted by the root-knot nematode *Meloidogyne*

graminicola binds rice OsHPP04 and suppresses a flg22-induced ROS burst [34]. Finally, the

potato mop-top virus (PMTV) movement protein TGB1 interacts with NbHIPP26 to promote the spread of the virus through the plant [29]. Together, these results suggest that the manipulation of H(I)PPs is required for successful infection by diverse pests and pathogens. In spite of this, little is known about the molecular mechanisms through which H(I)PPs affect plant resistance/susceptibility.

HMA domains typically coordinate metal ions via two cysteine residues in a MxCxxC motif positioned between α1 and β2. Intriguingly, the metal-binding motif is degenerate in all four HMA domains bound by AVR-Pia, with both cysteines replaced by other amino acids (MT<u>D</u>EK<u>T</u> in OsHPP09, MT<u>D</u>DK<u>I</u> in OsHPP10, MS<u>D</u>TK<u>M</u> in OsHPP11 and MT<u>D</u>ER<u>K</u> in OsHIPP21), presumably indicating a lack of metal-binding activity by these proteins. Consistent with this, no electron density suggestive of bound metal ions were observed in the crystal structure of OsHPP09-HMA in complex with AVR-Pia. Interestingly, OsHPP09^{E16}, which is important for AVR-Pia binding, is located in the degenerate metal binding motif. Of the clade A H(I)PPs bound by AVR-Pik, some contain an intact MxCxxC motif, while others (such as OsHIPP19 and OsHIPP20) lack one or both cysteine residues. OsHIPP43 has an intact MxCxxC motif, though it has not been established whether it binds metal ions in planta. In cases where HMA domains lack a complete MxCxxC motif, and presumably also lack the capacity to bind metals, the functions of the H(I)PPs are particularly intriguing. They may have acquired entirely new functions outside metal transport and homeostasis, with the stable core fold of the HMA domain supporting diversification of surface properties for new functions.

HIPPs and HPPs differ in the presence or absence of a CaaX isoprenylation motif at the C-terminus. Correspondingly, OsHIPP21 has a CaaX motif (CCQM) while no such motif is present in OsHPP09, OsHPP10 or OsHPP11. Protein prenylation can affect subcellular localisation, protein stability, and protein-protein interactions [61]. Several HIPPs have been found to localise to plasmodesmata, with the localisation being dependent on an intact CααX motif [29,32,62]. It is therefore plausible that OsHIPP21 differs in localisation and/or function from the three HPPs, and AVR-Pia interaction may also have different consequences. Further investigation is required to establish both the function of these proteins in planta, and the effect of AVR-Pia binding.

A foundation for precision engineering of effector targets for disease resistance

Pathogens require certain host factors to establish a successful infection, with the underlying genes typically referred to as S genes. Previous studies have shown that *OsHIPP05* (*Pi21*) and *OsHIPP20* are S genes for blast infection; *oshipp05* and *oshipp20* CRISPR-Cas9 knockout plants exhibited increased resistance. Further, a recent study reported that

overexpression of *HvHIPP43* increased susceptibility to *M. oryzae,* which suggests that
HvHIPP43 may also be a S gene [17,31,32,63–65]. In addition, other H(I)PPs have been reported to
be S genes for diverse pests and pathogens. *Arabidopsis thaliana* plants carrying loss-of-
function mutations in *AtHIPP27* or *AtHMAD* (also known as *AtHIPP40* [27]) exhibited reduced
susceptibility to the beet cyst nematode *Heterodera schachtii* [30] and the bacterial pathogen
Pseudomonas syringae [39], respectively. Silencing of *TaHIPP1* in wheat reduced sporulation
of the wheat stripe rust fungus *Puccinia striiformis f. sp. tritici* [37]. It is possible that *OsHPP09*,
OsHPP10, *OsHPP11* and/or *OsHIPP21* may also be S genes for rice blast disease, and
deletion of these genes may increase blast resistance.

While knocking out S genes has the potential to confer durable resistance, it can cause
undesirable pleiotropic effects on plant growth and development [66–69]. Advances in genome
editing technologies enable specific mutation of genes to alter amino acids in a target protein.
This precise approach reduces the likelihood of compromising the endogenous function of the
protein [70]. Furthermore, it was recently demonstrated that the virulence function of the MAX
effector Pwl2 depends on interaction with OsHIPP43 [32].The function of OsHPP09, and the
consequences of AVR-Pia binding, are unknown. However, preventing AVR-Pia binding to
OsHPP09 could compromise the function of the effector in promoting disease. In this study,
we have demonstrated that a single point mutation in OsHPP09, converting the glutamate in
position 16 to either alanine or arginine, is sufficient to prevent AVR-Pia binding. We also
found that OsHPP09 carrying an I35K mutation also lost interaction with AVR-Pia. However,
while we could produce and purify the recombinant HMA domain from *E. coli*, circular
dichroism experiments revealed that this mutation interfered with the correct folding of the
HMA domain, indicating that protein misfolding causes the observed loss of AVR-Pia binding.
This illustrates the importance of characterising the effects of mutations on protein structure
before commencing time-consuming and costly base editing experiments in plants.

In summary, we have characterised the interactions between AVR-Pia and a novel subset of
HMA domain-containing rice proteins. Biophysical and structural analyses guided
mutagenesis of these effector targets, and identified a mutation which blocks effector binding
without compromising structural integrity of the HMA domain. This provides a foundation for
targeted modification of these proteins for enhanced disease resistance.

Methods

Cloning

The coding sequences (CDS) of *OsH(I)PPs*, including the stop codon, were synthesised by Twist Bioscience. The CDS of *OsHIPP14*, *OsPi21*, *OsHIPP19*, *OsHIPP21* and *OsHPP09* were codon-optimized to enable synthesis.

For Y2H and split luciferase experiments, synthesised *OsH(I)PP* genes were used as template for amplification with primers containing *attB1* and *attB2* recombination sites (**Supplementary Table S6**) and Phusion High-Fidelity DNA Polymerase (Thermo Fisher Scientific). The resulting PCR products were subsequently introduced into the pDONR207 vector (Invitrogen) via Gateway® BP recombination (Thermo Fisher Scientific). Entry constructs for the MAX effectors dSP-AVR-Pia [13], dSP-AVR1-CO39 [10], dSP-AVR-PikD [44] and dSP-Pwl2 [11] were previously published. The dSP-AVR-PikC fragment was amplified from pCambia1300_dSP-AVR-PikC [50] and dSP-MAX58 from a synthesised gene, both using primers containing *attB1* and *attB2* recombination sites (Supplementary Table S6) and Phusion High-Fidelity DNA Polymerase (Thermo Fisher Scientific). The resulting PCR products were also cloned into the pDONR207 vector (Invitrogen) via Gateway® BP recombination (Thermo Fisher Scientific).

For recombinant protein production in *E. coli,* synthesised genes were used as templates for PCR amplification of the HMA domains (**Supplementary Table S1**) with primers containing *Bpi*I restriction sites (**Supplementary Table S6**) using CloneAmp™ HiFi PCR Premix (Clontech Laboratories). The resulting PCR products were introduced into the vector pICH41308 (Addgene #47998; [71]) by Golden Gate assembly [72] to generate level 0 CDS modules. The CDS of OsHIPP21-HMA was codon-optimised for expression in *E. coli* and supplied with the appropriate overhangs and restriction sites for direct assembly into the vector pICH41308 without PCR amplification. The level 0 CDS modules were used in Golden Gate reactions with the pOPIN-GG vector pPGN-C (Addgene #174578, [73]) and the level 0 module pICSL30015 (Addgene #174582, [73]) to generate *E. coli* expression constructs with a cleavable N-terminal 6xHis-MBP tag.

Site-directed mutagenesis of OsHPP09-HMA for Y2H, split luciferase, and recombinant protein production was carried out using a QuikChange Site-Directed Mutagenesis Kit (Agilent). To introduce mutations in the prenylation motif of OsHIPPs in the pDONR constructs to generate *OsHIPP14^{C187S}*, *OsHIPP21^{C133S}*, *OsHIPP39^{C190S}*, *OsHIPP41^{C152S},* and *AVR-Pia-H3*, site-directed mutagenesis PCR was carried out using Phusion High-Fidelity DNA Polymerase (Thermo Fisher Scientific) and mutation-specific primers listed in **Supplementary Table S6**.

Gateway® LR recombination using LR Clonase® II mix (Thermo Fisher Scientific) was used
to transfer *OsH(I)PPs* and *RGA5$_{Cter}$* [9] (883-1116 aa) into Gateway®-compatible destination
vectors, including a modified Y2H vector pGBKT7-GW [74] and the plant expression vector
pBIN-35S-3Flag-Cluc-GWY (Deslandes et al., unpublished). Similarly, MAX effector
pDONR constructs were recombined into a modified pGADT7-GW [74] and pBIN-35S-3HA-
Nluc-GWY (Deslandes et al., unpublished) destination vectors. The pBIN-35S-3HA-Nluc-
GWY and pBIN-35S-3Flag-Cluc-GWY vectors were generated by the ligation of a 3HA-Nluc-
FrameB or 3Flag-Cluc-FrameB fragment, respectively, into a XhoI/XbaI digested pBIN-35S
vector.

Previously published expression constructs for AVR1-CO39, AVR-Pia and AVR-Pia-H3 [4,10,11]
were used for protein production in the present study.

A complete list of plasmids used in this study is provided in **Supplementary Table S7.**

Yeast two-hybrid

The yeast strain Y2HGold (mating type a; Takara Bio) was transformed with the bait construct
expressed from the pGBKT7 vector, which contains the GAL4-BD and the *TRP1* marker gene.
These constructs included previously described HMA domains [47], as well as newly generated
constructs: OsHIPP21-HMA, OsH(I)PP^{WT} FL and OsH(I)PP$^{C/S}$ FL. The yeast strain Y187
(mating type α; Takara Bio) was transformed with the prey construct (MAX effectors)
expressed from the pGADT7 vector, which carries the GAL4-AD and the *LEU2* marker
gene. Yeast transformation was performed using a lithium acetate (LiAc)-based protocol
according to the Yeastmaker Yeast Transformation System (Takara Bio). Transformed yeast
cells were plated on synthetic defined (SD) media lacking either tryptophan (-W) or leucine (-
L) and incubated at 30 °C for 3 days.

Mating of the transformed yeast strains was carried out following the Matchmaker Gold Yeast
Two-Hybrid System User Manual (Takara Bio). Diploid yeast cells were selected on SD/-L-W
plates (SD medium lacking leucine and tryptophan) after incubation at 30 °C for 2 days. For
the Y2H assay, diploid yeast cells were inoculated in SD/-L-W medium, grown for 24 hours at
28°C with shaking at 100 rpm, then pelleted and washed twice with sterile water. Serial 10-
fold dilutions (10^{-1}, 10^{-2}, and 10^{-3}) were prepared and spotted onto SD selection plates: SD/-
L-W (growth control), SD/-L-W-H (lacking leucine, tryptophan, and histidine), SD/-L-W-H
supplemented with varying concentrations of 3-amino-1,2,4-triazole (3AT). Plates were
incubated at 30 °C for 7 days and subsequently photographed using the presentation station
UF-130DX (Samsung).

Yeast two-hybrid cDNA library screening

The Y2H cDNA library derived from the rice cultivar CO39, as well as the BD:dSP-AVR-Pia construct, have been previously described [9,44]. Library screening was conducted following the Matchmaker Gold Y2H system protocol (Takara Bio). Mating was performed between the Y2H Gold strain harbouring GAL4-BD:AVR-Pia and the Y187 strain carrying the CO39 cDNA library. Diploid yeasts were plated on SD/-L-W-H medium and supplemented with 20 mM 3AT. Colonies that grew were re-streaked onto fresh SD/-L-W-H medium containing 20 mM 3AT to confirm interaction-dependent growth. Yeast colony PCR was used to amplify the cDNA inserts from positive clones, and PCR products were sequenced. The resulting sequences were aligned to the Nipponbare rice genome using BLAST to identify the corresponding genes. Only genes for which the cDNA inserts (or fragments of cDNAs) were cloned in-frame with the GAL4-AD were retained as candidate AVR-Pia targets.

Agrobacterium-mediated transient expression in *N. benthamiana*

Agrobacterium tumefaciens strains AGL1 (carrying split-luciferase constructs) and GV3101-pMP90 (carrying 35S::P19) were grown overnight at 28°C in 5 mL LB medium supplemented with the appropriate antibiotics. Overnight cultures were pelleted at 4,000 x g for 10 minutes, and the pellets were resuspended in 2 mL infiltration buffer (10 mM $MgCl_2$, 10 mM MES pH 5.6, 150 µM acetosyringone). The bacterial suspensions were incubated in the dark for 2 hours. The optical density at 600 nm (OD_{600}) was adjusted to 0.05 for 35S::P19 and 0.1 for OsH(I)PPs and MAX effectors, respectively. In all infiltration mixes, 35S::P19 was co-infiltrated to suppress gene silencing. Agrobacterial suspensions were mixed accordingly and infiltrated into the leaves of 5–6-week-old *N. benthamiana* plants using a needleless syringe. Leaf discs were harvested 2 days post-infiltration for luminescence measurements or protein extraction. *N. benthamiana* plants were grown in a growth chamber on soil under a 16-hour light / 8-hour dark cycle, at 20 °C and 65 - 75% relative humidity.

Split luciferase assay

For luminescence measurements, 4-mm leaf discs were collected using a Miltex Biopsy Punch (Delta Microscopie) and individually placed into wells of a white 96-well plate containing water. The water was removed and replaced with a 1 mM Xenolight D-luciferin-K[+] salt bioluminescent substrate solution (Perkin Elmer). Luminescence was quantified using a Spark® microplate reader (Tecan) over three cycles of 10 minutes each. For all experiments, data from the third cycle were used for analysis. Each interaction was tested with eight technical replicates (eight leaf discs per plate from four individual plants per biological replicate, unless otherwise indicated in the figure legend). The luminescence assay was performed in three independent experiments. Statistical analyses from three independent replicates (typically involving 24 leaf

discs in total, unless otherwise indicated in the figure legend) and data visualization were done

using R (version 4.4.1). Multiple comparisons were performed using pairwise Wilcoxon tests

followed by a Benjamini-Hochberg p-value adjustment with the stats package. Boxplots were

generated using the ggplot2 package.

Protein extraction and Western blotting

Protein extraction from yeast was performed using a post-alkaline extraction protocol, as

described by Kushnirov [75]. For BD-OsHIPP14/19/20/39/41 and AD-MAX, yeast crude extracts

were resuspended in 50 µL of 1x LDS loading buffer (1x Bolt™ LDS sample buffer (Thermo

Fisher Scientific), 1x NuPAGE™ sample reducing agent (Thermo Fisher Scientific)), boiled at

95°C for 10 minutes, and loaded onto a 10% NuPAGE™ Bis-Tris gels (Invitrogen). For BD-

OsHPP09/10/11 and OsHIPP21, yeast crude extracts were resuspended in 50 µL of 1x SDS-

Tricine buffer (1x SDS-Tricine buffer (Thermo Fisher Scientific), 1x NuPAGE™ sample

reducing agent (Thermo Fisher Scientific)) and boiled at 85 °C for 2 minutes before loading

onto a 10-20% Tricine gel (Novax). Proteins were transferred to iBlot® nitrocellulose

membrane (Invitrogen) using the iBlot 2 Gel Transfer Device (program: 20 V 1 min, 23 V 4

min, 25 V, 2 min, Thermo Fisher Scientific), and analysed by immunoblotting.

For protein extraction from *N. benthamiana*, four leaf discs (0.8 mm diameter) were collected,

immediately frozen in liquid nitrogen and ground using a tissue homogenizer (Retsch GmbH).

The ground plant material was solubilized in 200 µL extraction buffer (20 mM Tris-Hcl pH 7.5,

150 mM NaCl, 1 mM EDTA pH 8.0, 1% Triton X-100, 0.1% SDS, 5 mM DTT, 1x Protease

inhibitor cocktail (Sigma-Aldrich), 1x EDTA-free cOmplete protease inhibitor (Roche)). Protein

extracts were centrifuged at 18,000 x *g* for 15 minutes at 4 °C, and the resulting supernatants

were collected. 2x LDS buffer (Bolt™ LDS sample buffer (Thermo Fisher Scientific) and

NuPAGE™ sample reducing agent (Thermo Fisher Scientific)) was added and samples were

denatured at 70 °C for 10 minutes. Proteins were then resolved by electrophoresis on

NuPAGE™ 4-12% Bis-Tris gels (Invitrogen). Proteins were transferred to iBlot® nitrocellulose

membrane (Invitrogen) using the iBlot 2 Gel Transfer Device (program: 20 V 1 min, 23 V 4

min, 25 V, 2 min, Thermo Fisher Scientific), and analysed by immunoblotting.

For immunodetection of proteins, rabbit anti-GAL4-BD (sc-577, Santa Cruz Biotechnology,

1:2,000), goat anti-rabbit IgG-HRP (1:20,000, Sigma-Aldrich, A0545), mouse anti-Flag M2-

HRP (1:60,000, Sigma-Aldrich) and rat anti-HA-HRP (clone 3F10, 1:1,1000, Roche) were

used. Chemiluminescence was detected using the Immobilon western kit (Millipore) or the

SuperSignal™ West Femto Maximum Sensitivity substrate (Thermo Fisher Scientific) and the

Syngene 680X EF GBOX imaging system (Syngene).

Protein production and purification

HMA domains

HMA domains were produced with a cleavable N-terminal 6xHis-MBP tag from *E. coli* BL21 (DE3) or BL21 (DE3) pLysS. 700 mL cell cultures were grown in LB media at 37 °C from a starting OD_{600} of 0.05-0.07 to an OD_{600} of 0.5-0.8. Protein production was induced by addition of IPTG (final concentration 1 mM) and cell cultures were grown for a further 14-16 hours at 20 °C. Cells were harvested by centrifugation and the pellets stored at -70 °C prior to purification. The pellets were then thawed and resuspended in 50 mM Tris-HCl pH 7, 300 mM NaCl, 30 mM imidazole supplemented with cOmplete EDTA-free protease inhibitor cocktail (Roche). Cells were lysed by sonication and the lysate clarified by centrifugation at 40,000 x *g* at 6 °C for 30 minutes to pellet insoluble material, then the supernatant was filtered (0.45 µm). All subsequent affinity chromatography and size exclusion chromatography steps were carried out with an ÄKTA Pure system (Cytiva) at 4 °C. The filtered supernatant was applied to a 5 ml HisTrap™ FF column (Cytiva) and, following washing with 50 mM Tris-HCl pH 7, 300 mM NaCl, 30 mM imidazole, bound protein was step-eluted with 50 mM Tris-HCl pH 7, 300 mM NaCl, 500 mM imidazole. The eluate was injected onto a Superdex 75 26/60 size exclusion chromatography column (Cytiva) pre-equilibrated in 50 mM Tris-HCl pH 7, 150 mM NaCl. Fractions containing the fusion protein were combined and incubated with 3C protease overnight at 4 °C. The HMA domains were separated from the 6xHis-MBP fusion tag and GST-tagged 3C protease by injecting the protease-treated sample onto a 5 ml HisTrap™ FF column (Cytiva), 5 ml MBPTrap™ HP column (Cytiva), and 5 ml GSTrap™ HP column (Cytiva) connected in tandem and equilibrated in 50 mM Tris-HCl pH 7, 300 mM NaCl, 30 mM imidazole. Fractions containing the HMA domains were pooled and concentrated to < 8 ml then injected onto a Superdex 75 26/60 size exclusion chromatography column (Cytiva) pre-equilibrated in 50 mM Tris-HCl pH 7, 150 mM NaCl. Fractions containing the purified HMA domains were combined and concentrated. At each stage, fractions containing the HMA domain were determined by SDS-PAGE. Samples were combined with 5X SDS loading buffer (60 mM Tris-HCl pH 6.8, 25% glycerol, 4% SDS, 0.1% bromophenol blue, 10 mM DTT) and heated at 95 °C for 5 minutes, then loaded onto a Bolt™ 10 %, Bis-Tris gel (Invitrogen). After migration, gels were stained in InstantBlue® Coomassie protein stain (Abcam). Final protein concentration was determined using a Pierce™ BCA Protein Assay Kit (ThermoFisher Scientific), due to the extreme molar extinction coefficients of the proteins (0 $M^{-1}cm^{-1}$ for OsHIPP21-HMA, 1490 $M^{-1}cm^{-1}$ for OsHPP-HMA domains) at 280 nm. Proteins were stored at -80 °C.

MAX effectors

AVR-Pia, AVR-Pia-H3 and AVR1-CO39 were produced and purified as previously described

[4,10,11]. Concentrations of the MAX effectors were determined from their molar extinction

coefficients and their absorbance at 280 nm.

Analytical size exclusion chromatography

Analytical size exclusion chromatography experiments were conducted using a Superdex 75

10/300 GL column (Cytiva) connected to an ÄKTA Pure system (Cytiva) at 4 °C. The running

buffer was 50 mM Tris pH 7, 150 mM NaCl, 1 mM DTT for experiments with AVR1-CO39 and

20 mM sodium citrate pH 5.6, 150 mM NaCl, 1 mM DTT for experiments with AVR-Pia and

AVR-Pia-H3. To investigate whether OsHPP09-HMA forms a complex with the effectors, the

two proteins were combined in a 1:1 molar ratio and incubated on ice for 1 h prior to analysis.

Each protein was also analysed alone, at a concentration equivalent to that present in the

mixture. For each experiment, 100 µl protein was injected onto the column. Elution was carried

out at a flow rate of 0.5 ml/min, and 500 µl fractions were collected for analysis by SDS-PAGE.

Samples were combined with 5X SDS loading buffer (60 mM Tris-HCl pH 6.8, 25% glycerol,

4% SDS, 0.1% bromophenol blue, 10 mM DTT) and heated at 95 °C for 5 minutes, then loaded

onto a Bolt™ 10 %, Bis-Tris gel (Invitrogen). After migration, gels were stained in InstantBlue®

Coomassie protein stain (Abcam).

Circular dichroism spectroscopy

Circular dichroism (CD) spectra were obtained using a Chirascan™ CD Spectrometer (Applied

Photophysics). The temperature of the sample compartment was set to 20 °C. Proteins were

dialysed into 10 mM sodium phosphate, pH 7.2 using Slide-A-Lyzer™ MINI dialysis devices

(Thermo Fisher Scientific). 40 µl protein was used to fill a 0.1 mm cuvette. Spectra were

obtained for wavelengths from 180 nm to 260 nm with a step size of 1 nm, bandwidth of 2 nm

and time-per-point of 2 s. Ten spectra were acquired for buffer and for each sample. The

average trace for the buffer was subtracted from the average trace for each protein. Units

were converted from millidegrees to mean residue molar ellipticity (MRME) using the equation:

981
$$MRME = \frac{millidegrees \times MRW}{10 \times L \times C}$$

982 where L = path length (cm), C = protein concentration (mg/ml), and MRW = mean residue

983 weight, determined as:

984
$$MRW = \frac{molar\ mass}{(number\ of\ amino\ acids - 1)}$$

Nuclear magnetic resonance titration

Samples of ^{15}N-labelled AVR-Pia (40 µM) alone and in a 1:1 molar ratio with unlabelled OsHPP09-HMA, or OsHPP09-HMA mutants, were prepared in 20 mM sodium citrate, pH 5.6, 150 mM NaCl and 1mM DTT. HSQC spectra were recorded at 305K on a Bruker Avance 800 MHz spectrometer equipped with a triple resonance (^{1}H, ^{15}N, ^{13}C) z-gradient cryo-probe. Chemical shift differences were measured from the HSQC spectra of AVR-Pia alone and the AVR-Pia/OsHPP09-HMA complex. All spectra are referenced to the internal reference DSS for the ^{1}H dimension and indirectly referenced for the ^{15}N dimension [76]. Spectra were processed using Topspin (v. 3.2) and analyzed using strip-plots with Cindy in house software and CCPN [77] [analysis v 2.3]. The ^{1}H, ^{15}N assignments were derived by analogy from the assignments of AVR-Pia wildtype. All the NMR samples were prepared with addition of 10% D_2O and 1 µM DSS as a reference.

Isothermal titration calorimetry

ITC experiments were performed using a MicroCal PEAQ-ITC (Malvern Panalytical). Proteins were dialysed overnight into the experiment buffer (20 mM sodium citrate pH 6.0, 150 mM NaCl, 1 mM TCEP). Experiments were carried out 25 °C with a reference power of 5 µcal/s. The calorimetric cell was filled with the purified HMA domain at 18 µM (OsHPP09-HMA), 35 µM (OsHIPP21-HMA and OsHPP11-HMA) or 60 µM (OsHPP10-HMA). This was titrated with AVR-Pia at a concentration 10X greater than that of the HMA domain (180 uM, 350 uM or 600 uM). Each experiment consisted of a single injection of 0.5 µL followed by 18 injections of 2 µL each at 150 s intervals with a stirring speed of 750 rpm. Data were processed with the MicroCal PEAQ-ITC analysis software (Malvern Panalytical). Experiments were carried out in triplicate.

Crystallisation of OsHPP09-HMA and AVR-Pia

Sitting drop, vapour diffusion crystallisation trials were set up in 96-well Swissci 3 lens crystallisation plates using dragonfly and mosquito dispensing robots (SPT Labtech). 100 nL of protein at a concentration of 8.6 mg/ml was combined with 100 nL reservoir solution. Needle-like crystals were observed in multiple conditions in the commercial Structure Screen 1 + 2 (HT-96; Molecular Dimensions) and PEGs Suite (NeXtal) screens, however these were unsuitable for X-ray diffraction. Optimisation around these conditions yielded multiple crystals suitable for diffraction studies. The X-ray diffraction data used to solve the structure was obtained from a crystal harvested from the condition containing 0.1 M MES pH 6.5, 0.2 M ammonium sulphate, 20% (w/v) PEG 4000. The crystal was mounted in a nylon loop and flash-frozen in liquid nitrogen prior to shipping to the European Synchrotron Radiation Facility.

X-ray diffraction data collection and processing and model refinement

X-ray diffraction data were collected on beamline MASSIF-1 (ID30A-1) at the European Synchrotron Radiation Facility. Data were processed with the EDNA autoprocessing pipeline [78]. The scaled, unmerged data file was passed to AIMLESS for data reduction [79,80] (implemented in CCP4i2 [81]). The AVR-Pia/OsHPP09-HMA structure was solved by molecular replacement with PHASER [82], using a monomer of AVR-Pia (PDB accession 6Q76, chain B) and a monomer of OsHPP09-HMA (modelled with ColabFold [83]). Iterative cycles of manual adjustment, refinement, and validation were carried out using COOT [84], REFMAC5 [85,86] and MolProbity [87,88], implemented in CCP4i2. Analysis of the interface was performed with qtPISA [52]. Interatomic distances were measured and structure superposition/RMSD calculations were performed with PyMOL v2.5.0 Open-Source [89]. Structure figures were prepared using the CCP4 molecular graphics (CCP4mg) software [90].

Structural modelling of AVR-Pia/HMA complexes

Structural modelling of AVR-Pia/HMA complexes was carried out with both ColabFold v1.5.5 ([83] (based on AlphaFold2 Multimer v3 [91,92]) and the AlphaFold3 web server (alphafoldserver.com; models generated between 2024-05-27 and 2024-07-02) [93]. For models generated with ColabFold, no template information from the PDB was used (template_mode: none) and the number of recycles was set to 20 (num_recycles: auto). Five models were generated for each complex and were relaxed using AMBER (num_relax: 5). Default settings were otherwise used. For models generated with AlphaFold3, default settings were used. At the time of use, AlphaFold3 did not support excluding template information from the PDB. Five models were generated for each complex using five different seed values. Interface analysis was carried out with qtPISA [52].

Data availability

The protein structure of the complex between OsHPP09-HMA and AVR-Pia has been deposited in the Protein Data Bank with accession number 9RSV (pdb_00009RSV). X-ray diffraction data are publicly available via the European Synchrotron Radiation Facility [94].

Figure Legends

Fig. 1. AVR-Pia interacts with OsH(I)PPs.

a Y2H interaction analysis between AVR-Pia (upper panel) or MAX58 (lower panel) and either the heavy metal-associated (HMA) domain or full-length (FL) proteins of OsH(I)PPs. The OsH(I)PP candidates were previously identified in a comprehensive pairwise Y2H analysis using AVR-Pia as prey and a library of HMA domains from OsH(I)PPs as bait. OsRGA5$_{Cter}$ (883 – 1116 aa) was included as a positive control in the experiment with AVR-Pia/OsH(I)PPs (top panel). After an interaction was observed between OsHPP09 and AVR-Pia, this interaction was used as a positive control in the subsequent experiment with MAX58 (bottom panel).. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH and SD/-LWH supplemented with 0.5 mM 3AT). Photos were taken after 7 days of incubation. Ø indicates positions where no yeast was spotted. AD, activating domain; BD, binding domain. **b** Quantitative measurement of luciferase-derived signals shows that AVR-Pia, but not MAX58, interacts with OsHPP09, OsHPP10, OsHPP11 and OsHIPP21 in a split-luciferase complementation assay. Neither AVR-Pia nor MAX58 interact with any other tested OsHIPP in planta. AVR-Pia and MAX58 were fused to the N-terminal part of luciferase (3xHA:NLuc:AVR-Pia, 3xHA:NLuc:MAX58) and co-expressed with OsH(I)PP FL proteins fused at their N-termini to the C-terminal part of luciferase (3xFlag:CLuc:OsH(I)PP). Co-expression of AVR-Pia with OsRGA5$_{Cter}$ (883 – 1116 aa) was used as a positive control. Leaves infiltrated with P19 alone served as background (BG) control. Leaf discs were collected 2 days post infiltration. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range. Values outside this range are shown as individual data points. Three independent replicates were performed (n = 8 per replicate). Due to experimental constraints, the following sample sizes apply to specific conditions: AVR-Pia/OsHPP09: n = 16 in replicates 1 and 2. Data points are individually plotted and shaped according to the replicate. Different letters indicate statistically significant differences based on a pairwise Wilcoxon test ($\alpha = 0.05$).

Fig. 2. AVR-Pia-H3 does not interact with OsHPP09, OsHPP10, OsHPP11 and HIPP21.

a Y2H interaction analysis between AVR-Pia-H3 and either the heavy metal-associated (HMA) domain or full-length (FL) proteins of OsH(I)PPs. OsHPP09/AVR-Pia served as positive control. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH). Photos were taken after 7 days of incubation. AD, activating domain; BD, binding domain. **b** Quantitative measurement of luciferase-derived signals shows that AVR-Pia-H3 does not interact with

OsHPP09, OsHPP10, OsHPP11 and OsHIPP21 in a split-luciferase complementation assay. MAX effectors were fused to the N-terminal part of luciferase (3xHA:NLuc:MAX) and co-expressed with OsH(I)PP FL proteins fused at their N-termini to the C-terminal part of luciferase (3xFlag:CLuc:OsH(I)PP). Co-expression of AVR-Pia with OsHPP09 served as positive control, co-expression of OsHPP09/MAX58 and OsRGA5$_{Cter}$ (883 – 1116 aa)/AVR-Pia-H3 served as negative controls. Leaves infiltrated with P19 alone served as background (BG) control. Leaf discs were collected 2 days post infiltration. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range. Values outside this range are shown as individual data points. Three independent replicates were performed, with n = 8 per combination per replicate. Due to experimental constraints, the following sample sizes apply to specific conditions: AVR-Pia/OsHPP09: n = 16 in replicates 1 and 2. Data points are individually plotted and shaped according to the replicate. Different letters indicate statistically significant differences based on a pairwise Wilcoxon test (α = 0.05). **c** Analytical gel filtration traces obtained from injection of OsHPP09-HMA alone (top panel), AVR-Pia-H3 alone (middle panel) and the two proteins in a 1:1 molar ratio (bottom panel). Significant peaks are indicated by coloured dashed lines with the elution volume labelled. SDS-PAGE gel inserts show fractions from peak elution volumes indicated by the grey shaded regions. OsHPP09-HMA absorbs light at 280 nm poorly (molar extinction coefficient of 1490 $M^{-1}cm^{-1}$) so the peak corresponding to OsHPP09-HMA is small. The elution volume observed for OsHPP09-HMA is consistent with dimerisation, as has been observed for other purified HMA domains [10,50]. There is no shift in the elution volume for AVR-Pia-H3 in the presence of OsHPP09-HMA indicating that the two proteins do not form a complex.

Fig. 3. OsH(I)PPs interacting with AVR-Pia do not interact with AVR1-CO39.

a Y2H interaction analysis between AVR1-CO39 and either the heavy metal-associated (HMA) domain or full-length (FL) proteins of OsH(I)PPs. OsRGA5$_{Cter}$ (883 – 1116 aa)/AVR1-CO39 served as positive control. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH). Photos were taken after 7 days of incubation. Ø indicates positions where no yeast was spotted. AD, activating domain; BD, binding domain. **b** Quantitative measurement of luciferase-derived signals shows that AVR1-CO39 does weakly interact with OsHPP09, OsHPP10 and OsHPP11 but not with OsHIPP21 in a split-luciferase complementation assay. MAX effectors were fused to the N-terminal part of luciferase (3xHA:NLuc:MAX) and co-expressed with OsH(I)PP FL proteins fused at their N-termini to the C-terminal part of luciferase (3xFlag:CLuc:OsH(I)PP). Co-expression of AVR-Pia or AVR1-CO39 with OsPi21

served as negative controls, while AVR-Pia/OsHPP09 served as a positive control. Leaf discs
were collected 2 days post infiltration. Box plots show the median (line), mean (cross) and
upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x
the interquartile range. Values outside this range are shown as individual data points. Three
independent replicates were performed, with n = 8 per combination per replicate. Data points
are individually plotted and shaped according to the replicate. Different letters indicate
statistically significant differences based on a pairwise Wilcoxon test (α = 0.05). **c** Analytical
gel filtration traces obtained from injection of OsHPP09-HMA alone (top panel), AVR1-CO39
alone (middle panel) and the two proteins in a 1:1 molar ratio (bottom panel). Significant peaks
are indicated by coloured dashed lines with the elution volume labelled. SDS-PAGE gel inserts
show fractions from peak elution volumes indicated by the grey shaded regions. OsHPP09-
HMA absorbs light at 280 nm poorly (molar extinction coefficient of 1490 $M^{-1}cm^{-1}$) so the peak
corresponding to OsHPP09-HMA is small. The small peak at ~13.7 ml in the trace for AVR1-
CO39 is likely due to some dimerisation of the AVR1-CO39 protein; the quantity of protein is
insufficient to observe a corresponding band on the SDS-PAGE gel. The elution volume
observed for OsHPP09-HMA is consistent with dimerisation, as has been observed for other
purified HMA domains [10,50]. There is no shift in the elution volume for AVR1-CO39 in the
presence of OsHPP09-HMA indicating that the two proteins do not form a complex.

1137

Fig. 4. OsH(I)PPs interacting with AVR-Pia do not interact with AVR-Pik or Pwl2.

a Y2H interaction analysis between AVR-PikC, AVR-PikD, Pwl2 and AVR-Pia and either the
heavy metal-associated (HMA) domain or full-length (FL) proteins of OsH(I)PPs.
OsHIPP19/AVR-PikC, OsHIPP20/AVR-PikC, OsHIPP19/AVR-PikD, OsHIPP20/AVR-PikD
and OsHIPP43/Pwl2 served as positive controls. Serial dilutions of diploid yeast were spotted
onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein
interactions (SD/-LWH). Photos were taken after 7 days of incubation. AD, activating domain;
BD, binding domain. **b** Quantitative measurement of luciferase-derived signals shows that
AVR-PikC and AVR-PikD do not interact with OsHPP09, OsHPP10, OsHPP11 and
OsHIPP21, while Pwl2 shows weak luminescence signals with these proteins, suggesting
weak interactions. AVR-Pia also interacts with OsHIPP19 and OsHIPP20 in planta. MAX
effectors were fused to the N-terminal part of luciferase (3xHA:NLuc:MAX) and co-expressed
with OsH(I)PP FL proteins fused at their N-termini to the C-terminal part of luciferase
(3xFlag:CLuc:OsH(I)PP). Co-expression of AVR-PikC/AVR-PikD with OsHIPP19/20 and Pwl2
with OsHIPP43 served as positive controls, AVR-Pia with OsHIPP43 served as negative
control. Leaves infiltrated with P19 alone served as background (BG) control. Leaf discs were
collected 2 days post infiltration. Box plots show the median (line), mean (cross) and

upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x

the interquartile range. Values outside this range are shown as individual data points. Unless

otherwise stated, three independent replicates were performed, with n = 8 per combination

per replicate. Due to experimental constraints, the following sample sizes apply to specific

conditions: AVR-Pia/OsHIPP19 and AVR-Pia/OsHIPP20 in AVR-PikC and AVR-PikD panels:

n = 4 in replicates 1 and 3. BG control in AVR-PikC and AVR-PikD panels: n = 3 in replicates

1 and 3 and BG control in Pwl2 panel: n = 2 in replicates 1 and 3. AVR-PikD/OsHIPP19 n =

16 in replicates 1 and 3 and n = 24 in replicate 2. Data points are individually plotted and

shaped according to the replicate. Different letters indicate statistically significant differences

based on a pairwise Wilcoxon test (α = 0.05).

1165

Fig. 5. AVR-Pia binds the HMA domains of OsH(I)PPs with varying affinities.

a Representative ITC experiments for each of the AVR-Pia/HMA interactions tested. Top panel

shows the raw thermogram obtained from titration of AVR-Pia into a solution containing the

purified HMA domains. Central panel shows the integrated heats (coloured dots) and binding

isotherm fitted to a single site model (black line) using the MicroCal PEAQ-ITC analysis

software (Malvern Panalytical). Bottom panel shows the differences (coloured crosses)

between the modelled and observed values (residuals). **b** Dot plot representation of the

apparent KD for each of the AVR-Pia/HMA interactions in three independent experiments.

Coloured dots represent the KD vales determined in each replicate with coloured lines

representing the error associated with the KD value as determined by the MicroCal PEAQ-

ITC analysis software (Malvern Panalytical).

1177

Fig. 6. Crystal structure of the HMA domain of OsHPP09 in complex with AVR-Pia.

The structures of OsHPP09-HMA and AVR-Pia are represented as teal and gold ribbons,

respectively. The amino acids comprising β2 of AVR-Pia and β2 of the HMA domain are

represented as cylinders. Side chains involved in intermolecular hydrogen bonds/salt bridges

are also represented as cylinders. Hydrogen bonds are represented by black dashed lines

labelled with the bond length. Left insert shows main chain hydrogen bonds between AVR-

Pia-β2 and OsHPP09-β2. Right insert shows intermolecular contacts between side chains of

residues in OsHPP09-α1 and AVR-Pia.

1186

Fig. 7. Mutation of OsHPP09 E16 reduces AVR-Pia binding.

a Y2H interaction analysis between AVR-Pia and full-length (FL) OsHPP09 mutant versions carrying single mutations in the binding interface. OsHPP09^{WT}/AVR-Pia served as positive control. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH and SD/-LWH supplemented with 5 mM 3AT). Photos were taken after 7 days of incubation. AD, activating domain; BD, binding domain. **b** Quantitative measurement of luciferase-derived signals shows that the E16A, E16R and I35K mutations in OsHPP09 significantly reduce its interaction with AVR-Pia in a split-luciferase complementation assay. AVR-Pia was fused to the N-terminal part of luciferase (3xHA:NLuc:AVR-Pia) and co-expressed with OsH(I)PP FL proteins fused at their N-termini to the C-terminal part of luciferase (3xFlag:CLuc:OsH(I)PP). Co-expression of AVR-Pia with OsHPP09 served as positive control, co-expression of OsHIPP43 with AVR-Pia served as negative control. Leaves infiltrated with P19 alone served as background (BG) control. Leaf discs were collected 2 days post infiltration. Box plots show the median (line), mean (cross) and upper/lower quartiles (box limits), whiskers extend to the most extreme data points within 1.5x the interquartile range. Values outside this range are shown as individual data points. Three independent replicates were performed, with n = 8 per combination per replicate. Data points are individually plotted and shaped according to the replicate. Different letters indicate statistically significant differences based on a pairwise Wilcoxon test (α = 0.05). **c** Circular dichroism (CD) spectra obtained for WT OsHPP09-HMA (pale blue) and variants carrying E16A, E16R, I35K or D33H mutations. Negative peaks at 222 nm and 208 nm and a positive peak at 193 nm are indicative of α-helices while a negative peak at 218 nm and positive peak at 195 nm are indicative of β-sheets, consistent with the expected βαββαβ topology of an HMA domain. In the case of the I35K mutant, low ellipticity above 210 nm and a negative peak at 200 nm is indicative of disorder. **d** Overlays of cross-peaks from HSQC spectra corresponding to residues L39, I44, E56 and Y85 of ^{15}N-labelled AVR-Pia alone (orange) and in the presence of either WT OsHPP09-HMA or variants carrying E16A, E16R or D33H mutations (blue/green/purple). Overlays of the full spectra are provided in Supp. Figures S19-S21. Chemical shift perturbation of ^{15}N-labelled AVR-Pia is indicative of an interaction with the HMA domain.

Supplementary Figure and Table Legends

Supplementary Fig. 1. AVR-Pia interacts with HMA domains from OsH(I)PPs.

Comprehensive pairwise Y2H analysis using AVR-Pia, MAX58, Pwl2, and AVR-PikD as prey, and a library of HMA domains from OsH(I)PPs as bait [47]. The left panel shows yeast growth on selective media, while the right panel identifies the rice H(I)PP proteins from which the HMA domains were derived. Diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH). Pictures were taken after 7 days of incubation. AD, activating domain; BD, binding domain; *, Autoactive BD construct leading to yeast growth regardless of the presence or identity of the AD construct; +++, strong yeast growth; ++, normal yeast growth, +, weak yeast growth; No HMA, proteins for which no HMA domain was identified by sequence analysis or structural prediction; Wrong annotation, constructs based on misannotated genes, resulting in missing or truncated HMA domains, thereby compromising Y2H assay.

Supplementary Fig. 2. Mutation of the isoprenylation motif of OsHIPPs does not promote AVR-Pia interaction.

Y2H interaction analysis between AVR-Pia and either wildtype (WT) full-length (FL) OsHIPPs, including OsHIPP14, OsHIPP21, OsHIPP39 and OsHIPP41, or the corresponding FL OsHIPP isoprenylation mutants, in which the cysteine (C) residue was replaced by a non-prenylatable serine (S) residue. OsHPP09/AVR-Pia and was included as positive control. Serial dilutions of diploid yeast were spotted onto synthetic defined (SD) media to monitor growth (SD/-LW) or to assess protein-protein interactions (SD/-LWH). Photos were taken after 7 days of incubation. AD, activating domain; BD, binding domain. OsHIPP$^{C/S}$ mutants: OsHIPP14^{C187S}, OsHIPP21^{C133S}, OsHIPP39^{C190S}, OsHIPP41^{C152S}.

Supplementary Fig. 3. Expression of bait and prey proteins in haploid yeast.

a Total yeast protein extracts from haploid yeast were separated on a 10 – 20% Tricine SDS-PAGE gel and analysed by immunoblot using an anti-BD antibody to detect Gal4-BD fusion proteins with either the heavy metal-associated (HMA) domain, full-length (FL) OsH(I)PPs and the OsHIPP21 isoprenylation mutant (C/S). Protein loading is indicated by Ponceau S staining (PS). **b** Total yeast protein extracts from haploid yeast were separated on a 10% NuPAGE™ Bis-Tris gel and analysed by immunoblot using an anti-BD antibody to detect Gal4-BD fusion proteins with either the HMA domain, FL OsHIPPs and their corresponding isoprenylation mutants (C/S). Protein loading is indicated by Ponceau S staining (PS). **c** Total yeast protein extracts from haploid yeast were separated on a NuPAGE™ Bis-Tris gel and analysed by immunoblot using an anti-HA antibody to detect Gal4-AD fusion proteins with MAX effectors.

Protein loading is indicated by Ponceau S staining (PS). **d** Total yeast protein extracts from

haploid yeast were separated on a 10% NuPAGE™ Bis-Tris gel and analysed by immunoblot

using an anti-BD antibody to detect Gal4-BD fusion proteins with FL OsHPP09 wildtype (WT)

and mutant versions. Protein loading is indicated by Ponceau S staining (PS). OsHIPP$^{C/S}$

mutants: OsHIPP14^{C187S}, OsHIPP21^{C133S}, OsHIPP39^{C190S}, OsHIPP41^{C152S}.

Supplementary Fig. 4. Amino acid alignment of OsHPP09, OsHPP10, OsHPP11,

OsHIPP21 and the C-terminus of OsRGA5.

Amino acid alignment of full-length OsHPP09, OsHPP10, OsHPP11 and OsHIPP21 and

residues 995-1116 of OsRGA5, including the HMA domain and the C-terminal region of the

protein which concludes with a putative CaaX isoprenylation motif (CSTM). Sequence

alignment was carried out with Clustal Omega. The blue shaded box highlights the HMA

domain, while the yellow shaded box highlights the CaaX motif present in OsHIPP21 and

OsRGA5.

Supplementary Fig. 5. Expression of CLuc- and NLuc fusion proteins (related to Fig. 1).

Immunoblot analysis of transiently expressed proteins in *N. benthamiana*, including

OsH(I)PPs N-terminally tagged with a 3xFlag epitope fused to the C-terminal part of luciferase

(CLuc), and AVR-Pia and MAX58 N-terminally tagged with 3xHA and the N-terminal part of

luciferase (NLuc). This experiment corresponds to replicate 1 of Fig. 1. Detection was

performed using anti-Flag and anti-HA antibodies. Membrane was stripped after anti-Flag

detection and re-probed with anti-HA. Protein loading is indicated by the Rubisco band

visualized by Ponceau S. staining (PS).

Supplementary Fig. 6. AlphaFold2 models do not differentiate between HMA domains

which interact with AVR-Pia and those which do not.

AlphaFold2 (ColabFold v1.5.5) models of AVR-Pia in complex with HMA domains which **a**

were experimentally shown to interact with AVR-Pia and **b** do not interact with AVR-Pia.

Structure models are represented as ribbons and coloured by pLDDT score using the classical

AlphaFold colour scheme as shown in the key. **c** Plots of ipTM, pTM and Multimer (0.8*iPTM

+ 0.2*pTM) scores for each of the five models generated for each HMA/AVR-Pia complex.

Points indicate scores of individual models; green and purple lines indicate mean values for

AVR-Pia-interacting and non-interacting HMA domains, respectively. **d** Comparison of

interface parameters determined by qtPISA [52]) for the top ranked model of each HMA/AVR-

Pia complex. Green and purple points indicate values for individual HMA/AVR-Pia models.

The centre line of the box represents the median and the limits of the box represent the upper

and lower quartiles. Whiskers extend to the smallest value within (Q1 − 1.5 × the interquartile

range (IQR)) and the largest value within (Q3 + 1.5 × IQR).

1287 Supplementary Fig. 7. AlphaFold3 models do not clearly differentiate between HMA

1288 domains which interact with AVR-Pia and those which do not.

AlphaFold3 models of AVR-Pia in complex with HMA domains which **a** were experimentally

shown to interact with AVR-Pia and **b** do not interact with AVR-Pia. Structure models are

represented as ribbons and coloured by pLDDT score using the classical AlphaFold colour

scheme as shown in the key. **c** Plots of ipTM, pTM and Multimer (0.8*iPTM + 0.2*pTM) scores

for each of the top ranked models from each of the five seeds used to model each HMA/AVR-

Pia complex. Points indicate scores of individual models; green and purple lines indicate mean

values for AVR-Pia-interacting and non-interacting HMA domains, respectively. **d** Comparison

of interface parameters determined by qtPISA [52]) for the top ranked model (selected from

models from all five seeds) of each HMA/AVR-Pia complex. Green and purple points indicate

values for individual HMA/AVR-Pia models. The centre line of the box represents the median

and the limits of the box represent the upper and lower quartiles. Whiskers extend to the

smallest value within (Q1 − 1.5 × the interquartile range (IQR)) and the largest value within

(Q3 + 1.5 × IQR).

1302 Supplementary Fig. 8. AVR-Pia H3 differs from AVR-Pia in two amino acid positions.

a Multiple sequence alignment of AVR-Pia variants from Magnaporthe oryzae isolates

described in [6]. Signal peptide is highlighted by a shaded grey box. Polymorphic residues 24

(F, S) and 46 (T, N) are indicated in a purple outlined box. Arrows above the alignment indicate

secondary structure elements. **b** The polymorphic residues F24 and T46 are located at the

HMA binding interface. Crystal structure of AVR-Pia in complex with Pikp-HMA (PDB 6Q76;

[45]). AVR-Pia and Pikp-HMA are represented as gold and teal ribbons, respectively, with the

side chains of AVR-Pia T46 and F24S shown in purple.

1310 Supplementary Fig. 9. Expression of CLuc- and NLuc fusion proteins (related to Fig. 2).

Immunoblot analysis of transiently expressed proteins in *N. benthamiana*, including

OsH(I)PPs and OsRGA5$_{Cter}$ (883 – 1116 aa) N-terminally tagged with a 3xFlag epitope fused

to the C-terminal part of luciferase (CLuc), and AVR-Pia-H3 N-terminally tagged with 3xHA

and the N-terminal part of luciferase (NLuc). This experiment corresponds to replicate 2 of Fig.

5. Detection was performed using anti-Flag and anti-HA antibodies. Membrane was stripped

after anti-Flag detection and re-probed with anti-HA. Protein loading is indicated by the

Rubisco band visualized by Ponceau S. staining (PS).

Supplementary Fig. 10. AVR-Pia binds to the HMA domain of OsHPP09 in analytical size exclusion chromatography experiments.

Analytical gel filtration traces obtained from injection of OsHPP09-HMA alone (top panel), AVR-Pia alone (middle panel) and the two proteins in a 1:1 molar ratio (bottom panel). Significant peaks are indicated by dashed lines with the elution volume labelled. SDS-PAGE gel inserts show fractions from the peak elution volumes indicated by the grey shaded regions. OsHPP09-HMA absorbs light at 280 nm poorly (molar extinction coefficient of 1490 M-1cm-1) so the peak corresponding to OsHPP09-HMA is small. The elution volume observed for OsHPP09-HMA is consistent with dimerisation, as has been observed for other purified HMA domains [10,50]. The shift in the elution volume for AVR-Pia in the presence of OsHPP09-HMA indicates that the two proteins form a complex.

Supplementary Fig. 11. Expression of CLuc- and NLuc fusion proteins (related to Fig. 3).

Immunoblot analysis of transiently expressed proteins in *N. benthamiana*, including OsH(I)PPs and OsRGA5$_{Cter}$ (883 – 1116 aa) N-terminally tagged with a 3xFlag epitope fused to the C-terminal part of luciferase (CLuc), and AVR1-CO39 N-terminally tagged with 3xHA and the N-terminal part of luciferase (NLuc). This experiment corresponds to replicate 1 of Fig. 4. Detection was performed using anti-Flag and anti-HA antibodies. Membrane was stripped after anti-Flag detection and re-probed with anti-HA. Protein loading is indicated by the Rubisco band visualized by Ponceau S. staining (PS).

Supplementary Fig. 12. Expression of CLuc- and NLuc fusion proteins (related to Fig. 4).

Immunoblot analysis of transiently expressed proteins in *N. benthamiana*, including OsH(I)PPs N-terminally tagged with a 3xFlag epitope fused to the C-terminal part of luciferase (CLuc), and AVR-Pia, AVR-PikC, AVR-PikD and Pwl2 N-terminally tagged with 3xHA and the N-terminal part of luciferase (NLuc). This experiment corresponds to replicate 1 of Fig. 2. Detection was performed using anti-Flag and anti-HA antibodies. Membrane was stripped after anti-Flag detection and re-probed with anti-HA. Protein loading is indicated by the Rubisco band visualized by Ponceau S. staining (PS).

Supplementary Fig. 13. Production and purification of OsHPP09-HMA.

a Size exclusion chromatography elution trace for 6xHis-MBP-OsHPP09-HMA (following the initial IMAC purification step). The SDS-PAGE gel shows fractions corresponding to the peak in the shaded area. **b** SDS-PAGE gel of pooled fractions following size exclusion chromatography before and after cleavage of the 6xHis-MBP tag with GST-tagged 3C

protease. **c** SDS-PAGE gel showing flow-through, wash and elution fractions from tandem

HisTrapTM, MBPTrapTM and GSTrapTM affinity columns. **d** Size exclusion chromatography

elution trace for OsHPP09-HMA. OsHPP09-HMA absorbs light at 280nm poorly (molar

extinction coefficient of 1490 M^{-1}cm^{-1}). The SDS-PAGE gel shows fractions corresponding to

the three peaks in the shaded areas. The peak elution volume of OsHPP09-HMA (193.9 ml)

is consistent with dimerisation of the HMA domain.

Supplementary Fig. 14. Replicates of the ITC experiments presented in Fig. 5.

Top panels show the raw thermograms obtained from titration of AVR-Pia into a solution

containing the purified HMA domains. Central panels show the integrated heats (coloured

dots) and binding isotherms fitted to a single site model (black line) using the MicroCal PEAQ-

ITC analysis software (Malvern Panalytical). Bottom panels show the differences (coloured

crosses) between the modelled and observed values (residuals).

Supplementary Fig. 15. Thermodynamic profiles from ITC experiments presented in

Fig. 5 and Supp. Fig. S14.

Bars represent the magnitude of the determined thermodynamic parameters ΔG (blue bar),

ΔH (green bar) and -TΔS (red bar).

Supplementary Fig. 16. Comparison of binding interfaces in the crystal structures of

AVR-Pia in complex with OsHPP09-HMA (PDB accession code 9RSV), AVR-Pia/Pikp-

HMA (6Q76, [45]) and AVR1-CO39/RGA5-HMA (5ZNG,[10])

a Global interface of the effector/HMA complex. Structures are represented as gold (AVR-

Pia), teal (OsHPP09-HMA), brown (AVR-Pia), pale blue (Pikp-HMA), orange (AVR1-CO39)

and green (RGA5-HMA) ribbons with relevant secondary structure elements indicated. **b**

Hydrogen bonds between residues in β2 of the effector and β2 of the HMA domain. Main chain

atoms are represented as cylinders (side chains not shown for clarity). Hydrogen bonds are

represented as black dashed lines with lengths (determined by qtPISA) indicated. **c-f**

Comparison of residues involved in forming intermolecular contacts at the effector/HMA

interface. Structures are presented in ribbon representation with relevant residues shown as

cylinders. Hydrogen bonds are represented as black dashed lines with lengths (determined

by qtPISA) indicated. Water molecules are represented as red spheres.

Supplementary Fig. 17. HSQC spectra of 15N-AVR-Pia (40 µM) in the absence (orange

cross-peaks) and presence (purple cross-peaks) of unlabelled OsHPP09-HMA (40 µM).

Cross-peak labels correspond to the ^{15}N-AVR-Pia spectra (previously assigned by [11]). Zoom

inset panels show cross-peaks corresponding to residues L39, I44, E56 and Y85 to illustrate

the chemical shift perturbations between the free (^{15}N-AVR-Pia alone) and bound (^{15}N-AVR-
Pia with OsHPP09-HMA). Zoomed areas are indicated by boxes on the spectra.

Supplementary Fig. 18. Expression of CLuc- and NLuc fusion proteins (related to Fig. 7).

Immunoblot analysis of transiently expressed proteins in *N. benthamiana*, including OsHPP09
wildtype (WT) and mutant versions N-terminally tagged with a 3xFlag epitope fused to the C-
terminal part of luciferase (CLuc), and AVR-Pia N-terminally tagged with 3xHA and the N-
terminal part of luciferase (NLuc). This experiment corresponds to replicate 2 of Fig. 7.
Detection was performed using anti-Flag and anti-HA antibodies. Membrane was stripped
after anti-Flag detection and re-probed with anti-HA. Protein loading is indicated by the
Rubisco band visualized by Ponceau S. staining (PS). Dashed lines between bands indicate
that lanes were cut from the same blot but were non-adjacent.

Supplementary Fig. 19. HSQC spectra of ^{15}N-AVR-Pia (40 µM) in the absence (orange cross-peaks) and presence (turquoise cross-peaks) of unlabelled OsHPP09^{D33H}-HMA (40 µM).

1400 Cross-peak labels correspond to the ^{15}N-AVR-Pia spectra (previously assigned [11]). Zoom inset
1401 panels show cross-peaks corresponding to residues L39, I44, E56 and Y85 to illustrate the
1402 chemical shift perturbations between the free (^{15}N-AVR-Pia alone) and bound (^{15}N-AVR-Pia
1403 with OsHPP09^{D33H}-HMA) as presented in Fig. 7. Zoomed areas are indicated by boxes on the
1404 spectra.

Supplementary Fig. 20. HSQC spectra of ^{15}N-AVR-Pia (40 µM) in the absence (orange cross-peaks) and presence (blue cross-peaks) of unlabelled OsHPP09^{E16A}-HMA (40 µM).

1407 Cross-peak labels correspond to the ^{15}N-AVR-Pia spectra (previously assigned [11]). Zoom inset
1408 panels show cross-peaks corresponding to residues L39, I44, E56 and Y85 to illustrate the
1409 chemical shift perturbations between the free (^{15}N-AVR-Pia alone) and bound (^{15}N-AVR-Pia
1410 with OsHPP09^{E16A}-HMA) as presented in Fig. 7. Zoomed areas are indicated by boxes on the
1411 spectra.

Supplementary Fig. 21. HSQC spectra of ^{15}N-AVR-Pia (40 µM) in the absence (orange cross-peaks) and presence (pale blue cross-peaks) of unlabelled OsHPP09^{E16R}-HMA (40 µM).

Cross-peak labels correspond to the ^{15}N-AVR-Pia spectra (previously assigned [11]). Zoom inset
panels show cross-peaks corresponding to residues L39, I44, E56 and Y85 to illustrate the
chemical shift perturbations between the free (^{15}N-AVR-Pia alone) and bound (^{15}N-AVR-Pia

with OsHPP09^{E16R}-HMA) as presented in Fig. 7. Zoomed areas are indicated by boxes on the
spectra.

Supplementary Fig. 22. Comparison of binding interfaces in the crystal structures of
MAX effectors bound to HMA domains of H(I)PPs.

The structures of AVR-Pia in complex with OsHPP09-HMA (PDB accession code 9RSV),
AVR-PikF in complex with OsHIPP19-HMA (PDB accession code 7B1I [41]) and Pwl2 in
complex with OsHIPP43-HMA (PDB accession code 8R7A [46]) are represented as gold (AVR-
Pia), teal (OsHPP09-HMA), brown (Pwl2), green (OsHIPP43-HMA), pink (AVR-PikF) and grey
(OsHIPP19-HMA) ribbons with relevant secondary structure elements indicated.
Superposition of MAX effectors (left) and HMA domains (right) was carried out in CCP4mg
using secondary structure matching.

Supplementary Table S1. Amino acid sequences and MSU/RGAP and RAP-DB/IRGSP
identifiers for the H(I)PPs used in the present study.

Supplementary Table S2. Rice proteins interacting with AVR-Pia in the Y2H screen.

Results of a Y2H screen using BD:AVR-Pia as bait and a cDNA library from the rice cultivar
CO39. The number of positive clones with insert sequences (cloned in frame with Gal4 AD)
corresponding to the designated proteins are shown. * Potential false positives found in >50%
of the Y2H screens performed in the lab using various bait proteins.

Supplementary Table S3. Thermodynamic parameters obtained from ITC experiments.

Supplementary Table S4. X-ray data collection and refinement statistics for OsHPP09-
HMA / AVR-Pia

Supplementary Table S5. Comparison of the binding interfaces of complexes between
HMA domains and MAX effectors.

Supplementary Table S6. List of primers used in this study.

Supplementary Table S7. List of constructs generated in this study.

1443

References

1. Baudin, M. *et al.* Pyricularia oryzae: Lab star and field scourge. *Mol Plant Pathol* **25**, (2024).

2. Savary, S. *et al.* The global burden of pathogens and pests on major food crops. *Nat Ecol Evol* **3**, 430–439 (2019).

3. Oliveira-Garcia, E., Yan, X., Oses-Ruiz, M., de Paula, S. & Talbot, N. J. Effector-triggered susceptibility by the rice blast fungus Magnaporthe oryzae. *New Phytologist* **241**, 1007–1020 (2024).

4. de Guillen, K. *et al.* Structure analysis uncovers a highly diverse but structurally conserved effector family in phytopathogenic fungi. *PLoS Pathog* **11**, (2015).

5. Lahfa, M. *et al.* The structural landscape and diversity of Pyricularia oryzae MAX effectors revisited. *PLoS Pathog* **20**, e1012176 (2024).

6. Le Naour--Vernet, M. *et al.* Adaptive evolution in virulence effectors of the rice blast fungus Pyricularia oryzae. *PLoS Pathog* **19**, e1011294 (2023).

7. Le Naour--Vernet, M. *et al.* Structure-guided insights into the biology of fungal effectors. *New Phytologist* **246**, 1460–1477 (2025).

8. Liao, J. *et al.* Pathogen effectors and plant immunity determine specialization of the blast fungus to rice subspecies. *Elife* **5**, (2016).

9. Cesari, S. *et al.* The rice resistance protein pair RGA4/RGA5 recognizes the Magnaporthe oryzae effectors AVR-Pia and AVR1-CO39 by direct binding. *Plant Cell* **25**, 1463–1481 (2013).

10. Guo, L. *et al.* Specific recognition of two MAX effectors by integrated HMA domains in plant immune receptors involves distinct binding surfaces. *Proc Natl Acad Sci U S A* **115**, 11637–11642 (2018).

11. Ortiz, D. *et al.* Recognition of the Magnaporthe oryzae effector AVR-Pia by the decoy domain of the rice NLR immune receptor RGA5. *Plant Cell* **29**, 156–168 (2017).

12. Cesari, S. Multiple strategies for pathogen perception by plant immune receptors. *New Phytologist* vol. 219 17–24 Preprint at https://doi.org/10.1111/nph.14877 (2018).

13. Cesari, S., Bernoux, M., Moncuquet, P., Kroj, T. & Dodds, P. N. A novel conserved
mechanism for plant NLR protein pairs: The 'integrated decoy' hypothesis. *Front Plant*
Sci **5**, (2014).

14. Kroj, T., Chanclud, E., Michel-Romiti, C., Grand, X. & Morel, J. B. Integration of decoy
domains derived from protein targets of pathogen effectors into plant immune receptors
is widespread. *New Phytologist* **210**, 618–626 (2016).

15. Sarris, P. F., Cevik, V., Dagdas, G., Jones, J. D. G. & Krasileva, K. V. Comparative
analysis of plant immune receptor architectures uncovers host proteins likely targeted
by pathogens. *BMC Biol* **14**, 1–19 (2016).

16. Bailey, P. C. *et al.* Dominant integration locus drives continuous diversification of plant
immune receptors with exogenous domain fusions. *Genome Biol* **19**, 1–18 (2018).

17. Oikawa, K. *et al.* The blast pathogen effector AVR-Pik binds and stabilizes rice heavy
metal-associated (HMA) proteins to co-opt their function in immunity. *PLoS Pathog* **20**,
e1012647 (2024).

18. Dykema, P. E. *et al.* A new class of proteins capable of binding transition metals. *Plant*
Mol Biol **41**, 139–150 (1999).

19. Suzuki, N., Yamaguchi, Y., Koizumi, N. & Sano, H. Functional characterization of a
heavy metal binding protein Cdl19 from Arabidopsis. *Plant Journal* **32**, 165–173 (2002).

20. De Abreu-Neto, J. B., Turchetto-Zolet, A. C., De Oliveira, L. F. V., Bodanese Zanettini,
M. H. & Margis-Pinheiro, M. Heavy metal-associated isoprenylated plant protein (HIPP):
characterization of a family of proteins exclusive to plants. *FEBS J* **280**, 1604–1616
(2013).

21. Tehseen, M., Cairns, N., Sherson, S. & Cobbett, C. S. Metallochaperone-like genes in
Arabidopsis thaliana. *Metallomics* **2**, 556–564 (2010).

22. Crowell, D. N. & Huizinga, D. H. Protein isoprenylation: the fat of the matter. *Trends*
Plant Sci **14**, 163–170 (2009).

23. Kay, B. K., Williamson, M. P. & Sudol, M. The importance of being proline: the
interaction of proline-rich motifs in signaling proteins with their cognate domains. *The*
FASEB Journal **14**, 231–241 (2000).

24. Khan, I. U. *et al.* Identification of novel rice (Oryza sativa) HPP and HIPP genes tolerant
to heavy metal toxicity. *Ecotoxicol Environ Saf* **175**, 8–18 (2019).

25. Li, J. *et al.* Heavy metal stress-associated proteins in rice and Arabidopsis: genome-
wide identification, phylogenetics, duplication, and expression profiles analysis. *Front*
Genet **11**, (2020).

26. Barr, Z. K., Werner, T. & Tilsner, J. Heavy Metal-Associated Isoprenylated Plant
Proteins (HIPPs) at plasmodesmata: Exploring the link between localization and
function. *Plants* vol. 12 Preprint at https://doi.org/10.3390/plants12163015 (2023).

27. Barth, O., Vogt, S., Uhlemann, R., Zschiesche, W. & Humbeck, K. Stress induced and
nuclear localized HIPP26 from Arabidopsis thaliana interacts via its heavy metal
associated domain with the drought stress related zinc finger transcription factor
ATHB29. *Plant Mol Biol* **69**, 213–226 (2009).

28. Barth, O., Zschiesche, W., Siersleben, S. & Humbeck, K. Isolation of a novel barley
cDNA encoding a nuclear protein involved in stress response and leaf senescence.
Physiol Plant **121**, 282–293 (2004).

29. Cowan, G. H. *et al.* Potato mop-top virus co-opts the stress sensor HIPP26 for long-
distance movement. *Plant Physiol* **176**, 2052–2070 (2018).

30. Radakovic, Z. S. *et al.* Arabidopsis HIPP27 is a host susceptibility gene for the beet
cyst nematode Heterodera schachtii. *Mol Plant Pathol* **19**, 1917–1928 (2018).

31. Fukuoka, S. *et al.* Loss of function of a proline-containing protein confers durable
disease resistance in Rice. *Science (1979)* **325**, 998–1001 (2009).

32. Were, V. M. *et al.* The Magnaporthe oryzae effector Pwl2 alters HIPP43 localization to
suppress host immunity. *Plant Cell* (2025) doi:10.1093/PLCELL/KOAF116.

33. Huang, Z. *et al.* Small secreted effector protein from Fusarium sacchari suppresses
host immune response by inhibiting ScPi21-induced cell death. *Mol Plant Pathol* **25**,
e13414 (2024).

34. Song, H. *et al.* The Meloidogyne graminicola effector MgMO289 targets a novel copper
metallochaperone to suppress immunity in rice. *J Exp Bot* **72**, 5638–5655 (2021).

1529 35. Huang, Q. *et al.* CRISPR/Cas9-mediated mutagenesis of the susceptibility gene
1530   OsHPP04 in rice confers enhanced resistance to rice root-knot nematode. *Front Plant*
1531   *Sci* **14**, (2023).

1532 36. Niu, X. *et al.* A small secreted protein, RsMf8HN, in Rhizoctonia solani triggers plant
1533   immune response, which interacts with rice OsHIPP28. *Microbiol Res* **266**, 127219
1534   (2023).

1535 37. Zhang, X. *et al.* Isolation and characterisation of cDNA encoding a wheat heavy metal-
1536   associated isoprenylated protein involved in stress responses. *Plant Biol* **17**, 1176–
1537   1186 (2015).

1538 38. Huang, G. *et al.* Genome-wide characterization of heavy metal-associated
1539   isoprenylated plant protein gene family from Citrus sinensis in response to
1540   huanglongbing. *Front Plant Sci* **15**, (2024).

1541 39. Imran, Q. M. *et al.* Nitric oxide responsive heavy metal-associated gene AtHMAD1
1542   contributes to development and disease resistance in Arabidopsis thaliana. *Front Plant*
1543   *Sci* **7**, 232346 (2016).

1544 40. Bentham, A. R. *et al.* A single amino acid polymorphism in a conserved effector of the
1545   multihost blast fungus pathogen expands host-target binding spectrum. *PLoS Pathog*
1546   **17**, e1009957- (2021).

1547 41. Maidment, J. H. R. *et al.* Multiple variants of the fungal effector AVR-Pik bind the HMA
1548   domain of the rice protein OsHIPP19, providing a foundation to engineer plant defense.
1549   *Journal of Biological Chemistry* **296**, (2021).

1550 42. Ashikawa, I. *et al.* Two adjacent nucleotide-binding site-leucine-rich repeat class genes
1551   are required to confer Pikm-specific rice blast resistance. *Genetics* **180**, 2267–2276
1552   (2008).

1553 43. Zdrzałek, R., Kamoun, S., Terauchi, R., Saitoh, H. & Banfield, M. J. The rice NLR pair
1554   Pikp-1/Pikp-2 initiates cell death through receptor cooperation rather than negative
1555   regulation. *PLoS One* **15**, (2020).

1556 44. Cesari, S. *et al.* New recognition specificity in a plant immune receptor by molecular
1557   engineering of its integrated domain. *Nat Commun* **13**, (2022).

45.　Varden, F. A. *et al.* Cross-reactivity of a rice NLR immune receptor to distinct effectors
from the rice blast pathogen Magnaporthe oryzae provides partial disease resistance.
J Biol Chem **294**, 13006–13016 (2019).

46.　Zdrzałek, R. *et al.* Bioengineering a plant NLR immune receptor with a robust binding
interface toward a conserved fungal pathogen effector. *Proc Natl Acad Sci U S A* **121**,
e2402872121 (2024).

47.　Langner, T., Kellner, R., Harant, A., Terauchi, R. & Kamoun, S. A clone resource of
cereal heavy-metal associated (HMA) domains— putative targets of plant pathogen
effector proteins. Preprint at https://doi.org/10.5281/zenodo.5148558 (2021).

48.　Gómez De La Cruz, D., Zdrzałek, R., Banfield, M. J., Talbot, N. J. & Moscou, M. J.
Molecular mimicry of a pathogen virulence target by a plant immune receptor. *bioRxiv*
(2024) doi:10.1101/2024.07.26.605320.

49.　Homma, F., Huang, J. & van der Hoorn, R. A. L. AlphaFold-Multimer predicts cross-
kingdom interactions at the plant-pathogen interface. *Nat Commun* **14**, 6040 (2023).

50.　Maqbool, A. *et al.* Structural basis of pathogen recognition by an integrated HMA
domain in a plant NLR immune receptor. *Elife* **4**, e08709 (2015).

51.　Ose, T. *et al.* Solution structure of an avirulence protein, AVR-Pia, from Magnaporthe
oryzae. *J Biomol NMR* **63**, 229–235 (2015).

52.　Krissinel, E. & Henrick, K. Inference of macromolecular assemblies from crystalline
state. *J Mol Biol* **372**, 774–797 (2007).

53.　Schymkowitz, J. *et al.* The FoldX web server: an online force field. *Nucleic Acids Res*
33, W382–W388 (2005).

54.　Park, C. H. *et al.* The Magnaporthe oryzae effector AvrPiz-t targets the RING E3
ubiquitin ligase APIP6 to suppress pathogen-associated molecular pattern–triggered
immunity in rice. *Plant Cell* **24**, 4748–4762 (2012).

55.　Park, C. H. *et al.* The E3 ligase APIP10 connects the effector AvrPiz-t to the NLR
receptor Piz-t in rice. *PLoS Pathog* **12**, (2016).

56.　Shi, X. *et al.* The fungal pathogen Magnaporthe oryzae suppresses innate immunity by
modulating a host potassium channel. *PLoS Pathog* **14**, e1006878 (2018).

57.   Wang, R. *et al.* Immunity to rice blast disease by suppression of effector-triggered
necrosis. *Current Biology* **26**, 2399–2411 (2016).

58.   Tang, M. *et al.* The Nup98 homolog APIP12 targeted by the effector AvrPiz-t is involved
in rice basal resistance against Magnaporthe oryzae. *Rice* **10**, 1–11 (2017).

59.   Couch, B. C. *et al.* Origins of host-specific populations of the blast pathogen
Magnaporthe oryzae in crop domestication with subsequent expansion of pandemic
clones on rice and weeds of rice. *Genetics* **170**, 613–630 (2005).

60.   Tosa, Y. *et al.* Evolution of an avirulence gene, AVR1-CO39, concomitant with the
evolution and differentiation of Magnaporthe oryzae. *Molecular Plant-Microbe*
Interactions **18**, 1148–1160 (2005).

61.   Hála, M., Žárský, V., Swiezewska, E. & Surmacz, L. Protein prenylation in plant stress
responses. *Molecules 2019, Vol. 24, Page 3906* **24**, 3906 (2019).

62.   Guo, T. *et al.* Arabidopsis HIPP proteins regulate endoplasmic reticulum-associated
degradation of CKX proteins and cytokinin responses. *Mol Plant* **14**, 1918–1934 (2021).

63.   Nawaz, G. *et al.* Knockout of Pi21 by CRISPR/Cas9 and iTRAQ-based proteomic
analysis of mutants revealed new insights into M. oryzae resistance in elite rice line.
Genes (Basel) **11**, 1–24 (2020).

64.   Tao, H. *et al.* Engineering broad-spectrum disease-resistant rice by editing multiple
susceptibility genes. *J Integr Plant Biol* **63**, 1639–1648 (2021).

65.   Zhou, Y. *et al.* Engineering of rice varieties with enhanced resistances to both blast and
bacterial blight diseases via CRISPR/Cas9. *Plant Biotechnol J* **20**, 876–885 (2022).

66.   Hanika, K. *et al.* Impairment of tomato WAT1 enhances resistance to vascular wilt fungi
despite severe growth defects. *Front Plant Sci* **12**, 721674 (2021).

67.   Kieu, N. P., Lenman, M., Wang, E. S., Petersen, B. L. & Andreasson, E. Mutations
introduced in susceptibility genes through CRISPR/Cas9 genome editing confer
increased late blight resistance in potatoes. *Sci Rep* **11**, 1–12 (2021).

68.   Wolter, M., Hollricher, K., Salamini, F. & Schulze-Lefert, P. The mlo resistance alleles
to powdery mildew infection in barley trigger a developmentally controlled defence
mimic phenotype. *MGG Molecular & General Genetics* **239**, 122–128 (1993).

69.　Kusch, S. *et al.* Arabidopsis mlo3 mutant plants exhibit spontaneous callose deposition
and signs of early leaf senescence. *Plant Mol Biol* **101**, 21–40 (2019).

70.　Garcia-Ruiz, H., Szurek, B. & Van den Ackerveken, G. Stop helping pathogens:
engineering plant susceptibility genes for durable resistance. *Curr Opin Biotechnol* **70**,
187–195 (2021).

71.　Weber, E., Engler, C., Gruetzner, R., Werner, S. & Marillonnet, S. A modular cloning
system for standardized assembly of multigene constructs. *PLoS One* **6**, e16765
(2011).

72.　Engler, C., Kandzia, R. & Marillonnet, S. A one pot, one step, precision cloning method
with high throughput capability. *PLoS One* **3**, (2008).

73.　Bentham, A. R. *et al.* pOPIN-GG: A resource for modular assembly in protein
expression vectors. *bioRxiv* 2021.08.10.455798 (2021)
doi:10.1101/2021.08.10.455798.

74.　Bernoux, M. *et al.* Structural and functional analysis of a plant resistance protein TIR
domain reveals interfaces for self-association, signaling, and autoregulation. *Cell Host*
Microbe **9**, 200–211 (2011).

75.　Kushnirov, V. V. Rapid and reliable protein extraction from yeast. *Yeast* **16**, 857–860
(2000).

76.　Wishart, D. S. *et al.* 1H, 13C and 15N chemical shift referencing in biomolecular NMR.
J Biomol NMR **6**, 135–140 (1995).

77.　Vranken, W. F. *et al.* The CCPN data model for NMR spectroscopy: Development of a
software pipeline. *Proteins: Structure, Function and Genetics* **59**, 687–696 (2005).

78.　Incardona, M. F. *et al.* EDNA: A framework for plugin-based applications applied to X-
ray experiment online data analysis. *J Synchrotron Radiat* **16**, 872–879 (2009).

79.　Evans, P. R. An introduction to data reduction: space-group determination, scaling and
intensity statistics. *Acta Crystallogr D Biol Crystallogr* **67**, 282 (2011).

80.　Evans, P. R. & Murshudov, G. N. How good are my data and what is the resolution?
urn:issn:0907-4449 **69**, 1204–1214 (2013).

81. Agirre, J. *et al.* The CCP4 suite: integrative software for macromolecular
crystallography. *urn:issn:2059-7983* **79**, 449–461 (2023).

82. McCoy, A. J. *et al.* Phaser crystallographic software. *urn:issn:0021-8898* **40**, 658–674
(2007).

83. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**,
679–682 (2022).

84. Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot.
Acta Crystallogr D Biol Crystallogr **66**, 486–501 (2010).

85. Murshudov, G. N., Vagin, A. A. & Dodson, E. J. Refinement of macromolecular
structures by the maximum-likelihood method. *Acta Crystallogr D Biol Crystallogr* **53**,
240–255 (1997).

86. Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal
structures. *urn:issn:0907-4449* **67**, 355–367 (2011).

87. Davis, I. W., Murray, L. W., Richardson, J. S. & Richardson, D. C. MolProbity: Structure
validation and all-atom contact analysis for nucleic acids and their complexes. *Nucleic*
Acids Res **32**, (2004).

88. Chen, V. B. *et al.* MolProbity: All-atom structure validation for macromolecular
crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12–21 (2010).

89. Schrödinger LLC. *The PyMOL Molecular Graphics System, Version 2.5.0 Open-*
Source. (2015).

90. McNicholas, S., Potterton, E., Wilson, K. S. & Noble, M. E. M. Presenting your
structures: the CCP4mg molecular-graphics software. *urn:issn:0907-4449* **67**, 386–394
(2011).

91. Evans, R. *et al.* Protein complex prediction with AlphaFold-Multimer. *bioRxiv* (2022)
doi:10.1101/2021.10.04.463034.

92. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*
596, 583–589 (2021).

93. Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions with
AlphaFold 3. *Nature* **630**, 493–500 (2024).

94.   Maidment, J. H. R., Bocquet, A., Gelin, M., De Guillen, K. & Cesari, S. BAG Montpellier

1674   [Dataset]. Preprint at https://doi.org/10.15151/ESRF-DC-2199982133 (2025).

1675

1676

Acknowledgements

We thank Mark Youles (The Sainsbury Laboratory, Norwich, UK) and Mark Banfield (John Innes Centre, Norwich UK) for generating and providing the pOPIN-GG vector and tag modules. Gateway-compatible split-luciferase vectors were kindly provided by Laurent Deslandes (INRAE Toulouse, France). We also thank Florian Veillet (Plant Health Institute of Montpellier, France) for providing the pFV79 construct used in this study. We are grateful to Julie Noell (Centre de Biologie Structurale, France) for assistance with protein purification, and to Corinne Lionne and Manon Mallet (Centre de Biologie Structurale, France) for helpful discussions about isothermal titration calorimetry methods. We thank Sebastien Ribeiro (Plant Health Institute of Montpellier, France) for valuable discussions and support throughout the project. We also thank Laura Mathieu (Plant Health Institute of Montpellier, France) for her support with the analysis of the split-luciferase data and Vincent Chochois (CIRAD, France) for his assistance with sequence analysis related to the Y2H cDNA library screen.

This work benefited from access to the NMR and RX facilities of the Integrated Biophysics and Structural Biology Platform (PIBBS) of the CBS. PIBBS is a GIS-IBISA platform and belongs to the French Infrastructure for Integrated Structural Biology (FRISBI), supported by the National Research Agency (ANR-10-INBS-05).

We acknowledge the European Synchrotron Radiation Facility (ESRF) for provision of synchrotron radiation facilities. We would like to thank the staff of the ESRF and EMBL Grenoble for assistance and support in using beamline ID30-A1 under proposal number mx-2409. This work was funded by the HORIZON EUROPE European Research Council grant ERC-2019-STG-852482-ii-MAX awarded to Stella Cesari.

Author contributions

Josephine H.R. Maidment: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing. **Svenja C. Saile**: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing. **Aurélien Bocquet**: Investigation; Methodology; Resources; Validation; Writing – review & editing. **Céline Thivolle**: Investigation; Methodology; Resources; Validation; Writing – review & editing. **Léo Bourcet**: Investigation; Methodology; Resources; Validation; Writing – review & editing. **Lisa-Fatimatou Planel**: Investigation; Methodology; Resources; Validation; Writing – review & editing. **Muriel Gelin**: Methodology; Resources; Writing – review & editing. **Thomas Kroj**: Conceptualization; Writing – review & editing. **André Padilla**: Conceptualization; Methodology; Writing – review & editing. **Karine de Guillen**: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing. **Stella Cesari**: Conceptualization; Data curation; Formal analysis; Funding acquisition; Investigation; Methodology; Project administration; Resources; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing.

1721 **Competing interests**

1722   The authors declare that they have no conflicts of interest with the contents of this article.