Cortical neural landscape captures mouse-to-mouse variability in anticipatory vs. inattentive decision making

Chang Yin [1], **Naoki Hiratani** [1*]
[1] Department of Neuroscience, Washington University in St Louis
*Correspondence: hiratani@wustl.edu

ABSTRACT

Understanding individual variability in behavior is crucial for both basic and clinical neuroscience, yet it remains challeng-
ing to study in traditional single laboratory experiments with small sample size. Leveraging standardized behavioral and
neural datasets from the International Brain Laboratory, comprising approximately 100 mice trained on a visual decision-
making task, we investigated the structure and neural correlates of inter-animal behavioral variability. Using reaction time
analysis and a deep learning-based embedding of individual animals, we uncovered large but low-dimensional differences
in behavioral traits. Some mice consistently exhibited anticipatory responses, marked by fast reaction times, while others
showed slower, more disengaged behavior. These behavioral profiles were consistent across sessions, with female mice
tending to show more anticipatory behavior than males. We hypothesized that this behavioral spectrum reflects differ-
ences in the depth of underlying cortical states, reflected in the temporal dynamics of neural activity. Supporting this idea,
we found that the characteristic timescale of population activity, measured during both inter-trial intervals and passive pe-
riods, correlated with an animal's anticipatory tendency across cortical areas, especially in medial visual areas. These find-
ings suggest that individual differences in the cortical dynamics may underlie distinct decision-making strategies.

INTRODUCTION

Individual variability is a hallmark of social animals. Even genetically homogeneous inbred mice and rats show considerable differ-
ences in stress resistance [1–3], task-acquisition speed [4–6], and decision-making behavior [7,8]. Traditionally, such inter-animal variabil-
ity has been viewed as a nuisance that undermines the reliability and reproducibility of experiments [9,10]. While efforts to reduce
variability have led to valuable guidelines for improving experimental rigor, variability itself can be informative rather than merely
problematic [11,12]. Natural variability offers a unique opportunity to test theories of brain function, which must ultimately account
for both behavioral and neural diversity. Moreover, understanding across-individual variability in behavior and its neural correlates is
crucial for elucidating the mechanisms underlying neurodevelopmental and psychiatric disorders.

Recent studies have begun leveraging animal-to-animal differences to probe how the brain generates diverse behaviors [6–8,11,13].
For example, one study found a large variability among rats in their responses to unfamiliar sounds during an auditory decision-
making task, and this variability led to mechanistic insights into how neural circuits resolve uncertainty [7]. Nevertheless, most pre-
vious investigations were single-laboratory studies with small sample sizes, inherently limiting statistical power and constraining
computational analyses of the origins of variability. Consequently, we still lack a clear understanding of how animal-to-animal vari-
ability is structured and how it is manifested in brain-wide neural activity. Furthermore, a scalable, data-driven method for charac-
terizing inter-animal variability across large cohorts is still missing.

Here, we address these challenges by leveraging data from the International Brain Laboratory (IBL), in which more than 100 C57BL/6
mice were trained on an identical visual decision-making task under fully standardized conditions [5], and brain-wide activity was
recorded with Neuropixels probes [14]. In this task, a mouse rotates a wheel clockwise or counter-clockwise depending on whether
a grating appears on the right or left side of a monitor, respectively. The experiment was conducted across twelve laboratories, yet
housing, training, and equipment were standardized to minimize extraneous variability. Behavioral performance and block-structure
dependence were consistent across labs, although the number of sessions required to reach proficiency differed slightly [5,15]. Like-
wise, Neuropixels recordings from identical brain sites showed similar neuronal yields, firing rates, and LFP power across laborato-
ries, with modest lab-to-lab differences in task-related encoding [16]. This small across-laboratory variability makes the IBL dataset

ideally suited for studying animal-to-animal variability, because the remaining differences are likely to reflect fundamental biological sources rather than experimental artifacts. However, inter-animal variability in the IBL data has not yet been systematically examined. Moreover, prior behavioral analyses have focused primarily on psychometric performance[5,15,17,18], leaving chronometric aspects of task engagement, where richer behavioral repertoires and stronger inter-animal differences may reside, largely unexplored.

In this work we begin by analyzing the distribution of reaction times (RTs) across animals and identify two interesting features. First, mice frequently make extremely fast responses by initiating wheel movement right before stimulus onset. Second, they occasionally respond very slowly, such that RTs form a bimodal distribution on a logarithmic scale, with a secondary peak around 3 second. Analyzing their behavioral characteristics, we show that early and late RTs likely correspond to anticipatory and inattentive responses, respectively. Although most mice display both behaviors, the proportion varies widely: some animals predominantly make early responses, whereas others favor late responses. To capture this difference we introduce an anticipatory-tendency index, defined as the difference between the fractions of early and late responses. This index is stable across sessions, and female mice exhibit a slightly but significantly higher anticipatory tendency relative to males.

To investigate inter-animal variability more deeply, we develop a data-driven method that learns a low-dimensional animal embedding jointly with a deep neural network that predicts choice and RT. The learned embeddings improve prediction in held out sessions and reveal a salient structure: the primary embedding axis captures individual differences in RT, while the secondary axis reflects left/right biases of animals.

Finally, we link behavioral variability to neural dynamics by analyzing brain-wide recordings. We hypothesized that inter-animal differences arise from variations in the depth of the underlying neural landscape, which predict systematic differences in autocorrelation timescales. We tested this prediction by measuring autocorrelation during post-task passive periods and inter-trial intervals (ITIs), using a bootstrap method to obtain unbiased estimates from short ITIs. Across both periods, the autocorrelation timescale in medial visual areas is negatively correlated with an animal's anticipatory-tendency index. Moreover, the cortex-wide relative timescales exhibits the same negative correlation, supporting our hypothesis.

RESULTS

Two oddities in the reaction time distribution

The IBL experiment is a decision-making task for mice. In each trial, a grating stimulus appears on either the left or right side of a screen positioned in front of a head-fixed mouse (Fig. 1A). The task requires the mouse to turn a wheel clockwise if the stimulus appears on the left, and counter-clockwise if it appears on the right. For brevity, we refer to the correct movement in response to a stimulus on the right (left) side as a right (left) motion. If the mouse performs the task correctly, it receives a small volume of sucrose water as a reward; otherwise, a burst of noise is delivered. If no choice is made within 60 seconds, the trial times out. The next trial begins automatically after a short interval (approximately 1 second) and a quiescent period (0.4–0.7 seconds). If the mouse moves the wheel more than 2° during the quiescent period, the period resets, and a new quiescent interval must be completed before the stimulus appears (see Supplementary Fig. S1A for the trial timeline). Stimulus contrast is randomly selected from 100%, 25%, 12.5%, 6%, or 0%. An auditory cue (5 kHz tone) signals the onset of the stimulus, allowing the mouse to detect the start of a trial even in low contrast trials. The probability of the stimulus appearing on the left or right is modulated in a block-wise fashion. In the initial ~100 trials of a session, stimulus locations are unbiased (50% left, 50% right), after which they become biased towards one side (80:20). Block changes occur every 20–100 trials and are not signaled to the animal.

Approximately half of the mice learned to perform the task with no significant performance differences across laboratories[5]. However, task performance as measured by psychometric curves does not fully capture the complexity of behavior[19,20]. In particular, reaction time (RT) may reveals additional behavioral variability across animals.

We examined RT distributions in expert mice, defined as those that met the criteria for electrophysiological recordings[5]. We defined the first movement onset time as the point when the mouse begins a significant wheel movement (orange vertical dotted line in Fig. 1B; see *Reaction time analysis* in Methods). RT was then defined as the interval between stimulus onset (blue line) and

Figure 1: Task description and RT statistics **(A)** Schematic of the task. **(B)** Definition of RT and ET. We defined the movement onset time as the point where a mouse started to make a significant wheel movement (orange line), and RT as the duration between the movement onset time and the stimulus onset time. Feedback time is the time point where the trial ends and the mouse receives either a water reward or a short wait time (green line). ET was defined as the duration between the feedback time and the stimulus onset time. **(C)** RT distribution across all animals and trials (n = 267,474 trials). **(D)** Psychometric curves of the wheel movement direction under early (RT < 0.08s) and normal (0.08s ≤ t < 1.25s) RTs. **(E)** RT plotted in a logarithmic scale. Note that bins are taken on a logarithmic scale, meaning that bins for a longer RT contain exponentially longer intervals. **(F)** The same as panel E, but divided into high (c = 25% and 100%) and low (c = 0%, 6%, and 12.5%) contrasts. Dark blue regions represent incorrect trials whereas light blue regions represent correct trials. **(G)** The ratio of early and late responses as a function of the relative position of a trial in a session. Early and late responses were defined as RT < 0.08 s and 1.25 s ≤ RT. Shaded areas in panels D and G represent the standard error of the means over trials.

the first movement onset. A trial ends when the mouse moves the wheel 35 degrees in either direction, at which point it receives a reward or a noise. We refer to this endpoint as the feedback time (green line), and the duration from stimulus onset to feedback time as the execution time (ET). We analyzed RTs from 267,474 trials across 415 sessions by 132 mice with electrophysiology recording. These sessions were chosen to study behavior of expert animals and to allow exploration of neural correlates in later analyses. We found two notable features in the RT distribution(Figs. 1C and 1E).

First, mice exhibited very fast, and often negative, RTs in approximately 10% of trials, indicating that some movements were initiated before stimulus onset (see Supp. Figs. S1B and S1C for example trajectories). In each trial, there is a 400–700 ms quiescent period during which the mouse must keep the wheel still. If the wheel is moved more than 2 degrees during this period, the quiescent period recurs (Supp. Figs. S1A). However, mice can circumvent this reset by making subtle movements or initiating movement just before stimulus onset, resulting in valid but negative RTs. These early responses were consistently observed across labs (yellow points in Supp. Fig. S2A).

To further investigate, we classified trials with RTs < 0.08 seconds as early response trials (red line in Fig. 1C), based on a clear inflection point in the RT distribution. Note that it is effectively impossible for a stimulus-driven response to occur within 80 ms of onset. As expected, in trials with normal RTs, choice behavior depended on both stimulus contrast and block structure (Fig. 1D, right). In contrast, in early RT trials, the probability of rightward movement was independent of stimulus side and contrast but strongly modulated by the block structure (Fig. 1D, left). This lack of stimulus dependence suggests that mice are making decisions before the stimulus is presented. Moreover, the strong influence of block structure implies that behavior is not random, but guided by their predictions about likely stimulus location. Note that we determined movement direction from the initial wheel movement, not the final choice, as mice sometimes changed their mind after an early movement (see Supp. Fig. S1C). These change-of-mind events were more frequent in early RT trials, particularly when the initial movement was incorrect (Supp. Fig. S1F), supporting the idea that early responses reflect predictive behavior. Additionally, early responses were slightly more common in the first half of a session (yellow line in Fig. 1G), likely due to higher motivation from water deprivation. The ITI distribution for early RT trials was also more long-

Figure 2: Individual variability in anticipatory tendency of movement initiation (A) Mouse-to-mouse variability in early/late response ratio ($N = 96$ animals). Each point represents one mouse, and points are color-coded by the value of the anticipatory tendency index. **(B)** The standard deviation (SD) of the anticipatory tendency over sessions calculated for each mouse, compared to random shuffle. In shuffling, we randomly attributed sessions to each animal and estimated the across-session variability for each animal. **(C, D)** The anticipatory tendency is not significantly correlated with the average task performance (C) or the number of attempted trials per session (D) across mice. Here, mice are color-coded by the lab identity specified in panel E. **(E)** Relationship between the anticipatory tendency and lab identity ($p = 0.002$; ANOVA over all labs; $p = 0.058$ ANOVA over labs A-K). **(F)** Sex dependence of the anticipatory tendency ($p = 0.024$; Welch's t-test on population means). **(G)** The relationship between the anticipatory tendency and median RT across mice.

tailed than for normal RTs (Supp. Fig. S1H), indicating frequent quiescent period resets. These findings suggest that early responses

are anticipatory actions exploiting task structure.

Second, we observed a subset of late RT trials (Fig. 1E; see Supp. Figs. S1D and S1E for example trajectories). When plotting posi-

tive RTs on a log scale, we found a bimodal distribution, with a secondary peak around 3 seconds. We defined late response trials as

those belonging to this second peak, using a threshold of 1.25 seconds (red dashed line in Fig. 1E). Late responses occurred pre-

dominantly during low-contrast trials (contrast $\leq 12.5\%$; Fig. 1F) and were associated with higher error rates, approaching chance

level (Supp. Fig. S1G), suggesting disengagement.

Although standard drift-diffusion models with urgency signals predict rising error rates with increased RT[21,22], they do not explain

the long-tailed RT distribution observed here. It is more likely that mice were inattentive during these trials, rather than deliber-

ating longer. Prior analysis of the same dataset showed larger pupil sizes before long RT trials[23], consistent with shifts in engage-

ment[24,25]. Similarly, GLM-HMM analyses have linked disengagement states to long RTs[17].

Late response trials were more frequent toward the end of sessions (purple line in Fig. 1G), likely due to satiation. To control for this

effect, we excluded the final 40 trials of each session in all subsequent analyses, except for the animal embedding analysis, where

within-session effects were explicitly modeled. The results above indicate that late responses primarily reflect disengagement or

inattention.

Animal-to-animal variability in fast-and-slow responses

Our analysis of the RT distribution revealed two notable behavioral patterns: early, anticipatory responses and late, disengaged re-

sponses. This raised the question of whether all mice exhibit both types of responses, or if these behaviors are specific to certain in-

dividuals. To investigate this, we computed the early and late response ratios for each animal, defined as the fraction of trials with

early or late RTs, respectively, out of all trials (Fig. 2A). Notably, some animals showed a strong bias toward early responses, while oth-

ers were more prone to late responses, revealing substantial inter-animal variability. To quantify this bias, we defined an Anticipatory

Tendency Index (ATI) as the difference between an animal's early and late response ratios (see Methods). A positive ATI indicates a

tendency toward anticipatory (early) responses (yellow points in Fig. 2A), whereas a negative ATI suggests a predominance of late,

disengaged responses (purple points).

Figure 3: Data-driven analysis of individual variability via animal embedding (A) Schematic figure of the network we trained for predicting the choice and reaction/execution time. See *Animal embedding algorithm* in Methods for the details. **(B)** Choice and RT/ET prediction performance in held-out session. The vanilla network only uses the stimulus contrast and block for predictions, while the '+Trial No.' model also uses the trial number, and the '+Animal ID' model incorporates animal ID for prediction. Error bars are the standard errors over 10 random training/test splits. **(C)** Normalized eigen-spectrum of the learned animal embedding. Thin lines represent results from 10 simulations with random train/test splits. **(D)** Projection of the embedding vector of each animal onto the space spanned by the two principal directions of the animal embedding matrix. **(E)** Schematic of the ideal animal psychometric and chronometric curves generation. **(F)** Expected psychometric and chronometric curves along the PC1 axis of the animal embedding. **(G)** Task performance and RT along the PC1 axis averaged over contrasts and blocks. **(H)** Projection of animal ID to PC1 of the animal embedding space correlates with the anticipatory tendency index ($r = 0.65$ and $p < 10^{-3}$). **(I)** Projection of PC1 and PC2 axes of the animal embedding (pink and yellow lines in the left panel) onto the space spanned by the mean RT and the mean choice bias (curves in the right panel).

Importantly, ATI values were generally consistent across sessions within the same animal. We assessed this by computing the stan-

dard deviation of ATI across sessions for each mouse. Most animals showed lower session-to-session variability than would be ex-

pected by chance, based on shuffled session data (Fig. 2B). ATI was not strongly correlated with overall task performance (Fig. 2C),

total number of trials performed in a session (Fig. 2D), or the number of sessions required to reach expert-level performance (Supp.

Fig. S2B).

Mice with a broad range of ATI values were observed in nearly all laboratories, although one lab (Lab L) exhibited a notably higher

fraction of animals with high ATIs ($p = 0.058$, one-way F-test excluding Lab L; Fig. 2E). Interestingly, female mice showed slightly

higher ATI values than males, though the effect size was modest (two-sided Welch's t-test, $p = 0.024$; $N_{male} = 67$, $N_{female} = 29$; Fig.

2F). This finding aligns with a recent study reporting faster average reaction times in female mice compared to males[4].

Reaction speeds, when measured using the median RT, were also consistent within individual animals across sessions (Supp. Fig.

S2C). However, differences in median RT were relatively small among animals with a positive ATI (Fig. 2G). This supports the notion

that ATI more effectively captures meaningful inter-animal variability in response patterns than simple metrics like median RT.

Animal embedding analysis

The results above indicate the presence of individual variability along the anticipatory-inattention axis. This raises the question of whether this axis is the dominant factor underlying animal-to-animal variability, or whether other dimensions also contribute substantially. Moreover, it remains unclear if the observed variability improves behavioral predictions.

To explore these questions, we conducted a data-driven analysis of the variability using a deep neural network. Specifically, we built a feedforward neural network to predict the choice, reaction time (RT), and execution time (ET) of each trial, given the stimulus contrast, block, trial number (i.e., its relative position within a session), and animal ID (Fig. 3A; see *Animal embedding algorithm* in Methods). Conceptually similar to a Generalized Linear Model for behavioral prediction, the model incorporates trainable animal embedding components that capture features that vary across individuals. It also uses nonlinear transformations to learn complex interactions among input variables. The animal ID is initially encoded as a 95-dimensional one-hot vector (corresponding to the 95 animals), which is then linearly projected into a four-dimensional animal embedding space (green layer in Fig. 3A). By training the network to optimize prediction performance, the embedding space learns low-dimensional representations that capture latent factors contributing to individual variability in choice, RT, and ET.

We trained this model on 334 sessions from 95 expert mice and evaluated its performance on held-out sessions. Sessions were split semi-randomly to ensure that each session in the test dataset had at least one corresponding session from the same animal in the training dataset. After training, the model incorporating animal ID outperformed the model without it in predicting both choices and RTs on held-out sessions (green vs. blue lines in Fig. 3B). This result highlights consistent behavior across sessions within individual animals and indicates that the model captures features enabling cross-session prediction. Additionally, the trial's relative position within a session was important for predicting RT (orange vs. blue line in Fig. 3B, right), consistent with the observed within-session shift in late-response rates (Fig. 1G).

To further investigate the structure of individual variability, we analyzed the learned animal embeddings. We performed eigen-decomposition on the weight matrix that maps one-hot animal IDs into the low-dimensional embedding space. We found that the first principal component (PC1) accounted for over 40% of the variance captured in the embedding space (Fig. 3C). When projecting all animals into the PC1–PC2 space, we observed a continuous distribution without distinct clusters (Fig. 3D).

A key advantage of our embedding approach is that it enables behavioral simulation of a hypothetical animal, allowing us to interpret the embedding axes. We aligned the embedding vector along the PC1 axis and used it to estimate average choice, RT, and ET across different stimuli and blocks, yielding psychometric and chronometric curves (Fig. 3E). Along the PC1 direction, we observed a drastic shift in chronometric curves: RTs exceeded one second for animals with negative PC1 projections (bottom-left panel, Fig. 3F) and became shorter or even negative for animals with positive projections (bottom-right). The psychometric curve showed a pronounced leftward bias in the high-RT region (top-left), a pattern observed in some mice. When averaging chronometric curves to estimate mean RT, we found that RT decreased monotonically, but nonlinearly, along the PC1 direction (right panel, Fig. 3G). Task performance also varied systematically: deviations in either direction along PC1 reduced accuracy (left panel, Fig. 3G). This suggests that both impulsivity and inattention are associated with poorer task performance, relative to a more balanced behavioral state. This relationship was less apparent when analyzing performance as a function of anticipatory tendency alone (Fig. 2C), possibly because higher-order components obscure this trend. Notably, projection onto PC1 was highly correlated with ATI defined earlier (Fig. 3H), suggesting that the anticipatory-inattentive axis is a major axis of individual variability.

In contrast, PC2 primarily captured variations in psychometric bias (left vs. right) with only minor effects on RT (Supp. Fig. S3B). When plotting average choice bias and RT across PC1 and PC2 axes, we observed nonlinear but approximately orthogonal trends (Fig. 3I). This suggests that individual variability is at least two-dimensional: although highly inattentive mice tend to exhibit left bias, animals across a broad range of anticipatory-inattentive profiles can display left, right, or no bias at all.

A neural landscape hypothesis on individual variability

The behavioral analyses above reveal a low-dimensional structure underlying animal-to-animal variability, with a principal axis corresponding to the anticipatory-inattentive tendency. In the following sections, we explore the potential neural origins of this variability.

Figure 4: A neural landscape model on individual variability (A) Schematic of the model. **(B)** First passage time (the duration it takes for the neural state to move from one to another) under three neural landscapes as a function of the input amplitude. We initiated the state at the center and estimated the time until it traverses to the right attractor as illustrated in the inset. **(C)** False-start ratio under the three models. Here, we defined the false-start as the transition to another state before the input is provided to the model. **(D)** autocorrelation of dynamics under the three models. See *Neural landscape model* in Methods for details.

One plausible hypothesis is that differences in the neural landscape underlie the observed behavioral differences, which has previously been proposed in the context of autism spectrum disorders[26,27] and sex differences in bandit task behaviors[4]. Figure 4A illustrates this conceptual model.

Neural population dynamics in the brain are often described as evolving within a landscape that contains attractor states, activity patterns corresponding to specific behavioral states, actions, or sensory inputs. In a shallow neural landscape (blue curve in Fig. 4A), the system can more easily transition between attractors. For example, if one attractor corresponds to a "hold" state and another to a "rightward movement preparation" state, then spontaneous transitions across the states may result in premature responses or false starts, as seen in anticipatory animals. In contrast, a deep landscape stabilizes neural states and reduces transitions, but may impair sensitivity to weak stimuli, potentially contributing to inattentiveness.

To test this idea, we implemented a simple dynamical model simulating state transitions under varying landscape depths. When constant input was applied, we found that the first-passage time, the time it takes for the network to transition between states, is longer in the deep landscape (Fig. 4B; see *Neural landscape model* in Methods). This effect was negligible when inputs were strong enough to reliably drive transitions (right side of Fig. 4B), but became noticeable under weak input conditions. Moreover, the false-start ratio, defined as the proportion of trials in which state transitions occurred before the input stimulus, was higher in the shallow landscape (Fig. 4C).

This toy model thus offers a mechanistic hypothesis for the neural basis of inter-animal variability. It also leads to a testable prediction: namely, that animals with shallow neural landscapes (i.e., anticipatory mice) should exhibit shorter autocorrelation timescales in their neural dynamics, while those with deeper landscapes (i.e., inattentive mice) should show longer timescales. This follows from the idea that shallow landscapes allow rapid transitions, causing autocorrelation to decay quickly, whereas deep landscapes promote state persistence and thus slower decay (blue vs. orange curves in Fig. 4D). We therefore hypothesize that autocorrelation timescales in neural activity will systematically differ between anticipatory and inattentive animals.

Comparison with autocorrelation during passive period

To test the predictions derived from the neural landscape model, we analyzed the neural activity of each animal across the brain recorded with Neuropixel probes[14] and compared it to their anticipatory tendency. We focused specifically on the passive period—the time following the task during which no external stimulus was presented (Fig. 5A) to minimize the influence of task-related behaviors on neural activity.

We estimated autocorrelation at the population level by calculating the population firing rate within each brain region. This approach allowed us to infer autocorrelation from the relatively short passive periods reliably. Because autocorrelation during this period often exhibited oscillatory components, we fitted a double-exponential function with an oscillatory term to the autocorrelation curves (Fig. 5B; see *Autocorrelation timescale estimation* in Methods); this improved the fitting consistently across areas (Supp. Fig. S4B). We defined the autocorrelation timescale as the longer of the two time constants from the fitted double-exponential terms. The estimated timescales were broadly consistent with those reported in previous work[29], although we did not observe a clear correlation with the cortical hierarchy scores reported in prior studies[30,31] (Supp. Fig. S4E). We excluded sessions with poor fit quality or timescale estimates that were clear outliers (see Methods for exclusion criteria).

Figure 5: Comparing the individual variability in anticipatory tendency and autocorrelation timescales during passive period. (A) Schematic of the autocorrelation analysis. **(B)** An example curve of the autocorrelation of population activity in one recording from the VISa region (black points) and its fitting (red curve). **(C, D)** Estimated autocorrelation timescales and their relationship with session-wise anticipatory tendency for VISam (blue), VISpm (orange), and VISa (green) regions. Each point represents one session. In panel D, we divided the autocorrelation timescales by the region-wise mean timescales (vertical lines in panel C). In panel C, p-values of correlations were $p = 0.09$, $p = 0.002$, and $p = 0.05$ for VISam, VISpm, and VISa, respectively. **(E)** Correlation between the anticipatory tendency and autocorrelation timescale of various regions (pale bars) and divisions (dark bars). The acronyms follow the Allen Brain Atlas[28]. The error bars represent standard errors without Bonferroni correction. **(F)** Correlation between the anticipatory tendency and the relative cortical timescale. Each point represents one mouse. **(G)** Histogram of the relative cortical timescale of male and female mice ($p = 0.71$, permutation test). **(H)** Relative cortical firing rate did not show correlation with the anticipatory tendency ($r = 0.08$, $p = 0.50$). **(I)** Correlation between wheel statistics and anticipatory tendency.

Figure 5E shows the correlation between the autocorrelation timescale of 29 cortical regions and each animal's anticipatory tendency during the preceding task. We excluded regions with fewer than five valid sessions. While many cortical areas exhibited negative correlations between autocorrelation timescale and anticipatory tendency, the variance was high. We thus grouped regions into functional divisions based on prior anatomical classifications[30], normalizing each session's estimated timescale by the average for that region (Figs. 5C and 5D). We split the medial division into visual and retrosplenial (RSP) areas, given their distinct roles and the potential importance of medial visual areas in this task[16].

With this grouping, we found significant negative correlations between session-wise anticipatory tendency index (ATI) and autocorrelation timescales in the medial visual areas ($p = 0.025$, Bonferroni-corrected across 7 divisions). This result aligns with our model's prediction: anticipatory animals exhibit faster neural dynamics (i.e., shorter timescales) compared to inattentive animals. We next computed a relative cortical timescale for each animal by averaging the normalized autocorrelation timescales across all recorded regions. Since Neuropixels recordings were performed acutely and the recording site varied daily for each animal[14], we aggregated data across days. For example, if VISam was recorded on Day 1 and MOp on Day 2, the animal's relative cortical timescale was taken as the mean of these two sessions' normalized timescales (see Methods). We found that the relative cortical timescale showed a significant negative correlation with ATI values across animals ($p = 0.002$, Pearson's r $= -0.33$, $N = 83$ animals; Fig. 5F), again supporting the neural landscape hypothesis. Although animals with longer cortical timescales were often male, the effect of sex on timescale was not statistically significant ($p = 0.71$, permutation test; Fig. 5G).

We also examined the relationship between relative cortical firing rates and anticipatory tendency. Using the same normalization

Figure 6: Comparing the individual variability in anticipatory tendency and autocorrelation timescales during ITI. (A) Schematic of the ITI and the selection of period for the autocorrelation estimation. **(B)** Autocorrelation estimation with vanilla or bootstrap methods. Here, the ground truth was generated artificially by sampling sequences from a Gaussian Process with an exponential decay kernel function. **(C)** The average autocorrelation curve of VISa population activity estimated with vanilla and bootstrap methods. The error bars are the standard error of the means over recordings. **(D)** Correlation between the anticipatory tendency and autocorrelation timescale of various regions (pale bars) and divisions (dark bars). The error bars are the standard errors without Bonferroni correction. **(E, F)** Correlation between the relative cortical timescale and the anticipatory tendency (E) and the projection to PC1 of the animal embedding (F). Each point represents one mouse. **(G)** Histogram of the relative cortical timescale. **(H)** Relative cortical firing rate did not show clear correlation with the anticipatory tendency ($r = -0.02, p = 0.85$). **(I)** Correlation between wheel statistics and anticipatory tendency.

and aggregation procedure, we found no clear correlation between firing rate and ATI (Fig. 5H), although relative firing rates of au-

ditory areas was correlated with session-wise ATI (Supp. Fig. S4D).

Finally, because mice often engaged with the wheel even during passive periods, we examined various wheel movement statistics

as proxies for spontaneous behavior. Total wheel movement distance, fraction of time spent moving, wheel energy, and the number

of movement onsets did not show significant correlations with ATI (Fig. 5I). However, we did find a weak but statistically significant

correlation between ATI and the number of trajectory change points in wheel movement ($p = 0.04$; Fig. 5I and Supp. Fig. S4I). This

suggests that anticipatory animals may be somewhat more fidgety even during the passive period.

Comparison with autocorrelation during inter-trial interval (ITI)

The results above suggest that animals with a higher anticipatory tendency exhibit faster neural dynamics during the passive pe-

riod, consistent with predictions from the neural landscape hypothesis. To further test this relationship, we next analyzed neural ac-

tivity during the inter-trial interval (ITI), as illustrated in Fig. 6A.

Analyzing neural dynamics during ITIs presents two key technical challenges. First, estimating autocorrelation from short-duration

ITIs is non-trivial. Standard autocorrelation methods become biased when the interval length is comparable to the underlying timescale,

due to intrinsic correlations between neural activity and its trial-wise mean [32–34]. Figure 6B illustrates the discrepancy between the

ground-truth autocorrelation and a naive ("vanilla") estimation method, even when averaging across thousands of ITIs (black vs.

blue lines; see *Autocorrelation timescale estimation* in Methods).

To address this, we developed a simple bootstrapping method in which the mean population activity is estimated using nearby ITIs rather than on a trial-by-trial basis. This enables computationally efficient estimation of autocorrelation curves and offers a practical alternative to more complex generative model-based methods[34,35]. We prove that this method yields asymptotically unbiased autocorrelation estimates when ITIs are independently drawn from a Gaussian process (see *Supplementary Text* for proof). Applied to synthetic data generated from a Gaussian process, the bootstrap method closely recovered the ground-truth autocorrelation (orange vs. black lines in Fig. 6B).

Applying this approach to neural recordings during ITIs, we observed that the vanilla method produced autocorrelation curves with a negative dip (blue line, Fig. 6C), as expected. In contrast, the bootstrap method, with a 30-trial window for mean and variance estimation, produced smooth, naturalistic autocorrelation decay curves (orange line, Fig. 6C). Using very large trial windows (e.g., all trials) introduced constant positive bias due to violation of the i.i.d. assumption. We fixed the trial window size at 30 for our analyses, though comparable results were obtained with 10- and 50-trial windows (Supp. Figs. S6C and S6D).

A second technical concern is that anticipatory mice tend to exhibit longer ITIs than inattentive ones (Supp. Fig. S2D), which could confound autocorrelation estimates. To control for this, we restricted analysis to the first 1.5 seconds of each ITI (Fig. 6A). For ITIs shorter than 2 seconds, we analyzed the period from ITI onset to 0.5 seconds before the stimulus onset, to reduce contamination from motor preparation.

Using the bootstrap method, we estimated autocorrelation timescales for each recording and assessed their correlation with ATI values at both the session and animal levels. Consistent with the passive period findings, we observed a significant negative correlation between relative autocorrelation timescale in medial visual areas and session-wise ATI (Fig. 6D; $p = 0.008$, Bonferroni-corrected across divisions; see also Supp. Fig. S5A for VISa-specific results). Furthermore, when averaging across all recorded regions per animal, the resulting relative cortical timescale also showed a significant negative correlation with animal-wise ATI ($p = 0.003$, Pearson's $r = -0.3$, $N = 103$ animals; Fig. 6E). In addition, this timescale was significantly correlated with each animal's projection onto PC1 of the animal embedding space (Fig. 6F). We confirmed that these results were robust to different neuron inclusion criteria and ITI definitions (Supp. Fig. S6).

As in the passive period analysis, sex differences in cortical timescales were not statistically significant (Fig. 6G), and cortex-wide firing rates did not correlate with anticipatory tendency (Fig. 6H, Supp. Fig. S5D). We also found no strong behavioral confounds: wheel movement metrics during ITIs, including movement distance, energy, active fraction, number of movement onsets and change points, showed no significant correlation with anticipatory tendency (Fig. 6I, Supp. Figs. S5F and S5G). Overall, the consistency between passive-period and ITI analyses reinforces our main conclusion: animals with higher anticipatory tendencies exhibit faster cortical dynamics, especially in medial visual areas, providing evidence for a neural correlate of individual behavioral variability.

DISCUSSION

In this study, we investigated behavioral variability among mice and its neural correlates in a decision-making task. By analyzing reaction times across more than quarter million trials from over 100 animals, we identified distinct properties of both negative-RT and very slow RT trials, trial types that are usually discarded in standard RT analyses, and uncovered substantial inter-animal variability in the proportion of these trials. Using an animal-embedding model, we further discovered a low-dimensional representation of this variability that predicts each animal's behavior. Finally, autocorrelation analysis of cortex-wide activity during passive and ITI periods revealed robust neural correlates of anticipatory tendency in autocorrelation timescale. Together, these findings support the hypothesis that differences in the "depth" of the landscape of cortical dynamics may underpin individual behavioral variability.

Our work also includes two methodological contributions. The first is the animal embedding algorithm (Fig. 3). While the use of deep neural networks is common in large-scale behavioral data analysis[36], previous studies either pooled data from all animals or trained a separate network for each individual animal. In contrast, our proposed approach captures inter-animal variability in a way that enables interpretable predictions of behavior. The second methodological contribution is an efficient algorithm for estimating autocorrelation from a short ITI period. We demonstrate that our method provides an unbiased estimate of autocorrelation even with limited data, unlike naive estimation approaches. This method also enables faster estimation than previous approaches based

on generative model estimation[34]. However, it assumes that ITIs are independently sampled, an assumption that is expected to be

violated over a long session (Fig. 6C).

Although we focused on passive and ITI periods to minimize task-related confounds, the observed negative relationship between

anticipatory bias and cortical time-scale may still have behavioral origins. Most animals moved the wheel significantly during fewer

than 5 % of passive periods (Suppl. Fig. S4H), yet we detected a weak positive correlation between the number of wheel-movement

change points and anticipatory tendency (Suppl. Fig. S4I). This suggests that highly anticipatory animals may fidget more even dur-

ing passive periods, modulating brain-wide dynamics[37]. We also observed that anticipatory mice exhibit a more heavy-tailed ITI

distribution than inattentive mice (Suppl. Fig. S2D). Although our analyses focused on the initial segment of each ITI to mitigate

this effect, this observation indicates cross-animal differences in mental states during ITIs. Moreover, the between-animal differ-

ences in cortical timescale we report may not reflect intrinsic circuit properties; Neuromodulatory signals are known to reshape

network autocorrelation[38,39], raising the possibility that animals with different ATI spend different amounts of duration in distinct

global brain states. Finally, while the medial visual cortex consistently showed the strongest neural–behavioral coupling, this may

partly reflect the denser sampling of VISa and VISam, which were targeted for reproducibility assessments[16].

Because all subjects were C57BL/6J mice, the variability we observed is unlikely to have a genetic origin. Prior studies have shown

that behavioral variability can be shaped by multiple factors[10], including the sex and age of the animals[1,4], the physical character-

istics of the vivarium and laboratory[40,41], in-cage social hierarchy[2,3], and implementation details of behavioral tasks[42,43]. Sex ex-

plained a modest fraction of the variance in our analysis, but the neural signature of this effect was not apparent. Housing condi-

tions present another plausible source: the mice were group-caged whenever possible, and social hierarchy within cages can in-

duce behavioral divergence even in genetically identical animals[2,3]. In particular, non-dominant mice tend to display more anxiety-

like behavior[2], which could influence their task engagement.

Structural and functional brain signatures of behavioral variability across individuals have been extensively studied in humans[44–46],

but a circuit-level understanding requires studies in animal models. In rodents, Kurikawa et al. reported behavioral variability in

response to unfamiliar auditory cues and linked it to individual differences in neural landscapes; however, their analysis was lim-

ited by low statistical power due to a small sample size[7]. Chen et al. identified sex differences in strategy during a bandit task in

mice and attributed them to differences in the depth of explore and exploit states[4], but they did not examine the underlying neural

correlates. Neural correlates of variability in contextual decision-making strategies[8] and learning trajectories[6] have also been ex-

plored recently. Nevertheless, systematic, data-driven accounts of behavioral variability and its brain-wide correlates remain lack-

ing. Our work addresses this gap by linking individual variability in cortical dynamics to behavioral variability along the anticipa-

tory–inattentive axis.

Several recent studies have analyzed behavior and neural activity in the IBL task[5,14–18,47,48]. Notably, fluctuations in task engage-

ment within single sessions have been reported[17,18], and concurrent work by Laranjeira et al. has identified multiple session-wise

behavioral phenotypes[49]. Our results complement these findings by revealing cross-animal variability in task engagement and

linking it to cortical neural dynamics.

Finally, our observations resonates with human resting-state fMRI studies reporting altered neural landscape dynamics in autism

spectrum conditions[26,50,51]. In particular, Watanabe et al. recently showed that autistic individuals exhibit more rigid network dy-

namics, resembling the inattentive landscape in our model[51]. The sex difference we observe in anticipatory bias may likewise relate

to the male predominance in autism[52]. Applying our framework to mouse models of autism-associated mutations therefore repre-

sents an important avenue for future research.

MATERIALS AND METHODS

Reaction time analysis

Definition of reaction time (RT)   For each trial, we defined the movement onset time as the first time point where the speed of

the wheel exceeds 0.5 [rad/s] after 300 ms before the stimulus onset time and following more than 50 ms of stationary wheel du-

ration during which speed is below 0.5. We picked 300 ms prior to the stimulus onset as the earliest movement onset of a trial be-

cause we found a large number of trials where a mouse starts moving the wheel right before the stimulus onset. RT was defined as the duration between the movement onset time and the stimulus onset time. If the movement onset occurs before the stimulus onset, it takes a negative value, otherwise, RT takes a positive value. Execution time (ET) was defined as the duration between the feedback time (the time when reward or noise feedback is provided) and the stimulus onset time. See Supp. Fig. S1A for the trial timeline.

Anticipatory tendency index   We defined the anticipatory tendency index by:

$$\text{Anticipatory tendency} = [\text{The ratio of trials with RT} < 0.08s] - [\text{The ratio of trials with RT} > 1.25s] \tag{1}$$

We excluded the last 40 trials from the estimation of the anticipatory tendency and other behavioral variables, to minimize the effect of satiation. A small number of trials did not have a valid RT either due to timeout without movement or potential mechanistic error; we did not exclude these trials from the denominator. Unless otherwise stated, we estimated the ratio for each animal by concatenating trials from all the sessions. In Figs. 5C-5E and 6D (excluding the last 'cortex' bars in 5E and 6D), we estimated the anticipatory tendency in a session-wise manner for a comparison with session-wise autocorrelation timescale.

Exclusion criteria   In our analysis, we only included sessions with electrophysiological recording to minimize the variability originating from task learning. Additionally, we only included sessions with more than 400 trials (exclusive). This is because a session is terminated at the 400th trial if the animal appears to be clearly disengaged [5]. In Fig. 2, to achieve a robust estimation of the animal-wise anticipatory tendency and other behavioral variables, we only included animals with two or more valid recording sessions ($N = 96$ animals).

Animal embedding algorithm

Datasets   We used behavioral data from recording sessions to study the behavioral variability among animals that have already learned the tasks. As before, we only included the sessions that had more than 400 trials (exclusive). For this embedding analysis, we defined a valid trial where task variables are not NaN, and RT and ET satisfy, $0.03s \leq RT - 0.2$ and $0.03s \leq ET$. The latter two conditions were introduced to make the logarithmic values of RT and ET valid and stable. We excluded one animal that had sessions where the number of valid trials was less than 80 %. We only included animals with two or more electrophysiology sessions as before. These criteria resulted in 371 sessions from 95 animals. We split the sessions into training and test sessions with a 90:10 split. We made sure that for each test session, there is at least one session from the same animal in the training dataset, but the training-test split was done randomly otherwise.

Inputs and outputs   Input vectors were constructed as a composition of the representation of the stimulus contrast, task block, trial number, and animal ID. Stimulus contrast vector is a nine-dimensional one-hot vector corresponding to one of nine stimulus contrasts (-1.0, -0.25, -0.125, -0.06, 0.0, 0.06, 0.125, 0.25, 1.0). Here, negative and positive contrasts represent stimuli on the left and right sides. Similarly, a block vector is a three-dimensional one-hot vector representing one of three task blocks (0.2, 0.5, 0.8). For the representation of the relative position of a given trial in a session, we used a Gaussian receptive field model. We set the activity of the i-th position encoding neuron given the t-th task as $\exp[-(\mu_i - t)^2/(2\sigma^2)]/\sqrt{2\pi\sigma^2}$, where μ_i is the selectivity of the i-th neuron and σ^2 is the variance. We used $N_{pos} = 4$ neurons for position encoding, and set $\mu_i = \frac{t_{tot}}{N_{pos}-1}i$ (for $i = 0, 1, 2, 3$), and $\sigma = \frac{t_{tot}}{2(N_{pos}-1)}$.

Animal embedding was constructed by assigning a one-hot vector to all 95 animals first, then projecting this 95-dimensional vector to a four-dimensional latent space using a learnable linear weight. The full input was then constructed by concatenating vectors representing the stimulus contrast, block, relative trial position, and animal embedding, resulting in a 20-dimensional vector.

As the target outputs, we consider two objectives simultaneously: choice and time for trial initiation and execution. For the choice, we consider the standard binary classifier with two-dimensional outputs corresponding to left and right choices. For the task time

prediction, we constructed two regression targets: RT and ET. Because the reaction time shows a bimodal distribution on the logarithmic scale, here we considered the prediction of time on the log-scale. To this end, we converted the reaction time and execution time to $\log(RT + 0.2)$ and $\log(ET)$. We excluded trials where these conversions are not well-defined.

Network architecture and learning algorithm As depicted in Fig. 3A, we used a feedforward neural network model to predict animal choice, RT, and ET. The model architecture is written as

$$\boldsymbol{h} = \phi\left(\boldsymbol{b}_2 + W_2\phi\left(\boldsymbol{b}_1 + W_1[\boldsymbol{v}_{task} \oplus \boldsymbol{v}_{pos} \oplus (W_0\boldsymbol{v}_a)]\right)\right), \tag{2a}$$

$$\hat{\boldsymbol{y}}_c = \text{softmax}[\boldsymbol{b}_c + W_c\boldsymbol{h}], \tag{2b}$$

$$\hat{y}_{rt} = \boldsymbol{w}_{rt} \cdot \boldsymbol{h} + b_{rt}, \quad \hat{y}_{et} = \boldsymbol{w}_{et} \cdot \boldsymbol{h} + b_{et}. \tag{2c}$$

Here, W_0, W_1, W_c are synaptic weight matrices, and $\boldsymbol{b}_1, \boldsymbol{b}_2, \boldsymbol{b}_c, b_{rt}, b_{et}$ are bias parameters. ϕ is the rectified linear function ($\phi(x) = \max(0, x)$) and $\boldsymbol{a} \oplus \boldsymbol{b}$ represents vector concatenation. We set the layer width of both hidden layers to be 32, thus the size of matrices W_1, W_2, and W_c are 32×20, 32×32, and 2×32, respectively. $\boldsymbol{w}_{et}$ and $\boldsymbol{w}_{et}$ are 32-dimensional vectors. The loss function was defined as

$$L = \sum_i \text{cross-entropy}(\boldsymbol{y}_i^*, \hat{\boldsymbol{y}}_{c(i)}) + \gamma \sum_i \left[(y_{rt,i}^* - \hat{y}_{rt,i})^2 + (y_{et,i}^* - \hat{y}_{et,i})^2\right]. \tag{3}$$

We set the balance parameter between the two losses γ as $\gamma = 1$, because the residual loss on the choice and RT/ET were roughly the same. We trained the network using Adam optimizer[53] with learning rate 0.001 and momentum 0.9. We used each session as a mini-batch and trained the network for 500 epochs. All weights were trained in an end-to-end fashion, meaning that the learned animal embedding is locally optimized for behavioral prediction. We implemented the algorithm using the Flax library[54].

Task generation and analysis For the latent analysis, we studied the structure of 4×95 dimensional embedding matrix W_0 using eigen-decomposition. In Figs. 3F and Supp. Figs. S3A and S3B, we generated the psychometric and chronometric curves using trained networks. In Fig. 3F, we fixed the trial number input to $t = 400$ with $t_{tot} = 800$, and set the animal embedding vector to $-0.5\boldsymbol{v}_1$, $\boldsymbol{0}$, and $0.5\boldsymbol{v}_1$ from the left to right where $\boldsymbol{v}_1$ is the principal eigenvector of the animal embedding. We then obtained the choice and RT predictions for all combinations of stimulus contrast and block to generate psychometric and chronometric curves. Similarly in Fig. S3A, we fixed the animal embedding vector to zero and set the trial number input to $t = 0, 400$, and 800 from the left to right while using $t_{tot} = 800$. This means that the left, middle, and right panels correspond to psychometric and chronometric curves at the beginning, middle, and end of a session.

In all the figures, we took the average over psychometric and chronometric curves generated from ten networks with different training/test splits and weight initialization. Because the learned animal embedding is sign invariant, we aligned the sign of PC1s across the ten obtained representations such that the negative sign corresponds to slower RT. Similarly, we aligned the sign of PC2s across ten networks such that the negative sign corresponds to the right bias.

Neural landscape model

In Fig. 4, we constructed a toy model of neural landscape $f(x)$ by

$$f(x) = \begin{cases} \frac{4b}{a^2}(x+a)^2 - b & \text{if } x < -a \\ -\frac{b}{2}\left(1 + \cos(\frac{2\pi x}{a})\right) & \text{if } -a \le x < a \\ \frac{4b}{a^2}(x-a)^2 - b & \text{if } a \le x \end{cases} \tag{4}$$

398 and simulated discretized dynamics by

$$x(t + \Delta t) = x(t) - \frac{\partial f(x)}{\partial x}\Delta t + \sigma\sqrt{\Delta t}\zeta(t), \tag{5}$$

where $\zeta(t)$ is a random variable sampled from a Gaussian distribution $\mathcal{N}(0,1)$, and $\Delta t = 0.01$ is the discretization bin size. We fixed a
at $a = 1$ and changed b across models as $b = 2, 4, 6$. Small b corresponds to a shallow landscape (blue curves), while large b yields a
deep landscape (orange curves).

In Figs. 4B and 4C, we initialized the dynamical system at $x = 0$, then ran the dynamics under $\sigma = 1.75$. After 50 time steps (at t =
0.5), we introduced a constant positive input v. We then evaluated the time it takes until the first time the state crosses the bump
on the right and achieves $x > a$. This is known as the first-passage time[55]. We subtracted the first 50 time steps from the estimated
duration. False-start rate was estimated by the ratio of trials where the dynamics reached $x > a$ before the input is provided (i.e.,
within 50 time steps).

In Fig. 4D, we set the noise amplitude $\sigma = 2.0$, ran the dynamics for 10^6 time steps, and estimated the autocorrelation curves.

Autocorrelation timescale estimation

409 **Autocorrelation fitting**   Autocorrelation curves typically contain multiple exponential factors corresponding to neuronal and network-
410 level autocorrelation timescales in addition to oscillatory components. We thus fitted the following function:

$$a(t) = \left(C_0 + C_1 e^{-t/\tau_1} + C_2 e^{-t/\tau_2}\right)\left(1 + C_3 \cos\omega t\right). \tag{6}$$

We binned population activity of a given region by 0.01-second bins, and calculated autocorrelation up to a 1.5-second delay. Pa-
rameters $(C_0, C_1, C_2, C_3, \tau_1, \tau_2, \omega)$ were estimated using a trust-region method implemented in the scipy curve-fit function. We sam-
pled initial values from a uniform distribution where the ranges were chosen as $[0.01, 1.0]$ for (C_0, C_1, C_2, C_3), $[0.001, 0.03]$ for τ_1, $[0.03, 1.0]$
for τ_2, and $[0.0, 2\pi/10]$ for ω. We picked different initial ranges for τ_1 and τ_2 to encourage the symmetry breaking. We bounded all co-
efficients to be non-negative, and τ_1 and τ_2 to be lower-bounded by 0.001 [second] for stability. We set the maximum iteration of
curve-fit to 10^6, and repeated fitting from 10^4 random initializations, then picked the parameters achieving the best fit. We picked
the $\max(\tau_1, \tau_2)$ as the autocorrelation timescale. This is because the exponential decay with faster timescale is expected to corre-
spond to single-neuron level timescale, not the network timescale[34].

In Supp Fig. 4B, we compared the fitting error of this function and a standard double exponential fitting:

$$a(t) = C_1 e^{-t/\tau_1} + C_2 e^{-t/\tau_2}. \tag{7}$$

The error was defined by $\epsilon = \sqrt{\frac{1}{T}\sum_{t=0}^{T-1}(a(t) - a^*(t))^2}$, where $a^*(t)$ is the empirical estimate of the autocorrelation curve.
In the ITI fitting, we again binned population activity of a given region by 0.01-second bin, but calculated autocorrelation only up
to 0.75-second delay, as the short ITI makes estimation of a long-delay autocorrelation infeasible. We used the double exponential
fitting (Eq. 7), because of this short delay and the oscillation was generally not strong during ITI compared to the passive period. We
used the same fitting protocol as above, but set the maximum iteration of curve-fit to 10^5, and repeated fitting from 10^3 random
initializations due to the presence of fewer free parameters.

Exclusion criteria   For the passive period analysis, we only included sessions where the passive period was clearly tagged in the
data, resulting in fewer sessions than the sessions used for the ITI analysis. Moreover, we only included sessions where the total num-
ber of neurons recorded from the given region is equal to or larger than 10, and the total number of spikes in the passive period is
larger than 10^4, to reduce the noise in the empirical autocorrelation estimation. For the ITI analysis, we set the same criteria for the
minimum number of neurons and the cumulative number of spikes during ITIs. We did not impose any criteria on the sorting qual-
ity for inclusion of a neuron for the analysis except for Supp. Fig. S4J and S6A.

After fitting, we imposed two more criteria: First, we only used the fitting where the fitting error is lower than 0.025 (orange verti-

cal line in Supp. Fig. S4B). Secondly, we excluded the fitting where the estimated relative timescale is outside of $\frac{1}{3}\tau_{median} \leq \tau <$

$3\tau_{median}$, where τ_{median} is the median timescale of the given region (see black vertical line in Supp. Fig. S4C). We imposed this sec-

ond criterion because it is unlikely to observe a ten-fold difference in autocorrelation timescale across individuals.

Unbiased autocorrelation estimation using bootstrapped mean and variance Given a sequence $\{x(t_i)\}_{i=1}^N$ of equally sampled

time points $(t_{i+j} - t_i = j\Delta t)$, the standard definition of autocorrelation follows [33,34]

$$\hat{a}_{corr}(j\Delta t) \equiv \frac{1}{(N-j)\hat{\sigma}^2} \sum_{i=1}^{N-j} \left(x(t_i) - \hat{\mu}_j^{(l)}\right)\left(x(t_{i+j}) - \hat{\mu}_j^{(r)}\right), \tag{8}$$

438 where the left and right means $\hat{\mu}_j^{(l)}, \hat{\mu}_j^{(r)}$ and the variance $\hat{\sigma}^2$ are given by

$$\hat{\mu}_j^{(l)} \equiv \frac{1}{N-j}\sum_{i=1}^{N-j} x(t_i), \quad \hat{\mu}_j^{(r)} \equiv \frac{1}{N-j}\sum_{i=j+1}^{N} x(t_i), \quad \hat{\sigma}^2 \equiv \frac{1}{N}\sum_{i=1}^{N}\left(x(t_i) - \tfrac{1}{N}\sum_{k=1}^{N} x(t_k)\right)^2. \tag{9}$$

However, this estimation is known to be biased when the total duration of the sequence $t_N - t_1$ is comparable to the typical auto-

correlation timescale [32,33]. This is because there exists an intrinsic correlation between $x(t_i)$ and $\hat{\mu}$ (see *Supplementary text*).

One way to overcome this bias is to consider inference of the underlying generative model [34]. However, this method is computa-

tionally expensive as it requires sample generation from the generative model at every iteration of optimization. An alternative ap-

proach we propose here is to bootstrap the mean and variance from nearby trials. More specifically, given a population activity se-

quence at the k-th trial $\{x_k(t_i)\}_{i=1}^N$, we estimate the autocorrelation at the k-th trial by

$$\hat{a}_{k,corr}(j\Delta t) \equiv \frac{1}{(N-j)\hat{\sigma}_k^2} \sum_{i=1}^{N-j} \left(x(t_i) - \hat{\mu}_k\right)\left(x(t_{i+j}) - \hat{\mu}_k\right), \tag{10}$$

445 where

$$\hat{\mu}_k \equiv \frac{1}{N(2L+1)} \sum_{k'=k-L}^{k+L} \sum_{i=1}^{N} x_{k'}(t_i), \quad \hat{\sigma}_k^2 \equiv \frac{1}{N(2L+1)} \sum_{k'=k-L}^{k+L} \sum_{i=1}^{N}(x_{k'}(t_i) - \hat{\mu}_k)^2. \tag{11}$$

Here, L specifies the range of trials we use to estimate the mean and variance. Because the true mean and variance underlying ITI

population activity are expected to shift during the session, it is not appropriate to use all trials for estimating $\hat{\mu}_k$ and $\hat{\sigma}_k$. By contrast,

if we only use one or two neighboring trials to estimate these values, the estimation would be unreliable. In the analysis, we fixed L

to be 15, which yielded autocorrelation curves without a dip or a constant factor (Fig. 6C).

Cortex-wide relative autocorrelation timescale We estimated the cortex-wide relative timescales plotted in Figs. 5F, 5G, and

6E-6G as follow. Let us denote the estimated autocorrelation timescale of region r in subject s by τ_s^r, the set of recorded regions

from subject s by Ω_s, and the set of subjects with recording from region r by Γ_r. Then the relative autocorrelation timescale of sub-

ject s is defined by

$$\bar{\tau}_s \equiv \frac{1}{|\Omega_s|} \sum_{r \in \Omega_s} \frac{\tau_s^r}{\frac{1}{|\Gamma_r|}\sum_{s' \in \Gamma_r} \tau_{s'}^r}, \tag{12}$$

454 where $|\Omega_s|$ and $|\Gamma_r|$ represents the size of the sets Ω_s and Γ_r, respectively. We used the same procedure to estimate the cortex-wide

455 relative firing rates.

456 **Firing rate analysis** In the firing rate analysis, we simply estimated the average firing rate per neuron during the passive period

457 and ITI. We applied the same criteria on the minimum total number of neurons per region and the total number of spikes.

Wheel statistics analysis In Figs. 5I and 6I, we calculated summary statistics of the wheel trajectories $w(t)$ during the passive period and ITI as follow. The wheel distance was defined as the total movement of the wheel per second, $\frac{1}{T} \int_0^T w(t)dt$. The active fraction is the ratio of active wheel movement period defined as $\frac{1}{T} \int_0^T$ heaviside$[\dot{w}(t) - v_{th}]dt$ where $\dot{w}(t)$ is the speed of the wheel at time t and $v_{th} = 0.5$ rad/s. The energy is given by $\frac{1}{T} \int_0^T \frac{1}{2} (\dot{w}(t))^2 \, dt$. The movement onsets were calculated in the same way we calculated the movement onset of a trial (see *Definition of reaction time* in Methods), and the value was defined as the number of movement onsets per second. Lastly, the change point is the point where the velocity changes sign. Given a discrete observation of the wheel trajectory at time $t_1, t_2, ...$, time point t_i is a change point if $(w(t_{i-1}) - w(t_i))(w(t_i) - w(t_{i+1})) < 0$. The number of change points was defined as the change points per second.

CODE AVAILABILITY

Codes for replicating all the results are available at **https://github.com/nhiratani/IBL_variability**.

DATA AVAILABILITY

Both the behavioral and neural data we analyzed are publicly available through The International Brain Laboratory.

ACKNOWLEDGMENTS

The authors thank Peter Latham for his help with the project conceptualization and discussion. This work was partially supported by the McDonnell Center for Systems Neuroscience.

AUTHOR COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

[1] Paola Palanza, Laura Gioiosa, and Stefano Parmigiani. Social stress in mice: gender differences and effects of estrous cycle and social dominance. *Physiology & behavior*, 73(3):411–420, 2001.

[2] Yasuyuki Horii, Tatsuhiro Nagasawa, Hiroyuki Sakakibara, Aki Takahashi, Akira Tanave, Yuki Matsumoto, Hiromichi Nagayama, Kazuto Yoshimi, Michiko T Yasuda, Kayoko Shimoi, et al. Hierarchy in the home cage affects behaviour and gene expression in group-housed c57bl/6 male mice. *Scientific reports*, 7(1):6991, 2017.

[3] Katherine B LeClair, Kenny L Chan, Manuella P Kaster, Lyonna F Parise, Charles Joseph Burnett, and Scott J Russo. Individual history of winning and hierarchy landscape influence stress susceptibility in mice. *Elife*, 10:e71401, 2021.

[4] Cathy S Chen, Evan Knep, Autumn Han, R Becket Ebitz, and Nicola M Grissom. Sex differences in learning from exploration. *Elife*, 10:e69748, 2021.

[5] International Brain Laboratory, Valeria Aguillon-Rodriguez, Dora Angelaki, Hannah Bayer, Niccolo Bonacchi, Matteo Carandini, Fanny Cazettes, Gaelle Chapuis, Anne K Churchland, Yang Dan, et al. Standardized and reproducible measurement of decision-making in mice. *Elife*, 10:e63711, 2021.

[6] Samuel Liebana, Aeron Laffere, Chiara Toschi, Louisa Schilling, Jessica Moretti, Jacek Podlaski, Matthias Fritsche, Peter Zatka-Haas, Yulong Li, Rafal Bogacz, et al. Dopamine encodes deep network teaching signals for individual learning trajectories. *Cell*, 2025.

[7] Tomoki Kurikawa, Tatsuya Haga, Takashi Handa, Rie Harukuni, and Tomoki Fukai. Neuronal stability in medial frontal cortex sets individual variability in decision-making. *Nature Neuroscience*, 21(12):1764–1773, 2018.

[8] Marino Pagan, Vincent D Tang, Mikio C Aoi, Jonathan W Pillow, Valerio Mante, David Sussillo, and Carlos D Brody. Individual variability of neural computations underlying flexible decisions. *Nature*, 639(8054):421–429, 2025.

[9] Editorial. Troublesome variability in mouse studies. *Nat. Neurosci.*, 12(1075), 2009.

[10] Rachel Michelle Saré, Abigail Lemons, and Carolyn Beebe Smith. Behavior testing in rodents: highlighting potential confounds affecting variability and reproducibility. *Brain sciences*, 11(4):522, 2021.

[11] Simon Musall, Anne E Urai, David Sussillo, and Anne K Churchland. Harnessing behavioral diversity to understand neural computations for cognition. *Current opinion in neurobiology*, 58:229–238, 2019.

[12] Bernhard Voelkl, Naomi S Altman, Anders Forsman, Wolfgang Forstmeier, Jessica Gurevitch, Ivana Jaric, Natasha A Karp, Martien J Kas, Holger Schielzeth, Tom Van de Casteele, et al. Reproducibility of animal research in light of biological variation. *Nature Reviews Neuroscience*, 21(7):384–393, 2020.

[13] Ariel Gilad, Yasir Gallero-Salas, Dominik Groos, and Fritjof Helmchen. Behavioral strategy determines frontal or posterior location of short-term memory in neocortex. *Neuron*, 99(4):814–828, 2018.

[14] International Brain Laboratory, Brandon Benson, Julius Benson, Daniel Birman, Niccolò Bonacchi, Kcénia Bougrova, Sebastian A Bruijns, Matteo Carandini, Joana A Catarino, Gaelle A Chapuis, et al. A brain-wide map of neural activity during complex behaviour. *biorxiv*, pages 2023–07, 2023.

[15] Sebastian A Bruijns, International Brain Laboratory, Kcénia Bougrova, Inês C Laranjeira, Petrina YP Lau, Guido T Meijer, Nathaniel J Miska, Jean-Paul Noel, Alejandro Pan-Vazquez, Noam Roth, et al. Dissecting the complexities of learning with infinite hidden markov models. *bioRxiv*, pages 2023–12, 2023.

[16] International Brain Laboratory, Kush Banga, Julius Benson, Jai Bhagat, Dan Biderman, Daniel Birman, Niccolò Bonacchi, Sebastian A Bruijns, Kelly Buchanan, Robert AA Campbell, et al. Reproducibility of in vivo electrophysiological measurements in mice. *bioRxiv*, pages 2022–05, 2022.

[17] Zoe C Ashwood, Nicholas A Roy, Iris R Stone, International Brain Laboratory, Anne E Urai, Anne K Churchland, Alexandre Pouget, and Jonathan W Pillow. Mice alternate between discrete strategies during perceptual decision-making. *Nature Neuroscience*, 25(2):201–212, 2022.

[18] Lenca I Cuturela, International Brain Laboratory, and Jonathan W Pillow. Internal states emerge early during learning of a perceptual decision-making task. *bioRxiv*, pages 2024–11, 2024.

[19] Rafal Bogacz, Eric Brown, Jeff Moehlis, Philip Holmes, and Jonathan D Cohen. The physics of optimal decision making: a formal analysis of models of performance in two-alternative forced-choice tasks. *Psychological review*, 113(4):700, 2006.

[20] Joshua I Gold and Michael N Shadlen. The neural basis of decision making. *Annu. Rev. Neurosci.*, 30(1):535–574, 2007.

[21] Jan Drugowitsch, Rubén Moreno-Bote, Anne K Churchland, Michael N Shadlen, and Alexandre Pouget. The cost of accumulating evidence in perceptual decision making. *Journal of Neuroscience*, 32(11):3612–3628, 2012.

[22] Gaurav Malhotra, David S Leslie, Casimir JH Ludwig, and Rafal Bogacz. Overcoming indecision by changing the decision boundary. *Journal of Experimental Psychology: General*, 146(6):776, 2017.

[23] Philippa A Johnson, Sander Nieuwenhuis, and Anne Urai. Pupil dynamics preceding switches in task engagement. In *Proceedings of the Annual Meeting of the Cognitive Science Society*, volume 46, 2024.

[24] Pauline Van der Wel and Henk Van Steenbergen. Pupil dilation as an index of effort in cognitive control tasks: A review. *Psychonomic bulletin & review*, 25:2005–2015, 2018.

[25] Nikola Grujic, Rafael Polania, and Denis Burdakov. Neurobehavioral meaning of pupil size. *Neuron*, 112(20):3381–3395, 2024.

[26] Takamitsu Watanabe, Geraint Rees, and Naoki Masuda. Atypical intrinsic neural timescale in autism. *elife*, 8:e42256, 2019.

[27] Norman H Lam, Thiago Borduqui, Jaime Hallak, Antonio Roque, Alan Anticevic, John H Krystal, Xiao-Jing Wang, and John D Murray. Effects of altered excitation-inhibition balance on decision making in a cortical circuit model. *Journal of Neuroscience*, 42(6):1035–1053, 2022.

[28] Quanxin Wang, Song-Lin Ding, Yang Li, Josh Royall, David Feng, Phil Lesnar, Nile Graddis, Maitham Naeemi, Benjamin Facer, Anh Ho, et al. The allen mouse brain common coordinate framework: a 3d reference atlas. *Cell*, 181(4):936–953, 2020.

[29] Andrei Khilkevich, Michael Lohse, Ryan Low, Ivana Orsolic, Tadej Bozic, Paige Windmill, and Thomas D Mrsic-Flogel. Brain-wide dynamics linking sensation to action during decision-making. *Nature*, 634(8035):890–900, 2024.

[30] Julie A Harris, Stefan Mihalas, Karla E Hirokawa, Jennifer D Whitesell, Hannah Choi, Amy Bernard, Phillip Bohn, Shiella Caldejon, Linzy Casal, Andrew Cho, et al. Hierarchical organization of cortical and thalamic connectivity. *Nature*, 575(7781):195–202, 2019.

[31] Xiao-Jing Wang. Macroscopic gradients of synaptic excitation and inhibition in the neocortex. *Nature reviews neuroscience*, 21(3):169–178, 2020.

[32] JA Pope and FHC Marriott. Errors in the estimation of serial correlations. *Nature*, 172(4382):778–778, 1953.

[33] FHC Marriott and JA Pope. Bias in the estimation of autocorrelations. *Biometrika*, 41(3/4):390–402, 1954.

[34] Roxana Zeraati, Tatiana A Engel, and Anna Levina. A flexible bayesian framework for unbiased estimation of timescales. *Nature computational science*, 2(3):193–204, 2022.

[35] Roxana Zeraati, Yan-Liang Shi, Nicholas A Steinmetz, Marc A Gieselmann, Alexander Thiele, Tirin Moore, Anna Levina, and Tatiana A Engel. Intrinsic timescales in the visual cortex change with selective attention and reflect spatial connectivity. *Nature communications*, 14(1):1858, 2023.

[36] Mackenzie Weygandt Mathis and Alexander Mathis. Deep learning tools for the measurement of animal behavior in neuroscience. *Current opinion in neurobiology*, 60:1–11, 2020.

[37] Carsen Stringer, Marius Pachitariu, Nicholas Steinmetz, Charu Bai Reddy, Matteo Carandini, and Kenneth D Harris. Spontaneous behaviors drive multidimensional, brain-wide activity. *Science*, 364(6437):eaav7893, 2019.

[38] Lucas Pinto, Michael J Goard, Daniel Estandian, Min Xu, Alex C Kwan, Seung-Hee Lee, Thomas C Harrison, Guoping Feng, and Yang Dan. Fast modulation of visual perception by basal forebrain cholinergic neurons. *Nature neuroscience*, 16(12):1857–1863, 2013.

[39] David A McCormick, Dennis B Nestvogel, and Biyu J He. Neuromodulation of brain state and behavior. *Annual review of neuroscience*, 43(1):391–415, 2020.

[40] Michael Nilsson, Ekaterina Perfilieva, Ulf Johansson, Owe Orwar, and Peter S Eriksson. Enriched environment increases neurogenesis in the adult rat dentate gyrus and improves spatial memory. *Journal of neurobiology*, 39(4):569–578, 1999.

[41] Jeremy G Turner, Jennifer L Parrish, Larry F Hughes, Linda A Toth, and Donald M Caspary. Hearing in laboratory animals: strain differences and nonauditory effects of noise. *Comparative medicine*, 55(1):12–23, 2005.

[42] Maria Gulinello, Heather A Mitchell, Qiang Chang, W Timothy O'Brien, Zhaolan Zhou, Ted Abel, Li Wang, Joshua G Corbin, Surabi Veeraragavan, Rodney C Samaco, et al. Rigor and reproducibility in rodent behavioral research. *Neurobiology of learning and memory*, 165:106780, 2019.

[43] Valeria Fascianelli, Aldo Battista, Fabio Stefanini, Satoshi Tsujimoto, Aldo Genovesio, and Stefano Fusi. Neural representational geometries reflect behavioral differences in monkeys and recurrent neural networks. *Nature Communications*, 15(1):6479, 2024.

[44] Michael J Frank, Bradley B Doll, Jen Oas-Terpstra, and Francisco Moreno. Prefrontal and striatal dopaminergic genes predict individual differences in exploration and exploitation. *Nature neuroscience*, 12(8):1062–1068, 2009.

[45] Emily S Finn, Xilin Shen, Dustin Scheinost, Monica D Rosenberg, Jessica Huang, Marvin M Chun, Xenophon Papademetris, and R Todd Constable. Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. *Nature neuroscience*, 18(11):1664–1671, 2015.

[46] Caterina Gratton, Timothy O Laumann, Ashley N Nielsen, Deanna J Greene, Evan M Gordon, Adrian W Gilmore, Steven M Nelson, Rebecca S Coalson, Abraham Z Snyder, Bradley L Schlaggar, et al. Functional brain networks are dominated by stable group and individual factors, not cognitive or daily variation. *Neuron*, 98(2):439–452, 2018.

[47] Charles Findling, Felix Hubert, International Brain Laboratory, Luigi Acerbi, Brandon Benson, Julius Benson, Daniel Birman, Niccolò Bonacchi, Sebastian Bruijns, Matteo Carandini, et al. Brain-wide representations of prior information in mouse decision-making. *BioRxiv*, pages 2023–07, 2023.

[48] Lorenzo Posani, Shuqi Wang, Samuel P Muscinelli, Liam Paninski, and Stefano Fusi. Rarely categorical, always high-dimensional: how the neural code changes along the cortical hierarchy. *bioRxiv*, pages 2024–11, 2025.

[49] Ines Laranjeira, Matthew Whiteway, International Brain Laboratory, Zach Mainen, and Daniel McNamee. The structure of individuality in micro-behavioral features of task performance. In *Computational and Systems Neuroscience (Cosyne)*, Montreal, Canada, March 2025.

[50] Lisa E Mash, Annika C Linke, Lindsay A Olson, Inna Fishman, Thomas T Liu, and Ralph-Axel Müller. Transient states of network connectivity are atypical in autism: A dynamic functional connectivity study. *Human brain mapping*, 40(8):2377–2389, 2019.

[51] Takamitsu Watanabe and Hidenori Yamasue. Noninvasive reduction of neural rigidity alters autistic behaviors in humans. *Nature Neuroscience*, pages 1–13, 2025.

[52] Sarah L Ferri, Ted Abel, and Edward S Brodkin. Sex differences in autism spectrum disorder: a review. *Current psychiatry reports*, 20:1–17, 2018.

[53] Diederik P Kingma and Jimmy Ba. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.

[54] Jonathan Heek, Anselm Levskaya, Avital Oliver, Marvin Ritter, Bertrand Rondepierre, Andreas Steiner, and Marc van Zee. Flax: A neural network library and ecosystem for JAX, 2024.

[55] Hannes Risken and Hannes Risken. *Fokker-planck equation*. Springer, 1996.

SUPPLEMENTARY FIGURES

Supplementary Figure 1: Reaction time statistics (A) Step-by-step description of a trial. Note that, below, we denote the combination of the interval period and quiescent period as the inter-trial interval. **(B-E)** Examples of wheel trajectories in early response trials (B,C) and late response trials (D,E). As in Fig. 1B, blue, orange, and green lines represent the stimulus onset, movement onset, and the feedback time. While ballistic responses shown in panels B and D are the most common, there were also trials where a mouse changes their mind in the middle (panel C), or stops the wheel in the middle and moves it again in the same direction (panel E). **(F)** The probability of the change of mind in early and normal response time trials. A trial was classified as a change of mind trial if the directions of the first movement and the final choice are different as in panel C. We estimated this probability for each signed stimulus contrast in congruent and incongruent trials. Congruent trials are defined as trials where the direction of the first movement is the correct direction, whereas incongruent trials are the ones where the first wheel movement is the opposite of the correct direction. **(G)** Error rate as a function of the reaction time in low contrast (gray) and high contrast (black) trials. **(H)** Distribution of inter-trial intervals preceding early, normal, and late trials. ITI was defined as the duration between the end of the previous trial and the stimulus onset of the new trial. In a reward trial, the trial ends one second after the feedback time, while in an error trial, the trial ends two seconds after the feedback time.

Supplementary Figure 2: Reaction time variability (A) Average early and late response ratios for each lab. As in Fig. 2, lab L is an outlier. **(B)** The inter-animal variability in the task acquisition speed, measured by the total number of sessions an animal performed before starting an electrophysiology session (including habituation sessions), was not correlated with the anticipatory tendency. Points are color-coded by the lab identity as in Fig. 2E. **(C)** The standard deviation of the median RT across sessions for each mouse. The gray histogram is a shuffle estimated by randomly re-assigning sessions to animals. **(D)** The ITI distributions of anticipatory (ATI > 0.15), normal, inattentive mice (ATI < -0.15). As in Supp. Fig. S1H, we included the quiescent period into ITI.

Supplementary Figure 3: Psychometric and chronometric curves generated from animal-to-vec analysis (A) Effect of within-task position on psychometric and chronometric curves. Positions 0, 0.5, and 1.0 correspond to the beginning, middle, and end of a session. **(B)** Shift in expected behavior along the second principal vector of the animal embedding. **(C)** Average task performance and RT along the PC2 of the animal embedding.

Supplementary Figure 4: Supplementary analysis of passive period autocorrelation timescale (A) Average autocorrelation of VISa population activity for inattentive (ATI < -0.15), normal, and anticipatory (ATI > 0.15) mice. Error bars represent the standard error of the mean across sessions. **(B)** The fitting error of the autocorrelation curve with or without oscillation components. The orange vertical line represents the cutoff for inclusion in the subsequent analysis. **(C)** The distribution of the estimated relative timescale. We excluded both overly small and large estimates from the analysis, as the inter-animal variance in autocorrelation is expectedly small. In panels B and C, if a session includes recording from multiple regions, they are counted independently. **(D)** Correlation between average firing rate of regions and divisions, and session-wise anticipatory tendency. **(E)** Estimated average cortical timescale and its relationship with the cortical hierarchy score [30]. **(F)** The same as Fig. 5E, but represented in a flattened cortical map. **(G)** Correlation between the relative cortical timescale of each animal and their behavioral statistics. We found a correlation with both early and late response ratios as they contribute equally to the anticipatory tendency. PC1 projection is the projection to the PC1 axis of the animal embedding. **(H, I)** Correlation between anticipatory tendency and the active fraction ($r = -0.03, p = 0.8$; panel H) and the change points ($r = 0.20, p = 0.04$; panel I) of wheel trajectory during the passive period (see *Wheel statistics analysis* in Methods for the definition of wheel statistics). **(J)** The same as Fig. 5F, but only the neurons with the sorting quality criteria = 1.0 were used. For the analysis in the main text, we included all neurons regardless of the sorting quality, as we analyzed population-level autocorrelation.

Supplementary Figure 5: Supplementary analysis of neural autocorrelation and behaviors during ITI (A) The correlation between the autocorrelation timescales at VISa and the session-wise anticipatory tendency ($p = 0.003$ without Bonferroni correction). Here, each point represents one session. **(B)** The same as Fig. 6D, but represented in a flattened cortical map. **(C)** Estimated average cortical timescale and its relationship with the hierarchy score [30]. **(D)** Correlation between average firing rate of regions and divisions, and session-wise anticipatory tendency. **(E)** Correlation between the relative cortical timescale of each animal and their behavioral statistics. **(F,G)** Correlation between anticipatory tendency and the active fraction ($r = -0.12$, $p = 0.16$; panel F) and change points ($r = 0.11$, $p = 0.2$; panel G) of wheel trajectory during ITI.

Supplementary Figure 6: Correlation between animal-wise anticipatory tendency and relative cortical timescale under various model variants. (A) Only the neurons with spike-sorting quality = 1.0 are included, as opposed to QC = 0.0 we used in the main text. **(B)** The entire ITI period, defined as the duration between the beginning of ITI to 0.5 seconds before the stimulus onset, was used instead of the initial fraction of ITI. **(C, D)** The mean and variance of autocorrelation were estimated using 10 (C) and 50 (D) trial windows, instead of 30 trial windows in the main text.

SUPPLEMENTARY TEXT: BIAS IN AUTOCORRELATION ESTIMATION

Here, we prove that autocorrelation estimation from a short interval is biased and the proposed method mitigates this bias, assuming that ITI activity is sampled independently from an Gaussian Process. Let us consider estimation of the autocorrelation from a set of M time series, each containing observation at $t = t_1, .., t_N$. We generate a set of M time series $x_i(t_1), ..., x_i(t_N)$ by a Gaussian Process with kernel function $k(\cdot, \cdot)$:

$$x_i \sim \mathcal{N}(\mu; K), \tag{13}$$

where $x_i \equiv [x(t_1), ..., x(t_N)]$. μ is an N-dimensional vector where all the elements are μ, and K is an $N \times N$ covariance matrix whose (i, j)-th element is $k(t_i, t_j)$. Below, we restrict kernel function to stationary kernel (i.e., $k(t_i, t_j) = k(t_i - t_j)$) for clarity, but the same argument holds for non-stationary kernels as well.

Suppose $t_1, .., t_N$ are sampled with the equal interval Δt (i.e., $t_k = k\Delta t$). Then, empirical estimation of auto-covariance for a single time series x is commonly defined by

$$\hat{a}_{cov}(j\Delta t) \equiv \frac{1}{(N-j)} \sum_{i=1}^{N-j} \left(x(t_i) - \hat{\mu}_j^{(l)} \right) \left(x(t_{i+j}) - \hat{\mu}_j^{(r)} \right), \tag{14}$$

where the empirical left and right means are defined by

$$\hat{\mu}_j^{(l)} \equiv \frac{1}{N-j} \sum_{i=1}^{N-j} x(t_i), \quad \hat{\mu}_j^{(r)} \equiv \frac{1}{N-j} \sum_{i=j+1}^{N} x(t_i). \tag{15a}$$

Below, we analyze autocovariance rather than autocorrelation for simplicity.

Let us first demonstrate that the above estimation of autocovariance is biased even at $M \to \infty$ limit. For the ease of notation, we define $\tilde{x}(t)$ as the zero-meaned $x(t)$:

$$\tilde{x}(t) \equiv x(t) - \mu. \tag{16}$$

Taking $M \to \infty$ limit and considering the expectation of $\hat{a}_{cov}(j\Delta t)$ over the Gaussian process, we get

$$
\begin{aligned}
\langle \hat{a}_{cov}(j\Delta t) \rangle &= \left\langle \frac{1}{(N-j)} \sum_{i=1}^{N-j} \left(x(t_i) - \hat{\mu}_j^{(l)} \right) \left(x(t_{i+j}) - \hat{\mu}_j^{(r)} \right) \right\rangle \\
&= \left\langle \frac{1}{(N-j)} \sum_{i=1}^{N-j} \left([x(t_i) - \mu] - [\hat{\mu}_j^{(l)} - \mu] \right) \left([x(t_{i+j}) - \mu] - [\hat{\mu}_j^{(r)} - \mu] \right) \right\rangle \\
&= \frac{1}{(N-j)} \sum_{i=1}^{N-j} \left\langle \tilde{x}(t_i)\tilde{x}(t_{i+j}) - \frac{1}{N-j} \sum_{k=1}^{N-j} [\tilde{x}(t_k)\tilde{x}(t_{i+j}) + \tilde{x}(t_i)\tilde{x}(t_{k+j})] + \frac{1}{(N-j)^2} \sum_{k=1}^{N-j} \sum_{l=1}^{N-j} \tilde{x}(t_k)\tilde{x}(t_{l+j}) \right\rangle \\
&= k(j\Delta t) - \frac{1}{(N-j)^2} \sum_{i=1}^{N-j} \sum_{k=1}^{N-j} [k(|i+j-k|\Delta t) + k(|k+j-i|\Delta t)] + \frac{1}{(N-j)^2} \sum_{k=1}^{N-j} \sum_{l=1}^{N-j} k(|l+j-k|\Delta t) \\
&= k(j\Delta t) - \frac{1}{(N-j)^2} \sum_{i=1}^{N-j} \sum_{k=1}^{N-j} k(|i+j-k|\Delta t). \tag{17}
\end{aligned}
$$

The equation above clearly shows that the empirical estimate of autocovariance $\hat{a}_{cov}$ is biased against the true auto-covariance $k(j\Delta t)$. This bias also results in the biased estimate of autocorrelation (trial-wise estimate of variance induces further bias). The bias term is $\mathcal{O}(1)$ when $k(|i+j-k|\Delta t)$ is non-zero for most (i, j, k) combination. On the other hand, if N is large, such that the kernel $k(|i+j-k|\Delta t)$ decays to zero for most (i, j, k) combination, then the bias scales with $\mathcal{O}\left(\frac{1}{N^2}\right)$.

A solution is to replace $\hat{\mu}$ with the average activity over nearby trials. Specifically, we can define auto-covariance of the μ-th time series $x_\mu = [x_\mu(t_1), ..., x_\mu(t_N)]$ by

$$\hat{a}_{\mu,cov}^M(j\Delta t) \equiv \frac{1}{(N-j)} \sum_{i=1}^{N-j} \left(x_\mu(t_i) - \hat{\mu}_j^{M,(l)} \right) \left(x_\mu(t_{i+j}) - \hat{\mu}_j^{M,(r)} \right), \tag{18}$$

where the empirical left and right means are defined by

$$\hat{\mu}_j^{M,(l)} \equiv \frac{1}{(N-j)M} \sum_{\nu=1}^{M} \sum_{i=1}^{N-j} x_\nu(t_i), \tag{19a}$$

$$\hat{\mu}_j^{M,(r)} \equiv \frac{1}{(N-j)M} \sum_{\nu=1}^{M} \sum_{i=j+1}^{N} x_\nu(t_i). \tag{19b}$$

In this definition, from a parallel calculation with the one above, the expectation over GP is given by

$$\hat{a}_{\mu,cov}^M(j\Delta t) = k(j\Delta t) - \frac{1}{M(N-j)^2} \sum_{i=1}^{N-j} \sum_{k=1}^{N-j} k(|i+j-k|\Delta t). \tag{20}$$

Thus, compared to Eq. 17, trial averaging reduces the bias by $1/M$ where M is the total number of trials used for the estimation of the mean.

This result indicates that it is best to use all the trials to minimize the bias. However, in the actual neural data, the i.i.d. assumption is violated. Thus, using all the trial may induce additional bias into the estimation of the mean activity. To navigate this balance, in the analysis, we used $M = 30$ for our analysis in the main result. Note that, by estimating the left and right means from different trials, we can further reduce the bias in principle, but it increases the variance in the estimate.