Heritable and environmental influences on innate immunity across three populations of threespine stickleback (*Gasterosteus aculeatus*)

Cole J. Wolf[1*], Panna A. Codner[1], Jesse N. Weber[1]

[1]Department of Integrative Biology, University of Wisconsin–Madison, Madison, WI 53706, USA

*Correspondence:
Corresponding Author
colejwolf@gmail.com

Keywords: threespine stickleback, innate immunity, ecoimmunology, microbial environment,
16S DNA, granulocytes, ROS.

Abstract

Environmental variation plays a key role in immune development and function; factors such as
pathogen exposure history, seasonality, and resource availability all affect an individual's immune
phenotype. However, the relative contributions of heritable and non-heritable factors remain unclear
for most immune phenotypes. We used populations of threespine stickleback (*Gasterosteus
aculeatus*) with heritable differences in immune function to investigate the relationship between
immunity, genetic divergence, and the environment. Fish were raised in tanks with different flow
rates (continuous or intermittent) that harbored differing microbial communities. After long-term
acclimation to one tank environment, a subset of adult fish were moved to the alternate flow regime
and allowed to acclimate for eighteen weeks. We then measured the effects of starting environment,
transfer between environments, and final environment across several immune parameters. Both
population and treatment significantly affected immune function. Sticklebacks from a population
previously found to display the highest heritable levels of innate immunity displayed the highest
oxidative burst capacity (ROS) regardless of tank environment. However, all fish in intermittent flow
tanks (both resident and transfers) tended to have higher ROS production, more granulocytes, and
greater spleen mass than in continuous flow. Variation in liver mass was mainly driven by population
effects. Together, this work demonstrates that a simple change in water flow dynamics can shift
aquarium microbial conditions, creating opportunities to explore immune flexibility. Perhaps more
importantly, we also highlight the need for further research examining how naturally evolved genetic
differences and environmental factors individually and jointly influence the magnitude and direction
of immune responses.

Introduction

Differential immune responses facilitate an organism's ability to maintain high fitness in the face of
changing pathogen risk. While some elements of immunity are highly innate, others are influenced
primarily by either environment or by genotype-by-environment interactions. Ecoimmunology has
traditionally focused on environmental factors such as seasonality, resource availability, and
exposure to pathogens (1). This is largely due to difficulty in identifying heritable differences in wild
organisms, which is further complicated when environmental influences vary across the life of an
organism. For example, many species have a critical window where immune system development is
sensitive to external inputs. In these cases, the adult immune function is contingent on early life
exposures (2,3). In contrast, other aspects of immunity display plastic shifts depending on current

environmental conditions. This includes the large body of work examining how exposure to toxins
impacts immunity and disease e.g., (4,5). However, a large research gap exists when considering
whether environmentally induced immune shifts are common to all individuals of a species or
parceled into genotype-specific effects where only some individuals respond to cues (i.e., genotype-
environment interaction; reviewed in Bolnick et al. (6).

The microbial environment surrounding an organism has strong impacts on host immunity. In
mammals, the gut microbiome plays an important role in immune development and function,
including maintenance of homeostasis and disease resistance (7). Much of what we know in this area
comes from laboratory work on captive, inbred lines of house mice (*Mus musculus*). In addition to
their limited genetic diversity, housing mice in sterile lab environments may limit our power to
establish links between microbes and host immune phenotypes. Indeed, lab mice lack certain immune
cell populations that are found in pet store mice (8) and have much lower levels of immune activation
than wild animals due to limited pathogen exposure (9). Wild immune phenotypes can also be
recapitulated through "rewilding" experiments that acclimate lab mice to natural outdoor enclosures.
For example, wild-acclimated mice greatly increase granulocyte production and immune activation
relative to laboratory animals (10,11). Similarly, inoculating lab mice with wild gut microbiomes
lowered levels of inflammation and improved resistance to viral infection (12). Although most
immunology research has focused on rodent models, teleost fish represent an emerging model for
connecting environmental and microbial variation with immune development and function.

Fish models offer several useful avenues for studying immune variation. One clear benefit of
these systems is that their mucosal organs are in constant contact with the aquatic environment and
can quickly respond to environmental changes (13). Mucosal tissues such as respiratory surfaces
harbor important microbial communities and act in the first line of immune defense (14). Numerous
studies have examined how environmental factors such as salinity, pH, temperature, and pollutants
affect fish mucosal immunity (reviewed in Morshed and Lee, (15)). Another large body of work
addresses connections between pathogens and innate immunity, mostly focusing on organ specific
immune responses (reviewed in Dezfuli et al, (16)). Apart from tissue or response specific studies,
the ability to easily breed and rear fish across a spectrum of environments also offers research
opportunities that are often not available with rodent models. For example, housing similar strains in
lab, controlled outdoor (i.e., aquaculture), and wild settings, offers great opportunities for
disentangling genetic and environmental effects on immunity. Moreover, in instances where genetics
differences drive immune variation, several fish systems offer the opportunity to understand how and
why these phenotypic differences evolved (17,18)

Threespine stickleback fish (*Gasterosteus aculeatus*) are well known as models for studying
the process of adaptation via natural selection. This ancestrally marine species repeatedly colonized
and adapted to freshwater ecosystems across the northern hemisphere as glaciers receded 11,000
years ago (19,20,21). Recent work suggests that population-specific immune divergence is an
important factor in the adaptive radiation of stickleback into freshwater environments. Differences in
innate immune function have been documented between numerous ecotypes, including marine-
freshwater (22), sympatric benthic-limnetic (23), parapatric lake-stream ecotypes (24), and variation
over larger, continental scales (25,26). While some of these studies include more modern measures of
immune gene expression (18,23,27), many rely on two classic flow cytometric assays of immune
cells collected from the head kidney (i.e., the primary hematopoietic organ). In one case, forward and
side scatter are used to categorize populations and relative ratios of granulocytes and lymphocytes in
the head kidney, with greater proportions of granulocytes suggesting greater investment in innate
immunity (26). The second assay measures the oxidative burst response of granulocytes, where

fluorescent dyes allow the quantification of reactive oxygen species (ROS) production after cells are
exposed to an immune stimulant. (24,26). In general, populations with high natural macroparasite
exposure tend to evolve higher proportions of granulocytes and larger oxidative burst responses
(23,24). In addition to flow cytometric assays, the splenosomatic index (spleen mass/body mass, SSI)
is frequently used to evaluate immune activity in fish, with an increased SSI corresponding to higher
pathogen related immune response (28,29), potentially resulting from a greater need to filter
circulating leukocytes. Another simple measure, the hepatosomatic index (liver mass/body mass,
HSI) provides a proxy for individual condition because the liver is used to store fatty acids in fish
(30).

We took advantage of threespine stickleback populations with recently evolved heritable
differences in immune function to examine the effects of environment, most specifically microbial
community, on host immunity. By rearing fish in a highly controlled laboratory setting, we were able
to focus on two experimental variables: the effect of ancestry (i.e., host population) and the effect of
water flow into tanks (i.e., constant versus intermittent flow). Based on previous results (18,26), we
expected the cestode-resistant population to exhibit the highest immune response. We also
hypothesized that variation in water flow rate would affect the composition of microbes in our fish
tanks; specifically, we predicted that intermittent flow tanks would have more diverse microbial
communities due to higher environmental stability. Finally, we predicted that housing adult fish in
different water flow environments would reveal flexibility in adult innate immunity, likely because of
different microbial communities. Specifically, if the intermittent flow tanks drive higher microbial
diversity, fish housed in those tanks should also exhibit the largest immune responses. As most of our
fish measurements focused on cellular immunity, to understand more generalized effects of water
flow we also compared plasticity in two organs. We predicted that increased immune investment
would correlate with increases in spleen size, and that decreased liver size would indicate increased
energetic expenditure, potentially connected to immune activation.

Materials and Methods

Animal Care & Husbandry

Wild fish (F0-generation) were obtained from two sites on Vancouver Island, British Columbia:
Roberts Lake ("resistant" population, 50.222, -125.541); Gosling Lake ("tolerant", 50.061, -
125.504); and from the Kenai River Estuary in southeast Alaska ("marine", 60.542, -151.218).
Offspring were generated via in-vitro fertilization (F1-generation) in the field in summer 2019 and
fertilized eggs were shipped to the University of Wisconsin-Madison (ADFG license P-19-005;
Canada: NA19-457335). F1 fish were bred in the lab after reaching maturity in 2021, and those
offspring (F2) were used in this experiment. This breeding design should minimize maternal effects
on measured phenotypes, as all parents shared the same laboratory environment. Fish were housed in
18L tanks in full-sibling groups (n/tank = 2 – 15) with a water temperature of 17°C. The room was
kept on a 16:8 light cycle. Fish were fed daily with a mixture of freeze-dried bloodworms and mysis
shrimp, and each tank had a section of PVC pipe and two artificial plants for environmental
enrichment.

Acclimation Design

We used two water flow regimes to create different tank environments. Continuous-flow tanks had a
volume of 18L and input flow rate of 1L/minute. The intermittent flow tanks had a volume of 23L,
and water changes involved 1L/minute for 5 minutes every 12 hours. The entire volume of water was

replaced every 30 minutes in continuous flow tanks, as opposed to every 48 hours in intermittent
flow tanks. The intermittent flow tanks were also provided standing bubbling filters to maintain
water quality and promote microbial growth on the filter sponge. All tanks in the experiment shared a
common water system, and water was run through a UV filter before recirculation to limit microbe
movement between tanks. Due to a limited number of intermittent flow tanks, fry and young fish
were all initially reared in constant flow tanks. All fish used for this experiment had been in their
current tank for at least 11 months.

Fish ranged in age from 14 – 24 months at the start of the experiment and were randomly
assigned to one of four treatments: continuous flow only ('c'), intermittent flow only ('i'), move from
continuous to intermittent flow ('ci'), and move from intermittent to continuous flow ('ic'). After the
latter two treatments were initiated, fish were maintained in the new tank for approximately eighteen
weeks. After acclimation, we fasted fish for 24 hours and then euthanized via a combination of MS-
222 overdose and pithing. Prior to dissection fish were weighed and standard length recorded. The
liver and spleen were removed and weighed with an analytical balance (±0.0001), and the head
kidneys (HKs) were removed for flow cytometric assays.

Flow Cytometry Assay

Maximum ROS production in HK cells was quantified via oxidative burst assays. Organs were
placed in ice cold media ($0.9\times$ Roswell Park Memorial Institute medium containing 10% fetal bovine
serum, 100 μM nonessential amino acids, 100 U/mL penicillin, 100 μg/mL streptomycin, and 55 μM
β-mercaptoethanol) immediately after removal. Single cell suspensions were generated by manually
disrupting and filtering the tissue on a cell strainer (35 μm) with a pipette. We washed the cell
suspension once in 4 mL of cold media (300g, 4°C, 10 min), poured off most of the supernatant, and
then resuspended the cell pellet in the remaining media. Live HK cells were counted using a
hemocytometer (Hausser Scientific 3520) via Trypan blue exclusion (Corning 25-900-CI). Each
individual had 2×10^5 HK cells distributed to three separate wells on a 96-well plate. Each well
received either (1) 200 μL of HK medium or (2 & 3) HK medium with DHR-123 (ROS indicator
fluorochrome, 2 μg/mL, Sigma D1054). Samples were incubated at 18°C, 3% CO_2 for 10 minutes.
Next, PMA (Phorbol 12-myristate-13-acetat, 130 ng/mL; Sigma P8139) was added to well 3 to
stimulate ROS production while wells 1 and 2 received equivalent volumes of HK media. Cells were
incubated for a final 20 min at 18°C, 3% CO_2, placed in a sealed bag on ice for transport, and run on
a ThermoFisher Attune NxT V6 flow cytometer.

Data Analysis

Flow cytometry data was analyzed in FlowJo (TreeStar). The magnitude of ROS production during
oxidative burst assays was determined by comparing median fluorescence intensity of PMA-
stimulated and unstimulated cells. The proportion of granulocytes and lymphocytes was determined
by flow cytometry of the unstained, unstimulated HK cells and forward and side scatter gating
described in Weber et al. (18).

All other data analysis was done in R v4.2. Because the amount of ROS produced by head
kidney cells depended on time since exposure to PMA, we included flow cytometry sample order as
a covariate. Specifically, we made separate linear models for PMA-stimulated and unstimulated cells
with sample order as a random effect with R package *lme4* v35.5 (31) and used the model residuals
as the corrected median fluorescence intensity. We then built a series of linear models for
associations between four traits of interest (ROS production, percent granulocytes, liver and spleen

mass) and a suite of potential predictor variables: population, sex, family, tank environment
(continuous/intermittent), flow treatment (c/ci/ic/i), movement between tanks (ic and ci groups were
moved, c and i were not), body mass (only used for spleen and liver), and all relevant combinations
of predictor variables and interaction terms. We used the R package *performance* v12.4 (32) to
compare models and selected the model with the lowest AIC score for each response variable. The R
package *emmeans* v1.11 (33) was used to estimate group averages and test for significant differences
between multiple factors.

Tank 16S Sequencing & Analysis

We selected a subset of tanks for 16S sequencing to characterize microbial community differences
between flow regimes. Five continuous and five intermittent flow tanks were sampled using a
microbiome testing kit from Aquabiomics LLC (Junction City, OR). Two samples were taken from
each tank. First, 60 mL of tank water was slowly pushed through a 22-micron mixed cellulose ester
filter, followed by 2 mL of 90% ethanol; the filter was allowed to dry then sealed in a small plastic
container. Second, each tank's water intake tube was swabbed vigorously with a cotton swab, the
swab was dunked in a 1.5 mL cryotube filled with 90% ethanol, allowed to dry, then sealed in a
plastic tube.

Sequence data were curated in R v4.2 with the *DADA2* pipeline (v1.16, 34). Reads were
filtered and trimmed (command: 'filterAndTrim') with parameters maxN=0, truncQ=2,
rm.phix=TRUE and maxEE=c(2,5). Paired reads were then merged (command: 'mergePairs') and
chimera sequences were removed (command: 'removeBimeraDenovo'). R package *DECIPHER*
(v.2.26, 35) was used to construct an amplicon sequence variants (ASVs) table (command: 'IdTaxa')
after training on a reference dataset (SILVA 138 SSU; 36). Analyses of species richness and
differences between communities were completed with the R package *phyloseq* (v1.42, 37).

Results

Tank Microbe Communities

Despite collecting and submitting 20 microbial samples for 16S library preparation and sequencing,
we only recovered usable data from four tanks: three continuous flow and one intermittent. We were
unable to acquire additional tank microbial samples because the experiment had concluded by the
time that we learned of the sequencing error (i.e., we no longer had access to the relevant fish
populations in each tank environment). Although the small number of tanks limits our confidence in
measurements of microbial differences between water flow environments, we nonetheless identified
several patterns. 2D ordination (Figure S1) separated the filter and swab samples along the first
NMDS axis, and the intermittent flow tank separated from the three continuous flow tanks along the
second axis. The intermittent flow tank also had the highest Shannon diversity index score (Figure
S2), though it did not diverge greatly from the continuous flow tanks. We generated a list of the top
ten ASVs that had the largest difference in read counts between the tank environments. ASVs in the
genera *Sediminibacterium*, *Polynucleobacter*, and *Undibacterium* were overrepresented in both filter
and swab samples in the intermittent tank (Table S1), whereas an ASV in the genus *Sphingopyxis*
was underrepresented in the intermittent samples compared to continuous flow. Together, this
suggests that the water flow regime likely altered the microbial communities where fish were housed.

Reactive Oxygen Species (ROS) Generation

The best fit regression model for ROS production included fixed effects for population and flow
treatment, and family as a random effect (Table 2). The resistant stickleback population produced
significantly higher ROS than tolerant (mean of 0.55 median fluorescence intensity vs -0.48 mfi,
Cohen's d =-3.56, t-test p<0.00001, Table 1, Figure 2) or marine populations across all treatments (d
=-1.72, p<0.00001). The tolerant population also produced less ROS than marine fish (d=-1.84,
p<0.00001). These data support a strong role of population ancestry and evolutionary divergence in
shaping oxidative burst capacity. Across all populations the c treatment had the lowest ROS
production, which was significantly different from the i (d=-0.27, p = 0.004) and ci treatments
(d=0.505, p = 0.0001) and marginally different from the ic treatment (d=0.37, p = 0.051). Fish in
intermittent flow tanks tended to have greater ROS production than those in continuous flow (Figure
S6, d=-0.11, p = 0.002). There was no effect of sex on ROS generation (p = 0.74).

Granulocyte:Lymphocyte Ratio

The best supported model for percent granulocytes included fixed effects for population, sex, and
flow regime, and family as a random effect. Intermittent flow tanks averaged significantly higher
proportions of granulocytes in the head kidney than continuous flow tanks (38.1% vs 34%; t-test p =
0.003, Cohen's d =-0.03, Table 1, Figure 3). The resistant population also had higher proportions of
granulocytes than the tolerant (41.2% vs 34.4%, p = 0.02, d=-0.83) or marine (32.6%, p = 0.001, d=-
1.05) populations. The latter two populations were not significantly different (p = 0.47).

Liver and Spleen Mass

Variation in spleen and liver mass was driven by fish mass, population, sex, and tank environment.
Fish from the tolerant population had significantly smaller livers than resistant fish after controlling
for mass (-0.44 vs 0.27 mass-scaled liver size, Cohen's d = -2.1, t-test p<0.00001, Table 1, Figure 4).
The marine population was intermediate in liver size and not significantly different from either
tolerant or resistant populations. Fish in the c treatment had significantly larger livers than the ci
(0.16 vs -0.12, d=0.81, p=0.03) and ic treatments (0.16 vs -0.13, d=0.83, p=0.02) and were
marginally larger than the ii treatment (0.16 vs -0.08, d=0.67, p=0.06). Males averaged smaller livers
than females, but this difference was not significant in multiple comparisons (-0.04 vs 0.04, d = 0.23,
p=0.16). The best-supported model for liver mass included population, sex, and flow treatment as
fixed effects and family as a random effect.

The marine population averaged larger spleens than both resistant (0.16 vs -0.08 mass
corrected spleen size, d = 0.49, p = 0.01, Figure 5) and tolerant populations (0.16 vs -0.08, d = 0.55, p
= 0.006). Male fish had larger spleens than females (0.11 vs -0.11, d=0.45, p=0.01), and fish in
intermittent flow had marginally larger spleens than continuous flow (0.06 vs -0.07, d = -0.28,
p=0.05). The best-supported model for spleen mass included population, sex, and flow regime. Mass-
corrected spleen size was not significantly correlated with either ROS production or
granulocyte:lymphocyte ratio (Pearson's correlation coefficient, p>0.1). Mass-corrected liver size
was positively correlated with ROS production (r=0.31, p=0.005), but not granulocyte:lymphocyte
ratio (p=0.18).

Discussion

We leveraged evolved differences in immunity across stickleback populations to investigate how
variation in the laboratory environment affects immune function. We found that simply transferring
fish between tanks with constant or intermittent water flow, a relatively modest environmental
change, led to significant changes in the inflammatory responses and composition of head kidney

cells. Intermittent water flow not only drove significantly higher levels of ROS production but also
greater proportion of granulocytes–a category that includes neutrophils likely to drive strong
oxidative burst responses. After eighteen weeks of exposure, the magnitude of both immune
measures approximated levels observed in fish that had been housed in each environment for a long
period. This demonstrates that environmentally induced flexibility in stickleback innate immune
responses is a relatively rapid process.

Apart from classic measures of innate immunity, we also tested the impacts of water flow
regime on liver and spleen size. Fish that ended the experiment housed in intermittent flow tanks had
significantly higher SSI, consistent with a plastic shift to greater innate immune allocation. In
contrast, there was no simple relationship between HSI and water flow treatment. Stable (non-
transplant) rearing under constant water flow yielded significantly higher HSI than the other
treatment groups. We had too few tanks available to perform sham transfers (i.e., c-c or i-i), so we
cannot rule out a handling effect in the ci or ic treatments. However, this seems unlikely given that
fish were allowed to acclimate to their new tanks for approximately eighteen weeks. The tolerant
population had the lowest HSI overall. This was likely due to this population's fast growth rate,
which is also associated with liver size in fish (38). Marine and resistant fish did not differ in relative
liver size even though marine fish are significantly larger.

All populations showed a similar magnitude of flexibility for ROS production, granulocyte
proportion, and SSI, and the heritable differences between populations were by far the strongest
predictor of immune response (Table 2). Population accounted for 89% of the variance in ROS
production, 42% of the variance in granulocyte proportion, and 43% of the variance in SSI. Several
recent studies have highlighted the utility of population level comparisons of immune variation (e.g.,
17, Robertson et al 2016, Weber et al 2022), but this remains a relatively neglected component of
most experimental designs (Sasser et al 2023).
Variation in ROS production in these populations evolved recently in response to parasitism
by the tapeworm *Schistocephalus solidus* which is not present in marine environments (18). We
found a similar magnitude of difference between resistant and tolerant populations compared to
previous work, although overall our measurements were lower than those reported in Weber et al
(18). Our measures of granulocyte proportion were comparable to our previous results (18) but
significantly higher than those reported in wild stickleback from Germany (24). This study was the
first to include measures of ROS production in ancestral marine stickleback and demonstrated that
the marine population is intermediate between the cestode resistant and tolerant freshwater
populations.

Because our experiment was performed in a laboratory with highly controlled environments,
we hypothesized that reducing the frequency of tank water changes would primarily lead to increased
microbial diversity. Our results, with the caveat that they represent only a few tanks, support this
prediction. The intermittent flow tank had the highest level of microbial diversity in swabs taken
from the tank outflow pipes, and the second highest diversity from filtered water samples (Figure
S2). Perhaps more striking was the divergence in microbe communities when accounting for ASV
abundance. Whether considering microbes filtered directly from the water or from a swab of the tank
outflow, 16S divergence along the first NMDS axis was strongly captured by flow regime (Fig S1).
While the present study was a preliminary examination of whether variation in water flow could
separately influence tank microbial environment and fish immunity, our positive results beg several
questions: Do changes in tank microbial environment alter the fish microbiome and, if so, are the
effects general or specific to areas such as the skin or gut? And are shifts in host immunity caused by
microbial exposure, alterations in microbiome, or both events? More directed experiments are needed
to address these questions, but our results do at least support a microbe-immune connection. The

higher ROS and granulocyte levels of head kidney cells from stickleback in intermittent flow tanks suggests an increased systemic immune stress (40). This is consistent with pathogen response, potentially mediating via migration of antigen presenting cells from the periphery to head kidney (41).

Ethics Statement

This experiment was reviewed and approved by the University of Wisconsin-Madison IACUC under protocol L006460-A01.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

1. Author Contributions

Conceived and designed the experiments: JW and CW. Performed the experiments, analyzed the data, and prepared the figures: CW. Wrote the first draft: CW and PC. Provided funding: JW. All authors contributed to the article and approved the submitted version.

2. Funding

This work was funded by a startup award from the University of Wisconsin-Madison Department of Integrative Biology (JNW) and the National Institute of General Medical Sciences (5R35GM142891; JNW)

3. Acknowledgments

We thank Dr J Benavente Paredes and M Chotlos for assistance with data collection; J Kosmopoulos and Dr K Anantharaman for advice on 16S data analysis; and E Kerns, K Yu, J Abels, and L Rippin for helpful feedback on this manuscript.

4. Data Availability Statement

The datasets generated for this study can be found on Dryad (https://doi.org/10.5061/dryad.0vt4b8hb8) and GenBank (link).

References

1.   Schoenle, Laura A., Cynthia J. Downs, and Lynn B. Martin. "An introduction to ecoimmunology." Advances in comparative immunology (2018): 901-932.

2.   Stinson LF. Establishment of the early-life microbiome: a DOHaD perspective. J Dev Orig Health Dis. 2020 Jun;11(3):201–10.

3.   MacGillivray DM, Kollmann TR. The Role of Environmental Factors in Modulating Immune Responses in Early Life. Front Immunol [Internet]. 2014 Sep 12

4.   Campolim CM, Schimenes BC, Veras MM, Kim YB, Prada PO. Air pollution accelerates the
development of obesity and Alzheimer's disease: the role of leptin and inflammation - a mini-review.
Front Immunol. 2024 Jun 12;15:1401800.

5.   Kreitinger JM, Beamer CA, Shepherd DM. Environmental Immunology: Lessons Learned
from Exposure to a Select Panel of Immunotoxicants. The Journal of Immunology. 2016 Apr
15;196(8):3217–25.

6.   Bolnick DI, Fuess LE, Graham AL, Khan I, Steinel NC, Vaziri GJ. Evolutionary
Immunology. Annual Review of Ecology, Evolution, and Systematics [Internet]. 2025 May 6

7.   Zheng D, Liwinski T, Elinav E. Interaction between microbiota and immunity in health and
disease. Cell Res. 2020 Jun;30(6):492–506.

8.   Beura LK, Hamilton SE, Bi K, Schenkel JM, Odumade OA, Casey KA, et al. Normalizing the
environment recapitulates adult human immune traits in laboratory mice. Nature. 2016
Apr;532(7600):512–6.

9.   Abolins S, King EC, Lazarou L, Weldon L, Hughes L, Drescher P, et al. The comparative
immunology of wild and laboratory mice, Mus musculus domesticus. Nat Commun. 2017 May
3;8(1):14811.

10.   Lin JD, Devlin JC, Yeung F, McCauley C, Leung JM, Chen YH, et al. Rewilding Nod2 and
Atg16l1 Mutant Mice Uncovers Genetic and Environmental Contributions to Microbial Responses
and Immune Cell Composition. Cell Host & Microbe. 2020 May;27(5):830-840.e4.

11.   Yeung F, Chen YH, Lin JD, Leung JM, McCauley C, Devlin JC, et al. Altered Immunity of
Laboratory Mice in the Natural Environment Is Associated with Fungal Colonization. Cell Host &
Microbe. 2020 May;27(5):809-822.e6.

12.   Rosshart SP, Vassallo BG, Angeletti D, Hutchinson DS, Morgan AP, Takeda K, et al. Wild
Mouse Gut Microbiota Promotes Host Fitness and Improves Disease Resistance. Cell. 2017
Nov;171(5):1015-1028.e13.

13.   Cabillon NAR, Lazado CC. Mucosal Barrier Functions of Fish under Changing
Environmental Conditions. Fishes. 2019 Jan 10;4(1):2.

14.   Nochi T, Jansen CA, Toyomizu M, Eden WV. The Well-Developed Mucosal Immune
Systems of Birds and Mammals Allow for Similar Approaches of Mucosal Vaccination in Both
Types of Animals. Front Nutr. 2018 Jul 12;5:60.

15.   Morshed SM, Lee TH. The role of the microbiome on fish mucosal immunity under changing
environments. Fish & Shellfish Immunology. 2023 Aug;139:108877.

16.   Sayyaf Dezfuli B, Lorenzoni M, Carosi A, Giari L, Bosi G. Teleost innate immunity, an
intricate game between immune cells and parasites of fish organs: who wins, who loses. Front
Immunol. 2023 Oct 16;14:1250835.

17.   Peuß R, Box AC, Chen S, Wang Y, Tsuchiya D, Persons JL, et al. Adaptation to low parasite
abundance affects immune investment and immunopathological responses of cavefish. Nat Ecol
Evol. 2020 Jul 20;4(10):1416–30.

18.   Weber JN, Steinel NC, Peng F, Shim KC, Lohman BK, Fuess LE, et al. Evolutionary gain
and loss of a pathological immune response to parasitism. Science. 2022 Sep 9;377(6611):1206–11.

19.   Bell, M. A., and S. A. Foster, eds. 1994. The evolutionary biology of the threespine
stickleback. Oxford Univ. Press, Oxford, U.K

20.   McKinnon JS, Rundle HD. Speciation in nature: the threespine stickleback model systems.
Trends in Ecology & Evolution. 2002 Oct;17(10):480–8.

21.   Reid K, Bell MA, Veeramah KR. Threespine Stickleback: A Model System For Evolutionary
Genomics. Annu Rev Genom Hum Genet. 2021 Aug 31;22(1):357–83.

22.   Whiting JR, Magalhaes IS, Singkam AR, Robertson S, D'Agostino D, Bradley JE, et al. A
genetics□based approach confirms immune associations with life history across multiple populations
of an aquatic vertebrate (Gasterosteus aculeatus). Molecular Ecology. 2018 Aug;27(15):3174–91.

23.   Stutz WE, Lau OL, Bolnick DI. Contrasting Patterns of Phenotype-Dependent Parasitism
within and among Populations of Threespine Stickleback. The American Naturalist. 2014
Jun;183(6):810–25.

24.   Scharsack JP, Kalbe M, Harrod C, Rauch G. Habitat-specific adaptation of immune responses
of stickleback (Gasterosteus aculeatus) lake and river ecotypes. Proc R Soc B. 2007 Jun
22;274(1617):1523–32.

25.   Hamley M, Franke F, Kurtz J, Scharsack JP. An experimental approach to the immuno-
modulatory basis of host-parasite local adaptation in tapeworm-infected sticklebacks. Experimental
Parasitology. 2017 Sep;180:119–32.

26.   Weber JN, Steinel NC, Shim KC, Bolnick DI. Recent evolution of extreme cestode growth
suppression by a vertebrate host. Proc Natl Acad Sci USA. 2017 Jun 20;114(25):6575–80.

27.   Robertson S, Bradley JE, MacColl ADC. Measuring the immune system of the three□spined
stickleback – investigating natural variation by quantifying immune expression in the laboratory and
the wild. Molecular Ecology Resources. 2016 May;16(3):701–13.

28.   Kaufmann J, Lenz TL, Kalbe M, Milinski M, Eizaguirre C. A field reciprocal transplant
experiment reveals asymmetric costs of migration between lake and river ecotypes of three□spined
sticklebacks (Gasterosteus aculeatus). J of Evolutionary Biology. 2017 May;30(5):938–50.

29.   Rajkov J, El Taher A, Böhne A, Salzburger W, Egger B. Gene expression remodelling and
immune response during adaptive divergence in an African cichlid fish. Molecular Ecology. 2021
Jan;30(1):274–96.

30.   Chellappa S, Huntingford FA, Strang RHC, Thomson RY. Condition factor and
hepatosomatic index as estimates of energy status in male three□spined stickleback. Journal of Fish
Biology. 1995 Nov;47(5):775–87.

31.   Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. J
Stat Soft. 2015 67(1). Available from: http://www.jstatsoft.org/v67/i01/

32.   Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D. performance: An R Package
for Assessment, Comparison and Testing of Statistical Models. JOSS. 2021 Apr 21;6(60):3139.

33.   Lenth RV. emmeans: Estimated Marginal Means, aka Least-Squares Means. 2025.

34.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods. 2016 Jul;13(7):581–3.

35.   Murali A, Bhargava A, Wright ES. IDTAXA: a novel approach for accurate taxonomic
classification of microbiome sequences. Microbiome. 2018 Dec;6(1):140.

36.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal
RNA gene database project: improved data processing and web-based tools. Nucleic Acids Research.
2012 Nov 27;41(D1):D590–6.

37.   McMurdie PJ, Holmes S. phyloseq: An R Package for Reproducible Interactive Analysis and
Graphics of Microbiome Census Data. Watson M, editor. PLoS ONE. 2013 Apr 22;8(4):e61217.

38.   Pelletier D, Dutil JD, Blier P, Guderley H. Relation between growth rate and metabolic
organization of white muscle, liver and digestive tract in cod, Gadus morhua. J Comp Physiol B.
1994 Jul;164(3):179–90.

39.   Sasser KT, Weber JN. A call for more ecologically and evolutionarily relevant studies of
immune costs. Evol Ecol. 2023 Feb;37(1):203–14.

40.   Speirs ZC, Loynes CA, Mathiessen H, Elks PM, Renshaw SA, Jørgensen LVG. What can we
learn about fish neutrophil and macrophage response to immune challenge from studies in zebrafish.
Fish & Shellfish Immunology. 2024 May;148:109490.

41.   Iliev DB, Thim H, Lagos L, Olsen R, Jørgensen JB. Homing of Antigen-Presenting Cells in
Head Kidney and Spleen – Salmon Head Kidney Hosts Diverse APC Types. Front Immunol. 2013
June;137:1-13

Population	Treatment	ROS (mfi)	% granulocytes	Rel. liver mass	Rel. spleen mass
Marine	c	11.58±0.28 (11)	34.13±3.36 (15)	0.22±0.13 (22)	-0.06±0.12 (22)
	ci	11.88±0.17 (11)	31.42±6.08 (10)	0.06±0.12 (12)	0.34±0.37 (12)
	ic	11.96±0.22 (13)	29.93±3.97 (13)	-0.08±0.08 (17)	0.07±0.12 (17)
	i	11.68±0.20 (14)	35.02±2.76 (16)	0.01±0.07 (20)	0.19±0.17 (20)
Cestode tolerant	c	10.84±0.20 (21)	32.25±4.98 (17)	-0.27±0.11 (25)	-0.02±0.26 (25)
	ci	11.56±1.08 (4)	42.23±21.74 (3)	-0.42±0.24 (4)	-0.49±0.37 (4)
	ic	11.13±0.30 (8)	31.06±9.50 (5)	-0.28±0.11 (8)	-0.28±0.23 (8)
	i	11.57±2.02 (3)	41.38±14.88 (4)	-0.33±0.26 (6)	-0.16±0.19 (6)
Cestode resistant	c	12.51±0.41 (8)	37.09±8.25 (11)	0.34±0.47 (13)	-0.10±0.20 (13)
	ci	12.64±0.35 (8)	50.70±10.06 (8)	0.66±0.51 (7)	0.21±0.50 (7)
	ic	12.20±0.33 (3)	33.33±4.84 (3)	0.05±0.55 (3)	-0.05±0.26 (3)
	i	12.35±0.50 (9)	39.08±6.28 (10)	-0.07±0.07 (16)	-0.13±0.10 (16)

434 Table 1: trait means with 95% confidence interval by population and flow treatment (c = constant
435 flow, i = intermittent flow, ci = moved from constant to intermittent, ic = moved from intermittent to
436 constant). Sample size for each group in parenthesis. Relative masses are residuals from regression of
437 organ mass on body mass.

Trait	Predictor	Beta	SE	p	R^2
ROS	intercept	-0.06 (-0.38,0.24)	0.14	0.63	(0.61,0.70)
	pop-tolerant	-0.53 (-0.88,-0.18)	0.15	0.01	
	pop-resistant	0.50 (0.14,0.85)	0.15	0.01	
	treatment-ci	0.22 (0.05,0.39)	0.09	0.01	
	treatment-ic	0.18 (-0.01,0.38)	0.10	0.06	
	treatment-i	0.07 (-0.09,0.25)	0.09	0.38	
% Gran	intercept	-0.19 (-0.69,0.29)	0.19	0.35	(0.18,0.39)
	pop-tolerant	0.09 (-0.47,0.66)	0.23	0.70	
	pop-resistant	0.44 (-0.12,1.01)	0.22	0.10	
	sex-m	-0.1 (-0.27,0.05)	0.08	0.19	
	flow-int	0.21 (0.03,0.38)	0.09	0.02	
Liver mass	intercept	0.33 (-0.06,0.73)	0.16	0.09	(0.53,0.70)
	fish mass	0.95 (0.77,1.12)	0.09	0.00	
	pop-tolerant	-0.7 (-1.17,-0.23)	0.20	0.01	
	pop-resistant	0.1 (-0.36,0.56)	0.19	0.61	
	sex-m	-0.11 (-0.21,-0.02)	0.05	0.02	
	treatment-ci	-0.18 (-0.34,-0.02)	0.08	0.02	
	treatment-ic	-0.21 (-0.36,-0.06)	0.08	0.01	
	treatment-i	-0.15 (-0.29,-0.01)	0.07	0.03	
Spleen mass	intercept	0.07 (-0.06,0.20)	0.07	0.29	(0.40,0.40)
	fish mass	0.8 (0.62,0.98)	0.09	0.00	
	pop-tolerant	-0.43 (-0.64,-0.22)	0.11	0.00	
	pop-resistant	-0.24 (-0.4,-0.08)	0.08	0.00	
	sex-m	0.13 (0.00,0.26)	0.07	0.04	
	flow-int	0.13 (0.00,0.26)	0.07	0.05	

440 Table 2: coefficients from top-ranked model of each trait of interest. 95% confidence interval in
441 parenthesis. 2