1

Characterisation of β-tubulin isotypes in *Uncinaria stenocephala* and implications for

benzimidazole resistance in hookworms

Thomas Stocker[1] and Jan Šlapeta[1,2,*]

[1] Sydney School of Veterinary Science, Faculty of Science, The University of Sydney, New South

Wales 2006, Australia

[2] Sydney Infectious Diseases Institute, New South Wales 2006, Australia

*Author for correspondence: jan.slapeta@sydney.edu.au

Abstract:

Uncinaria stenocephala is a widespread hookworm of dogs across Europe, Canada, southern

Australia, and other temperate regions, where it often outnumbers infections caused by

Ancylostoma caninum. Although a putative β-tubulin isotype-1 mutation associated with

resistance has been detected in *U. stenocephala*, clinical resistance to benzimidazoles has not

yet been confirmed. Benzimidazole resistance is linked to single nucleotide polymorphisms

(SNPs) in the β-tubulin isotype-1 gene; however, the β-tubulin genes of *U. stenocephala* have

not been fully characterised. We aimed to identify β-tubulin genes and confirm the coding

sequences for key residues (Q134, F167, E198, and F200) in the β-tubulin isotype-1 gene of the

U. stenocephala genome. Two *U. stenocephala* specimens were subjected to Illumina

sequencing, and species identity was confirmed through morphological and molecular analysis

using ITS rDNA and cox-1 markers. Genome assembly revealed the presence of β-tubulin

isotype-1 (10 exons) and isotype-2 (9 exons), both homologous to β-tubulins from other

hookworms (*A. caninum*, *A. ceylanicum*, *A. duodenale* and *Necator americanus*). The β-tubulin

isotype-1 protein sequence of *U. stenocephala* contained two variable residues (S37Q and

G441A) compared to other hookworm sequences. While the isotype-2 protein sequence was

conserved among *Ancylostoma* species, *U. stenocephala* exhibited six distinct polymorphisms

(E39D, T40S, N115S, L130I, A287S, T439G). The benzimidazole-susceptible residues (Q134,

F167, E198, F200) were present in the β-tubulin isotype-1 protein sequence. Characterisation of

the complete coding regions of β-tubulin isotypes 1 and 2 enables population-level screening

for benzimidazole resistance-associated SNPs and provides a foundation for future

epidemiological studies in *U. stenocephala*.

Keywords:

tubb-1, *tubb-2*, whole genome sequence, WGS, mutation, exon, Ancylostomatidate

Highlights

41 *Uncinaria stenocephala* outnumbers *Ancylostoma caninum* in temperate regions globally

42 β-tubulin isotypes-1 and -2 were fully characterised in *Uncinaria stenocephala*

43 Key benzimidazole-susceptible residues were confirmed in β-tubulin isotype-1

44 β-tubulin isotypes-1 and -2 enables SNP screening and resistance surveillance efforts

1. Introduction

Hookworm nematodes (Ancylostomatidae) cause anaemia in humans, cats, and dogs through gastrointestinal haemorrhages. Until recently, however, anthelmintic resistance had not been a major concern (Geary et al., 2025). Benzimidazoles have been used effectively for over 50 years in companion animals to treat hookworm infections. Their mode of action involves disrupting the assembly of the nematode cytoskeleton and so impairing development. Clinical resistance to benzimidazoles was first documented in the canine hookworm (*Ancylostoma caninum*) using both *in vivo* (faecal egg count reduction test, FECRT) and *in vitro* (egg hatch assay, EHA) methods (Jimenez Castro et al., 2019; Kitchen et al., 2019). While FECRT and EHA are invaluable, they are inherently low-throughput. High-throughput amplicon sequencing has emerged as a powerful alternative for screening known single nucleotide polymorphisms (SNPs) in tubulin genes, which are critical components of the nematode cytoskeleton. Amplicon metabarcoding and sequencing of the β-tubulin isotype-1 (*tubb-1*) gene have revealed widespread mutations at amino acid positions 134 and 167, which confer benzimidazole resistance in *A. caninum* populations from dogs in North America and Australia (Abdullah et al., 2025; Venkatesan et al., 2023).

While mutations in the *tubb-1* gene have been strongly associated with benzimidazole resistance in hookworms, the broader genetic context of β-tubulin genes in nematodes warrants further investigation. Strongyle nematodes, including hookworms, possess four genes encoding β-tubulins (*tubb-1*, *tubb-2*, *tubb-3* and *tubb-4*), in contrast to the model organism *Caenorhabditis elegans*, which also contains the *ben-1* gene and is amenable to genetic studies (Collins et al., 2024; Saunders et al., 2013). Although non-synonymous variants at codons 134, 167, 198 and 200 of the *tubb-1* gene have been associated with benzimidazole resistance, *tubb-1* may not be the sole gene involved (Collins et al., 2024). The causal role of these codon changes was demonstrated through mutations in the *ben-1* gene of *C. elegans*, which confer resistance to benzimidazoles (Dilks et al., 2020; Dilks et al., 2021; Kitchen et al., 2019;

Venkatesan et al., 2023). In *C. elegans*, *ben-1* confers the highest level of resistance, and the

loss of a second β-tubulin gene does not alter this resistance level (Collins et al., 2024). In

nematodes lacking *ben-1*, resistance is conferred by other β-tubulin genes, with *tubb-1* playing

a major, though not exclusive, role. Therefore, it is important to define the full complement of β-

tubulin genes, particularly *tubb-1* and *tubb-2*, which are ubiquitously expressed in nematode

cells, and to screen for mutations that may either confer or contribute to benzimidazole

resistance (Collins et al., 2024; Gandasegui et al., 2024; Saunders et al., 2013; Venkatesan et

al., 2023).

Hookworm nematodes are among the common intestinal parasites of companion

animals, particularly dogs and cats, alongside roundworms and whipworms (Lightner et al.,

1978). In these hosts, the primary pathogenic hookworms are *A. caninum* in dogs and

Ancylostoma tubaeforme in cats (Miller, 1968). However, in cooler temperate regions, *Uncinaria*

stenocephala, traditionally considered less pathogenic, predominates. Unlike the blood-

feeding *Ancylostoma* species that cause anaemia, *U. stenocephala* infections are associated

with protein-losing enteropathy (Walker and Jacobs, 1985). Recognised as the 'northern

hookworm', *U. stenocephala* is widely distributed across Europe, Canada, southern Australia,

and New Zealand, where it often outnumbers *Ancylostoma* spp. infections (Beveridge, 2002;

Gibbs, 1961; Xhaxhiu et al., 2011).

Given the recent evidence of benzimidazole resistant-associated mutation in *U.*

stenocephala, a deeper molecular characterisation of its β-tubulin genes is needed. Building on

molecular tools developed for *A. caninum*, Stocker et al. (2023) demonstrated that a *tubb-1*-

targeting assay could also amplify *U. stenocephala tubb-1*, making it suitable for screening

potential genetic markers of benzimidazole resistance in this species. A study from Australia

reported the presence of the canonical F167Y benzimidazole resistance mutation in *U.*

stenocephala tubb-1 (Abdullah et al., 2025). Although clinical treatment failure with

benzimidazoles in *U. stenocephala* has not yet been documented, proactive molecular

surveillance is essential. To anticipate emerging resistance, comprehensive characterisation of

β-tubulin genes, particularly full-length *tubb-1*, across hookworm species is critical for enabling

comparative analyses and early detection of resistance-associated mutations.

The aim of this study was to characterise the *tubb-1* and *tubb-2* genes of *U.*

stenocephala and to identify key amino acid residues potentially associated with benzimidazole

resistance. To achieve this, we isolated adult *U. stenocephala* specimens and performed

whole-genome sequencing. Using a comparative approach with available hookworm genomic

data from *A. caninum*, *Ancylostoma ceylanicum*, *Ancylostoma duodenale*, and *Necator*

americanus, we identified and characterised the coding sequences of *tubb-1* and *tubb-2*. To

facilitate high-throughput screening of benzimidazole-susceptible residues in TUBB-1 (Q134,

F167, E198, and F200), we reviewed existing primer sequences and in silico assessed their

suitability for *U. stenocephala* and other hookworm species.

2. Materials and Methods

2.1. Adult *Uncinaria stenocephala* material

To obtain *Uncinaria stenocephala*, we decided to consider the invasive red fox (*Vulpes*

vulpes) that a pest animal in Australia and is regularly culled (Stobo-Wilson et al., 2021). Red

foxes are commonly infected by *U. stenocephala* in Australia (Ryan, 1976). Single red fox

intestinal tract was donated by a registered shooter in September 2024 (Mardi; 33.2990° S,

151.4084° E); the fox was culled under an ongoing fox control program in Central Coast region of

New South Wales, Australia. The gastrointestinal tract was transported on ice to the University

of Sydney, stored for 10 days at -80°C prior to dissection. During dissection the gastrointestinal

track was thawed, incised longitudinally and content scraped through the researcher's thumb

and index finger. The scrapings and small intestinal contents washed with lukewarm tap water

into a 150 μm nylon mesh sieve by means of water jet. The washed content was examined in a

white metal tray with the aid of stereomicroscope. Nematodes were manually picked against

the white tray, washed in phosphate buffered saline (pH=8.0) and stored at -80°C. Hookworms,

such as *U. stenocephala*, were identified in accordance with the description of Ransom (1924).

2.2. Genomic DNA extraction and molecular identification

Genomic DNA (gDNA) was isolated using the Monarch® Spin gDNA Extraction Kit (New

England Biolabs, Australia). The gDNA was stored at -20°C. The isolated DNA was subject to two

separate PCR assays. The PCR amplifying ITS rDNA used NC16 [S0863] (5'- AGT TCA ATC GCA

ATG GCT T -3') and NC2 [S0864] (5'- TTA GTT TCT TTT CCT CCG CT -3') (Chilton et al., 2003) The

partial *cox*1 was amplified with JB3 [S0361] (5'- TTT TTT GGG CAT CCT GAG GTT TA -3') and JB4.5

[S0362] (5'- TAA AGA AAG AAC ATA ATG AAA ATG -3') (Bowles et al., 1992). Amplicons of

expected size (~800 bp, ITS rDNA; ~450 bp, *cox*1) were submitted for Sanger sequencing

(Macrogen, Korea). Sequences were assembled and compared to available sequences in CLC

Main Workbench 25.0.1 (CLC bio, Qiagen, Australia). As a reference of *U. stenocephala* ITS

rDNA sequence we used (Nadler et al., 2000) and *cox*1 we used AJ407939 (Hu et al., 2002).

2.3. Whole genome sequencing and assembly

The gDNA obtained from two single *U. stenocephala* was aliquoted (30 µL) into a 1.5 mL

Eppendorf tube and submitted to Novogene (HK) Co., Ltd for indexing, library preparation and

whole genome sequencing to generate approximately 30 Gb of sequencing data per sample.

Sequencing was performed using the Illumina NovaSeq 6000 sequencing platform using 150 bp

paired-end sequencing chemistry. Raw FastQ files (JS6871 and JS6872) generated from the

same sample were processed in Galaxy Australia (https://usegalaxy.org.au/) (Galaxy

Community, 2024)**.** Raw paired-end sequencing reads were examined for quality using *FastQC*

(version 0.12.1) (Andrews, 2010) and trimmed using *Trimmomatic* (version 0.36.6) (Bolger et al.,

2014) (parameters: "ILLUMINACLIP:TruSeq3-PE.fa:2:30:10 SLIDINGWINDOW:4:20"). Read were

mapped using *bwa-mem2* (version 2.2.1) (Li and Durbin, 2009) using simple Illumina mode onto

red fox genome (VulVul3) and all mapper reads were removed. The FastQ files were assembled

using *SPAdes* (version 4.2.0) (Prjibelski et al., 2020) (parameters: "--cov-cutoff 40.0 -k 77").

Contigs were classified using *Kraken* (version 1.1.1) (database: minikraken) (Wood and

Salzberg, 2014) and only unclassified contigs were retained. Assembly statistics were

generated using *Fasta Statistics* (Seemann and Gladman, 2012).

2.4. Identification of β-tubulin genes in the genome assemblies

The *tubb-1* and *tubb-2* genes and their corresponding proteins were retrieved from

hookworm genomes at 'WormBase ParaSite' (Howe et al., 2016; Howe et al., 2017). The genes

were identified in *A. caninum, A. ceylanicum, A. duodenale* and *Ne. americanus* with and

addition of an outgroup (*Ni. braziliensis*) (International Helminth Genomes, 2019; Schwarz et

al., 2015). The reference sequences for *tubb-1* (ANCCAN_08153, Acey_s0035.v2.g13228,

ANCDUO_05511, Necator_chrII.pre1.g8352 and NBR_0000799801) and *tubb-2*

(ANCCAN_23405, Acey_s0432.v2.g1106, Necator_chrII.pre1.g8352 and Nippo_chrII.g5566)

were aligned and genes structure identified. There was no *tubb-2* for *A. duodenale* localised due

to partial coverage of the genome.

Local BLAST databases from the assembled contigs of *U. stenocephala* were created and

queried with the reference *tubb-1* and *tubb-2* genes sequences. Candidate contigs were aligned

to the reference alignments and exon-intron boundaries identified. Alignments and local BLAST

searches were performed in CLC Main Workbench v 25.0.1 (CLC bio, Qiagen, Australia).

Sequence reads were mapped to the putative *tubb-1* and *tubb-2* using *bwa-*

mem2 (version 2.2.1) (Li and Durbin, 2009) using simple Illumina mode. *Samtools mpileup*

(Version 1.15.1) (Danecek et al., 2021) was used to input variants into *VarScan* (Version 2.4.2)

(Koboldt et al., 2012) (parameters: --min-coverage 20 --min-reads2 2 --min-avg-qual 15 --min-

var-freq 0.2 --min-freq-for-hom 0.75 --p-value 0.05). The resulting BAM and VCF files were

visualised using *IGV* (version 2.18.1). VCF file was processed to select only SNP within the

exons of *tubb-1* and *tubb-2*. The variant detection was performed in Galaxy Australia

(https://usegalaxy.org.au/) (Galaxy Community, 2024).

To evaluate the binding of the primers targeting *A. caninum tubb-1* (ACB1_167_F: 5'-GGY

GCA GGA AAC AAC TG -3', ACB1_167_R: 5'- CTT TGG TGA GGG GAC AAC A -3', ACB1_200_F: 5'-

GTR GTG GAG CCA TAC AAT GC -3' and ACB1_200_R: 5'- GGC ATG AAG AAG TGA AGA CGT -3')

we first mapped them onto the reference *A. caninum tubb-1* and then in an multiple sequence

alignment that included the *U. stenocephala tubb-1* we evaluated the number of mismatches

across the primer binding region. The primers were originally development for amplicon

metabarcoding to consider the frequency of the SNPs that code for the nonsynomynous

mutations at 134, 167, 198 and 200 amino acid residue of *A. caninum tubb-1* responsible for

benzimidazole resistance (Jimenez Castro et al., 2019; Venkatesan et al., 2023).

2.5. Data availability

The *U. stenocephala* sequences were submitted to GenBank under the following

accession numbers: PV848420 and PV848421 (ITS rDNA), PV848422 and PV848423 (*cox*1),

PV133906 (*tubb-1*) and PV862047 (*tubb-2*). Raw FastQ sequence data were deposited at SRA

NCBI BioProject: PRJNA1281746 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1281746).

The additional data are available at LabArchives (https://dx.doi.org/10.25833/ffcf-2175).

3. Results

3.1. Morphological and molecular confirmation of *Uncinaria stenocephala*

Four hookworms (5-7 mm) were extracted from the small intestinal contents of a red fox

(*V. vulpes*, TSF_1). The hookworms (2 males, 2 females) were consistent with the morphology of

U. stenocephala (**Figure 1**; Ransom (1924))

Two *U. stenocephala* (TSF_1.1, TSF_1.2) were molecular characterised at ITS rDNA and

*cox*1 gene. Sequenced ITS rDNA (818 nt) were identical to each other and to the reference

sequence of *U. stenocephala* (AF194145). Comparison of the new *U. stenocephala* ITS rDNA

with *U. stenocephala* ITS2 rDNA data from amplicon metabarcoding data from dog faeces

(Abdullah et al., 2025) revealed 95.8% to 100% match across 260 nt. The sequenced partial

*cox*1 (393 nt) sequence from the two *U. stenocephala* differed by a single synonymous mutation

(99.8% identity), and they were 99.2-99.5% identical to the reference *U. stenocephala* sequence

(AJ407939).

3.2 Sequencing and assembly of *Uncinaria stenocephala* genome

Sequencing of TSF_1.1 (JS6871) and TSF_1.2 (JS6872) generated 117,718,741 and

102,660,825 paired-end reads (150 bp), respectively, with average GC contents of 46% and

45%. This yielded a total of 35.2 Gb for TSF_1.1 and 30.6 Gb for TSF_1.2. After quality trimming,

117,358,176 (99.7%) and 102,353,896 (99.7%) high-quality paired-end reads were retained.

There were 4% and 23% sequence reads that mapped to the red fox (VulVul3) genome that were

removed from TSF_1.1 and TSF_1.2, respectively. The final datasets comprised 31.0 Gb for

TSF_1.1 and 21.6 Gb for TSF_1.2, both with a GC content of 45-46%. Each sample was

assembled independently. The TSF_1.1 assembly had N50 of 2,203 bp and L50 of 32,460

contigs. We then removed potential bacterial or viral contaminant contigs (TSF_1.1: 0.2% of the

contigs) to produce final assembly contigs (TSF_1.1: N50 of 2,090 bp and L50 of 32,766 contigs)

(**Table 1**). The TSF_1.2 assembly had N50 of 8,683 bp and L50 of 2,153 contigs, following

decontamination it had N50 of 8,431 bp and L50 of 2,065 contigs (**Table 1**).

The TSF_1.1 assembly yielded a contig (NODE_5499; length 7,645 nt, coverage 19.8) that

contained complete *tubb-1* gene using blast searches with other *tubb-1* from hookworms as a

query. The *tubb-2* was represented by three contigs, one contig (NODE_15465; length 4,291 nt,

coverage 30.3) included all but first exon and was truncated at the first intron. The first exon with

partial first intron was found on two contigs, where the first exon was 100% identical between

the contigs (NODE_82476; length 558 nt, coverage 29.6; NODE_38650; length 1,780 nt,

coverage 24.1). The *tubb-2* complete coding sequence was fused from the contigs, the

nucleotide sequence is truncated in the first exon. There were no contigs that would contain

putative *tubb-1* and *tubb-2* in the TSF_1.1 assembly.

3.3 Characterisation of *tubb-1* and *tubb-2* of *Uncinaria stenocephala*

The putative *tubb-1* and *tubb-2* of *Uncinaria stenocephala* genes were used to reference

map the available sequence reads. A total of 1,774 and 1,256 reads were successfully mapped

to the *tubb-1*, resulting in an average coverage of 59.7 and 41.6 bases per position across the

covered regions from TSF_1.1 and TSF_1.2, respectively. Similarly, a total of 2,335 and 1,610

reads were successfully mapped to the *tubb-2*, resulting in an average coverage of 68.7 and

47.2 bases per position across the covered regions from TSF_1.1 and TSF_1.2, respectively.

The *U. stenocephala tubb-1* gene was 4,011 nt long including 10 exons homologous to

other *btub-1* form hookworms (**Figure 1A**). The largest intron between the exon 4 and 5 was

1,688 nt, shorter compared to the intron from *A. caninum tubb-1* (46 nt: ANCCAN_08153 and

1217 nt: DQ459314). The *U. stenocephala btub-2* gene has 9 exons homologous to those of

other hookworms (**Figure 1B**). VarScan has detected 9 independent heterozygous nucleotide

position in TSF_1.1 and TSF_1.2 within *tubb-1* of *U. stenocephala*. Within *tubb-2* of *U.*

stenocephala there were 48 and 32 heterozygous positions in TSF_1.1 and TSF_1.2,

respectively.

The coding protein sequence of *U. stenocephala* TUBB-1 had two amino acid changes

when compared to other hookworms (S37Q and G441A), the remaining animo acid sequence

was identical with other hookworms. The TUBB-2 protein of *U. stenocephala* had five amino

acid changes compared to the TUBB-2 from *Ancylostoma* species: E39D, T40S, N115S, L130I

and A287S. In addition, there is a single change shared with *N. americanus* TUBB-2 (T439G).

Both TUBB-1 and TUBB-2 proteins included tubulin and β-tubulins protein motifs:

VPRAVLVDLEP, SLGGGTGSG, VVEPYN and the β-tubulin specific motif FKR (**Figure 1C**).

We then compared the primers designed by (Jimenez Castro et al., 2019) for the

amplification of the canonical SNPs in *tubb-1* of *A. caninum* with that of *U. stenocephala* (**Figure**

2A). The ACB1_167_F primer sequence was a perfect match, while ACB1_167_F had single,

G>A, mismatch in the middle of the primer sequence compared to the *U. stenocephala tubb-1*.

Both ACB1_200_F and ACB1_200_R had single mismatch each compared to the *U.*

stenocephala tubb-1. The mismatch in the ACB1_200_R is at the 3' terminal nucleotide position,

T>C.

4. Discussion

We successfully characterised the coding sequences of two representatives of the β-

tubulin gene family in *U. stenocephala*. Both the *tubb-1* and *tubb-2* genes are highly conserved

among hookworms and retain an identical exon–intron structure. Access to the complete *U.*

stenocephala tubb-1 sequence enabled us to confirm the presence of key residues (Q134,

F167, E198, and F200) that are associated with benzimidazole susceptibility in hookworms

(Venkatesan et al., 2023). The corresponding TUBB-1 protein sequence was found to be

homologous to the partial *U. stenocephala* sequences previously identified in amplicon

metabarcoding studies (Abdullah et al., 2025; Stocker et al., 2023; Stocker et al., 2024).

Anthelmintic resistance in companion animals is an emerging concern, and in the canine

hookworm *A. caninum*, it has become an established reality (Geary et al., 2025; von Samson-

Himmelstjerna et al., 2021). This study represents the first successful sequencing of the full

tubb-1 gene in the northern hookworm *U. stenocephala*. Characterisation of both *tubb-1* and

tubb-2 now enables population-level screening for resistance-associated SNPs in *U.*

stenocephala, similar to the foundational work for *A. caninum* by (Schwenkenbecher et al.,

2007).

The European red fox is a widespread invasive species in Australia (Stobo-Wilson et al.,

2021). The prevalence of *U. stenocephala* in red foxes was reported to range from 18.2% to

30.6% (Coman, 1973; Dybing et al., 2013; Ryan, 1976). Using ITS rDNA and

partial *cox*1 sequences, we demonstrated that *U. stenocephala* isolates from red foxes are

genetically indistinguishable from those found in domestic dogs (Abdullah et al., 2025; Górski

et al., 2023). Given that wild red foxes are not exposed to anthelmintic treatments, their high

infection rates could represent a substantial refugium for benzimidazole-susceptible

hookworms. However, the detection of the *tubb-1* F167Y SNP, associated with benzimidazole

resistance, in *U. stenocephala* from an urban dog in Australia by Abdullah et al. (2025) suggests

that the epidemiology of this parasite in dogs and foxes may be more complex and distinct than

currently assumed.

Although there is currently no conclusive clinical evidence of benzimidazole treatment

failure in *U. stenocephala*, the detection of the F167Y SNP in the *tubb-1* gene by Abdullah et al.

(2025) provides early warning of potential resistance. This finding suggests that the conditions

for resistance are present, and it may only be a matter of time before the frequency of this SNP

increases to levels associated with clinical drug failure. In the absence of phenotypic validation

of the F167Y SNP as a resistance marker in *U. stenocephala*, we must remain vigilant and

anticipate the spread of benzimidazole resistance, as has occurred in *A. caninum* populations

in Australia and North America (Abdullah et al., 2025; Geary et al., 2025; Leutenegger et al.,

2024; Leutenegger et al., 2023; Venkatesan et al., 2023).

DNA polymerase fidelity is highly sensitive to mismatches at the 3' end of PCR primers,

particularly when the 3' terminal nucleotide is incorrectly base-paired, as this can hinder the

initiation of extension by the enzyme (Kwok et al., 1990). In the case of the ACB1_200_R primer,

a T>C mismatch at the 3' end poses a potential risk for PCR failure due to impaired primer

extension. However, previous studies have shown that this mismatch did not interfere with

successful amplification of the *U. stenocephala tubb-1* amplicon using the

ACB1_200_F/ACB1_200_R primer pair (Abdullah et al., 2025; Stocker et al., 2023; Stocker et al.,

2024; Venkatesan et al., 2023). With our characterisation of the *U. stenocephala tubb-1* gene

and comparative analyses with other hookworm species, there is now an opportunity to refine

these primers, either through degeneracy or redesign, to better accommodate sequence

variability. The availability of genome sequencing data and expanded hookworm reference

datasets further enhances the potential for improved diagnostic accuracy and data reliability in

high- to medium-throughput molecular applications (Ilík et al., 2024).

We concluded that the identified protein sequnces belong to *U. stenocephala* β-tubulin,

specifically TUBB-1 and TUBB-2. Burns (1991) established three protein motifs common to all

tubulins, the VPRA(V/I)(F/L)(V/L)DLEP, S(F/L/A)AGGTGSG and VV(E/N)PYN motifs. These motifs

were detected in all TUBB-1 and TUBB-2 from hookworms, with a sinlge exception. There was a

deviation at the third residue of the second motif (A140G) aparently shared across all

hookworms. All β-tubulin posses the motive F(K/R)R motif that was conserved across all

hookworms. Whereas α- and γ-tubulin have KFD or RKR motif instead, respectively (Burns,

1991). Burns (1991) considered the hypervariable regions of the tubulin genes to infer functional

residues. The divergent TUBB-1 residues and four of the six divergent TUBB-2 residues (E39D,

T40S, A287S, T349G) of *U. stenocephala* were located in hypervariable regions and so unlikely

to impact the function of the β-tubulins. The benzimidazole-susceptible residues Q134, F167,

E198, and F200 were present in *U. stenocephala* TUBB-1 and were homologues to those in other

hookworm sequences.

5. Conclusion

Benzimidazole- and multidrug-resistant *A. caninum* are present in domestic dog

populations across North America and Australia (Abdullah et al., 2025; Kitchen et al., 2019;

Venkatesan et al., 2023)**.** In contrast, although *U. stenocephala* is the predominant hookworm

species in canines from cooler regions such as continental Europe, southern Australia, and

likely other under-sampled areas, data on its distribution and resistance profile remain scarce.

By characterising the complete coding regions of *tubb-1* and *tubb-2*, we enable population-level

screening for benzimidazole resistance-associated SNPs and lay the groundwork for

epidemiological studies in the 'northern hookworm' *U. stenocephala*. Genetic screening

revealed widespread SNPs in *A. caninum tubb-1* linked to benzimidazole resistance, indicating

that these alleles circulated within populations long before clinical drug failures were

recognised by veterinary parasitologists (Geary et al., 2025). Geary et al. (2025)recently

proposed a set of action points to address resistance in *A. caninum*; we argue that these

recommendations are equally relevant and urgently needed for both *A. caninum* and *U. stenocephala*.

Acknowledgements

We acknowledge the Coastal Wildlife Management, Australia, that for donating the red fox material. T. Stocker would like to acknowledge the support and encouragement from Megan and Heidi. The computational part was supported by Galaxy Australia, a service provided by Australian BioCommons and its partners; the service receives NCRIS funding through Bioplatforms Australia, as well as The University of Melbourne and Queensland Government RICF funding.

Author contributions: CRediT

Thomas Stocker: Conceptualization, Formal analysis, Investigation, Writing – original draft. **Jan Šlapeta**: Conceptualization, Funding acquisition, Project administration, Validation, Writing – review and editing.

Funding sources

The work was funded by the Betty & Keith Cook Canine Research Fund (The University of Sydney).

References

Abdullah, S., Stocker, T., Kang, H., Scott, I., Hayward, D., Jaensch, S., Ward, M.P., Jones, M.K.,

Kotze, A.C., Šlapeta, J., 2025. Widespread occurrence of benzimidazole resistance

single nucleotide polymorphisms in the canine hookworm, *Ancylostoma caninum*, in

Australia. Int. J. Parasitol. 55, 173-182. https://doi.org/10.1016/j.ijpara.2024.12.001.

Andrews, S. 2010. A Quality Control Tool for High Throughput Sequence Data. In FastQC

(Babraham Bioinformatics).

Beveridge, I., 2002. Australian hookworms (Ancylostomatoidea): a review of the species

present, their distributions and biogeographical origins. Parassitologia 44, 83-88.

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinform. 30, 2114-2120.

https://doi.org/10.1093/bioinformatics/btu170.

Bowles, J., Blair, D., McManus, D.P., 1992. Genetic variants within the genus *Echinococcus*

identified by mitochondrial DNA sequencing. Mol. Biochem. Parasitol. 54, 165-173.

https://doi.org/10.1016/0166-6851(92)90109-w.

Burns, R.G., 1991. α-, β-, and γ-tubulins: sequence comparisons and structural constraints.

Cell Motil. Cytoskeleton 20, 181-189. https://doi.org/10.1002/cm.970200302.

Chilton, N.B., Huby-Chilton, F., Gasser, R.B., 2003. First complete large subunit ribosomal RNA

sequence and secondary structure for a parasitic nematode: phylogenetic and

diagnostic implications. Mole. Cell. Probes 17, 33-39. https://doi.org/10.1016/S0890-

8508(02)00107-X.

Collins, J.B., Stone, S.A., Koury, E.J., Paredes, A.G., Shao, F., Lovato, C., Chen, M., Shi, R., Li, A.Y.,

Candal, I., Al Moutaa, K., Moya, N.D., Andersen, E.C., 2024. Quantitative tests of

albendazole resistance in *Caenorhabditis elegans* beta-tubulin mutants. Int. J. Parasitol.

Drugs Drug Resist. 25, 100556. https://doi.org/10.1016/j.ijpddr.2024.100556.

Coman, B.J., 1973. Helminth parasites of the fox (*Vulpes vulpes*) in Victoria. Aust. Vet. J. 49, 378-384. https://doi.org/10.1111/j.1751-0813.1973.tb09345.x.

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane, T., McCarthy, S.A., Davies, R.M., Li, H., 2021. Twelve years of SAMtools and BCFtools. Gigascience 10. https://doi.org/10.1093/gigascience/giab008.

Dilks, C.M., Hahnel, S.R., Sheng, Q., Long, L., McGrath, P.T., Andersen, E.C., 2020. Quantitative benzimidazole resistance and fitness effects of parasitic nematode beta-tubulin alleles. Int. J. Parasitol. Drugs Drug Resist. 14, 28-36. https://doi.org/10.1016/j.ijpddr.2020.08.003.

Dilks, C.M., Koury, E.J., Buchanan, C.M., Andersen, E.C., 2021. Newly identified parasitic nematode beta-tubulin alleles confer resistance to benzimidazoles. Int. J. Parasitol. Drugs Drug Resist. 17, 168-175. https://doi.org/10.1016/j.ijpddr.2021.09.006.

Dybing, N.A., Fleming, P.A., Adams, P.J., 2013. Environmental conditions predict helminth prevalence in red foxes in Western Australia. Int. J. Parasitol. Parasites Wildl. 2, 165-172. https://doi.org/10.1016/j.ijppaw.2013.04.004.

Galaxy Community, 2024. The Galaxy platform for accessible, reproducible, and collaborative data analyses: 2024 update. Nucleic Acids Res. 52, W83-W94. https://doi.org/10.1093/nar/gkae410.

Gandasegui, J., Grau-Pujol, B., Novela, V., Muchisse, O., Cambra-Pellejà, M., Cossa, A., Jamine, J.C., Sacoor, C., Brienen, E.A.T., Catala-Moll, F., van Lieshout, L., Martínez-Valladares, M., Paredes, R., Muñoz, J., Doyle, S.R., 2024. Deep-amplicon sequencing of the complete beta-tubulin gene in *Trichuris trichiura* before and after albendazole treatment. Int. J. Parasitol. Drugs Drug Resist. 26, 100570. https://doi.org/10.1016/j.ijpddr.2024.100570.

Geary, T.G., Drake, J., Gilleard, J.S., Chelladurai, J.R.J.J., Jimenez Castro, P.D., Kaplan, R.M., Marsh, A.E., Reinemeyer, C.R., Verocai, G.G., 2025. Multiple anthelmintic drug

401 resistance in the canine hookworm *Ancylostoma caninum*: AAVP position paper and

402 research needs. Vet. Parasitol., 110536. https://doi.org/10.1016/j.vetpar.2025.110536.

403 Gibbs, H.C., 1961. Studies on the life cycle and developmental morphology of *Dochmoides*

404 *stenocephala* (Railliet 1884) (Ancylostomidae: Nematoda). Can. J. Zool. 39, 325-348.

405 https://doi.org/10.1139/z61-037.

406 Górski, P., Długosz, E., Bartosik, J., Bąska, P., Łojek, J., Zygner, W., Karabowicz, J., Wiśniewski,

407 M., 2023. Morphological and molecular comparison of nematodes from the family

408 Ancylostomatidae isolated from selected species of carnivorous mammals in central

409 Poland. Med. Weter. 79, 223-226. https://doi.org/10.21521/MW.6766.

410 Howe, K.L., Bolt, B.J., Cain, S., Chan, J., Chen, W.J., Davis, P., Done, J., Down, T., Gao, S., Grove,

411 C., Harris, T.W., Kishore, R., Lee, R., Lomax, J., Li, Y., Muller, H.M., Nakamura, C., Nuin,

412 P., Paulini, M., Raciti, D., Schindelman, G., Stanley, E., Tuli, M.A., Van Auken, K., Wang,

413 D., Wang, X., Williams, G., Wright, A., Yook, K., Berriman, M., Kersey, P., Schedl, T., Stein,

414 L., Sternberg, P.W., 2016. WormBase 2016: expanding to enable helminth genomic

415 research. Nucleic Acids Res 44, D774-780. https://doi.org/10.1093/nar/gkv1217.

416 Howe, K.L., Bolt, B.J., Shafie, M., Kersey, P., Berriman, M., 2017. WormBase ParaSite - a

417 comprehensive resource for helminth genomics. Mol. Biochem. Parasitol. 215, 2-10.

418 https://doi.org/10.1016/j.molbiopara.2016.11.005.

419 Hu, M., Chilton, N.B., Zhu, X., Gasser, R.B., 2002. Single-strand conformation polymorphism-

420 based analysis of mitochondrial cytochrome c oxidase subunit 1 reveals significant

421 substructuring in hookworm populations. Electrophoresis 23, 27-34.

422 https://doi.org/10.1002/1522-2683(200201)23:1%3C27::aid-elps27%3E3.0.co;2-7.

423 Ilík, V., Schwarz, E.M., Nosková, E., Pafčo, B., 2024. Hookworm genomics: dusk or dawn? Trends

424 Parasitol. 40, 452-465. https://doi.org/10.1016/j.pt.2024.04.003.

425 International Helminth Genomes, C., 2019. Comparative genomics of the major parasitic

426 worms. Nat. Genet. 51, 163-174. https://doi.org/10.1038/s41588-018-0262-1.

Jimenez Castro, P.D., Howell, S.B., Schaefer, J.J., Avramenko, R.W., Gilleard, J.S., Kaplan, R.M.,

2019. Multiple drug resistance in the canine hookworm *Ancylostoma caninum*: an

emerging threat? Parasit. Vectors 12, 576. https://doi.org/10.1186/s13071-019-3828-6.

Kitchen, S., Ratnappan, R., Han, S., Leasure, C., Grill, E., Iqbal, Z., Granger, O., O'Halloran, D.M.,

Hawdon, J.M., 2019. Isolation and characterization of a naturally occurring multidrug-

resistant strain of the canine hookworm, *Ancylostoma caninum*. Int. J. Parasitol. 49,

397-406. https://doi.org/10.1016/j.ijpara.2018.12.004.

Koboldt, D.C., Zhang, Q., Larson, D.E., Shen, D., McLellan, M.D., Lin, L., Miller, C.A., Mardis,

E.R., Ding, L., Wilson, R.K., 2012. VarScan 2: somatic mutation and copy number

alteration discovery in cancer by exome sequencing. Genome Res. 22, 568-576.

https://doi.org/10.1101/gr.129684.111.

Kwok, S., Kellogg, D.E., McKinney, N., Spasic, D., Goda, L., Levenson, C., Sninsky, J.J., 1990.

Effects of primer-template mismatches on the polymerase chain reaction: human

immunodeficiency virus type 1 model studies. Nucleic Acids Res. 18, 999-1005.

https://doi.org/10.1093/nar/18.4.999.

Leutenegger, C.M., Evason, M.D., Willcox, J.L., Rochani, H., Richmond, H.L., Meeks, C., Lozoya,

C.E., Tereski, J., Loo, S., Mitchell, K., Andrews, J., Savard, C., 2024. Benzimidazole F167Y

polymorphism in the canine hookworm, *Ancylostoma caninum*: Widespread

geographic, seasonal, age, and breed distribution in United States and Canada dogs.

Int. J. Parasitol. Drugs Drug Resist. 24, 100520.

https://doi.org/10.1016/j.ijpddr.2024.100520.

Leutenegger, C.M., Lozoya, C.E., Tereski, J., Savard, C., Ogeer, J., Lallier, R., 2023. Emergence of

Ancylostoma caninum parasites with the benzimidazole resistance F167Y

polymorphism in the US dog population. Int. J. Parasitol. Drugs Drug Resist. 21, 131-140.

https://doi.org/10.1016/j.ijpddr.2023.01.001.

Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows-Wheeler

transform. Bioinform. 25, 1754-1760. https://doi.org/10.1093/bioinformatics/btp324.

Lightner, L., Christensen, B.M., Beran, G.W., 1978. Epidemiologic findings on canine and feline

intestinal nematode infections from records of the Iowa state University Veterinary

Clinic. J. Am. Vet. Med. Assoc. 172, 564-567.

https://doi.org/10.2460/javma.1978.172.05.564.

Miller, T.A., 1968. Pathogenesis and immunity in hookworm infection. Trans. Royal Soc. of

Trop.Med. Hyg 62, 473-489. https://doi.org/10.1016/0035-9203(68)90130-2.

Nadler, S.A., Hoberg, E.P., Hudspeth, D.S., Rickard, L.G., 2000. Relationships of *Nematodirus*

species and *Nematodirus battus* isolates (Nematoda: Trichostrongyloidea) based on

nuclear ribosomal DNA sequences. J. Parasitol. 86, 588-601.

https://doi.org/10.1645/0022-3395(2000)086[0588:ronsan]2.0.co;2.

Prjibelski, A., Antipov, D., Meleshko, D., Lapidus, A., Korobeynikov, A., 2020. Using SPAdes De

Novo Assembler. Curr. Protoc. in Bioinform. 70, e102. https://doi.org/10.1002/cpbi.102.

Ransom, B.H., 1924. Hookworms of the genus *Uncinaria* of the dog, fox, and badger. Proc. U. S.

Natl. Mus. 65. https://doi.org/10.5479/si.00963801.65-2533.1.

Ryan, G.E., 1976. Helminth parasites of the fox (*Vulpes vulpes*) in New South Wales. Aust. Vet. J.

52, 126-131. https://doi.org/10.1111/j.1751-0813.1976.tb05445.x.

Saunders, G.I., Wasmuth, J.D., Beech, R., Laing, R., Hunt, M., Naghra, H., Cotton, J.A.,

Berriman, M., Britton, C., Gilleard, J.S., 2013. Characterization and comparative analysis

of the complete *Haemonchus contortus* beta-tubulin gene family and implications for

benzimidazole resistance in strongylid nematodes. Int. J. Parasitol. 43, 465-475.

https://doi.org/10.1016/j.ijpara.2012.12.011.

Schwarz, E.M., Hu, Y., Antoshechkin, I., Miller, M.M., Sternberg, P.W., Aroian, R.V., 2015. The

genome and transcriptome of the zoonotic hookworm *Ancylostoma ceylanicum* identify

infection-specific gene families. Nat. Genet. 47, 416-422.

https://doi.org/10.1038/ng.3237.

Schwenkenbecher, J.M., Albonico, M., Bickle, Q., Kaplan, R.M., 2007. Characterization of beta-

tubulin genes in hookworms and investigation of resistance-associated mutations using

real-time PCR. Mol. Biochem. Parasitol. 156, 167-174.

https://doi.org/10.1016/j.molbiopara.2007.07.019.

Seemann, T., Gladman, S. 2012. Fasta Statistics: Display summary statistics for a fasta file

(GitHub, Galaxy Project).

Stobo-Wilson, A.M., Murphy, B.P., Crawford, H.M., Dawson, S.J., Dickman, C.R., Doherty, T.S.,

Fleming, P.A., Gentle, M.N., Legge, S.M., Newsome, T.M., Palmer, R., Rees, M.W., Ritchie,

E.G., Speed, J., Stuart, J.M., Thompson, E., Turpin, J., Woinarski, J.C.Z., 2021. Sharing

meals: Predation on Australian mammals by the introduced European red fox

compounds and complements predation by feral cats. Biol. Conserv. 261, 109284.

https://doi.org/10.1016/j.biocon.2021.109284.

Stocker, T., Scott, I., Šlapeta, J., 2023. Unambiguous identification of *Ancylostoma caninum* and

Uncinaria stenocephala in Australian and New Zealand dogs from faecal samples. Aust.

Vet. J. 101, 373-376. https://doi.org/10.1111/avj.13272.

Stocker, T., Ward, M.P., Šlapeta, J., 2024. Nationwide USA re-analysis of amplicon

metabarcoding targeting β-tubulin isoform-1 reveals absence of benzimidazole resistant

SNPs in *Ancylostoma braziliense*, *Ancylostoma tubaeforme* and *Uncinaria*

stenocephala. Vet. Parasitol. 327, 110118.

https://doi.org/10.1016/j.vetpar.2024.110118.

Venkatesan, A., Jimenez Castro, P.D., Morosetti, A., Horvath, H., Chen, R., Redman, E., Dunn,

K., Collins, J.B., Fraser, J.S., Andersen, E.C., Kaplan, R.M., Gilleard, J.S., 2023. Molecular

evidence of widespread benzimidazole drug resistance in *Ancylostoma caninum* from

domestic dogs throughout the USA and discovery of a novel beta-tubulin benzimidazole

resistance mutation. PLoS Pathog. 19, e1011146.

https://doi.org/10.1371/journal.ppat.1011146.

von Samson-Himmelstjerna, G., Thompson, R.A., Krucken, J., Grant, W., Bowman, D.D.,

Schnyder, M., Deplazes, P., 2021. Spread of anthelmintic resistance in intestinal

helminths of dogs and cats is currently less pronounced than in ruminants and horses -

Yet it is of major concern. Int. J. Parasitol. Drugs Drug Resist. 17, 36-45.

https://doi.org/10.1016/j.ijpddr.2021.07.003.

Walker, M., Jacobs, D., 1985. Pathophysiology of *Uncinaria stenocephala* infections of dogs.

Vet. Annu. 25, 263-271.

Wood, D.E., Salzberg, S.L., 2014. Kraken: ultrafast metagenomic sequence classification using

exact alignments. Genome Biol. 15, R46. https://doi.org/10.1186/gb-2014-15-3-r46.

Xhaxhiu, D., Kusi, I., Rapti, D., Kondi, E., Postoli, R., Rinaldi, L., Dimitrova, Z.M., Visser, M.,

Knaus, M., Rehbein, S., 2011. Principal intestinal parasites of dogs in Tirana, Albania.

Parasitol. Res. 108, 341-353. https://doi.org/10.1007/s00436-010-2067-8.

Table 1. Assembly of the whole genome sequencing of *Uncinaria stenocephala*

	TSF_1.1	TSF_1.1*	TSF_1.2	TSF_1.2*
L50	32,460	32,766	2,152	2,065
N50	2,203	2,090	8,683	8,431
Number of bp	341,439,219	333,472,904	62,113,770	57,967,112
Number of sequences	1,039,638	1,037,453	50,108	49,593
GC content (%)	45.5	45.6	44.6	44.7

520   * putative bacterial and viral contaminants removed via Kraken

Figures:

Figure 1: Morphology of sampled *Uncinaria stenocephala*. (A) Dorso-ventral view of the anterior end. Bicuspid cutting plates are visualised with an arrow for each cutting plate. (B) The lateral view of the anterior end. The internal cephalic papilla is labelled with a black arrow. The white arrows signify the boundary line between the thick ventral portion and the thinner dorsal portion of the lateral wall of the oral capsule. This boundary crosses through the internal cephalic papilla which is consistent with *U. stenocephala*. Scale bar is representative of 100 µm.

Figure 2: Comparison of *Uncinaria stenocephala* structure of *tubb-1* and *tubb-2* genes with other hookworms. (A) Gene sequence showing exon-intron boundaries of *tubb-1* of hookworms. Regions where primers targeting *A. caninum tubb-1* (ACB1_167_F: 5'-GG**Y** GCA GGA AAC AAC TG -3', ACB1_167_R: 5'- CTT TGG TGA GGG GAC AAC A -3', ACB1_200_F: 5'- GT**R** GTG GAG CCA TAC AAT GC -3' and ACB1_200_R: 5'- GGC ATG AAG AAG TGA AGA CGT -3') and benzimidazole-susceptible residues in *tubb-1* (Q134, F167, E198, and F200) are enlarged to show nucleotide sequences (identical residues to the top sequence as dots). (B) Gene sequence showing exon-intron boundaries of *tubb-2* of hookworms. Names with (*) indicate

that sequence data were obtained from WormBase Parasite, *tubb-1* (ANCCAN_08153,

Acey_s0035.v2.g13228, ANCDUO_05511, Necator_chrII.pre1.g8352 and NBR_0000799801) and

tubb-2   (ANCCAN_23405,   Acey_s0432.v2.g1106,   Necator_chrII.pre1.g8352   and

Nippo_chrII.g5566), otherwise sequences are labelled by GenBank accession number. (C)

Amino acid alignment regions of TUBB-1 and TUBB-2 showing the tubulin motifs

VPRA(V/I)(F/L)(V/L)DLEP, S(F/L/A)AGGTGSG and VV(E/N)PYN motifs. TUBB-1 second motive

devaiates from the standard motive (A140G, arrowed). The β-tubulin conserved protein motive

F(K/R)R motif in preset across all sequences.