Perceived time drives physical fatigue

Pierre-Marie Matta[1,2,*], Julien Duclay[2], Robin Baurès[1§], Andrea Alamia[1§]

[1]Univ Toulouse, CNRS, CerCo, Toulouse, France

[2]Univ Toulouse, Inserm, ToNIC, Toulouse, France

*Corresponding author: Pierre-Marie Matta (pierre-marie.matta@inserm.fr)

[§]These authors contributed equally to this work.

Conflict of interest: The authors declare that they have no competing interests.

Data availability statement: The datasets will be made publicly available on the Open Science Framework (OSF) upon publication of the manuscript.

Ethical statement: The study procedure was approved by the French Ethical 'Comité de Protection des Personnes' (number: 2023-A00412-43).

Acknowledgements: This project has received funding from the H2020 European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreements No 101075930 – Andrea Alamia). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them. Pierre-Marie Matta received funding from the French Ministère de l'Enseignement Supérieur et de la Recherche (MESR). We thank Maxime Picquet, Romain Martinie, Khiara Vorzillo, and Nina Bertho for their valuable help with the data acquisition. We are also grateful to Laurent Gonthier for his technical support in setting up the online force acquisition card, and to Nathalie Vayssiere for her assistance in obtaining the ethical agreement.

Abstract

Recent studies suggest that fundamental physiological processes, such as physical fatigue, rely on perceived rather than actual time. However, the neural correlates underlying this effect and its disentanglement from motivational factors (i.e., performance goals) remain unknown. To investigate the time deception effect on fatigue, we developed a novel EEG design in which participants (N = 24) performed 100 isometric contractions at a fixed pace and resistance in four distinct sessions. The actual contraction duration (short or long) and the calibration of the displayed clock (normal or biased toward acceleration or deceleration) were independently manipulated across sessions to examine whether fatigue and its neural correlates evolved in response to perceived or actual time. Our results show an accumulation of physical fatigue that follows the perceived time, irrespective of motivational factors. This effect was consistently observed only when the clock was slowed down. This time-deception effect involved frontal theta- and beta-band dynamics: theta modulated only under the slowed clock and beta robustly shaped by perceived time across both slowed and accelerated clocks, while motor beta showed no modulation. Further analyses highlighted the key role of frontal oscillatory dynamics in the effectiveness of the time-deception effect on physical fatigue.

Significance statement

Can a clock change the course of physical fatigue? In this study, we addressed this question while controlling for motivational confounds and monitoring EEG-associated power modulations. Our findings demonstrate a slowed-down fatigue accumulation in the presence of a slowed-down clock. This time manipulation effect was driven by a frontal oscillatory dynamic that largely followed the perceived time. These results highlight the direct influence of psychological factors on physiological processes and unveil the neural correlates underlying this effect.

Introduction

The relationship between the mind and body has been a subject of inquiry for centuries (Dijkerman & Lenggenhager, 2018), raising fundamental questions about how mental states influence bodily processes. A prevalent approach to exploring this relationship involves deceptively manipulating participants' beliefs while recording psychophysiological variables (Crum et al., 2011; Crum & Langer, 2007; H. S. Jones et al., 2013; Williams et al., 2014). Among the various psychological parameters that can be successfully deceived, manipulating the subjective passage of time has been suggested as particularly relevant because of the trust we routinely place in its measurement (Thönes et al., 2018). This manipulation is typically achieved by covertly altering the clock's speed displayed to participants during a task, either slowing it down or accelerating it (Thönes et al., 2018). For instance, when the passage of time is artificially slowed down, studies have reported reduced perceived pain in response to thermal stimulation (Pomares et al., 2011) but also a smaller decrease in blood glucose level (Park et al., 2016) and slower healing (Aungle & Langer, 2023). Similarly, studies interested in physical activity have shown that, in the presence of a slowed-down clock, there is a slower spontaneous motor activity (Craik & Sarbin, 1963) and a slower accumulation of physical fatigue (Matta et al., 2024). Conversely, accelerating the perceived passage of time has been associated with opposite effects, with the notable exception of physical fatigue, which, to our knowledge, remains unexplored in this context.

Physical fatigue, broadly defined as the decline in the ability to generate force or sustain performance (Bigland-Ritchie & Woods, 1984; Gandevia, 2001), is a central determinant of endurance, productivity, and daily functioning (Enoka & Duchateau, 2016). Beyond its role in limiting athletic and occupational performance, it is also closely linked to injury risk (Chan, 2011; Cunningham et al., 2022; Ibrahim et al., 2023; C. M. Jones et al., 2017; Kreher & Schwartz, 2012; Rodrigues et al., 2023) and impaired quality of life (Behrens et al., 2023). Understanding how it may be shaped by psychological factors is therefore of both theoretical and practical significance, with implications for athletes, workers in physically demanding professions, and individuals in everyday life. This makes physical fatigue a particularly relevant phenomenon for investigating deceptive manipulations. Furthermore, physical fatigue develops progressively as a function of task duration, reflecting the cumulative physiological and neural demands imposed by sustained activity (Bigland-Ritchie & Woods, 1984; Edwards, 1981; Enoka & Duchateau, 2008; Gandevia, 2001). As a result, fatigue is intrinsically linked to temporal representations, as information about elapsed and remaining time provides a key

reference for regulating effort throughout a task. In most goal-directed behaviors, time constitutes a fundamental performance constraint, shaping pacing strategies, effort allocation, and expectations about task completion (Brehm & Self, 1989; de Koning et al., 2011; Noakes, 2011; Tucker, 2009). Consequently, beliefs about time serve as a critical reference signal during sustained physical activity, pointing out the importance of examining how these temporal representations may influence the development of physical fatigue.

Nevertheless, important gaps remain in our understanding of how time perception influences physical fatigue: although these perceived time-dependent effects have been demonstrated for several variables (Thönes et al., 2018), their underlying neural mechanisms remain poorly understood. Moreover, motivational factors, such as performance goals, are critical confounds often unaccounted for when measuring deceptive effects. These confounds are particularly important in physical fatigue, in which previous studies have consistently employed deceptive manipulations to enhance physical performance (i.e., the amount of time a task-to-exhaustion is maintained or the time needed to complete a given distance) (Ansdell et al., 2018; Ducrocq et al., 2017; Matta et al., 2024). Consequently, previous results on physical fatigue may be alternatively explained by performance-related goals rather than time perception.

In the current study, we investigate the neural correlates of the time deception effect on the accumulation of physical fatigue and carefully isolate this effect from motivational confounds. In addition, we also test its temporal symmetry, that is, when the clock is either slowed down or accelerated. To this end, we used an original 2×2 design, in which each session consisted of 100 isometric contractions at a given pace against a fixed, individualized resistance. The contraction's actual duration (short or long) and the calibration of the displayed clock (normal or biased toward acceleration or deceleration) were manipulated between sessions to test whether physical fatigue was evolving according to the perceived (i.e., displayed) or actual time (Fig. 1A). Unlike previous studies, we tested all four possible combinations for the first time in a within-subject design. We recorded EMG and EEG signals to assess fatigue accumulation and associated neural correlates. We specifically focused on oscillatory dynamics at the sensor level in the motor and prefrontal cortex, as both regions are involved in physical fatigue and effort maintenance (Bigliassi & Filho, 2022; Matta et al., 2025; McMorris et al., 2018; Robertson & Marino, 2016).

Figure 1 – Experimental protocol and EMG/EEG computation

A. Experimental Protocol. All participants performed 100 contractions during four separate sessions. While the
rest time between contractions was identical (5s) for each session, each contraction's real (R) and perceived (P)
time were manipulated. In each session, the contractions' time was either short (10s) or long (12s), and the digital
clock displayed in front of the participants was manipulated to induce either a short (10 s) or long (12 s) perceived
duration. This resulted in a within-subject, partially crossed four-condition design manipulating Real duration (10
s vs. 12 s) and Perceived duration (10 s vs. 12 s), yielding four counterbalanced conditions: P10R10 (normal clock),
P12R12 (normal clock), P10R12 (fast clock), and P12R10 (slowed clock). Before and after the task, Maximal
Voluntary Contractions (MVC) were performed to assess fatigue induced by the 100 contractions. Resting State
(RS) EEG (3 min) was recorded before and after the task to quantify the associated EEG correlates. B. EMG/EEG
Computation. All panels represent the typical trace of a single contraction, except the upper-right panel, which
illustrates the entire task. EMG and EEG were recorded throughout the task and explicitly analyzed over a 2-10 s
window centered on each contraction (0 s = contraction onset), from which we extracted a single average value
per contraction. The EMG was computed in normalized Root Mean Square (RMS), while we computed the mean
power of the theta (θ) and an individualized beta-band (Iβ) for the EEG, in frontal electrodes and Cz (illustrated in
black and yellow respectively on the topographical plots) (see Methods for details).

Results

Design Summary

Participants performed four sessions (each spaced 3 to 10 days apart) of 100 quadriceps isometric contractions, during which EMG and EEG were continuously recorded. For each participant, the contraction intensity was identical across sessions ($\leq 20\%$ of the participant's maximal force; see Methods for details). While the rest time between contractions was identical for each session, each contraction's real (R) and perceived (P) time were independently manipulated. Each contraction could be either short (10 s) or long (12 s) in real time, and the digital clock displayed in front of the participants was manipulated to induce either a short (10 s) or long (12 s) perceived duration. This resulted in a 2 × 2 design, with the factors Real duration (10 s vs. 12 s) and Perceived duration (10 s vs. 12 s), yielding four counterbalanced conditions: P10R10 (normal clock), P12R12 (normal clock), P10R12 (fast clock), and P12R10 (slowed clock) (Fig. 1). Before and after the task, participants performed Maximal Voluntary Contractions (MVC) and 3-minute EEG resting states.

Neuromuscular fatigue

We first assessed the effects of real and perceived time on physical fatigue, quantified as the pre-post change in maximal voluntary torque (MVT). Analysis of the pre-post difference in MVT revealed no significant main effect of perceived time (p = .90), real time (p = .47), nor their interaction (p = .23). This lack of effect may stem from the relatively small variation in workload across sessions (Matta et al., 2025) combined to a limited sensitivity of pre-post MVT measurements to accurately detect physical fatigue (Lebesque et al., 2022).

We next examined whether perceived time modulated fatigue accumulation during the task, quantified as the EMG Root Mean Square (RMS) activity recorded across contractions (Kim et al., 2007; Merletti et al., 1991). Model comparison showed that allowing for a nonlinear effect of contraction number did not improve model fit relative to a linear model, ($\chi^2_{(4)} = 7.10$, p = .13), and the linear model was therefore retained. This model revealed a significant effect of contraction number ($F_{(1, 23)} = 45.97$, p < .0001), as well as significant main effects of perceived time ($F_{(1, 9546)} = 16.26$, p < .0001). The interaction between contraction number and perceived time was significant ($F_{(1, 9546)} = 15.12$, p = .0001), as was the interaction between contraction number and real time ($F_{(1, 9546)} = 32.08$, p < .0001). The three-way interaction between contraction number, perceived time, and real time was also significant ($F_{(1, 9546)} = 8.77$, p = .003), while no significant main effect of real time alone nor interaction between perceived and

real time was observed (all p > .59). Post-hoc contrasts on the estimated marginal slopes

indicated that perceived time did not affect EMG rate when real time was short (p = 1, d_z =

0.13, BF = 0.22). When real time was long, however, EMG increased significantly less for

shorter perceived time (p < .0001), highlighting a strong (d_z = 0.99) slowing of fatigue when

the clock was slowed. Consistent with this effect, real time did not show a significant effect

when perceived time was short (p = .22, d_z = 0.39, BF = 0.30). Conversely, when perceived

time was long, EMG increased less for short real time (p < .0001), indicating here a strong (d_z

= 1.24) dependency on the real time when the clock was accelerated (Fig. 2A, 2B). Together,

these results suggest that perceived time modulates the rate of physical fatigue accumulation,

effectively slowing it when the task feels shorter (Matta et al., 2024), whereas, conversely,

fatigue accumulation cannot be accelerated when the task appears longer.

Figure 2 – Physical Fatigue

A. EMG modulations over the main task. EMG is expressed in normalized Root Mean Square (RMS) over

contractions, whose increase reflects neuromuscular fatigue accumulation. The lines represent the averaged

distribution, and the shaded areas are the standard error of the mean (SEM). B. Fatigue differences computed for

each contrast. The upper panel illustrates perceived-time effects by holding real duration constant and contrasting

long versus short perceived durations, whereas the lower panel illustrates real-time effects by holding perceived

duration constant and contrasting long versus short real durations. Both panels represent the difference in slope of

the EMG-RMS over the bins. The bar plots represent model-based estimates of condition differences, with error

bars indicating 95% confidence intervals. Intervals not crossing zero indicate statistically significant effects at a

Bonferroni-corrected alpha of 0.05.

Power modulations in motor areas

After showing the influence of perceived time on physical fatigue, we investigated the neural correlates underlying this effect. We first examined oscillatory activity in the Cz electrode, commonly used as an approximation of activity in the motor cortex (Dal Maso et al., 2012; Forrester et al., 2006; Gordon et al., 2023; Meier et al., 2008), a key region involved in executing voluntary contractions (Brown, 2000; Pearson, 2000). Focusing on this sensor, we assessed relative spectral power changes in each participant's individualized beta-band (Iβ) (Matta et al., 2025) (i.e., a 4 Hz spectral window in the beta band with the largest desynchronization, see Methods for details).

We compared two linear mixed-effects models differing in the complexity of the contraction term (natural spline with 1 vs. 2 degrees of freedom). Model comparison indicated that the second-degree spline model provided a better fit ($\chi^2_{(4)} = 11.18$, p = .025), and was therefore retained. Analysis of variance on the selected model revealed a significant main effect of contraction number ($F_{(2,\ 45.4)} = 8.01$, p = .001), as well as a significant interaction between contraction number and perceived time ($F_{(2,\ 9542)} = 5.28$, p = .005), showing a larger decrease in Iβ Cz power when perceived time was shorter. To better understand this effect, we compared the slope differences between the two normal sessions (P10R10 vs. P12R12) but found no difference (p = 0.60, uncorrected, $d_z = 0.11$, BF = 0.23), which may reflect the relatively small discrepancy in workload between the two sessions (Matta et al., 2025) and limits the interpretability of these results. No significant main effects of perceived or real time alone, nor significant interactions involving real time or the three-way interaction, were observed (all p > .05).

We next investigated the effects of perceived and real time on the pre-post difference in Iβ Cz, reflecting overall changes in beta activity across the session. A linear mixed-effects model including participant as a random intercept revealed no significant main effects of perceived time (p = .91) or real time (p = .89), and no perceived × real time interaction (p = .26). This lack of effect may relate to the relatively small discrepancy in workload between the sessions (Matta et al., 2025).

Power modulations in frontal electrodes

Next, we investigated oscillatory activity at frontal electrodes (F3, F4, F7, F8) – commonly used as an approximation of prefrontal cortex (PFC) activity (Okamoto et al., 2004; Robertson & Marino, 2015), as this region has been proposed to play a key role in effort maintenance

(Bigliassi & Filho, 2022; Denis, 2020; McMorris et al., 2018; Robertson & Marino, 2016). We
analyzed relative spectral power changes in the theta (θ) and Iβ bands. While frontal θ is
assumed to reflect cognitive control (Cavanagh & Frank, 2014), the β band has also been
proposed to have a similar function (Stoll et al., 2016).

To assess how perceived time modulates θ frontal activity, we first compared two linear
mixed-effects models differing in the complexity of the contraction term (natural spline with 1
vs. 2 degrees of freedom). The more complex model did not improve fit ($\chi^2_{(4)} = 4.91$, p = .30),
and the linear model was retained. The selected model revealed no significant main effects of
perceived time ($F_{(1, 9546)} = 0.90$, p = .344), real time ($F_{(1, 9546)} = 0.59$, p = .443), or contraction
number ($F_{(1, 23)} = 0.39$, p = .540). A significant interaction was observed between contraction
number and perceived time ($F_{(1, 9546)} = 29.65$, p < .0001), as well as between perceived and real
time ($F_{(1, 9546)} = 12.56$, p < .0001). The three-way interaction among contraction number,
perceived time, and real time was also significant ($F_{(1, 9546)} = 5.49$, p = .019), while the
interaction between contraction number and real time was not significant (p = .65). At short real
time, post-hoc contrast on the estimated marginal slopes was inconclusive, showing no reliable
difference in the rate of increase in θ frontal for long versus short perceived time (p = .11, d_z =
0.45, BF = 0.41). However, when real time was long, the rate of increase in θ frontal was
significantly (p < .0001) steeper for long perceived time, with a strong effect size ($d_z = 1.12$).
In parallel, real-time influence on θ slopes was inconclusive at short perceived duration (p =
.19, $d_z = 0.40$, BF = 0.46) and showed evidence for an absence of difference at long perceived
time duration (p = .63, $d_z = 0.27$, BF = 0.28) (Fig. 3A, 3B). Overall, these results indicate that
frontal theta changes across the task robustly follow perceived time when the clock is slowed
down, while the results are inconclusive regarding the accelerated clock.
247

Figure 3 – Neural correlates of time deception

A. EEG Power modulations in frontal electrodes. Relative power changes in θ (panel 1) and Iβ (panel 2) over

contractions (relative to the first contractions). For both panels, the line represents the averaged distribution, and

the shaded areas are the SEM. B. Power differences in the frontal electrodes. Power differences were computed

for each contrast. The left panels illustrate perceived-time effects by holding real duration constant and contrasting

long versus short perceived durations, whereas the right panels illustrate real-time effects by holding perceived

duration constant and contrasting long versus short real durations. The upper panels represent the difference in the

slope of the frontal θ power, while the lower panels present the difference in the means of Iβ power over the bins.

The bar plots represent model-based estimates of condition differences, with error bars indicating 95% confidence

intervals. Intervals not crossing zero indicate statistically significant effects at a Bonferroni-corrected alpha of

0.05.

For Iβ frontal activity, model comparison showed that a second-degree spline model

provided a significantly better fit than the linear model ($\chi^2_{(4)}$ = 207.24, p < .0001) and was

therefore retained. In the selected model, a significant main effect of contraction number was

observed ($F_{(2, 45.1)}$ = 92.07, p < .0001). No significant main effects of perceived time ($F_{(1, 9542)}$ =

0.41, p = .523) or real time ($F_{(1, 9542)}$ = 0.00, p = .970) were found. A significant interaction

between contraction number and perceived time was observed ($F_{(2, 9542)}$ = 39.53, p < .0001),

while the interaction between perceived and real time showed a trend ($F_{(1, 9542)}$ = 3.76, p = .052).

All other interactions were non-significant (all p > .18). Although the interaction did not reach

conventional significance, post-hoc contrasts on the estimated marginal means were conducted

to test our a priori hypothesis regarding the influence of perceived time (Schad et al., 2020).

They revealed a robust effect of perceived time: Iβ was greater for longer perceived time, both

when real time was short (p < .0001) and when it was long (p < .0001), in both cases with very

strong effect sizes (respectively, d_z = 1.77 and d_z = 3.73). Real-time influence on Iβ at short

perceived time was significant ($p < .0001$) and large ($d_z = 1.46$), though Iβ was greater in the

short real duration compared to the long real duration. In other words, the difference is contrary

to the expected pattern if real duration were to play a role. Finally, there was no effect of real

time at long perceived time ($p = .051$, $d_z = 0.51$, BF $= 0.26$) (Fig. 3A, 3B). Together, these

results suggest that frontal beta dynamic is strongly shaped by perceived time, both when the

clock is slowed down and accelerated.

EEG-EMG Connectivity

After independently investigating the EMG and EEG power modulations, we examined their

relationship by computing subject-by-subject correlations (see Methods for details). We first

found consistently across participants a negative correlation between EMG and Iβ Cz for all

sessions ($p < .045$, $d_z > 0.43$) (Fig. 4A), indicating more EMG activity for larger Iβ

desynchronization. We then investigated the link between EMG and frontal oscillatory activity.

Similarly, we found a negative correlation between EMG activity and frontal θ-band oscillations

in P10R12 ($t_{(23)} = 2.70$, $p = .013$, $d_z = 0.46$), and as a trend in P10R10 ($t_{(23)} = 1.99$, $p = .059$,

$d_z = 0.41$, BF $= 1.14$), but no correlations in the P12R12 and P12R10 sessions ($p > 0.36$, $d_z <$

0.19, BF < 0.32) (Fig. 4B). Conversely, we found a positive correlation between the EMG

activity and the Iβ in the P12R10 session ($t_{(23)} = 2.25$, $p = .035$, $d_z = 0.46$) and, as a trend, in the

P12R12 one ($t_{(23)} = 1.75$, $p = .094$, $d_z = 0.36$, BF $= 0.80$), while there was no correlation in the

sessions whose perceived time was shorter (P10, $p > 0.75$, $d_z < 0.66$, BF < 0.23, for both) (Fig.

4C). These results mirror the trends observed in frontal power modulations (Fig. 3A, 3B),

showing lower Frontal-EMG correlations for short perceived times and larger correlations for

longer ones, suggesting their dependency on perceived rather than actual time.

Figure 4 – Brain-muscle correlations

Within-subject correlations were performed between EMG and θ Frontal (A), Iβ Cz (B), and Iβ Frontal (C),

respectively. The resulting r values were then compared against 0 using t-tests. The bar plots show the average

distribution, and the error bars represent the 95% confidence intervals. Intervals that do not cross zero indicate

statistically significant effects with $p < .05$.

Relationship between frontal oscillatory dynamics and fatigue accumulation

Our results show that perceived time can significantly influence the accumulation of physical

fatigue and that the frontal oscillatory dynamics are robustly modulated by perceived rather

than actual time. Starting from the assumption that PFC is involved in belief representation

(Geuter et al., 2017), we hypothesized that participants with the strongest deceptive effect in

the frontal electrodes would also exhibit the largest EMG effect. To test our hypothesis, we split

the participants into two groups via a median split computed on the normalized θ-power

difference between the two sessions with the same perceived time. The first split separated

participants based on the normalized θ-power difference computed between sessions that shared

a perceived short duration (P10) but differed in real duration. In a second median split, the same

variable was computed for sessions that shared a perceived long duration (P12) but differed in

real duration. These power differences were normalized by the difference between the P12R12

and P10R10 conditions to account for the power difference observed under normal clock

conditions. In both cases, smaller differences correspond to a larger deceptive effect, as the θ

power modulation in frontal electrodes follows the perceived rather than the actual time

('Subjective-time' (ST) group). Conversely, participants with larger power differences would

rely more on actual than perceived time ('Objective-time' (OT) group). Note that we used the

θ power difference to perform our median split because of its well-established relationship with
cognitive control (Cavanagh & Frank, 2014).
323

Figure 5 – Physical fatigue accumulation in the 'Subjective-time' subgroup

A. EMG modulations over the main task. EMG is expressed in normalized RMS over contractions, whose increase
reflects neuromuscular fatigue accumulation. Panels contain the 'Subjective-time' participants only (N = 12)
(based on the deception effects measured in the θ-band, see text for details). Panel on the left was computed on
frontal θ absolute sum power difference between sessions sharing the same perceived short duration (P10) and
panel on the right between sessions sharing the same perceived long duration (P12). For both panels, the lines
represent the averaged distribution, and the error bars are the standard error of the mean. B. EMG differences.
EMG differences were computed for each contrast. The left panel illustrates perceived-time effects by holding real
duration constant and contrasting long versus short perceived durations, whereas the right panel illustrates real-
time effects by holding perceived duration constant and contrasting long versus short real durations. All panels
represent differences in the slope of the EMG-RMS across bins. For all panels, the bar plots represent model-based
estimates of condition differences, with error bars indicating 95% confidence intervals. Intervals not crossing zero
indicate statistically significant effects at a Bonferroni-corrected alpha of 0.05.

For all subgroups, a linear model with a simple contraction effect (linear) was sufficient, as
adding a second-degree spline did not improve model fit (all p > .15). In the ST subgroup
defined on the θ P10 difference, significant main effects of contraction number ($F_{(1, 11)}$ = 15.91,
p = .0021) and perceived time ($F_{(1, 4770)}$ = 17.03, p < .0001) were observed. Significant
interactions were found between contraction number and perceived time ($F_{(1, 4770)}$ = 50.40, p <
.0001), between contraction number and real time ($F_{(1, 4770)}$ = 7.05, p = .0079), and a three-way
interaction among contraction number, perceived time, and real time ($F_{(1, 4770)}$ = 8.77, p = .0031).
Post-hoc contrasts on the estimated marginal slopes indicated that EMG increased more for

long perceived time, both when real time was short (p = .014, d_z = 0.84) and long (p < .001, d_z = 2.05), indicating that EMG can be both slowed down and accelerated, according to the clock calibration. Accordingly, there was no significant influence of real duration when perceived time was short (p = 1, d_z = 0.06, BF = 0.29). Finally, when perceived time was long, EMG rate was still modulated by real time (p = <.0001, d_z = 1.15) (Fig. 5A, 5B).

In the ST subgroup defined on the θ P12 difference, contraction number ($F_{(1, 11)}$ = 25.13, p < .0001), perceived time ($F_{(1, 4770)}$ = 32.40, p < .0001), and real time ($F_{(1, 4770)}$ = 23.38, p < .0001) showed significant main effects, while interactions between contraction number and perceived time ($F_{(1, 4770)}$ = 38.89, p < .0001), between contraction number and real time ($F_{(1, 4770)}$ = 35.14, p < .0001), and the three-way interaction ($F_{(1, 4770)}$ = 10.39, p = .0013) were also significant. EMG increased more at long perceived time when real time was long (p < .0001, d_z = 1.93), supporting a modulation by perceived time under slowed clocks, while the evidence was inconclusive at short real time (p = .13, d_z = 0.61, BF = 0.32). Finally, real duration provided inconclusive evidence when perceived time was short (p = 0.22, d_z = 0.55, BF = 0.46) and had a significant influence under long perceived time (p < .0001, d_z = 1.87) (Fig. 5A, 5B).

In the OT subgroup defined on the θ P10 difference, significant effects of contraction number ($F_{(1, 11)}$ = 33.09, p < .0001), real time ($F_{(1, 4770)}$ = 5.97, p = .015), and an interaction between contraction number and real time ($F_{(1, 4770)}$ = 30.48, p < .0001) were observed, highlighting a dependency on real time only.

Finally, in the OT subgroup defined on the θ P12 difference, main effects of real time ($F_{(1, 4770)}$ = 21.91, p < .0001) and a significant interaction between perceived and real time ($F_{(1, 4770)}$ = 9.91, p = .0017) were observed, along with a significant interaction between contraction number and real time ($F_{(1, 4770)}$ = 4.77, p = .029). Post-hoc contrasts on the estimated marginal means showed that EMG was greater with short perceived time at long real time (p < .0001, d_z = 1.31) – that is, in the opposite direction if perceived duration were to play a role, while there was no difference at short real time (p = 0.25, d_z = 0.54, BF = 0.29). Similarly, EMG was greater with short real duration at long perceived time (p < .0001, d_z = 2.06), while there was no difference at short perceived time (p = 1, d_z = 0.21, BF = 0.29).

While this analysis remains exploratory and is partially limited by a median split and a small sample size (n = 12 per group), these results suggest that the frontal power changes play a key role in the effectiveness of the deceptive effect on fatigue accumulation, as measured by EMG. Interestingly, it reveals that, in the ST group, fatigue can be slowed down and accelerated depending on the clock calibration.

Link between interoceptive awareness and deception

To further explore the individual differences that may affect the strength of the deceptive effect, we considered participants' interoceptive awareness, measured via the Multidimensional Assessment of Interoceptive Awareness (MAIA) (Mehling et al., 2012) after their last session, as a potential moderating factor. We specifically focused on the 'Noticing' and 'Body Listening' subscales, which are more closely linked to bodily processes (Iodice et al., 2019). We then tested the correlation between the Noticing and Body Listening scores with the normalized absolute difference of neuromuscular fatigue accumulation (i.e., EMG) for the two sessions with the same subjective time (P10 or P12) but a different real time. In short, a low normalized index indicates no difference between the two sessions, suggesting a large deceptive effect, while a high index reflects the opposite. We hypothesized a positive correlation between the interoceptive score and the normalized index, i.e., the lower the interoceptive awareness, the higher the deceptive effect, and vice-versa. Using non-parametric Kendall's tau correlation, we found a positive correlation between the Body Listening score and the normalized difference in sessions with the same short perceived time only ($\tau = 0.29$, $p = 0.029$, unilateral test, H1: positive correlation). No significant correlation was found with the Noticing score ($p > 0.27$, $BF < 0.47$). These results suggest that the participant's body awareness might predict the deceptive effect on neuromuscular fatigue accumulation in the slow-down clock configuration.

As our previous findings suggested a role for frontal oscillatory dynamics in the deceptive effect, we tested the correlation between the Noticing and Body Listening scores and the normalized absolute difference in θ frontal power modulations across the two sessions with the same perceived time. We found a positive correlation between the Body Listening score and the θ normalized difference from sessions with the same long perceived time only ($\tau = 0.31$, $p = 0.020$, unilateral test, H1: positive correlation). While this result supports a link between interoception awareness and the deceptive effect, it remains unclear why this effect appears only when time is accelerated, but not when it is slowed down, as no other significant correlations were found ($p > 0.11$, $BF < 1.02$, unilateral test, H1: positive correlation).

Discussion

In the current study, we used a novel design to test the influence of time deception on physical fatigue accumulation while monitoring associated EEG power modulations. We found that physical fatigue accumulation was modulated by perceived time, specifically accumulating more slowly when the perceived duration was short, independent of real time. Importantly, this deceptive effect was observed in the absence of motivational confounds, demonstrating, for the first time, the direct influence of time deception on the course of physical fatigue. Regarding the neural correlates of the deceptive effect, we found that frontal oscillatory dynamics was largely shaped by perceived time. We then showed that frontal dynamics predict the effectiveness of time deception on fatigue accumulation, suggesting its key role in the time deception effect.

Physical fatigue follows the perceived time

Our findings reveal that physical fatigue, as indexed by EMG activity increasing across contractions, follows the subjective rather than the objective time when the clock is slowed down, echoing previous work emphasizing the mind's influence on bodily processes (Craik & Sarbin, 1963; Crum et al., 2011; Crum & Langer, 2007; Langer et al., 2010; London & Monello, 1974; Pomares et al., 2011; Schachter & Gross, 1968; Snyder et al., 1974; Thönes et al., 2021), such as blood glucose level (Park et al., 2016) and healing (Aungle & Langer, 2023). While physical fatigue has been proposed to depend on subjective time (Matta et al., 2024), the current study is the first to accurately disentangle the motivational effects from the temporal deception. Furthermore, our results show an asymmetrical effect of clock manipulation, which might reflect mechanisms of efficiency and homeostatic regulation. Indeed, slowing perceived time may promote more economical effort allocation, delaying fatigue accumulation, and facilitating task completion (Meyniel et al., 2013). In contrast, accelerating perceived time does not produce a corresponding benefit, likely because fatigue functions as a protective signal that limits performance to prevent physiological harm and maintain homeostasis (Noakes, 2012; St Clair Gibson et al., 2003, 2013). As a result, even when perceived time is deceptively accelerated, compensatory mechanisms may be engaged to mitigate fatigue and preserve performance. Yet, complementary analyses (Fig. 5) nuance this effect, showing that some of the participants rely more on perceived time, both when the clock was slowed down and accelerated, while others follow objective time. Our findings suggest that interoceptive awareness may modulate this effect: lower interoceptive awareness is associated with a larger

deceptive effect, and vice versa. However, other factors likely influence this perceived-time

dependency, such as the strength of participants' beliefs (e.g., expectations that shorter times

lead to less fatigue) (Shiv & Carmon, 2005), the effectiveness of external clock manipulations

on the perception of time (Craik & Sarbin, 1963; Thönes et al., 2018), but also participants'

backgrounds or personality (Zhou et al., 2019). Further research is needed to better characterize

the factors driving time deception's effectiveness on fatigue.

Neural correlates of the time deception effect

In the current study, we found no differences between the normal sessions in the Cz

oscillatory dynamics, reflecting activity in motor areas, likely due to a relatively small workload

difference (Matta et al., 2025), making any potential difference due to the deceptive

manipulation even more challenging to observe. Yet, these parameters were carefully chosen to

ensure the implicit plausibility of the biased clock calibration, as increasing the manipulation's

magnitude could have raised awareness of the effect, potentially negating and confounding its

impact (Ducrocq et al., 2017; Williams et al., 2014).

Even though the task constraints imposed a relatively short temporal difference between

conditions, frontal oscillatory dynamics were significantly shaped by perceived time. Notably,

theta-band activity was modulated by perceived time in the slowed-clock condition, whereas

beta-band activity was sensitive to perceived time across both slowed and accelerated clocks.

Interestingly, both theta and beta-band activity increases have been previously associated with

prolonged engagement in demanding tasks (Boksem et al., 2005; Stoll et al., 2016). Specifically,

enhanced theta power has been extensively linked to increasing mental (Baumeister et al., 2012;

Borghini et al., 2014; Craig et al., 2012; Fan et al., 2015; Käthner et al., 2014; Liu et al., 2010;

Shigihara et al., 2013; Tanaka et al., 2014; Wascher et al., 2014) and, albeit less consistently,

physical (Bailey et al., 2008; Hottenrott & Gronwald, 2013; Mechau et al., 1998) fatigue – both

supposedly sharing common neural resources (Bray et al., 2012; Jin et al., 2024; Marcora et al.,

2009; Mehta & Parasuraman, 2014; Van Cutsem et al., 2017). This increase in theta power has

been proposed as reflecting a fatigue-related compensatory mechanism (Wascher et al., 2014)

to allow sustaining performance, seemingly driven by subjective rather than actual

performance. Moreover, given that frontal theta is a well-established marker of cognitive

control (Arnau et al., 2024; Cavanagh et al., 2012; Cavanagh & Frank, 2014; Ehrhardt et al.,

2022; Howells et al., 2010; Huster et al., 2013; Ishii et al., 1999; Lopez-Gamundi et al., 2024;

McFerren et al., 2021; Yu et al., 2022), our results suggest that the cognitive control required

to sustain a physical task depends primarily on perceived rather than actual time. Finally, several

studies also reported an increase in frontal beta with the rise in fatigue (Bailey et al., 2008;
Enders et al., 2016; Hottenrott & Gronwald, 2013; Kubitz & Mott, 1996; Mechau et al., 1998).
This increase might also reflect a top-down cognitive control (Buschman & Miller, 2007; Stoll
et al., 2016) allowing one to continue the task despite uncomfortable sensations, which might
follow perceived rather than actual time. However, our study did not include continuous
measurements of subjective experience, such as perceived effort (Pageaux, 2016), which limits
our ability to fully characterize these processes. Future studies should incorporate such
measures to better understand the relationship between perceived time and participants'
subjective experience of effort. Together, these results highlight the reliance of specific
cognitive control time-related correlates to perceived rather than actual time. Additionally, it
emphasizes the importance of considering subjective components in cognitive experimental
paradigms, as various subjective factors may significantly influence physiological measures.
487

A predictive coding perspective
We found that frontal oscillatory dynamics (supposedly reflecting PFC's activity) and
physical fatigue evolve according to perceived rather than actual time. The influence of frontal
areas and the PFC on physical fatigue may be explained through predictive coding (Kaptchuk,
2018), a framework suggesting that the brain continuously generates expectations about
incoming sensory information (Friston, 2010). When there is a mismatch between expected and
actual sensations – a prediction error – the brain can resolve this discrepancy either by updating
its expectations (priors) or by actively adjusting bodily states (active inference) (Friston et al.,
2009; Pezzulo et al., 2015; Seth, 2013). If we apply this framework to our findings, we may
consider beliefs about time as prior factors that regulate physical fatigue through an active
inference process. In this situation, the temporal deceptive manipulation induces an inaccurate
temporal estimate, which generates a prediction error in the participants about their current
fatigue level. In order to minimize such prediction error – the discrepancy between expected
and actual fatigue – the brain may modulate interoceptive input (i.e., interoceptive signals
related to physical fatigue, later influencing EMG signals), thereby adjusting the evolution of
physical fatigue rather than revising its priors (Pagnini et al., 2023). In line with this framework,
frontal oscillatory dynamics could reflect temporal priors, as the PFC has been proposed to play
a key role in this process (Geuter et al., 2017). Furthermore, previous research has demonstrated
that beliefs can influence physiological effector systems such as the autonomic nervous system
(Amigo et al., 1993; Meissner, 2009, 2011; Nakamura et al., 2012; Wager et al., 2009). In line
with this interpretation, one may speculate that the change in the physical fatigue course is

mediated by an indirect pathway that originates in the PFC and regulates peripheral organs (e.g.,

cardiac, respiratory) (Hassan et al., 2013; Thayer et al., 2009, 2012), impacting muscle fatigue

as an outcome to reduce the prediction error. Furthermore, our complementary analysis,

splitting the participants based on their PFC power difference between sessions with the same

subjective time, support the hypothesis that frontal areas modulate the time manipulation effect.

Indeed, as shown in Figure 5, only participants exhibiting strong deceptive effects in the PFC

also experienced the deceptive effect on fatigue, as revealed in the EMG recordings. While this

observation should be interpreted with caution, given the limited sample size and the

exploratory use of a median split, it may pinpoint the PFC as one of the core structures involved

in mediating the time deception effects on physical fatigue. Future studies with larger samples

and continuous, model-based approaches will be necessary to confirm this relationship and to

characterize individual differences in susceptibility to time-deception effects more robustly.

Conclusion

Our study demonstrates the direct influence of time deception on physical fatigue,

irrespective of motivational factors and performance goals, and reveals key neural correlates of

this effect. We show that this influence is primarily associated with frontal oscillatory dynamics

that closely track perceived time and are pivotal to the effectiveness of such deceptive time

manipulation. Building on these findings, we propose potential mechanisms underlying this

mind-body interaction. By showing that psychological factors can modulate physical fatigue,

our results have broad implications for performance enhancement, but also everyday life and

occupational settings, suggesting ways to optimize endurance, productivity, and safety in

workers and other physically demanding contexts. Future research should further investigate

the neural mechanisms driving this phenomenon from different perspectives (e.g., cardiac,

respiratory or fMRI recordings) to further deepen its theoretical understanding. It may also

explore applications across various contexts (e.g., sport, work, daily life) and their potential

translation into clinical conditions.

Methods

Participants

Twenty-eight healthy volunteers with no known history of neurological diseases or physical injuries and regular physical practice (once to several times a week) participated in this study. Out of the 28 individuals, 24 were able to complete the entire protocol and were kept on for the analysis (13 women, age: 24 ± 4, weight: 70 ± 19, mean ± SD). Before participating in the study, all individuals provided informed consent and were informed of their right to withdraw from the study at any time. The four sessions were justified to the participants on the grounds that they allowed (i) the study of fatigue evolution between two different levels of effort (short and long contractions) and (ii) each level to be repeated twice to assess the reproducibility of fatigue measurements at the same level of effort. No further details were provided. At the end of the study, participants completed a survey addressing various aspects of their experience during the sessions (e.g., whether some sessions felt more difficult than others, whether the load seemed consistent across the four sessions, whether recovery time felt the same between sessions, and whether the duration of effort was similar across sessions). No participant reported any specific remarks suggesting awareness of the temporal manipulation. Participants were only informed of the deception after the study was completed. The study procedure was approved by the French Ethical 'Comité de Protection des Personnes' (number: 2023-A00412-43).

Data Acquisition

During each session, all participants were seated on a Biodex isokinetic ergometer with their hips and knees bent to a 90° angle. Their right leg was securely fastened to the ergometer accessory at the ankle with a non-compliant strap just above the malleolus. Two shoulder and one abdominal harnesses guaranteed a reduction in their range of motion and the involvement of other muscles besides the quadriceps. The installation parameters were optimized for each subject and kept identical within subjects across sessions.

Torque and electromyography (EMG) were recorded at 2000 Hz using a Biopac MP150 system and AcqKnowledge software. EMG was recorded through three pairs of electrodes (AG-Cl, diameter: 11 mm, interelectrode distance: 2 cm) placed on the Vastus Medialis, Vastus Lateralis, and Rectus Femoris according to SENIAM recommendations. Before placement, the skin was carefully prepared through shaving and alcohol to ensure low impedance (<5 KΩ). The reference electrode was positioned on the patella of the contralateral knee. EEG was

recorded using a 64 electrodes Biosemi Active 2 system with 64 Ag-AgCl sintered active

electrodes at 2048 Hz. The EEG electrode locations followed the 10–20 international system.

The offset was kept below 20 mV using an active gel. All acquisitions were performed with the

light off to minimize electrical noise.

The main task was programmed in Python and displayed through a Windows computer

(Resolution: 1920 × 1080; Refresh rate: 60 Hz), using a NiDAQ acquisition card (NI USB-

6218, Sample rate: 2000 Hz) to read the torque data (low-pass filtered at 10 Hz) in real-time.

The same computer also generated the triggers used to mark the onset and offset of each

contraction towards BIOPAC and BIOSEMI, allowing us to synchronize all our signals offline.

Experimental Protocol

All participants came to the lab on four days (each spaced 3 to 10 days apart) to perform four

sessions. On each day, the participants performed the following steps (Fig. 1):

- Warm-up (consisting of 15-20 contractions with increasing intensity, approximately

from 20 to 80% of their estimated Maximal Voluntary Torque (MVT))

- 2 Maximal Voluntary Contractions (MVC) (3-4 s each, 30 s spaced)

- 1 MVC, 20 s of rest, 1 Long MVC (30 s), 30 s of rest, 3 minutes of EEG recording at

rest (eyes opened)

- 5 minutes of rest

- Main task: 100 contractions (see below)

- 1 MVC (5 s after the 100th contraction of the main task), 20 s of rest, 1 Long MVC (30

s), 30 s of rest, 3 minutes of EEG recording at rest (eyes opened)

Regarding the main task, participants performed 100 contractions at a fixed individualized

intensity, defined as 20% of their maximal force, or 40 N.m if their maximal force was larger

than 200 N.m. The intensity was the same among the four sessions. While the rest time between

contractions was identical for each session (5 s), each contraction's real (R) and perceived (P)

time were independently manipulated. In each session, the contractions' time was either short

(10s) or long (12s), while the clock displayed to the participants was also either short (10 s) or

long (12 s). This resulted in a 2 × 2 design with Perceived duration (10 s vs. 12 s) nested within

Real duration (10 s vs. 12 s). An accelerated clock was only possible at the shortest real duration,

whereas a slowed-down clock was only possible at the longest real duration. This yielded four

counterbalanced conditions: P10R10, P12R12, P10R12, and P12R10 (Fig. 1). In practice, a

sound indicated to the participants when to begin and end each contraction. Simultaneously

with the start sound, a rectangle was displayed in the middle of a black screen to provide online

feedback to participants on their actual force relative to the individualized target. The

rectangle's color was updated online based on the ongoing force: red indicated participants were

below the targeted force, blue above, and green was in the aimed range (-5% to +5% around

the targeted force). A digital clock was displayed in the middle of the colored rectangle 0.75 s

after the start of the contraction (preparation time left to the participants to reach the targeted

force). This clock allowed us to perform our manipulation, combining real time (R10 or R12)

and perceived time (P10 or P12). The clock was therefore either normal (P10R10 and P12R12),

slowed down (P10R12), or accelerated (P12R10) without the participants knowing it. After the

ending sound, the final perceived time (10.0 or 12.0 s) remained on the screen for 1 s, then the

number of contractions already performed was displayed during the resting time. Note that part

of the data (only from the two normal sessions, P10R10 and P12R12) has been previously

reported (Matta et al., 2025) in the context of a different question; Long MVC were specifically

used there.

Data pre-processing

All data pre-processing and analyses were conducted offline using custom MATLAB

(version 2022b) scripts along with EEGLAB functions (Delorme & Makeig, 2004). Torque and

EMG data were down-sampled offline to 1000 Hz. Torque data were then low-pass filtered at

10 Hz, while EMG data were band-pass filtered between 20 and 500 Hz and notch filtered

between 47 and 53 Hz.

EEG data were down-sampled offline to 250 Hz. A notch filter (47 to 53 Hz) and a high-pass

filter with a 1 Hz cut-off frequency were applied. Channels with excessive noise were

automatically rejected using the CleanRawData plugin, configured with a 5-second 'flat line

criterion,' a 0.6 'minimum channel correlation,' and a kurtosis threshold exceeding five times

the mean. On average, 61.7 ± 1.8 (mean $\pm$ SD) channels were retained for analysis. The

electrodes were then re-referenced using a common average reference. Independent Component

Analysis (ICA) was performed with the Infomax algorithm, and the resulting components were

labeled using the ICLabel plugin (Pion-Tonachini et al., 2019). Components labeled as muscle,

eye, heart, line noise, channel noise, or other with a probability above 90% were automatically

rejected, averaging 12.9 ± 5.5 components removed. Previously excluded channels were

subsequently interpolated.

EEG data from the two resting-state periods were segmented into 2-second non-overlapping

windows. For the main task, EEG and EMG data were segmented into 18-second epochs,

starting 3 seconds before the contraction onset (as marked by its trigger) and ending 15 seconds

after. A final epoch rejection procedure was applied to the main task EEG data to remove additional noise. Epochs with a standard deviation greater than 1.5 times the interquartile range (IQR) above the 3rd quartile or below 1.5 times the IQR under the 1st quartile were rejected for each subject and electrode. This procedure eliminated 3.9% of the epochs over the four sessions.

Data analysis

Force and EMG

Pre-Post. The highest torque from the three MVCs performed before the task was used to determine the pre-test Maximal Voluntary Torque (MVT). The post-test measurement was taken as the highest torque recorded during the post-task MVC. The variation rate was calculated as (post – pre)/pre ×100.

Main Task. EMG was quantified as the root-mean-square (RMS) during each contraction of the main task from 2 to 10 seconds after the onset of the contraction (Fig. 1B). This period was chosen to exclude the initial contraction phase, when torque stabilization has not yet occurred, and to ensure identical comparison times across all sessions. Within this period, EMG was quantified as the RMS over 180 ms non-overlapping windows, normalized by the RMS EMG associated with the previously selected maximal torque (centered from -90 ms to +90 ms around it) for the respective muscle and session. The three muscle EMG values were averaged over time to yield one EMG value per contraction in the main task.

EEG Spectral Power

Spectral power was obtained by applying a continuous wavelet transform to the pre-processed, epoched EEG data within the 1 to 50 Hz frequency range. The wavelet length was linearly increased from one cycle at 1 Hz to twenty cycles at 50 Hz, optimizing the tradeoff between temporal and frequency resolution across the entire frequency range. Contractions that were removed were subsequently linearly interpolated.

Spectral power was specifically extracted for the Theta (θ, 4-8 Hz) and an Individualized Beta band (Iβ), which was defined for each subject to account for inter-individual variability in the beta desynchronization-specific frequency band. Specifically, the frequency with the largest desynchronization was identified for each subject, and Iβ was defined as the range from -2 to +2 Hz around this frequency (Matta et al., 2025). Power from the lateral-frontal (F7-F8) and mid-frontal (F3-F4) electrodes was averaged to create a 'Frontal' cluster, chosen to reflect activity in the ventrolateral and dorsolateral prefrontal cortex (Okamoto et al., 2004; Robertson

& Marino, 2015). The Cz electrode was used to reflect motor cortex activity associated with quadriceps contraction, a location confirmed by topographically visualizing beta desynchronization (Dal Maso et al., 2012; Forrester et al., 2006; Gordon et al., 2023; Meier et al., 2008).

Pre-Post Analysis. Raw spectral powers from pre- and post-EEG resting state recordings were independently averaged over epochs (90) and time (0 to 2 s). The variation rate was calculated between the pre- and post-values for each frequency band and region of interest.

Main Task Analysis. For each contraction, raw spectral power bands during the period of interest (2 to 10 s) were normalized to the first contractions of the session, averaged over the same 2 to 10 s period (Fig. 1B). This normalization, focusing on the fatigue effect, involved subtracting the baseline and dividing by it, then multiplying by 100 to express power values as a percentage of a non-fatigued state. The baseline was defined as the average of contractions 2 to 5, excluding the first contraction due to its significant torque variability in all sessions. The resulting values were averaged over time and frequencies, yielding one power value per frequency band, contraction, and region of interest.

EEG-EMG Connectivity

To investigate the EMG-EEG relationships, we computed Pearson's correlation between EMG and Iβ Cz, θ Frontal, and Iβ Frontal, for each subject, respectively. Each variable was independently averaged in bins of 10 contractions. We then performed bilateral one-sample t-tests to determine whether the r values significantly differed from zero.

Statistical analyses

Dependent variables were analyzed using linear mixed-effects models (LMMs) to account for repeated measurements across participants. For variables measured longitudinally across 100 contractions, contraction number was included as a continuous predictor, z-scored within participants, and modeled either linearly or using a second-degree natural spline to capture potential non-linear trends. Real time and perceived time were included as fixed factors, and participants were modeled with a random intercept and an uncorrelated random slope for contraction number. Model comparisons were performed using likelihood-ratio tests to select the best degree of complexity for the contraction term (i.e., the degree of the natural splines function), yielding models of the form: *DependentVar ~ ns(NumContraction_z, df = d) * Perceived * Real + (1 + NumContraction_z || Participant)*, where *ns* represents the natural cubic spline function applied to the *NumContraction_z* factor, and d represents its degree, which

was set to either 1 or 2 according to the best model fit (note that d = 1 represents a linear model,

d > 1 a non-linear model). In the above formulation, the asterisks denote that the fixed terms

are considered with all interactions, while we consider the slope of the *NumContration_z* factor

for the random terms.

Our primary aim was to assess the rate of change (slopes) across the session. Specifically,

we were interested in the effect of perceived time at short and long real durations, and

conversely, the effect of real time at short and long perceived durations, yielding to four

contrasts of interest. To test these effects, we adopted a hierarchical approach to post-hoc

contrasts. When a three-way interaction between contraction number, perceived time, and real

time was significant, this indicated that the rate of change depended jointly on both perceived

and real time; in this case, post-hoc contrasts were conducted using estimated marginal trends

to extract slopes for the four contrasts of interest, directly addressing our primary hypothesis

concerning changes in rate. If only the two-way interaction between perceived and real time

was significant, indicating a modulation of overall levels rather than slope differences, post-hoc

contrasts were conducted on estimated marginal means for the corresponding four contrasts. If

neither interaction was significant, post-hoc contrasts were not reported. In all cases, contrasts

were adjusted for multiple comparisons using a Bonferroni correction.

For variables measured as pre-post differences, LMMs included perceived and real time as

fixed factors, with participants modeled as a random intercept, resulting in models of the form:

*DependentVar ~ Perceived * Real + (1 | Participant)*.

In the case of non-significant results in the post hoc contrasts, Bayes Factors (BFs) were also

computed to provide additional evidence regarding the presence or absence of an effect. In the

Bayesian framework, the BF represents the ratio of the likelihood of the data under the

alternative hypothesis to that under the null hypothesis. Conventionally, BFs > ~3 provide

evidence in favor of the alternative hypothesis, while BFs < ~0.3 indicate evidence in favor of

the null hypothesis (van Doorn et al., 2021).

Statistical analyses were conducted using R (CranR, version 4.3.1).

References

Amigo, I., Cuesta, V., Fernández, A., & González, A. (1993). The effect of verbal instructions on blood pressure measurement. *Journal of Hypertension*, *11*(3), 293-296. https://doi.org/10.1097/00004872-199303000-00010

Ansdell, P., Thomas, K., Howatson, G., Amann, M., & Goodall, S. (2018). Deception Improves Time Trial Performance in Well-trained Cyclists without Augmented Fatigue. *Medicine and Science in Sports and Exercise*, *50*(4), 809-816. https://doi.org/10.1249/MSS.0000000000001483

Arnau, S., Liegel, N., & Wascher, E. (2024). Frontal midline theta power during the cue-target-interval reflects increased cognitive effort in rewarded task-switching. *Cortex; a Journal Devoted to the Study of the Nervous System and Behavior*, *180*, 94-110. https://doi.org/10.1016/j.cortex.2024.08.004

Aungle, P., & Langer, E. (2023). Physical healing as a function of perceived time. *Scientific Reports*, *13*(1), 22432. https://doi.org/10.1038/s41598-023-50009-3

Bailey, S. P., Hall, E. E., Folger, S. E., & Miller, P. C. (2008). Changes in EEG during graded exercise on a recumbent cycle ergometer. *Journal of Sports Science & Medicine*, *7*(4), 505-511.

Baumeister, J., Reinecke, K., Schubert, M., Schade, J., & Weiss, M. (2012). Effects of induced fatigue on brain activity during sensorimotor control. *European Journal of Applied Physiology*, *112*(7), 2475-2482. https://doi.org/10.1007/s00421-011-2215-6

Behrens, M., Gube, M., Chaabene, H., Prieske, O., Zenon, A., Broscheid, K.-C., Schega, L., Husmann, F., & Weippert, M. (2023). Fatigue and Human Performance : An Updated Framework. *Sports Medicine (Auckland, N.Z.)*, *53*(1), 7-31. https://doi.org/10.1007/s40279-022-01748-2

Bigland-Ritchie, B., & Woods, J. J. (1984). Changes in muscle contractile properties and neural control during human muscular fatigue. *Muscle & Nerve*, *7*(9), 691-699. https://doi.org/10.1002/mus.880070902

Bigliassi, M., & Filho, E. (2022). Functional significance of the dorsolateral prefrontal cortex during exhaustive exercise. *Biological Psychology*, *175*, 108442. https://doi.org/10.1016/j.biopsycho.2022.108442

Boksem, M. A. S., Meijman, T. F., & Lorist, M. M. (2005). Effects of mental fatigue on attention : An ERP study. *Brain Research. Cognitive Brain Research*, *25*(1), 107-116. https://doi.org/10.1016/j.cogbrainres.2005.04.011

Borghini, G., Astolfi, L., Vecchiato, G., Mattia, D., & Babiloni, F. (2014). Measuring

neurophysiological signals in aircraft pilots and car drivers for the assessment of mental workload, fatigue and drowsiness. *Neuroscience and Biobehavioral Reviews*, *44*, 58-75. https://doi.org/10.1016/j.neubiorev.2012.10.003

Bray, S. R., Graham, J. D., Martin Ginis, K. A., & Hicks, A. L. (2012). Cognitive task performance causes impaired maximum force production in human hand flexor muscles. *Biological Psychology*, *89*(1), 195-200. https://doi.org/10.1016/j.biopsycho.2011.10.008

Brehm, J. W., & Self, E. A. (1989). The intensity of motivation. *Annual Review of Psychology*, *40*, 109-131. https://doi.org/10.1146/annurev.ps.40.020189.000545

Brown, P. (2000). Cortical drives to human muscle : The Piper and related rhythms. *Progress in Neurobiology*, *60*(1), 97-108. https://doi.org/10.1016/s0301-0082(99)00029-5

Buschman, T. J., & Miller, E. K. (2007). Top-down versus bottom-up control of attention in the prefrontal and posterior parietal cortices. *Science (New York, N.Y.)*, *315*(5820), 1860-1862. https://doi.org/10.1126/science.1138071

Cavanagh, J. F., Figueroa, C. M., Cohen, M. X., & Frank, M. J. (2012). Frontal theta reflects uncertainty and unexpectedness during exploration and exploitation. *Cerebral Cortex (New York, N.Y.: 1991)*, *22*(11), 2575-2586. https://doi.org/10.1093/cercor/bhr332

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control. *Trends in Cognitive Sciences*, *18*(8), 414-421. https://doi.org/10.1016/j.tics.2014.04.012

Chan, M. (2011). Fatigue : The most critical accident risk in oil and gas construction. *Construction Management & Economics*, *29*, 341-353. https://doi.org/10.1080/01446193.2010.545993

Craig, A., Tran, Y., Wijesuriya, N., & Nguyen, H. (2012). Regional brain wave activity changes associated with fatigue. *Psychophysiology*, *49*(4), 574-582. https://doi.org/10.1111/j.1469-8986.2011.01329.x

Craik, K. H., & Sarbin, T. R. (1963). Effect of covert alteratons of clock rate upon time estimations and personal tempo. *Perceptual and Motor Skills*, *16*, 597-610. https://doi.org/10.2466/pms.1963.16.2.597

Crum, A. J., Corbin, W. R., Brownell, K. D., & Salovey, P. (2011). Mind over milkshakes : Mindsets, not just nutrients, determine ghrelin response. *Health Psychology: Official Journal of the Division of Health Psychology, American Psychological Association*, *30*(4), 424-429; discussion 430-431. https://doi.org/10.1037/a0023467

Crum, A. J., & Langer, E. J. (2007). Mind-set matters : Exercise and the placebo effect. *Psychological Science*, *18*(2), 165-171. https://doi.org/10.1111/j.1467-9280.2007.01867.x

Cunningham, T. R., Guerin, R. J., Ferguson, J., & Cavallari, J. (2022). Work-related fatigue :

A hazard for workers experiencing disproportionate occupational risks. *American journal of*

industrial medicine, *65*(11), 913-925. https://doi.org/10.1002/ajim.23325

Dal Maso, F., Longcamp, M., & Amarantini, D. (2012). Training-related decrease in

antagonist muscles activation is associated with increased motor cortex activation : Evidence

of central mechanisms for control of antagonist muscles. *Experimental Brain Research*,

220(3-4), 287-295. https://doi.org/10.1007/s00221-012-3137-1

de Koning, J. J., Foster, C., Bakkum, A., Kloppenburg, S., Thiel, C., Joseph, T., Cohen, J.,

& Porcari, J. P. (2011). Regulation of pacing strategy during athletic competition. *PloS One*,

6(1), e15863. https://doi.org/10.1371/journal.pone.0015863

Delorme, A., & Makeig, S. (2004). EEGLAB : An open source toolbox for analysis of single-

trial EEG dynamics including independent component analysis. *Journal of Neuroscience*

Methods, *134*(1), 9-21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Denis, G. (2020). *Implication des dimensions neurocognitives dans le maintien de l'effort*

physique au travers du rôle endossé par le cortex préfrontal et de la perspective

coûts/bénéfices. Université Côte d'Azur.

Dijkerman, C., & Lenggenhager, B. (2018). The body and cognition : The relation between

body representations and higher level cognitive and social processes. *Cortex; a Journal*

Devoted to the Study of the Nervous System and Behavior, *104*, 133-139.

https://doi.org/10.1016/j.cortex.2018.06.001

Ducrocq, G. P., Hureau, T. J., Meste, O., & Blain, G. M. (2017). Increased Fatigue Response

to Augmented Deceptive Feedback during Cycling Time Trial. *Medicine and Science in Sports*

and Exercise, *49*(8), 1541-1551. https://doi.org/10.1249/MSS.0000000000001272

Edwards, R. H. (1981). Human muscle function and fatigue. *Ciba Foundation Symposium*,

82, 1-18. https://doi.org/10.1002/9780470715420.ch1

Ehrhardt, N. M., Fietz, J., Kopf-Beck, J., Kappelmann, N., & Brem, A.-K. (2022). Separating

EEG correlates of stress : Cognitive effort, time pressure, and social-evaluative threat. *The*

European Journal of Neuroscience, *55*(9-10), 2464-2473. https://doi.org/10.1111/ejn.15211

Enders, H., Cortese, F., Maurer, C., Baltich, J., Protzner, A. B., & Nigg, B. M. (2016).

Changes in cortical activity measured with EEG during a high-intensity cycling exercise.

Journal of Neurophysiology, *115*(1), 379-388. https://doi.org/10.1152/jn.00497.2015

Enoka, R. M., & Duchateau, J. (2008). Muscle fatigue : What, why and how it influences

muscle function. *The Journal of Physiology*, *586*(1), 11-23.

https://doi.org/10.1113/jphysiol.2007.139477

Enoka, R. M., & Duchateau, J. (2016). Translating Fatigue to Human Performance. *Medicine*

and Science in Sports and Exercise, *48*(11), 2228-2238. https://doi.org/10.1249/MSS.0000000000000929

Fan, X., Zhou, Q., Liu, Z., & Xie, F. (2015). Electroencephalogram assessment of mental fatigue in visual search. *Bio-Medical Materials and Engineering*, *26 Suppl 1*, S1455-1463. https://doi.org/10.3233/BME-151444

Forrester, L. W., Hanley, D. F., & Macko, R. F. (2006). Effects of treadmill exercise on transcranial magnetic stimulation-induced excitability to quadriceps after stroke. *Archives of Physical Medicine and Rehabilitation*, *87*(2), 229-234. https://doi.org/10.1016/j.apmr.2005.10.016

Friston, K. (2010). The free-energy principle : A unified brain theory? *Nature Reviews. Neuroscience*, *11*(2), 127-138. https://doi.org/10.1038/nrn2787

Friston, K. J., Daunizeau, J., & Kiebel, S. J. (2009). Reinforcement learning or active inference? *PloS One*, *4*(7), e6421. https://doi.org/10.1371/journal.pone.0006421

Gandevia, S. C. (2001). Spinal and supraspinal factors in human muscle fatigue. *Physiological Reviews*, *81*(4), 1725-1789. https://doi.org/10.1152/physrev.2001.81.4.1725

Geuter, S., Koban, L., & Wager, T. D. (2017). The Cognitive Neuroscience of Placebo Effects : Concepts, Predictions, and Physiology. *Annual Review of Neuroscience*, *40*, 167-188. https://doi.org/10.1146/annurev-neuro-072116-031132

Gordon, E. M., Chauvin, R. J., Van, A. N., Rajesh, A., Nielsen, A., Newbold, D. J., Lynch, C. J., Seider, N. A., Krimmel, S. R., Scheidter, K. M., Monk, J., Miller, R. L., Metoki, A., Montez, D. F., Zheng, A., Elbau, I., Madison, T., Nishino, T., Myers, M. J., … Dosenbach, N. U. F. (2023). A somato-cognitive action network alternates with effector regions in motor cortex. *Nature*, *617*(7960), 351-359. https://doi.org/10.1038/s41586-023-05964-2

Hassan, S. F., Cornish, J. L., & Goodchild, A. K. (2013). Respiratory, metabolic and cardiac functions are altered by disinhibition of subregions of the medial prefrontal cortex. *The Journal of Physiology*, *591*(23), 6069-6088. https://doi.org/10.1113/jphysiol.2013.262071

Hottenrott, K., & Gronwald, T. (2013). Cortical Brain Activity is Influenced by Cadence in Cyclists. *The Open Sports Sciences Journal*, *6*, 9-14. https://doi.org/10.2174/1875399X01306010009

Howells, F. M., Stein, D. J., & Russell, V. A. (2010). Perceived mental effort correlates with changes in tonic arousal during attentional tasks. *Behavioral and Brain Functions: BBF*, *6*, 39. https://doi.org/10.1186/1744-9081-6-39

Huster, R. J., Enriquez-Geppert, S., Lavallee, C. F., Falkenstein, M., & Herrmann, C. S. (2013). Electroencephalography of response inhibition tasks : Functional networks and

cognitive contributions. *International Journal of Psychophysiology: Official Journal of the*
International Organization of Psychophysiology, *87*(3), 217-233.
https://doi.org/10.1016/j.ijpsycho.2012.08.001

Ibrahim, A., Nnaji, C., Namian, M., Koh, A., & Techera, U. (2023). Investigating the impact
of physical fatigue on construction workers' situational awareness. *Safety Science*, *163*, 106103.
https://doi.org/10.1016/j.ssci.2023.106103

Iodice, P., Porciello, G., Bufalari, I., Barca, L., & Pezzulo, G. (2019). An interoceptive
illusion of effort induced by false heart-rate feedback. *Proceedings of the National Academy of*
Sciences of the United States of America, *116*(28), 13897-13902.
https://doi.org/10.1073/pnas.1821032116

Ishii, R., Shinosaki, K., Ukai, S., Inouye, T., Ishihara, T., Yoshimine, T., Hirabuki, N.,
Asada, H., Kihara, T., Robinson, S. E., & Takeda, M. (1999). Medial prefrontal cortex generates
frontal midline theta rhythm. *Neuroreport*, *10*(4), 675-679. https://doi.org/10.1097/00001756-
199903170-00003

Jin, H., Xiao, M., Liu, L., Kan, S., Fu, Y., & Zhang, D. (2024). Relationship between
physical fatigue and mental fatigue based on multimodal measurement under different load
levels. *Ergonomics*, *67*(11), 1748-1763. https://doi.org/10.1080/00140139.2024.2364667

Jones, C. M., Griffiths, P. C., & Mellalieu, S. D. (2017). Training Load and Fatigue Marker
Associations with Injury and Illness : A Systematic Review of Longitudinal Studies. *Sports*
Medicine (Auckland, N.z.), *47*(5), 943-974. https://doi.org/10.1007/s40279-016-0619-5

Jones, H. S., Williams, E. L., Bridge, C. A., Marchant, D., Midgley, A. W., Micklewright,
D., & Mc Naughton, L. R. (2013). Physiological and psychological effects of deception on
pacing strategy and performance : A review. *Sports Medicine (Auckland, N.Z.)*, *43*(12),
1243-1257. https://doi.org/10.1007/s40279-013-0094-1

Kaptchuk, T. J. (2018). Open-Label Placebo : Reflections on a Research Agenda.
Perspectives in Biology and Medicine, *61*(3), 311-334. https://doi.org/10.1353/pbm.2018.0045

Käthner, I., Wriessnegger, S. C., Müller-Putz, G. R., Kübler, A., & Halder, S. (2014). Effects
of mental workload and fatigue on the P300, alpha and theta band power during operation of
an ERP (P300) brain-computer interface. *Biological Psychology*, *102*, 118-129.
https://doi.org/10.1016/j.biopsycho.2014.07.014

Kim, G., Ahad, M. A., Ferdjallah, M., & Harris, G. F. (2007). Correlation of muscle fatigue
indices between intramuscular and surface EMG signals. *Proceedings 2007 IEEE*
SoutheastCon, 378-382. https://doi.org/10.1109/SECON.2007.342928

Kreher, J. B., & Schwartz, J. B. (2012). Overtraining Syndrome. *Sports Health*, *4*(2),

128-138. https://doi.org/10.1177/1941738111434406

Kubitz, K. A., & Mott, A. A. (1996). EEG power spectral densities during and after cycle ergometer exercise. *Research Quarterly for Exercise and Sport*, *67*(1), 91-96. https://doi.org/10.1080/02701367.1996.10607929

Langer, E., Djikic, M., Pirson, M., Madenci, A., & Donohue, R. (2010). Believing is seeing : Using mindlessness (mindfully) to improve visual acuity. *Psychological Science*, *21*(5), 661-666. https://doi.org/10.1177/0956797610366543

Lebesque, L., Scaglioni, G., & Martin, A. (2022). The impact of submaximal fatiguing exercises on the ability to generate and sustain the maximal voluntary contraction. *Frontiers in Physiology*, *13*, 970917. https://doi.org/10.3389/fphys.2022.970917

Liu, J.-P., Zhang, C., & Zheng, C.-X. (2010). Estimation of the cortical functional connectivity by directed transfer function during mental fatigue. *Applied Ergonomics*, *42*(1), 114-121. https://doi.org/10.1016/j.apergo.2010.05.008

London, H., & Monello, L. (1974). Cognitive manipulation of boredom. In *Thought and feeling : Cognitive alteration of feeling states.* (p. 239-239). Aldine.

Lopez-Gamundi, P., Mas-Herrero, E., & Marco-Pallares, J. (2024). Disentangling effort from probability of success : Temporal dynamics of frontal midline theta in effort-based reward processing. *Cortex; a Journal Devoted to the Study of the Nervous System and Behavior*, *176*, 94-112. https://doi.org/10.1016/j.cortex.2024.03.014

Marcora, S. M., Staiano, W., & Manning, V. (2009). Mental fatigue impairs physical performance in humans. *Journal of Applied Physiology (Bethesda, Md.: 1985)*, *106*(3), 857-864. https://doi.org/10.1152/japplphysiol.91324.2008

Matta, P.-M., Baurès, R., Duclay, J., & Alamia, A. (2025). Modulation of beta oscillatory dynamics in motor and frontal areas during physical fatigue. *Communications Biology*, *8*(1), 687. https://doi.org/10.1038/s42003-025-08122-8

Matta, P.-M., Glories, D., Alamia, A., Baurès, R., & Duclay, J. (2024). Mind over muscle ? Time manipulation improves physical performance by slowing down the neuromuscular fatigue accumulation. *Psychophysiology*, *61*(4), e14487. https://doi.org/10.1111/psyp.14487

McFerren, A., Riddle, J., Walker, C., Buse, J. B., & Frohlich, F. (2021). Causal role of frontal-midline theta in cognitive effort : A pilot study. *Journal of Neurophysiology*, *126*(4), 1221-1233. https://doi.org/10.1152/jn.00068.2021

McMorris, T., Barwood, M., & Corbett, J. (2018). Central fatigue theory and endurance exercise : Toward an interoceptive model. *Neuroscience and Biobehavioral Reviews*, *93*, 93-107. https://doi.org/10.1016/j.neubiorev.2018.03.024

Mechau, D., Mücke, S., Weiss, M., & Liesen, H. (1998). Effect of increasing running
velocity on electroencephalogram in a field test. *European Journal of Applied Physiology and*
Occupational Physiology, *78*(4), 340-345. https://doi.org/10.1007/s004210050429

Mehling, W. E., Price, C., Daubenmier, J. J., Acree, M., Bartmess, E., & Stewart, A. (2012).
The Multidimensional Assessment of Interoceptive Awareness (MAIA). *PLoS ONE*, *7*(11),
e48230. https://doi.org/10.1371/journal.pone.0048230

Mehta, R. K., & Parasuraman, R. (2014). Effects of mental fatigue on the development of
physical fatigue : A neuroergonomic approach. *Human Factors*, *56*(4), 645-656.
https://doi.org/10.1177/0018720813507279

Meier, J. D., Aflalo, T. N., Kastner, S., & Graziano, M. S. A. (2008). Complex organization
of human primary motor cortex : A high-resolution fMRI study. *Journal of Neurophysiology*,
100(4), 1800-1812. https://doi.org/10.1152/jn.90531.2008

Meissner, K. (2009). Effects of placebo interventions on gastric motility and general
autonomic activity. *Journal of Psychosomatic Research*, *66*(5), 391-398.
https://doi.org/10.1016/j.jpsychores.2008.09.004

Meissner, K. (2011). The placebo effect and the autonomic nervous system : Evidence for
an intimate relationship. *Philosophical Transactions of the Royal Society of London. Series B,*
Biological Sciences, *366*(1572), 1808-1817. https://doi.org/10.1098/rstb.2010.0403

Merletti, R., Lo Conte, L. R., & Orizio, C. (1991). Indices of muscle fatigue. *Journal of*
Electromyography and Kinesiology: Official Journal of the International Society of
Electrophysiological Kinesiology, *1*(1), 20-33. https://doi.org/10.1016/1050-6411(91)90023-X

Meyniel, F., Sergent, C., Rigoux, L., Daunizeau, J., & Pessiglione, M. (2013).
Neurocomputational account of how the human brain decides when to have a break.
Proceedings of the National Academy of Sciences of the United States of America, *110*(7),
2641-2646. https://doi.org/10.1073/pnas.1211925110

Nakamura, Y., Donaldson, G. W., Kuhn, R., Bradshaw, D. H., Jacobson, R. C., & Chapman,
R. C. (2012). Investigating dose-dependent effects of placebo analgesia : A
psychophysiological approach. *Pain*, *153*(1), 227-237.
https://doi.org/10.1016/j.pain.2011.10.024

Noakes, T. D. (2011). Time to move beyond a brainless exercise physiology : The evidence
for complex regulation of human exercise performance. *Applied Physiology, Nutrition, and*
Metabolism, *36*(1), 23-35. https://doi.org/10.1139/H10-082

Noakes, T. D. (2012). Fatigue is a Brain-Derived Emotion that Regulates the Exercise
Behavior to Ensure the Protection of Whole Body Homeostasis. *Frontiers in Physiology*, *3*, 82.

https://doi.org/10.3389/fphys.2012.00082

Okamoto, M., Dan, H., Sakamoto, K., Takeo, K., Shimizu, K., Kohno, S., Oda, I., Isobe, S., Suzuki, T., Kohyama, K., & Dan, I. (2004). Three-dimensional probabilistic anatomical cranio-cerebral correlation via the international 10-20 system oriented for transcranial functional brain mapping. *NeuroImage*, *21*(1), 99-111. https://doi.org/10.1016/j.neuroimage.2003.08.026

Pageaux, B. (2016). Perception of effort in Exercise Science : Definition, measurement and perspectives. *European Journal of Sport Science*, *16*(8), 885-894. https://doi.org/10.1080/17461391.2016.1188992

Pagnini, F., Barbiani, D., Cavalera, C., Volpato, E., Grosso, F., Minazzi, G. A., Vailati Riboni, F., Graziano, F., Di Tella, S., Manzoni, G. M., Silveri, M. C., Riva, G., & Phillips, D. (2023). Placebo and Nocebo Effects as Bayesian-Brain Phenomena : The Overlooked Role of Likelihood and Attention. *Perspectives on Psychological Science: A Journal of the Association for Psychological Science*, *18*(5), 1217-1229. https://doi.org/10.1177/17456916221141383

Park, C., Pagnini, F., Reece, A., Phillips, D., & Langer, E. (2016). Blood sugar level follows perceived time rather than actual time in people with type 2 diabetes. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(29), 8168-8170. https://doi.org/10.1073/pnas.1603444113

Pearson, K. (2000). Motor systems. *Current Opinion in Neurobiology*, *10*(5), 649-654. https://doi.org/10.1016/s0959-4388(00)00130-6

Pezzulo, G., Rigoli, F., & Friston, K. (2015). Active Inference, homeostatic regulation and adaptive behavioural control. *Progress in Neurobiology*, *134*, 17-35. https://doi.org/10.1016/j.pneurobio.2015.09.001

Pion-Tonachini, L., Kreutz-Delgado, K., & Makeig, S. (2019). ICLabel : An automated electroencephalographic independent component classifier, dataset, and website. *NeuroImage*, *198*, 181-197. https://doi.org/10.1016/j.neuroimage.2019.05.026

Pomares, F. B., Creac'h, C., Faillenot, I., Convers, P., & Peyron, R. (2011). How a clock can change your pain? The illusion of duration and pain perception. *Pain*, *152*(1), 230-234. https://doi.org/10.1016/j.pain.2010.10.047

Robertson, C. V., & Marino, F. E. (2015). Prefrontal and motor cortex EEG responses and their relationship to ventilatory thresholds during exhaustive incremental exercise. *European Journal of Applied Physiology*, *115*(9), 1939-1948. https://doi.org/10.1007/s00421-015-3177-x

Robertson, C. V., & Marino, F. E. (2016). A role for the prefrontal cortex in exercise tolerance and termination. *Journal of Applied Physiology (Bethesda, Md.: 1985)*, *120*(4),

464-466. https://doi.org/10.1152/japplphysiol.00363.2015

Rodrigues, F., Monteiro, D., Ferraz, R., Branquinho, L., & Forte, P. (2023). The Association between Training Frequency, Symptoms of Overtraining and Injuries in Young Men Soccer Players. *International Journal of Environmental Research and Public Health*, *20*(8), 5466. https://doi.org/10.3390/ijerph20085466

Schachter, S., & Gross, L. P. (1968). Manipulated time and eating behavior. *Journal of Personality and Social Psychology*, *10*(2), 98-106. https://doi.org/10.1037/h0026285

Schad, D. J., Vasishth, S., Hohenstein, S., & Kliegl, R. (2020). How to capitalize on a priori contrasts in linear (mixed) models : A tutorial. *Journal of Memory and Language*, *110*, 104038. https://doi.org/10.1016/j.jml.2019.104038

Seth, A. K. (2013). Interoceptive inference, emotion, and the embodied self. *Trends in Cognitive Sciences*, *17*(11), 565-573. https://doi.org/10.1016/j.tics.2013.09.007

Shigihara, Y., Tanaka, M., Ishii, A., Kanai, E., Funakura, M., & Watanabe, Y. (2013). Two types of mental fatigue affect spontaneous oscillatory brain activities in different ways. *Behavioral and Brain Functions: BBF*, *9*, 2. https://doi.org/10.1186/1744-9081-9-2

Shiv, B., & Carmon, Z. (2005). Placebo Effects of Marketing Actions : Consumers May Get What They Pay For. *Journal of Marketing Research*, *42*, 383-393. https://doi.org/10.2139/ssrn.707541

Snyder, M., Schulz, R., & Jones, E. E. (1974). Expectancy and apparent duration as determinants of fatigue. *Journal of Personality and Social Psychology*, *29*(3), 426-434. https://doi.org/10.1037/h0035917

St Clair Gibson, A., Baden, D. A., Lambert, M. I., Lambert, E. V., Harley, Y. X. R., Hampson, D., Russell, V. A., & Noakes, T. D. (2003). The conscious perception of the sensation of fatigue. *Sports Medicine (Auckland, N.Z.)*, *33*(3), 167-176. https://doi.org/10.2165/00007256-200333030-00001

St Clair Gibson, A., De Koning, J. J., Thompson, K. G., Roberts, W. O., Micklewright, D., Raglin, J., & Foster, C. (2013). Crawling to the finish line : Why do endurance runners collapse? Implications for understanding of mechanisms underlying pacing and fatigue. *Sports Medicine (Auckland, N.Z.)*, *43*(6), 413-424. https://doi.org/10.1007/s40279-013-0044-y

Stoll, F. M., Wilson, C. R. E., Faraut, M. C. M., Vezoli, J., Knoblauch, K., & Procyk, E. (2016). The Effects of Cognitive Control and Time on Frontal Beta Oscillations. *Cerebral Cortex (New York, N.Y.: 1991)*, *26*(4), 1715-1732. https://doi.org/10.1093/cercor/bhv006

Tanaka, M., Ishii, A., & Watanabe, Y. (2014). Neural effects of mental fatigue caused by continuous attention load : A magnetoencephalography study. *Brain Research*, *1561*, 60-66.

https://doi.org/10.1016/j.brainres.2014.03.009

Thayer, J. F., Ahs, F., Fredrikson, M., Sollers, J. J., & Wager, T. D. (2012). A meta-analysis
of heart rate variability and neuroimaging studies : Implications for heart rate variability as a
marker of stress and health. *Neuroscience and Biobehavioral Reviews*, *36*(2), 747-756.
https://doi.org/10.1016/j.neubiorev.2011.11.009

Thayer, J. F., Hansen, A. L., Saus-Rose, E., & Johnsen, B. H. (2009). Heart rate variability,
prefrontal neural function, and cognitive performance : The neurovisceral integration
perspective on self-regulation, adaptation, and health. *Annals of Behavioral Medicine: A*
Publication of the Society of Behavioral Medicine, *37*(2), 141-153.
https://doi.org/10.1007/s12160-009-9101-z

Thönes, S., Arnau, S., & Wascher, E. (2018). Cognitions about time affect perception,
behavior, and physiology—A review on effects of external clock-speed manipulations.
Consciousness and Cognition, *63*, 99-109. https://doi.org/10.1016/j.concog.2018.06.014

Thönes, S., Arnau, S., Wascher, E., & Schneider, D. (2021). Boosting working memory with
accelerated clocks. *NeuroImage*, *226*, 117601.
https://doi.org/10.1016/j.neuroimage.2020.117601

Tucker, R. (2009). The anticipatory regulation of performance : The physiological basis for
pacing strategies and the development of a perception-based model for exercise performance.
British Journal of Sports Medicine, *43*(6), 392-400. https://doi.org/10.1136/bjsm.2008.050799

Van Cutsem, J., Marcora, S., De Pauw, K., Bailey, S., Meeusen, R., & Roelands, B. (2017).
The Effects of Mental Fatigue on Physical Performance : A Systematic Review. *Sports*
Medicine (Auckland, N.Z.), *47*(8), 1569-1588. https://doi.org/10.1007/s40279-016-0672-0

van Doorn, J., van den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., Etz, A.,
Evans, N. J., Gronau, Q. F., Haaf, J. M., Hinne, M., Kucharský, Š., Ly, A., Marsman, M.,
Matzke, D., Gupta, A. R. K. N., Sarafoglou, A., Stefan, A., Voelkel, J. G., & Wagenmakers,
E.-J. (2021). The JASP guidelines for conducting and reporting a Bayesian analysis.
Psychonomic Bulletin & Review, *28*(3), 813-826. https://doi.org/10.3758/s13423-020-01798-5

Wager, T. D., Waugh, C. E., Lindquist, M., Noll, D. C., Fredrickson, B. L., & Taylor, S. F.
(2009). Brain mediators of cardiovascular responses to social threat : Part I: Reciprocal dorsal
and ventral sub-regions of the medial prefrontal cortex and heart-rate reactivity. *NeuroImage*,
47(3), 821-835. https://doi.org/10.1016/j.neuroimage.2009.05.043

Wascher, E., Rasch, B., Sänger, J., Hoffmann, S., Schneider, D., Rinkenauer, G., Heuer, H.,
& Gutberlet, I. (2014). Frontal theta activity reflects distinct aspects of mental fatigue.
Biological Psychology, *96*, 57-65. https://doi.org/10.1016/j.biopsycho.2013.11.010

Williams, E. L., Jones, H. S., Sparks, S., Marchant, D. C., Micklewright, D., & McNaughton,
L. R. (2014). Deception studies manipulating centrally acting performance modifiers : A
review. *Medicine and Science in Sports and Exercise*, *46*(7), 1441-1451.
https://doi.org/10.1249/MSS.0000000000000235

Yu, S., Mückschel, M., & Beste, C. (2022). Superior frontal regions reflect the dynamics of
task engagement and theta band-related control processes in time-on task effects. *Scientific*
Reports, *12*(1), 846. https://doi.org/10.1038/s41598-022-04972-y

Zhou, L., Wei, H., Zhang, H., Li, X., Bo, C., Wan, L., Lu, X., & Hu, L. (2019). The Influence
of Expectancy Level and Personal Characteristics on Placebo Effects : Psychological
Underpinnings. *Frontiers in Psychiatry*, *10*, 20. https://doi.org/10.3389/fpsyt.2019.00020

1082