Perceived time drives physical fatigue

Pierre-Marie Matta[1,2,*], Robin Baurès[1], Julien Duclay[2], Andrea Alamia[1]

1Univ Toulouse, CNRS, CerCo, Toulouse, France

2Univ Toulouse, Inserm, ToNIC, Toulouse, France

*Corresponding author: Pierre-Marie Matta (pierre-marie.matta@inserm.fr)

Conflict of interest: The authors declare that they have no competing interests.

Keywords: Time deception, False-clock paradigm, Frontal oscillations, Fatigue, Mind-body connection

Abstract

Recent studies suggested that fundamental physiological processes, such as physical fatigue, rely on the perceived rather than the actual time. However, the neural correlates underlying this effect and its disentanglement from motivational factors (i.e., performance goals) remain unknown. To investigate the time deception effect on fatigue, we developed a novel EEG design in which participants performed 100 isometric contractions at a fixed pace and resistance per session. The actual contraction duration (short or long) and the calibration of the displayed clock (normal or biased to an acceleration or deceleration) were independently manipulated between sessions to examine whether fatigue and its neural correlates evolved based on perceived or actual time. Our results show an accumulation of physical fatigue that follows the perceived time, irrespective of motivational factors. This effect was consistently observed only when the clock was slowed down. This time deception effect was mediated by an oscillatory dynamic that followed perceived time in frontal (theta- and beta-bands) but not motor areas (beta-band). Further analyses highlighted the key role of the frontal oscillatory dynamics in the effectiveness of the time deception effect on physical fatigue.

Significance statement

Can a clock change the course of physical fatigue? In this study, we addressed this question while controlling for motivational confounds and monitoring EEG-associated power modulations. Our findings demonstrate a slowed-down fatigue accumulation in the presence of a slowed-down clock. This time manipulation effect was driven by a frontal oscillatory dynamic that thoroughly followed the perceived time. These results highlight the direct influence of psychological factors on physiological processes and unveil the neural correlates underlying this effect.

Introduction

The relationship between the mind and body has been a subject of inquiry for centuries (Dijkerman & Lenggenhager, 2018), raising fundamental questions about how mental states influence bodily processes. A prevalent approach to exploring this relationship involves deceptively manipulating participants' beliefs while recording psychophysiological variables (Crum et al., 2011; Crum & Langer, 2007; H. S. Jones et al., 2013; Williams et al., 2014). Among the various psychological parameters that can be successfully deceived, manipulating the subjective passage of time has been suggested as particularly relevant because of the trust we routinely place in its measurement (Thönes et al., 2018). This manipulation is typically achieved by covertly altering the clock's speed displayed to participants during a task, either slowing it down or accelerating it (Thönes et al., 2018). For instance, when the passage of time is artificially slowed down, studies have reported a reduced perceived pain in response to thermal stimulation (Pomares et al., 2011) but also a smaller decrease in blood glucose level (Park et al., 2016) and slower healing (Aungle & Langer, 2023). Similarly, studies interested in physical activity have shown that, in the presence of a slowed-down clock, there is a slower spontaneous motor activity (Craik & Sarbin, 1963) and a slower accumulation of physical fatigue (Matta et al., 2024). Conversely, accelerating the perceived passage of time has been associated with opposite effects, with the notable exception of physical fatigue, which remains, to our knowledge, unexplored in this context.

Physical fatigue, broadly defined as the decline in the ability to generate force or sustain performance (Bigland-Ritchie & Woods, 1984; Gandevia, 2001), is a central determinant of endurance, productivity, and daily functioning (Enoka & Duchateau, 2016). Beyond its role in limiting athletic and occupational performance, it is also closely linked to injury risk (Chan, 2011; Cunningham et al., 2022; Ibrahim et al., 2023; C. M. Jones et al., 2017; Kreher & Schwartz, 2012; Rodrigues et al., 2023) and impaired quality of life (Behrens et al., 2023). Understanding how it may be shaped by psychological factors is therefore of both theoretical and practical significance, with implications for athletes, workers in physically demanding professions, and individuals in everyday life. This makes physical fatigue a particularly relevant phenomenon through which to investigate deceptive time manipulations.

Nevertheless, important gaps remain in our understanding of how time perception influences physical fatigue: although these perceived time-dependent effects are being demonstrated for several variables (Thönes et al., 2018), their underlying neural mechanisms remain poorly understood. Moreover, motivational factors, such as performance goals, are critical confounds

often unaccounted for when measuring deceptive effects. These confounds are particularly important in physical fatigue, in which previous studies have consistently employed deceptive manipulations to enhance physical performance (i.e., the amount of time a task to exhaustion is maintained or the time needed to complete a given distance) (Ansdell et al., 2018; Ducrocq et al., 2017; Matta et al., 2024). Consequently, previous results on physical fatigue may be alternatively explained by performance-related goals rather than time perception.

In the current study, we investigate the neural correlates associated with the time deception effect on the accumulation of physical fatigue, and we carefully isolate this effect from motivational confounds. In addition, we also test its temporal symmetry, that is, when the clock is either slowed down or accelerated. To this end, we used an original 2×2 design, where each session consisted of performing 100 isometric contractions at a given pace against a fixed individualized resistance. The contraction's actual duration (short or long) and the calibration of the displayed clock (normal or biased toward acceleration or deceleration) were manipulated between sessions to test whether physical fatigue was evolving according to the perceived (i.e., displayed) or actual time (Fig 1A). Unlike previous studies, we tested all four possible combinations for the first time in a within-subject design. We recorded EMG and EEG signals to assess fatigue accumulation and associated neural correlates. We specifically focused on oscillatory dynamics at the sensor level in the motor and prefrontal cortex, as both regions are involved in physical fatigue and effort maintenance (Bigliassi & Filho, 2022; Matta et al., 2025; McMorris et al., 2018; Robertson & Marino, 2016).

Figure 1 – Experimental protocol and EMG/EEG computation

A. Experimental Protocol. All participants performed 100 contractions during four separate sessions. While the rest time between contractions was identical (5s) for each session, each contraction's real (R) and perceived (P) time were manipulated. In each session, the contractions' time was either short (10s) or long (12s), while the digital clock displayed in front of the participants was either Normal (N) or Biased (B). Before and after the task, Maximal Voluntary Contractions (MVC) were performed to assess the fatigue generated by the 100 contractions. Resting State (RS) EEG (3 min) was recorded before and after the task to quantify the associated EEG correlates. B. EMG/EEG Computation. All panels represent the typical trace of one contraction except the upper right panel, which illustrates the whole task. EMG and EEG were recorded throughout the task and explicitly analyzed over a 2 to 10 s window within each contraction (0 s being the contraction onset), from which we extracted one average value for each contraction. The EMG was computed in normalized Root Mean Square (RMS), while we computed the mean power of the theta (θ) and an individualized beta-band (Iβ) for the EEG, in frontal electrodes and Cz (illustrated in black and yellow respectively on the topographical plots) (see Methods for details).

Results

Design Summary

Participants performed four sessions (each spaced 3 to 10 days apart) of 100 quadriceps isometric contractions, during which EMG and EEG were continuously recorded. For each participant, the contraction intensity was identical across sessions ($\leq 20\%$ of the participant's maximal force; see Methods for details). While the rest time between contractions was identical for each session, each contraction's real (R) and perceived (P) time were independently manipulated. In each session, the contractions' time was either short (10 s) or long (12 s), while the digital clock displayed in front of the participants was either Normal (N) or Biased (B), leading to 4 counterbalanced conditions: N10 (10s P, 10s R), N12 (12s P, 12s R), B10 (10s P, 12s R), B12 (12s P, 10s R) (Fig 1). Before and after the task, participants performed Maximal Voluntary Contractions (MVC) and 3-minute EEG resting states.

Neuromuscular fatigue

We first compared the physical fatigue generated in the two sessions with unbiased clocks (i.e., N12 and N10 sessions) to confirm that a longer effort generated a measurable difference in neuromuscular fatigue. To this end, linear mixed-effects models were fitted on the pre-post variation rate of maximal voluntary torque, with Condition as a fixed factor and Participant as a random intercept (see Methods for details). As expected, we found a significantly negative N12-N10 contrast ($t_{(69)} = 2.26$, $p = .013$, $d_z = 0.38$, $BF = 1.75$), indicating greater physical fatigue in the N12 session compared to the N10 session (Fig 2A, 2C). We then applied the same analysis to the B10-N10 and N12-B12 contrasts, thus comparing the sessions with the same subjective time but different real times. Remarkably, we found no difference between the B10 and N10 sessions ($t_{(69)} = 0.71$, $p = .48$, $d_z = 0.14$, $BF = 0.26$), suggesting similar levels of fatigue between the two, despite the B10 session having a longer amount of actual work, identical to the N12 one. However, this subjective time-dependent relationship was non-symmetric, as evidenced by a significantly negative contrast between the N12 session and the one perceived as long but identical to the short session, namely B12 ($t_{(69)} = 2.38$, $p = .020$, $d_z = 0.58$, $BF = 4.99$). This comparison indicates more fatigue in the N12 session (Fig 2A, 2C), consistent with its higher actual workload.

Figure 2 – Physical Fatigue

A. Torque Pre-Post Variation Rate (VR). A decrease in Maximal Voluntary Torque (MVT) reflects a decline in the

maximal force production. B. EMG modulations over the main task. EMG is expressed in normalized Root Mean

Square (RMS) over contractions, whose increase reflects neuromuscular fatigue accumulation. The dots represent

the averaged distribution, and the error bars are the standard error of the mean (SEM). C. Fatigue differences

computed for each pair of interest (N12-N10; B10-N10; N12-B12). The upper panel represents the difference for

the Pre-Post VR of the MVT, while the lower one contains the difference for the sum of the EMG-RMS over the

bins. The bar plots represent the averaged distribution; the error bars are the SEM. Asterisks denote a significant

effect with p<0.05. The Bayes Factor (BF), rounded to the decimal, was reported when there was no significant

difference.

We then investigated whether the biased clock influenced physical fatigue during the task,

as observed in the pre-post measurements. To this end, we considered the sum of the EMG Root

Mean Square (RMS) activity recorded during each session (Kim et al., 2007; Merletti et al.,

1991). As in the pre-post case, we found a significantly positive contrast between the N12 and

N10 sessions ($t_{(69)} = 2.23$, $p = .015$, $d_z = 0.47$, BF = 3.64), supporting a greater fatigue

accumulation in the longer, normal session. Concerning the comparison between the sessions

with the same perceived time but a different real one, we found no differences between the

subjectively short ones (B10-N10: $t_{(69)} = 0.93$, $p = .35$, $d_z = 0.26$, BF = 0.43), as well as between

the subjectively long ones (N12-B12: $t_{(69)} = 1.33$, $p = .19$, $d_z = 0.29$, BF = 0.53) (Fig 2B, 2C).

Together, these results indicate that physical fatigue follows perceived rather than actual time

when the clock is slowed down (Matta et al., 2024), whereas the results appear less compelling

when the clock is accelerated.

Power modulations in motor areas

After demonstrating that physical fatigue depends on perceived time, consistently when the clock is slowed down but less clearly when it is accelerated, we investigated the neural correlates underlying this effect. We first examined oscillatory activity in the Cz electrode, commonly used as an approximation of activity in the motor cortex (Dal Maso et al., 2012; Forrester et al., 2006; Gordon et al., 2023; Meier et al., 2008), a key region involved in executing voluntary contractions (Brown, 2000; Pearson, 2000). Focusing on this sensor, we assessed relative spectral power changes in each participant's individualized beta-band (Iβ) (Matta et al., 2025) (i.e., a 4 Hz spectral window in the beta band with the largest desynchronization, see Methods for details). Importantly, we found an absence of difference between the N12 and N10 sessions ($t_{(69)} = 0.29$, $p = .39$, $d_z = 0.094$, BF = 0.32), possibly due to the relatively small discrepancy between the two sessions' workload (Matta et al., 2025). Similarly, we found no difference between all pairs of sessions with the same subjective time ($p > .54$, $d < 0.093$, BF < 0.24) (Fig 3A, 3B).

Figure 3 – Neural correlates of time deception

A. EEG Power modulations in the Cz electrode. Panel 1 represents relative power changes over contractions (respective to the first contractions) quantified in an individualized beta-band (Iβ) (see Methods for details). The dots represent the averaged distribution, and the error bars are the standard error of the mean (SEM). Panel 2 represents the Resting State (RS) power Pre-Post Variation Rate (VR) quantified in Iβ. B. Power differences between conditions in the Cz electrodes (N12-N10; B10-N10; N12-B12). The upper panel represents the difference for the sum of the Iβ Cz over the bins, while the lower one contains the difference for the Pre-Post VR of the Iβ Cz RS. C. EEG Power modulations in frontal electrodes. Relative power changes in θ (panel 1) and Iβ (panel 2) over contractions (respective to the first contractions). For both panels, the dots represent the averaged distribution, and the error bars are the SEM. D. Power differences in the frontal electrodes. Power differences were computed for each pair of interests (N12-N10; B10-N10; N12-B12). The upper and lower panels represent the difference in the sum of the frontal θ and Iβ power over the bins, respectively. For B. and D., the barplots represent the averaged distribution, and the error bars are the SEM. Asterisks denote a significant effect with p<0.05. The Bayes Factor (BF), rounded to the decimal, was reported when there was no significant difference.

We additionally looked at the Cz electrode's pre- to post-resting states (RS) Iβ power changes. Aligning with our previous results showing a consistent absence of physical fatigue difference between the sessions with a perceived short time, we found no difference between B10 and N10 Iβ power pre-post changes ($t_{(69)} = 0.13$, $p = .90$, $d_z = 0.030$, $BF = 0.22$). Other comparisons were found to be statistically inconclusive (N12-N10: $t_{(69)} = 1.59$, $p = .06$, $d_z = 0.27$, $BF = 0.81$; N12-B12: $t_{(69)} = 1.70$, $p = .093$, $d_z = 0.36$, $BF = 0.80$), though both showed a trend towards a significant difference (Fig 3A, 3B), overall reflecting the same pattern observed in the MVT results (Fig 1A, Fig 1C).

Power modulations in frontal electrodes

Next, we investigated oscillatory activity at frontal electrodes (F3, F4, F7, F8) – commonly used as an approximation of prefrontal cortex (PFC) activity (Okamoto et al., 2004; Robertson & Marino, 2015), as this region has been proposed to play a key role in effort maintenance (Bigliassi & Filho, 2022; Denis, 2020; McMorris et al., 2018; Robertson & Marino, 2016). We analyzed relative spectral power changes in the theta (θ) and Iβ bands. While frontal θ is assumed to reflect cognitive control (Cavanagh & Frank, 2014), the β band has also been proposed to have a similar function (Stoll et al., 2016). We found a significantly positive N12-N10 contrast in both the θ ($t_{(69)} = 2.06$, $p = .022$, $d_z = 0.40$, $BF = 2.10$) and Iβ bands ($t_{(69)} = 2.06$, $p = .021$, $d_z = 0.45$, $BF = 3.29$). Remarkably, we also found that the sessions with the same perceived time did not exhibit any power difference for both bands (B10-N10, θ: $t_{(69)} = 0.81$, $p = .42$, $d_z = 0.15$, $BF = 0.27$; Iβ: $t_{(69)} = 0.55$, $p = .58$, $d_z = 0.11$, $BF = 0.25$); B12-N12, θ: $t_{(69)} = 0.26$, $p = .79$, $d_z = 0.066$, $BF = 0.23$; Iβ: $t_{(69)} = 0.39$, $p = .70$, $d_z = 0.085$, $BF = 0.23$)) (Fig

213 3C, 3D), indicating a strong dependency of these variables on perceived rather than actual time.
214 Differently than in Cz, for which we did not observe an influence of the time manipulation,
215 these results suggest a key role of the frontal oscillatory activity in the perceived time-dependent
216 fatigue accumulation.
217
EEG-EMG Connectivity
After independently investigating the EMG and the EEG power modulations, we considered
the link between them by computing subject-by-subject correlations (see Methods for details).
We first found consistently across participants a negative correlation between EMG and Iβ Cz
for all sessions ($p < .045$, $d > 0.43$, $BF > 1.41$) (Fig 4A), indicating more EMG activity for
larger Iβ desynchronization. We then investigated the link between EMG and frontal oscillatory
activity. Similarly, we found a negative correlation between EMG activity and frontal θ-band
oscillations in B10 ($t_{(23)} = 2.70$, $p = .013$, $d_z = 0.46$, $BF = 3.96$), and as a trend in N10
($t_{(23)} = 1.99$, $p = .059$, $d_z = 0.41$, $BF = 1.14$), but no correlations in the N12 and B12 sessions (p
> 0.36, $d < 0.19$, $BF < 0.32$) (Fig 4B). Conversely, we found a positive correlation between the
EMG activity and the Iβ in the B12 session ($t_{(23)} = 2.25$, $p = .035$, $d_z = 0.46$, $BF = 1.74$) and, as
a trend, in the N12 one ($t_{(23)} = 1.75$, $p = .094$, $d_z = 0.36$, $BF = 0.80$), while there was no
correlation in the sessions whose perceived time was shorter (N10 and B10, $p > 0.75$, $d < 0.66$,
$BF < 0.23$, for both) (Fig 4C). These results mirror the trends observed in frontal power
modulations (Fig 3C), showing lower Frontal-EMG correlations for short perceived times and
larger correlations for longer ones, suggesting their dependency on perceived rather than actual
time.
235

Figure 4 – Brain-muscle correlations

Within-subject correlations were performed between EMG and θ Frontal (A), Iβ Cz (B), and Iβ Frontal (C),

respectively. The resulting r values were then compared against 0 using t-tests. The barplots represent the averaged

distribution, and the error bars are the standard error of the mean. Asterisks denote a significant effect with p<0.05.

The Bayes Factor (BF), rounded to the decimal, was reported when there was no significant difference.

Relationship between frontal oscillatory dynamics and fatigue accumulation

Our results show a perceived time-dependent accumulation of physical fatigue and a frontal

oscillatory dynamic that thoroughly follows perceived rather than actual time. Starting from the

assumption that PFC is involved in belief representation (Geuter et al., 2017), we hypothesized

that participants with the strongest deceptive effect in the frontal electrodes would also exhibit

the most substantial effect in the EMG. To test our hypothesis, we split the participants twice

into two groups, via a median split, computed on the normalized θ power difference between

the two sessions with the same perceived time. Such differences were computed as the absolute

sum of the θ power difference between the B10-N10 and the N12-B12 sessions, normalized by

the difference between the N12 and N10 conditions. In both cases, smaller differences

correspond to a larger deceptive effect, as the θ power modulation in frontal electrodes follows

the perceived rather than the actual time ('Subjective-time' (ST) group in Fig 5). Conversely,

participants with larger power differences would rely more on the actual rather than the

perceived time ('Objective-time' (OT) group in Fig 5). Note that we used the θ power difference

to perform our median split due to its well-established relationship to cognitive control

(Cavanagh & Frank, 2014).

Figure 5 – Physical fatigue accumulation regarding frontal oscillatory dynamics

A. EMG modulations over the main task. EMG is expressed in normalized RMS over contractions, whose increase reflects neuromuscular fatigue accumulation. Panels 1 and 2 contain the 'Subjective-time' participants, and panels 3 and 4 the 'Objective-time' ones (based on the deception effects measured in the θ-band, see text for details). Panels 1 and 3 were computed on the B10-N10 frontal θ absolute sum power difference and Panels 2 and 4 on the N12-B12 one. For all panels, the dots represent the averaged distribution, and the error bars are the standard error of the mean (SEM). B. EMG differences. EMG differences were computed for each pair of interest (N12-N10; B10-N10; N12-B12). Each panel is computed based on the sum of the EMG represented above it. For all panels, the barplots represent the averaged distribution, and the error bars are the SEM. Asterisks denote significant effect with p<0.05. The Bayes Factor (BF), rounded to the decimal, was reported when there was no significant difference.

As for the main analyses, we fitted linear mixed-effects models on the EMG, with Condition as a fixed factor and Participant as a random intercept, for each group separately, and focused on the three pre-defined contrasts. The ST group was the only one showing a significant positive difference between N12 and N10 in the fatigue accumulation as reflected by the EMG activity (ST: p < .047, BF > 2.58, OT: p > 0.076, BF < 0.86). Remarkably, when considering the θ-band power difference between B10-N10 conditions to split participants, we found an absence of difference between B10 and N10 ($t_{(33)}$ = 0.054, p = .96, d_z = 0.033, BF = 0.29), as well as between N12 and B12 ($t_{(33)}$ = 0.47, p = .64, d_z = 0.13, BF = 0.32) for the ST group in their EMG response. Further considering the θ-band power contrast between N12-B12 conditions to split participants, we also found a trend for an absence of difference for the two sessions with the

same perceived time (p > .22, BF < 0.63 for both B10-N10 and N12-B12) for the ST group.

Conversely, the OT group showed no difference for any pairs of sessions, regardless of the

power differences considered to split participants (p > 0.11, BF < 1.25 for all) (Fig 5A,5B).

Together, these results suggest that the frontal power changes occurring play a key role in the

effectiveness of the deceptive effect on fatigue accumulation, as measured by the EMG.

Interestingly, it also reveals that, in the ST group, fatigue can be slowed down and accelerated.

Link between interoceptive awareness and deception

To further explore the individual differences that may affect the strength of the deceptive

effect, we considered as a mediating parameter the participants' interoceptive awareness, which

was measured via the Multidimensional Assessment of Interoceptive Awareness (MAIA)

(Mehling et al., 2012) after their last session. We specifically focused on the 'Noticing' and

'Body Listening' subscales, which are more closely linked to bodily processes (Iodice et al.,

2019). We then tested the correlation between the Noticing and Body Listening scores with the

normalized absolute difference of neuromuscular fatigue accumulation (i.e., EMG) for the two

sessions with the same subjective time (B10-N10 or N12-B12). In short, a low normalized index

indicates no difference between the two sessions, suggesting a large deceptive effect, while a

high index reflects the opposite. We hypothesized a positive correlation between the

interoceptive score and the normalized index, i.e., the lower the interoceptive awareness, the

higher the deceptive effect, and vice-versa. Using non-parametric Kendall's tau correlation, we

found a positive correlation between the Body Listening score and the B10-N10 normalized

difference only ($\tau = 0.29$, p = 0.029, BF = 3.16, unilateral test, H1: positive correlation). No

correlation was found with the Noticing score (p > 0.27, BF < 0.47). These results suggest that

the participant's body awareness might predict the deceptive effect on neuromuscular fatigue

accumulation in the slow-down clock configuration.

As our findings previously suggested a role of the frontal oscillatory dynamics in the

deceptive effect, we tested the correlation between the Noticing and Body Listening scores with

the normalized absolute difference of θ frontal power modulations for the two sessions with the

same perceived time. We found a positive correlation between the Body Listening score and the

N12-B12 θ normalized difference only ($\tau = 0.31$, p = 0.020, BF = 4.30, unilateral test, H1:

positive correlation). While this result supports a link between interoception awareness and the

deceptive effect, it remains unclear why it only appears when time is accelerated but not slowed

down, as no other significant correlations were found (p > 0.11, BF < 1.02, unilateral test, H1:

positive correlation).

Discussion

In the current study, we used a novel design to test the influence of time deception on physical fatigue accumulation while monitoring associated EEG power modulations. We found that physical fatigue consistently evolved according to the perceived rather than the actual time when the clock was slowed down, while it was not systematic for the accelerated clock. Importantly, this deceptive effect was found in the absence of motivational confounds, demonstrating for the first time the direct influence of time deception on the course of physical fatigue. Regarding the neural correlates of the deceptive effect, we observed no significant impact in Cz, possibly due to a relatively small difference in workload between the sessions. However, we found that frontal oscillatory dynamics thoroughly followed the perceived time. We then showed that frontal dynamics predict the effectiveness of time deception on fatigue accumulation, suggesting its key role in the time deception effect.

Physical fatigue follows the perceived time

Our findings reveal that physical fatigue follows the subjective rather than the objective time when the clock is slowed down, echoing previous work emphasizing the mind's influence on bodily processes (Craik & Sarbin, 1963; Crum et al., 2011; Crum & Langer, 2007; Langer et al., 2010; London & Monello, 1974; Pomares et al., 2011; Schachter & Gross, 1968; Snyder et al., 1974; Thönes et al., 2021), such as blood glucose level (Park et al., 2016) and healing (Aungle & Langer, 2023). While physical fatigue has been proposed to depend on subjective time (Matta et al., 2024), the current study is the first to accurately disentangle the motivational effects from the temporal deception. Furthermore, our results seem to show an asymmetrical effect regarding clock manipulation. A possible explanation may involve efficiency considerations, as slowing down the process that conducts to exhaustion appears effective in fulfilling the required task (Meyniel et al., 2013), while accelerating it may be counterproductive. However, complementary analyses (Fig 5) nuance this effect, showing that some of the participants rely more on perceived time, both when the clock was slowed down and accelerated, while others follow objective time. Our findings suggest that interoceptive awareness mediates this effect: lower interoceptive awareness is associated with a larger deceptive effect, and vice versa. However, other factors likely influence this perceived-time dependency, such as the strength of participants' beliefs (e.g., expectations that shorter times lead to less fatigue) (Shiv & Carmon, 2005), the effectiveness of external clock manipulations on the perception of time (Craik & Sarbin, 1963; Thönes et al., 2018), but also participants'

backgrounds or personality (Zhou et al., 2019). Further research is needed to better characterize

the factors driving time deception's effectiveness on fatigue.

Neural correlates of the time deception effect

In the current study, we found no differences between sessions in the Cz oscillatory

dynamics, reflecting activity in motor areas, likely due to a relatively small workload difference

(Matta et al., 2025), making any potential difference due to the deceptive manipulation even

more challenging to observe. Yet, these parameters were carefully chosen to ensure the implicit

plausibility of the biased clock calibration, as increasing the manipulation's magnitude could

have raised awareness of the effect, potentially negating and confounding its impact (Ducrocq

et al., 2017; Williams et al., 2014).

Even though the task constraints imposed a relatively short temporal difference between

conditions, we nevertheless found a significant effect reflected in theta and beta frontal

oscillatory dynamics, which follow perceived rather than actual time. Interestingly, both theta

and beta-band activity increases have been previously associated with prolonged engagement

in demanding tasks (Boksem et al., 2005; Stoll et al., 2016). Specifically, enhanced theta power

has been extensively linked to increasing mental (Baumeister et al., 2012; Borghini et al., 2014;

Craig et al., 2012; Fan et al., 2015; Käthner et al., 2014; Liu et al., 2010; Shigihara et al., 2013;

Tanaka et al., 2014; Wascher et al., 2014) and, albeit less consistently, physical (Bailey et al.,

2008; Hottenrott & Gronwald, 2013; Mechau et al., 1998) fatigue – both supposedly sharing

common neural resources (Bray et al., 2012; Jin et al., 2024; Marcora et al., 2009; Mehta &

Parasuraman, 2014; Van Cutsem et al., 2017). This increase in theta power has been proposed

as reflecting a fatigue-related compensatory mechanism (Wascher et al., 2014) to allow

sustaining performance, seemingly driven by subjective rather than actual performance.

Moreover, given that frontal theta is a well-established marker of cognitive control (Arnau et

al., 2024; Cavanagh et al., 2012; Cavanagh & Frank, 2014; Ehrhardt et al., 2022; Howells et

al., 2010; Huster et al., 2013; Ishii et al., 1999; Lopez-Gamundi et al., 2024; McFerren et al.,

2021; Yu et al., 2022), our results suggest that the cognitive control required to sustain a

physical task depends on perceived rather than actual time. Finally, several studies also reported

an increase in frontal beta with the rise in fatigue (Bailey et al., 2008; Enders et al., 2016;

Hottenrott & Gronwald, 2013; Kubitz & Mott, 1996; Mechau et al., 1998). This increase might

also reflect a top-down cognitive control (Buschman & Miller, 2007; Stoll et al., 2016) allowing

one to continue the task despite uncomfortable sensations, which might follow perceived rather

than actual time. Together, these results highlight the reliance of specific cognitive control time-

related correlates to perceived rather than actual time. Additionally, it emphasizes the
importance of considering subjective components in cognitive experimental paradigms, as
various subjective factors may significantly influence physiological measures.

A predictive coding perspective
We found that frontal oscillatory dynamics (supposedly reflecting PFC's activity) and
physical fatigue evolve according to perceived rather than actual time. The influence of frontal
areas and the PFC on physical fatigue may be explained through predictive coding (Kaptchuk,
2018), a framework suggesting that the brain continuously generates expectations about
incoming sensory information (Friston, 2010). When there is a mismatch between expected and
actual sensations – a prediction error – the brain can resolve this discrepancy either by updating
its expectations (priors) or by actively adjusting bodily states (active inference) (Friston et al.,
2009; Pezzulo et al., 2015; Seth, 2013). If we apply this framework to our findings, we may
consider beliefs about time as prior factors that regulate physical fatigue through an active
inference process. In this situation, the temporal deceptive manipulation induces an inaccurate
temporal estimate, which generates a prediction error in the participants about their current
fatigue level. In order to minimize such prediction error – the discrepancy between expected
and actual fatigue – the brain may modulate interoceptive input (i.e., interoceptive signals
related to physical fatigue, later influencing EMG signals), thereby adjusting the evolution of
physical fatigue rather than revising its priors (Pagnini et al., 2023). In line with this framework,
frontal oscillatory dynamics could reflect temporal priors, as the PFC has been proposed to play
a key role in this process (Geuter et al., 2017). Furthermore, previous research has demonstrated
that beliefs can influence physiological effector systems such as the autonomic nervous system
(Amigo et al., 1993; Meissner, 2009, 2011; Nakamura et al., 2012; Wager et al., 2009). In line
with this interpretation, one may speculate that the change in the physical fatigue course is
mediated by an indirect pathway that originates in the PFC and regulates peripheral organs (e.g.,
cardiac, respiratory) (Hassan et al., 2013; Thayer et al., 2009, 2012), impacting muscle fatigue
as an outcome to reduce the prediction error. Furthermore, our complementary analysis,
splitting the participants based on their PFC power difference between sessions with the same
subjective time, support the hypothesis that frontal areas modulate the time manipulation effect.
Indeed, as shown in Figure 5, only participants exhibiting strong deceptive effects in the PFC
also experienced the deceptive effect on fatigue, as revealed in the EMG recordings. This result
may pinpoint the PFC as one of the core structure involved in mediating the time deception
effects on physical fatigue.

Conclusion

Our study demonstrates the direct influence of time deception on physical fatigue, irrespective of motivational factors and performance goals, and reveals key neural correlates of this effect. We show that this influence is primarily mediated by frontal oscillatory dynamics, which thoroughly follow the perceived time and are pivotal to the effectiveness of such deceptive time manipulation. Building on these findings, we propose potential mechanisms underlying this mind-body interaction. By showing that psychological factors can modulate physical fatigue, our results have broad implications for performance enhancement, but also everyday life and occupational settings, suggesting ways to optimize endurance, productivity, and safety in workers and other physically demanding contexts. Future research should further investigate the neural mechanisms driving this phenomenon from different perspectives (e.g., cardiac, respiratory or fMRI recordings) to further deepen its theoretical understanding. It may also explore applications across various contexts (e.g., sport, work, daily life) as well as its potential translantion to clinical conditions.

Methods

Participants

Twenty-eight healthy volunteers (16 women, age: 24 ± 4, weight: 70 ± 18, mean $\pm$ SD) with no known history of neurological diseases or physical injuries and regular physical practice (once to several times a week) participated in this study. Out of the 28 individuals, 24 were able to complete the entire protocol and were kept on for the analysis. Before participating in the study, all individuals provided informed consent and were informed of their right to withdraw from the study at any time. The four sessions were justified to the participants on the grounds that they allowed (i) the study of fatigue evolution between two different levels of effort (short and long contractions) and (ii) each level to be repeated twice to assess the reproducibility of fatigue measurements at the same level of effort. No further details were provided. At the end of the study, participants completed a survey addressing various aspects of their experience during the sessions (e.g., whether some sessions felt more difficult than others, whether the load seemed consistent across the four sessions, whether recovery time felt the same between sessions, and whether the duration of effort was similar across sessions). No participant reported any specific remarks suggesting awareness of the temporal manipulation. Participants were only informed of the deception after the study was completed. The study procedure was approved by the French Ethical 'Comité de Protection des Personnes' (number: 2023-A00412-43).

Data Acquisition

During each session, all participants were seated on a Biodex isokinetic ergometer with their hips and knees bent to a 90° angle. Their right leg was securely fastened to the ergometer accessory at the ankle with a non-compliant strap just above the malleolus. Two shoulder and one abdominal harnesses guaranteed a reduction in their range of motion and the involvement of other muscles besides the quadriceps. The installation parameters were optimized for every subject individually and kept identical within subjects for each session.

Torque and electromyography (EMG) were recorded at 2000 Hz using a Biopac MP150 system and AcqKnowledge software. EMG was recorded through three pairs of electrodes (AG-Cl, diameter: 11 mm, interelectrode distance: 2 cm), which were placed on the Vastus Medialis, Vastus Lateralis, and Rectus Femoris according to SENIAM recommendations. Before placement, the skin was carefully prepared through shaving and alcohol to ensure low impedance (<5 KΩ). The reference electrode was positioned on the patella of the contralateral

knee. EEG was recorded using a 64 electrodes Biosemi Active 2 system with 64 Ag-AgCl sintered active electrodes at 2048 Hz. The EEG electrode locations followed the 10–20 international system. The offset was kept below 20 mV using an active gel. All acquisitions were performed with the light off to minimize electrical noise.

The main task was programmed in Python and displayed through a Windows computer (Resolution: 1920 × 1080; Refresh rate: 60 Hz), using a NiDAQ acquisition card (NI USB-6218, Sample rate: 2000 Hz) to read the torque data (low-pass filtered at 10 Hz) in real-time. The same computer also generated the triggers used to mark the onset and offset of each contraction towards BIOPAC and BIOSEMI, allowing us to synchronize all our signals offline.

Experimental Protocol

All participants came to the lab on four days (each spaced 3 to 10 days apart) to perform four sessions. On each day, the participants performed the following steps (Fig 1):
- Warm-up (consisting of 15-20 contractions with increasing intensity, approximately from 20 to 80% of their estimated Maximal Voluntary Torque (MVT))
- 2 Maximal Voluntary Contractions (MVC) (3-4 s each, 30 s spaced)
- 1 MVC, 20 s of rest, 1 Long MVC (30 s), 30 s of rest, 3 minutes of EEG recording at rest (eyes opened)
- 5 minutes of rest
- Main task: 100 contractions (see below)
- 1 MVC (5 s after the 100th contraction of the main task), 20 s of rest, 1 Long MVC (30 s), 30 s of rest, 3 minutes of EEG recording at rest (eyes opened)

Regarding the main task, participants performed 100 contractions at a fixed individualized intensity, defined as 20% of their maximal force, or 40 N.m if their maximal force was larger than 200 N.m. The intensity was the same among the four sessions. While the rest time between contractions was identical for each session (5 s), each contraction's real (R) and perceived (P) time were independently manipulated. In each session, the contractions' time was either short (10s) or long (12s), while the clock displayed to the participants was either Normal (N) or Biased (B). This manipulation led to 4 counterbalanced conditions: N10 (10s P, 10s R), N12 (12s P, 12s R), B10 (10s P, 12s R), B12 (12s P, 10s R). In practice, a sound indicated to the participants when to begin and end each contraction. Simultaneously with the starting sound, a rectangle was displayed in the middle of a black screen to provide online feedback to the participants on their actual force compared to the individualized targeted one. The rectangle's color was updated online according to the ongoing force produced: red indicated that

participants were below the targeted force, blue above, and green in the aimed range (-5% to

+5% around the targeted force). A digital clock was displayed in the middle of the colored

rectangle 0.75 s after the start of the contraction (preparation time left to the participants to

reach the targeted force). This clock allowed us to perform our manipulation: it was either

normal (N10 and N12), slowed down (B10), or accelerated (B12) without the participants

knowing it. After the ending sound, the final perceived time (10.0 or 12.0 s) remained on the

screen for 1 s, then the number of contractions already performed was displayed during the

resting time. Note that data from the two normal sessions (N10 and N12) have been previously

reported (Matta et al., 2025) in the context of a different question; Long MVC were specifically

used there.

Data pre-processing

All data pre-processing and analyses were conducted offline using custom MATLAB

(version 2022b) scripts along with EEGLAB functions (Delorme & Makeig, 2004). Torque and

EMG data were down-sampled offline to 1000 Hz. Torque data were then low-pass filtered at

10 Hz, while EMG data underwent band-pass filtering between 20 and 500 Hz and notch

filtering between 47 and 53 Hz.

EEG data were down-sampled offline to 250 Hz. A notch filter (47 to 53 Hz) and a high-pass

filter with a 1 Hz cut-off frequency were applied. Channels with excessive noise were

automatically rejected using the CleanRawData plugin, configured with a 5-second 'flat line

criterion,' a 0.6 'minimum channel correlation,' and a kurtosis threshold exceeding five times

the mean. On average, 61.7 ± 1.8 (mean $\pm$ SD) channels were retained for analysis. The

electrodes were then re-referenced using a common average reference. Independent Component

Analysis (ICA) was performed with the Infomax algorithm, and the resulting components were

labeled using the ICLabel plugin (Pion-Tonachini et al., 2019). Components labeled as muscle,

eye, heart, line noise, channel noise, or other with a probability above 90% were automatically

rejected, averaging 12.9 ± 5.5 components removed. Previously excluded channels were

subsequently interpolated.

EEG data from the two resting-state periods were segmented into 2-second non-overlapping

windows. For the main task, EEG and EMG data were segmented into 18-second epochs,

starting 3 seconds before the onset of the contraction (as marked by its trigger) and ending 15

seconds after. A final epoch rejection procedure was applied to the main task EEG data to

remove additional noise. Epochs with a standard deviation greater than 1.5 times the

interquartile range (IQR) above the 3rd quartile or below 1.5 times the IQR under the 1st

quartile were rejected for each subject and electrode. This procedure eliminated 3.9% of the epochs over the four sessions.

Data analysis

Force and EMG

<u>Pre-Post.</u> The highest torque from the three MVCs performed before the task was used to determine the pre-test Maximal Voluntary Torque (MVT). The post-test measurement was taken as the highest torque recorded during the post-task MVC. The variation rate was calculated as (post − pre)/pre ×100.

<u>Main Task.</u> EMG was quantified as the root-mean-square (RMS) during each contraction of the main task from 2 to 10 seconds after the onset of the contraction (Fig 1). This period was chosen to exclude the beginning of the contraction, where torque stabilization has not yet occurred, and to ensure identical comparison times for all sessions. Within this period, EMG was quantified as the RMS over 180 ms non-overlapping windows, normalized by the RMS EMG associated with the previously selected maximal torque (centered from -90 ms to +90 ms around it) for the respective muscle and session. The three muscle EMG values were averaged over time to yield one EMG value per contraction in the main task. These EMG values were then averaged in bins of 10 contractions, with the first bin zero-centered, and the sum of the EMG evolution over the task was computed, resulting in a single representative EMG value for each variable and session.

EEG Spectral Power

Spectral power was obtained by applying a continuous wavelet transform to the pre-processed, epoched EEG data within the 1 to 50 Hz frequency range. The wavelet length was linearly increased from one cycle at 1 Hz to twenty cycles at 50 Hz, optimizing the tradeoff between temporal and frequency resolution across the entire frequency range. Contractions that were removed were subsequently linearly interpolated.

Spectral power was specifically extracted for the Theta (θ, 4-8 Hz) and an Individualized Beta band (Iβ), which was defined for each subject to account for inter-individual variability in the beta desynchronization-specific frequency band (Fig 1). Specifically, the frequency with the largest desynchronization was identified for each subject, and Iβ was defined as the range from -2 to +2 Hz around this frequency (Matta et al., 2025). Power from the lateral-frontal (F7-F8) and mid-frontal (F3-F4) electrodes was averaged to create a 'Frontal' cluster, chosen to reflect

the ventrolateral and dorsolateral prefrontal cortex (Okamoto et al., 2004; Robertson & Marino, 2015) activity. The Cz electrode was used to reflect motor cortex activity associated with quadriceps contraction, a location confirmed by topographically visualizing beta desynchronization (Dal Maso et al., 2012; Forrester et al., 2006; Gordon et al., 2023; Meier et al., 2008).

Pre-Post Analysis. Raw spectral powers from pre- and post-EEG resting state recordings were independently averaged over epochs (90) and time (0 to 2 s). The variation rate was calculated between the pre- and post-values for each frequency band and region of interest.

Main Task Analysis. For each contraction, raw spectral power bands during the period of interest (2 to 10 s) were normalized to the first contractions of the session, averaged over the same 2 to 10 s period (Fig 1). This normalization, focusing on the fatigue effect, involved subtracting the baseline and dividing by it, then multiplying by 100 to express power values as a percentage of a non-fatigued state. The baseline was defined as the average of contractions 2 to 5, excluding the first contraction due to its significant torque variability in both sessions. The resulting values were averaged over time and frequencies, yielding one power value per frequency band, contraction, and region of interest. As for the EMG, values were then averaged in bins of 10 contractions, with the first bin zero-centered. Finally, the sum of the power evolution over the task was eventually extracted, yielding one representative value for each variable and session.

EEG-EMG Connectivity

To investigate the EMG-EEG relationships, we computed Pearson's correlation between EMG and Iβ Cz, θ Frontal, and Iβ Frontal, for each subject respectively. We then performed bilateral one-sample t-tests to determine whether the r values significantly differed from zero.

Statistical analyses

Dependent variables were analyzed using linear mixed-effects models, with real and subjective times included as fixed factors and a random intercept for each participant. The N10 condition was set as the reference condition to which all other conditions were initially compared. As only specific pairs of conditions were of interest, estimated marginal means were subsequently analyzed with contrasts to compare three pairs of real and perceived times.

First, we focused on N12–N10 to assess the effect of a longer effort without time manipulation, for which a significant difference was expected (one-sided t-test), thereby isolating the effect of fatigue. Second, we compared N12–B12, representing conditions in

which perceived duration was long but actual duration varied, highlighting the potential acceleration of fatigue. Finally, we analyzed B10–N10, representing conditions in which perceived duration was short but actual duration differed, highlighting the potential slowing of fatigue. Two-sided t-tests were applied for these contrasts, as we did not have directional a priori hypotheses.

To properly evaluate support for the null hypothesis, Bayesian paired t-tests were also conducted for these three contrasts. In the Bayesian framework, the Bayes Factor (BF) represents the ratio of the likelihood of the data under the alternative hypothesis to that under the null hypothesis. Conventionally, BFs > ~3 provide evidence in favor of the alternative hypothesis, while BFs < ~0.3 indicate evidence in favor of the null hypothesis (van Doorn et al., 2021).

Statiscal analyses were conducted using R (CranR, version 4.3.1) and JASP (2023, version 0.18.2) (Love et al., 2019), with default uniform prior distributions for the Bayesian analyses.

Funding

This project has received funding from the H2020 European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreements No 101075930 – Andrea Alamia). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them. Pierre-Marie Matta received funding from the French Ministère de l'Enseignement Supérieur et de la Recherche (MESR).

Acknowledgements

We thank Maxime Picquet, Romain Martinie, Khiara Vorzillo, and Nina Bertho for their valuable help with the data acquisition. We are also grateful to Laurent Gonthier for his technical support in setting up the online force acquisition card, and to Nathalie Vayssiere for her assistance in obtaining the ethical agreement.

References

Amigo, I., Cuesta, V., Fernández, A., & González, A. (1993). The effect of verbal instructions on blood pressure measurement. *Journal of Hypertension*, *11*(3), 293-296. https://doi.org/10.1097/00004872-199303000-00010

Ansdell, P., Thomas, K., Howatson, G., Amann, M., & Goodall, S. (2018). Deception Improves Time Trial Performance in Well-trained Cyclists without Augmented Fatigue. *Medicine and Science in Sports and Exercise*, *50*(4), 809-816. https://doi.org/10.1249/MSS.0000000000001483

Arnau, S., Liegel, N., & Wascher, E. (2024). Frontal midline theta power during the cue-target-interval reflects increased cognitive effort in rewarded task-switching. *Cortex; a Journal Devoted to the Study of the Nervous System and Behavior*, *180*, 94-110. https://doi.org/10.1016/j.cortex.2024.08.004

Aungle, P., & Langer, E. (2023). Physical healing as a function of perceived time. *Scientific Reports*, *13*(1), 22432. https://doi.org/10.1038/s41598-023-50009-3

Bailey, S. P., Hall, E. E., Folger, S. E., & Miller, P. C. (2008). Changes in EEG during graded exercise on a recumbent cycle ergometer. *Journal of Sports Science & Medicine*, *7*(4), 505-511.

Baumeister, J., Reinecke, K., Schubert, M., Schade, J., & Weiss, M. (2012). Effects of induced fatigue on brain activity during sensorimotor control. *European Journal of Applied Physiology*, *112*(7), 2475-2482. https://doi.org/10.1007/s00421-011-2215-6

Behrens, M., Gube, M., Chaabene, H., Prieske, O., Zenon, A., Broscheid, K.-C., Schega, L., Husmann, F., & Weippert, M. (2023). Fatigue and Human Performance : An Updated Framework. *Sports Medicine (Auckland, N.Z.)*, *53*(1), 7-31. https://doi.org/10.1007/s40279-022-01748-2

Bigland-Ritchie, B., & Woods, J. J. (1984). Changes in muscle contractile properties and neural control during human muscular fatigue. *Muscle & Nerve*, *7*(9), 691-699. https://doi.org/10.1002/mus.880070902

Bigliassi, M., & Filho, E. (2022). Functional significance of the dorsolateral prefrontal cortex during exhaustive exercise. *Biological Psychology*, *175*, 108442. https://doi.org/10.1016/j.biopsycho.2022.108442

Boksem, M. A. S., Meijman, T. F., & Lorist, M. M. (2005). Effects of mental fatigue on attention : An ERP study. *Brain Research. Cognitive Brain Research*, *25*(1), 107-116. https://doi.org/10.1016/j.cogbrainres.2005.04.011

Borghini, G., Astolfi, L., Vecchiato, G., Mattia, D., & Babiloni, F. (2014). Measuring

neurophysiological signals in aircraft pilots and car drivers for the assessment of mental workload, fatigue and drowsiness. *Neuroscience and Biobehavioral Reviews*, *44*, 58-75. https://doi.org/10.1016/j.neubiorev.2012.10.003

Bray, S. R., Graham, J. D., Martin Ginis, K. A., & Hicks, A. L. (2012). Cognitive task performance causes impaired maximum force production in human hand flexor muscles. *Biological Psychology*, *89*(1), 195-200. https://doi.org/10.1016/j.biopsycho.2011.10.008

Brown, P. (2000). Cortical drives to human muscle : The Piper and related rhythms. *Progress in Neurobiology*, *60*(1), 97-108. https://doi.org/10.1016/s0301-0082(99)00029-5

Buschman, T. J., & Miller, E. K. (2007). Top-down versus bottom-up control of attention in the prefrontal and posterior parietal cortices. *Science (New York, N.Y.)*, *315*(5820), 1860-1862. https://doi.org/10.1126/science.1138071

Cavanagh, J. F., Figueroa, C. M., Cohen, M. X., & Frank, M. J. (2012). Frontal theta reflects uncertainty and unexpectedness during exploration and exploitation. *Cerebral Cortex (New York, N.Y.: 1991)*, *22*(11), 2575-2586. https://doi.org/10.1093/cercor/bhr332

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control. *Trends in Cognitive Sciences*, *18*(8), 414-421. https://doi.org/10.1016/j.tics.2014.04.012

Chan, M. (2011). Fatigue : The most critical accident risk in oil and gas construction. *Construction Management & Economics*, *29*, 341-353. https://doi.org/10.1080/01446193.2010.545993

Craig, A., Tran, Y., Wijesuriya, N., & Nguyen, H. (2012). Regional brain wave activity changes associated with fatigue. *Psychophysiology*, *49*(4), 574-582. https://doi.org/10.1111/j.1469-8986.2011.01329.x

Craik, K. H., & Sarbin, T. R. (1963). Effect of covert alteratons of clock rate upon time estimations and personal tempo. *Perceptual and Motor Skills*, *16*, 597-610. https://doi.org/10.2466/pms.1963.16.2.597

Crum, A. J., Corbin, W. R., Brownell, K. D., & Salovey, P. (2011). Mind over milkshakes : Mindsets, not just nutrients, determine ghrelin response. *Health Psychology: Official Journal of the Division of Health Psychology, American Psychological Association*, *30*(4), 424-429; discussion 430-431. https://doi.org/10.1037/a0023467

Crum, A. J., & Langer, E. J. (2007). Mind-set matters : Exercise and the placebo effect. *Psychological Science*, *18*(2), 165-171. https://doi.org/10.1111/j.1467-9280.2007.01867.x

Cunningham, T. R., Guerin, R. J., Ferguson, J., & Cavallari, J. (2022). Work-related fatigue : A hazard for workers experiencing disproportionate occupational risks. *American journal of industrial medicine*, *65*(11), 913-925. https://doi.org/10.1002/ajim.23325

Dal Maso, F., Longcamp, M., & Amarantini, D. (2012). Training-related decrease in antagonist muscles activation is associated with increased motor cortex activation : Evidence of central mechanisms for control of antagonist muscles. *Experimental Brain Research*, *220*(3-4), 287-295. https://doi.org/10.1007/s00221-012-3137-1

Delorme, A., & Makeig, S. (2004). EEGLAB : An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9-21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Denis, G. (2020). *Implication des dimensions neurocognitives dans le maintien de l'effort physique au travers du rôle endossé par le cortex préfrontal et de la perspective coûts/bénéfices*. Université Côte d'Azur.

Dijkerman, C., & Lenggenhager, B. (2018). The body and cognition : The relation between body representations and higher level cognitive and social processes. *Cortex; a Journal Devoted to the Study of the Nervous System and Behavior*, *104*, 133-139. https://doi.org/10.1016/j.cortex.2018.06.001

Ducrocq, G. P., Hureau, T. J., Meste, O., & Blain, G. M. (2017). Increased Fatigue Response to Augmented Deceptive Feedback during Cycling Time Trial. *Medicine and Science in Sports and Exercise*, *49*(8), 1541-1551. https://doi.org/10.1249/MSS.0000000000001272

Ehrhardt, N. M., Fietz, J., Kopf-Beck, J., Kappelmann, N., & Brem, A.-K. (2022). Separating EEG correlates of stress : Cognitive effort, time pressure, and social-evaluative threat. *The European Journal of Neuroscience*, *55*(9-10), 2464-2473. https://doi.org/10.1111/ejn.15211

Enders, H., Cortese, F., Maurer, C., Baltich, J., Protzner, A. B., & Nigg, B. M. (2016). Changes in cortical activity measured with EEG during a high-intensity cycling exercise. *Journal of Neurophysiology*, *115*(1), 379-388. https://doi.org/10.1152/jn.00497.2015

Enoka, R. M., & Duchateau, J. (2016). Translating Fatigue to Human Performance. *Medicine and Science in Sports and Exercise*, *48*(11), 2228-2238. https://doi.org/10.1249/MSS.0000000000000929

Fan, X., Zhou, Q., Liu, Z., & Xie, F. (2015). Electroencephalogram assessment of mental fatigue in visual search. *Bio-Medical Materials and Engineering*, *26 Suppl 1*, S1455-1463. https://doi.org/10.3233/BME-151444

Forrester, L. W., Hanley, D. F., & Macko, R. F. (2006). Effects of treadmill exercise on transcranial magnetic stimulation-induced excitability to quadriceps after stroke. *Archives of Physical Medicine and Rehabilitation*, *87*(2), 229-234. https://doi.org/10.1016/j.apmr.2005.10.016

Friston, K. (2010). The free-energy principle : A unified brain theory? *Nature Reviews.*

Neuroscience, *11*(2), 127-138. https://doi.org/10.1038/nrn2787

Friston, K. J., Daunizeau, J., & Kiebel, S. J. (2009). Reinforcement learning or active

inference? *PloS One*, *4*(7), e6421. https://doi.org/10.1371/journal.pone.0006421

Gandevia, S. C. (2001). Spinal and supraspinal factors in human muscle fatigue.

Physiological Reviews, *81*(4), 1725-1789. https://doi.org/10.1152/physrev.2001.81.4.1725

Geuter, S., Koban, L., & Wager, T. D. (2017). The Cognitive Neuroscience of Placebo

Effects : Concepts, Predictions, and Physiology. *Annual Review of Neuroscience*, *40*, 167-188.

https://doi.org/10.1146/annurev-neuro-072116-031132

Gordon, E. M., Chauvin, R. J., Van, A. N., Rajesh, A., Nielsen, A., Newbold, D. J., Lynch,

C. J., Seider, N. A., Krimmel, S. R., Scheidter, K. M., Monk, J., Miller, R. L., Metoki, A.,

Montez, D. F., Zheng, A., Elbau, I., Madison, T., Nishino, T., Myers, M. J., … Dosenbach, N.

U. F. (2023). A somato-cognitive action network alternates with effector regions in motor

cortex. *Nature*, *617*(7960), 351-359. https://doi.org/10.1038/s41586-023-05964-2

Hassan, S. F., Cornish, J. L., & Goodchild, A. K. (2013). Respiratory, metabolic and cardiac

functions are altered by disinhibition of subregions of the medial prefrontal cortex. *The Journal*

of Physiology, *591*(23), 6069-6088. https://doi.org/10.1113/jphysiol.2013.262071

Hottenrott, K., & Gronwald, T. (2013). Cortical Brain Activity is Influenced by Cadence in

Cyclists. *The Open Sports Sciences Journal*, *6*, 9-14.

https://doi.org/10.2174/1875399X01306010009

Howells, F. M., Stein, D. J., & Russell, V. A. (2010). Perceived mental effort correlates with

changes in tonic arousal during attentional tasks. *Behavioral and Brain Functions: BBF*, *6*, 39.

https://doi.org/10.1186/1744-9081-6-39

Huster, R. J., Enriquez-Geppert, S., Lavallee, C. F., Falkenstein, M., & Herrmann, C. S.

(2013). Electroencephalography of response inhibition tasks : Functional networks and

cognitive contributions. *International Journal of Psychophysiology: Official Journal of the*

International Organization of Psychophysiology, *87*(3), 217-233.

https://doi.org/10.1016/j.ijpsycho.2012.08.001

Ibrahim, A., Nnaji, C., Namian, M., Koh, A., & Techera, U. (2023). Investigating the impact

of physical fatigue on construction workers' situational awareness. *Safety Science*, *163*, 106103.

https://doi.org/10.1016/j.ssci.2023.106103

Iodice, P., Porciello, G., Bufalari, I., Barca, L., & Pezzulo, G. (2019). An interoceptive

illusion of effort induced by false heart-rate feedback. *Proceedings of the National Academy of*

Sciences of the United States of America, *116*(28), 13897-13902.

https://doi.org/10.1073/pnas.1821032116

Ishii, R., Shinosaki, K., Ukai, S., Inouye, T., Ishihara, T., Yoshimine, T., Hirabuki, N.,
Asada, H., Kihara, T., Robinson, S. E., & Takeda, M. (1999). Medial prefrontal cortex generates
frontal midline theta rhythm. *Neuroreport*, *10*(4), 675-679. https://doi.org/10.1097/00001756-
199903170-00003

Jin, H., Xiao, M., Liu, L., Kan, S., Fu, Y., & Zhang, D. (2024). Relationship between
physical fatigue and mental fatigue based on multimodal measurement under different load
levels. *Ergonomics*, *67*(11), 1748-1763. https://doi.org/10.1080/00140139.2024.2364667

Jones, C. M., Griffiths, P. C., & Mellalieu, S. D. (2017). Training Load and Fatigue Marker
Associations with Injury and Illness : A Systematic Review of Longitudinal Studies. *Sports*
Medicine (Auckland, N.z.), *47*(5), 943-974. https://doi.org/10.1007/s40279-016-0619-5

Jones, H. S., Williams, E. L., Bridge, C. A., Marchant, D., Midgley, A. W., Micklewright,
D., & Mc Naughton, L. R. (2013). Physiological and psychological effects of deception on
pacing strategy and performance : A review. *Sports Medicine (Auckland, N.Z.)*, *43*(12),
1243-1257. https://doi.org/10.1007/s40279-013-0094-1

Kaptchuk, T. J. (2018). Open-Label Placebo : Reflections on a Research Agenda.
Perspectives in Biology and Medicine, *61*(3), 311-334. https://doi.org/10.1353/pbm.2018.0045

Käthner, I., Wriessnegger, S. C., Müller-Putz, G. R., Kübler, A., & Halder, S. (2014). Effects
of mental workload and fatigue on the P300, alpha and theta band power during operation of
an ERP (P300) brain-computer interface. *Biological Psychology*, *102*, 118-129.
https://doi.org/10.1016/j.biopsycho.2014.07.014

Kim, G., Ahad, M. A., Ferdjallah, M., & Harris, G. F. (2007). Correlation of muscle fatigue
indices between intramuscular and surface EMG signals. *Proceedings 2007 IEEE*
SoutheastCon, 378-382. https://doi.org/10.1109/SECON.2007.342928

Kreher, J. B., & Schwartz, J. B. (2012). Overtraining Syndrome. *Sports Health*, *4*(2),
128-138. https://doi.org/10.1177/1941738111434406

Kubitz, K. A., & Mott, A. A. (1996). EEG power spectral densities during and after cycle
ergometer exercise. *Research Quarterly for Exercise and Sport*, *67*(1), 91-96.
https://doi.org/10.1080/02701367.1996.10607929

Langer, E., Djikic, M., Pirson, M., Madenci, A., & Donohue, R. (2010). Believing is seeing :
Using mindlessness (mindfully) to improve visual acuity. *Psychological Science*, *21*(5),
661-666. https://doi.org/10.1177/0956797610366543

Liu, J.-P., Zhang, C., & Zheng, C.-X. (2010). Estimation of the cortical functional
connectivity by directed transfer function during mental fatigue. *Applied Ergonomics*, *42*(1),
114-121. https://doi.org/10.1016/j.apergo.2010.05.008

London, H., & Monello, L. (1974). Cognitive manipulation of boredom. In *Thought and*

feeling : Cognitive alteration of feeling states. (p. 239-239). Aldine.

Lopez-Gamundi, P., Mas-Herrero, E., & Marco-Pallares, J. (2024). Disentangling effort

from probability of success : Temporal dynamics of frontal midline theta in effort-based reward

processing. *Cortex; a Journal Devoted to the Study of the Nervous System and Behavior*, *176*,

94-112. https://doi.org/10.1016/j.cortex.2024.03.014

Love, J., Selker, R., Marsman, M., Jamil, T., Dropmann, D., Verhagen, A. J., Ly, A., Gronau,

Q., Šmíra, M., Epskamp, S., Matzke, D., Wild, A., Knight, P., Rouder, J., Morey, R., &

Wagenmakers, E.-J. (2019). JASP : Graphical Statistical Software for Common Statistical

Designs. *Journal of Statistical Software*, *88*. https://doi.org/10.18637/jss.v088.i02

Marcora, S. M., Staiano, W., & Manning, V. (2009). Mental fatigue impairs physical

performance in humans. *Journal of Applied Physiology (Bethesda, Md.: 1985)*, *106*(3),

857-864. https://doi.org/10.1152/japplphysiol.91324.2008

Matta, P.-M., Baurès, R., Duclay, J., & Alamia, A. (2025). Modulation of beta oscillatory

dynamics in motor and frontal areas during physical fatigue. *Communications Biology*, *8*(1),

687. https://doi.org/10.1038/s42003-025-08122-8

Matta, P.-M., Glories, D., Alamia, A., Baurès, R., & Duclay, J. (2024). Mind over muscle?

Time manipulation improves physical performance by slowing down the neuromuscular fatigue

accumulation. *Psychophysiology*, *61*(4), e14487. https://doi.org/10.1111/psyp.14487

McFerren, A., Riddle, J., Walker, C., Buse, J. B., & Frohlich, F. (2021). Causal role of

frontal-midline theta in cognitive effort : A pilot study. *Journal of Neurophysiology*, *126*(4),

1221-1233. https://doi.org/10.1152/jn.00068.2021

McMorris, T., Barwood, M., & Corbett, J. (2018). Central fatigue theory and endurance

exercise : Toward an interoceptive model. *Neuroscience and Biobehavioral Reviews*, *93*,

93-107. https://doi.org/10.1016/j.neubiorev.2018.03.024

Mechau, D., Mücke, S., Weiss, M., & Liesen, H. (1998). Effect of increasing running

velocity on electroencephalogram in a field test. *European Journal of Applied Physiology and*

Occupational Physiology, *78*(4), 340-345. https://doi.org/10.1007/s004210050429

Mehling, W. E., Price, C., Daubenmier, J. J., Acree, M., Bartmess, E., & Stewart, A. (2012).

The Multidimensional Assessment of Interoceptive Awareness (MAIA). *PLoS ONE*, *7*(11),

e48230. https://doi.org/10.1371/journal.pone.0048230

Mehta, R. K., & Parasuraman, R. (2014). Effects of mental fatigue on the development of

physical fatigue : A neuroergonomic approach. *Human Factors*, *56*(4), 645-656.

https://doi.org/10.1177/0018720813507279

Meier, J. D., Aflalo, T. N., Kastner, S., & Graziano, M. S. A. (2008). Complex organization

of human primary motor cortex : A high-resolution fMRI study. *Journal of Neurophysiology*,

100(4), 1800-1812. https://doi.org/10.1152/jn.90531.2008

Meissner, K. (2009). Effects of placebo interventions on gastric motility and general

autonomic activity. *Journal of Psychosomatic Research*, *66*(5), 391-398.

https://doi.org/10.1016/j.jpsychores.2008.09.004

Meissner, K. (2011). The placebo effect and the autonomic nervous system : Evidence for

an intimate relationship. *Philosophical Transactions of the Royal Society of London. Series B,*

Biological Sciences, *366*(1572), 1808-1817. https://doi.org/10.1098/rstb.2010.0403

Merletti, R., Lo Conte, L. R., & Orizio, C. (1991). Indices of muscle fatigue. *Journal of*

Electromyography and Kinesiology: Official Journal of the International Society of

Electrophysiological Kinesiology, *1*(1), 20-33. https://doi.org/10.1016/1050-6411(91)90023-X

Meyniel, F., Sergent, C., Rigoux, L., Daunizeau, J., & Pessiglione, M. (2013).

Neurocomputational account of how the human brain decides when to have a break.

Proceedings of the National Academy of Sciences of the United States of America, *110*(7),

2641-2646. https://doi.org/10.1073/pnas.1211925110

Nakamura, Y., Donaldson, G. W., Kuhn, R., Bradshaw, D. H., Jacobson, R. C., & Chapman,

R. C. (2012). Investigating dose-dependent effects of placebo analgesia : A

psychophysiological approach. *Pain*, *153*(1), 227-237.

https://doi.org/10.1016/j.pain.2011.10.024

Okamoto, M., Dan, H., Sakamoto, K., Takeo, K., Shimizu, K., Kohno, S., Oda, I., Isobe, S.,

Suzuki, T., Kohyama, K., & Dan, I. (2004). Three-dimensional probabilistic anatomical cranio-

cerebral correlation via the international 10-20 system oriented for transcranial functional brain

mapping. *NeuroImage*, *21*(1), 99-111. https://doi.org/10.1016/j.neuroimage.2003.08.026

Pagnini, F., Barbiani, D., Cavalera, C., Volpato, E., Grosso, F., Minazzi, G. A., Vailati

Riboni, F., Graziano, F., Di Tella, S., Manzoni, G. M., Silveri, M. C., Riva, G., & Phillips, D.

(2023). Placebo and Nocebo Effects as Bayesian-Brain Phenomena : The Overlooked Role of

Likelihood and Attention. *Perspectives on Psychological Science: A Journal of the Association*

for Psychological Science, *18*(5), 1217-1229. https://doi.org/10.1177/17456916221141383

Park, C., Pagnini, F., Reece, A., Phillips, D., & Langer, E. (2016). Blood sugar level follows

perceived time rather than actual time in people with type 2 diabetes. *Proceedings of the*

National Academy of Sciences of the United States of America, *113*(29), 8168-8170.

https://doi.org/10.1073/pnas.1603444113

Pearson, K. (2000). Motor systems. *Current Opinion in Neurobiology*, *10*(5), 649-654.

https://doi.org/10.1016/s0959-4388(00)00130-6

Pezzulo, G., Rigoli, F., & Friston, K. (2015). Active Inference, homeostatic regulation and adaptive behavioural control. *Progress in Neurobiology*, *134*, 17-35. https://doi.org/10.1016/j.pneurobio.2015.09.001

Pion-Tonachini, L., Kreutz-Delgado, K., & Makeig, S. (2019). ICLabel : An automated electroencephalographic independent component classifier, dataset, and website. *NeuroImage*, *198*, 181-197. https://doi.org/10.1016/j.neuroimage.2019.05.026

Pomares, F. B., Creac'h, C., Faillenot, I., Convers, P., & Peyron, R. (2011). How a clock can change your pain? The illusion of duration and pain perception. *Pain*, *152*(1), 230-234. https://doi.org/10.1016/j.pain.2010.10.047

Robertson, C. V., & Marino, F. E. (2015). Prefrontal and motor cortex EEG responses and their relationship to ventilatory thresholds during exhaustive incremental exercise. *European Journal of Applied Physiology*, *115*(9), 1939-1948. https://doi.org/10.1007/s00421-015-3177-x

Robertson, C. V., & Marino, F. E. (2016). A role for the prefrontal cortex in exercise tolerance and termination. *Journal of Applied Physiology (Bethesda, Md.: 1985)*, *120*(4), 464-466. https://doi.org/10.1152/japplphysiol.00363.2015

Rodrigues, F., Monteiro, D., Ferraz, R., Branquinho, L., & Forte, P. (2023). The Association between Training Frequency, Symptoms of Overtraining and Injuries in Young Men Soccer Players. *International Journal of Environmental Research and Public Health*, *20*(8), 5466. https://doi.org/10.3390/ijerph20085466

Schachter, S., & Gross, L. P. (1968). Manipulated time and eating behavior. *Journal of Personality and Social Psychology*, *10*(2), 98-106. https://doi.org/10.1037/h0026285

Seth, A. K. (2013). Interoceptive inference, emotion, and the embodied self. *Trends in Cognitive Sciences*, *17*(11), 565-573. https://doi.org/10.1016/j.tics.2013.09.007

Shigihara, Y., Tanaka, M., Ishii, A., Kanai, E., Funakura, M., & Watanabe, Y. (2013). Two types of mental fatigue affect spontaneous oscillatory brain activities in different ways. *Behavioral and Brain Functions: BBF*, *9*, 2. https://doi.org/10.1186/1744-9081-9-2

Shiv, B., & Carmon, Z. (2005). Placebo Effects of Marketing Actions : Consumers May Get What They Pay For. *Journal of Marketing Research*, *42*, 383-393. https://doi.org/10.2139/ssrn.707541

Snyder, M., Schulz, R., & Jones, E. E. (1974). Expectancy and apparent duration as determinants of fatigue. *Journal of Personality and Social Psychology*, *29*(3), 426-434. https://doi.org/10.1037/h0035917

Stoll, F. M., Wilson, C. R. E., Faraut, M. C. M., Vezoli, J., Knoblauch, K., & Procyk, E.
(2016). The Effects of Cognitive Control and Time on Frontal Beta Oscillations. *Cerebral*
Cortex (New York, N.Y.: 1991), *26*(4), 1715-1732. https://doi.org/10.1093/cercor/bhv006

Tanaka, M., Ishii, A., & Watanabe, Y. (2014). Neural effects of mental fatigue caused by
continuous attention load : A magnetoencephalography study. *Brain Research*, *1561*, 60-66.
https://doi.org/10.1016/j.brainres.2014.03.009

Thayer, J. F., Ahs, F., Fredrikson, M., Sollers, J. J., & Wager, T. D. (2012). A meta-analysis
of heart rate variability and neuroimaging studies : Implications for heart rate variability as a
marker of stress and health. *Neuroscience and Biobehavioral Reviews*, *36*(2), 747-756.
https://doi.org/10.1016/j.neubiorev.2011.11.009

Thayer, J. F., Hansen, A. L., Saus-Rose, E., & Johnsen, B. H. (2009). Heart rate variability,
prefrontal neural function, and cognitive performance : The neurovisceral integration
perspective on self-regulation, adaptation, and health. *Annals of Behavioral Medicine: A*
Publication of the Society of Behavioral Medicine, *37*(2), 141-153.
https://doi.org/10.1007/s12160-009-9101-z

Thönes, S., Arnau, S., & Wascher, E. (2018). Cognitions about time affect perception,
behavior, and physiology—A review on effects of external clock-speed manipulations.
Consciousness and Cognition, *63*, 99-109. https://doi.org/10.1016/j.concog.2018.06.014

Thönes, S., Arnau, S., Wascher, E., & Schneider, D. (2021). Boosting working memory with
accelerated   clocks.   *NeuroImage*,   *226*,   117601.
https://doi.org/10.1016/j.neuroimage.2020.117601

Van Cutsem, J., Marcora, S., De Pauw, K., Bailey, S., Meeusen, R., & Roelands, B. (2017).
The Effects of Mental Fatigue on Physical Performance : A Systematic Review. *Sports*
Medicine (Auckland, N.Z.), *47*(8), 1569-1588. https://doi.org/10.1007/s40279-016-0672-0

van Doorn, J., van den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., Etz, A.,
Evans, N. J., Gronau, Q. F., Haaf, J. M., Hinne, M., Kucharský, Š., Ly, A., Marsman, M.,
Matzke, D., Gupta, A. R. K. N., Sarafoglou, A., Stefan, A., Voelkel, J. G., & Wagenmakers,
E.-J. (2021). The JASP guidelines for conducting and reporting a Bayesian analysis.
Psychonomic Bulletin & Review, *28*(3), 813-826. https://doi.org/10.3758/s13423-020-01798-5

Wager, T. D., Waugh, C. E., Lindquist, M., Noll, D. C., Fredrickson, B. L., & Taylor, S. F.
(2009). Brain mediators of cardiovascular responses to social threat : Part I: Reciprocal dorsal
and ventral sub-regions of the medial prefrontal cortex and heart-rate reactivity. *NeuroImage*,
47(3), 821-835. https://doi.org/10.1016/j.neuroimage.2009.05.043

Wascher, E., Rasch, B., Sänger, J., Hoffmann, S., Schneider, D., Rinkenauer, G., Heuer, H.,

& Gutberlet, I. (2014). Frontal theta activity reflects distinct aspects of mental fatigue.
Biological Psychology, *96*, 57-65. https://doi.org/10.1016/j.biopsycho.2013.11.010

Williams, E. L., Jones, H. S., Sparks, S., Marchant, D. C., Micklewright, D., & McNaughton,
L. R. (2014). Deception studies manipulating centrally acting performance modifiers : A
review. *Medicine and Science in Sports and Exercise*, *46*(7), 1441-1451.
https://doi.org/10.1249/MSS.0000000000000235

Yu, S., Mückschel, M., & Beste, C. (2022). Superior frontal regions reflect the dynamics of
task engagement and theta band-related control processes in time-on task effects. *Scientific*
Reports, *12*(1), 846. https://doi.org/10.1038/s41598-022-04972-y

Zhou, L., Wei, H., Zhang, H., Li, X., Bo, C., Wan, L., Lu, X., & Hu, L. (2019). The Influence
of Expectancy Level and Personal Characteristics on Placebo Effects : Psychological
Underpinnings. *Frontiers in Psychiatry*, *10*, 20. https://doi.org/10.3389/fpsyt.2019.00020
947