Title

Endothelial Trauma Depends on Surface Charge and Extracellular Calcium Levels

Authors

Jade H. Cleary[1], Kara J. Wentzel[1], Abigail J. Howard[1], Adrian M. Sackheim[1], Sophia H. Piffard[1], J. Cory

Benson[2], Grant Hennig[4], Dev Majumdar[3], Mohamed Trebak[2], Mark T. Nelson[4,5], Kalev Freeman[1,4]

Affiliations

Departments of [1]Emergency Medicine, [3]Surgery, and [4]Pharmacology, University of Vermont, Burlington

VT; [2]Department of Pharmacology and Chemical Biology and Vascular Medicine Institute, University of

Pittsburgh, Pittsburg PA; [5]Division of Cardiovascular Sciences, University of Manchester, Manchester UK

Corresponding Author

Kalev Freeman, Kalev.Freeman@med.uvm.edu, 802-338-1622

Sources of Support

NIH/NIGMS R35 GM144099, P20 GM125498 and P20 GM135007 (Core C: Cus- tomized Physiology

and Imaging Core); NIH/NHLBI R35 HL140027, R35 HL150778 and R01 HL166944; NIH/NIA/NINDS RF

NS128963, R01 NS110656, and R01 NS119971; NIH Office of the Director (S10-OD026843); European

Union Horizon 2020 Research and Innovation Programme Grant Agreement 666881; Leducq Foundation

Transatlantic Network of Excellence, Stroke-IMPaCT 19CVD01; the Totman Medical Research Trust.

Conflicts of Interest

There are no conflicts of interest for any of the authors.

Significance

In acute critical illness, the rapid collapse of vascular endothelial functions drives aberrant blood clotting and organ failure through mechanisms that are not understood. The recent breakthrough in transfusion medicine showing that administration of donor plasma improves survival of trauma patients has transformed the massive transfusion protocols used in surgical settings, but the sodium citrate used to prevent coagulation often produces significant and severe hypocalcemia. Here, we demonstrate that cytotoxic trauma factors that are elevated in the blood during resuscitation bind endothelial cell phospholipids, and that low Ca^{2+} exacerbates toxicity by increasing this interaction. These experiments lead to our surprising discovery of unexpected endothelial cell membrane theatrics that occur in response to injury, with direct visualization of fast plasma membrane movements and the release and reuptake of extracellular vesicles. These findings provide important insights into the nature of shock-induced endotheliopathy and highlight the potential cardiovascular risk associated with chelation-induced hypocalcemia during resuscitation.

Abstract

We hypothesized that cationic histones bind to endothelial cell membranes through electrostatic interactions with negatively charged phospholipids, and consequently, that low Ca^{2+} exacerbates toxicity by increasing binding and uptake of histones into cells. The role of the ubiquitous store-operated Ca^{2+} entry (SOCE) in histone responses is also unknown. Here, studying endothelial cells in surgical preparations and in culture, we observed clinically relevant histone concentrations, produced fast plasma membrane movements including vesiculation, blebbing, ruffling, and cellular collapse. The cell membrane theatrics observed were markedly different from the uniform pattern of exocytosis and blebbing produced by calcium overload with ionomycin. Interestingly, membrane permeabilization produced by histones (and not ionomycin) was transient and a subset of cells recovered membrane integrity within 1 hour. A role for SOCE in histone responses was ruled out by genetic ablation of the ORAI1/2/3 channel trio. Removal of extracellular Ca^{2+} prevented histone-induced intracellular Ca^{2+} overload while surprisingly exacerbating plasma membrane deformation. Conversely, decreasing the density of the negative charge surface by

adding calcium or multivalent cations gadolinium or increasing extracellular Ca^{2+} levels effectively

screened common membrane phospholipids from interactions with labeled histones and prevented

endothelial damage in cells exposed to histones. Collectively, these results indicate that low extracellular

Ca^{2+} levels enhance interactions between histones and endothelial cell membrane phospholipids to

increase cytotoxicity. Importantly, this supports the concept of aggressive Ca^{2+} repletion during

resuscitation to prevent hypocalcemia, stabilize the endothelial cell membranes and improve

cardiovascular recovery from shock.

Introduction

Cardiovascular shock secondary to trauma and sepsis is among the most common causes of

morbidity and mortality worldwide[1,2]. Over 60 trillion interconnected vascular endothelial cells (ECs)

connect all organs and regulate function in the body[3]. The sudden and devastating disruption of endothelial

function that occurs locally and in distant organs after a severe insult, disrupting blood clotting, barrier

function, and blood flow, known as shock-induced endotheliopathy, is a common denominator of acute

critical illness that drives progressive organ failure and mortality[4–6]. Despite the emerging consensus that

the release of cytotoxic histone proteins into the bloodstream, both from injured cells and activated

neutrophils in the form of extracellular traps, is the inciting event that drives acute endothelial injury in

trauma and severe infection[7–13], there are no proven treatments, and the fundamental mechanisms that

underlie endothelial cell responses to severe injury remain largely unknown.

The results of clinical trials showing that the rapid transfusion healthy donor plasma to in the early

prehospital setting (before arrival to the hospital) decreases biomarkers of endothelial injury and improves

the survival of severely injured patients has transformed the emergency care of surgical trauma, and

clinical protocols used at trauma centers for massive transfusion of blood products are evolving to these

changes. Hypocalcemia is a common complication during the resuscitation of critically injured patients,

resulting from factors including the infusion of blood products containing sodium citrate (which has an

anticoagulant effect by chelating Ca^{2+} ions), hypoperfusion, and decreased hepatic clearance of the citrate

due to impaired perfusion[13]. Low plasma Ca^{2+} levels in trauma are predictive of mortality and need for

multiple transfusions[14,15]. Despite the prevalence of hypocalcemia in trauma patients, there is controversy

regarding the appropriate thresholds, timing and extent of Ca^{2+} administration used for massive transfusion

protocols[16,17].

Here, we hypothesized that cationic histones interact electrostatically s with negatively charged

phospholipids, and consequently, that low Ca^{2+} exacerbates toxicity by increasing binding of histones to

cell membranes. We studied ECs both in culture and in surgically opened vascular preparations using live

cell video microscopy, to address the following questions: (1) Are histone-induced Ca^{2+} oscillations SOCE-

dependent? (2) Is Ca^{2+} influx necessary for histone-induced toxicity? (3) Do extracellular Ca^{2+} levels

modulate the membrane effects of histones? (4) To what extent are histone-induced membrane changes

dependent on the negative charge density of the local electric field near the cell surface?

Results

Gadolinium prevents histone-induced endothelial cell calcium events. To establish the contribution

of SOCE to Ca^{2+} elevations, we first employed a pharmacological approach using Gd^{3+}, a trivalent

lanthanide that is useful when applied at low concentrations (<10 µM) as a SOCE inhibitor[18]. Freshly

harvested murine mesenteric arteries, surgically opened to expose the endothelium, were loaded with a

fluorescent Ca^{2+} indicator and exposed to clinically relevant concentrations of histones (50 µg/mL) with

Ca2+ (1.2 mM) in the presence or absence of Gd^{3+} (10 µM) (Figure 1 A-C, Supplemental Videos 1 and 2).

Rapid (<5 minutes) and irregular increases in endothelial cell Ca^{2+} fluorescence were observed as

expected. Surprisingly, these effects were completely blocked if histones were applied following pre-

treatment, but not after, with Gd^{3+}.

These Ca^{2+} oscillations were recapitulated in HEK-293 cells loaded with the ratiometric Ca^{2+}

indicator Fura-2 and stimulated with increasing concentrations of histones. Ca^{2+} signals were present in

response to histones starting at concentrations of 25 µg/mL, and high amplitude, frequent calcium

oscillations were observed in response to 50 µg/mL of histones (Figure 1 D,E). As we previously

demonstrated in surgical vascular EC preparations[11], extracellular Ca^{2+} is necessary for histone-induced

Ca^{2+} events. When HEK-293s were exposed to histones (50 µg/mL) in a solution of 0 mM Ca^{2+}, no Ca^{2+}

activity was observed. After 5 minutes, the extracellular solution was replaced with 2 mM Ca^{2+} and a

significant increase in activity immediately after the extracellular Ca^{2+} levels were restored. Similar to our

findings in ECs, we found that pre-treatment with Gd^{3+} (10 µM) completely blocked histone-induced Ca^{2+}

signals in HEK-293s (Figure 1 D,F). Interestingly, post-treatment with Gd^{3+} (10 µM) within 5 minutes of

histone (50 µg/mL) exposure was unable to quench histone-induced Ca^{2+} influx (Figure 1 D,G). We

additionally tested a lower concentration of Gd^{3+} (5 µM) which should be effective at inhibition of SOCE-

dependent Ca^{2+} activity[18]. In contrast to the complete block achieved by 10 µM Gd^{3+}, this lower

concentration was unable to inhibit histone-induced Ca^{2+} increases in either pre-treatment or post-

treatment conditions (Supplemental Figure 1A,B).

Store operated calcium entry is not required for histone-induced calcium events. Since

pharmacological approaches to blocking SOCE, such as Gd^{3+}, can have non-selective effects on other

cation channels, we applied a complementary genetic approach[19]. ORAI channels are key mediators of

SOCE through their binding to STIM1, and knocking out all three ORAI homologs completely prevents

functional SOCE responses[19–21]. We therefore utilized ORAI triple knockout (TKO) HEK293 cells to

determine the role of SOCE in histone-mediated Ca^{2+} signals[18]. While the blockade of histone responses

by Gd3+ (Fig 1A-C) had suggested a role for ORAI channels, surprisingly, there was no apparent difference

in histone-induced Ca^{2+} activity between wildtype and ORAI-TKO HEK-293 cells (Figure 1D-G). These

results effectively rule out ORAI channels as a contributor to histone-induced Ca^{2+} influx in ECs, and forced

us to reassess the Gd^{3+} results. Since the inhibitory activity of Gd^{3+} for SOCE is in the nM range, with an

IC50 of less than 100nM[22], and 5 µM Gd^{3+} pre-treatment was insufficient to achieve a complete block of

histone-induced Ca^{2+} signals (Supplemental Figure 1), we therefore concluded that the Gd^{3+} effect must

be due to a mechanism of action independent of SOCE.

Histones bind and deform endothelial cell plasma membranes. Because histones-induced Ca^{2+} entry

was independent of ORAI channels, but blocked by 10 µM Gd^{3+}, we reasoned that the Gd^{3+} may be acting

via a surface charge screening effect. Multivalent cations such as Gd^{3+} can neutralize the negative charge

at the membrane surface, altering the local electric field near the membrane, thereby shielding the plasma

membrane lipids. Preventing histones from binding directly to the plasma membrane might prevent

localized membrane injury or pore-formation that allows entry of extracellular Ca^{2+}. To evaluate the impact of histones on plasma membrane structure and morphology in ECs, we performed electron microscopy on *en face* murine mesenteric arteries following stimulation with histones (50 µg/mL) for 30 minutes in HEPES-PSS. We found that untreated arteries appeared to have intact cell-to-cell junctions and a flat, uniform cell morphology. The plasma membranes appeared smooth and minimal extracellular debris or vesicles were noted (Figure 2A). In contrast, ex vivo ECs treated with histones showed significant alterations in cell morphology. Cells appeared to have retracted, causing a breakdown of cell-to-cell junctions and the exposure of retraction fibers. Numerous blebs and extracellular vesicles were also visible on histone-treated cells (Figure 2A).

Due to the dramatic membrane changes seen on electron microscopy, we next wanted to elucidate the dynamic effect of histones on the endothelial cell plasma membrane. The styryl pyridinium dye FM1-43, which fluoresces when inserted in lipid bilayers, was used to stain the plasma membranes of histone-stimulated ECs. Furthermore, FM1-43 can also reveal cell membrane disruption and permeability. While normally impermeable to cells, FM1-43 floods the cytoplasm and stains internal membrane structures of damaged cells[23]. We compared the membrane effects of histones to those of ionomycin at a concentration sufficient to both mobilize intracellular stores and produce sustained Ca^{2+} influx[24]. Both stimuli resulted in cellular blebbing and membrane permeability over a period of 40 minutes, but the pattern and characteristics of plasma membrane responses to these stimuli were markedly different (Supplemental Videos 3 and 4). The endothelial response to ionomycin was characterized by synchronized cytoplasmic filling of the membrane dye between 20 minutes to 30 minutes of exposure, culminating in 92% ($\pm$ 12%) of cells showing a 2-fold or greater increase in fluorescence at 40 minutes. In comparison, the endothelial response to histones also showed membrane dye internalization within 40 minutes of exposure, but in a different pattern. Most importantly, histone exposure resulted in extensive FM1-43 dye uptake in only a subset of ECs 34% ($\pm$ 7%) within 40 minutes at normal Ca^{2+} levels (1.2 mM). The timing of dye entry was also not uniform across cells, instead occurring in a few cells at a time over the course of 10 to 20 minutes. The peak timing of the membrane changes (~30-40 minutes) was much later than when we observed the Ca^{2+} surges (~3-5 minutes). We also noted that dye uptake originated from hotspots within a single cell

then spreads to the rest of the cytoplasm. After initial dye uptake in these hot spots, FM1-43 staining

spreads from the outer membrane to internal membrane components, until the entire cell cytoplasm is

rapidly filled with dye while the nucleus remains unstained. In the remaining ~ ⅔ of cells, there were either

no changes in cell permeability or dye entry was sequestered to hot spots within cell protrusions without

progressing further.

Close inspection of the video recordings demonstrated fast plasma membrane movements and

release of extracellular vesicles in response to both ionomycin and histone, but different patterns of

responses were evident. To compare the patterns of membrane changes, we present two dramatic

examples of fields of endothelial cells recorded over 1 hour with zoomed-in regions of interest, in order to

demonstrate specific features (Figure 3). The patterns of membrane movements caused by ionomycin

included extension of filopodia and blebbing (Figure 3A, Supplemental Videos 5, 6). Histone exposure

resulted in dynamic cell responses in a different pattern. Membrane movements produced by histones and

not ionomycin included cell rounding with exposure of retraction fibers and ruffling sheets of plasma

membrane at the edges of cells particularly at surfaces near adjacent cells (Figure 3B, Supplemental

Videos 7, 8). Histone exposure also resulted in an end-stage pattern of blebbing and cessation of cell

movements in some cells(Figure 3B, Supplemental Videos 9). Additional examples of histone-induced

plasma membrane ruffling and blebbing are also provided (Supplemental Videos 10, 11).

The patterns of EV release produced by histones and ionomycin was also different. Histone-

induced membrane fragments showed a wide range of sizes, including large fragments consistent with

apoptotic bodies (Supplemental Video 4). In contrast, the endothelial response to ionomycin was disticn

resulted in the synchronous appearance of punctate, stippled, rapidly vibrating symmetrical membrane

particles <5 μm. In some cases, extracellular vesicles were visualized being released by cells absorbed

by both adjacent and remote cells, surfing on nanotubes, or passed between cells directly across

connected lamellipodia (Supplemental Video 12).

Plasma membrane changes are not due to calcium overload. Based upon the earlier and more rapid

occurrence of histone-induced endothelial cell Ca^{2+} influx than plasma membrane changes, we considered

the possibility that Ca^{2+} overload was the trigger for subsequent membrane disruption. We previously

established extracellular Ca^{2+} is required for histone-induced intracellular Ca^{2+} entry into vascular

endothelial cells[11], so we reasoned that, if histone-induced endothelial cell Ca^{2+} overload drives cell

membrane changes, the removal of extracellular Ca^{2+} should prevent the membrane responses. We

therefore tested the impact of low Ca^{2+} on histone-induced endothelial cell membrane responses visualized

with FM1-43. Surprisingly, exposure to histones in low Ca^{2+} exhibited an exaggerated response compared

to normal extracellular Ca^{2+} conditions (1.2 mM). While histone-exposure caused significant dye uptake in

33.6% ($\pm$ 6.8%) of cells 1.2 mM Ca^{2+} levels, this value increased to 60.1% ($\pm$ 11.3%) of cells at 0 mM Ca^{2+}

concentrations. Similarly, the average highest change in fluorescence at 40 minutes increased from 6.56

AU at 1.2 mM Ca^{2+} to 18.72 AU at 0 mM Ca^{2+} (Figure 4). Furthermore, the pattern of cell membrane

deformation produced by histones under low Ca^{2+} was similar to that produced in normal Ca^{2+} containing

solutions, but with more extensive of membrane thickening, blebbing, and release of membrane fragments

(Figures 3-4).

Neutralizing cell membrane surface charge prevents histone interactions. We then investigated the

impact of high Ca^{2+} and Gd^{3+} on histone-induced membrane disruption in live ECs. Both high Ca^{2+} (12 mM)

and Gd^{3+} (10 µM) were protective. The fast plasma membrane movement and deformation produced by

histones was markedly diminished in the presence of high Ca^{2+}, and pretreatment with Gd^{3+} rendered

histone-stimulated cells practically indistinguishable from unstimulated cells. Quantitative spatiotemporal

analysis of video recordings showed that high Ca^{2+} and Gd^{3+} prevented FM1-43 uptake into cells produced

by histones (Figure 4).

To confirm our microscopy findings, we used flow cytometry to quantify endothelial cell FM1-43

update after histone exposure under conditions of normal or high Ca^{2+} compared to Ca^{2+} free conditions.

Flow cytometry results recapitulated the pronounced internalization of FM1-43 in a subset of cells following

exposure to histones (50 µg/mL) in both normal and low Ca^{2+} conditions. The mean fluorescent intensity

(MFI) of FM1-43 increased 1.23-fold following histone exposure in 1.2 mM Ca^{2+} and 1.33-fold in 0 mM

Ca^{2+}. Conversely, cells stimulated with histones in 12 mM Ca^{2+} showed a 1.36-fold decrease in MFI relative

to unstimulated cells at the same Ca^{2+} concentration (Figure 5A). These findings further support that

histone interactions with endothelial cell membranes can be modulated by extracellular Ca^{2+}

concentrations.

The pronounced effects of extracellular Ca^{2+} and Gd^{3+} on blocking histone cytotoxicity indicated

that histone-induced membrane effects depend on strength of the plasma membrane surface charge and

occur independently of SOCE and Ca^{2+} entry. Thus, we hypothesized that these cations protect ECs from

histones by directly neutralizing negatively-charged phospholipids and reducing electrostatic interactions

between histones and the plasma membrane. To test this, we measured the binding affinity of a tagged-

histone protein (GST-H3) to individual phospholipids found commonly in biological membranes in the

presence or absence of elevated Ca^{2+} or Gd^{3+}. Like prior studies[11], we observed that in the absence of

Ca^{2+}, histones strongly bind to anionic not zwitterionic membrane phospholipids, consistent with an

electrostatic interaction (Figure 5B,C). We found that Gd^{3+} (30 µM) completely blocked all histone-

phospholipid interactions, removing the preferential binding motifs. Similarly, high Ca^{2+} (12 mM)

significantly reduced histone binding to most phospholipids. Interestingly, the presence of Ca^{2+} appeared

to increase binding affinity for lysophosphocholine.

Phosphatidyl serine translocation to external leaflet occurs in response to histone stimulation even

in the absence of external calcium. Histone stimulation of endothelial cells promotes thrombosis in part

due to translocation of phosphatidyl serine (PS) to the membrane surface through a process that is

inhibited by blocking the calcium-activated phospholipid scramblase TMEM16F[26]. However, increased PS

on the external leaflet of the plasma membrane may also indicate programmed cell death, so it is not clear

if the Ca^{2+} overload produced by histones activated the scramblase directly to produce transient changes

in PS or if the cells were pre-apoptotic and destined for cell death. We were therefore interested to

determine whether histone-exposed PS translocation can be modulated by extracellular Ca^{2+}

concentrations, and the extent to which these cells also permeable to to the viability dye propidium iodide

(PI). Following 1 hour of histone exposure in HEPES-PSS containing varying concentrations of Ca^{2+} (0,

1.2, 12 mM), cells were stained with Annexin V for PS exposure and PI for permeability then evaluated by

flow cytometry. At a moderate histone dose of 50 µg/mL, cells at normal Ca^{2+} levels of 1.2 mM showed a

5-fold increase in PS^+/PI^- cells to 3.85% ($\pm$ 0.22%) relative to unstimulated cells at the same Ca^{2+} level.

This percentage dropped to 2.73% ($\pm$ 0.55%) at 0 mM Ca^{2+} and 1.16% ($\pm$ 0.33%) at 12 mM Ca^{2+}, which

was almost identical to unstimulated cells at all Ca^{2+} levels (Supplemental Figure 2). Similarly, 50 µg/mL

histones resulted in 8.40% ($\pm$ 1.69%) of cells that were double positive (PS^+/PI^+) at 1.2 mM Ca^{2+} which

represents a 2-fold increase relative to unstimulated cells. Like the PS^+/PI^- findings, this percentage

dropped to 6.19% ($\pm$ 2.97%) at 0 mM Ca^{2+} concentrations. Notably, unstimulated cells in the 12 mM Ca^{2+}

condition showed an elevated PS^+/PI^+ percentage at 8.18% ($\pm$ 0.28), matching the respective histone-

stimulated population of 8.29% ($\pm$ 2.76%). Thus, high Ca^{2+} levels were protective from histone-induced

cytotoxicity, but caused increase PS exposure and PI uptake at baseline (Supplemental Figure 2).

Since 50 µg/mL of histones did not cause dramatic changes in PS exposure or PI uptake, we next

tested a higher dose of 100 µg/mL histones which we have previously shown was sufficient to produce PS

translocation in endothelial cells[26]. At this higher dose, 32.15% ($\pm$ 7.37%) of cells were PS^+/PI^- at 1.2 mM

Ca^{2+} which dropped to 24.20% ($\pm$ 4.95%) at 0 mM Ca^{2+} and 3.52% ($\pm$ 1.32%) at 12 mM Ca^{2+}. When

compared to unstimulated controls at the various Ca^{2+} levels, these changes represent 16-fold, 14-fold,

and 1.5-fold increases in PS^+/PI^- percentages for 1.2 mM Ca^{2+}, 0 mM Ca^{2+}, and 12 mM levels, respectively.

Similarly, the PS^+/PI^+ percentages were 13.38% ($\pm$ 3.02%), 12.93% ($\pm$ 1.82%), and 4.11% ($\pm$ 1.05%)

following 1 hour of 100 µg/mL histone stimulation at 12 mM, 0 mM, and 12 mM Ca^{2+} concentrations,

respectively. Interestingly, since the percentage of PS^+/PI^+ cells in the unstimulated 12 mM Ca^{2+} condition

was elevated from 5.51% ($\pm$ 1.01%) in normal Ca^{2+} to 8.14% ($\pm$ 4.13%), the histone-stimulated cells at the

12 mM Ca^{2+} level appeared to have less cell death than their respective control. These results indicate that

both moderate and high doses of histones lead to both PS exposure and PI uptake in a subset of cells.

Membrane permeabilization produced by histones is reversible. Although these findings show that

high doses of histones can kill ECs, we also considered the possibility that a subset of PS-flipped and

permeabilized cells could recover their membrane integrity. Hence, we next sought to evaluate cell

recovery following histone-induced injury using a novel strategy involving live-cell imaging with different

colored viability dyes. Live ECs were initially stained with Hoechst nuclear dye and SYTOX Green to

determine baseline viability. The adherent monolayers were then washed multiple times to remove

unbound dye and subsequently incubated in HEPES-PSS containing no stimulant, histones (100 µg/mL),

or ionomycin (10 µM). Following 1 hour of stimulation, the cells were stained and imaged with SYTOX

Green again to determine post-stimulation viability. Finally, the cells were recovered in fresh DMEM with

10% FBS at 37 °C and 5% CO_2 for 3 hours then stained with red SYTOX AADvanced for post-recovery

viability (Figure 7A). Through this experimental design, we were able to use microscope filters to

distinguish between 4 distinct cell phenotypes: 1. Intact (blue alone), 2. Late Permeability (blue and red),

3. Sustained Permeability (blue, red, and green), and 4. Recovered (blue and green).

In the histone-treated group, a subset of cells exhibited early membrane permeability to SYTOX

Green, but excluded the red SYTOX AADvanced, indicating regained membrane integrity (Figure 7C,E).

In contrast, cells treated with ionomycin, a positive control, demonstrated sustained and uniform

permeability to both SYTOX Green and red SYTOX AADvanced, consistent with irreversible membrane

damage. Panel D shows cumulative permeability over time, based on the combined percentage of cells

positive for either dye at each timepoint. The increase in permeability observed in the histone group may

be partially influenced by variability in imaging fields across timepoints. Panel E quantifies the distribution

of cells in each condition according to dye uptake classification (as outlined in Panel A): SYTOX Green

only ("recovered"), SYTOX Green and red SYTOX AADvanced double-positive ("sustained permeability"),

or red SYTOX AADvanced only (not observed). At the final timepoint, 15.0% ($\pm$ 2.9%) of histone-treated

cells were SYTOX Green-positive only, indicating recovery. Conversely, 41.5% ($\pm$ 10.7%) were double-

positive, indicating sustained permeability. In the ionomycin-treated condition, 94.8% ($\pm$ 2.6%) of cells were

double-positive, and 0% ($\pm$ 0%) of cells had recovered membrane integrity. These results indicate that,

although histones are highly cytotoxic, a subset of permeabilized endothelial cells can repair their plasma

membranes and recover from histone exposure.

Discussion

The identification of endotheliopathy as a driver of coagulopathy and poor outcomes in shock states

of sepsis (e.g., bacterial infection, COVID-19) and trauma (e.g., hemorrhage, brain or multisystem injury,

burns)[9,27,28] was an important advance in critical care medicine. The paradigm of SHINE provides a unifying

pathophysiologic mechanism that explains how different shock states converge to produce similar patterns

of endothelial activation and injury[4,5]. In this study, we investigated mechanisms of histone-induced endotheliopathy that demonstrate the importance of extracellular Ca^{2+} in shielding electronegative endothelial cell surfaces from histone cytotoxicity. We have previously demonstrated that histones initiate large EC Ca^{2+} oscillations that are pro-inflammatory, pro-thrombotic, and promote barrier breakdown[11,29,30]. Store-operated calcium entry (SOCE) is the most ubiquitous Ca^{2+} entry in non-excitable cells, such as ECs, that is triggered in response to a variety of stimuli such as vascular endothelial growth factor (VEGF) and thrombin[31]. This led to our initial hypothesis that SOCE contributes to histone-induced Ca^{2+} influx; however, our experimental results did not support this model. We then considered an alternate hypothesis: histones may induce Ca^{2+} entry by binding and deforming EC membranes, independently of SOCE. In addition, we aimed to understand the extent to which intracellular Ca^{2+} overload drives histone toxicity[32–34]. We also considered the possibility that removal of Ca^{2+} would increase exposure of negatively charged phospholipids on EC membranes to interact with cationic histones while surface charge screening with the addition of Ca^{2+} or multivalent cations would decrease this interaction. We report 4 novel findings: 1) Activation of plasma membrane SOCE channels ORAI1/2/3 either directly or indirectly through release of intracellular Ca^{2+} stores does *not* contribute to histone-induced cytosolic Ca^{2+} influx; 2) histones bind to and deform ECs, producing fast irregular plasma membrane Ca^{2+} flickers or spikes in a pattern distinctly different from the regenerative Ca^{2+} oscillations induced by physiological agonists[19] or the large plateaus caused by ionomycin-induced Ca^{2+} influx; 3) Ca^{2+} influx is *not* required for histone-induced endothelial cell deformation, and 4) histone-induced membrane permeability depends on electrostatic interactions with negatively charged membrane phospholipids which are exacerbated by low Ca^{2+} and blocked by mM Ca^{2+} or μM Gd^{3+}. These findings provide important insights into the nature of SHINE and highlight the potential cardiovascular risk associated with hypocalcemia during resuscitation.

The release of intracellular Ca^{2+} stores and subsequent activation of ORAI channels is an important mechanism for endothelial and immune cell signaling. Our initial experiments which showed that Gd^{3+} blocked histone-induced endothelial cell Ca^{2+} influx (Figure 1) suggested SOCE may also play a role in histone responses, as Gd^{3+} is a known ORAI channel antagonist[22]. However, we were surprised to discover that the ablation of all three SOCE channels (ORAI1, ORAI2, and ORAI3) had no impact on histone-

induced endothelial cell Ca^{2+} overload (Figure 2). This effectively ruled out SOCE as a mechanism of

histone-induced Ca^{2+} signaling. Moreover, these results are consistent with recent observations that

histones and LPS activate ECs by distinct, additive pathways[30].

Our observation that histone-induced Ca^{2+} entry was independent of ORAI but blocked by Gd^{3+}

provided support for an alternate model in which histones allow direct entry of extracellular Ca^{2+} by creating

pores and deforming the outer membrane. We visualized the time course and extent of endothelial cell

membrane deformation produced by histone exposure at high spatial and temporal resolution for the first

time using video imaging of endothelial cell plasma membranes labeled with the fluorescent dye FM1-43.

This dye has been used to study endothelial cell membrane injury produced by other insults such as laser

injury[23,35], but we are not aware of prior studies using this dye to study histone responses. We found that,

indeed, histones deform endothelial cell membranes and produce a large increase in membrane

fluorescence within cells. Membrane dye uptake after histone exposure was statistically significant,

measured both by spatiotemporal analysis of video results and by separate flow cytometry experiments.

Utilizing both electron microscopy and live cell confocal microscopy, we identified novel patterns of

membrane dynamics in response to histones (Figure 3). First, we were surprised by the dramatic and fast

plasma membrane movements that histones produced, which included examples of 1) membrane ruffling,

2) blebbing of the cell surface, 3) release of cell membrane fragments, 4) retractions of lamellipodia, and

5) cellular collapse with rapid membrane dye internalization. Second, over a period of 40 minutes, only a

subset of approximately one-third to one-half of visible cells in each experiment responded to histones

while other cells appeared unaffected. Moreover, the pattern of membrane response even at the

subcellular level was not uniform, with membrane changes initially isolated to "hot spots" which spread

inward over several minutes until the entire cell rapidly filled with dye. In the remaining cells, there were

either no changes in cell permeability or dye entry was sequestered to hot spots within cell protrusions

without progressing further. This heterogeneity is consistent with other studies of histone responses in both

cultured cells and native endothelium, in which only a fraction of cells showed Ca^{2+} entry or PI uptake after

histone exposure[11]. Endothelial cell heterogeneity, even within a single blood vessel, has also been

recognized in other systems; for example, only a subset of aortic ECs respond to laminar flow[36].

Third, we were also surprised to find that Ca^{2+} is not required for histone-induced endothelial cell

deformation. Intracellular Ca^{2+} overload produced by any means, whether by SOCE or disruption of

membrane integrity, can destabilize cell membranes and lead to activation of cell death pathways[37]. The

peak timing of the histone-induced membrane changes (30-40 minutes) was later than when we observed

the Ca^{2+} surges (3-5 minutes). However, exposure to ionomycin, which elevates Ca^{2+} to a maximum level,

produced a different pattern of membrane activation from that of histones. Cells that are activated by

ionomycin did not show extensive membrane ruffling or retractions of lamellipodia, that we observed with

histone exposure. Rather, the endothelial response to ionomycin was characterized by a dramatic,

synchronous release of symmetrical (~5 μm diameter) membrane fragments across all cells. This pattern

appears consistent with exocytosis of preformed vesicles which likely represented Ca^{2+} dependent

exocytosis of Weibel-Palade bodies[38,39]. We also visualized extracellular vesicles being re-absorbed by

both adjacent and remote cells; extracellular vesicles were also passed between cells directly across

connected lamellipodia (Supplemental Video 12).

Surprisingly, exposure to histones in low Ca^{2+} exhibited an exaggerated response compared to

normal Ca^{2+} conditions (Figure 5). The patterns of dye entry and cellular deformation produced by histones

under low Ca^{2+} conditions matched those of normal Ca^{2+} but they were more extensive. The extent of

membrane thickening, blebbing, release of membrane fragments appeared visibly more extensive. This

was unexpected, because removal of extracellular Ca^{2+} prevents histone-induced endothelial cell Ca^{2+}

events[11] so it seemed plausible that Ca^{2+} overload would contribute to membrane effects. However, these

results were additionally confirmed by flow cytometry using FM1-43. Thus, low extracellular Ca^{2+}

concentrations make ECs more susceptible to histone-mediated membrane permeability.

In contrast, we also observed that elevated levels of extracellular Ca^{2+} could entirely block

membrane permeability in ECs. We also found that both high Ca^{2+} or Gd^{3+} blocked binding of GST-tagged

histone H3 to various membrane lipids, indicating that these compounds can interfere with histones binding

to the plasma membrane. Interestingly, only μM concentrations of Gd^{3+} were required to completely block

membrane permeability following histone exposure. In comparison, there was not a substantial block

observed by 1.2 mM Ca^{2+} levels which indicates the vast difference in neutralizing ability between these

two compounds. While both are highly cationic, Gd^{3+} binds more readily and strongly due to its higher

charge and greater charge density. These results support a model in which cationic histones bind the

negatively charged phospholipids in the cell surface through electrostatic interactions exacerbated when

during hypocalcemia due to loss of surface charge screening by calcium.

Finally, we found that high doses of histones lead to exposure of PS and uptake of viability dyes

after 1 hour. These findings align with prior literature on the endothelial cell toxicity of histones and their

ability to increase PS and activate pro-coagulatory changes [7,8,11,40,26]. Most notably, the exposure of PS

not only indicates endothelial activation, but could also represent a positive feedback loop in which histones

are more readily able to bind to the membranes of activated or pre-apoptotic cells. Since PS is a highly

negatively charged phospholipid that avidly binds histones (Figure 5), its presence on the outer leaflet of

the plasma membrane may cause pro-coagulatory or pre-apoptotic cells to become even more highly

susceptible to histone binding.

In addition to these findings, we also present novel evidence that a subset of endothelial cells can

recover from histone exposure. Although only a subset of histone-treated cells appeared to recover

membrane integrity (~15%) after 3 hours, this population is notable given that ionomycin-treated cells

consistently showed 0% recovery across all trials. These observations point to the existence of functionally

distinct subpopulations within the histone-treated group. It seems that histones may induce severe,

irreversible damage in a subset of highly susceptible cells while others experience a more moderate,

transient injury that permits membrane repair. Interestingly, a similar mosaic pattern of endothelial cell

calcium responses within a monolayer was reported in response to shear stress supporting the emerging

concept that a distinct subpopulation of cells within endothelial monolayers serve as sentinals that detect

and respond to local environmental cues[36].

In conclusion, our studies demonstrate that low Ca^{2+} levels enhance the interactions between

histones and endothelial cell membranes, leading to increased cytotoxicity. Conversely, elevated Ca^{2+}

levels – or treatment with agents that alter provide surface charge screening effects such as Gd^{3+} – block

histone-phospholipid interactions and mitigate endothelial damage. The effects of Ca^{2+} and Gd^{3+}, and lack

of effect with knocking out ORAI channels, support a model in which histone binding to cells occurs

primarily due to electrostatic interactions with negatively charged membranes.

Intravenous Ca^{2+} salts are used clinically, including during cardiac resuscitation and in

hyperkalemia. The stabilizing effects of Ca^{2+} on nerve and cardiac muscle cell membranes are usually

explained by the surface charge theory, in which higher Ca^{2+} levels decrease the density of the negative

charge surface near the sodium channel of myelinated nerve fibers, making it harder for sodium ions to

enter the cell and trigger an action potential[41,42]. For all cell types, the plasma membrane surface itself is

more electronegative than the lipid surface of cellular organelles, as shown by measurement of the binding

of prenylated cation probes[43]. That is because electric fields generated by cells depend not only on lipids

but also on ionizable components embedded in the external membrane surface. In vascular endothelial

cells, sugar moieties of the specialized glycocalyx confer additional net negative charge to the outer

surface[44]. This is critical to physiologic regulation of blood clotting, because the strong electronegative field

of the inner surface of blood vessels serves to repel the negatively charged circulating platelets and blood

components. In fact, early experiments showed that reversal of transmural potential across the aorta

produced gradual intravascular thrombus produces in complete aortic occlusion within 9 days, and

resulting biophysical models of charge characteristics were important in the development of vascular grafts

and surgical procedures[45,46]. However, the electric field generated by the net negative surface charge of

the vascular endothelium also serves to recruit extrinsic cationic proteins such as cytotoxic histones. We

show here that changes in the endothelial cell surface charge produced by elevated Ca^{2+} or by the addition

of Gd^{3+} to an extent sufficient to decrease the density of negative charge surface can screen membrane

phsopholipids from attracting cytotoxic histone proteins. This conclusion is supported by prior findings on

membrane dynamics and the mechanisms of histone cytotoxicity. For example, some proteins can bind to

the curved surface of cell membranes and thereby bend them, with resulting membrane deformations that

indirectly attract each other, producing membrane invaginations that lead to vesiculation[47]. This

phenomenon may partially explain the mode of action of histones. It is well known that histones are highly

cationic proteins that bind negatively charged phospholipids in bilayers and cell membranes[40,48–51], and

direct interactions between histones and cell membranes have been implicated in their toxic effects[8,52].

Recently, the globular domain of histones alone was shown to deform and lyse the lipid membranes of

liposomes and cultured ECs[40].

While the evidence strongly supports a model in which histone interacts with lipid bilayers directly,

independent of protein factors, the differences in responses among various cell types has not been

explained. For example, neurons are relatively insensitive to histones, compared to macrophages or ECs[53].

Our findings do not eliminate the possibly that extracellular histones have other important roles in innate

immune responses, for example, neutrophil-extracellular traps can serve to tether neutrophils to vascular

walls by binding von Willebrand factor[10,54]. Additionally, it has been shown that downstream immune or

pyroptotic responses can vary due to histone post-translational modifications, composition, DNA content,

and the cell type[8,48,52,55–57]. Prior literature has also implicated toll-like receptors 2 and 4 (TLR2 and TLR4),

the purinergic receptor P2X7, the receptor for advanced glycosylation end-products (RAGE), and/or direct

interaction between histones and membrane phospholipids[11,40,48,52,53,55,58–65]. It seems likely that histones

damage cells through multiple pro-inflammatory mechanisms, creating a cacophony of danger signals

which ultimately orchestrate cell death. Only through elucidating these synergistic pathways can we

understand why some ECs are susceptible to histones while others can withstand or even recover.

Our findings have clinical relevance to coagulopathy and endotheliopathy in shock and trauma.

Hypocalcemia occurs frequently during trauma resuscitation due to transfusion of citrated blood

products[14,15], but clinical protocols for massive transfusions diverge, with some recommending routine Ca^{2+}

administration while others recommend supplementation only as needed[16,17]. Moreover, it is important to

consider that, while we focused on ECs, other vascular cells are also sensitive to histones[63,52,56,58]. Future

studies will need to evaluate whether the previously reported histone toxicity to vascular smooth muscle

cells, erythrocytes, and platelets may also be increased in the presence of low Ca^{2+} conditions[63,64,64]; this

might have important implications in the context of resuscitation and massive transfusion protocols. In sum,

our results suggest that one of the dangers of untreated hypocalcemia is its potential to amplify histone-

induced endothelial injury, underscoring the importance of early and aggressive Ca^{2+} correction during

resuscitation. The development of standardized protocols to optimize Ca^{2+} administration in critically

injured patients has the potential to reduce the incidence of endotheliopathy and improve outcomes.

Materials and Methods

Animal Husbandry. Mice were kept on a 12 h light/dark cycle with ad libitum access to food and water and were housed in groups of five. All studies were conducted in accordance with the Guidelines for the Care and Use of Laboratory Animals (National Institute of Health) and approved by the Institutional Animal Care and Use Committee of the University of Vermont.

En Face Ca^{2+} Imaging. En face murine vascular endothelial preparations were utilized for imaging as previously described[11,30]. Briefly, adult male wildtype C57BL6j (C57) mice (Jackson Laboratories) were euthanized using 5% Isoflurane, 95% O2, and decapitated. The mesentery was immediately harvested and placed in 4°C HEPES-buffered physiological saline solution (HEPES-PSS: 10 mM HEPES, 134 mM NaCl, 6 mM KCl, 1 mM MgCl$_2$, 2 mM CaCl$_2$, 7 mM glucose, pH 7.4). Mesenteric arteries (100-150 μm internal diameter) were excised in sections of approximately 500 μm, cut longitudinally to expose the lumen, and pinned flat with the endothelial surface exposed en face for imaging. Vessels were loaded with green-fluorescent Ca^{2+} indicator (5 μM Cal-520 AM, Thermo Fisher Scientific) and incubated at 37 °C for 30 minutes prior to imaging

Scanning Electron Microscopy. Third order mesenteric arteries (~200 μm diameter) from male C57BL6/j mice were cut open on one side longitudinally ("en face") and pinned down flat on a small silicone dish to expose the intact endothelium. Arteries were incubated with 50 μg/mL of unfractionated histones (Sigma-Aldrich, USA) for 30 minutes in HEPES-PSS (pH 7.4) at 37 °C. The arteries were then fixed in paraformaldehyde (2%)-glutaraldehyde (2.5%) cacodylate buffer solution (0.1 M; pH 7.2) for 24 h, washed with cacodylate buffer, post-fixed in a solution of 2.5% potassium ferrocyanide for 1 h, and washed again in cacodylate buffer and ultrapure water. Sections were dehydrated in ascending grades of ethanol, subjected to critical point drying in CO$_2$ (Samdri-PVT-3D, Tousimis, Rockville MD), coated with 10 nm of pure gold in a vacuum sputter coater (Quorum EMS150R S, Electron Microscopy Sciences, Hatfield, PA) and studied in a direct mode using a JEOL 6060 scanning electron microscope (Oxford Instruments, UK).

Cultured cells. Human endothelial (EA.hy926, ATCC) and kidney epithelial (HEK293, ATCC) cells were cultured according to manufacturer instructions. Genetically engineered HEK 293 cells with a triple

knockout of ORAI1, ORAI2, and ORAI3 store-operated Ca^{2+} entry (SOCE) channels generated using

CRISPR-Cas9 gene editing as previously described[19].

Live cell Ca^{2+} imaging. Ca^{2+} events were recorded in cells before and after histone stimulation using the

Ca^{2+} indicators. HEK cells were loaded with Fura-2 and imaged with a plate reader. Cells were manually

identified and Ca^{2+} levels within each cellular region of interest were reported as the ratio of F_{340}/F_{380}. ECs

(EA.hy926 or mesenteric artery vascular preparations) were loaded with Cal-520 imaged using spinning

disk confocal microscopy. Ca^{2+} oscillations from movies were analyzed using automated custom software

(Volumetry, Grant Hennig) to create 2D and 3D prevalence maps allowing quantification of the numbers

of cells activated per field and the frequency, amplitude, and duration of Ca^{2+} events, as previously

described[30]. Ionomycin was applied at the conclusion of each experiment as a positive control which

maximally increased Ca^{2+} levels in all cells within the field of view[24].

Live cell membrane imaging. EA.hy926 cells were plated on glass coverslips and grown to ~70%

confluence. The coverslips were transferred to individual wells containing HEPES-PSS with no added Ca^{2+},

1.2 mM Ca^{2+}, or 12 mM Ca^{2+} as indicated. Membrane dye FM1-43 (2 µM) and Gd^{3+} (10 µM) – where

indicated – were added directly to the wells immediately prior to imaging. A baseline 3D Z-stack was then

acquired before the careful addition of 1 mL extra HEPES-PSS (vehicle control), 50 µg/mL histones or 10

µM ionomycin. Continuous 2D recordings were acquired at 1 frame per second for 40 minutes. Finally, a

second Z-stack was obtained. Individual cell cytoplasm ROIs were drawn manually in ImageJ, excluding

outer membrane signal at all timepoints. Fluorescence intensity was quantified for each cell as ΔF/F0 using

the following formula: $(F_t - F_{0min}) / F_{0min}$. For representative Z-stack images, .png files were loaded into

ImageJ and converted to 8-bit. Contrast was enhanced by lowering the maximum to 150 for each image

within ImageJ.

Live cell viability imaging. EA.hy926 cells were seeded at a density of 800,000 on 12 mm glass

coverslips and allowed to grow for 1-2 days prior to imaging. Hoechst 33342 (1 µg/mL) and SYTOX Green

(50 nM) were added to each well and incubated for 10 minutes at 37 °C in 5% CO_2. Coverslips were

washed three times with HEPES-PSS following staining. For baseline and post-stim timepoints, three fields

were imaged per coverslip, each consisting of a z-stack of 50 slices covering approximately 15-20 µm in

depth. Following baseline imaging, coverslips were transferred into stimulation wells containing either

HEPES-PSS control, histones (100 µg/mL) in HEPES-PSS, or ionomycin (10 µM) in HEPES-PSS. Cells

were incubated at 37 °C for 1 hour. At 50 minutes, SYTOX Green was added to the stimulation wells and

returned to the incubator. At the 1-hour mark, cells were imaged immediately (within 2-4 minutes),

acquiring three non-overlapping fields per coverslip. After imaging, coverslips were washed and transferred

into a recovery well pre-filled with 2 mL of media and returned to the incubator for a 3-hour recovery period.

After the 3-hour recovery in media, coverslips were removed, washed with HEPES-PSS, and placed in

fresh HEPES-PSS containing SYTOX AADvanced for 10 minutes before final imaging. For the final

timepoint, three fields per coverslip were imaged using a reduced stack depth of three slices (~15 nm).

Flow Cytometry. Cultured ECs were plated in 24-well tissue culture plates. Once cells were grown to

confluency, the media was aspirated off and replaced with HEPES-PSS buffer solution containing histones

as indicated. HEPES-PSS solutions contained no added Ca^{2+}, 1.2 mM Ca^{2+}, or 12 mM Ca^{2+} depending on

the condition. For membrane permeability assays, FM1-43 (2 µM) was added concurrently with histone

exposure and incubated with the cells for 40 minutes at 37 °C with 5%. The cells were then trypsinized,

centrifuged, and washed with phosphate buffered solution (PBS). For experiments evaluating cell viability,

cells were trypsinized and washed with PBS then resuspended in Annexin V binding buffer and stained

with Annexin V (Biolegend #640920) and Propidium Iodide (Thermo Fisher #2301111). Cells were

incubated at room temperature for 15 minutes prior to 1:5 dilution in binding buffer. Samples were run on

a flow cytometer (MACSQuant Analyzer 10, Miltenyi Biotec). Overall sample fluorescence was quantified

as the median fluorescent intensity (MFI); n of 3-4 for all conditions.

Protein-Lipid Binding Assay. Mega Lipid Strips (Echelon-Inc, Salt Lake City, UT) kits were used in

binding assays according to the manufacturer's instructions. Briefly, strips were blocked for one hour at

room temperature with 3% bovine serum albumin (BSA: Sigma-Aldrich, St. Louis, MO) in PBS-T

(phosphate buffered saline with 1% Tween-20). Strips were then incubated with Histone H3-GST (0.5

µg/mL; BPS Biosciences, San Diego, CA) for one hour at room temperature on an orbital shaker (170

RPM). Strips were then washed in PBS-T three times for 7 minutes each wash. Strips were then incubated

in 75 µL of a GST secondary antibody (Echelon-Inc, Salt Lake City, UT) for one hour at room temperature.

Strips were washed twice in PBS-T and twice in PBS for 5 minutes each. Strips were then incubated in 2

mL of TMB precipitating solution (Echelon-Inc, Salt Lake City, UT) for 2 minutes before imaging. Gd^{3+} (30

μM) was added during the blocking step prior to the addition of histones where indicated. Following

background subtraction, dot blot intensity was then quantified as area under the curve (AUC) using ImageJ

software.

Data analysis and statistics. GraphPad Prism software (ver. 10.4.0; GraphPad Software, USA) was

used for x–y graphing and analysis. Data were tested for normality using the Kolmogorov-Smirnov test.

Parametric t-test or ANOVA, or appropriate non-parametric alternatives were applied for comparisons

between groups, and differences were considered significant if $P < 0.05$.

Ca^{2+} Fluorescence (Fura-2) Intensity. Max peak intensity was quantified for each cell for the period of

stimulation. Since multiple measurements were taken from the same cell and compared across multiple

stimuli, matched values for each cell were stacked. For each trace, a two-way ANOVA was conducted to

compare the means across WT and ORAI TKO for each period of stimulation. Geisser-Greenhouse

correction was used for non-sphericity and Šidák correction was used to correct for multiple comparisons.

Spatiotemporal Analysis of Ca^{2+} Fluorescence (Cal-520). Data was quantified as the total area with Ca^{2+}

signals, normalized as a percent of the ionomycin response. Thus, each independent replicate generated

a percentage for baseline, Gd^{3+}, and histones. Two individual Friedman Tests for nonparametric, matched

data were conducted for the cultured and *en face* preps respectively.

Lipid Binding. The area under the curve (AUC) was quantified in ImageJ for each lipid dot. A two-way

ANOVA was used to compare histone-lipid binding for each lipid across different ionic conditions (0 mM

Ca^{2+}, 12 mM Ca^{2+}, 30 μM Gd^{3+}). Data points were stacked by replicate (n=3). Geisser-Greenhouse

correction was used for non-sphericity and Tukey correction was used to correct for multiple comparisons.

Membrane Imaging. Cell fluorescence (FM1-43) at 40 minutes was quantified as $\Delta F_{40min}/F0$. All individual

cell values were combined across replicates (n=3) and a Kruskal-Wallis test for non-parametric data was

performed. A threshold of greater than 1.0 $\Delta F_{40min}/F0$, representing 2-fold increase in fluorescence relative

to baseline, was used to calculate the percent of cells filled with dye at 40 minutes.

<u>Flow Cytometry.</u> MFI was quantified using the median fluorescent value for each histone replicate (n=3 or

4) and compared across ionic conditions (0 mM, 1.2 mM, 12 mM Ca^{2+}) using a Kruskal-Wallis test.

Recovery Imaging Analysis. Raw images were acquired using Fusion software and post-processed in

Imaris. Standard parameters were applied uniformly across all replicates to ensure consistency, including

four iterations of deconvolution and baseline subtraction using a 0.218 µm spatial threshold, as specified

in the software's baseline correction settings. Nuclear regions were identified based on the Hoechst 33342

signal. Using the Imaris classification tool, each nucleus was then assigned to a viability category – double-

positive (SYTOX Green[+]/SYTOX AADvanced[+]), SYTOX Green[+] only, SYTOX AADvanced[+] only, or

negative – based on fluorescence intensity and channel overlap within the defined nuclear region. This

approach ensured each cell was counted once and accurately categorized based on nuclear uptake alone.

References

1. Haagsma JA, Graetz N, Bolliger I, Naghavi M, Higashi H, Mullany EC, et al. The global burden of injury: incidence, mortality, disability-adjusted life years and time trends from the Global Burden of Disease study 2013. Inj Prev J Int Soc Child Adolesc Inj Prev. 2016 Feb;22(1):3–18.

2. Rudd KE, Johnson SC, Agesa KM, Shackelford KA, Tsoi D, Kievlan DR, et al. Global, regional, and national sepsis incidence and mortality, 1990–2017: analysis for the Global Burden of Disease Study. The Lancet. 2020 Jan 18;395(10219):200–11.

3. Aird WC. Phenotypic heterogeneity of the endothelium: I. Structure, function, and mechanisms. Circ Res. 2007 Feb 2;100(2):158–73.

4. Johansson PI, Stensballe J, Ostrowski SR. Shock induced endotheliopathy (SHINE) in acute critical illness - a unifying pathophysiologic mechanism. Crit Care Lond Engl. 2017 Feb 9;21(1):25.

5. Henriksen HH, McGarrity S, SigurÐardóttir RS, Nemkov T, D'Alessandro A, Palsson BO, et al. Metabolic Systems Analysis of Shock-Induced Endotheliopathy (SHINE) in Trauma: A New Research Paradigm. Ann Surg. 2020 Dec;272(6):1140–8.

6. Bunch CM, Chang E, Moore EE, Moore HB, Kwaan HC, Miller JB, et al. SHock-INduced Endotheliopathy (SHINE): A mechanistic justification for viscoelastography-guided resuscitation of traumatic and non-traumatic shock. Front Physiol. 2023;14:1094845.

7. Xu J, Zhang X, Pelayo R, Monestier M, Ammollo CT, Semeraro F, et al. Extracellular histones are major mediators of death in sepsis. Nat Med. 2009 Nov;15(11):1318–21.

8. Abrams ST, Zhang N, Manson J, Liu T, Dart C, Baluwa F, et al. Circulating Histones Are Mediators of Trauma-associated Lung Injury. Am J Respir Crit Care Med. 2013 Jan 15;187(2):160–9.

9. Johansson PI, Windeløv NA, Rasmussen LS, Sørensen AM, Ostrowski SR. Blood levels of histone-complexed DNA fragments are associated with coagulopathy, inflammation and endothelial damage early after trauma. J Emerg Trauma Shock. 2013 Jul;6(3):171–5.

10. Kolaczkowska E, Jenne CN, Surewaard BGJ, Thanabalasuriar A, Lee WY, Sanz MJ, et al. Molecular mechanisms of NET formation and degradation revealed by intravital imaging in the liver vasculature. Nat Commun. 2015 Mar 26;6:6673.

11. Collier DM, Villalba N, Sackheim A, Bonev AD, Miller ZD, Moore JS, et al. Extracellular histones induce calcium signals in the endothelium of resistance-sized mesenteric arteries and cause loss of endothelium-dependent dilation. Am J Physiol Heart Circ Physiol. 2019 01;316(6):H1309–22.

12. Shaw RJ, Abrams ST, Austin J, Taylor JM, Lane S, Dutt T, et al. Circulating histones play a central role in COVID-19-associated coagulopathy and mortality. Haematologica. 2021 Sep 1;106(9):2493–8.

13. Moore EE, Moore HB, Kornblith LZ, Neal MD, Hoffman M, Mutch NJ, et al. Trauma-induced coagulopathy. Nat Rev Dis Primer. 2021 Apr 29;7(1):30.

14. Magnotti LJ, Bradburn EH, Webb DL, Berry SD, Fischer PE, Zarzaur BL, et al. Admission ionized calcium levels predict the need for multiple transfusions: a prospective study of 591 critically ill trauma patients. J Trauma. 2011 Feb;70(2):391–5; discussion 395-397.

15. Moore HB, Tessmer MT, Moore EE, Sperry JL, Cohen MJ, Chapman MP, et al. Forgot calcium?
Admission ionized-calcium in two civilian randomized controlled trials of prehospital plasma for
traumatic hemorrhagic shock. J Trauma Acute Care Surg. 2020 May;88(5):588–96.

16. DeBot M, Sauaia A, Schaid T, Moore EE. Trauma-induced hypocalcemia. Transfusion (Paris). 2022
Aug;62 Suppl 1:S274–80.

17. MacKay EJ, Stubna MD, Holena DN, Reilly PM, Seamon MJ, Smith BP, et al. Abnormal Calcium
Levels During Trauma Resuscitation Are Associated With Increased Mortality, Increased Blood
Product Use, and Greater Hospital Resource Consumption: A Pilot Investigation. Anesth Analg. 2017
Sep;125(3):895–901.

18. Shinde AV, Motiani RK, Zhang X, Abdullaev IF, Adam AP, González-Cobos JC, et al. STIM1 controls
endothelial barrier function independently of Orai1 and Ca2+ entry. Sci Signal. 2013 Mar
19;6(267):ra18.

19. Yoast RE, Emrich SM, Zhang X, Xin P, Johnson MT, Fike AJ, et al. The native ORAI channel trio
underlies the diversity of Ca2+ signaling events. Nat Commun. 2020 May 15;11(1):2444.

20. Emrich SM, Yoast RE, Xin P, Arige V, Wagner LE, Hempel N, et al. Omnitemporal choreographies of
all five STIM/Orai and IP3Rs underlie the complexity of mammalian Ca2+ signaling. Cell Rep. 2021
Mar 2;34(9):108760.

21. Emrich SM, Yoast RE, Trebak M. Physiological Functions of CRAC Channels. Annu Rev Physiol.
2022 Feb 10;84:355–79.

22. Trebak M, Bird GSJ, McKay RR, Putney JW. Comparison of human TRPC3 channels in receptor-
activated and store-operated modes. Differential sensitivity to channel blockers suggests
fundamental differences in channel composition. J Biol Chem. 2002 Jun 14;277(24):21617–23.

23. Koerdt SN, Gerke V. Annexin A2 is involved in Ca2+-dependent plasma membrane repair in primary
human endothelial cells. Biochim Biophys Acta Mol Cell Res. 2017 Jun;1864(6):1046–53.

24. Morgan AJ, Jacob R. Ionomycin enhances Ca2+ influx by stimulating store-regulated cation entry
and not by a direct action at the plasma membrane. Biochem J. 1994 Jun 15;300 (Pt 3)(Pt 3):665–
72.

25. Pereira LF, Marco FM, Boimorto R, Caturla A, Bustos A, De la Concha EG, et al. Histones interact
with anionic phospholipids with high avidity; its relevance for the binding of histone-antihistone
immune complexes. Clin Exp Immunol. 1994 Aug;97(2):175–80.

26. Bonson G, Lambert AR, Sackheim AM, Howard AJ, Piffard SH, Lescieur-Garcia C, et al.
ENDOTHELIAL-SPECIFIC KNOCKOUT OF THE SCRAMBLASE TMEM16F IMPAIRS IN VIVO
CLOT FORMATION. Shock Augusta Ga. 2025 May 1;63(5):788–95.

27. Johansson PI, Henriksen HH, Stensballe J, Gybel-Brask M, Cardenas JC, Baer LA, et al. Traumatic
Endotheliopathy: A Prospective Observational Study of 424 Severely Injured Patients. Ann Surg.
2017 Mar;265(3):597–603.

28. Rahbar E, Cardenas JC, Baimukanova G, Usadi B, Bruhn R, Pati S, et al. Endothelial glycocalyx
shedding and vascular permeability in severely injured trauma patients. J Transl Med. 2015 Apr
12;13:117.

29. Villalba N, Sackheim AM, Lawson MA, Haines L, Chen YL, Sonkusare SK, et al. The Polyanionic
Drug Suramin Neutralizes Histones and Prevents Endotheliopathy. J Immunol Baltim Md 1950. 2023
Aug 15;211(4):648–57.

30. Piffard SH, Hennig GW, Sackheim AM, Howard AJ, Lambert A, Majumdar D, et al. DISTINCT
PATTERNS OF ENDOTHELIAL CELL ACTIVATION PRODUCED BY EXTRACELLULAR
HISTONES AND BACTERIAL LIPOPOLYSACCHARIDE. Shock Augusta Ga. 2024 Nov
1;62(5):728–35.

31. Abdullaev IF, Bisaillon JM, Potier M, Gonzalez JC, Motiani RK, Trebak M. Stim1 and Orai1 mediate
CRAC currents and store-operated calcium entry important for endothelial cell proliferation. Circ Res.
2008 Nov 21;103(11):1289–99.

32. Fujii T, Sakata A, Nishimura S, Eto K, Nagata S. TMEM16F is required for phosphatidylserine
exposure and microparticle release in activated mouse platelets. Proc Natl Acad Sci. 2015 Oct
13;112(41):12800–5.

33. Suzuki J, Fujii T, Imao T, Ishihara K, Kuba H, Nagata S. Calcium-dependent Phospholipid
Scramblase Activity of TMEM16 Protein Family Members*. J Biol Chem. 2013 May
10;288(19):13305–16.

34. Schmaier AA, Anderson PF, Chen SM, El-Darzi E, Aivasovsky I, Kaushik MP, et al. TMEM16E
regulates endothelial cell procoagulant activity and thrombosis. J Clin Invest. 133(11):e163808.

35. Ashraf APK, Gerke V. Plasma membrane wound repair is characterized by extensive membrane lipid
and protein rearrangements in vascular endothelial cells. Biochim Biophys Acta Mol Cell Res. 2021
Jun;1868(7):118991.

36. Hong SG, Ashby JW, Kennelly JP, Wu M, Steel M, Chattopadhyay E, et al. Mechanosensitive
membrane domains regulate calcium entry in arterial endothelial cells to protect against
inflammation. J Clin Invest. 2024 May 21;134(13):e175057.

37. Orrenius S, Zhivotovsky B, Nicotera P. Regulation of cell death: the calcium-apoptosis link. Nat Rev
Mol Cell Biol. 2003 Jul;4(7):552–65.

38. Zupancic G, Ogden D, Magnus CJ, Wheeler-Jones C, Carter TD. Differential exocytosis from human
endothelial cells evoked by high intracellular Ca(2+) concentration. J Physiol. 2002 Nov
1;544(3):741–55.

39. Erent M, Meli A, Moisoi N, Babich V, Hannah MJ, Skehel P, et al. Rate, extent and concentration
dependence of histamine-evoked Weibel-Palade body exocytosis determined from individual fusion
events in human endothelial cells. J Physiol. 2007 Aug 15;583(Pt 1):195–212.

40. Pan Y, Peng M, Tong M, He Y, Hao M, Gao HL, et al. The Globular Domain of Extracellular Histones
Mediates Cytotoxicity via Membrane Disruption Mechanism. J Biol Chem. 2024 Nov 28;108038.

41. Hille B. Charges and potentials at the nerve surface. Divalent ions and pH. J Gen Physiol. 1968
Feb;51(2):221–36.

42. Hille B, Ritchie JM, Strichartz GR. The effect of surface charge on the nerve membrane on the action
of tetrodotoxin and saxitoxin in frog myelinated nerve. J Physiol. 1975 Aug;250(1):34P-35P.

43. Eisenberg S, Haimov E, Walpole GFW, Plumb J, Kozlov MM, Grinstein S. Mapping the electrostatic
profiles of cellular membranes. Mol Biol Cell. 2021 Feb 1;32(3):301–10.

44. Schmidt EP, Yang Y, Janssen WJ, Gandjeva A, Perez MJ, Barthel L, et al. The pulmonary
endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. Nat
Med. 2012 Aug;18(8):1217–23.

45. Sawyer PN, Pate JW. Bio-electric phenomena as an etiologic factor in intravascular thrombosis. Am
J Physiol. 1953 Oct;175(1):103–7.

46. Sawyer PN. BIOELECTRIC PHENOMENA AND INTRAVASCULAR THROMBOSIS: THE FIRST 12
YEARS. Surgery. 1964 Nov;56:1020–6.

47. Reynwar BJ, Illya G, Harmandaris VA, Müller MM, Kremer K, Deserno M. Aggregation and
vesiculation of membrane proteins by curvature-mediated interactions. Nature. 2007 May
24;447(7143):461–4.

48. Kleine TJ, Lewis PN, Lewis SA. Histone-induced damage of a mammalian epithelium: the role of
protein and membrane structure. Am J Physiol. 1997 Dec;273(6):C1925-1936.

49. Pereira LF, Marco FM, Boimorto R, Caturla A, Bustos A, De la Concha EG, et al. Histones interact
with anionic phospholipids with high avidity; its relevance for the binding of histone-antihistone
immune complexes. Clin Exp Immunol. 1994 Aug;97(2):175–80.

50. Fürnrohr BG, Groer GJ, Sehnert B, Herrmann M, Voll RE. Interaction of histones with phospholipids--
implications for the exposure of histones on apoptotic cells. Autoimmunity. 2007 Jun;40(4):322–6.

51. Hariton-Gazal E, Rosenbluh J, Graessmann A, Gilon C, Loyter A. Direct translocation of histone
molecules across cell membranes. J Cell Sci. 2003 Nov 15;116(Pt 22):4577–86.

52. Silvestre-Roig C, Braster Q, Wichapong K, Lee EY, Teulon JM, Berrebeh N, et al. Externalized
histone H4 orchestrates chronic inflammation by inducing lytic cell death. Nature. 2019
May;569(7755):236–40.

53. Villalba N, Baby S, Cha BJ, Yuan SY. Site-specific opening of the blood-brain barrier by extracellular
histones. J Neuroinflammation. 2020 Sep 22;17(1):281.

54. Ward CM, Tetaz TJ, Andrews RK, Berndt MC. Binding of the von Willebrand factor A1 domain to
histone. Thromb Res. 1997 Jun 15;86(6):469–77.

55. Allam R, Scherbaum CR, Darisipudi MN, Mulay SR, Hägele H, Lichtnekert J, et al. Histones from
dying renal cells aggravate kidney injury via TLR2 and TLR4. J Am Soc Nephrol JASN. 2012
Aug;23(8):1375–88.

56. Vulliamy P, Gillespie S, Armstrong PC, Allan HE, Warner TD, Brohi K. Histone H4 induces platelet
ballooning and microparticle release during trauma hemorrhage. Proc Natl Acad Sci U S A. 2019
Aug 27;116(35):17444–9.

57. Marsman G, Zeerleder S, Luken BM. Extracellular histones, cell-free DNA, or nucleosomes:
differences in immunostimulation. Cell Death Dis. 2016 Dec 8;7(12):e2518.

58. Tsourouktsoglou TD, Warnatsch A, Ioannou M, Hoving D, Wang Q, Papayannopoulos V. Histones,
DNA, and Citrullination Promote Neutrophil Extracellular Trap Inflammation by Regulating the
Localization and Activation of TLR4. Cell Rep. 2020 May 5;31(5):107602.

59. Xu J, Zhang X, Monestier M, Esmon NL, Esmon CT. Extracellular histones are mediators of death
through TLR2 and TLR4 in mouse fatal liver injury. J Immunol Baltim Md 1950. 2011 Sep
1;187(5):2626–31.

60. Yang X, Li L, Liu J, Lv B, Chen F. Extracellular histones induce tissue factor expression in vascular
endothelial cells via TLR and activation of NF-κB and AP-1. Thromb Res. 2016 Jan 1;137:211–8.

61. Kleine TJ, Gladfelter A, Lewis PN, Lewis SA. Histone-induced damage of a mammalian epithelium:
the conductive effect. Am J Physiol. 1995 May;268(5 Pt 1):C1114-1125.

62. Kim J, Baalachandran R, Li Y, Zhang CO, Ke Y, Karki P, et al. Circulating extracellular histones
exacerbate acute lung injury by augmenting pulmonary endothelial dysfunction via TLR4-dependent
mechanism. Am J Physiol Lung Cell Mol Physiol. 2022 Sep 1;323(3):L223–39.

63. Semeraro F, Ammollo CT, Morrissey JH, Dale GL, Friese P, Esmon NL, et al. Extracellular histones
promote thrombin generation through platelet-dependent mechanisms: involvement of platelet TLR2
and TLR4. Blood. 2011 Aug 18;118(7):1952–61.

64. Semeraro F, Ammollo CT, Esmon NL, Esmon CT. Histones induce phosphatidylserine exposure and
a procoagulant phenotype in human red blood cells. J Thromb Haemost JTH. 2014 Oct;12(10):1697–
702.

65. Al-Aqtash R, Ross MS, Collier DM. Extracellular histone proteins activate P2XR7 channel current. J
Gen Physiol. 2023 Jul 3;155(7):e202213317.

Figure Legends

Figure 1. Pre-treatment with Gd^{3+} blocks histone-induced Ca^{2+} signals independent of ORAI channels. (A) En face murine arteries were stained with fluorescent Ca^{2+} indicator (Cal-520, 5 μM) and stimulated with histones (50 μg/mL) with or without Gd^{3+} (10 μM) pre-treatment- Representative, cumulative Ca^{2+} prevalence images from continuous fields of view (FOV) showing surgically-opened murine endothelium at baseline followed by 5 minutes of histone stimulation (top) or after pretreatment withGd^{3+} followed by 5 minutes of histone stimulation. (B) Summary of Ca^{2+} activity in en face cells at baseline compared to histone stimulation. (C) Summary of Ca^{2+} activity at baseline, after Gd^{3+} pre-treatment, and during subsequent histone stimulation. Ca^{2+} signal was quantified as percent of the FOV area with oscillating cells using a specialized standard deviation method, normalized to the ionomycin-induced maximal response. (D) Wildtype (WT, left) and ORAI triple knockout (TKO, right) HEK 293 cells were stained with Ca^{2+} indicator (Fura-2) and stimulated with histones (n=1). Ionomycin (10 μM) was added at the end of each experiment as a positive control. Representative traces of fluorescence over time in response to increasing concentrations of histones (top), pretreatment with Gd^{3+} (10 μM) followed by histones (50 μg/mL) (middle), and Gd^{3+} (10 μM) post-treatment (bottom). Quantification of maximum peak intensity for each cell following histone titration (E), Gd^{3+} pre-treatment (F), and Gd^{3+} post-treatment (G).

Figure 2. Cultured ECs exhibit morphological changes and FM1-43 dye internalization in response to histones and ionomycin over 40 minutes of exposure. (A) Electron microscopy of native endothelial cells exposed en face in murine blood vessels following exposure to histones (50 μg/mL) with arrows pointing to areas of interest, including potential Weibel Palade bodies, retraction fibers, and blebs or extracellular vesicles. (B) Representative images of cultured cells stained with membrane dye FM1-43 during exposure to buffered saline, histones (50 μg/mL), or ionomycin (10 μM). The same field of view is presented at baseline (0 minutes), 20 minutes, and 40 minutes after exposure. (C) Magnification of histone-stimulated and ionomycin-stimulated cells at 40 minutes, with arrows pointing to examples of bleb formation on cell membranes.

Figure 3. The patterns of plasma membrane movements produced by histones are different than those caused by Ca^{2+} overload with ionomycin. Examples of the distinct patterns of membrane

movements observed during video imaging of EC monolayers over a 60-minute time frame are presented as both microscopy images and cartoons, with accompanying video supplements. (A) Ca^{2+} overload with ionomycin caused similar changes in almost all cells across the field of view. The initial response consisted of rapid membrane dye incorporation into the cytoplasm concomitant with the formation of small punctate vesicular bodies located both intracellularly and in the bathing media. Following dye incorporation, numerous filopodia were formed that elongated outwards from the cell body. These filopodia were motile and not attached to the surface of the slide (A1: Supplemental Video 5). The formation of large blebs marked the final response of cells before plasma membrane integrity was irrevocably lost (A2: Supplemental Video 6). (B) Histone exposure produced a heterogenous, mosaic pattern in which some cells respond rapidly and dramatically while others appear largely unaffected (Supplemental Video 7). Dye uptake occurred in a patchy fashion in cells, often initiating at one or more regions before spreading throughout the cell body. Extracellular vesicles appear to be both released and reabsorbed. Unique to the histone response, membrane ruffling behavior was observed at the edges of some cells, with undulating movements and expansion of the membrane border away from the cell body (B1: Supplemental Video 8) whilst other cells underwent rapid retraction of the outer membrane (rounding), leaving behind retraction fibers that remain connected to the media surface and adjacent cells (B2: Supplemental Video 9). A subset of cells showed similar end-stage membrane movements with large amounts of blebbing and dye uptake (B3: Supplemental Video 10). FOV Width = 490μm. Sub panel width = 70 – 97 μm.

Figure 4. Histone-induced endothelial cell membrane deformation and lysis are exacerbated by low Ca^{2+} and blocked by high Ca^{2+} or Gd^{3+}. (A) Representative Z-stacks of cultured ECs after 40-minute exposure to histones (50 μg/mL) under conditions of low (0 mM), normal (1.2 mM), or high (12 mM) Ca^{2+}, or with normal Ca^{2+} following Gd^{3+} (10 μM) pre-treatment. (B) Representative traces of individual cell cytoplasmic fluorescence ($\Delta F/F0$) over 40 minutes for each experimental condition, quantified using ImageJ. (C) Final fluorescence ($\Delta F_{40min}/F0$) for each cell with replicates displayed separately (n=3) for each condition. *Kruskal-Wallis test; P<0.05.*

Figure 5. Histone-induced membrane permeabilization and membrane lipid binding can be modulated by extracellular cation concentrations. (A) Flow cytometry of histones on ECs shows

exacerbation of membrane permeabilization by low Ca^{2+} (0 mM) and block by high Ca^{2+} (12 mM).

Representative flow cytometry histograms showing FM1-43 fluorescence (PE) following incubation of

EA.hy926 cells with FM1-43 dye and 50 µg/mL histones or 10 µM ionomycin for 40 minutes in buffered

saline solution (HEPES-PSS) with zero added Ca^{2+} (0 mM), normal Ca^{2+} levels (1.2 mM), or high Ca^{2+} (12

mM). . The mean fluorescent intensity (MFI) was quantified using the median fluorescence for cells

exposed to 50 µg/mL histones at different Ca^{2+} concentrations. *Kruskal-Wallis test; P<0.05. (B)* Histone

H3 selectively binds to membrane bound lipids in a charge dependent manner. Representative images of

lipid binding strips exposed to histone H3 alone, in the presence of Gd^{3+} (30 µM), or in the presence of

Ca^{2+} (12 mM) (left). Schematic diagram of lipid identities, adapted from the manufacturer (Right; Echelon

Biosciences, Alabama, USA). (C) Summary data for histone H3 binding to cardiolipin, Phosphatidylinositol

4-phosphate (PtdIns(4)P), phosphatidic acid (PA), phosphatidylethanolamine (PE), Phosphatidylinositol

4,5-bisphosphate (PtdIns(4,5)P2), and phosphatidylserine (PS). phosphatidylcholine (PC), sphingomyelin,

and lysophosphocholine alone, or in the presence of Gd^{3+} (30 µM), or Ca^{2+} (12 mM). Binding was quantified

in ImageJ as Area Under the Curve (AUC). N=3 for each group. *Two-Way ANOVA with Bonferroni's*

Correction for Multiple Comparisons; P<0.05.

Figure 6. High dose histone exposure induces a pre-apoptotic phenotype in ECs, which is

enhanced by low Ca^{2+} and blocked by high Ca^{2+}. EA.hy926 cells were treated with histones (100 µg/mL)

in HEPES-PSS with 0, 1.2, or 12 mM Ca^{2+} for 1 hour, followed by Annexin V (PS) and Propidium Iodide

(PI) staining. (A) Representative flow cytometry histograms showing fluorescence of Annexin V (top) and

PI (bottom) for cells incubated in HEPES-PSS alone (blue) or with the addition of histones (pink). (B, C)

Median fluorescence intensity (MFI) for Annexin V and PI shown (n = 4). (D) Representative dot plots of

Annexin V and PI fluorescence. (E, F) Quantification of percent PS^+/PI^- cells and PS^+/PI^+ cells using the

gating strategy shown in D.

Figure 7. A subset of ECs with histone-induced membrane permeability can recover their

membrane integrity after 3 hours. (A) Schematic diagram of experimental design allowing the

determination of four cell phenotypes following histone (100 µg/mL) or ionomycin (10 µM) stimulation: 1.

Intact (Hoeschst$^+$), 2. Late Permeability (Hoeschst$^+$ SYTOX-AADvanced$^+$), 3. Sustained Permeability

(Hoeschst[+] SYTOX-AADvanced[+] SYTOX Green[+]), and 4. Recovered (Hoeschst[+] SYTOX Green[+]). (B)

Magnification of one representative field of view showing multiple permeability phenotypes following

histone stimulation for 1 hour and recovery for 3 hours. (C) Representative images from the post-recovery

timepoint for HEPES-PSS, histones, and ionomycin. Individual fluorescence channels are shown for

Hoeschst, SYTOX Green, and SYTOX AADvanced followed by a merge of all channels. (D) Cumulative

percentages of cells permeable to either SYTOX dye (SYTOX Green and/or SYTOX AADvanced) at

baseline, post-stimulation, and post-recovery for each treatment condition. (E) Percent of cells in each

phenotypic category at the post-recovery timepoint following stimulation with HEPES-PSS alone, histones,

or ionomycin.

Supplemental Figure Legends

Supplemental Figure 1. Low-dose Gd^{3+} pre-treatment only partially blocks histone-induced Ca^{2+}

oscillations in HEK-293 cells, independent of ORAI channels. Wildtype (WT, left) and ORAI triple

knockout (TKO, right) HEK-293 cells were stained with Ca^{2+} indicator (Fura-2) and stimulated with histones

(50 µg/mL, n=1). Ionomycin (10 µM) was added at the end of each experiment as a positive control. (A)

Representative traces of fluorescence over time in response to pretreatment with Gd^{3+} (5 µM) followed by

histones (50 µg/mL) for WT (left) and ORAI-TKO (middle). Quantification of maximum peak intensity for

each cell (right). (B) Representative traces of fluorescence over time in response to post-treatment with

Gd^{3+} (5 µM) followed by histones (50 µg/mL) for WT (left) and ORAI-TKO (middle). Quantification of

maximum peak intensity for each cell (right). (C) Representative traces of fluorescence over time in

response to histones (50 µg/mL) in 0 mM Ca^{2+} followed by 2 mM Ca^{2+} for WT (left) and ORAI-TKO (middle).

Quantification of maximum peak intensity for each cell (right). 2-way ANOVA for significance (n = 1 for

each experiment).

Supplemental Figure 2. Moderate dose histones induce a subtle, pre-apoptotic phenotype in ECs,

which is enhanced by low Ca^{2+} enhances and blocked by high Ca^{2+} blocks. EA.hy926 cells were

treated with histones (50 µg/mL) in HEPES-PSS with 0, 1.2, or 12 mM Ca^{2+} for 1 hour, followed by Annexin

V (PS) and PI staining. (A) Representative flow cytometry histograms showing fluorescence of Annexin V

(top) and PI (bottom) for cells incubated in HEPES-PSS alone (blue) or with the addition of histones (pink).

(B, C) Median fluorescence intensity (MFI) for Annexin V and PI shown (n = 4). (D) Representative dot

plots of Annexin V and PI fluorescence. (E, F) Quantification of percent PS^+/PI^- cells and PS^+/PI^+ cells

using the gating strategy shown in D.

Supplemental Video Legends

Supplemental Video 1. Histones induce Ca^{2+} influx in en face murine ECs. 3D representation of

cumulative Ca^{2+} oscillations over the 5 minute period after exposure to histones (50 µg/mL). Spatio-

temporal representation overlaid on 2D image of the same field of view with ionomycin-induced maximal

Ca^{2+} response.

Supplemental Video 2. Pretreatment with Gd^{3+} blocks histone-induced Ca^{2+} influx in en face murine ECs.

3D representation of cumulative Ca^{2+} oscillations after exposure to histones (50 µg/mL) in the presence of

Gd^{3+} (10 µM). Spatio-temporal representation overlaid on 2D image of the same field of view with

ionomycin-induced maximal Ca^{2+} response.

Supplemental Video 3. Video recording (25x) of ionomycin-induced endothelial cell membrane changes

visualized using the membrane dye FM1-43.

Supplemental Video 4. Video recording (25x) of histone-induced endothelial cell membrane deformation

visualized using the membrane dye FM1-43.

Supplemental Video 5. Full field of view with inset video of endothelial cell filopodia extrusion and

movement after application of ionomycin (30 minute recording: speed 96x).

Supplemental Video 6. Full field of view with inset video of endothelial cell membrane blebbing after

application of ionomycin (30 minute recording, speed 96x).

Supplemental Video 7. Full field of view with inset video of endothelial cell membrane ruffles after

application of histones (20 minute recording, speed 96x).

Supplemental Video 8. Full field of view with inset video of endothelial cell rounding after application of

histones (20 minute recording, speed 96x).

Supplemental Video 9. Full field of view with inset video of endothelial cell membrane blebbing after

application of histones (14 minute recording, speed 96x).

Supplemental Video 10. Video of histones-induced endothelial cell membrane ruffling (30 minute

recording, speed 96x).

Supplemental Video 11. Video of histone-induced endothelial cell membrane blebbing (30 minute

recording, speed 96x).

Supplemental Video 12. Video of ionomycin-induced release and reuptake of extracellular vesicles (30

minute recording, speed 96x).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 6

Figure 7