Rainwater-driven transport of matter and microbes from phyllosphere to soil in a temperate beech forest

Markus Krüger[1], Karin Potthast[2,3], Beate Michalzik[2,3], Alexander Tischer[2,3,4], Martina Herrmann[1,3,5*]

[1]Institute of Biodiversity, Friedrich Schiller University Jena, Jena, Germany; [2]Department of Soil Science, Friedrich Schiller University Jena, Jena, Germany; [3]German Center for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany, [4]ThüringenForst, Forestry Research and Competence Centre, 99867 Gotha, Germany, [5]Cluster of Excellence Balance of the Microverse, Friedrich Schiller University Jena, Jena, Germany,

*Corresponding author: martina.herrmann@uni-jena.de

Abstract

In forest ecosystems, the transport of organic matter including microbial cells by throughfall

and stemflow represents a central linkage between the phyllosphere and soil. However, the

extent and taxon-specificity of rainwater-mediated vertical transfer of microorganisms, and its

relevance for matter fluxes is sparsely understood. We investigated the rainwater-driven

transport of bacteria and its contribution to matter transport along bulk rainfall, throughfall,

stemflow and seepage flow paths from the phyllosphere to soil in a temperate deciduous

forest. Throughfall-mediated cell fluxes equalled those in seepage below the litter layer,

pointing to a substantial above-ground translocation of bacteria. Bacterial cells contributed up

to 11 % and 33 % to the exported total organic carbon and total nitrogen, respectively. Bacteria

affiliated   with   *Pseudomonadaceae*,   *Oxalobacteraceae*,   *Hymenobacteraceae*,   and

Sphingomonadaceae were differentially transmitted from the canopy and enriched in

throughfall and stemflow. 117 Amplicon Sequence Variants (ASVs) (1.7 %) occurred in all

rainwater flow paths, suggesting a vertical connectivity from the phyllosphere through the soil

that allows the continuous import, proliferation, and mobilization of selected bacterial taxa.

The genetic potential of the transported bacterial communities was strongly linked to organic

matter degradation particularly of carbohydrates and proteins. Our findings extend our

understanding of rainwater-mediated vertical matter transport in forest ecosystems and

highlight the contribution of microbial cells to this transport especially in above-ground

compartments. Strong shifts in community composition and abundance of the transported taxa

between throughfall, stemflow and soil seepage are linked to differences in source

communities but may also point to different mechanisms of cell detachment across

compartments.

Graphical abstract

1. Introduction

Rainwater represents an important carrier of various particles including organic matter and

microorganisms across long distances (Hu et al., 2018). Such wet deposition may harbor tens

of millions of bacteria m^{-2} d^{-1} (Reche et al., 2018) and other microorganisms such as archaea,

fungi and protists (Després et al., 2012; Sánchez-Parra et al., 2021), and thus plays a crucial

role in microbial dispersal within ecosystems. Microbes from atmospheric sources may

contribute directly to the microbiome of the plant phyllosphere (Ponette-González et al., 2020),

which exhibits a global surface area of 6.4 x 10^8 km^2 as microbial habitat (Vorholt, 2012).

During rain events, huge amounts of rainwater can pass the forest canopy and emerge as

throughfall and stemflow water fluxes (Levia et al., 2011). Throughfall represents that part of

precipitation which falls through the canopy, partially along leaf surfaces, while stemflow is

defined as the flow path of rainwater which runs down the stem (Sadeghi et al., 2020). In

temperate forests, both water flow paths can transport a substantial amount of organic matter

ranging between 0.2 – 200 kg C ha^{-1} y^{-1} (Ponette-González et al., 2020; Van Stan et al., 2017)

and bacteria in the range of 1.5 x 10^{16} cells ha^{-1} yr^{-1} (Bittar et al., 2018). As consequence, the

forest canopy may act as a considerable source of microorganisms, influencing the bacterial

composition of rainwater as it is transported down to the forest floor (Sadeghi et al., 2020; Van

Stan et al., 2017). A certain proportion of this rainwater further percolates through the soil as

seepage, containing mobile organic matter including microorganisms (Dibbern et al., 2014;

Krüger et al. 2021; Herrmann et al., 2023; Zhang et al., 2018), resulting in continued microbial

transport to subsoils and ultimately to the groundwater (Friesen, 2020).

In the phyllosphere, bacteria perform a variety of matter transformation processes such as

carbon- and nitrogen fixation, nitrification and methanol oxidation (Andrews and Harris, 2000;

Guerrieri et al., 2020, 2024) . Methanotrophic bacteria such as *Methylomonas* and

Methylobacter, Flavobacterium and *Novosphingobium* are common members of the

microbiome on tree leaf and bark surfaces (Iguchi et al., 2015; Jeffrey et al., 2021). In addition,

microbial extracellular enzymes which degrade hydrocarbons and secondary plant

metabolites can be exported from the phyllosphere by throughfall and stemflow (Mori et al.,

2019), contributing to exoenzymatic matter degradation in the soil. In contrast to litter fall in

autumn, which acts as a periodic mass transfer of organic matter, microorganisms, and their

degradation capacity to the forest floor, throughfall and stemflow represent season-

independent, high-frequency episodic flow events from the phyllosphere, ensuring

connectivity between above-ground and below-ground compartments throughout the year.

Assuming a certain fraction of the microbial cells remains viable during transport, rainwater

delivers air- and phyllosphere-derived bacteria which constitute an essential source of new

genetic material and functional potential to the sink communities. In fact, previous work

demonstrated that some bacteria are selectively washed off from leaves and stems and get

enriched along the water flow paths (Guerrieri et al., 2020). Stemflow creates funneling of

water along with matter and microbes which are rapidly transported along roots into the soil

(Carlyle-Moses et al., 2020; Tischer et al., 2020). In addition, rainwater provides fresh

particulate and biolabile dissolved carbon and nitrogen to the forest floor (Michalzik et al.,

2001; Van Stan and Stubbins, 2018), which can affect soil nutrient pools (Van Stan et al.,

2017) and soil microbial activity (Deng et al., 2017).

Previous studies investigated the microbial diversity and mass fluxes associated with aerosol

particles (Després et al., 2012), tree phyllosphere, throughfall and stemflow (Van Stan et al.,

2020). However, research is scarce on the quantity and structure of microbial communities

transported by rainwater, throughfall, and stemflow, and which fraction of the transported

matter originates from microbial cells and their functional potential across different

compartments including soil seepage. Seepage-mediated transport of microbes through soils

was previously studied at the Hainich Critical Zone Exploratory using lysimeters installed in a

beech forest (Krüger et al., 2021; Herrmann et al., 2023). In this study, we used zero tension

lysimeters along with bulk rainwater, throughfall and stemflow collectors (Levia et al., 2023;

Potthast et al., 2017) to investigate the abundance and community composition of bacterial

communities along the flow path from the phyllosphere to soil seepage, along with an

estimation of their functional potential and contributions to matter fluxes. We aimed to (1)

estimate bacterial fluxes and contributions to carbon and nitrogen transport in throughfall,

stemflow and seepage, comparing an early and late growing season rain event; (2)

characterize the bacterial composition of the respective compartments and their predicted

functional potential, and (3) assess core and shared communities for each intersection of

neighboring compartments to provide insights into the fraction of bacterial communities that

can be mobilized along the vertical flow path.

2. Materials and methods

2.1 Study site and sampling procedure

Sampling was conducted in spring and autumn on a managed deciduous temperate forest plot (N51°06'56.5' E10°24'45.2) and on an open area site (N51°07'63.5' E10°23'68.2') in the Hainich-Dün area comprising the Hainich Critical Zone Exploratory, Germany (**Figure 1**A). Forest stands are dominated by European beech (*Fagus sylvatica* L.) with minor admixtures of sycamore maple (*Acer pseudoplatanus* L.) and ash (*Fraxinus excelsior* L.), a common forest composition in the Hainich-Dün area. A nearby, extensively managed grassland site (open area) was used for measurements of bulk precipitation (rainwater). The climate is characterized by annual precipitation of 600 mm and mean air temperature of 9°C (Potthast et al., 2017).

Ten throughfall and four rainwater collectors with a sampling area of 314 cm^2 at approximately 1.2 m height were deployed at a forest and an open area plot 2 km apart, respectively (**Figure 1**B). Stemflow collars were installed at two tree individuals of each *F. sylvatica* and *A. pseudoplatanus* at 1.3 m height with a basal area in breast height between 346 to 740 cm^2. Rainwater, throughfall and stemflow plastic containers were cleaned and surface-disinfected with H_2O_2 prior to exposure and were placed in the field 1 to 2 days before two major rain events in May and September 2020, respectively. All three collector types were equipped with a 2 mm PE mash to avoid the introduction of large particles or insects during the collection period. Seepage was obtained from 16 zero-tension lysimeters with an area of 284 cm^2 accessing soil leachate which percolated the soil through the litter, and the upper 4 cm, 16 cm and 30 cm (Krüger et al., 2021). Details on lysimeter properties and installation were described by Potthast et al., 2017. Residual solution from the lysimeters was emptied before the rain events to obtain only freshly percolated soil leachate. Throughfall, stemflow and seepage solutions at the forest site and bulk rainwater at the pasture site were sampled on 13[th] May 2020 and 28[th] September 2020 after rain events of 12 and 19 mm, respectively (**Figure 1**C). Triplicate leaf samples of the phyllosphere from two individuals of *F. sylvatica* and *A.*

pseudoplatanus at low canopy height of 4 to 5 m were clipped off with ethanol cleaned scissors

and stored in autoclaved plastic container after each rain event. In addition, soil samples as

reference to the mobilized microbial communities were collected in triplicates close to the

lysimeters at 0 to 4 cm, 4 to 16 cm and 16 to 30 cm depths in May 2020. All samples were

stored in cooling boxes at approximately 8°C during transport and were immediately

processed at the laboratory within 2 h. Rainwater, throughfall, stemflow, and seepage samples

were weighed to determine the volume and 10 to 100 ml was used for determination of pH,

electrical conductivity, carbon, nitrogen, and elemental analysis. For molecular analyses,

water samples were filtered through 0.2 µm polyethersulfone membrane filters (Supor, Pall

Corporation, USA) and stored at -80°C. Filtrate was stored at -20°C for determination of

inorganic nitrogen. Phyllosphere samples in plastic bottles were treated with 1 min of mild

sonification in 250 ml NaCl buffer (0.15 M NaCl + 0.1 % Tween 20) to detach microorganisms

from leaf surfaces, followed by 20 min shaking at 100 rpm (Herrmann et al., 2021). Resulting

suspensions were filtered through 0.2 µm pore size filters and stored at -80°C until further

processing. The remaining leaves were dried at 50°C for 1 week to measure the dry weight.

Soil samples for molecular analysis were stored at -80°C while those for determination of dry

weight and pH were stored at 4°C.

2.2 Physicochemical analyses and calculation of matter fluxes

Rainwater, throughfall, stemflow, and seepage samples were analyzed for total organic

carbon (TOC), inorganic carbon (IC) and total nitrogen (TN) as well as dissolved organic

carbon (DOC) and total dissolved nitrogen (TDN) concentrations. Analysis involved filtration

through <0.45 µm prewashed membrane filters (cellulose acetate, Sartorius, Göttingen,

Germany)   for   dissolved   elements   and   catalytic   thermal   oxidation   and

NDIR/chemiluminescence detection of both non-filtrated and filtrated samples using a TOC-

VCPN/TNM-1 (Shimadzu, Duisburg, Germany) (Potthast et al., 2017; Visco et al., 2005).

Particulate organic carbon (POC) and total particulate nitrogen (TPN) were calculated by

subtracting the dissolved fractions from the total concentrations. Inorganic nitrogen (IN),

mainly consisting of NH_4^+ and NO_3^- was determined by standard colorimetric approaches

(Scheiner, 1974; Seelmeyer, 1954).

Carbon (TOC) and nitrogen (TN) fluxes per rain event were calculated from the concentration

of the respective parameters from water samples (rainwater, throughfall, stemflow, seepage)

multiplied by the respective sample volume and the collector area and were expressed in mg

C or N m^{-2}. Stemflow fluxes were calculated for the individual basal area of each tree specimen

and scaled up to the tree stand area in ha from both species.

Electrical conductivity and pH were determined using a WTW pH / Cond meter (pH Cond 3320

with SenTix81 and TetraCond, Weilheim, Germany) (Potthast et al., 2017). The soil water

content was determined by drying the samples over 48 h at 105°C. Soil pH was determined

from slurries prepared using sterile deionized water (ratio 1:2.5) and measured with a pH glass

electrode (Krüger et al., 2021).

2.3 DNA extraction, quantitative PCR, and amplicon sequencing

Microbial biomass from a filter half (leaves, rainwater, throughfall, stemflow, seepage) and

from 0.25 g of soil of each sample were extracted using the DNeasy PowerSoil kit (Qiagen,

Germany) following the manufacturer's protocol and stored at -20°C. Bacterial and archaeal

16S rRNA gene abundances were quantified with a CFX96 real-time PCR thermocycler (Bio-

Rad, USA) using primer pair Bakt_0341F/Bakt_799R (Chelius and Triplett, 2001; Klindworth

et al., 2013) which discriminates against chloroplast-derived 16S rRNA genes, and

Arch806F/Arch958R (DeLong, 1992; Takai and Horikoshi, 2000), respectively. Details on PCR

and cycling conditions are described elsewhere (Herrmann et al., 2021, 2012). Plasmids

containing cloned bacterial and archaeal 16S rRNA gene fragments were used to construct

qPCR standards, which were linear across the concentration range of 50 to 5 $\times10^8$ copies per

reaction. Gene abundances in rain, throughfall, stemflow and seepage were calculated as per

L, while gene abundance on leaf surfaces and soil was expressed as genes per g of dry

weight.

Library construction for 16S rRNA gene targeted amplicon sequencing assessing the bacterial

community composition across samples was conducted with the same primer pair as for

qPCR, targeting the V3-V4 region of the bacterial 16S rRNA gene. Reactions were performed

with 1 ng DNA template per sample in 15 µl volume reactions under previously reported

conditions (Herrmann et al., 2021). Subsequent steps are described in Krüger et al. (2021),

and sequencing was performed on an Illumina MiSeq platform using v3 chemistry.

2.4 Sequence processing and PICRUSt2 analysis

PCR primer sequences were trimmed from raw reads using Cutadapt (Martin, 2011) and reads

were further processed with the R package DADA2 1.22.0 following the standard sequence

analysis pipeline for 16S rRNA genes (Callahan et al., 2016). Trimmed reads were filtered and

denoised using the filterAndTrim function under the following conditions: (truncLen = c(270,

240), maxEE = c(2,3), truncQ = 2, maxN = 0). The filtered reads of each sample were

dereplicated, Amplicon Sequencing Variants (ASVs) were deduced for forward and reverse

reads separately, and then the forward and reverse reads were merged at overlapping

positions. Chimeric sequences were removed and taxonomic assignment of amplicon

sequence variants (ASV) was performed using the SILVA taxonomy reference database

release version 138 (Quast et al., 2013). ASV sequences were extracted and aligned using

DECIPHER (Wright, 2020, 2016), and a phylogenetic tree was constructed using phangorn

(Schliep, 2011). In depth analysis of the bacterial community was performed in R using the

packages phyloseq (McMurdie and Holmes, 2013), vegan (Oksanen et al., 2008) and

tidyverse (Wickham et al., 2019) and their dependencies. Non-bacterial sequences assigned

to chloroplasts, mitochondria, eukaryotes, and archaea were excluded from further analysis,

resulting in 15659 unique ASVs and a median of 20,464 reads per sample. One seepage

sample was excluded from analysis due to very low amounts of sequencing reads per sample.

Normalization was done by rarefying all reads from each sample using the phyloseq function

rarefy_even_depth at a minimum of 4426 sequencing reads per sample. Subsequently, ASVs

with only 1 read were excluded from analysis. Shannon and Chao alpha diversity indexes

were calculated using the estimate_richness function in phyloseq.

2.5 Prediction of microbial functions

To approximate the functional potential of the microbial communities across the different

compartments, PICRUSt2 (Phylogenetic Investigation of Communities by Reconstruction of

Unobserved States) analysis was applied to predict microbial functions based on the obtained

amplicon sequencing data using v2.3.0 beta following the standard procedure for 16S rRNA

(Douglas et al., 2020). ASV sequences were first aligned and placed into a reference tree and

ASVs were corrected for multiple 16S rRNA genes by the 16S rRNA operon number of closely

related representatives. Gene families for each ASV were predicted based on KEGG orthologs

(KO) as identifiers for metabolic pathways (Kanehisa et al., 2023). The PICRUSt2 output of

7,523 KOs was subset for KO abundance of enzymes involved in the following pathways:

carbon fixation, nitrogen fixation, methane oxidation, carbon degradation and protein

hydrolysis. Metagenomic predictions of the selected pathways were based on the relative

abundance of the KO IDs across each sample. To assess the proportion of the bacterial

community associated with the selected metabolic functions across samples, we used the

relative abundance of each ASV from amplicon sequencing data for which a corresponding

KO was predicted.

2.6 Determination of absolute cell abundances and cell fluxes

Studies have shown that combining amplicon sequencing of marker genes and qPCR can be

used as an accurate method to infer cell numbers similar to direct cell determination by flow

cytometry (Bittar et al., 2018) and to enhance bacterial community profiling (Jian et al., 2020;

Lou et al., 2018). The quantity of exported cells was calculated using the 16S rRNA qPCR-

based gene abundances per liter and the amplicon sequencing reads on family-level. To

account for diverging operon numbers of the 16S rRNA gene across the 303 observed

bacterial families, gene abundances were corrected using the number of operons of the

respective family-level taxa obtained from the ribosomal RNA operon database

(https://rrndb.umms.med.umich.edu/) (Stoddard et al., 2015). The estimated absolute cell

count per liter for each sample of rainwater, throughfall, stemflow and seepage was

determined based on the relative read abundance from each taxon multiplied by the total 16S

rRNA gene abundance and divided by the family-specific operon numbers. Bacterial cell fluxes

per m^{-2} were calculated as described for TOC and TN using the estimated absolute cell

abundance per total sample volume and collector area for each sample from the two sampling

time points.

2.7 Estimation of bacterial contributions to TOC and TN

The fraction of bacterial cell carbon and nitrogen contributing to TOC and TN in rainwater,

throughfall, stemflow and seepage was estimated as previously described (Herrmann et al.,

2023; Krüger et al., 2021). Briefly, a mean carbon content of 39 fg per cell and mean nitrogen

content of 12 fg per cell was assumed as estimated under carbon-limited and nitrogen-limited

conditions (Vrede et al., 2002), respectively. The contribution of bacterial cell carbon and

nitrogen to TOC and TN, respectively, was estimated by first multiplying the operon number

corrected absolute bacterial abundances as well as the TOC/ TN content per liter by the

sample volume. The estimated number of bacterial cells was multiplied by the assumed

carbon and nitrogen content per cell. The resulting estimates of microbial cell carbon and

nitrogen were then set in relation to the total organic carbon (TOC) and total nitrogen (TN) in

the corresponding samples to estimate the bacterial contribution.

2.8 Statistical analysis

All statistical tests were performed using R version 4.12 (R Core Team, 2020) unless

otherwise described. Normal distribution of gene abundances and element fluxes were tested

with Shapiro-Wilk test. Non-parametric Kruskal-Wallis rank sum test and Dunn's multiple

comparison post-hoc test with R package FSA (Ogle et al., 2020) was used to test data for

significant differences between the sample types. To assess the contributing factors

accounting for the similarity/ dissimilarity of the bacterial community, a Permutational

Multivariate Analysis of Variance Using Distance Matrices (PERMANOVA) was conducted

using a Bray-Curtis distance matrix and the adonis2 function with 9999 permutations in R

package vegan (Oksanen et al., 2008). Nonmetric multidimensional scaling using Bray-Curtis

distance matrix was performed to test dissimilarity of the bacterial community between the

sample types, the growing seasons and the tree species. To visualize the distribution patterns

of abundant bacterial families across different rainwater flow paths and significant

environmental factors, a redundancy analysis (RDA) was performed using the decostand

function in vegan and Hellinger method for data standardization.

The significance of the observed differentiation was tested with ANalysis Of SIMilarity

(ANOSIM) using the anosim2 function from vegan. Correlations between microbial gene

abundance, TOC, TN and the sample volumes were assessed and visualized using ggpubr

(Kassambara, 2023). Relationships between the absolute abundance of the most abundant

17 bacterial families and the matter content of water solution samples were determined by

pairwise correlation using Spearman's rank coefficient and visualization using

ComplexHeatmap (Gu et al., 2016) and its dependencies. The number of shared ASV across

the water solution samples was assessed using packages phyloseq (McMurdie and Holmes,

2013), MicrobiotaProcess (Xu et al., 2023) and VennDiagram (Chen and Boutros, 2011). The

prevalence of dominant ASV representing the core microbiome across the samples was

determined using the metagMisc package including phyloseq, vegan, ALDEx2 (Fernandes et

al., 2013), metagenomeSeq (Paulson et al., 2013) and DESeq2 (Love et al., 2014). The

shared core microbiome of leaf, rain, throughfall and stemflow was assessed by filtering ASV

which were present in at least 70 % of the samples using the function

phyloseq_filter_prevalence from metagMisc. The sequences of core AVSs from leaf, rain,

throughfall and stemflow samples were extracted and placed in a phylogenetic tree using

ggtree (Xu et al., 2022). Lastly, the relative abundances of those ASV during both growing

seasons were plotted next to the tree as described in (Rolando et al., 2022).

The differential abundance of abundant ASV between phyllosphere and canopy-derived flow

paths was assessed using the package DESeq2. Unrarefied sequence reads with > 1

sequence across all samples were subset for leaf, throughfall and stemflow sample types. The

phyloseq object was converted to a DESeq2 data file and the ASV table was transformed

using the median ratio method for differential abundance analysis. In addition, we used the

microbiomeMarker package (Cao et al., 2022) to investigate the bacterial community for

biomarker ASV on genus level. Linear discriminant analysis (LDA) Effect Size (LEFSe) was

performed across all samples using CPM normalization method with a p value of 0.05 and

LDA cutoff of 4.

3. Results

3.1 Carbon and nitrogen fluxes across compartments

The export of total organic carbon (TOC) ranged from 1.4 mg C m^{-2} in stemflow to 1,600 mg C m^{-2} in seepage right below the litter layer and was significantly different between sample types ($p < 0.001$) (**Figure 2**A). Inorganic carbon (IC) fluxes were on average 2 to 4 times lower across all sample types (**Figure S 1**). Total nitrogen (TN) flux varied from as low as 0.1 mg N m^{-2} in stemflow to a maximum of 1,405 mg N m^{-2} in seepage collected after percolation through the upper 16 cm of the soil profile (**Figure 2**A). C:N ratios showed significantly higher values in throughfall and stemflow ($p < 0.001$) compared to rainwater and seepage (**Figure S 1**). Ammonium flux decreased from rainwater, throughfall and stemflow to the seepage flow path while fluxes of nitrate were highest in seepage (**Figure S 1**).

3.2 Microbial gene abundances, estimated cell fluxes, and contributions to C and N transport across compartments

Bacterial 16S rRNA gene abundances (**Figure S 2**A) differed significantly between the different sample types ($p < 0.001$) and were highest in throughfall and stemflow. Rainwater exhibited on average three orders of magnitude lower gene abundances. In seepage, bacterial 16S rRNA gene abundances decreased with increasing collection depth from below the litter layer down to 30 cm soil depth. Archaeal 16S rRNA gene abundances were consistently two to three orders of magnitude lower than bacterial 16S rRNA gene abundances and exhibited an opposing depth-dependent trend in seepage.

We further estimated bacterial cell abundances based on qPCR data and amplicon sequencing data, applying family-level corrections for multiple 16S rRNA gene operons per cell (**Table S 1**). Estimated bacterial abundances on leaf surfaces ranged from 1.4 x 10^5 to 5.4 x 10^8 cells g^{-1} dry leaf material while soil bacterial abundances ranged from 9.6 x 10^8 to 7.8 x 10^9 cells g^{-1} dry soil. In rainwater, estimated bacterial cell abundance ranged from 1.2 x 10^6 to 1.6 x 10^8 cells L^{-1} while throughfall and stemflow cell abundances were about 100 times

higher, ranging from 5.8 x 10^8 to 3.1 x 10^{11} cells L^{-1} and 7.9 x 10^9 to 2.3 x 10^{11} cells L^{-1},

respectively. Seepage bacterial abundance ranged from 3.8 x 10^8 to 3.9 x 10^{10} cells L^{-1} with

a decreasing trend with soil depth ($p < 0.01$). Based on estimated cell abundances, we also

determined the flux of bacterial cells transported by rain, throughfall, stemflow and seepage

of both sampling events. Across all samples and the two sampling time points, cell fluxes

ranged from lowest values in rainwater with 2.3 x 10^7 cells m^{-2}, 9.3 x 10^{11} cells m^{-2} by litter

seepage to highest export of 2.8 x 10^{12} cells m^{-2} per event by throughfall with significant

differences between sample types ($p < 0.001$) (**Figure 2**A). On average, cell fluxes via

throughfall and seepage litter were three orders of magnitude higher than in rainwater.

Seepage-mediated cell fluxes of seepage collected below 4 cm soil depth were one order of

magnitude lower than in throughfall. No differences in estimated cell fluxes were observed

between early and late growing season for a given sample type.

The transported amount of TOC and TN by rainwater, throughfall, stemflow and seepage were

positively correlated with the estimated total bacterial cell abundance per rain event across all

sample types and sampling months ($p < 0.05$) (**Figure 2**B), underlining the strong relationship

between microbial transport dynamics and matter transport. Estimation of the bacterial

contribution to matter transport indicated a significant bacterial contribution across the different

water flow pathways ($p < 0.001$) with bacterial cells contributing between 0.006 % of TOC and

0.007 % TN in rainwater, and up to 11.3 % TOC and 33.6 % TN in throughfall (**Figure 2**A).

3.3 Bacterial community composition from phyllosphere to soil

The ASV richness and alpha diversity (Shannon index) differed significantly between all

sample types (Kruskal-Wallis, $p < 0.001$) (**Figure 3**A). The highest number of different

bacterial taxa was observed in seepage, followed by stemflow, soil, leaf surface and

throughfall samples, while rainwater samples were the least diverse. Leaf, seepage, and soil

communities were on average more diverse than communities from rain, throughfall and

stemflow samples ($p < 0.05$). Alpha diversity and species richness of throughfall and stemflow

communities were lower during early growing season compared to late growing season ($p <$

0.05) while this effect was not observed for the other sample types.

Beta diversity analysis of bacterial communities revealed a clear differentiation based on the

sampled compartment (ANOSIM, R^2 = 0.52, $p <$ 0.001) with phyllosphere, seepage and soil

communities being distinct, while rainwater, throughfall and stemflow communities appeared

very variable and were not clearly separated (**Figure 3**B). Sample type was the strongest

driver for the community differentiation, accounting for 25.2 % of the total community variation

(PERMANOVA, $p <$ 0.001) (**Figure 3**B). The sampling month contributed 2.1 % of the

variation. Influence of tree species (European beech versus sycamore maple) explained only

less than 1 % of the variation and was not included as influencing factor in further analysis.

On family-level, 23 bacterial taxa each comprised more than 1 % of the total community

(**Figure S 2**B), which exhibited distinct distribution patterns across rain, throughfall, stemflow

and seepage (ANOSIM: R^2 = 0.54, $p <$ 0.001). Gammaproteobacteria, Alphaproteobacteria,

Bacteroidia, Actinobacteria and Acidobacteriae accounted for at least 90 % of all sequence

reads across all samples (**Figure 3**C). Across rainwater, throughfall and stemflow samples,

Pseudomonadaceae accounted for a large proportion of between 24 – 42 % together with

common plant pathogens such as *Erwiniaceae* and *Yersiniaceae*, particularly during the early

growing season. In phyllosphere samples, *Hymenobacteraceae* (32 mean ± 7 % standard

deviation) dominated at the late growing season sampling point together with

Sphingomonadaceae (16 ± 7 %) while *Beijerinckiaceae* (13 ± 6 %) were more abundant at the

early growing season. In contrast to rainwater, throughfall and stemflow,

Gammaproteobacteria were less abundant in phyllosphere samples and were here mostly

represented by *Oxalobacteraceae* (5 ± 3 %) and *Comamonadaceae* (6 ± 4 %) in the early and

late growing season samples, respectively. In soil communities, Alphaproteobacteria

accounted on average for 24 ± 3 % with *Xanthobacteraceae* being the dominant family (19 ±

3 %). Seepage communities were dominated by *Flavobacteriaceae, Oxalobacteraceae* and

Comamonadaceae.

The taxonomic composition derived from estimated cell abundances confirmed the patterns

revealed by relative read abundances (**Figure S 2**C). Analysis based on absolute abundance

revealed that most of rainwater-derived bacterial families, including *Pseudomonadaceae*,

Yersiniaceae and *Sphingomonadaceae*, were enriched by two to five orders of magnitude in

throughfall and stemflow. In contrast to those communities, the bacteria in seepage depicted

a two to three orders of magnitude increase in cell abundance of taxa such as

Xanthobacteraceae, *Spirosomaceae* and *Diplorickettsiaceae,* which showed also high relative

abundances in soil communities.

3.4 Predicted metagenomic functions of bacterial communities

The prediction of the bacterial genetic repertoire for selected biogeochemical transformations

unveiled distinct patterns across the sample types (**Figure S 3**, ANOSIM: $R^2 = 0.42$, $p < 0.001$).

The predicted relative abundance of KOs showed that most pathways had higher prevalence

towards soil and leaf communities except for protein hydrolysis, which was more prevalent in

bacteria transported by seepage.

The functional metagenomic prediction was based on the presence and absence of functional

modules of each pathway (**Figure 4**A). Hydrolytic enzymes showed high relative abundances

of methionyl aminopeptidase (0.083 %) and leucyl aminopeptidase (0.062 %) across the

samples. Potential for the degradation of carbon sources was also highly prevalent across the

samples depicted by beta-N-acetylhexosaminidase (0.046 %), hexosaminidase (0.033 %),

beta-glucosidase (0.029 %), chitinase (0.022 %) and alpha-amylase (0.017 %). Potential for

carbon fixation was mostly associated with the Calvin–Benson–Bassham pathway,

represented by *cbbS* and *cbbL* genes with an average relative abundance of 0.002 % and

0.006 %. Nitrogenases as primary enzymes for nitrogen fixation were prevalent in soil (0.0037

%), phyllosphere (0.0022 %) and seepage communities (0.0018 %). Methane oxidation was

primary related to methanol dehydrogenase (0.0014 %) across all samples, with the highest

relative fraction in phyllosphere (0.0037 %) and rainwater communities (0.0024 %).

A similar pattern of the functional profiles across the sample types was observed based on

relative abundance of bacterial classes for which predicted KO were present (**Figure 4**B). The

prevalence of bacteria which showed functional potential differed significantly between the

sample types for each pathway ($p < 0.05$). The fraction of bacterial communities showing

capability of utilizing enzymes involved in various carbon and protein decomposition

processes ranged, on average, from 52 % in throughfall to 75 % in seepage and 60 % in

throughfall to 80 % in leaf samples, respectively. Carbon fixation made up by a fraction of 2 %

to 13 % of the bacterial community with lowest potential in throughfall and stemflow, and

highest potential in seepage samples. Pathways related to biological nitrogen fixation and

methane oxidation showed the highest prevalence of 2.5 % and 1.1 %, respectively, in leaf

communities, while rain and throughfall communities showed functional potential for both

pathways of less than 1 %.

3.5 Connectivity of bacterial communities along the rainwater flow path

Only a small proportion of 1.7 % (117 ASVs of all ASVs) of the bacterial community was shared

across all water sample types. The major fraction of 62 % of all ASVs were exclusively present

in seepage, while less than 10 % of the total ASVs were only found in either rain, throughfall

or stemflow communities (**Figure S 4**). Stemflow communities shared 5.9 % of the ASVs with

rainwater, 10.7 % with throughfall, and 6.9 % with seepage.

We defined bacterial core taxa based on occurrence in at least 70 % of all samples and tracked

the enrichment or decline of single core ASVs from rainwater to phyllosphere and transport by

throughfall, stemflow to seepage. Core taxa which constituted a large community fraction on

leaves included members of the *Hymenobacteraceae*, *Acetobacteraceae*, *Bejierinckiaceae*

and *Sphingomonadaceae*. ASVs affiliated with the genera *Hymenobacter*, *Acidiphilium*,

Methylorubrum and *Sphingomonas* accounted for up to 1 % relative abundance of the total

community on leaves but were less abundant in rainwater. Some ASVs classified as

Sphingomonas were also observed in throughfall and stemflow (**Figure 5**). Among the

phyllosphere community, *Acidiphilium*, *Methylorubrum* and *Sphingomonas* were also

observed in rainwater. In contrast, *Pseudomonadaceae* and *Oxalobacteraceae* including the

genera *Pseudomonas* and *Massilia*, respectively, were abundant in rainwater, transferred via

throughfall and stemflow and ultimately exported by seepage but were less abundant on

leaves.

In addition to the preferential association of core ASVs with a specific sample type, we

determined the differential abundance of genus-level taxa which showed either the tendency

to stick on leaf surfaces or to be washed off by throughfall and stemflow (**Figure S 5**),

indicating their potential as dispersal agents between microbial habitats. Members of

Pseudomonadaceae, *Yersiniaceae* and *Oxalobacteraceae* were highly enriched in throughfall

and stemflow while *Sphingomonadaceae* and *Hymenobacteraceae* showed stronger

association with the phyllosphere. Instead, members of *Xanthobacteraceae* and

Flavobacteriaceae were more enriched in seepage compared to throughfall and stemflow

(**Figure S 6**), and *Flavobacteriaceae* and *Sphingomonadaceae* were strongly enriched in

seepage compared to soil. LEFSe analysis confirmed the specific association of those

bacterial taxa with either the tree phyllosphere or the different mobile fractions along the

vertical water flow path (**Figure S 7**).

4. Discussion

Precipitation events transmit a vast number of organic particles, including microbes, representing a major environmental agent connecting the atmosphere with the phyllosphere, pedosphere and lithosphere (Ponette-González et al., 2020; Van Stan et al., 2020). The transported matter and microbial cells can potentially impact microbial communities and nutrient cycling along the water flow path from the plant phyllosphere to soil (Van Stan and Stubbins, 2018; Van Stan et al., 2020). However, the extent of rainwater-driven microbial mobilization within the forest phyllosphere and from bark surfaces, and its contribution to carbon and nitrogen fluxes have so far remained unknown. Here, we assessed the diversity and abundance of bacteria along this flow path, starting from intercepting rain via tree canopy-derived throughfall and stemflow to soil seepage in a temperate forest area at an early and late stage of the growing season.

Our results clearly demonstrate that canopy-derived throughfall and stemflow microbial communities are quantitatively important carriers of organic matter to the forest floor, contributing up to 11 and 34 % to total carbon and nitrogen fluxes, respectively. Thus, they play a central role in the throughfall- and stemflow-mediated provision of fresh organic carbon and nitrogen to the forest floor (Van Stan et al., 2021). Throughfall and stemflow obtained in this study from a beech forest during a three-days rain event were highly enriched with bacteria with 10 to 1000 higher estimated cell numbers than previously reported from a subtropical oak-cedar forest (Bittar et al., 2018). Given the fact that throughfall and stemflow samplers were directly sampled at the end of a three-days rain event at overall low ambient temperatures between 0-20°C, we can exclude that massive growth of bacterial cells in the samplers may have led to a strong overestimation of these bacterial loads. Despite on average similar cell concentrations L^{-1} in throughfall and stemflow solutions, stemflow-mediated cell fluxes normalized to tree area were two orders of magnitude lower than in throughfall, which is typically also observed for TOC and TN fluxes when comparing these two water flow paths (Van Stan and Stubbins, 2018). Throughfall cell fluxes were comparable to those from

seepage collected below the litter layer, indicating a substantial above-ground translocation of

bacteria to soil.

In contrast to the high estimated proportions of bacterial derived carbon and nitrogen in

throughfall and stemflow, bacterial contributions to TOC and TN were three times lower in

seepage solutions, matching with previously reported values from the same lysimeters (Krüger

et al., 2021) and tension-controlled lysimeters from the same field site (Herrmann et al., 2023).

Total estimated cell fluxes were even ten times lower in seepage below 16 cm compared to

throughfall. Apparently, a large fraction of the cells mobilized by throughfall and stemflow get

retained in the soil, which acts as a strong filter for this above-ground transport. Turnover

processes in the soil result in carbon and nitrogen export primarily as dissolved compounds

with a lower contribution of mobilized microbial cells. During rain events, throughfall and

stemflow mediated inputs might supplement the nutrient pool of the forest litter and soil and

could additionally affect microbial nutrient cycling capabilities by introducing active

microorganisms with their respective biodegradation capabilities or exoenzymes from

upstream communities. Pathway predictions indeed indicated that bacterial communities from

throughfall, stemflow and seepage exhibited strong potential for hydrolytic degradation of

carbohydrates and proteins. However, since PICRUSt2 only predicts metagenomic functions

from marker gene abundance and phylogenetic placement to reference sequences (Douglas

et al., 2020), speculations about the full degradation potential need to be treated with caution.

The type of canopy-derived water flow path (throughfall versus stemflow) was the strongest

effector for bacterial community similarity/ dissimilarity, suggesting that the composition of the

phyllosphere versus bark-associated source communities and potential differences in

microbial detachment mechanisms between these two habitats could underlie the observed

community differences. Phyllosphere communities were more similar to rain, throughfall and

stemflow communities than to seepage and soil, suggesting an exchange of microbes

between phyllosphere, throughfall and stemflow in a taxon-specific manner. Rainwater can

potentially support the colonization but also the wash-off of epiphytic microbes from leaf

surfaces (Guerrieri et al., 2020). The observed fraction of 10 % of shared taxa between

throughfall and stemflow communities suggested that runoff from the canopy exported a basic,

selective fraction of bacteria independent of the phyllosphere location, which was further

modulated by exchange with the specific colonizers of either leaves or bark (Guerrieri et al.,

2020; Teachey et al., 2018). Similarly, Hudson et al. (2023) demonstrated that the bacterial

community composition in stemflow was distinct from that found on tree bark. Bark-water

interactions during rainfall may lead to the detachment of loosely attached microbes and

enable their accumulation further down as major infiltration source in soil (Carlyle-Moses et

al., 2020).

Weather conditions and tree species were identified as important factors shaping throughfall

bacterial communities in a mediterranean holm oak forest (Guerrieri et al., 2020) and a

subtropical Southern live oak forest (Teachey et al., 2018) with Alphaproteobacteria,

Gammaproteobacteria and Bacteroidia representing main bacterial classes. Following the

above-ground   flow   path   at   our   study   site,   *Pseudomonadaceae*,   *Erwiniaceae*,

Oxalobacteraceae and *Sphingobacteriaceae* were introduced into the phyllosphere by rainfall

and depicted 10 to 100 times enrichment in relative abundance on leaf surfaces compared to

rainwater before being selectively transported by throughfall and stemflow to the soil. Our

results further demonstrated a substantial difference in the microbial communities of the

above-ground flow paths when comparing an early and late time point of the growing season.

During May sampling, *Pseudomonadaceae* accounted on average for over 50 % of the total

community in bulk rain, throughfall and stemflow, indicating that at this time point, only a single

taxonomic   group   contributed   substantially   to   matter   transport   and   potential   carbon

degradation capabilities. *Pseudomonas* species act as common pathogens and epiphytic

plant commensals in the phyllosphere (Vorholt, 2012). However, they are also typically

observed among microbes attached to plant pollen (Manirajan et al., 2018; Williams, 2013),

which can be transmitted by dry deposition (Sánchez-Parra et al., 2021) and rainwater (Cho

and Jang, 2014). Higher pollen bioaerosols or pollen deposits derived from the flowering of

beech or nearby Norway spruce stands in May might have constituted a large microbial source

of pollen-associated *Pseudomonadaceae* in rainwater, throughfall and stemflow (Guidone et

al., 2021).

The observed distinction of bacterial communities in seepage versus throughfall and stemflow

may not only be linked to differences in the bacterial source communities but also to

differences in the general bacterial detachment mechanisms between above-ground and

below-ground habitats. Preferential, taxon-specific microbial transport through soil by

seepage-mediated translocation is mainly attributed to cellular characteristics such as cell

size, shape and surface charge (i.e., Zeta potential) (Dibbern et al., 2014; Herrmann et al.,

2023; Zhang et al., 2018). In the phyllosphere, the main factor controlling the preferential

transfer from leaf surfaces to throughfall might be the strength of attachment to the surface

rather than cell size. Members of the *Pseudomonadaceae* can increase local water availability

of leaf surfaces by releasing surfactants, allowing proliferation (Koskella, 2020), which might

also be a reason for preferential mobilization and detachment of members of this family by

throughfall and stemflow. In addition, surfactants may also support the detachment of other

leaf-associated microorganisms. Prevalent bacterial epiphytes from the phyllosphere beyond

Pseudomonadaceae are *Methylobacteriaceae* and *Sphingomonadaceae* (Bashir et al., 2022;

Delmotte et al., 2009; Lindow and Brandl, 2003; Vacher et al., 2016; Vorholt, 2012), which

were also observed in rainwater at our study site, depicting the connectivity between

atmosphere, phyllosphere and pedosphere. Especially *Pseudomonadaceae* could also

constitute an important source of extracellular exoenzymes performing hydrolytic degradation,

affecting decomposition of soil organic matter (Mori et al., 2019).

Despite the high numbers of cells transported by throughfall and stemflow, it remains an open

question whether cells remain viable throughout the whole vertical passage, and how many

living microbes ultimately reach the soil. The fact that we were able to trace individual ASVs

from the phyllosphere to throughfall, stemflow and seepage samples demonstrates that

distinct members of the bacterial community at least remained intact during this continuous

passage or were successfully transferred between two neighboring compartments. A seepage

study using electron microscopy further indicated that living microbes remained intact during

soil passage across 60 cm depth by seepage transmission (Dibbern et al., 2014).

Our results further suggest that multiple taxa were able to colonize and thrive in all of the

investigated ecosystem compartments from atmosphere to soil. Establishment and

proliferation in the respective compartments probably ensure that the taxa can also be

released again from these compartments and be transported again along the flow path.

However, with our approach, we cannot distinguish between continuous cross-compartment

transport of individual cells or discontinuous transport, resulting from establishment in each

compartment, proliferation, and subsequent secondary mobilization of microbial taxa.

Consequently, occurrence of the same ASVs across the vertical flow path does not necessarily

indicate an uninterrupted flow of individual cells through the system but the ability of certain

taxa to be continuously mobilized from one compartment and transported to the other, and to

establish stable populations in the respective sink communities, from whereon cells can be

mobilized again, resulting in the observed continuity and connectivity of microbial communities

and their functional potential.

5. Conclusions

Our study clearly demonstrates that microorganisms transported by rainwater, throughfall,

stemflow and seepage contribute substantially to carbon and nitrogen fluxes in a deciduous

forest ecosystem. Contributions and cell fluxes were larger in above-ground compared to

below-ground compartments, suggesting that a considerable fraction of mobilized cells was

retained in the soil and not further transported. Throughfall- and stemflow-mediated flow paths

act as substantial carriers of microbial biomass, carbon and nitrogen which might affect soil

matter transformation by introducing a fresh source of nutrients but also new genetic variability

to the respective microbial sink communities along the flow path. In contrast to periodic cell

and nutrient inputs via litter fall, the throughfall- and stemflow-mediated vertical microbial

transport to the forest floor throughout the year, acting as central connection between

phyllosphere and forest floor in terms of carbon and nitrogen inputs but also as contribution to

the functional diversity of soil microbiomes. Our results further expand a previously observed

taxon-specificity in seepage-mediated microbial mobilization in soil to throughfall and

stemflow, presumably along with different mechanisms of microbial detachment across these

flow paths.

Data availability

Raw sequence files were deposited at the European Nucleotide Sequence Archive with project number PRJEB92466 under accessions ERS25231161 to ERS25231303.

Acknowledgements

We thank Falko Gutmann, Stefan Riedel and Linda Gorniak for support of field and laboratory work, and Martin Stephan from the Otto Schott Institute for Materials Research at Friedrich Schiller University Jena for producing customized soil lysimeters. We are further grateful for the background graphics provided by Clemens Schulze.

Funding

This study was part of the Collaborative Research Centre 1076 AquaDiva (CRC AquaDiva) of the Friedrich Schiller University Jena, funded by the Deutsche Forschungsgemeinschaft under project number 218627073. The infrastructure for MiSeq Illumina sequencing was financially supported by the Thüringer Ministerium für Wirtschaft, Wissenschaft und Digitale Gesellschaft (TMWWDG project 2016 FGI 0024 "BIODIV").

Author contributions: CRediT

Markus Krüger: Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Writing – original draft, Writing – review and editing. Karin Potthast: Conceptualization, Methodology, Writing – review and editing. Beate Michalzik: Conceptualization, Writing – review and editing. Alexander Tischer: Conceptualization, Methodology, Writing – review and editing. Martina Herrmann: Funding acquisition, Conceptualization, Sample processing, Writing – review and editing.

Declaration of competing interest

The authors declare that the work of this study was achieved without financial or personal conflict of interest.

References

Andrews, J.H., Harris, R.F., 2000. The ecology and biogeography of microorganisms on plant surfaces. Annu. Rev. Phytopathol. 38, 145-180. https://doi.org/10.1146/annurev.phyto.38.1.145

Bashir, I., War, A.F., Rafiq, I., Reshi, Z.A., Rashid, I., Shouche, Y.S., 2022. Phyllosphere microbiome: Diversity and functions. Microbiol. Res. 254, (126888). https://doi.org/10.1016/j.micres.2021.126888

Bittar, T.B., Pound, P., Whitetree, A., Moore, L.D., Van Stan, J.T., 2018. Estimation of Throughfall and Stemflow Bacterial Flux in a Subtropical Oak-Cedar Forest. Geophys. Res. Lett. 45, 1410–1418. https://doi.org/10.1002/2017GL075827

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nat. Methods 13, 581–583. https://doi.org/10.1038/nmeth.3869

Cao, Y., Dong, Q., Wang, D., Zhang, P., Liu, Y., Niu, C., 2022. microbiomeMarker: an R/Bioconductor package for microbiome marker identification and visualization. Bioinformatics 38, 4027–4029. https://doi.org/10.1093/bioinformatics/btac438

Carlyle-Moses, D.E., Iida, S., Germer, S., Llorens, P., Michalzik, B., Nanko, K., Tanaka, T., Tischer, A., Levia, D.F., 2020. What We Know About Stemflow's Infiltration Area. Front. For. Glob. Chang. 3, (532805). https://doi.org/10.3389/ffgc.2020.577247

Chelius, M.K., Triplett, E.W., 2001. The diversity of archaea and bacteria in association with the roots of Zea mays L. Microb. Ecol. 41, 252–263. https://doi.org/10.1007/s002480000087

Chen, H., Boutros, P.C., 2011. VennDiagram: A package for the generation of highly-customizable Venn and Euler diagrams in R. BMC Bioinformatics 12, 35. https://doi.org/10.1186/1471-2105-12-35

Cho, B.C., Jang, G. II, 2014. Active and diverse rainwater bacteria collected at an inland site in spring and summer 2011. Atmos. Environ. 94, 409–416. https://doi.org/10.1016/j.atmosenv.2014.05.048

Delmotte, N., Knief, C., Chaffron, S., Innerebner, G., Roschitzki, B., Schlapbach, R., Von Mering, C., Vorholt, J.A., 2009. Community proteogenomics reveals insights into the physiology of phyllosphere bacteria. Proc. Natl. Acad. Sci. U. S. A. 106, 16428–16433. https://doi.org/10.1073/pnas.0905240106

DeLong, E.F., 1992. Archaea in coastal marine environments. Proc. Natl. Acad. Sci. 89, 5685–5689. https://doi.org/10.1073/pnas.89.12.5685

Deng, Q., Hui, D., Chu, G., Han, X., Zhang, Q., 2017. Rain-induced changes in soil CO_2 flux and microbial community composition in a tropical forest of China. Sci. Rep. 7, 5539. https://doi.org/10.1038/s41598-017-06345-2

Després, V.R., Alex Huffman, J., Burrows, S.M., Hoose, C., Safatov, A.S., Buryak, G., Fröhlich-Nowoisky, J., Elbert, W., Andreae, M.O., Pöschl, U., Jaenicke, R., 2012. Primary biological aerosol particles in the atmosphere: A review. Tellus, Ser. B Chem. 64, (15598). Phys. Meteorol. https://doi.org/10.3402/tellusb.v64i0.15598

Dibbern, D., Schmalwasser, A., Lueders, T., Totsche, K.U., 2014. Selective transport of plant root-associated bacterial populations in agricultural soils upon snowmelt. Soil Biol. Biochem. 69, 187–196. https://doi.org/10.1016/j.soilbio.2013.10.040

Douglas, G.M., Maffei, V.J., Zaneveld, J.R., Yurgel, S.N., Brown, J.R., Taylor, C.M.,
Huttenhower, C., Langille, M.G.I., 2020. PICRUSt2 for prediction of metagenome
functions. Nat. Biotechnol. 38, 685-688. https://doi.org/10.1038/s41587-020-0548-6

Fernandes, A.D., Macklaim, J.M., Linn, T.G., Reid, G., Gloor, G.B., 2013. ANOVA-Like
Differential Expression (ALDEx) Analysis for Mixed Population RNA-Seq. PLoS One 8,
e67019. https://doi.org/10.1371/journal.pone.0067019

Friesen, J., 2020. Flow pathways of throughfall and stemflow through the subsurface, in:
Precipitation Partitioning by Vegetation: A Global Synthesis. Springer International
Publishing, pp. 214–227. https://doi.org/10.1007/978-3-030-29702-2_13

Gu, Z., Eils, R., Schlesner, M., 2016. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics 32, 2847–2849.
https://doi.org/10.1093/bioinformatics/btw313

Guerrieri, R., Cáliz, J., Mattana, S., Barceló, A., Candela, M., Elustondo, D., Fortmann, H.,
Hellsten, S., Koenig, N., Lindroos, A.J., Matteucci, G., Merilä, P., Michalski, G., Nicolas,
M., Thimonier, A., Turroni, S., Vanguelova, E., Verstraeten, A., Waldner, P., Watanabe,
M., Casamayor, E.O., Peñuelas, J., Mencuccini, M., 2024. Substantial contribution of
tree canopy nitrifiers to nitrogen fluxes in European forests. Nat. Geosci. 17, 130–136.
https://doi.org/10.1038/s41561-023-01364-3

Guerrieri, R., Lecha, L., Mattana, S., Cáliz, J., Casamayor, E.O., Barceló, A., Michalski, G.,
Peñuelas, J., Avila, A., Mencuccini, M., 2020. Partitioning between atmospheric
deposition and canopy microbial nitrification into throughfall nitrate fluxes in a
Mediterranean forest. J. Ecol. 108, 626–640. https://doi.org/10.1111/1365-2745.13288

Guidone, M., Gordon, D.A., Van Stan, J.T., 2021. Living particulate fluxes in throughfall and
stemflow during a pollen event. Biogeochemistry 153, 323–330.
https://doi.org/10.1007/s10533-021-00787-7

Herrmann, M., Geesink, P., Richter, R., Küsel, K., 2021. Canopy Position Has a Stronger
Effect than Tree Species Identity on Phyllosphere Bacterial Diversity in a Floodplain
Hardwood Forest. Microb. Ecol. 81, 157–168. https://doi.org/10.1007/s00248-020-
01565-y

Herrmann, M., Hädrich, A., Küsel, K., 2012. Predominance of thaumarchaeal ammonia
oxidizer abundance and transcriptional activity in an acidic fen. Environ. Microbiol. 14,
3013–3025. https://doi.org/10.1111/j.1462-2920.2012.02882.x

Herrmann, M., Lehmann, K., Totsche, K.U., Küsel, K., 2023. Seepage-mediated export of
bacteria from soil is taxon-specific and driven by seasonal infiltration regimes. Soil Biol.
Biochem. 187, 109192. https://doi.org/10.1016/J.SOILBIO.2023.109192

Hu, W., Niu, H., Murata, K., Wu, Z., Hu, M., Kojima, T., Zhang, D., 2018. Bacteria in
atmospheric waters: Detection, characteristics and implications. Atmos. Environ. 179,
201-221. https://doi.org/10.1016/j.atmosenv.2018.02.026

Hudson, J.E., Levia, D.F., Yoshimura, K.M., Gottel, N.R., Hudson, S.A., Biddle, J.F., 2023.
Mapping bark bacteria: initial insights of stemflow-induced changes in bark surface
phyla. Microbiol. Spectr. 11, e03562-23. https://doi.org/10.1128/spectrum.03562-23

Iguchi, H., Yurimoto, H., Sakai, Y., 2015. Interactions of methylotrophs with plants and other
heterotrophic bacteria. Microorganisms. 3, 137-151.
https://doi.org/10.3390/microorganisms3020137

Jeffrey, L.C., Maher, D.T., Chiri, E., Leung, P.M., Nauer, P.A., Arndt, S.K., Tait, D.R.,
Greening, C., Johnston, S.G., 2021. Bark-dwelling methanotrophic bacteria decrease

methane emissions from trees. Nat. Commun. 12, 2127.
https://doi.org/10.1038/s41467-021-22333-7

Kanehisa, M., Furumichi, M., Sato, Y., Kawashima, M., Ishiguro-Watanabe, M., 2023. KEGG
for taxonomy-based analysis of pathways and genomes. Nucleic Acids Res. 51, 587–
592. https://doi.org/10.1093/nar/gkac963

Kassambara, A., 2023. "ggplot2" Based Publication Ready Plots.

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., Glöckner, F.O.,
2013. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and
next-generation sequencing-based diversity studies. Nucleic Acids Res. 41, e1.
https://doi.org/10.1093/nar/gks808

Koskella, B., 2020. The phyllosphere. Curr. Biol. 30, 1143-1146.
https://doi.org/10.1016/j.cub.2020.07.037

Krüger, M., Potthast, K., Michalzik, B., Tischer, A., Küsel, K., Deckner, F.F.K., Herrmann, M.,
2021. Drought and rewetting events enhance nitrate leaching and seepage-mediated
translocation of microbes from beech forest soils. Soil Biol. Biochem. 154, 108153.
https://doi.org/10.1016/j.soilbio.2021.108153

Levia, D., Bischoff, S., Gruselle, M.-C., Näthe, K., Legates, D.R., Lutgen, A.N., Boyer, E.W.,
Michalzik, B., 2023. Geometric configurations of particulate matter in terrestrial
solutions of a temperate beech forest. J. Aerosol Sci. 106196.
https://doi.org/10.1016/J.JAEROSCI.2023.106196

Levia, D.F., Keim, R.F., Carlyle-Moses, D.E., Frost, E.E., 2011. Throughfall and Stemflow in
Wooded Ecosystems, in: Forest Hydrology and Biogeochemistry. pp. 425–443.
https://doi.org/10.1007/978-94-007-1363-5_21

Lindow, S.E., Brandl, M.T., 2003. Microbiology of the Phyllosphere. Appl. Environ. Microbiol.
69, 1875–1883. https://doi.org/10.1128/AEM.69.4.1875

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 15. https://doi.org/10.1186/s13059-014-
0550-8

Manirajan, B.A., Maisinger, C., Ratering, S., Rusch, V., Schwiertz, A., Cardinale, M., Schnell,
S., 2018. Diversity, specificity, co-occurrence and hub taxa of the bacterial-fungal pollen
microbiome. FEMS Microbiol. Ecol. 94, 112. https://doi.org/10.1093/femsec/fiy112

Martin, M., 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet.journal 17, 10. https://doi.org/10.14806/ej.17.1.200

McMurdie, P.J., Holmes, S., 2013. Phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. PLoS One 8, e61217.
https://doi.org/10.1371/journal.pone.0061217

Michalzik, B., Kalbitz, K., Park, J.H., Solinger, S., Matzner, E., 2001. Fluxes and
concentrations of dissolved organic carbon and nitrogen - A synthesis for temperate
forests. Biogeochemistry 52, 173–205. https://doi.org/10.1023/A:1006441620810

Mori, T., Wang, S., Zhang, W., Mo, J., 2019. A potential source of soil ecoenzymes: From
the phyllosphere to soil via throughfall. Appl. Soil Ecol. 139, 25–28.
https://doi.org/10.1016/j.apsoil.2019.02.004

Ogle, D., Wheeler, P., Dinno, A., 2020. FSA: Fisheries Stock Analysis. R package version
0.8.31.

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Simpson, G.L., Solymos, P.M., Stevens,

M.H.H., & Wagner, H., 2008. The vegan package. Community Ecol. Packag. 190.

Paulson, J.N., Colin Stine, O., Bravo, H.C., Pop, M., 2013. Differential abundance analysis
for microbial marker-gene surveys. Nat. Methods 10, 1200–1202.
https://doi.org/10.1038/nmeth.2658

Ponette-González, A.G., Van Stan, J.T., Magyar, D., 2020. Things seen and unseen in
throughfall and stemflow, in: Precipitation Partitioning by Vegetation: A Global
Synthesis. Springer, Cham, pp. 70–87. https://doi.org/10.1007/978-3-030-29702-2_5

Potthast, K., Meyer, S., Crecelius, A.C., Schubert, U.S., Tischer, A., Michalzik, B., 2017.
Land-use and fire drive temporal patterns of soil solution chemistry and nutrient fluxes.
Sci. Total Environ. 605–606, 514–526.
https://doi.org/10.1016/J.SCITOTENV.2017.06.182

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner,
F.O., 2013. The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. Nucleic Acids Res. 41, 590–596.
https://doi.org/10.1093/nar/gks1219

Reche, I., D'Orta, G., Mladenov, N., Winget, D.M., Suttle, C.A., 2018. Deposition rates of
viruses and bacteria above the atmospheric boundary layer. ISME J. 12, 1154–1162.
https://doi.org/10.1038/s41396-017-0042-4

Rolando, J.L., Kolton, M., Song, T., Kostka, J.E., 2022. The core root microbiome of Spartina
alterniflora is predominated by sulfur-oxidizing and sulfate-reducing bacteria in Georgia
salt marshes, USA. Microbiome 10. https://doi.org/10.1186/s40168-021-01187-7

Sadeghi, S.M.M., Gordon, D.A., Van Stan, J.T., 2020. A global synthesis of throughfall and
stemflow hydrometeorology, in: Precipitation Partitioning by Vegetation: A Global
Synthesis. Springer International Publishing, pp. 48–69. https://doi.org/10.1007/978-3-
030-29702-2_4

Sánchez-Parra, B., Núñez, A., García, A.M., Campoy, P., Moreno, D.A., 2021. Distribution of
airborne pollen, fungi and bacteria at four altitudes using high-throughput DNA
sequencing. Atmos. Res. 249, 105306. https://doi.org/10.1016/j.atmosres.2020.105306

Scheiner, D., 1974. A modified version of the sodium salicylate method for analysis of
wastewater nitrates. Water Res. 8, 835–840. https://doi.org/10.1016/0043-
1354(74)90030-X

Schliep, K.P., 2011. phangorn: Phylogenetic analysis in R. Bioinformatics 27, 592–593.
https://doi.org/10.1093/bioinformatics/btq706

Seelmeyer, G., 1954. Deutsche Einheitsverfahren zur Wasseruntersuchung.
Herausgegeben im Auftrage der Fachgruppe Wasserchemie in der Gesellschaft
Deutscher Chemiker E. V. Bearbeitet von L. W. Haase, H. Stooff, G. Gad, W. Wesly
u.a. Verlag Chemie, Weinheim/Bergstraße 1954, 180. Mater. Corros. 5, 357–357.
https://doi.org/10.1002/maco.19540050819

Stoddard, S.F., Smith, B.J., Hein, R., Roller, B.R.K., Schmidt, T.M., 2015. rrnDB: Improved
tools for interpreting rRNA gene abundance in bacteria and archaea and a new
foundation for future development. Nucleic Acids Res. 43, 593–598.
https://doi.org/10.1093/nar/gku1201

Takai, K., Horikoshi, K., 2000. Rapid detection and quantification of members of the archaeal
community by quantitative PCR using fluorogenic probes. Appl. Environ. Microbiol. 66,
5066–5072. https://doi.org/10.1128/AEM.66.11.5066-5072.2000

Teachey, M.E., Pound, P., Ottesen, E.A., Van Stan, J.T., 2018. Bacterial community

composition of throughfall and stemflow. Front. For. Glob. Chang. 1.
https://doi.org/10.3389/ffgc.2018.00007

Team, R.C., 2020. A Language and Environment for Statistical Computing. R Found. Stat.
Comput.

Tischer, A., Michalzik, B., Lotze, R., 2020. Nonuniform but highly preferential stemflow
routing along bark surfaces and actual smaller infiltration areas than previously
assumed: A case study on European beech (Fagus sylvatica L.) and sycamore maple
(Acer pseudoplatanus L.). Ecohydrology 13, e2230. https://doi.org/10.1002/eco.2230

Vacher, C., Hampe, A., Porté, A.J., Sauer, U., Compant, S., Morris, C.E., 2016. The
Phyllosphere: Microbial Jungle at the Plant-Climate Interface. Annu. Rev. Ecol. Evol.
Syst. 47, 1–24. https://doi.org/10.1146/annurev-ecolsys-121415-032238

Van Stan, J.T., Morris, C.E., Aung, K., Kuzyakov, Y., Magyar, D., Rebollar, E.A., Remus-
Emsermann, M., Uroz, S., Vandenkoornhuyse, P., 2020. Precipitation partitioning—
hydrologic highways between microbial communities of the plant microbiome?, in:
Precipitation Partitioning by Vegetation: A Global Synthesis. Springer International
Publishing, pp. 228–251. https://doi.org/10.1007/978-3-030-29702-2_14

Van Stan, J.T., Ponette-González, A.G., Swanson, T., Weathers, K.C., 2021. Throughfall
and stemflow are major hydrologic highways for particulate traffic through tree
canopies. Front. Ecol. Environ. 19, 404–410. https://doi.org/10.1002/fee.2360

Van Stan, J.T., Stubbins, A., 2018. Tree-DOM: Dissolved organic matter in throughfall and
stemflow. Limnol. Oceanogr. Lett. 3, 199–214. https://doi.org/10.1002/lol2.10059

Van Stan, J.T., Wagner, S., Guillemette, F., Whitetree, A., Lewis, J., Silva, L., Stubbins, A.,
2017. Temporal Dynamics in the Concentration, Flux, and Optical Properties of Tree-
Derived Dissolved Organic Matter in an Epiphyte-Laden Oak-Cedar Forest. J.
Geophys. Res. Biogeosciences 122, 2982–2997.
https://doi.org/10.1002/2017JG004111

Visco, G., Campanella, L., Nobili, V., 2005. Organic carbons and TOC in waters: An
overview of the international norm for its measurements, in: Microchemical Journal.
Elsevier, 79, pp. 185–191. https://doi.org/10.1016/j.microc.2004.10.018

Vorholt, J.A., 2012. Microbial life in the phyllosphere. Nat. Rev. Microbiol. 10, 828-840.
https://doi.org/10.1038/nrmicro2910

Vrede, K., Heldal, M., Norland, S., Bratbak, G., 2002. Elemental composition (C, N, P) and
cell volume of exponentially growing and nutrient-limited bacterioplankton. Appl.
Environ. Microbiol. 68, 2965–2971. https://doi.org/10.1128/AEM.68.6.2965-2971.2002

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,
Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller,
K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., Takahashi, K., Vaughan, D., Wilke,
C., Woo, K., Yutani, H., 2019. Welcome to the Tidyverse. J. Open Source Softw. 4,
1686. https://doi.org/10.21105/joss.01686

Williams, C.G., 2013. Forest tree pollen dispersal via the water. Am. J. Bot. 100, 1184–1190.
https://doi.org/10.3732/ajb.1300085

Wright, E.S., 2020. RNAcontest: Comparing tools for noncoding RNA multiple sequence
alignment based on structural consistency. RNA 26, 531–540.
https://doi.org/10.1261/rna.073015.119

Wright, E.S., 2016. Using DECIPHER v2.0 to analyze big biological sequence data in R. R J.
8, 352–359. https://doi.org/10.32614/rj-2016-025

Xu, S., Li, L., Luo, X., Chen, M., Tang, W., Zhan, L., Dai, Z., Lam, T.T., Guan, Y., Yu, G.,
2022. Ggtree: A serialized data object for visualization of a phylogenetic tree and
annotation data. iMeta 1, 56. https://doi.org/10.1002/imt2.56

Xu, S., Zhan, L., Tang, W., Wang, Q., Dai, Z., Zhou, L., Feng, T., Chen, M., Wu, T., Hu, E.,
Yu, G., 2023. MicrobiotaProcess: A comprehensive R package for deep mining
microbiome. Innovation 4, 100388. https://doi.org/10.1016/j.xinn.2023.100388

Zhang, L., Lehmann, K., Totsche, K.U., Lueders, T., 2018. Selective successional transport
of bacterial populations from rooted agricultural topsoil to deeper layers upon extreme
precipitation events. Soil Biol. Biochem. 124, 168–178.
https://doi.org/10.1016/j.soilbio.2018.06.012

Figures

Figure 1: Study site and study design for sample collection at the Hainich Critical Zone Exploratory in western

Thuringia, Germany. A) Sampling was conducted at an open area-site for rainwater representative of

above-canopy precipitation and at a mixed beech forest site for phyllosphere, throughfall, stemflow, and

seepage and soil sampling at different depths. B) Number of samples for each sample type across two

sampling time points. C) Conceptional representation of samples spanning from the tree canopy down to

the soil, and photographs of the two tree species, sample collectors and a typical soil core.

Figure 2: Bacterial transport across the different compartments, and contributions and correlations to

transported carbon and nitrogen. A) Fluxes of TOC and TN expressed per m^2, bacterial contributions to

TOC and TN content, and fluxes of bacterial cells per m^2 across the different compartments during an

early and late growing season rain event. Each symbol represents a single sample, and the crossbar

depicts the average for each sample type. Significant differences are shown in letter code tested by post

hoc Dunn's multiple comparing ($p < 0.05$). B) Spearman's rank correlation (r_s) between estimated

bacterial cell numbers and TOC and TN of water samples per sampling event in May and September.

Figure 3: Bacterial communities across rainwater (n = 8), leaves (n = 48), throughfall (n = 20), stemflow (n = 8),

seepage (n = 25) and soil (n = 34) samples. A) Observed ASV richness (left panel) and Shannon index (right

panel). Single data points are shown as dots for samples from May and triangles from September

(crossbar represents mean of values). Letter code indicates significant differences between types and

indexes after Dunn's post hoc test ($p < 0.05$). B) Distribution of most abundant bacterial families depicted

as NMDS based on Bray-Curtis dissimilarities across all samples. Results of PERMANOVA (9999

permutations) are displayed as Venn diagram, asterisks depict confidence interval at $p < 0.001$ for each

accounting factor. C) Bacterial community composition of most abundant bacterial families with more

than 1 % absolute read abundance per event. Single stacked bars show the relative read abundance of

16S rRNA gene sequences on family level.

Figure 4: Prediction of metabolic functions of bacterial communities across the different sample types and the

two time points. A) Abundance of KEGG orthologues (KO) corresponding to genes for enzymes of

selected metabolic pathways. The size of the bubbles depicts the mean relative KO abundance of each

sample type. B) Proportions of bacterial communities employing the predicted KO. Small dots depict the

mean relative abundance of bacterial classes per sample and the bigger dots represent the mean relative

abundance for a given sample type. Letter code displays significant differences between sample types

within each pathway based on post hoc Dunn's multiple comparing ($p < 0.05$).

Figure 5: Phylogeny and relative abundance of bacterial core taxa across rainwater, phyllosphere, throughfall,

stemflow and seepage samples during both sampling time points. The tree is based on a neighbor-joining

model of a 16S rRNA gene alignment. The color and shape of each node show the presence of an ASV on

family-level and on genus-level, respectively, at a given sample type. The size of nodes depicts the mean

relative abundance of each ASV across the sample types. Stacked bars display the mean relative

abundance of each ASV across all samples from both sampling points of rain, phyllosphere, throughfall,

stemflow and seepage communities.

Figure 1

Figure 1: Study site and study design for sample collection at the Hainich Critical Zone Exploratory in
western Thuringia, Germany. **A)** Sampling was conducted at an open area-site for rainwater
representative of above-canopy precipitation and at a mixed beech forest site for phyllosphere,
throughfall, stemflow, and seepage and soil sampling at different depths. **B)** Number of samples for
each sample type across two sampling time points. **C)** Conceptional representation of samples
spanning from the tree canopy down to the soil, and photographs of the two tree species, sample
collectors and a typical soil core.

Figure 2

Figure 2: Bacterial transport across the different compartments, and contributions and correlations to
transported carbon and nitrogen. **A)** Fluxes of TOC and TN expressed per m^2, bacterial contributions
to TOC and TN content, and fluxes of bacterial cells per m^2 across the different compartments during
an early and late growing season rain event. Each symbol represents a single sample, and the crossbar
depicts the average for each sample type. Significant differences are shown in letter code tested by
post hoc Dunn's multiple comparing ($p < 0.05$). **B)** Spearman's rank correlation (r_s) between estimated
bacterial cell numbers and TOC and TN of water samples per sampling event in May and September.

Figure 3

Figure 3: Bacterial communities across rainwater (n = 8), leaves (n = 48), throughfall (n = 20), stemflow
(n = 8), seepage (n = 25) and soil (n = 34) samples. **A)** Observed ASV richness (left panel) and Shannon
index (right panel). Single data points are shown as dots for samples from May and triangles from
September (crossbar represents mean of values). Letter code indicates significant differences between
types and indexes after Dunn's post hoc test (p < 0.05). **B)** Distribution of most abundant bacterial
families depicted as NMDS based on Bray-Curtis dissimilarities across all samples. Results of
PERMANOVA (9999 permutations) are displayed as Venn diagram, asterisks depict confidence interval
at p < 0.001 for each accounting factor. **C)** Bacterial community composition of most abundant bacterial
families with more than 1 % absolute read abundance per event. Single stacked bars show the relative
read abundance of 16S rRNA gene sequences on family level.

Figure 4

Figure 4: Prediction of metabolic functions of bacterial communities across the different sample types
and the two time points. **A)** Abundance of KEGG orthologues (KO) corresponding to genes for enzymes
of selected metabolic pathways. The size of the bubbles depicts the mean relative KO abundance of
each sample type. **B)** Proportions of bacterial communities employing the predicted KO. Small dots
depict the mean relative abundance of bacterial classes per sample and the bigger dots represent the
mean relative abundance for a given sample type. Letter code displays significant differences between
sample types within each pathway based on post hoc Dunn's multiple comparing ($p < 0.05$).

Figure 5

Figure 5: Phylogeny and relative abundance of bacterial core taxa across rainwater, phyllosphere,
throughfall, stemflow and seepage samples during both sampling time points. The tree is based on a
neighbor-joining model of a 16S rRNA gene alignment. The color and shape of each node show the
presence of an ASV on family-level and on genus-level, respectively, at a given sample type. The size
of nodes depicts the mean relative abundance of each ASV across the sample types. Stacked bars
display the mean relative abundance of each ASV across all samples from both sampling points of rain,
phyllosphere, throughfall, stemflow and seepage communities.

bulk precipitation

F. sylvatica   *A. pseudoplatanus*

24x   leaves

2x   throughfall

2x   stemflow

4x   litter · 0-4 cm · 0-16 cm · 0-30 cm   seepage

12x   0-4 cm · 4-16 cm · 16-30 cm   soil

open area site

forest site

© bing maps

700 m

C)

early rain
late
early leaves
late
early throughfall
late
early stemflow
late
early seepage
late
early soil

Hymenobacteraceae   Microbacteriaceae
Sphingomonadaceae   Spirosomaceae
Acetobacteraceae
Beijerinckiaceae
Morganellaceae
Acidobacteriaceae_(Subgroup_1)
Acidothermaceae
Comamonadaceae   Bryobacteraceae
Xanthobacteraceae
Erwiniaceae
Diplorickettsiaceae
Sphingobacteriaceae   Solibacteraceae
Oxalobacteraceae   Chitinophagaceae
Unknown_Family
Flavobacteriaceae   Caulobacteraceae
Yersiniaceae
Pseudomonadaceae

ASV richness   Shannon

stress: 0.097
Bray ~ Type, $p < 0.001$, R^2 = 0.89

Type
25.2%*** Growing season
2.6%***2.1%***
0.4%*
0.5%* 0.1%
Tree species
0.2%**
residuals: 68.9%