Germline de novo mutation rate of the highly heterozygous amphioxus genome

Jing Xue[1,2*], Lei Tao[1,*],Junwei Cao[3], Guang Li[3,#], Cai Li[1,#]

[1]State Key Laboratory of Biocontrol, School of Life Sciences, Guangdong Provincial Key Laboratory for Aquatic Economic Animals, Sun Yat-sen University, Guangzhou, Guangdong, China

[2]Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, Guangdong, China

[3]School of Life Sciences, Xiamen University, Xiamen, Fujian, China

[*]contributed equally

[#]corresponding authors

Abstract

Germline de novo mutations (DNMs) are the ultimate source of heritable variation,

yet their occurrence in highly heterozygous genomes remains poorly understood.

Amphioxus, an early-branching chordate with exceptionally high genomic

heterozygosity (3.2~4.2% in sequenced species), offers a unique model to explore

mutational dynamics in such contexts. It is unclear whether the high heterozygosity in

amphioxus is due to a large effective population size, an increased mutation rate, or

both. Here, we perform deep short-read whole genome sequencing of a two-

generation pedigree of the amphioxus *Branchiostoma floridae* comprising two parents

and 104 offspring, and develop a framework based on allele-aware parental

assemblies as the reference to accurately identify DNMs. We detect 256 high-

confidence DNMs, yielding a genome-wide mutation rate of 4.41×10^{-9} per base

per generation which is comparable to that of vertebrates. By combining this estimate

with observed genomic heterozygosity of *B. floridae*, we obtain an effective

population size of ~2.2 million, indicating that its extreme heterozygosity mainly

results from a large effective population size. Furthermore, we observed a maternal

mutation bias and increased mutation rates in coding regions, suggesting unique

mutational mechanisms in amphioxus. Our study not only provides the first mutation

rate estimate for amphioxus but also offers a scalable strategy for studying DNMs in

highly heterozygous genomes, shedding light on the evolution of mutation rates

across chordates.

Main text

Introduction

Germline de novo mutations (DNMs) differ from somatic mutations in that they can be transmitted across generations, serving as the ultimate source of genetic variation and fueling biological evolution (Lynch 2010). These mutations arise as a consequence of DNA damage or replication errors, and because most mutations are relatively deleterious to the individual, organisms incur considerable adaptive costs to minimize such errors and damage, thereby maintaining the number of DNMs at an extremely low level, approximately 10^{-8} to 10^{-9} per site per generation (Lynch 2010).

The de novo mutation rate (μ) measures how often DNMs occur. Its variation across genomic positions, developmental stages, parental origins, and mutation types are closely linked to the underlying mutational mechanisms. For example, in mammals the C>T mutation rate at CpG sites is about one order of magnitude higher than the genome-wide average, which is mainly caused by spontaneous deamination of methylated cytosines (Hodgkinson and Eyre-Walker 2011). A recent comparative study in amniotes showed that while both sexes exhibit similar mutation rates during early development, the male germline exhibits a higher mutation rate following sexual differentiation (de Manuel et al. 2022), indicating influence of different developmental stages and gametogenesis of different sexes on mutational occurrence.

Amphioxus (cephalochordates) represents the earliest-branching lineage of extant chordates, diverging from the common ancestor of vertebrates and tunicates over 500 million years ago, serving as a key model for understanding vertebrate origins. Amphioxus possesses a body plan that is similar to vertebrates but markedly more simplified, and its chromosomal and genomic sequences have undergone far fewer lineage-specific changes compared with those of tunicates (Putnam et al. 2008).

Intriguingly, amphioxus exhibits an extremely high level of genomic heterozygosity (3.2~4.2% in sequenced species), one of the highest among sequenced chordates (Bi et al. 2020). Under the neutral theory of molecular evolution, the equilibrium level of heterozygosity is proportional to both mutation rate and effective population size. In the absence of mutation rate data, it is unclear whether the high heterozygosity in

amphioxus is due to a large effective population size, an increased mutation rate, or
both.

Therefore, measuring the magnitude and distribution of DNMs in the amphioxus
genome is crucial for precisely estimating the mutation rate and understanding
underlying mutational mechanisms. With an accurate estimate of the mutation rate,
we can also uncover the reason why the amphioxus genome has such an astonishingly
high level of heterozygosity.

To date, no study has investigated DNMs in amphioxus. A major obstacle is its high
heterozygosity, which complicates key genomic analysis steps such as genome
assembly, read alignment, and variant detection. Studies in macaques have shown that
even gold-standard pipelines for DNM detection produce elevated error rates in
highly heterozygous species (Bergeron et al. 2022).

In this study, we sequenced a two-generation pedigree (two parents and 104 F1
offspring) of *Branchiostoma floridae*, comprising 104 trios, to systematically study
DNMs in the amphioxus. To address the challenges in alignment and variant detection,
we developed a workflow leveraging the high heterozygosity to assemble the contig-
level haplotypes of parental genomes, thereby allowing mapping offspring reads to
the parental genomes for identifying DNMs. Using this approach, we successfully
identified 256 high-confidence DNMs in amphioxus. We estimated its mutation rate
to be 4.41×10^{-9} per base per generation, which is comparable to those reported in
fish and other vertebrates. With the mutation rate, we estimated the effective
population size of *B. floridae* wild populations to be 2,199,546, one of the largest in
surveyed chordates. Therefore, the high heterozygosity in amphioxus is mainly caused
by the large effective population size. Furthermore, we discovered that multiple
aspects of DNMs in amphioxus differ from those observed in vertebrates, implying
some unique mutational mechanisms in this lineage.

Results

A strategy based on parental genome assembly for detecting DNMs

To identify DNMs, we performed deep short-read whole genome sequencing for two
parental individuals and 104 F1 offspring of a *B. floridae* family, yielding an average
coverage of 81× per sample (ranging from 50× to 144×) and high sequencing quality

(Supplementary Figure 1). The high sequencing depth enabled us to do contig-level

de novo assembly (Figure 1 A).

Figure 1 Schematic of the workflow for detecting DNMs. First, we collected DNA
samples from a *B. floridae* family of two parents and 104 F1 offspring. We performed
deep short-read sequencing (>50X) for all individuals. Using the parental WGS data,
we assembled allele-aware parental genomes with Platanus-allee and combined the
resulting assemblies to form a parental reference genome for downstream analyses.
Short reads from both parents and offspring were aligned to this custom reference
using BWA-MEM2. We then computed the read depth distribution for each contig in
both parents and offspring. An example is shown for a pair of paternal bubble contigs.
Based on the read depth distributions in parents and an offspring, we identified the
inherited contig from the father in the offspring. The inheritance states of all parental
contigs and read alignments were further used to identify DNMs in each offspring,
with an example shown in the bottom right panel.

k-mer analysis revealed that the heterozygosity of the parental genomes were

approximately 3% (Supplementary Figure 2). The high level of heterozygosity in the

B. floridae genome poses a challenge for existing DNM detection pipelines (Bergeron

et al. 2022), which struggle to accurately genotype both offspring and parents

simultaneously. Because of low sequence similarity between our sequencing data and

the public reference genome, many reads in our sequencing data cannot be correctly

mapped to the reference genome by common short-read aligners (e.g. BWA), leading

to unreliable variant calling results.

We reasoned that the heterozygosity of *B. floridae* can facilitate allele-aware

assembly of the parental genomes, as many divergent regions in the parental genomes

can be assembled into separated contigs. Using the parental genome assembly as a

custom reference, we should be able to align most offspring and parental reads to the

parental genomes, thereby obtaining high-quality genotype information of offspring

and parents for DNM detection (**Figure 1**).

Therefore, we constructed the genome assembly for each parent using the allele-aware

assembler Platanus_allee (v2.2.2) (Kajitani et al. 2019). We obtained a maternal

genome assembly of 773.8 Mb and a paternal genome assembly of 763.7 Mb,

approximately 1.6 times the length of the previously published haploid genome (489

Mb), indicating that a large fraction of the diploid parental genomes was assembled

into phased alleles. In Platanus-allee, heterozygous regions are assembled as "bubble"

contigs with two allelic sequences, named as "primary" and "secondary" respectively.

Homozygous regions with identical sequences in an individual are assembled as "non-

bubble" contigs (**Figure 1**).

By merging the two parental genome assemblies, we generated a composite parental

reference genome, with detailed assembly statistics provided in **Supplementary**

Table 1. The parental reference genome has a total length of approximately 1.4 Gb,

consisting of 263,358 bubble contigs covering around 856 Mb and 758,976 non-

bubble contigs spanning approximately 780 Mb. The contig N50 metrics of bubble

contigs and non-bubble contigs are 6.5 kb and 7.6 kb, respectively. Most

sequencing reads of the two parents can be aligned to this parental reference genome

(**Supplementary Figure 1**), indicating that the vast majority of genomic segments

have been successfully assembled and are available for downstream analysis.

Figure 2 Estimates of mutation rate and effective population size. (**A**) Per site per generation *de novo* mutation rate (μ) estimates from this study and previous studies (Bergeron et al. 2023). Error bars denote 95% confidence intervals. (**B**) Comparison of mutation rates and effective population sizes across diverse taxonomic groups, using data for other taxa from Bergeron et al. (2023).

Next, we aligned sequencing reads from both offspring and parents to this parental reference genome and detected variants using bcftools (Danecek et al. 2021). A custom filtering pipeline, accounting for factors such as read depth and variant quality (**Supplementary Table 2**), was employed to ensure the accuracy of DNMs (see Methods). After applying the filtering criteria, 1,102 candidate DNMs were retained. Among these DNMs, some false positives that could only be ruled out through visualization of IGV identified by manual check (see Methods). After careful check, a total of 256 DNMs were confirmed across 104 offspring, corresponding to an average of 2.46 DNMs per individual. With an average callable genome size of 558.4 Mb, the mean mutation rate across the genome was calculated to be 4.4×10^{-9} mutations per base per generation, with a 95% confidence interval (CI) of $3.90\text{-}4.98 \times 10^{-9}$, as shown

in **Figure 2A**. The mutation rate of *B. floridae* is comparable to those reported in fish

and other vertebrates (Bergeron et al. 2023).

High heterozygosity of *B. floridae* is mainly attributed to its large effective

population size

It has long been recognized that the heterozygosity of *B. floridae* is much higher than

in other species (Putnam et al. 2008). Yet, since the mutation rate in *B. floridae* was

not accurately quantified, the cause(s) of its high heterozygosity was uncertain. The

expected heterozygosity per nucleotide, expressed as π, can be calculated by $\pi = 4N_e\mu$

under neutral evolution (Kimura 1985), where N_e represents the effective population

size and μ the germline mutation rate. The effective population size ($\pi/4\mu$) calculated

with our mutation rate and the π estimate (3.88%) from Huang et al. (2023) was

approximately 2,199,546. According to Bergeron et al. (Bergeron et al. 2023), the

harmonic mean of N_e over multiple time points estimated by PSMC correlated very

well with N_e estimated by $\pi/4\mu$ ($r^2 = 0.81$). Therefore, we placed *B. floridae*'s N_e and

μ within the context of N_e and μ derived from Bergeron et al. to show its unique

position (**Figure 2B**), indicating that *B. floridae*'s N_e is largest among chordates with

known measurements. Collectively, we concluded that the high heterozygosity of

amphioxus is not due to an elevated mutation rate but a very large effective

population size.

Figure 3 Characteristics of germline de novo mutation in *B. floridae*. (A) Parental
origins of DNMs. **(B)** The male-to-female ratios (α) of DNMs for different groups of
chordates. **(C)** Mutation rates in different gene annotation contexts, separating
paternal (blue) and maternal (red) DNMs. **(D)** Proportions of DNMs in different gene
annotation contexts, separating paternal (blue) and maternal (red) DNMs. Parental
origin differences assessed by χ^2 tests. Parental origin effects in 1mer de
novo mutation spectrum: **(E)** Mutation rates of different substitution types, separating
paternal (blue) and maternal (red) DNMs. **(F)** Proportions of DNMs of different
substitution types, separating paternal (blue) and maternal (red) DNMs. Parental
origin differences assessed by χ^2 tests. **(G)** Counts of sibling-shared and unique
DNMs. **(H)** Proportion of DNMs across shared and unique DNM. Parental origin
differences assessed by χ^2 tests. **(I)** Proportions of DNMs of different substitution
types, separating shared and unique DNMs. Shared and unique DNMs differences for
each substitution type assessed by χ^2 tests. All error bars are represented with 95%
confidence intervals computed using binomial variance, and all p-values are labeled
above the bars.

Characteristics of germline *de novo* mutation in *B. floridae*.

Distinguishing the parental origins of DNMs is crucial for understanding mutational mechanisms; while previous studies could only assign parental origin for 30%–80% of DNMs based on nearby polymorphic markers (Bergeron et al. 2023; Wooldridge et al. 2025), our method inherently provides parental origin information for all detected DNMs in *B. floridae*.

Among the 256 DNMs, 154 originated from the father and 102 from the mother (**Figure 3A**), yielding a male mutation bias coefficient (α) of 0.66. For a cross-species comparison, we integrated several representative vertebrates with known α values (**Figure 3B**). The results show that the α value of *B. floridae* is lower than most vertebrates. Bergeron et al. (2023) reported that on average fish tend to have apparently lower α values (mean $\alpha = 0.8$, 95% $CI = 0.5$–1.4) than other vertebrates (mammals, birds and reptiles). They suggested that difference in gametogenesis seasonal breeding and synchronous ovulation could lead to the lower α values in fish. The low α of amphioxus might be partly due to these reasons.

Due to the fragmented nature of the parental genome assemblies, direct genome annotation was challenging; therefore, we mapped the parental genome coordinates to the published haploid reference genome from Huang et al. (Huang et al. 2023) and used its annotation for downstream functional analyses. Of the total 489 Mb of the haploid reference genome, 363.21 Mb (74.27%) was covered by the contigs of the assembled parental contigs. The parental contigs that were mapped to the haploid reference genome were defined as mappable regions. Among the DNMs located in these regions, 198 were identified, accounting for 77.34% of all DNMs.

Based on the haploid genome annotation (Huang et al. 2023), the genome was divided into four functional regions: coding sequences (CDS), untranslated regions (UTR), intronic regions, and intergenic regions. The results showed that the mutation rate in CDS regions was significantly higher than in other regions for both paternal and maternal DNMs (**Figure 3C**). In terms of DNM proportions, the intergenic region showed a significantly higher proportion of maternal DNMs compared to paternal DNMs, while the opposite trend tends to be in intronic regions (**Figure 3D**). These findings demonstrate that both parental origin and functional genomic context exert complex influences on the accumulation of de novo mutations, likely due to species-

specific differences in reproductive strategies or DNA damage/repair processes in *B.*

floridae.

We further investigate DNM counts of different substitution types. The results

indicate that maternal mutation rates of all substitution types are consistently higher

than paternal ones (**Figure 3E**), and there is no significant difference between sexes

when looking at proportions of DNMs of different substitution types (**Figure 3F**).

Cross-species comparisons of mutational spectra (**Supplementary Figure 3**) revealed

that *B. floridae* exhibits a unique pattern, with a CpG site C>T mutation fraction of

only 5%, significantly lower than the 16–20% observed in typical vertebrates,

whereas its A>G mutation frequency of 31% is consistently higher than the 16-26%

seen in other chordate groups; the lower mutation frequency at CpG sites is probably

due to lower methylation levels at CpG sites in amphioxus (Marletaz et al. 2018).

60 (23%) of DNMs are shared among siblings (named ssDNMs, **Figure 3G**) and the

proportions of shared DNMs in the two parents are similar (**Fig. 3H**). The mutational

spectrum shows a significant increase in the proportions of A>G and C>G mutations

in ssDNMs (**Fig. 3I**).

Discussion

In this study, we generated a pedigree comprising both parents and 104 offspring and

conducted high-depth short-read sequencing for all individuals. Standard approaches

for identifying DNMs typically rely on detecting heterozygous genotypes in offspring

that are homozygous in both parents. However, the high heterozygosity of the *B.*

floridae genome complicates such identification. The large difference between the

commonly used reference genome and re-sequenced individuals increases the chance

of misalignment and false negatives.

However, high heterozygosity can also facilitate allele-aware diploid genome

assembly without requiring long-read or Hi-C data. We assembled haplotype-resolved

diploid genomes at the contig level for each parent and developed a custom mapping

and variant calling pipeline using the parental assemblies as reference. This strategy

resolved the challenges of detecting DNMs in highly heterozygous species like *B.*

floridae. It also allowed us to assign the parent-of-origin for all detected DNMs.

Furthermore, based on the built parental assemblies and read alignments of offsprings,

we were also able to comprehensively investigate the meiotic recombination

landscape in a companion study (Tao et al. 2025).

With an average callable genome size of 558 Mb, we identified 256 DNMs and

estimated the germline mutation rate in *B. floridae* to be 4.41×10^{-9} (95% confidence

interval: $3.90\text{-}4.98\times10^{-9}$) per base per generation. Based on this mutation rate, the

effective population size of *B. floridae* is estimated to be over two million, suggesting

that the high heterozygosity results from a large effective population size. Our study

did not estimate the germline mutation rate for INDELs, as generally poor alignments

around these variants make it challenging to detect de novo INDELs. Future studies

incorporating long-read sequencing technologies may facilitate estimating INDEL

mutation rates.

We further analyzed the parental origin, genomic distribution, mutation spectrum, and

ssDNMs in *B. floridae*, revealing distinctive features in this basal chordate. We

observed a strong maternal bias in DNM origin, with a male-to-female mutation ratio

(α) of 0.66. This is lower than most reported values for vertebrates (de Manuel et al.

2022; Bergeron et al. 2023). Interestingly, we found that coding sequences have

higher mutation rates than intronic and intergenic regions. The overall mutation

spectrum showed no obvious difference between sexes. The low frequency of C>T

transitions at CpG sites is consistent with previously reported low levels of CpG

methylation in amphioxus (Marletaz et al. 2018). The presence of ssDNMs provided a

valuable window into early germline mutational events. The relative proportions of

different parental origins in ssDNMs did not differ significantly from that of non-

shared DNMs, a pattern distinct from that in humans. Taken together, it suggests that

mutational mechanisms involved in early embryo development and later

gametogenesis shaping parental biases in amphioxus are quite different from other

chordate lineages. Future DNM studies spanning different ages and developmental

stages of parents could further clarify the mechanisms of mutagenesis in amphioxus.

In addition, functional genomics studies focused on DNA damage and repair

pathways will be essential to understanding the mutational mechanisms.

In summary, through high-depth whole-genome sequencing and the development of a

novel analysis pipeline, we quantified the germline mutation rate in *B. floridae* using

trio-based sequencing for the first time and resolved the long-standing question of

what causes its high heterozygosity. We also explored DNMs from multiple

perspectives, including parental origin, gene annotation contexts, mutation spectrum
and shared DNMs among siblings. As amphioxus represents the most basal living
chordate, some of the unique DNM features observed in this study may reflect
ancestral characteristics of germline mutagenic processes. The DNM analysis
framework based on assembling parental genomes for read alignment and variant
calling is cost effective and can be generalized to other highly heterozygous species.

Materials and Methods

Amphioxus Culture

Amphioxus (*B. floridae*) strains obtained from a stock maintained by Dr. Jr-Kai Yu
originating from Tampa, Florida, were housed in a temperature-controlled facility at
Xiamen University, which maintained a 14 h light and 10 h dark cycle and ambient
seawater temperature at 19°C (Li et al. 2012). A pair of parental individuals was
selected for crossing. Fertilized eggs were subsequently cultured following the
cultivation strategy described previously until they reached the designated sampling
stage. A total of 104 F1 offspring were generated through this process. To capture
distinct developmental time points, the offspring were snap-frozen in liquid nitrogen
at two specific post-growth intervals: 6 months and 1 year. The first batch, comprising
35 individuals, was preserved at the 6-month time point, and the second batch,
consisting of 71 individuals, was preserved at the 1-year time point.

Sample collection and library construction

We extracted high molecular weight genomic DNAs from the muscle tissues of a
single individual using the DNeasy Blood & Tissue Kit (QIAGEN, Valencia, CA),
and inspected the DNA quality by Qubit 2.0 Fluorometer (Thermo Fisher Scientific,
Waltham, MA) and 2100 Agilent Bioanalyzer (Agilent). After Sample QC, gDNA
was fragmented by ultrasound on Covaris E220 (Covaris, Brighton, UK), then
selected to 300 bp - 500 bp using magnet beads size selection. The selected DNA
fragments were then end-repaired and A-tailed. The indexed adaptors were ligated to
both ends of the DNA fragments. The ligation product was amplified by PCR, and
circularized to get single-stranded circular (ssCir) library. The ssCir library was then
amplified through rolling circle amplification (RCA) to obtain DNA nanoball (DNB).

The DNB was then loaded to flowcell, and sequenced by DNBSEQ Platform.

Parental genome assembly

Genome heterozygosity of parental individuals was estimated via k-mer analysis of
Illumina short-read data using GenomeScope (v2.0) (Ranallo-Benavidez et al. 2020).
Raw short-read sequences were quality-processed with Fastp (v0.24.0) (Chen 2023),
implementing: (1) elimination of reads with >10% ambiguous bases ('N'), and (2)
retention of reads exhibiting a minimum mean Phred quality score of 20 (Q20).
Adapter sequences were concurrently trimmed. These processed reads were subjected
to allele-aware diploid genome assembly using Platanus_allee (v2.2.2) (Kajitani et al.
2019) through a two-stage workflow: First, the *de novo* assembly phase was executed
with the command "assemble -m 200 -t 20"; second, haplotype resolution was
performed via the phase module using "phase -i 3 -t 20". Contigs shorter than 150 bp
in length were subsequently filtered from both parental phased assemblies. Initial
assembly quality metrics, including contig N50, total length, and GC content, were
comprehensively evaluated with QUAST (v5.3.0) (Gurevich et al. 2013) and seqkit
(v2.9.0) (Shen et al. 2016).

Following the phased assembly procedure, contigs underwent automated
classification by Platanus_allee. In the contig-level assemblies generated by
Platanus_allee, contigs were classified into two distinct categories: (1) *bubble*
contigs harboring allelic variants, which were further stratified as *primary contigs*
(longer allelic sequences) and *secondary contigs* (shorter counterparts), and (2) *non-*
bubble contigs designated by the assembler as lacking allelic diversity. However,
through comprehensive whole-genome alignment analyses, we discovered that a
subset of non-bubble contigs exhibited unexpected sequence overlaps with other
contigs across the assembly, indicating residual heterozygous regions or potential
mis-assemblies unresolved by the phasing algorithm.

Finally, the filtered phased diploid contig sets derived from paternal and maternal
genomes were concatenated to construct a custom reference genome (hereafter termed
parental reference genome), which were used in many downstream analyses.

Alignment of offspring and parental reads to parental reference genome and
variant calling

Processed short reads from two parents and offspring were aligned to the parental

reference genome assembly using BWA-MEM2 (v2.2.1) (Vasimuddin et al. 2019)

with parameters "-M -k 19". Post-alignment processing entailed coordinate-based

sorting and removal of PCR duplicates via samtools (v1.21) (Li et al. 2009) and

Picard Tools in GATK4 (v4.6.1.0) (Van der Auwera and O'Connor 2020) with default

parameters. Per-base depths across the whole reference genome and individual-level

average sequencing depth metrics were subsequently computed using mosdepth

(v0.3.10) (Pedersen and Quinlan 2018) with default parameters.

Based on the read alignments above, we used bcftools (v1.21) (Danecek et al. 2021)

to identify small variants. First, we generated "pileup" files for the set of all samples

by running "bcftools mpileup" with parameters "–annotate FORMAT/AD,

FORMAT/ADF, FORMAT/ADR, FORMAT/DP, FORMAT/SP, INFO/AD,

INFO/ADF, INFO/ADR" on all input bam files. With the "pileup" files as input, we

ran "bcftools call -mv" to generate VCF files, which contain SNVs and short

insertions/deletions, for 104 offspring and two parents.

386 Inferring inheritance states of parental contigs in offspring genomes based on

387 read alignments against the parental reference genome

According to the principles of Mendelian inheritance, genomic reads in the offspring

are derived from the gene pool of two parental genomes. As described in previous

sections, we obtained the paternal and maternal diploid genome assemblies as sets of

haplotype-resolved contigs. By mapping sequencing reads of an offspring back to the

parental reference genome (combining paternal and maternal genome assemblies), we

can determine which allele of a specific locus (bubble or non-bubble contigs) of a

given parent is inherited by an offspring. More specifically, to determine whether a

particular contig from a given parent is inherited by a specific offspring, we compare

the similarity of read depth distribution profiles of that contig between the offspring

and each parent. For a contig with specific parental origin, the depth distribution of

the offspring's reads aligned to this contig is expected to more closely resemble the

depth distribution of the reads from the parent-of-origin rather than that from the

opposite parent.

To quantify depth distribution, each contig was divided into 50-bp bins, discarding

any terminal bin shorter than 50 bp. The mean depth within each bin was calculated

for each individual, generating a depth distribution vector per individual across the

contig. The similarity between offspring and parental vectors was quantified using a

weighted measure combining: (1) Cosine similarity for bins with non-zero depth in

both individuals being compared, and (2) Jaccard similarity for bins with zero

coverage in at least one of the two individuals (accounting for regions of no aligned

reads). A parental contig was considered inherited by an offspring if its depth

distribution vector in that offspring showed greater similarity to the vector of its

specific parental source than to the other parent. Parental contigs exhibiting a depth

less than 1/4 of an offspring's mean depth or a depth less than 10 were classified as

low-depth and thus non-inherited in the specific offspring.

However, the similarity of some contigs between the parental genomes, probably

arising from shared ancestry, could confound the inheritance calls derived from the

offspring-parent depth distribution similarity algorithm (**Supplementary Figure 4**).

Since direct pairwise genome sequence alignment of two parental genomes is difficult

for assessing parental contig similarities due to potential length disparities of

orthologous contigs in different parental assemblies, we used the similarity of read

depth distributions (calculated using the same algorithm employed for offspring-

parent comparisons) when mapping each parent's reads *to the same contig* as a robust

proxy. If the depth distribution similarity between the two parents for a specific contig

exceeds 0.9, the inheritance calls for that contig in the offspring are deemed unreliable

due to indistinguishable parental origins, necessitating further validation with

additional data.

Because a given diploid offspring can inherit only one allele from each parent at a

locus, determining the inheritance status of one contig within a pair of bubble contigs

from a parent inherently resolves the inheritance status of both contigs in the pair. In

other words, the inheritance of one contig in a bubble pair can be inferred based on

the inheritance of its counterpart. Therefore, we further took advantage of the very

low read depths in some loci to conclude that an allele has not been inherited; under

such circumstances, the alternative allele with aligned reads must be inherited by the

offspring. In contrast, for non-bubble contigs, the inability to identify a second allele

precludes this method of assessing similarity between parental contigs, and therefore

these contigs are excluded from further analysis. Finally, all scenarios arising from the

interplay between inheritance and contig similarity, which lead to differences in depth

distributions in offspring, are comprehensively visualized in Supplementary Figure 4.

Based on the inheritance states of parental contigs in each offspring, we in fact have
reconstructed the offspring genomes with phased alleles in the majority of regions.
Such phased offspring genomes can facilitate downstream analyses.

Determining the callable genome

Estimating mutation rates requires determining the length of genomic regions within
each offspring in which DNMs can be confidently identified, defined as the callable
genome size. In each offspring, the callable genome is primarily composed of the
inherited genomic segments defined in the previous section. However, the accurate
inference of the inheritance of non-bubble contigs with high similarity between the
two parents proved challenging. For non-bubble contigs lacking pre-existing allelic
data in the parental genomes, the allele-depth-distribution-based method for inferring
contig inheritance patterns is inapplicable. Consequently, determining the inheritance
of homologous contigs within non-bubble contigs becomes challenging. We therefore
implemented stricter filtering criteria, excluding both non-bubble contigs with depth
distribution similarity ≥0.5 from the callable genome regions.

Additionally, we removed sites in repetitive regions, as they are prone to erroneous
alignments and often exhibit abnormally high alignment depths. Sites with too many
reads may represent problematic repetitive regions (Li 2014). Specifically, based on
the observed depth distribution in each offspring, any regions with a read depth
exceeding three times the individual's average were excluded.

Finally, by integrating these filters based on both inheritance and sequencing depth,
we obtained the callable genome size for each offspring ranging from 521.70 Mb to
580.88   Mb, accounting for approximately 53–59% of the diploid genome size.

Identifying the candidate DNMs

For each parent-offspring trio, the variants in three VCF files of each trio were filtered
to a subset of single nucleotide variants (SNVs) by bcftools (v1.21) (Danecek et al.
2021).

Variants were retained only if they met all computational criteria: (1) homozygous
alternative allele status in the offspring with read depth >10×; (2) the variant was

located exclusively within the predefined callable genome regions; (3) there were no

more than two alternative allele observations in each parental BAM file, as verified by

our custom Python allelic-depth quantification pipeline; and (4) the variant had a

minimum QUAL score 220, ensuring high-confidence base calls; (5) no proximity to

indels (±10 bp), (6) not located in terminal contig regions (within 200 bp of contig

ends); (7) variants not at the sites experiencing putative gene conversions, as gene

conversions lead to read alignments resembling DNMs.

After applying these filters, the resulting set of candidate DNMs was further evaluated

by importing both parental and offspring BAM files into IGV (v2.19.2)

(Thorvaldsdottir et al. 2013), and using IGVtools to generate screenshots for each

candidate DNM in both the offspring and the parents. Subsequent manual inspection

led to the removal of any candidate DNMs if they met one or more of the following

criteria (1) structural variant/recombination breakpoint neighborhoods, and (2)

complex repetitive domains in short contigs (<5 kb) with ambiguous alignments. All

filtered variants were categorized by exclusion criteria, with counts of discarded sites

per filter tier provided in Figure S2. These steps yielded a DNM dataset for each

individual. Individual mutation rates were calculated by dividing the number of

DNMs detected in each offspring by the corresponding callable genome size, and the

final genome-wide mutation rate was calculated by the number of all DNMs divided

by sum of callable genome sizes in offspring. Notably, the callable genome size was

not multiplied by two, since the reconstructed inherited genomic segments effectively

represent an approximately diploid genome for each offspring.

Because the coordinates of DNMs are located on the parental genomic contigs, the

parental origin of the contig harboring each DNM naturally reveals its source,

allowing for unequivocal determination of the DNM's parental origin without any

further processing. This same principle applies to the callable genome as well.

Analysis of the effective population size

The average long-term N_e was estimated using the formula $N_e = \pi / 4\mu$ (Kimura

1985), where μ is the mutation rate estimated by our work (4.41×10^{-9} per site per

generation) and π from a previous study (Huang et al. 2023). For comparison, the

harmonic mean of historical effective population size in vertebrates calculated using

PSMC (Li and Durbin 2011) was obtained from data published by Bergeron et al.

(Bergeron et al. 2023). The rationale for choosing this dataset is that the authors found

a strong correlation (r^2 = 0.83) between effective population sizes computed via

PSMC and long-term effective population sizes, demonstrating that this dataset can

serve as a robust proxy for long-term effective population size.

Aligning parental genome assemblies to the published *B. floridae* genome

Owing to inherent limitations of short-read sequence, the assembly of parental

genomes could not achieve chromosome-scale or high continuity, posing challenges

for some downstream analyses. Furthermore, defining orthologous relationships

between contigs of two parents in a specific locus based solely on pairwise alignments

of contigs, as previously mentioned, was complicated by the variability in the lengths

of assembled contigs derived from that locus. To address these issues, we aligned the

contigs of parental genomes to a published *B. floridae* haploid reference genome

(Huang et al. 2023). After testing multiple alignment tools under various parameter

configurations, Minimap2 (v2.24) (Li 2021) was selected as the optimal aligner, with

empirically optimized parameters: "-x asm20 --secondary=no". Resultant alignments

were processed through sequential refinement: coordinate sorting via samtools (v1.21)

(Li et al. 2009), (2) removal of multi-mapping regions using Sambamba (v0.8.2)

(Tarasov et al. 2015). A custom script was employed to establish the mapping

relationship between each parental genome and the haploid reference genome. This

approach yielded a multiple genome alignment that aligned positions across the

paternal, maternal, and reference genomes. As a result, the majority of parental

genomic positions were successfully anchored onto the reference, thereby obtaining

orthologous genomic coordinates across the parental genomes and facilitating

subsequent analyses.

Comparative analysis of mutation rates in different contexts

To compare mutation rates in different contexts (e.g., different gene features), we

utilized the coordinate mapping relationships described in the previous section. We

mapped the gene functional annotations from the published *B. floridae* reference

genome (Huang et al. 2023) onto the parental genomes. In doing so, the coordinates

for CDS, intronic, UTRs, and intergenic regions were converted to those of the

parental genomes. This allowed us to determine the lengths of each gene feature

where DNMs and callable genome regions were present, enabling the calculation of

gene feature-specific mutation rates for each individual.

To generate the mutation spectrum, the DNMs were also stratified into six groups

according to their mutation types (A > C, A > T, A > G, C > A, C > G, C > T), with a

special inspection on C > T mutations at CpG sites (the 7th group). Using custom

scripts, we extracted CpG positions from the parental genomes, thereby allowing us to

distinguish between CpG and non-CpG sites.

References

Bergeron LA, Besenbacher S, Turner T, Versoza CJ, Wang RJ, Price AL, Armstrong E, Riera
M, Carlson J, Chen HY et al. 2022. The Mutationathon highlights the importance of
reaching standardization in estimates of pedigree-based germline mutation rates. *Elife*
11.
Bergeron LA, Besenbacher S, Zheng J, Li P, Bertelsen MF, Quintard B, Hoffman JI, Li Z, St
Leger J, Shao C et al. 2023. Evolution of the germline mutation rate across
vertebrates. *Nature* **615**: 285-291.
Bi C, Lu N, Han T, Huang Z, Chen JY, He C, Lu Z. 2020. Whole-Genome Resequencing of
Twenty Branchiostoma belcheri Individuals Provides a Brand-New Variant Dataset
for Branchiostoma. *Biomed Res Int* **2020**: 3697342.
Chen S. 2023. Ultrafast one-pass FASTQ data preprocessing, quality control, and
deduplication using fastp. *Imeta* **2**: e107.
Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
McCarthy SA, Davies RM et al. 2021. Twelve years of SAMtools and BCFtools.
Gigascience **10**.
de Manuel M, Wu FL, Przeworski M. 2022. A paternal bias in germline mutation is
widespread in amniotes and can arise independently of cell division numbers. *Elife* **11**.
Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* **29**: 1072-1075.
Hodgkinson A, Eyre-Walker A. 2011. Variation in the mutation rate across mammalian
genomes. *Nat Rev Genet* **12**: 756-766.
Huang Z, Xu L, Cai C, Zhou Y, Liu J, Xu Z, Zhu Z, Kang W, Cen W, Pei S et al. 2023. Three
amphioxus reference genomes reveal gene and chromosome evolution of chordates.
Proc Natl Acad Sci U S A **120**: e2201504120.
Kajitani R, Yoshimura D, Okuno M, Minakuchi Y, Kagoshima H, Fujiyama A, Kubokawa K,
Kohara Y, Toyoda A, Itoh T. 2019. Platanus-allee is a de novo haplotype assembler
enabling a comprehensive access to divergent heterozygous regions. *Nat Commun* **10**:
1702.
Kimura M. 1985. *The neutral theory of molecular evolution*. Cambridge university press.
Li G, Yang X, Shu Z, Chen X, Wang Y. 2012. Consecutive spawnings of Chinese amphioxus,
Branchiostoma belcheri, in captivity. *PLoS One* **7**: e50838.
Li H. 2014. Toward better understanding of artifacts in variant calling from high-coverage
samples. *Bioinformatics* **30**: 2843-2851.
Li H. 2021. New strategies to improve minimap2 alignment accuracy. *Bioinformatics* **37**:
4572-4574.
Li H, Durbin R. 2011. Inference of human population history from individual whole-genome
sequences. *Nature* **475**: 493-496.
Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
Genome Project Data Processing S. 2009. The Sequence Alignment/Map format and

SAMtools. *Bioinformatics* **25**: 2078-2079.

Lynch M. 2010. Evolution of the mutation rate. *Trends Genet* **26**: 345-352.

Marletaz F, Firbas PN, Maeso I, Tena JJ, Bogdanovic O, Perry M, Wyatt CDR, de la Calle-
Mustienes E, Bertrand S, Burguera D et al. 2018. Amphioxus functional genomics
and the origins of vertebrate gene regulation. *Nature* **564**: 64-70.

Pedersen BS, Quinlan AR. 2018. Mosdepth: quick coverage calculation for genomes and
exomes. *Bioinformatics* **34**: 867-868.

Putnam NH, Butts T, Ferrier DE, Furlong RF, Hellsten U, Kawashima T, Robinson-Rechavi
M, Shoguchi E, Terry A, Yu JK et al. 2008. The amphioxus genome and the
evolution of the chordate karyotype. *Nature* **453**: 1064-1071.

Ranallo-Benavidez TR, Jaron KS, Schatz MC. 2020. GenomeScope 2.0 and Smudgeplot for
reference-free profiling of polyploid genomes. *Nat Commun* **11**: 1432.

Shen W, Le S, Li Y, Hu F. 2016. SeqKit: A Cross-Platform and Ultrafast Toolkit for
FASTA/Q File Manipulation. *PLoS One* **11**: e0163962.

Tao L, Xue J, Cao JW, Li G, Li C. 2025. Direct detection of meiotic recombination events in
the highly heterozygous amphioxus genome. *bioRxiv*.

Tarasov A, Vilella AJ, Cuppen E, Nijman IJ, Prins P. 2015. Sambamba: fast processing of
NGS alignment formats. *Bioinformatics* **31**: 2032-2034.

Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV):
high-performance genomics data visualization and exploration. *Brief Bioinform* **14**:
178-192.

Van der Auwera GA, O'Connor BD. 2020. *Genomics in the Cloud: Using Docker, GATK,*
and WDL in Terra. O'Reilly Media.

Vasimuddin M, Misra S, Li H, Aluru S. 2019. Efficient Architecture-Aware Acceleration of
BWA-MEM for Multicore Systems. In *2019 IEEE International Parallel and*
Distributed Processing Symposium (IPDPS), doi:10.1109/IPDPS.2019.00041, pp.
314-324.

Wooldridge TB, Ford SM, Conwell HC, Hyde J, Harris K, Shapiro B. 2025. Direct
Measurement of the Mutation Rate and Its Evolutionary Consequences in a Critically
Endangered Mollusk. *Mol Biol Evol* **42**.

Zhang C, Reid K, Sands AF, Fraimout A, Schierup MH, Merila J. 2023. De Novo Mutation
Rates in Sticklebacks. *Mol Biol Evol* **40**.

611