Direct detection of meiotic recombination events in the highly heterozygous amphioxus genome

Lei Tao[1,*], Jing Xue[1,2*], Junwei Cao[3], Guang Li[3,#], Cai Li[1,#]

[1]State Key Laboratory of Biocontrol, School of Life Sciences, Guangdong Provincial Key Laboratory for Aquatic Economic Animals, Sun Yat-sen University, Guangzhou, Guangdong, China

[2]Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou, Guangdong, China

[3]School of Life Sciences, Xiamen University, Xiamen, Fujian, China

[*]contributed equally

[#]corresponding authors

Abstract

Amphioxus, a basal chordate with highly heterozygous genomes (3.2~4.2% in

sequenced species), represents a key model for understanding vertebrate origins.

However, the extreme heterozygosity poses challenges for many genomic analyses,

including studying meiotic recombination. Here, we present a novel bioinformatic

pipeline that enables direct detection of crossover (CO) and non-crossover (NCO)

recombination events using short-read whole-genome sequencing of a two-generation

pedigree (two parents and 104 F1 offspring) of the amphioxus *Branchiostoma*

floridae. Using parental assemblies generated by Platanus-allee as a custom reference

for read alignment, we tracked inheritance patterns in offspring and phased

contig-level haplotypes in parents, allowing us to detect recombination events. We

identified 2,329 paternal and 2,288 maternal COs, yielding recombination rates of

4.57 cM/Mb and 4.49 cM/Mb, respectively. We found CO coldspots spanning >140

Mb in each parent and these are likely associated with large-scale heterozygous

inversions. CO rates were positively correlated with transposable element and gene

density in both sexes, but showed weak or no correlation with GC content. We further

identified ~10,000 paternal and ~5,800 maternal NCO events, predominantly <200 bp

in length, and found evidence of GC-biased gene conversion. This work provides the

first direct and genome-wide measurement of recombination in amphioxus and

demonstrates how high heterozygosity, often considered a barrier, can be leveraged

for fine-scale recombination mapping. Our findings illuminate conserved and

divergent features of recombination in chordates and establish a framework for

studying recombination in other highly heterozygous organisms.

Keywords: amphioxus, recombination, crossover, non-crossover, highly heterozygous

genomes

Introduction

Cephalochordate amphioxus is the sister group of tunicates and vertebrates, and represents the most primitive living chordates[1]. This key phylogenetic position, together with its simple and slowly evolving morphology and genome structure, makes amphioxus an essential model for studying the origin and evolution of many aspects of vertebrates[2-6].

Genomic research of amphioxus species can offer crucial insights into chordate origin and evolution. The sequenced genomes of amphioxus species span about 500 Mb [7], with chromosome numbers varying among species (18-20 chromosomes) due to chromosomal rearrangements[8]. Notably, amphioxus genomes have a high heterozygosity rate of 3.2-4.2%, significantly higher than humans and most vertebrates, making it one of the most heterozygous known species[8]. This high heterozygosity leads to substantial genomic difference among individuals, posing challenges for traditional variant detection methods which are usually based on short-read alignments to a commonly used reference genome. As a result, despite its extensive use in evolutionary studies, population genomic analysis within amphioxus species remains limited.

Meiotic recombination is a fundamental biological process that ensures the proper segregation of homologous chromosomes during gametogenesis and generates novel allele combinations, thereby fueling genetic diversity and facilitating adaptive evolution across eukaryotes[9]. Our current understanding of amphioxus recombination rates was based on population genomic variations[8], which reflect historical recombination events averaged over sexes and many generations. As a result, direct experimental observations and quantitative measurements of meiotic recombination—including the absolute number of crossovers per meiosis and their distribution along individual chromosomes—remain completely unexplored in amphioxus.

In this study, we generated short-read whole genome sequencing data for a two-generation family of the amphioxus species *Branchiostoma floridae*, including two parents and 104 F1 offspring. Considering the high heterozygosity of the *B.*

floridae genome, we developed a novel bioinformatic pipeline for detecting

recombination events in meiosis of the two parents, including crossover (CO) and

non-crossover (NCO) gene conversion events. Based on the detected recombination

events, we further investigated potential factors influencing recombination in the

amphioxus genome. Our work provides novel insights into the recombination

landscape of amphioxus and facilitate recombination analysis in species with

high-heterozygosity genomes like amphioxus.

Results

The workflow for detecting meiotic recombination events

We generated >50X short-read sequencing data for each of the 106 individuals

(two parents and 104 offspring). The sequencing depth is not only sufficient for

resequencing analysis, but also for high enough for contig-level *de novo* assembly.

Given the high heterozygosity of the amphioxus genome, the existing

chromosome-level genome assembly[7, 8] differs significantly from the sequenced

genomes in our study. When aligning short reads in our study against the previously

used reference genome using short-read aligners, because of low sequence similarity,

many reads can not be correctly mapped, leading to unreliable variant calling results.

To address this issue, we developed a novel strategy for read alignment and

variant calling. Because high heterozygosity can facilitate haplotype assembly, we

first used Platanus-allee[10], an haplotype assembler designed for highly heterozygous

regions, to assemble the parental genomes separately and merged them as a custom

reference genome for read alignment of offspring (**Fig. 1a**). As offspring inherit

genetic information from parents, most reads from offspring can be correctly mapped

to the Platanus-allee-assembled parental genomes, which are though comprised of

relatively short contigs. In Platanus-allee, heterozygous regions are assembled as

"bubble" contigs with two allelic sequences, named as "primary" and "secondary"

respectively. Homozygous regions with identical sequences in an individual are

assembled as "non-bubble" contigs (**Fig. 1a**). More details about the assembly and

alignment processes are given in a companion paper focusing on *de novo* mutations[11].

Based on alignments of offspring reads against parental genomes, we were able to call
variants and determine which parental contigs are inherited by each offspring (more
details are given in the companion paper[11]).

Since non-bubble regions in the parents have identical haplotypes across all
offspring and are not informative for recombination analysis, we thus only focused on
bubble contigs. The paternal genome assembly has 42,378 bubble contig pairs
(~466.3 Mb in total, average contig length of ~5.5 kb), and the maternal one has
43,292 pairs (~389.9 Mb in total, average contig length of ~4.5 kb). For bubble
regions of a specific parent, each offspring can only inherit either the primary or
secondary contig, so such regions can be considered bi-allele markers for phasing
analysis. Using hapi[12] with offspring states of bubble contigs as markers, we
performed parent-level phasing of bubble contigs and reconstructed parental
haplotypes for each parent (**Fig. 1a**). Our method is analogous to that of using gamete
sequencing data to reconstruct parental haplotypes[13, 14], though our method has much
easier sample preparation and affordable sequencing cost. After phasing, by
comparing offspring haplotypes and reconstructed haplotypes from a specific parent,
we can detect CO recombination events occurring in the meiosis of that parent (**Fig.
1a**).

Fig.1 Workflow for detecting recombination events. **a**. Detection workflow for CO

events using bubble contigs as markers. By aligning sequencing reads of offspring

against the assembled parental genomes, the inherited state of each bubble contig pair

of parents in each offspring was determined. Subsequently, the phased contig

haplotypes of each parent were determined with contig haplotypes of offspring

inherited from that parent. Finally, the CO events in meiosis of a specific parent were

identified by comparing parental and offspring haplotypes. The position where an

exemplary CO event occurs in Offspring5 is marked with a red arrow. **b**. Detection of

CO breakpoints within a bubble contigs pair. A breakpoint caused by a CO event will

form four haplotypes in the offspring according to short-reads alignment. **c**. Detection

workflow for NCO events. Within a pair of bubble contigs, a portion of one contig

replaces the allelic region of the other, giving rise to three observable haplotypes in

offspring.

Unlike CO detection based on single nucleotide polymorphism (SNP) markers,

bubble contigs can have internal COs, especially for longer contigs. Therefore, we

also analyzed putative CO-induced breakpoints within contigs based the read

alignment of offspring reads against parental bubble contigs (**Fig. 1b**). After

recombination, related sequences from a bubble contig pair form four possible

haplotypes in the offspring. Though the number of breakpoints that can be found

within contigs is limited, this analysis can complement the CO detection method that

used bubble contigs as markers.

In addition to CO events, we also detected NCO gene conversion events in the

genome. NCO events usually have a smaller impact sequence range. For instance, the

average tract length of NCO events in humans is 459 bp[15]. Due to few informative

markers, we cannot identify NCO events in non-bubble regions, so we focused on

those in bubble regions. To detect NCO events, we compared the primary and

secondary contig in each bubble contig pair and identified high-quality heterozygous

variants as informative markers. An NCO event was considered to have occurred

when the parental phase of first and last markers on a contig remained unchanged

while the middle marker(s) switched to another allelic type (**Fig. 1c**). For NCO events,

we also required that only three haplotypes were observed in the offspring.

Characterization of CO events

Based on the inheritance patterns of bubble contigs in offspring samples, we

reconstructed parental haplotypes for each parent, using bubble contigs as markers.

This allowed us to identify CO events during meiosis in the father and the mother,

respectively. In total, we identified 2,329 CO events from paternal meioses across all

offspring, averaging 22.4 per meiosis (range: 13-37) (**Fig. 2a, Supplementary Table**

1). For maternal meioses, we identified 2,288 CO events, averaging 22.0 per meiosis

(range: 10-37) (**Fig. 2a, Supplementary Table 1**). In most diploid sexually

reproducing species, there is usually at least one CO event per meiosis on

average[16-18]. The CO frequency in this amphioxus family aligns well with this

pattern. As the haploid genome of *B. floridae* is about 490.4 Mb[8], the CO

recombination rate is 4.57 cM/Mb in the father and 4.49 cM/Mb in the mother, higher

than that in mammals (~0.5 cM/Mb) and birds (~2 cM/Mb)[19, 20]. Unlike in humans

and some other vertebrates where a certain specific sex has higher CO rates[21, 22], we

found no significant sex biased CO rate in this *B. floridae* family.

Fig.2 Characterization of CO events in paternal and maternal genomes. **a**.

Numbers of CO events observed in 104 offspring, separating paternal and maternal

COs. **b**. An example of a CO breakpoint within a bubble contigs pair. Sequencing

coverage depth plots are shown for four offspring aligned to the bubble contigs region.

From top to bottom, these four samples respectively inherited the primary contig, the

secondary contig, a new contig formed by the front segment of the primary contig

combined with the rear segment of the secondary contig, and another new contig

formed by the rear segment of the primary contig combined with the front segment of

the secondary contig, representing 4 distinct haplotypes. **c**. Distribution of distances

between all identified within-contig CO breakpoints and their nearest detected CO

events using bubble contigs as markers. **d**. Numbers of identified CO events per

chromosome in the father and the mother. **e**. Correlation between the genomic

distributions of CO events identified in the father and those in the mother. The

window size was set to 5 Mb, with a step size of 1 Mb. Linear regression was

employed to fit the data, with the 95% confidence interval as shaded area. **f**.

Correlation between the distribution of CO events and recombination rates estimated

by Huang et al.[8] The window size was set to 5 Mb, with a step size of 1 Mb. Linear

regression was employed to fit the data, with the 95% confidence interval as shaded

area.

We further identified putative CO breakpoints within bubble contigs. These

recombined regions swapped parts of two paired bubble contigs, creating four

haplotypes in the offspring (**Fig. 2b**). We identified 171 putative breakpoints (95

paternal and 76 maternal). Most breakpoints (~90.6%) were within 1 Mb of the

nearest CO event detected by the bubble contig inheritance method (**Fig. 2c**),

suggesting they represent the same CO events. This also implies that our strategy

using bubble contigs as markers effectively detected most genomic CO events.

Subsequently, we analyzed the distribution of CO events across different

chromosomes. We observed a higher number and density of CO events on certain

chromosomes, such as chromosomes 11, 12, and 14 (**Fig. 2d**). This may relate to

specific genomic characteristics, a topic we will discuss later. Most chromosomes

showed limited sexual dimorphism in the number of CO events (**Fig. 2d**).

Additionally, the distribution of CO events on chromosomes in the father and the

mother exhibited a high correlation (**Fig. 2e**, Pearson's r = 0.66, in windows of 5 Mb).

This suggests that in this amphioxus pedigree, sex difference has limited impact on

both the quantity and chromosomal distribution of CO events. We also compared our

CO results with the population-level CO rates estimated by Huang et al.[8] based on

population genomic variations. The correlation between our pedigree-based CO rates

and previous population-level CO rates is weak (Pearson's r = 0.09, p = 0.045, in

windows of 5 Mb) (**Fig. 2f**), likely due to the fact that the two methods focus on

different CO events. Huang et al. focused on the historical CO events at the
population level which can be affected by evolutionary processes such as selection,
but our data focus on contemporary CO events derived from a single family.

A sex-determining region of *B. floridae* spanning 0–3.1 Mb of chromosome 1620
(resulting from a chr16-chr20 fusion) was reported previously[8]. This species exhibits
a ZW sex determination system, with males being homogametic (ZZ) and females
heterogametic (ZW)[23]. In the analyzed pedigree, CO events were detected near the
sex-determining region in the paternal genome but absent in the maternal genome,
consistent with suppressed recombination in the heterogametic sex.

Impact of heterozygous inversions on the distribution of CO events

When analyzing CO distribution across the genome, we noticed that although the
overall correlation of CO events on chromosomes in the father and the mother was
relatively high, there are large genomic regions without any CO events, which differ
among paternal and maternal genomes (**Fig. 3a**). For each parent, we defined regions
spanning 5 Mb or more with no CO events in any offspring as coldspot regions. The
paternal genome had ~209.1 Mb of CO coldspot regions (genomic coordinates in
Supplementary Table 2), and the maternal genome had ~141.9 Mb (**Supplementary
Table 2**). One of the reasons for the formation of coldspot regions might be the
inhibitory effect on peripheral CO events near the centromere[24, 25], but this cannot
explain the existence of coldspot regions at positions far from the centromere.
Another point that easily comes to mind is that the sparsity of markers (bubble contigs)
in these regions leads to a decrease in the sensitivity of detecting CO events. However,
in many cases, coldspot regions had high marker densities, and COs were also
detected in marker-sparse regions. For example, on chromosome 3 of the maternal
genome, CO events were mainly observed in the marker-sparse half (16.9-32.3 Mb),
while the marker-dense half (0-16.9 Mb) had few COs (**Fig. 3a**). This suggests that
coldspot regions are not simply due to marker sparsity.

Previous studies suggest that large-scale structural variations, especially
heterozygous inversions, can suppress CO events[26-28]. The amphioxus genome

exhibits substantially elevated rates of structural variations, with the rate of

translocations and inversions reported to be approximately 30 times of that observed

in the human-chimpanzee comparison[29]. To check if such variations cause CO

coldspots in amphioxus meiosis, we searched for large-scale inversions in parents

using phased haplotype information. We found that some coldspot regions indeed

harbor large-scale heterozygous inversions (**Supplementary Table 3**). For example,

around the position of 1.4 Mb on maternal chromosome 3, the secondary contig of a

bubble contig pair aligns with this region nicely, but the corresponding primary contig

can be mapped to this region only partially, with the other half being inversely aligned

to a region around 13.5 Mb on the same chromosome (**Fig. 3b**). This implies a

large-scale inversion (involving 1.4 Mb to 13.5 Mb of chromosome 3) in the primary

contig-carrying haplotype, likely suppressing COs during maternal meiosis and

forming a maternal CO coldspot. A similar case was found in chromosome 7 of the

paternal genome (**Fig. 3c**). However, due to the limitations of genome assemblies

from short read sequencing data, we cannot confirm whether all coldspot regions

harbor large heterozygous inversions. Nonetheless, we propose that such

heterozygous inversions are likely one main cause for CO coldspots in the amphioxus

genome.

Fig.3 Distribution of COs on chromosomes and examples of large inversions potentially suppressing COs. **a**. Densities of identified CO events and bubble contig markers along the paternal and maternal genomes, respectively. A window size of 1 Mb was used. Centromeric regions (genomic coordinates from Huang et al.[8]) are

shaded. Centromere positions for chromosomes 7 and 17 were not definitively

identified. (**b**) and (**c**) respectively illustrate examples of heterozygous inversions in

the maternal and paternal genomes. Shown are pairwise alignments of primary and

secondary contigs from one set of bubble contigs against the chromosome-level

reference genome assembly[8].

Association between CO events and genomic features

We further investigated the relationship between CO rates and genomic features,

including transposable element (TE) density, gene density, and GC content[30]. In both

paternal and maternal genomes, TE density showed significant positive correlations

with CO events (Pearson's r of 0.43 in father and 0.40 in mother, **Fig. 4a**), suggesting

that TEs are a likely factor influencing COs in amphioxus. Further analysis of

different TE types showed that SINEs, LINEs, LTRs, MITEs and DNA transposons

all correlate with CO distribution respectively (**Fig. 4b**), without a clear preference for

any specific TE class. Many previous studies found a negative correlation between TE

density and CO rate[31, 32], though some examples of positive correlation were also

reported[33-35], implying the complex relationship between TEs and CO events. Gene

density also correlated with CO events (Pearson's r of 0.28 in father and 0.26 in

mother, **Fig. 4c**), but less strongly than TE density. Regions with high gene density

tend to have higher CO rates, possibly due to more accessible chromatin structure in

gene-dense regions[36], which facilitates the binding of recombination-associated

proteins and promotes recombination. No significant correlation was found between

GC content and paternal COs, while a weak negative correlation was observed in

maternal genome (Pearson's r = -0.12, p-value = 0.0164, **Fig. 4d**). Overall, paternal

and maternal genomes showed similar patterns: higher TE and gene densities tend to

be associated with higher CO rates.

Fig.4 Association between CO events and genomic features. a. Scatter plots displaying the relationship between CO density and TE density, with Pearson's correlation coefficients in the figures. The window size for calculating densities was set to 5 Mb, with a step size of 1 Mb. The blue scatter plot at the top represents paternal data, and the red scatter plot at the bottom represents maternal data. Linear regression was employed to fit the data, with the 95% confidence interval as shaded area. **b**. Pearson's correlation coefficients between CO density and the densities of specific TE types. All correlations shown have p-values < 0.001. **c**. Scatter plots displaying the relationship between CO density and gene density. Window size, dot color and data fitting are the same as in panel a. **d**. Scatter plots displaying the relationship between CO density and GC content. Window size, dot color and data fitting are the same as in panel a.

Characterization of NCO events

Apart from CO events, recombination can generate NCO gene conversion events.
As NCO tract lengths are usually short, detecting NCO events is highly dependent on
the density of informative heterozygous SNP markers surrounding NCOs. The high
heterozygosity of *B. floridae* individuals provides a unique advantage for identifying
NCOs. Due to insufficient SNP markers in non-bubble regions and some bubble
regions with few differences between primary and secondary contigs, we first
identified the genomic regions where NCO events could be detected (see Methods). In
the paternal genome, an average of 73.0 Mb of the genome for each offspring was
analyzable for NCO events, with 9,992 NCOs detected (63-141 per offspring,
averaging 96.1, **Fig. 5a, Supplementary Table 4**). In the maternal genome, an
average of 61.1 Mb for each offspring was analyzable, with 5,850 NCOs found
(25-114 per offspring, averaging 56.3, **Fig. 5a, Supplementary Table 4**). Overall,
more NCO events were observed in the paternal genome. However, unlike CO event
detection with bubble contigs, NCO detection is more affected by marker density.
Thus, it is unclear whether the higher NCO count in the paternal meiosis reflects a
true biological difference like humans[37] or is due to more informative paternal
markers (6.71 million versus 5.78 million markers). Assuming the analyzable regions
are representative of the whole genome, the genome wide NCO:CO ratios are 28.8
and 20.5 for paternal and maternal meiosis respectively. These ratios in humans are
7.84 in male and 3.91 in female[37]. It is also higher than some other vertebrates, such
as zebra finch (5.4)[38]. The high ratios in amphioxus could be due to the amphioxus
genome's high heterozygosity that enhances NCO detection sensitivity. We note that
although it is impossible to distinguish an NCO event from a *de novo* substitution at
the parental heterozygous sites with our method, given the low mutation rate (4.4×
10^{-9} per base per generation[11]), the level of NCO false positives due to *de novo*
substitutions at heterozygous sites should be very low (more information given in the
Methods).
345
346

Fig.5 Detection of NCO events. **a**. Numbers of NCO events identified in the paternal and maternal alleles of 104 offspring. **b**. Length distributions of NCO events in paternal (blue barplots, top) and maternal genomes (red barplots, bottom). **c**. Barplots showing the number of NCO events covered by a specific number of SNP markers. **d**. Conversion spectrum of NCO events. A>C includes A>C in the positive strand and T>G in the negative strand, similarly for other types. **e**. NCO conversion bias. NCO events are divided into single-SNP tracts and multi-SNPs tracts according to the numbers of covered markers. X-axis depicts the proportion of A/T > C/G events among ([A/T > C/G] + [C/G > A/T]) events. The p-values and 95% confidence intervals (error bars) were obtained from an exact binomial test. P-values are

displayed on the right, with '***' representing for p-value<0.001 and '**'

representing for 0.001<p-value<0.01. **f**. The relationship between local CO rates and

NCO rates in the paternal and maternal genomes, with a sliding window size of 5 Mb.

Linear regression was employed to fit the data, with the 95% confidence interval as

shaded area. Scatterplots are shown separately for data with and without CO coldspots.

We further analyzed the impacted sequence lengths and conversion types of

NCO events. Most NCO events in paternal and maternal genomes involve small

genomic regions, with 90.9% being under 200 bp, with the average length of NCO

events being 99 bp (**Fig. 5b**). Most NCO events (83.6%) only covered one converted

marker (**Fig. 5c**). These NCO events were considered to be single-SNP NCOs, with

the rest to be multi-SNP ones. Regarding the conversion types (**Fig. 5d**), we found

that in both paternal and maternal genomes, A>G (or T>C) and C>T (G>A)

conversions were the most frequent. GC-biased gene conversion (gBGC), common in

mammals, favors the fixation of C/G alleles over A/T alleles and mainly occurs in

single-SNP conversion tracts[39]. Except for the maternal multi-SNP NCOs which

showed an AT bias, we discovered GC bias in all other types of NCOs to varying

degrees (**Fig. 5e**), with paternal genomes exhibiting higher GC bias. This indicates

that gBGC also contributes to NCO events in amphioxus.

We also investigated the relationship between local CO and NCO rates in the

paternal and maternal genomes (**Fig. 5f**), with a sliding window size of 5 Mb. We

found no or very weak correlation between local CO and NCO rates when including

the CO coldspot regions. When excluding CO coldspots, we found a relatively weak

positive correlation (Pearson's r = 0.29, p = 1.536e-7) between local CO and NCO

rates in the paternal genome, but no correlation in the maternal genome. This suggests

that the local frequencies of CO and NCO events in this species appear to be largely

independent.

Discussion

In this study, we conducted whole genome sequencing for 106 individuals from a

two-generation *B. floridae* family to systematically analyze CO and NCO events

during male and female meioses. The high heterozygosity of *B. floridae* individuals

allowed us to assemble the paternal and maternal genomes into relatively long contigs

(average length of ~5 kb) with only short read data. Based on the offspring read

alignments with the parental genomes (assembled contigs), we phased and

reconstructed parental haplotypes at the contig level. This method enables phasing of

diploid parents without direct gamete sequencing. This approach leverages the high

heterozygosity of the amphioxus genome, turning a potential analytical challenge into

an advantage. The high genomic heterozygosity also provides a high density of

markers for detecting NCO recombination events. If not sequencing gametes,

previous genomic research directly detecting recombination events usually needed to

sequence pedigrees of three or more generation[30, 40], but our approach only needs

two generations. Thus, our strategy offers a new perspective for studying

recombination in species with a highly heterozygous genome.

Using this approach, we analyzed CO events in paternal and maternal genomes.

The estimated CO recombination rate was higher (paternal: 4.57 cM/Mb, maternal:

4.49 cM/Mb) than that of vertebrates such as mammals (~0.5 cM/Mb) or birds (~2

cM/Mb)[19, 20]. However, this discrepancy may be partially attributed to the compact

genome size of amphioxus relative to vertebrates. In fact, the average number of CO

events per haploid chromosome per meiosis was approximately 1.2, close of that of

vertebrates[19]. Unlike many vertebrates, no obvious sexual dimorphism was found in

CO rates in paternal and maternal genomes of this family, but the reason behind this is

unclear. Both paternal and maternal genomes have many CO coldspots, likely due to

large-scale heterozygous inversions that can suppress CO events[28]. Like the high

density of SNPs, there are likely many polymorphic heterozygous inversions and

other structural variations in the wild amphioxus populations[29]. Given that

large-scale heterozygous inversions can suppress CO events, there may be large

difference in genomic distribution of COs between different individuals, leading to a

relatively weak correlation between the directly-measured individual CO

recombination map and the historical CO recombination map estimated from

population genomic data. Besides, we found that TE and gene densities are positively

correlated with local CO rates, though the underlying molecular mechanisms leading

to such correlations are unclear. In many vertebrates, the genomic binding profile of

PRDM9 (the protein with zinc-finger structures binding specific DNA sequences and

triggering histone modifications like H3K4me3/H3K36me3) determines CO

hotspots[41]. Due to the lack of PRDM9 ChIP-seq data in amphioxus species, we did

not perform any PRDM9 related analysis, so its potential role in amphioxus COs

remains unknown.

Our analysis of NCO events showed that in amphioxus NCO events occur much

more frequently than CO events and have short impact sequence ranges. Like

vertebrates, amphioxus also exhibits gBGC, which tends to preserve C/G alleles

during conversions[42, 43]. Many characteristics of the NCO events in amphioxus are

similar to those of vertebrates, suggesting that the mechanisms underlying NCO

events may be highly conserved during chordate evolution.

We acknowledge several limitations in our analysis. First, while using bubble

contigs as markers for detecting CO recombination events offers higher specificity,

bubble contigs are derived from highly heterozygous regions of the genome. This can

lead to an uneven genomic distribution of bubble contigs, potentially causing an

underestimation of recombination rates in regions with lower heterozygosity. Second,

our study sequenced a single family, and thus there are possible biases due to the

small sample size. This is especially relevant given that there may be substantial

individual-specific heterozygous inversions that can suppress recombination.

Consequently, we cannot exclude the possibility that such limitations may mask some

differences existing at the population level, such as those linked to sex.

In summary, by sequencing 106 individuals from a *B. floridae* family, we have

comprehensively mapped CO and NCO events in amphioxus male and female

individuals. This study fills important gaps in our understanding of amphioxus

recombination and enhances our knowledge of genetic diversity in amphioxus

populations. Our work can facilitate future research of this important model organism

in biology.

Methods

Sample collection and sequencing

Amphioxus (*B. floridae*) was obtained from a stock maintained by Jr-Kai Yu

originating from Tampa, Florida, and the colony and their offspring were raised under

previously described conditions[44]. 106 individuals from a two-generation family,

including two parents and 104 offspring, were adopted in the analysis. DNA was

extracted and used to construct paired-end 150 bp sequencing libraries via a shotgun

strategy. Whole genome sequencing was performed using the BGISEQ-500 platform.

More details can be found in a companion study on amphioxus *de novo* mutation

rates[11].

Genome assembly and determining inherited parental haplotype contigs in

offspring

We performed *de novo* haplotype-aware assembly of the parental genomes using

Platanus-allee[10] and merged the resultant haplotype contigs into a custom parental

reference genome. In Platanus-allee, heterozygous regions are assembled as 'bubble'

contigs with two allelic sequences, with the longer one named as 'primary' and the

shorter as 'secondary'. Homozygous regions with identical sequences in an individual

are assembled as 'non-bubble' contigs. By aligning offspring sequencing reads to the

custom reference genome using BWA[45], we determined which haplotype contig of a

specific parent at a divergent locus is inherited in each offspring. Detailed methods

can be found in a companion study on amphioxus *de novo* mutation rates[11].

Identification of CO events

As non-bubble contigs derived from homozygous regions are not informative for

recombination analysis, we used only bubble contigs as markers to detect CO events

during meiosis in the father and the mother.

The ideal marker bubble contigs should clearly show haplotype inheritance from

parents to most offspring. For each bubble contig pair, the ratio of offspring inheriting

primary or secondary contigs should be about 1:1. Hence we removed bubble contigs

where inheritance couldn't be determined in over 20% of the offspring or those

significantly deviating from a binomial test (P < 0.05), which might introduce bias

and inaccurately estimate CO events. We then took the homologous regions from the

remaining bubble contig pairs and used nucmer in MUMmer v4.0.0beta2[46] to map

them onto the chromosome-level genome assembly[8] to determine their chromosomal

positions. We excluded the bubble contigs mapped to multiple chromosomes (possible

rearrangement sites) and those unmapped. After filtering, 28,795 bubble contig pairs

remained in the paternal genome and 27,673 in the maternal genome for CO detection.

We grouped these contigs by chromosome, combined inheritance results from all

offspring, and used the R package hapi[12] with maximum parsimony to reconstruct

parental chromosomal haplotypes for each parent. By comparing offspring

chromosomal haplotypes with parental ones, we identified CO events in the gametes

of each offspring.

To identify CO coldspots across the genome, a sliding window approach with a

window size of 5 Mb and a step size of 1 Mb was employed. A region was defined as

a coldspot if no CO events were detected in any offspring within that region.

Successive coldspot regions were subsequently merged to delineate genome-wide

coldspot distributions.

Identification of breakpoints within contigs induced by CO events

To verify and complement our strategy using bubble contigs as markers, we

analyzed putative CO-induced breakpoints within bubble contigs. Based on the

alignment of offspring sequencing reads to the Platanus-allee parental reference

genome, a potential CO breakpoint was identified when an offspring's reads aligned to

both the primary and secondary haplotypes of a bubble contig pair, with the aligned

regions being complementary and forming a new haplotype. This indicates a partial

sequence exchange between the two original haplotypes. To confirm the breakpoint

was from a CO rather than an NCO event, we checked read alignments of all

offspring in that locus. If four haplotypes were present in the offspring population, the

breakpoint was confirmed as a CO breakpoint.

Detection of heterozygous inversions

To identify large-scale heterozygous inversions in parents that may suppress CO

events, we first used the Platanus-allee[10] assembly and bubble contig phasing results

to generate haplotype sequences for each parental chromosome. Then, we aligned

these parental haplotypes to a more complete chromosome-level assembly[8] using

Minimap2 v2.26-r1175[47] with parameters '-a -x asm20 --cs -r2k'. Next, we used

SVIM-asm v1.0.3[48] in haploid mode to detect inversions in each haplotype. By

comparing variations between the two haplotypes, we identified heterozygous

inversions in the parental genomes. Subsequently, the contigs carrying heterozygous

inversions were obtained, and the regions with a mapping length of > 500 bp were

selected for further analysis and visualization.

Correlation analysis between CO events and genomic features

To assess the correlation between CO events and genomic features, we scanned

the genome using a sliding window approach. Given the sparse distribution of CO

events, we used a large window size of 5 Mb with a step size of 1 Mb to reduce

random fluctuations from low CO counts. We excluded CO coldspot regions, which

are potentially affected by parental heterozygous inversions, from the analysis.

Genomic annotations for features like gene density were from Huang et al.[8], and

repeat or transposon regions were annotated with RepeatMasker.

Analysis of NCO events

As non-bubble regions lack markers for NCO detection, we focused on NCO

events in bubble regions. By aligning primary and secondary contigs of each bubble

contig pair, we identified candidate SNPs. Regions with excessive SNPs may suffer

from poor alignment quality, so based on the previously reported heterozygosity

(3.2-4.2%) of the amphioxus genome, we selected bubble regions with 1-5 SNPs per

100 bp and used those SNPs inside as NCO markers. We also excluded homologous

markers shared by the father and the mother to avoid potential alignment errors across

different parental alleles, which could lead to false positive NCOs. Also, as it needs at

least three markers to determine if there is an NCO event, we removed bubble contig

pairs with fewer than three SNPs.

Based on the alignment data of offspring and parental genomes, we used

HaplotypeCaller module of GATK v4.2.0.0[49] with the '-ERC GVCF' parameter to

call variants and determine the genotypes of all offspring at the marker sites. An NCO

event was identified, when an inherited contig from a bubble contig pair had the same

parental phase of markers at both ends but another parental phase in the middle

marker(s), with only three haplotypes observed in offspring. Regarding the impacted

sequenced range of NCO events, given the inability to obtain the precise length, we

used the average of the minimum length (distance between the first and last converted

markers, or 1 bp if only one marker was converted) and the maximum length

(distance between the nearest markers flanking the converted region) as the

approximate length of the NCO events. Although our method cannot distinguish an

NCO event from a *de novo* substitution, given the low mutation rate (4.4×10^{-9} per base

per generation[11]), the level of NCO false positives due to *de novo* substitutions at the

heterozygous sites should be very low. For example, of the detected 9,992 paternal

NCOs, the number of potential *de novo* substitutions in the surveyed genomic space

of 6.71 million (heterozygous markers) ×104 is estimated to be about 1.02 ($4.4 \times 10^{-9} \times$

$6.71 \times 10^{6} \times 104 \times 1/3$, with $1/3$ assuming equal probabilities of three possible

substitutions for a nucleotide), yielding a false positive rate of 0.01%.

Data Availability

The raw sequence data reported in this paper have been deposited in the Genome

Sequence Archive (GSA: CRA027444) that are publicly accessible at

https://ngdc.cncb.ac.cn/gsa. Genome assemblies produced by Platanus-allee have

been deposited in Science Data Bank (ScienceDB) repository

(https://doi.org/10.57760/sciencedb.27648).

Code availability

The custom code used for analysis is deposited at

https://github.com/tale1209/bf_recom.

Author contribution

Lei Tao: Investigation, Methodology, Formal analysis, Writing - Original draft,

Writing - Review & Editing. **Jing Xue**: Investigation, Methodology, Formal analysis,

Writing - Review & Editing. **Junwei Cao**: Data Curation, Investigation. **Guang Li**:

Conceptualization, Data Curation, Resources, Writing - Review & Editing,

Supervision. **Cai Li**: Conceptualization, Resources, Methodology, Writing - Review

& Editing, Supervision.

Conflict of interest

The authors have declared no competing interests.

Acknowledgments

We thank Dr. Luohao Xu for his assistance in obtaining previous genomic data of

amphioxus populations. This work was supported by National Natural Science

Foundation of China (32470690), Natural Science Foundation of Fujian Province of

China (2022J06004), State Key Laboratory of Biocontrol and Guangdong Provincial

Key Laboratory for Aquatic Economic Animals.

Reference

[1] Delsuc F, Brinkmann H, Chourrout D, et al. Tunicates and not cephalochordates are the closest

living relatives of vertebrates[J]. Nature,2006,439(7079):965-968.

[2] Schubert M, Escriva H, Xavier-Neto J, et al. Amphioxus and tunicates as evolutionary model

systems[J]. Trends Ecol Evol,2006,21(5):269-277.

[3] Yuan S, Tao X, Huang S, et al. Comparative immune systems in animals[J]. Annu Rev Anim

Biosci,2014,2:235-258.

[4] Holland L Z. Chordate roots of the vertebrate nervous system: expanding the molecular toolkit[J].
Nat Rev Neurosci,2009,10(10):736-746.

[5] D'Aniello S, Bertrand S, Escriva H. Amphioxus as a model to study the evolution of development
in chordates[J]. Elife,2023,12.

[6] Tao X, Yuan S, Chen F, et al. Functional requirement of terminal inverted repeats for efficient
ProtoRAG activity reveals the early evolution of V(D)J recombination[J]. Natl Sci
Rev,2020,7(2):403-417.

[7] Putnam N H, Butts T, Ferrier D E, et al. The amphioxus genome and the evolution of the chordate
karyotype[J]. Nature,2008,453(7198):1064-1071.

[8] Huang Z, Xu L, Cai C, et al. Three amphioxus reference genomes reveal gene and chromosome
evolution of chordates[J]. Proc Natl Acad Sci U S A,2023,120(10):e2093463176.

[9] Ritz K R, Noor M, Singh N D. Variation in Recombination Rate: Adaptive or Not?[J]. Trends
Genet,2017,33(5):364-374.

[10] Kajitani R, Yoshimura D, Okuno M, et al. Platanus-allee is a de novo haplotype assembler
enabling a comprehensive access to divergent heterozygous regions[J]. Nat Commun,2019,10(1):1702.

[11] Xue J, Tao L, Cao J, et al. Germline de novo mutation rate of the highly heterozygous amphioxus
genome[J]. bioRxiv,2025.

[12] Li R, Qu H, Chen J, et al. Inference of Chromosome-Length Haplotypes Using Genomic Data of
Three or a Few More Single Gametes[J]. Mol Biol Evol,2020,37(12):3684-3698.

[13] Zhang W, Tariq A, Jia X, et al. Plant sperm cell sequencing for genome phasing and determination
of meiotic crossover points[J]. Nat Protoc,2025,20(3):690-708.

[14] Xie H, Li W, Guo Y, et al. Long-read-based single sperm genome sequencing for
chromosome-wide haplotype phasing of both SNPs and SVs[J]. Nucleic Acids
Res,2023,51(15):8020-8034.

[15] Masaki N, Browning S R. Mean gene conversion tract length in humans estimated to be 459 bp
from UK Biobank sequence data[J]. bioRxiv,2025.

[16] Haag C R, Theodosiou L, Zahab R, et al. Low recombination rates in sexual species and sex-asex
transitions[J]. Philos Trans R Soc Lond B Biol Sci,2017,372(1736).

[17] Dumont B L. Variation and Evolution of the Meiotic Requirement for Crossing Over in
Mammals[J]. Genetics,2017,205(1):155-168.

[18] Lenormand T, Engelstadter J, Johnston S E, et al. Evolutionary mysteries in meiosis[J]. Philos
Trans R Soc Lond B Biol Sci,2016,371(1706).

[19] Szasz-Green T, Shores K, Vanga V, et al. Comparative Phylogenetics Reveal Clade-specific
Drivers of Recombination Rate Evolution Across Vertebrates[J]. Mol Biol Evol,2025,42(5).

[20] Stapley J, Feulner P, Johnston S E, et al. Variation in recombination frequency and distribution
across eukaryotes: patterns and processes[J]. Philos Trans R Soc Lond B Biol Sci,2017,372(1736).

[21] Kong A, Thorleifsson G, Gudbjartsson D F, et al. Fine-scale recombination rate differences
between sexes, populations and individuals[J]. Nature,2010,467(7319):1099-1103.

[22] Cooney C R, Mank J E, Wright A E. Constraint and divergence in the evolution of male and
female recombination rates in fishes[J]. Evolution,2021,75(11):2857-2866.

[23] Shi C, Wu X, Su L, et al. A ZZ/ZW Sex Chromosome System in Cephalochordate Amphioxus[J].
Genetics,2020,214(3):617-622.

[24] Vincenten N, Kuhl L M, Lam I, et al. The kinetochore prevents centromere-proximal crossover

recombination during meiosis[J]. Elife,2015,4.

[25] Limborg M T, Mckinney G J, Seeb L W, et al. Recombination patterns reveal information about

centromere location on linkage maps[J]. Mol Ecol Resour,2016,16(3):655-661.

[26] Morgan A P, Gatti D M, Najarian M L, et al. Structural Variation Shapes the Landscape of

Recombination in Mouse[J]. Genetics,2017,206(2):603-619.

[27] Harringmeyer O S, Hoekstra H E. Chromosomal inversion polymorphisms shape the genomic

landscape of deer mice[J]. Nat Ecol Evol,2022,6(12):1965-1979.

[28] Li H, Berent E, Hadjipanteli S, et al. Heterozygous inversion breakpoints suppress meiotic

crossovers by altering recombination repair outcomes[J]. PLoS Genet,2023,19(4):e1010702.

[29] Huang S, Chen Z, Yan X, et al. Decelerated genome evolution in modern vertebrates revealed by

analysis of multiple lancelet genomes[J]. Nat Commun,2014,5:5896.

[30] Penalba J V, Wolf J. From molecules to populations: appreciating and estimating recombination

rate variation[J]. Nat Rev Genet,2020,21(8):476-492.

[31] Kent T V, Uzunovic J, Wright S I. Coevolution between transposable elements and

recombination[J]. Philos Trans R Soc Lond B Biol Sci,2017,372(1736).

[32] Dolgin E S, Charlesworth B. The effects of recombination rate on the distribution and abundance

of transposable elements[J]. Genetics,2008,178(4):2169-2177.

[33] Wang X, Li Z, Feng T, et al. Chromosome-level genome and recombination map of the male

buffalo[J]. Gigascience,2022,12.

[34] Jiang X, Li D, Du H, et al. Genomic features of meiotic crossovers in diploid potato[J]. Hortic

Res,2023,10(6):d79.

[35] Schnable P S, Ware D, Fulton R S, et al. The B73 maize genome: complexity, diversity, and

dynamics[J]. Science,2009,326(5956):1112-1115.

[36] Marand A P, Jansky S H, Zhao H, et al. Meiotic crossovers are associated with open chromatin

and enriched with Stowaway transposons in potato[J]. Genome Biol,2017,18(1):203.

[37] Palsson G, Hardarson M T, Jonsson H, et al. Complete human recombination maps[J].

Nature,2025,639(8055):700-707.

[38] Prentout D, Bykova D, Hoge C, et al. Germline mutation rates and fine-scale recombination

parameters in zebra finch[J]. PLoS Genet,2025,21(4):e1011661.

[39] Li R, Bitoun E, Altemose N, et al. A high-resolution map of non-crossover events reveals impacts

of genetic diversity on mammalian meiotic recombination[J]. Nat Commun,2019,10(1):3900.

[40] Johnston S E. Understanding the Genetic Basis of Variation in Meiotic Recombination: Past,

Present, and Future[J]. Mol Biol Evol,2024,41(7).

[41] Paigen K, Petkov P M. PRDM9 and Its Role in Genetic Recombination[J]. Trends

Genet,2018,34(4):291-300.

[42] Capra J A, Hubisz M J, Kostka D, et al. A model-based analysis of GC-biased gene conversion in

the human and chimpanzee genomes[J]. PLoS Genet,2013,9(8):e1003684.

[43] Glemin S, Arndt P F, Messer P W, et al. Quantification of GC-biased gene conversion in the

human genome[J]. Genome Res,2015,25(8):1215-1228.

[44] Li G, Shu Z, Wang Y. Year-round reproduction and induced spawning of Chinese amphioxus,

Branchiostoma belcheri, in laboratory[J]. PLoS One,2013,8(9):e75461.

[45] Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform[J].

Bioinformatics,2009,25(14):1754-1760.

[46] Marcais G, Delcher A L, Phillippy A M, et al. MUMmer4: A fast and versatile genome alignment

system[J]. PLoS Comput Biol,2018,14(1):e1005944.

[47]   Li   H.   Minimap2:   pairwise   alignment   for   nucleotide   sequences[J].
Bioinformatics,2018,34(18):3094-3100.

[48] Heller D, Vingron M. SVIM-asm: structural variant detection from haploid and diploid genome
assemblies[J]. Bioinformatics,2021,36(22-23):5519-5521.

[49] Mckenna A, Hanna M, Banks E, et al. The Genome Analysis Toolkit: a MapReduce framework
for analyzing next-generation DNA sequencing data[J]. Genome Res,2010,20(9):1297-1303.

695