Title: Early detection of ampicillin susceptibility in *Enterococcus faecium* with MALDI-TOF
MS and machine learning

Thomas Pichl[1], Lucas Miranda[2], Nina Wantia[1], Karsten Borgwardt[2], Janko Sattler[2]

[1]Institute of Medical Microbiology, Immunology and Hygiene, TUM School of Medicine and
Health, Department of Preclinical Medicine, Technical University of Munich, Germany

[2]Department of Machine Learning and Systems Biology, Max Planck Institute of
Biochemistry, Martinsried, Germany

Abstract

Background

Enterococcus faecium can cause severe infections and is often resistant to the first-line

antibiotic ampicillin. Consequently, clinicians usually prescribe broad-spectrum antibiotics,

promoting the selection of multidrug-resistant bacteria. In this study, we investigate the

application of machine learning techniques to detect ampicillin susceptibility directly from

MALDI-TOF mass spectrometry. This technique could enable an earlier optimised treatment

in infections with ampicillin-susceptible *E. faecium*.

Methods

Two datasets of clinical *E. faecium* MALDI-TOF spectra and their resistance phenotype were

analysed: our own Technical University of Munich (TUM) dataset and the publicly available

MS-UMG dataset. We tested logistic regression (LR) and LightGBM models on each dataset

via nested cross-validation and explored transferability on the respective other dataset.

Results

LightGBM demonstrated slightly better performance than LR in identifying susceptible

isolates in the TUM dataset (area under the precision-recall curve (AUPRC) 0.907 ± 0.016 vs

0.902 ± 0.030) as well as in the MS-UMG dataset (AUPRC 0.902 ± 0.029 vs 0.899 ± 0.054).

External validation demonstrated good model transferability (AUPRC of 0.784 ± 0.039 when

trained on MS-UMG; 0.804 ± 0.013 when trained on TUM). SHAP analysis consistently

identified a top-ranked spectral feature corresponding to a peak at an m/z of 5091 in resistant

isolate spectra.

Conclusion

This study demonstrates that LR and LightGBM models can identify ampicillin-susceptible *E.*

faecium isolates from MALDI-TOF spectra and generalise well to unseen datasets. While

clinical implementation currently still requires confirmatory testing, the addition of larger

datasets in the future will support the development of more robust machine learning models.

1   Introduction

Enterococcus faecium is an important opportunistic pathogen that can cause severe infections such as endocarditis, hepatobiliary sepsis, and urinary tract infections. It is a Gram-positive bacterium within the *Enterococcaceae* family and part of the gastrointestinal microbiota [1]. While enterococcal species such as *E. faecalis* rarely exhibit resistance against ampicillin and vancomycin, *E. faecium* is a World Health Organisation-designated priority pathogen with multidrug-resistant strains being a global public health concern [2, 3].

Aminopenicillins such as ampicillin are generally well-tolerated and highly effective against susceptible *Enterococcus* strains. Mutations in the penicillin-binding protein 5 (PBP5) are considered the main cause of ampicillin resistance in *E. faecium,* though additional mechanisms remain less well characterised [4]. Due to the high prevalence of ampicillin resistance in healthcare-associated *E. faecium* isolates, glycopeptide antibiotics like vancomycin are commonly administered. However, the use of these broad-spectrum antibiotics promotes the selection and spread of resistant clones, e.g., *vanA*-mediated vancomycin-resistant *E. faecium*. Early identification of ampicillin-susceptible isolates would facilitate prompt targeted therapy and could therefore reduce unnecessary broad-spectrum antibiotic use and resistance dissemination.

Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) is a commonly used method for bacterial identification [5]. In recent years, several studies investigated its potential for antimicrobial resistance prediction with machine learning (ML) models [6-8]. While most studies focused on *Staphylococcus aureus*, publications on *E. faecium* are scarce [9]. Public databases, such as the *MALDI-TOF Mass Spectra and Resistance Information on Antimicrobials from University Medical Center Göttingen* (MS-UMG), provide MALDI-TOF spectra and corresponding antimicrobial resistance data, facilitating research projects in this emerging field [10].

The objective of this study is to predict ampicillin susceptibility in *E. faecium* isolates using MALDI-TOF MS spectra in combination with ML. By analysing spectra derived from clinical samples at our microbiology laboratory at the Technical University of Munich (TUM), along with spectra provided by the MS-UMG dataset, we aim to develop ML models that can be seamlessly integrated into the routine workflows of diagnostic laboratories that already use MALDI-TOF MS for species identification.

2   Methods

2.1   Acquisition of spectra and susceptibility profiles

The TUM dataset comprises clinical MALDI-TOF spectra of *E. faecium* isolates collected at the TUM microbiology laboratory between August 2021 and May 2025 and the corresponding antimicrobial resistance profile. The majority of clinical samples originate from the TUM University Hospital Rechts der Isar and the TUM University Hospital German Heart Centre.

Spectral acquisition and bacterial identification were performed with the MALDI Biotyper® sirius one System (Bruker Daltonics, Bremen, Germany) using the manufacturer's software solutions (flexcontrol 3.4, MBT Compass 4.1, MBT Compass Reference Library 2023). The direct transfer method was applied to bacterial isolates, followed by treatment with 70% formic acid (Merck, Darmstadt, Germany) and matrix application (α-Cyano-4-hydroxycinnamic acid, Bruker Daltonics, Bremen, Germany).

Antimicrobial resistance testing was conducted with the automated microdilution system VITEK® 2 (bioMérieux, Marcy-l'Étoile, France), disc diffusion tests (Becton Dickinson, Franklin Lakes, USA; MAST Group, Bottle, UK), or strip tests (Liofilchem, Roseto degli Abruzzi, Italy). *E. faecium* isolates were classified as ampicillin-resistant or -susceptible using the EUCAST breakpoints effective at the time of spectral acquisition. Only spectra with a MALDI-TOF score of 2.00 or higher and a single corresponding antibiogram were included.

E. faecium spectra and their corresponding antimicrobial susceptibility profiles were extracted from the MS-UMG dataset. We only included spectra from the diagnostic routine section of the database, matching the characteristics of the TUM dataset.

In both the TUM and MS-UMG datasets, spectra of isolates described as "susceptible, increased exposure (I)" in ampicillin susceptibility, as well as those with implausible antimicrobial susceptibility profiles, duplicate entries, and empty spectra were excluded from further analysis. Vancomycin-resistant *E. faecium* (VRE), which are generally ampicillin-resistant [11], were also excluded to avoid confounding.

2.2   Spectral preprocessing and feature alignment

Spectral preprocessing followed the protocol established by Weis et al. [6] employing the MALDIquant package version 1.22.2 [12] and included variance stabilization via square root transformation, signal smoothing using the Savitzky–Golay algorithm (half-window size: 10), baseline correction with 100 iterations of the SNIP algorithm, total ion current normalization, and trimming to the predefined m/z range of 2,000–20,000.

Spectral alignment was applied to correct for m/z shifts using a reference-based linear

alignment approach adapted from publicly available code [13]. For internal validation, each

dataset was aligned to its own reference peaks, defined as occurring in at least 90% of spectra

within that dataset, with a tolerance of 0.2% m/z. For spectral alignment during external

validation, the *external* dataset was aligned to the reference peaks of the *internal* training

dataset. Aligned spectra were subsequently binned into 3 m/z windows, resulting in a feature

vector of 6,000 dimensions per spectrum.

2.3   Model development and evaluation

The spectral feature vectors served as input for supervised ML models, with ampicillin

susceptibility status as the dependent classification target (susceptible = positive class;

resistant = negative class).

Two classification algorithms were employed in this study: LightGBM, a gradient boosting

algorithm with strong performance in antibiotic resistance prediction from MALDI-TOF

spectra [7], and logistic regression (LR), included for its interpretability and established use

[9].

Models were developed using the scikit-learn (version 1.5.0) and lightgbm (version 4.4.0)

Python libraries with hyperparameter tuning performed via RandomizedSearchCV. The

Matthews Correlation Coefficient (MCC) was selected as the optimisation metric due to its

suitability for imbalanced datasets.

Internal model performance was assessed using a stratified five-fold nested cross-validation

approach (80/20 train/test split in the outer loop; 75/25 split on the inner loop). Performance

metrics were computed and averaged across the outer folds, with macro means and standard

deviations reported at a classification threshold of 0.5. Additionally, the area under the

precision-recall curve (AUPRC) was calculated to provide a threshold-independent metric.

Model transferability was evaluated by training stratified five-fold cross-validation models on

the MS-UMG and TUM datasets, with the respective other serving as the external test set.

Results were averaged over five different random states to obtain reliable performance

estimates.

SHAP (Shapley Additive exPlanations) values were calculated to determine the most relevant

spectral features contributing to model predictions [14]. For further implementation details,

please refer to the code in the supplemental material.

3   Results

3.1   Dataset characteristics

Extraction of MALDI-TOF MS spectra and corresponding susceptibility profiles of *E. faecium* isolates from the TUM dataset yielded 2109 spectra of ampicillin-resistant and 303 of susceptible isolates. The MS-UMG dataset comprises 1136 spectra of resistant isolates and 158 spectra of susceptible ones. Class distribution was similar in both datasets, with 14% and 12% susceptible isolates in the TUM and MS-UMG datasets, respectively.

3.2   5-fold nested cross-validation model performance on the TUM dataset

We first trained and evaluated a LR model using a 5-fold nested cross-validation approach on the TUM dataset. At a classification threshold of 0.5, the LR model showed good performance in determining ampicillin-susceptible isolates, achieving a positive predictive value (PPV) of 0.829 ± 0.048, a sensitivity of 0.878 ± 0.042, and an area under the precision-recall curve (AUPRC) of 0.902 ± 0.03 (**Table 1**). The model also performed strongly in identifying ampicillin-resistant isolates with a negative predictive value (NPV) of 0.982 ± 0.006 and a specificity of 0.973 ± 0.009.

Metric	Logistic regression		LightGBM	
	Macro Mean	SD	Macro Mean	SD
PPV	0.829	0.048	0.887	0.027
Sensitivity	0.878	0.042	0.855	0.067
NPV	0.982	0.006	0.979	0.010
Specificity	0.973	0.009	0.984	0.004
MCC	0.830	0.038	0.852	0.048
AUPRC	0.902	0.030	0.907	0.016

Table 1. Average performance metrics (arithmetic mean) and standard deviations (SD) across folds for logistic Regression and LightGBM models using stratified 5-fold nested cross validation on the TUM dataset. All metrics, except for AUPRC (area under the Precision-Recall Curve), were calculated at a threshold of 0.5. PPV = Positive Predictive Value, NPV = Negative Predictive Value, MCC = Matthews Correlation Coefficient.

We then investigated whether model performance could be improved by employing LightGBM, as previously described [7]. Performance metrics of the LightGBM model showed a better PPV compared to LR with 0.887 ± 0.027, a slightly lower sensitivity of 0.855 ± 0.067, and a slightly higher AUPRC of 0.907 ± 0.016 (**Table 1**). Metrics for the detection of ampicillin-resistant isolates were comparable to LR.

3.3   5-fold nested cross-validation model performance on the MS-UMG dataset

To evaluate performance metrics of logistic regression and LightGBM on another, independent dataset, we performed nested cross-validation on the publicly available MS-UMG dataset.

163　While the LR model achieved a PPV of 0.781 ± 0.044 and a sensitivity of 0.811 ± 0.113, the

164　LightGBM model showed a higher PPV (0.911 ± 0.092) but a lower sensitivity (0.671 ±

165　0.150) at the same classification threshold of 0.5 (**Table 2**). However, the threshold-

166　independent AUPRC was 0.899 ± 0.054 for LR and slightly higher with 0.902 ± 0.029 for

167　LightGBM, indicating strong performance in identifying ampicillin-susceptible isolates.

| Metric | *Logistic regression* | | *LightGBM* | |
	Macro Mean	SD	Macro Mean	SD
PPV	0.781	0.044	0.911	0.092
Sensitivity	0.811	0.113	0.671	0.150
NPV	0.974	0.015	0.956	0.019
Specificity	0.968	0.008	0.989	0.012
MCC	0.766	0.077	0.750	0.073
AUPRC	0.899	0.054	0.902	0.029

Table 2. Average performance metrics (arithmetic mean) and standard deviations (SD) across folds for logistic
regression and LightGBM models using 5-fold nested cross validation on the MS-UMG dataset. All metrics,
except for AUPRC (area under the Precision-Recall Curve), were calculated at a threshold of 0.5. PPV = Positive
Predictive Value, NPV = Negative Predictive Value, MCC = Matthews Correlation Coefficient.

3.4　External model validation with LightGBM

We assessed model transferability by training LightGBM 5-fold cross-validation models on

the TUM and MS-UMG datasets, with the respective other dataset serving as the test set

across five different random states.

Both models demonstrated good overall transferability with an AUPRC of 0.784 ± 0.039 for

the model trained on the TUM dataset and tested on the MS-UMG dataset, and an AUPRC of

0.804 ± 0.013 for the model trained on the MS-UMG dataset and tested on the TUM dataset

(**Table 3**). PPV was high for the TUM-trained model (0.895 ± 0.089) at a classification

threshold of 0.5, but sensitivity was low (0.410 ± 0.084). The opposite trend was observed for

the MS-UMG trained model at the same threshold, which showed a sensitivity of 0.930 ±

0.012 and a PPV of 0.397 ± 0.017.

| Metric | *training TUM; testing MS-UMG* | | *training MS-UMG; testing TUM* | |
	Macro Mean	SD	Macro Mean	SD
PPV	0.895	0.089	0.397	0.017
Sensitivity	0.410	0.084	0.930	0.012
NPV	0.924	0.009	0.988	0.002
Specificity	0.992	0.009	0.797	0.013
MCC	0.569	0.034	0.529	0.019
AUPRC	0.784	0.039	0.804	0.013

183　**Table 3**. Average performance metrics (arithmetic mean) and standard deviations (SD) across five random states
184　for LightGBM models, using stratified 5-fold nested cross validation (80/20 train/test split) on the TUM dataset
185　(training) and MS-UMG dataset (testing), and vice versa. All metrics, except for AUPRC (area under the

186 Precision-Recall Curve), were calculated at a threshold of 0.5. PPV = Positive Predictive Value, NPV = Negative
187 Predictive Value, MCC = Matthews Correlation Coefficient.

3.5 Model interpretability via SHAP value analysis

In both LR and LightGBM internal validation models, SHAP value analysis of the ten most

impactful spectral features (i.e., bins) from the outer folds' test sets of the TUM dataset

consistently identified a top-ranked feature in the m/z range of 5093-5096 (**Figure 1**).

Interestingly, the same feature - or directly adjacent features – was also top-ranked in all

internal LightGBM and LR models applied to the MS-UMG dataset. This trend persisted in

the external validation models, where the feature was top-ranked across random states when

trained on the TUM dataset and tested on the MS-UMG dataset and in four out of five random

states when trained on MS-UMG and tested on TUM. All SHAP plots for internal and

external model validation are provided in the supplementary material.

Figure 1. Top SHAP Features Predicting Ampicillin Susceptibility Across Folds. Mass-to-charge (m/z)
ranges (x-axis) derived from the top ten most influential SHAP (SHapley Additive exPlanations) features of each
fold in determining ampicillin susceptibility in *E. faecium* isolates from the TUM dataset, using stratified 5-fold
nested cross-validation. Only features that appeared in the top ten SHAP ranking of all five folds are shown.
Numbers within the boxes indicate the SHAP rank of the feature within the respective fold. Each fold is
represented by a distinct colour and a corresponding number on the y-axis. Results are shown separately for
logistic regression (left) and LightGBM (right).

Corresponding to the top-ranked feature at the m/z bin 5093-5096, we observed a distinct

peak at an m/z of 5093.9 in the averaged spectra of unbinned, warped ampicillin-resistant *E.*

faecium isolates from the TUM dataset. The peak was absent in the averaged spectra of

susceptible isolates. By extrapolating from two conserved *E. faecium* ribosomal proteins in

close proximity at m/z 4424 and 5353, we estimated the corresponding protein mass of the

m/z 5093.9 peak at 5091 Da.

4   Discussion

Machine learning as a means to directly predict antimicrobial susceptibility from MALDI-

TOF spectra has attracted growing interest in recent years [9]. In this context, *E. faecium*

received comparatively little attention, and studies focused mostly on vancomycin resistance

prediction [15, 16].

E. faecium is phylogenetically divided into clades which correlate with ampicillin

susceptibility: Clade A1 (mostly resistant) is primarily found in healthcare settings, A2

(mostly susceptible) is linked to livestock, and B (also mostly susceptible) is frequently

encountered in community environments [1, 17]. Faury et al. successfully differentiated *E.*

faecium clades A1/A2 from clade B by constructing a MALDI-TOF spectra library but were

unable to distinguish the 52 included isolates based on ampicillin susceptibility alone, with

only 50% identified correctly [18]. In the present study, we set out to investigate ampicillin

susceptibility prediction in *E. faecium* isolates from MALDI-TOF spectra derived from two

clinical datasets - the TUM dataset and the MS-UMG dataset.

Clinical outcome data comparing ampicillin to glycopeptide antibiotics for ampicillin-

susceptible enterococcal infections are scarce, with some retrospective studies reporting a

lower mortality with ampicillin-based therapy [19, 20]. International guidelines recommend

ampicillin, either as monotherapy or in combination, as the first-line treatment in severe

infections caused by ampicillin-susceptible *E. faecium* [21]. Empiric treatment with

glycopeptide antibiotics may be de-escalated once culture-based antibiotic susceptibility

testing confirms ampicillin susceptibility.

Our model can support clinical decision-making in two potential scenarios. First,

classification thresholds could be adjusted to maximise sensitivity, thereby increasing the

chance of detecting ampicillin-susceptible isolates. This approach would allow for the early

addition of ampicillin to empiric broad-spectrum antibiotic therapy, at the cost of accepting a

higher number of false positives (lower PPV). This is analogous to the South Australia Health

guidelines for empiric treatment of *Staphylococcus aureus* bloodstream infections, where

cefazolin (optimal for oxacillin-susceptible *S. aureus)* and vancomycin (necessary for

resistant strains) are both administered initially until susceptibility testing results become

available [22]. Conversely, a classification threshold that prioritises the PPV may allow for

earlier step-down from broad-spectrum antibiotics to ampicillin, reducing the unnecessary use

of glycopeptide antibiotics. This approach accepts a lower sensitivity, whereby some

susceptible isolates will be missed. However, with the current model performance as

determined in this study, ampicillin susceptibility would still require confirmation via an

additional, highly specific test. The precision-recall curves (supplementary **Figures S1 and**

S7) highlight the trade-off between sensitivity and PPV in the context of predicting

ampicillin-susceptibility.

SHAP value analysis revealed an adjusted m/z peak at 5091 as highly relevant for

differentiating between the ampicillin-susceptible and -resistant classes, consistently across

both datasets and different ML algorithms. Brackmann et al. identified oxidised bacteriocin

T8 – also known as bacteriocin 43 or hiracin JM79 (Uniprot Q0Z8B6) – as the source of an

m/z peak at 5092 in MALDI-TOF spectra of vancomycin-resistant *E. faecium* isolates using

MALDI TOF/TOF (MS/MS) analysis [23]. Bacteriocin T8 presence has been shown to confer

a competitive advantage relevant to the emergence of dominant *E. faecium* lineages [24].

Notably, a recent study found bacteriocin T8 to occur frequently in healthcare-associated *E.*

faecium isolates, but rarely in community-acquired strains [25]. The spectral acquisition mode

used by Brackmann et al. differed from the one used in this study, likely explaining the 1Da

difference between the observed peaks. Whole-genome sequencing data were not available for

the TUM or MS-UMG datasets; however, Faury et al. published genome sequences for 26

ampicillin-resistant and 26 -susceptible *E. faecium* isolates (Bioproject PRJEB56579) [18].

We detected bacteriocin T8 in 1 of 26 translated protein sequences from susceptible isolates

and in 17 of 26 from resistant ones, based on alignments with ≥90% sequence identity and

≥90% query coverage. In summary, the peak observed in ampicillin-resistant *E. faecium*

isolates in this study likely represents bacteriocin T8, which, while not directly mediating

ampicillin resistance, may serve as a useful surrogate marker.

This study has several strengths. We used two independent clinical datasets from major

German university hospitals, with spectra generated during routine diagnostic procedures. We

assessed the performance of two machine learning algorithms -LightGBM and logistic

regression- following established best practices to reduce overfitting and evaluate model

interpretability. Finally, we demonstrated model transferability across datasets, evaluating

broader diagnostic implementation.

However, there are important limitations. The retrospective design of this study limits its
immediate applicability to prospective diagnostic workflows. The datasets used lacked
genomic information and clade classification, preventing association between genes and the
spectral peaks as well as clade-stratified analysis. Both datasets originate from German
hospitals, and it remains uncertain whether similar results would be observed in other
geographic settings with a different strain epidemiology.

Overall, our findings indicate good performance in the detection of ampicillin-susceptible *E.*
faecium isolates using MALDI-TOF MS in combination with ML. This study contributes to
the rapidly evolving field of machine learning for antibiotic susceptibility testing. In the
future, the availability of additional and better characterised datasets will allow the
development of more robust and generalisable models.

Data availability

The Python and R scripts developed in this study are available on GitHub at
https://github.com/MicrobeTom/E_faecium_ampicillin. MALDI-TOF spectra of *Enterococcus*
faecium isolates used in this study are available from the Zenodo repository
https://doi.org/10.5281/zenodo.15769315 distributed under a Creative Commons Zero v1.0
Universal licence.

Declaration of Generative AI and AI-assisted technologies

ChatGPT (OpenAI) was used to assist with code generation and proofreading of the
manuscript. All AI-assisted contributions were reviewed and verified by the authors.

Ethical Approval

This study was reviewed by the TUM Ethics Committee (reference number 2024-543-W-SB),
which determined that formal ethical approval was not required in accordance with Paragraph
15 of the Professional Code for Physicians in Bavaria ("Berufsordnung für Ärzte in Bayern").

Funding: None

Competing interests: None declared

References

1   Wei Y, Palacios Araya D, Palmer KL. Enterococcus faecium: evolution,
adaptation, pathogenesis and emerging therapeutics. Nat Rev Microbiol
2024;22(11):705-721. https://doi.org/10.1038/s41579-024-01058-6.

2   Huycke MM, Sahm DF, Gilmore MS. Multiple-drug resistant enterococci: the
nature of the problem and an agenda for the future. Emerg Infect Dis
1998;4(2):239-249. https://doi.org/10.3201/eid0402.980211.
3   WHO. WHO Bacterial Priority Pathogens List: bacterial pathogens of public
health importance to guide research, development and strategies to prevent
and control antimicrobial resistance. 2024.
4   Zhang X, Paganelli FL, Bierschenk D et al,. Genome-wide identification of
ampicillin resistance determinants in Enterococcus faecium. PLoS genetics
2012;8(6). https://doi.org/10.1371/JOURNAL.PGEN.1002804.
5   van Belkum A, Welker M, Pincus D, Charrier JP, Girard V. Matrix-Assisted
Laser Desorption Ionization Time-of-Flight Mass Spectrometry in Clinical
Microbiology: What Are the Current Issues? Ann Lab Med 2017;37(6):475-483.
https://doi.org/10.3343/alm.2017.37.6.475.
6   Weis C, Cuenod A, Rieck B et al,. Direct antimicrobial resistance prediction
from clinical MALDI-TOF mass spectra using machine learning. Nat Med
2022;28(1):164-174. https://doi.org/10.1038/s41591-021-01619-9.
7   Wang WY, Chiu CF, Tsao SM, Lee YL, Chen YH. Accurate prediction of
antimicrobial resistance and genetic marker of Staphylococcus aureus clinical
isolates using MALDI-TOF MS and machine learning - across DRIAMS and
Taiwan database. Int J Antimicrob Agents 2024;64(5):107329.
https://doi.org/10.1016/j.ijantimicag.2024.107329.
8   Nguyen HA, Peleg AY, Song J et al,. Predicting Pseudomonas aeruginosa
drug resistance using artificial intelligence and clinical MALDI-TOF mass
spectra. mSystems 2024;9(9):e0078924.
https://doi.org/10.1128/msystems.00789-24.
9   López-Cortés XA, Manríquez-Troncoso JM, Kandalaft-Letelier J, Cuadros-
Orellana S. Machine learning and matrix-assisted laser desorption/ionization
time-of-flight mass spectra for antimicrobial resistance prediction: A systematic
review of recent advancements and future development. J Chromatogr A
2024;1734:465262. https://doi.org/10.1016/j.chroma.2024.465262.
10   Park Y, Weig M, Noll C, Hauschild A-C, Bader O. MS-UMG: MALDI-TOF Mass
Spectra and Resistance Information on Antimicrobials from University Medical
Center Göttingen. 2024. https://doi.org/10.5281/zenodo.13911744.
11   Shah KJ, Cherabuddi K, Shultz J, Borgert S, Ramphal R, Klinker KP. Ampicillin
for the treatment of complicated urinary tract infections caused by
vancomycin-resistant Enterococcus spp (VRE): a single-center university
hospital experience. Int J Antimicrob Agents 2018;51(1):57-61.
https://doi.org/10.1016/j.ijantimicag.2017.06.008.
12   Gibb S, Strimmer K. MALDIquant: a versatile R package for the analysis of
mass spectrometry data. Bioinformatics 2012;28(17):2270-2271.
https://doi.org/10.1093/BIOINFORMATICS/BTS447.
13   BorgwardtLab. maldi_amr: Code for "Direct Antimicrobial Resistance
Prediction from MALDI-TOF mass spectra through Machine Learning".
GitHub, 2020.
14   Lundberg S, Lee S-I. A Unified Approach to Interpreting Model Predictions.
2017. https://doi.org/10.48550/arXiv.1705.07874.
15   Wang HY, Hsieh TT, Chung CR et al,. Efficiently Predicting Vancomycin
Resistance of Enterococcus Faecium From MALDI-TOF MS Spectra Using a
Deep Learning-Based Approach. Front Microbiol 2022;13:821233.
https://doi.org/10.3389/fmicb.2022.821233.

16 Candela A, Arroyo MJ, Sánchez-Molleda Á et al,. Rapid and Reproducible
MALDI-TOF-Based Method for the Detection of Vancomycin-Resistant
Enterococcus faecium Using Classifying Algorithms. Diagnostics (Basel)
2022;12(2). https://doi.org/10.3390/diagnostics12020328.
17 Gao W, Howden BP, Stinear TP. Evolution of virulence in Enterococcus
faecium, a hospital-adapted opportunistic pathogen. Curr Opin Microbiol
2018;41:76-82. https://doi.org/10.1016/j.mib.2017.11.030.
18 Faury H, Le Guen R, Demontant V et al,. Ampicillin-susceptible Enterococcus
faecium infections: clinical features, causal clades, and contribution of MALDI-
TOF to early detection. Microbiol Spectr 2023;11(5):e0454522.
https://doi.org/10.1128/spectrum.04545-22.
19 Fletcher JM, Kram SJ, Sarubbi CB, Anderson DJ, Kram BL. Effectiveness of
Vancomycin or Beta-Lactam Therapy in Ampicillin-Susceptible Enterococcus
spp. Bloodstream Infections. J Pharm Pract 2019;32(4):375-381.
https://doi.org/10.1177/0897190017751208.
20 Seong YJ, Song JE, Lee E et al,. Clinical outcome of ampicillin or
ampicillin/sulbactam versus glycopeptides in ampicillin-susceptible
Enterococcus faecalis/faecium bacteremia: a 10-year retrospective cohort
study. BMC Infect Dis 2024;24(1):906. https://doi.org/10.1186/s12879-024-
09824-w.
21 Delgado V, Ajmone Marsan N, de Waha S et al,. 2023 ESC Guidelines for the
management of endocarditis. Eur Heart J 2023;44(39):3948-4042.
https://doi.org/10.1093/eurheartj/ehad193.
22 Staphylococcus aureus Bacteraemia (SAB) Management Clinical Guideline
(Adult). In: Health S, ed. Government of South Australia, 2023.
23 Brackmann M, Leib SL, Tonolla M, Schurch N, Wittwer M. Antimicrobial
resistance classification using MALDI-TOF-MS is not that easy: lessons from
vancomycin-resistant Enterococcus faecium. Clin Microbiol Infect
2020;26(3):391-393. https://doi.org/10.1016/j.cmi.2019.10.027.
24 Mills EG, Hewlett K, Smith AB et al,. Bacteriocin production facilitates
nosocomial emergence of vancomycin-resistant Enterococcus faecium. Nature
Microbiology 2025 10:4 2025-03-21;10(4). https://doi.org/10.1038/s41564-025-
01958-0.
25 Tedim AP, Almeida-Santos AC, Lanza VF et al,. Bacteriocin distribution
patterns in Enterococcus faecium and Enterococcus lactis: bioinformatic
analysis using a tailored genomics framework. Appl Environ Microbiol
2024;90(10):e0137624. https://doi.org/10.1128/aem.01376-24.