Reliability and signal comparison of OPM-MEG, fMRI & iEEG in a repeated movie viewing paradigm

Olivia R. Christiano & Sebastian Michelmann

Department of Psychology, New York University, New York, NY, United States
Corresponding Authors: Sebastian Michelmann (s.michelmann@nyu.edu) & Olivia R. Christiano (olivia.christiano@nyu.edu)

ABSTRACT

Optically pumped magnetometers (OPMs) offer a promising advancement in noninvasive neuroimaging via magnetoencephalography (MEG), but establishing their reliability and comparability to existing methods remains an ongoing endeavor. Here, we evaluated OPM recordings by assessing their test-retest reliability and comparing them to functional magnetic resonance imaging (fMRI) and intracranial electroencephalography (iEEG) recordings in a repeated movie viewing paradigm. In 7 frequency bands (δ: 0.5–4 Hz, θ: 4–8 Hz, α: 8–12 Hz, β: 12–28 Hz, γ_1: 28–46 Hz, γ_2: 55–70 Hz, and HF: 64–116 Hz) we quantified the signal consistency (1) within individuals, (2) across subjects, and (3) across modalities. OPM exhibited widespread reliability, particularly in lower frequency bands; spatial patterns resembled those of fMRI and iEEG in visual and auditory regions. Cross-modal analyses revealed robust OPM-fMRI correspondence, including inverse correlations at low frequencies and positive correlations at higher frequencies, consistent with known relationships between oscillatory power and BOLD responses. Comparisons of signal-to-noise (SNR) estimates further revealed that in some regions, the SNR of cross-modal alignment exceeded within-modality reliability, suggesting that bridging between modalities can sometimes enhance SNR by attenuating reliably shared noise. Our findings demonstrate that OPM consistently captures stimulus-locked neural dynamics that converge with established modalities.

Keywords: OPM, MEG, fMRI, iEEG, reliability, intersubject correlation, cross-modal correlation

1. INTRODUCTION

Advancing our understanding of the neural activity underlying human cognition depends on tools that reliably capture neurophysiological signals with high temporal precision. Optically pumped magnetometers (OPMs) represent a promising innovation in magnetoencephalography (MEG), a noninvasive technique with high temporal resolution (milliseconds) that can (under favorable conditions) measure cortical sources with up to 2–3 mm precision (Brookes et al., 2021; D. Cohen, 1968, 1972; Hämäläinen et al., 1993; Proudfoot et al., 2014). OPMs address many of the practical limitations associated with superconducting quantum interference devices (SQUIDs) used in conventional MEG systems, whose cryogenic requirements impose substantial operational costs and technical constraints (Andersen et al., 2017; Boto et al., 2017; D. Cohen, 1972; Hämäläinen et al., 1993; Hill et al., 2020): Unlike SQUIDs, OPMs operate at room temperature, eliminating the need for cryogenic cooling and the associated costs (Boto et al., 2017). Their lightweight, compact design allows for direct placement on the scalp, which minimizes the sensor-to-brain distance and may consequently enhance signal amplitude and spatial resolution (Hill et al., 2020; Iivanainen et al., 2017; Nugent et al., 2022). Additionally, OPMs can be configured into wearable, adjustable helmets, accommodating naturalistic movement during data acquisition, and enabling greater adaptability to different head sizes and shapes (Boto et al., 2018). These features not only broaden MEG's accessibility and versatility in research but also have the potential to improve signal quality by ensuring consistent and close sensor-to-scalp proximity across individuals (Boto et al., 2016, 2017, 2018; Brookes et al., 2021; Hill et al., 2020; Nugent et al., 2022; Rea et al., 2022).

OPM–MEG (hereafter referred to as OPM) is consequently positioned to be a major advancement in neuroimaging, yet its utility hinges on validating whether it captures reliable and meaningful neural signals. While it has been shown that OPMs can detect canonical MEG signals such as evoked responses, oscillatory activity, functional localization, and large-scale connectivity (An et al., 2022; Boto et al., 2021; Rier et al., 2023; Safar et al., 2024; Sun et al., 2024; Tierney et al., 2018; Wang et al., 2024), their reliability has not been as thoroughly established as the reliability of SQUID based MEG (e.g., Lankinen et al., 2014; Lew et al., 2021; Liu et al., 2022; Martín-Buro et al., 2016).

Moreover, while OPMs have demonstrated strong concordance with SQUIDs across sensor- and source-level analyses, sometimes achieving comparable or greater signal quality with fewer sensors (Iivanainen et al., 2023; Marhl et al., 2022; Rhodes et al., 2023), comparative evidence with other neuroimaging and electrophysiological modalities remains limited. Because different measurement techniques are sensitive to distinct neural

Figure 1: Schematic of correlation analyses. Within-subject correlations (blue) were computed between the first and second viewing for each subject and averaged across the respective cohort; for iEEG, electrodes were treated as a single subject. Between-subject correlations (green) were computed across viewings, averaging correlations from viewing 1 to viewing 2 (solid) and vice versa (dashed), comparing each subject to the rest of their cohort. Between-method correlations (red) were computed similarly, comparing each subject *i* in one modality to each subject *j* in another, or to the single-subject iEEG dataset; fMRI to iEEG comparisons were conducted but are not depicted here for simplicity. Within-viewing analyses were also conducted and are reported in the Supplementary Materials. Notes: Movie was presented in color. Diagrams created in https://BioRender.com/epg2wkb. Image used in this schematic: Warner Bros. (1974). [Al Pacino and Penelope Allen in Dog Day Afternoon] [Photograph]. Wikimedia Commons.

processes and noise sources, direct comparisons are needed to clarify both shared and modality-specific components. For instance, EEG and iEEG primarily detect radially oriented electrical currents in cortical gyri, while MEG is most sensitive to tangentially oriented sources located in the sulci. Furthermore, MEG is particularly sensitive in regions closer to the sensors because magnetic field strength diminishes rapidly as it travels away from the source (Hämäläinen et al., 1993). In contrast, fMRI measures vascular responses associated with neural activity, capturing slow hemodynamic changes rather than direct electrophysiological signals. As a novel technique, it is important to understand which neurophysiological signals OPM captures, and critically, to determine whether it does so reliably across individuals and in alignment with signals measured by other established neuroimaging methods.

Here, we systematically compare neural activity recorded with OPM, fMRI, and iEEG during repeated viewings of a continuous audio-visual movie stimulus. We *(i)* quantify OPM signal reliability within individuals, *(ii)* quantify shared OPM signal between individuals, *(iii)* determine the extent to which OPM captures the same stimulus-locked neural activity as fMRI and iEEG, and *(iv)* characterize how its signal reliability varies across these comparisons. To answer these questions, we compute test–retest reliability within individual participants, between different subjects

(Intersubject correlations, ISC; Hasson et al., 2004), and between recording modalities. (See Figure 1 for an overview of our analysis framework.) Finally, we compare signal-to-noise ratios (SNR) within OPM and between OPM and fMRI to characterize their variation across levels of comparison.

2. METHODS

Neural activity was recorded from three independent subject groups as they watched the same segment of the feature film Dog Day Afternoon (DDA; Lumet, 1975) twice. The datasets used in this study comprised recordings from ten OPM subjects (Rier et al., 2023), eleven fMRI subjects (Haufe et al., 2018), and five iEEG subjects (Honey et al., 2012). Comprehensive details on study subjects, data acquisition, experimental settings, and preprocessing steps can be found in the corresponding articles; however, a succinct description of each study's methodology is provided below. Unless otherwise specified, all data were processed using custom MATLAB scripts and the FieldTrip toolbox (Oostenveld et al., 2010).

2.1 OPM (Rier et al., 2023)

2.1.1 Subjects and experimental settings

OPM data were originally collected at the University of Nottingham as part of a separate study (Rier et al., 2023). The OPM cohort consisted of ten subjects (4 female, 6 male, all right-

handed; age: M = 31 years, SD = 8) who provided written informed consent to participate in the study. The study protocol was approved by the University of Nottingham Medical School Research Ethics Committee.

Subjects were presented with a 600 s clip of DDA twice, with approximately 1–2 minutes between each viewing. The movie clips were presented at a visual angle of ~13° horizontally and ~9° vertically, approximately 80 cm from the subjects' heads, and audio was delivered via a waveguide connected to speakers outside of the scanning room (Rier et al., 2023).

2.1.2 Signal acquisition

OPM data were recorded using triaxial OPM sensors (QuSpin, Inc., Colorado, USA) mounted in a 3D-printed helmet (Cerca Magnetics Ltd., Nottingham, UK), selected to best fit each participant. Each sensor has a noise floor of ~13 fT/sqrt (Hz), a bandwidth of ~150 Hz, and provides three independent channels. Sensors were distributed evenly across the scalp, enabling approximately whole-head coverage, which resulted in an average of ~168 channels per subject (range: 165–174). Recordings took place in an OPM-optimized magnetically shielded room (Cerca Magnetics Ltd., Nottingham, UK) equipped with background magnetic field control via reference OPMs, biplanar electromagnetic coils, and a motion tracking camera system. Once magnetic field stabilization procedures were complete, data were recorded as participants viewed the movie segment (600 s per run, sampled at 1200 Hz). Co-registration of sensors to anatomy was performed by aligning 3D digitizations of participants' heads, collected with and without the helmet on, to MRI-derived surfaces (Rier et al., 2023).

2.1.3 Preprocessing

OPM data were minimally preprocessed. Channels were excluded based on the initial preprocessing performed by Rier et al., 2023, which removed sensors with near-zero variance, resulting in an average of 164 usable channels per subject (range: 161–166). To minimize edge artifacts, MEG data were padded with 1 s before and after the 600 s movie segment, then bandpass filtered (0.5–150 Hz), notch filtered at 50, 100, and 120 Hz, and down-sampled to 256 Hz.

2.1.4 Source reconstruction

Source reconstruction was performed using linear constrained minimum variance (LCMV) beamforming (Veen et al., 1997). Individual T1-weighted MRIs were segmented and used to construct single-shell head models (Nolte, 2003), co-registered with subject-specific sensor locations. An 8 mm-spaced template grid in MNI space was nonlinearly warped to each participant's anatomy to obtain individualized source positions for corresponding MNI coordinates. Sources were estimated using the data covariance computed separately for each movie segment across the 600 s time window. The dipole orientation was fixed to the direction of maximum amplitude. Separate filters were computed for each movie repetition in each of the 10 participants.

For comparison to iEEG, source reconstruction of the OPM data was repeated using a shared grid defined by the MNI coordinates of 656 iEEG electrodes. These coordinates were nonlinearly warped to each participant's anatomical space, and source activity was reconstructed directly at these locations using the same preprocessed MEG data, head models, and beamforming procedure described above.

2.1.5 Amplitude of the analytic signal

To extract frequency-specific activity, the source-reconstructed OPM data were bandpass filtered according to the following frequency bands: δ (0.5–4 Hz), θ (4–8 Hz), α (8–12 Hz), β (12–28 Hz), two ranges within the γ band (γ_1: 28–46 Hz, γ_2: 55–70 Hz), and a high frequency band (HF: 64–116 Hz) using a 4th order Butterworth IIR filter. We then estimated the log-transformed amplitude at each source location by taking the logarithm of the absolute value of the analytic signal obtained via the Hilbert transform. This computation was performed separately for each subject, movie viewing, and frequency band.

2.1.6 Cross-modal temporal alignment

OPM data were trimmed to match the adjusted 297 s time axis of the fMRI data (see Section 2.2) by removing the first 16.74 s and the final 286.26 s of the original 600 s recording. Time courses of log-transformed amplitude were resampled to 20 Hz to enable subsequent cross-modal analyses and ensure temporal alignment across modalities.

2.2 fMRI (Haufe et al., 2018)

2.2.1 Subjects and experimental settings

fMRI data were originally collected at Princeton University for the purpose of another study (Haufe et al., 2018). The fMRI sample consisted of eleven subjects (6 female, 5 male; age range: 20-35 years) who provided written informed consent to participate in the study, approved by the Princeton University Institutional Review Board. All subjects were fluent in English, with normal or corrected-to-normal vision, and no history of psychiatric or neurological disorders.

Subjects viewed a 325 s clip from DDA twice, with approximately 11 minutes between repetitions, as part of a larger study. Movie clips were displayed (subtended 20° horizontally and 16° vertically) using the Psychophysics Toolbox in MATLAB, synchronized with MRI data acquisition, and presented with audio delivered via in-ear headphones (Haufe et al., 2018).

2.2.2 Signal acquisition

MRI data were collected using a 3T Skyra scanner (Siemens, Munich, Germany) with a 16-channel head coil. Functional images were acquired with a gradient-echo EPI sequence (3 × 3 × 4 mm resolution, 27 interleaved slices, TR = 1.5 s, TE = 30 ms, flip angle = 72°, FOV = 192 × 192 mm², GRAPPA iPAT = 2); high-resolution structural scans were obtained in each session using an MPRAGE sequence (0.9375 mm isotropic, TR = 1.9 s, TE = 2.1 ms, flip angle = 9°, FOV = 240 × 240 mm²) (Haufe et al., 2018).

2.2.3 Spatial registration

Each subject's data were aligned to MNI space using AFNI's linear (3dAllineate) and nonlinear (3dQWarp) registration tools (Cox, 1996), resampled to a 4 mm grid using linear interpolation, and spatially smoothed using a 10 mm full-width-at-half-maximum Gaussian kernel, resulting in the retained volumetric data (32,798 voxels; Haufe et al., 2018).

2.2.4 Preprocessing

Functional data were preprocessed by the original authors as follows: slice-time- and motion-corrected (AFNI's 3dvolreg) data were linearly detrended, and the discrete Fourier transform (and its inverse) were used to apply a high-pass filter at 0.01 Hz. To ensure signal stability and exclude the blank screen period at the conclusion of the recording, the first 15 s and final 13 s were removed, yielding 297 s of data (Haufe et al., 2018).

2.2.5 Cross-modal alignment

To align the fMRI data to the OPM source grid, we interpolated fMRI activity from MNI-space coordinates onto the template source model used in the OPM beamforming analysis. For each grid point inside the brain volume, we identified the nearest fMRI coordinate(s) based on the minimum Euclidean distance and averaged the corresponding time series. This yielded spatially aligned fMRI data on the same cortical grid used for OPM source projection.

For analyses comparing fMRI and iEEG, we aligned the fMRI data to the MNI-space coordinates of the 656 iEEG electrodes.

For each electrode, we identified all fMRI voxels within a 6 mm radius based on Euclidean distance and averaged their time series. This yielded a version of the fMRI dataset spatially down-sampled to the iEEG grid, allowing direct comparison across modalities at matched anatomical locations. Voxels outside this radius were excluded from the average, and voxel counts per electrode were tracked to normalize the resulting time series.

To ensure temporal alignment across imaging modalities, the fMRI time axis was adjusted to account for hemodynamic delay. The signal was shifted by the lag that maximized the cross-correlation between the amplitude envelope of the movie audio and the time series of the 20 locations nearest to an anatomical mask of auditory cortex regions defined by the AAL atlas (Tzourio-Mazoyer et al., 2002). Preprocessed data were then linearly interpolated to a resolution of 20 Hz.

2.3 iEEG (Honey et al., 2012)

2.3.1 Subjects and experimental settings

iEEG data were collected at New York University for the purpose of a separate study (Honey et al., 2012). The iEEG cohort consisted of five subjects (4 female, 1 male; age range: 20–47 years) from the Comprehensive Epilepsy Center of the New York University School of Medicine experiencing pharmacologically refractory complex partial seizures. All patients provided informed consent both before and after electrode implantation in accordance with National Institutes of Health guidelines administered by the local Institutional Review Board.

Patients viewed the same intact 325 s clip of DDA twice, each followed by a 30 s fixation period, with approximately 12 minutes between intact repetitions. Stimuli presented using PsychToolbox Extensions for MATLAB were displayed approximately 40–60 cm from the subjects' eyes, on a bedside laptop.

2.3.2 Signal acquisition

Recordings were obtained from 922 subdural electrodes in implanted arrays (8 × 8, 4 × 8, or 1 × 8 configurations) spaced 10 mm apart and 2.3 mm in exposed diameter; signals were referenced to skull screws and acquired at 30 kHz using a custom system built on the NSpike framework (L.M. Frank and J. MacArthur, Harvard University Instrument Design Laboratory, Cambridge, MA), which incorporates a 0.6 Hz hardware high-pass filter (Honey et al., 2012).

2.3.3 Spatial registration

Electrode positions were determined using a combination of T1 images (collected pre- and post-implantation) labeling, intra-operative photos, and a MATLAB tool incorporating known electrode geometry (Yang et al., 2012)); SPM's DARTEL algorithm (Ashburner, 2007) was then used to warp each subject's anatomy and individual electrode coordinates to MNI space (Honey et al., 2012). Note that, unlike Haufe et al. (2018) and Honey et al. (2012), we did not exclude electrodes based on hemisphere location; thus, all 656 artifact-free, labeled, and MNI-registered electrodes were retained.

Figure 2: Location of iEEG electrodes on cortical surface. iEEG electrode placements are color-coded by subject: Subject 1 (red), Subject 2 (green), Subject 3 (blue), Subject 4 (orange), Subject 5 (purple).

2.3.4 Preprocessing

Data from all five subjects were first aggregated into a single dataset. Signals were high-pass (0.5 Hz), low-pass (150 Hz), and notch filtered at 50, 100, and 120 Hz using 4th-order Butterworth IIR filters implemented with FieldTrip. The resulting data were then down-sampled to 256 Hz.

2.3.5 Amplitude of the analytic signal

To estimate the time-resolved amplitude of iEEG activity, broadband signals were bandpass filtered into the same frequency bands as the OPM data using 4th-order Butterworth IIR filters, and the log-transformed amplitude was computed as described in Section 2.1.5.

2.3.6 Cross-modal temporal alignment

For the 338 s of iEEG data, we validated the temporal alignment with the stimulus based on the lag that maximized the cross-correlation between the envelope of the movie's audio-trace and the auditory high-frequency response. To this end, high-frequency activity from the aggregated dataset was averaged across the 20 locations nearest to an anatomical mask of auditory cortex regions defined by the AAL atlas (i.e., we followed the same procedure used to temporally align the fMRI to the movie). Data were then trimmed to the same 297 s as the OPM and fMRI.

Time courses of log-transformed amplitude were, again, resampled to 20 Hz.

2.4 Analyses

2.4.1 Overview and methodological considerations

To identify the shared neural responses elicited across repeated movie viewings, we evaluated the consistency of activity within individuals, between individuals, and across recording modalities. All correlation analyses were performed within separate frequency bands; for fMRI analyses the BOLD signal was used. For OPM and fMRI, coefficients were computed using Pearson product-moment correlations, whereas for iEEG, Spearman rank correlations were used to account for potential outliers in signal amplitudes due to epileptic activity. Because iEEG electrode locations varied across subjects, between-subject comparisons could not be performed within the iEEG dataset.

2.4.2 Within-subject comparisons

For each of the three modalities, subject-level empirical correlations were computed between the time series from the first and second movie viewings.

2.4.3 Between-subject comparisons

Pairwise correlations were computed bidirectionally between one viewing from each subject (first/second) with the alternate viewings (second/first) of the remaining subjects in the same cohort (OPM or fMRI).

2.4.4 Between-method comparisons

2.4.4.1 OPM to fMRI

Cross-modal consistency between OPM and fMRI was assessed by computing Pearson correlations between each movie viewing (first/second) from each OPM subject and the alternate viewing from each fMRI subject (second/first). Fisher z-transformed correlations were then averaged across OPM and fMRI subjects.

2.4.4.2 OPM to iEEG

Spearman correlations were computed between each viewing from each OPM subject (first/second viewing) and the alternate viewing from the single-subject iEEG dataset (second/first viewing). Fisher z-transformed correlation coefficients were averaged across OPM subjects.

2.4.4.3 fMRI to iEEG

To comprehensively compare all three imaging modalities, we evaluated the consistency between iEEG and fMRI to reproduce

the findings of Haufe et al. (2018) with our analysis approach. Spearman correlations were computed bidirectionally between one viewing from each fMRI subject and the alternate viewing from the single-subject iEEG dataset and averaged across fMRI subjects.

2.4.5 Statistical significance and analysis framework

Correlation estimates are sensitive to the autocorrelational structure of neural time series (Schaworonkow et al., 2015; Woolrich et al., 2001), which may be magnified by temporal downsampling and smoothing (James et al., 2019; Mejia et al., 2015; Pajula & Tohka, 2014; Zarahn et al., 1997). To demonstrate this effect, we first correlated fMRI and OPM data at 0.67 Hz (see Table S1, for correlation coefficients). Crucially, because we aimed to quantify signal consistency rather than to maximize correlation coefficients, we resampled all estimates to 20 Hz, a resolution commonly used in the analysis of power spectra in human electrophysiology (e.g., Michelmann et al., 2016; see also: M. X. Cohen, 2014). We then computed surrogate null distributions using the iterative amplitude-adjusted Fourier transform (iAAFT; Schreiber and Schmitz, 1996), and standardized empirical correlations by deriving z-scores relative to the null distribution. We further assessed the significance of observed correlations based on the distribution of averages of surrogate correlation. Specifically, we performed the following steps:

1. Empirical correlations were computed from pairs of real-time series at a given channel (i.e., source position, voxel, or electrode) in a comparison of interest (between viewings, subjects, or modalities). Subject-level empirical correlations were then Fisher z-transformed and averaged across subjects, producing a grand-averaged (GA) empirical correlation coefficient for each channel.

2. We generated 100 surrogate datasets for each time series (movie viewing, subject, frequency band, and modality) and computed surrogate correlations, where one of the empirical time series in a given comparison was replaced with a phase-randomized surrogate time series. The 100 surrogate correlations that correspond to an empirical correlation were also normalized using the Fisher z-transform.

3. Null distributions of averages were then bootstrapped by randomly sampling one surrogate correlation per subject and averaging across subjects to form a single set of GA surrogate correlation coefficients analogous to the empirical GA correlations. This averaging was repeated 10,000 times per comparison, randomly resampling surrogates (i.e., selecting 1 out of

100 options per subject for each surrogate GA). This yielded a null distribution of 10,000 GA surrogate correlation coefficients per channel (Stelzer et al., 2013).

3a. For the iEEG within-subject analysis, subject-level bootstrapping could not be performed. Instead, we repeated step 2 with 1,000 surrogate datasets per channel, using the resulting surrogate correlations directly as the null distribution.

4. p-values were calculated for each channel as the proportion of the surrogate GA correlations that exceeded the empirical GA coefficient. The Benjamini-Hochberg correction was applied to the empirically derived p-values across channels in each analysis, effectively controlling the false discovery rate (FDR) at a level of q = .05.

5. For visualization and interpretation, the empirical GA correlations at each channel were standardized relative to the channel's surrogate distributions by computing z-scores: $z = \frac{(r - \mu_{surr})}{\sigma_{surr}}$, where μ_{surr} and σ_{surr} denote the M and SD of the surrogate null distribution. These z-scores reflect estimates of SNR and enable comparison of SNR on a common scale across modalities and analyses.

2.4.6 Signal-to-noise (SNR) comparisons

To evaluate the relative SNR of OPM signals within and across subjects, we compared the channel-wise SNR estimates by subtracting the within-subject z-scores from the between-subject z-scores for each frequency band. Positive values in the resulting maps indicate greater consistency of neural responses across individuals than within, whereas negative values indicate greater consistency within than across individuals. Cluster-based thresholding was used to identify contiguous channels exceeding the 97.5th or falling below the 2.5th percentile of the observed SNR distribution. The sum of z-scores for each cluster was used to identify the maximum clusters for both the positive and negative tails. We then assessed the significance of the maximum clusters relative to 10,000 surrogate SNR maps. These were computed as the difference between randomly sampled surrogate-within- and between-subject correlation sets derived from the null distributions computed in previous analyses. Empirical p-values were calculated as the proportion of surrogate absolute cluster sums that were bigger than the empirical maximum cluster sums. We controlled the false discovery rate at q = .025 per frequency band.

We performed the same analysis to compare between-subject correlations in OPM to between-method correlations in OPM and fMRI. For each frequency band, we computed channel-wise

SNR maps by subtracting the OPM's between-subject z-scores from the corresponding z-scores in the OPM to fMRI comparison. Because cross-modal comparisons may yield meaningful inverse relationships, we used the absolute value of the between-method z-scores to represent the magnitude of the correlation, regardless of its direction. Thus, resulting positive values indicate greater consistency between OPM and fMRI than across OPM subjects alone, whereas negative values indicate greater consistency across OPM subjects than between OPM and fMRI. Cluster-based permutation testing and significance evaluation followed the same procedure as described above. It is important to note that clusters reflect regions with significant differences, not areas where reliability was equally strong across comparisons.

3. RESULTS

3.1 Within-subject comparisons

Across all three methods, we observed significant reliability in spatially distinct but functionally meaningful areas. For OPM, reliable signal, indicated by channels surviving correction for multiple comparisons (MC), was observed across all frequency bands and spanned regions involved in visual processing, higher-order associative functions, and motor control (Figure 3A). Peak z-scores ranged from 4.65 (γ_1) to 12.20 (δ), with the highest proportion of channels surviving FDR correction in the α band (93.6%; M_z = 6.0, SD_z = 2.3). Signal reliability was broadly distributed in the lower frequency bands (δ, θ, α, and β), with maxima localized to the right precentral gyrus, left middle occipital gyrus, right cuneus,

and right inferior parietal cortex, respectively; in contrast, higher frequency bands (γ_1, γ_2, and HF) displayed more focal distributions, with peaks in the postcentral gyrus, inferior temporal cortex, and paracentral lobule (see Table 1, OPM).

For fMRI, reliable signal was more spatially localized, with 55.9% of voxels surviving MC correction (Table 1, fMRI). Significant effects were observed primarily in regions associated with auditory and visual processing (see Figure 3B), with a peak z-score of 13.43 observed in the left middle temporal gyrus (M_z = 4.4, SD_z = 2.4; see also Table 1, fMRI).

iEEG showed strong within-subject reliability, though spatial extent was more limited, with 11.1–34.1% (refer Table 1, iEEG) of electrodes surviving correction for MC across frequency bands. In the lower frequency bands, reliability was strongest in temporal, occipital, and sensorimotor areas (see Figure 3C); the highest FDR corrected z-score was observed for the high-frequency band concentrated in the left supramarginal gyrus (z = 20.89, M_z = 5.6, SD_z = 3.7).

Although absolute correlation values were highest for fMRI (r_{max} = .32) and iEEG (r_{max} = .33) compared to OPM (r_{max} = .06, compare Table S1), we found that both temporally downsampling the data and "grand averaging" time series across subjects substantially increased OPM correlations (see Table S1, OPM). When re-computed at 0.67 Hz, matching the temporal resolution of fMRI, OPM peak correlations more than doubled across frequency bands (r_{max} = .14), and further increased after

Table 1: Within-subject reliability statistics across modalities and frequency bands.

Modality	Frequency		% MC	M_z	SD_z	Maxima		
						z	MNI Coord.	AAL Label (Left/Right)
OPM	δ	0.5–4 Hz	81.9%	4.2	1.7	12.20	48, –16, 48	Precentral (R)
	θ	4–8 Hz	91.7%	4.0	1.4	9.17	–8, –104, 0	Occipital Mid (L)
	α	8–12 Hz	93.6%	6.0	2.3	11.98	8, –88, 40	Cuneus (R)
	β	12–28 Hz	83.8%	3.8	1.3	8.76	48, –40, 56	Parietal Inf (R)
	γ_1	28–46 Hz	19.7%	3.0	0.5	4.65	–48, –16, 56	Postcentral (L)
	γ_2	55–70 Hz	8.4%	3.1	0.4	5.04	64, –40, –24	Temporal Inf (R)
	HF	64–116 Hz	51.5%	2.8	0.6	4.70	–16, –24, 64	Paracentral Lobule (L)
fMRI	–	–	55.9%	4.4	2.4	13.43	–56, –16, 0	Temporal Mid (L)
iEEG	δ	0.5–4 Hz	32.5%	3.8	1.9	15.87	–58, –14, 48	Postcentral (L)
	θ	4–8 Hz	34.1%	3.8	1.8	11.87	–72, –10, 7	SupraMarginal (R)[a]
	α	8–12 Hz	25.8%	3.6	1.4	9.09	–68, –11, 8	Temporal Sup (L)
	β	12–28 Hz	17.2%	3.7	1.3	8.29	–67, –21, 16	SupraMarginal (L)
	γ_1	28–46 Hz	11.1%	3.7	1.1	6.72	–48, –54, –24	Temporal Inf (L)
	γ_2	55–70 Hz	12.5%	4.2	1.7	9.38	–72, –38, 9	Temporal Mid (L)
	HF	64–116 Hz	26.7%	5.6	3.7	20.89	–67, –21, 16	SupraMarginal (L)

Note. All statistics are reported for data surviving correction for multiple comparisons (MC). For each modality and frequency band, we report the percentage of FDR-correction surviving channels/voxels/electrodes (% MC), the M and SD of z-scores, the peak z-score, its MNI coordinate, and the anatomical label of that location. Superscript a denotes that the first label was found for the electrode with the second highest z-score.

Figure 3: Within-subject reliability for different imaging modalities and frequency bands. FDR-corrected *z*-scores are shown for: (A) OPM, 3581 source locations across 10 subjects; (B) fMRI, 3581 voxels across 11 subjects; (C) iEEG, 656 electrodes pooled across five subjects. For iEEG, *z*-scores were computed relative to a null distribution of 1,000 surrogate correlations, without bootstrapping (see Methods). (D) Schematic of within-subject correlation analysis. Data were bandpass filtered to seven frequency bands: δ (0.5–4 Hz), θ (4–8 Hz), α (8–12 Hz), β (12–28 Hz), γ_1 (28–46 Hz), γ_2 (55–70 Hz), and HF (64–116 Hz). Color scales reflect *z*-scores surviving correction for MC of within-subject correlations, plotted on sensor-specific locations (iEEG) or MNI coordinates (OPM, fMRI).

grand averaging measurements across subjects before computing correlations (r_{max} = .44). These findings illustrate that high test-retest correlations can be achieved with OPM by downsampling and averaging across subjects. Importantly, estimating the SNR via *z*-scores (see above) confirmed robust statistical reliability.

3.2 Between-subject comparisons

OPM exhibited reliable signal across all frequency bands, with effects most pronounced in occipitotemporal and cerebellar regions implicated in perceptual and linguistic processing (see Figure 4A; MacEvoy and Epstein, 2011; Nakatani et al., 2022). Peak *z*-scores ranged from 5.45 (γ_1) to 9.37 (θ), with the greatest proportion of channels surviving correction for MC in the β band (99.9%; M_z = 5.3, SD_z = 1.4; see Table 2, OPM). In lower frequencies, regions of strongest reliability were concentrated in the left postcentral gyrus (δ), right Crus II of the cerebellum (θ), and the left middle occipital gyrus for both α and β bands, suggesting pronounced reliability in posterior visual areas. High-frequency bands exhibited more focal effects, with peak values in the inferior temporal cortex (γ_1), right inferior parietal cortex (γ_2), and fusiform gyrus (HF).

Table 2: Between-subject reliability statistics for OPM and fMRI.

Modality	Frequency Band		% MC	M_z	SD_z	Maxima		
						z	MNI Coord.	AAL Label (Left/Right)
	δ	0.5–4 Hz	53.9%	3.8	1.4	8.97	−64, −16, 32	Postcentral (L)
	θ	4–8 Hz	88.4%	4.1	1.4	9.37	16, −88, −32	Crus II (R)
	α	8–12 Hz	58.6%	3.8	1.4	8.31	−8, −104, 0	Occipital Mid (L)
OPM	β	12–28 Hz	99.9%	5.3	1.4	9.23	−8, −104, −8	Occipital Mid (L)
	γ_1	28–46 Hz	13.9%	3.1	0.5	5.45	−48, −64, −8	Temporal Inf (L)
	γ_2	55–70 Hz	15.6%	3.2	0.6	6.09	56, −56, 40	Parietal Inf (R)
	HF	64–116 Hz	45.4%	3.0	0.6	5.47	32, −64, −8	Fusiform (R)
fMRI	–	–	24.3%	4.5	2.0	11.88	−48, −24, 8	Temporal Sup (L)

Note. All statistics are reported for data surviving MC correction. For each modality and frequency band, this table reports the percentage of voxels/channels/electrodes surviving FDR correction (% MC), the *M* and *SD* of *z*-scores, and the peak *z*-score with its MNI coordinate and anatomical label.

Figure 4: Between-subject reliability for OPM and fMRI. FDR-corrected *z*-scores are shown for: (A) OPM, 3,581 source locations across 10 subjects for 7 frequency bands; (B) fMRI: 3581 voxels, across 11 subjects. (C) Schematic of between-subject correlation analysis. Color bars reflect *z*-scores (surviving correction for MC).

For fMRI, although fewer voxels survived correction for MC (24.3%; Table 2, fMRI) compared to the within-subject analysis, the spatial pattern was largely preserved, with effects localized to auditory and visual regions (M_z = 4.5, SD_z = 2.0; see Figure 4B). The maximum was observed in the left superior temporal gyrus (peak *z* = 11.88), consistent with its role in processing complex audio-visual stimuli (Ghinst et al., 2016; Park et al., 2018; Ye et al., 2017).

3.3 Between-method comparisons

3.3.1 OPM to fMRI.

Consistent cross-modal alignment between OPM and fMRI was observed across all frequencies, with the most pronounced effects localized to visual processing regions in the mid-range and high-frequency bands (see Figure 5A). After correction for MC, peak *z*-scores ranged from -9.36 (β) to 6.50 (HF), with the largest proportion of surviving channels observed in the β band (12.1%; M_z = -3.3, SD_z = 2.9; Table 5, OPM to fMRI). Importantly, we observed negative cross-modal correlations in the δ, θ, α, and β bands, and positive correlations in the γ_1, γ_2, and high-frequency bands, in line with previous demonstrations of the relationship between MEG power spectra and BOLD signal (Zumer et al., 2010).

Of the frequency bands that demonstrated broader topographies (θ, α and β), peak *z*-scores were located in the left superior parietal cortex (θ) and the left middle occipital cortex (α and β), with additional focal peaks observed for γ_2 and HF in bilateral cuneus regions. These spatial patterns are consistent with prior

findings demonstrating that MEG and fMRI exhibit the greatest intersubject consistency in occipital areas during naturalistic movie viewing (Lankinen et al., 2018).

3.3.2 OPM to iEEG.

Although overall cross-modal consistency between OPM and iEEG was modest, significant effects were observed in lower frequency bands, spanning frontal, parietal, and ventral visual regions (see Figure 5B). As reported in Table 5, OPM to iEEG, peak *z*-scores ranged from 3.68 (γ_2) to 4.69 (θ), with relatively few electrodes surviving MC correction at each frequency (maximum of 1.1% for θ; M = 4.1, SD = 0.5).

Spatially, peak cross-modal correlations in the lower bands were localized to the left middle frontal gyrus (δ), precuneus (θ), and fusiform gyrus (α). At higher frequencies, significant effects were sparse, with only one isolated electrode in the left supplementary motor area surviving MC correction (γ_2). No electrodes survived correction for MC in the β, γ_1, or HF bands.

3.3.3 fMRI to iEEG.

To provide a comprehensive evaluation of the consistency between all three imaging modalities, we reproduced the results of Haufe et al. (2018) comparing iEEG to fMRI.

We observed significant cross-modal correlations between fMRI and iEEG across all frequency bands, with effects spanning temporal, parietal, and occipital regions (Figure 5C), consistent with Haufe et al. (2018). As in our fMRI to OPM comparison, negative correlations were observed in lower-frequency bands (δ-β), while (γ) and high-frequency bands showed positive associations

with the BOLD signal (see Table 5, fMRI to iEEG). Peak *z*-scores ranged from -7.98 (θ) to 8.86 (HF), with the largest proportion of channels surviving correction for MC in the HF band (8.7%, M_z = 4.8, SD_z = 1.7).

Peak effects were largely located in areas associated with audi-tory and linguistic processing (Bhaya-Grossman & Chang, 2022; Wernicke, 1974), including the middle (θ and HF) and superior (α and γ_1) temporal gyri. This pattern is consistent with prior work demonstrating strong BOLD coupling to oscillatory activity in au-ditory cortex during movie viewing (Mukamel et al., 2005).

Table 3: Between-method reliability statistics across frequency bands for OPM, fMRI, and iEEG.

Comparison	Frequency Band		% MC	M_z	SD_z	Maxima		
						z	MNI Coord.	AAL Label (Left/Right)
OPM to fMRI	δ	0.5–4 Hz	0.8%	−3.6	2.3	−5.69	24, –8, 64	Frontal Sup (R)
	θ	4–8 Hz	11.9%	−2.8	3.3	−7.75	–24, –56, 64	Parietal Sup (L)
	α	8–12 Hz	8.9%	−3.8	2.3	−8.97	–40, –88, 8	Occipital Mid (L)
	β	12–28 Hz	12.1%	−3.3	2.9	−9.36	–32, –88, 16	Occipital Mid (L)
	γ_1	28–46 Hz	0.1%	3.7	0.1	3.77	–64, –32, 0	Temporal Mid (L)
	γ_2	55–70 Hz	1.6%	3.9	0.5	5.36	16, –88, 24	Cuneus (R)
	HF	64–116 Hz	3.5%	3.9	1.2	6.50	–8, –80, 24	Cuneus (L)
OPM to iEEG	δ	0.5–4 Hz	0.3%	4.2	0.2	4.30	–55, 19, 38	Frontal Mid (L)
	θ	4–8 Hz	1.1%	4.1	0.5	4.69	–8, –56, 70	Precuneus (L)
	α	8–12 Hz	0.8%	4.1	0.3	4.48	–44, –61, –21	Fusiform (L)
	β	12–28 Hz	0%	–	–	–	–	–
	γ_1	28–46 Hz	0%	–	–	–	–	–
	γ_2	55–70 Hz	0.2%	3.7	0.0	3.68	–9, 8, 50	Supp Motor Area (L)
	HF	64–116 Hz	0%	–	–	–	–	–
fMRI to iEEG	δ	0.5–4 Hz	4.1%	−4.6	1.4	−7.71	–11, –64, 75	Precuneus (L)
	θ	4–8 Hz	7.8%	−4.3	1.6	−7.98	–74, –28, 3	Temporal Mid (L)
	α	8–12 Hz	8.1%	−4.3	1.3	−7.84	–68, –11, 8	Temporal Sup (L)
	β	12–28 Hz	5.2%	−4.3	1.1	−7.36	–35, –91, 22	Occipital Mid (L)
	γ_1	28–46 Hz	4.0%	3.5	2.7	6.11	–67, –8, 3	Temporal Sup (L)
	γ_2	55–70 Hz	4.7%	4.2	2.4	7.73	–21, –55, 78	Parietal Sup (L)
	HF	64–116 Hz	8.7%	4.8	1.7	8.86	–74, –28, 3	Temporal Mid (L)

Note. All statistics are reported for data surviving correction for MC. For each comparison and band, we report the percent of vox-els/electrodes surviving correction for MC (% MC), *M* and *SD* of *z*-scores, and the peak *z*-score with its MNI coordinate and anatomical label.

Figure 5: Consistent between-method alignment. (A) OPM to fMRI: *z*-scores from OPM subjects (n = 10) and fMRI subjects (n = 11) are shown for 3581 locations. (B) OPM to iEEG: *z*-scores from OPM subjects (n = 10) and single-subject iEEG from 656 electrodes pooled across five subjects. (C) fMRI to iEEG: *z*-scores from fMRI subjects (n = 11) and single-subject iEEG from 656 electrodes pooled across five subjects. Color bars reflect *z*-scores (see above).

3.4 Signal-to-noise (SNR) comparisons

3.4.1 Comparison of within-subject SNR to between-subject SNR in OPM.

We observed clusters of significantly reduced between-subject SNR relative to within-subject SNR for OPM across all frequency bands ($p < .001$, controlling the false discovery rate at $q < .025$, as per cluster-based permutation; See methods). Maximum negative clusters were spatially distributed across sensorimotor, parietal, and frontal regions, reflecting areas with greater variabil-

ity between subjects (Figure 6). The most pronounced SNR loss was observed in the α band, which yielded the largest significant negative cluster (90 channels, sum = -874.69), corresponding in the observed data to a region centered on the right superior frontal gyrus (peak z = -11.16, M_z = -9.7, SD_z = 0.5). As shown in the top panel of Table 4, significant decreases in SNR were also found across δ (sum = -211.61, max in precentral gyrus), θ (sum = -118.97, max in anterior cingulate cortex), β (sum = -143.82, max in precentral gyrus), γ_1 (sum = -276.67, max in postcentral gyrus), γ_2 (sum = -170.30, max in superior parietal lobule), and HF (sum

= -255.48, max in supramarginal gyrus).

Additionally, all frequencies revealed clusters of significant positive SNR, representing areas where neural responses were more consistent across the OPM cohort than within individuals. These clusters were consistently left-lateralized and largely localized to frontal and posterior association areas (see Figure 6), consistent with previous findings of intersubject synchrony during emotionally engaging movie viewing (Jääskeläinen et al., 2008; Maffei, 2020). Frontal peaks spanned the gyrus rectus (γ_1, sum = 129.72), as well as the superior (δ, sum = 236.62) and middle frontal gyri (γ_2, sum = 231.19; HF, sum = 191.87), with the largest cluster observed in the orbital part of the middle frontal gyrus (β band, 51 channels, sum = 342.16, z = 7.82; Table 4, bottom panel). Additional positive clusters were found in θ (sum = 137.57, max in middle occipital gyrus) and α (sum = 132.54, max in angular gyrus). Such higher SNR between different subjects can stem from a decrease in shared noise that exceeds the decrease in shared signal.

Table 4: Cluster statistics for SNR differences in within- and between-subject OPM comparisons.

Cluster Type	Frequency Band		Size	Sum	M_z	SD_z	Cluster Maxima		
							z	MNI Coord.	AAL Label (L/R)
Negative	δ	0.5–4 Hz	38	−211.61	−5.6	0.9	−7.54	40, −16, 64	Precentral (R)
	θ	4–8 Hz	28	−118.97	−4.3	0.7	−5.55	0, 24, −8	Cingulum Ant (L)
	α	8–12 Hz	90	−874.69	−9.7	0.5	−11.16	16, −8, 64	Frontal Sup (R)
	β	12–28 Hz	48	−143.82	−3.0	0.5	−4.20	24, −24, 72	Precentral (R)
	γ_1	28–46 Hz	48	−276.67	−5.8	0.7	−7.25	−64, −16, 24	Postcentral (L)
	γ_2	55–70 Hz	42	−170.30	−4.1	0.7	−5.81	−16, −64, 64	Parietal Sup (L)
	HF	64–116 Hz	75	−255.48	−3.4	0.3	−4.45	−56, −48, 32	SupraMarginal (L)
Positive	δ	0.5–4 Hz	77	236.62	3.1	0.6	4.74	−24, −8, 72	Frontal Sup (L)
	θ	4–8 Hz	30	137.57	4.6	0.8	6.84	−32, −96, −8	Occipital Mid (L)
	α	8–12 Hz	57	132.54	2.3	0.6	3.71	−32, −48, 0	Angular (L)[a]
	β	12–28 Hz	51	342.16	6.7	0.5	7.82	−16, 32, −8	Frontal Mid Orb (L)[b]
	γ_1	28–46 Hz	33	129.72	3.9	0.4	5.39	0, 48, −24	Rectus (L)
	γ_2	55–70 Hz	53	231.19	4.4	0.8	6.02	−24, 24, 16	Frontal Mid (L)[a]
	HF	64–116 Hz	51	191.87	3.7	0.5	5.10	16, 40, 56	Frontal Mid (L)[b]

Note. All statistics are reported for the largest cluster per frequency band surviving MC correction (q < .025). Negative clusters (top) reflect higher within- than between-subject reliability. Positive clusters (bottom) reflect higher between- than within-subject reliability. For each cluster, we report its size (number of channels), the sum, *M*, and *SD* of *z*-scores, and the peak *z*-score along with its MNI coordinate and anatomical label. Superscript a denotes that the first label was found in the channel with the third highest *z*-score. Superscript b denotes that the first label was found for the voxel with the second highest *z*-score.

Figure 6: SNR differences in within- and between-subject comparisons for OPM. Within-modality SNR differences in OPM were computed by subtracting within-subject from between-subject *z*-scores at each channel. Negative values (blue) indicate greater signal consistency within individuals than across individuals; positive values (green) indicate greater signal consistency across individuals than within individuals. Significant clusters surviving MC correction (FDR *q* < .025) are outlined in the respective color. SNR changes are shown across frequency bands from δ (0.5- 4 Hz) to high-frequency (64-116 Hz). Diagrams below illustrate the comparison logic. Maps of uncorrected clusters identified by the initial cluster-forming threshold are shown in Figure S3A.

3.4.2 SNR comparison of intersubject correlations in OPM and inter-method correlations between OPM and fMRI.

Cluster-based permutation testing revealed significant negative clusters surviving MC correction in all frequency bands, indicating greater signal consistency within modality (OPM) than across modalities (OPM to fMRI). These effects spanned frontal, parietal, occipital, and cerebellar regions (see Figure 7), with the largest cluster appearing in the δ band (69 channels, sum = -451.38), corresponding to a region centered on the left postcentral gyrus (z = -8.68, M_z = -6.5, SD_z = 0.7; Table 5, top panel). Interestingly, negative clusters were also observed in the angular gyrus (α, sum = -527.20; γ_2, sum = -162.03), an area which previously showed weak intersubject coherence in fMRI (Hasson et al., 2004). Moreover, greater within-modality consistency in cerebellar regions (θ, sum = -313.54; β, sum = -291.23) aligns with prior work demonstrating that OPM is capable of capturing activity in the cerebellum (Lin et al., 2019).

Surprisingly, significant positive clusters also emerged across all frequency bands, indicating that, in some regions, signal correspondence between OPM and fMRI exceeded within-modality reliability. These clusters were largely concentrated in auditory and visual processing areas (see Figure 7), likely indicating convergent stimulus-evoked activity captured by both OPM and fMRI during audio-visual movie viewing. Consistent with prior findings on high-frequency oscillations in visual processing (Martinovic & Busch, 2011; Mukamel et al., 2005), the largest HF-band cluster (83 channels, sum = 309.68, Table 5, bottom panel) localized to the left superior occipital gyrus. A similar pattern emerged in the γ_2 band, where, despite OPM's fragmentary spatial pattern, alignment with fMRI was robust in visual cortex, peaking in the left lingual gyrus (sum = 155.04). Complementary to these effects, positive clusters for θ and γ_1 were localized to the left middle temporal pole (sum = 74.16) and supramarginal gyrus (sum = 184.74), respectively, in line with proposed oscillatory dynamics during speech processing (Giraud & Poeppel, 2012). These findings indicate that, although within-modality reliability is generally higher, cross-modal correspondence may exceed it in anatomically and spectrally specific patterns.

Table 5: Cluster statistics for SNR differences in between-subject OPM and between-method OPM to fMRI comparisons.

Cluster Type	Frequency Band		Size	Sum	M_z	SD_z	Cluster Maxima		
							z	MNI Coord.	AAL Label (L/R)
Negative	δ	0.5–4 Hz	69	−451.38	−6.5	0.7	−8.68	−56, −16, 48	Postcentral (L)
	θ	4–8 Hz	44	−313.54	−7.1	0.7	−8.91	24, −88, −24	Cerebellum Crus I (R)
	α	8–12 Hz	47	−285.01	−6.1	0.5	−7.32	−40, −64, 32	Angular (L)
	β	12–28 Hz	37	−291.23	−7.9	0.4	−8.98	8, −64, −56	Cerebellum VIII (R)[a]
	γ_1	28–46 Hz	17	−58.75	−3.5	0.4	−4.15	−32, −64, 0	Occipital Mid (L)[a]
	γ_2	55–70 Hz	42	−162.03	−3.9	0.6	−5.14	40, −64, 48	Angular (R)
	HF	64–116 Hz	44	−170.67	−3.9	0.3	−4.78	16, 8, 8	Frontal Mid (R)[b]
Positive	δ	0.5–4 Hz	43	126.08	2.9	0.6	4.17	−24, −80, −8	Fusiform (L)
	θ	4–8 Hz	38	74.16	2.0	0.4	2.68	−48, 16, −32	Temporal Pole Mid (L)
	α	8–12 Hz	43	134.24	3.1	0.5	4.18	48, 24, −24	Temporal Pole Sup (R)
	β	12–28 Hz	48	66.24	1.4	0.2	4.00	−16, −72, 48	Parietal Sup (L)
	γ_1	28–46 Hz	40	184.74	4.6	0.8	7.22	−56, −24, 24	SupraMarginal (L)
	γ_2	55–70 Hz	35	155.04	4.4	0.7	6.28	24, −64, 0	Lingual (R)
	HF	64–116 Hz	83	309.68	3.7	0.8	5.83	−16, −80, 32	Occipital Sup (L)

Note. All statistics are reported for the largest cluster per frequency band surviving MC correction (q < .025). Negative clusters (top) reflect greater reliability between OPM subjects than when compared to fMRI. Positive clusters (bottom) reflect greater cross-modal correspondence between OPM and fMRI than within the OPM group alone. For each cluster, we report its size (number of channels), the sum, M, and SD of z-scores, and the peak z-score along with its MNI coordinate and anatomical label. Superscript a denotes that the first label was found in the channel with the second highest z-score. Superscript b denotes that the first label was found in the channel with the third highest z-score.

Figure 7: SNR differences within and across OPM and fMRI. Between-modality SNR differences contrast the SNR from between-subject OPM comparisons with the SNR from between-method OPM-to-fMRI comparisons. Negative values (green) indicate greater signal reliability within the OPM sample than in the cross-modal comparison. Positive values (red) indicate greater signal reliability in the cross-modal comparison (fMRI-to-OPM) than within the OPM sample. Significant clusters surviving MC correction (FDR q < .025) are outlined in the respective color. SNR changes are shown across frequency bands from δ (0.5- 4 Hz) to high-frequency (64-116 Hz). Diagrams below illustrate the comparison logic for each panel. Maps of uncorrected clusters identified by the initial cluster-forming threshold are shown in Figure S3B.

4. DISCUSSION

We systematically evaluated the reliability of time-resolved neural signals in 7 frequency bands captured by OPM, and their cross-modal alignment with fMRI and iEEG during repeated viewings of a movie stimulus. OPM demonstrated widespread within-subject reliability, especially at lower frequencies; these results were consistent with SQUID-MEG studies of naturalistic movie viewing (Lankinen et al., 2018), but showed a broader spatial distribution that extended beyond occipital regions. We further observed reliable correlations *between* subjects in our OPM sample; their SNR exceeded, in some cases, that of within-subject analyses, which suggests that a shared signal consistently exceeded noise across individuals. Moreover, spatial maps became increasingly focused to sensory regions, mirroring patterns reported in previous intersubject analyses (Chang et al., 2015; Hasson et al., 2004; Haufe et al., 2018; Lankinen et al., 2014).

Correlations of OPM signals produced SNR estimates comparable in magnitude to those observed in fMRI and iEEG and with convergent spatial patterns in visual and auditory regions. OPM signals also directly correlated with both modalities; the cross-modal comparisons revealed substantially stronger OPM-fMRI alignment than OPM-iEEG, with the largest effects localized to visual areas. Contrary to previous SQUID–MEG based findings (Lankinen et al., 2018), OPM achieved reliable correlations with fMRI and showed frequency-dependent shifts in correlation-direction that are consistent with the inverse relationship be-

tween low-frequency oscillatory power and BOLD signals (Zumer et al., 2010).

Still, shared activity between two modalities does not imply overall cross-modal congruence. For instance, although both OPM and iEEG demonstrated strong alignment with fMRI, their correspondence with each other was comparatively modest. This discrepancy may reflect a combination of iEEG's sparse electrode coverage, sensitivity to localized high-frequency activity, and modality-specific capacity to detect different neural generators: MEG and iEEG, which are most sensitive to sulcal and gyral sources, respectively, may capture distinct yet complementary neural signals that uniquely contribute to the BOLD response (Zumer et al., 2010). Nevertheless, our results suggest that OPM and fMRI converge on temporally structured neural activity, which, in the context of our movie viewing paradigm, likely relates to modulations of power during ongoing visual processing (Brookes et al., 2005). Collectively, these results demonstrate that OPM reliably captures stimulus-locked neural dynamics shared across individuals and imaging modalities.

One advantage of our analysis approach is that it directly quantifies the reliability of shared signals relative to shared noise. In our analyses, Pearson correlation coefficients (*r*-values) increased substantially when down-sampling to 0.67 Hz. The computed *z*-scores, on the other hand, account for temporal autocorrelation and noise structure, offering an appropriate metric of signal reliability relative to surrogate data (Haufe et al., 2018). Fur-

thermore, by conducting pairwise analyses at a common sampling rate of 20 Hz, we preserved intersubject variance and temporal precision and were able to demonstrate the reliability of OPM both within and between individuals. Significant effects emerged despite minimal preprocessing and early-generation sensor hardware, which underscores the efficacy of OPM that can likely be improved with future advances in sensor design and denoising techniques (Rier et al., 2023).

Importantly, estimating SNR via *z*-scores places all modalities on a shared statistical scale, allowing for comparisons across datasets with differing temporal resolution, spatial coverage, and sample sizes. Leveraging this approach, we observed that within-subject analyses yielded more widespread reliability than between-subject comparisons. Indeed, between-subject comparisons appeared spatially focused in primary sensory regions, consistent with neural responses commonly evoked during movie viewing (Hasson et al., 2004; Lankinen et al., 2014). I.e., while it is often considered a limitation of a design when data are acquired from separate participant groups, because of the introduction of subject-specific variability into comparisons, our findings suggest that intersubject analyses can also mitigate idiosyncratic biases by attenuating shared noise (e.g., spatial leakage in source reconstruction) and task-unrelated fluctuations. In our data, between-subject comparisons effectively isolated stimulus-driven responses generalizable across individuals (Hasson et al., 2004, 2010; Nastase et al., 2019). Our SNR comparisons further suggest that this principle can extend across imaging modalities. Speculatively, within-subject designs with cross-modal comparisons may also overestimate reliability by capturing stable but characteristic individual signal components such as subject-specific lateralization.

Finally, we observed that SNR can – in some regions – increase when comparisons are drawn between different subjects undergoing different imaging modalities. While differences in modality sensitivity often reduce overall alignment, they may also support the suppression of shared noise and can consequently sometimes lead to increased SNR. In particular, while high-frequency OPM activity appeared spatially diffuse, its correlation with fMRI was tightly localized to regions expected to support visual processing. I.e., while high-frequency measurements in OPM may not be stable enough to be reliably localized across subjects or measurements within OPM, its similarity to fMRI signal allowed for the demonstration that it is reliably present. Notably, such preserved shared signal can only be detected in regions where the modalities functionally align. In the future, cross-modal convergence may be leveraged to guide denoising and source reconstruction, effectively integrating the spatial precision of fMRI with the temporal resolution of MEG.

In conclusion, our findings demonstrate that during audio-visual processing, OPM achieves reliability levels comparable to fMRI and iEEG; these effects were most robust in low-frequency bands. OPM showed strong correspondence with BOLD signals, following the established patterns of inverse relationships between high- and low-frequency neural activity. These results underscore the potential of OPM as a flexible, noninvasive tool capable of resolving complex brain activity across multiple temporal and spatial scales.

DATA AND CODE AVAILABILITY

Upon publication of this manuscript, all data and analysis code will be made openly available at https://osf.io/urjvb.

AUTHOR CONTRIBUTIONS

Olivia R. Christiano: Conceptualization; Data curation; Formal analysis; Project administration; Software; Visualization; Writing original draft. Sebastian Michelmann: Conceptualization; Data curation; Formal analysis; Methodology; Resources; Software; Supervision; Writing review & editing.

DECLARATION OF COMPETING INTERESTS

The authors declare that there are no competing interests.

ACKNOWLEDGMENTS

We would like to thank Uri Hasson and Christopher J. Honey, who generously shared the data that made this work possible, as well as Stefan Haufe, who made the preprocessed fMRI data from Arcaro et al. (2015) available and pioneered the analysis approach. We also thank Lukas Rier and coauthors, notably Elena Boto and Matthew J. Brookes, for making the OPM data publicly available and for valuable comments on early conceptualization and analyses of this study.

SUPPLEMENTARY MATERIAL

Supplementary material for this article is included at the end of this document.

REFERENCES

An, K. M., Shim, J. H., Kwon, H., Lee, Y. H., Yu, K. K., Kwon, M., Chun, W. Y., Hirosawa, T., Hasegawa, C., Iwasaki, S., Kikuchi, M., & Kim, K. (2022). Detection of the 40 hz auditory steady-state

response with optically pumped magnetometers. *Scientific Reports*, *12*(1). https://doi.org/10.1038/s41598-022-21870-5

Andersen, L. M., Oostenveld, R., Pfeiffer, C., Ruffieux, S., Jousmäki, V., Hämäläinen, M., Schneiderman, J. F., & Lundqvist, D. (2017). Similarities and differences between on-scalp and conventional in-helmet magnetoencephalography recordings. *PLoS ONE*, *12*(7), e0178602. https://doi.org/10.1371/journal.pone.0178602

Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *NeuroImage*, *38*(1), 95–113. https://doi.org/10.1016/j.neuroimage.2007.07.007

Bhaya-Grossman, I., & Chang, E. F. (2022). Speech computations of the human superior temporal gyrus. *Annual Review of Psychology*, *73*(1), 79–102. https://doi.org/10.1146/annurev-psych-022321-035256

Boto, E., Bowtell, R., Krüger, P., Fromhold, T. M., Morris, P. G., Meyer, S. S., Barnes, G. R., & Brookes, M. J. (2016). On the potential of a new generation of magnetometers for meg: A beamformer simulation study. *PLoS ONE*, *11*(8), e0157655. https://doi.org/10.1371/journal.pone.0157655

Boto, E., Hill, R. M., Rea, M., Holmes, N., Seedat, Z. A., Leggett, J., Shah, V., Osborne, J., Bowtell, R., & Brookes, M. J. (2021). Measuring functional connectivity with wearable meg. *NeuroImage*, *230*, 117815. https://doi.org/10.1016/J.NEUROIMAGE.2021.117815

Boto, E., Holmes, N., Leggett, J., Roberts, G., Shah, V., Meyer, S. S., Muñoz, L. D., Mullinger, K. J., Tierney, T. M., Bestmann, S., Barnes, G. R., Bowtell, R., & Brookes, M. J. (2018). Moving magnetoencephalography towards real-world applications with a wearable system. *Nature*, *555*(7698), 657–661. https://doi.org/10.1038/nature26147

Boto, E., Meyer, S. S., Shah, V., Alem, O., Knappe, S., Kruger, P., Fromhold, T. M., Lim, M., Glover, P. M., Morris, P. G., Bowtell, R., Barnes, G. R., & Brookes, M. J. (2017). A new generation of magnetoencephalography: Room temperature measurements using optically-pumped magnetometers. *NeuroImage*, *149*, 404–414. https://doi.org/10.1016/j.neuroimage.2017.01.034

Brookes, M. J., Boto, E., Rea, M., Shah, V., Osborne, J., Holmes, N., Hill, R. M., Leggett, J., Rhodes, N., & Bowtell, R. (2021). Theoretical advantages of a triaxial optically pumped magnetometer magnetoencephalography system. *NeuroImage*, *236*, 118025. https://doi.org/10.1016/j.neuroimage.2021.118025

Brookes, M. J., Gibson, A. M., Hall, S. D., Furlong, P. L., Barnes, G. R., Hillebrand, A., Singh, K. D., Holliday, I. E., Francis, S. T., & Morris, P. G. (2005). Glm-beamformer method demonstrates stationary field, alpha erd and gamma ers co-localisation with fmri bold response in visual cortex. *NeuroImage*, *26*(1), 302–308. https://doi.org/10.1016/j.neuroimage.2005.01.050

Chang, W. T., Jääskeläinen, I. P., Belliveau, J. W., Huang, S., Hung, A. Y., Rossi, S., & Ahveninen, J. (2015). Combined meg and eeg show reliable patterns of electromagnetic brain activity during natural viewing. *NeuroImage*, *114*, 49–56. https://doi.org/10.1016/j.neuroimage.2015.03.066

Cohen, D. (1968). Magnetoencephalography: Evidence of magnetic fields produced by alpha-rhythm currents. *Science*, *161*(3843), 784–786. https://doi.org/10.1126/science.161.3843.784

Cohen, D. (1972). Magnetoencephalography: Detection of the brain's electrical activity with a superconducting magnetometer. *Science*, *175*(4022), 664–666. https://doi.org/10.1126/science.175.4022.664

Cohen, M. X. (2014). *Analyzing neural time series data*. The MIT Press. https://doi.org/10.7551/mitpress/9609.001.0001

Cox, R. W. (1996). Afni: Software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research*, *29*(3), 162–173. https://doi.org/10.1006/cbmr.1996.0014

Ghinst, M. V., Bourguignon, M., de Beeck, M. O., Wens, V., Marty, B., Hassid, S., Choufani, G., Jousmäki, V., Hari, R., Bogaert, P. V., Goldman, S., & Tiège, X. D. (2016). Left superior temporal gyrus is coupled to attended speech in a cocktail-party auditory scene. *Journal of Neuroscience*, *36*(5), 1596–1606. https://doi.org/10.1523/JNEUROSCI.1730-15.2016

Giraud, A.-L., & Poeppel, D. (2012). Cortical oscillations and speech processing: Emerging computational principles and operations. *Nature Neuroscience*, *15*(4), 511–517. https://doi.org/10.1038/nn.3063

Hämäläinen, M., Hari, R., Ilmoniemi, R. J., Knuutila, J., & Lounasmaa, O. V. (1993). Magnetoencephalographytheory, instrumentation, and applications to noninvasive studies of the working human brain. *Reviews of Modern Physics*, *65*(2), 413–497. https://doi.org/10.1103/RevModPhys.65.413

Hasson, U., Malach, R., & Heeger, D. J. (2010). Reliability of cortical activity during natural stimulation. *Trends in Cognitive Sciences*, *14*(1), 40–48. https://doi.org/10.1016/j.tics.2009.10.011

Hasson, U., Nir, Y., Levy, I., Fuhrmann, G., & Malach, R. (2004). Inter-subject synchronization of cortical activity during natural vision. *Science*, *303*(5664), 1634–1640. https://doi.org/10.1126/science.1089506

Haufe, S., DeGuzman, P., Henin, S., Arcaro, M., Honey, C. J., Hasson, U., & Parra, L. C. (2018). Elucidating relations between fmri, ecog, and eeg through a common natural stimulus. *NeuroImage*, *179*, 79–91. https://doi.org/10.1016/j.neuroimage.2018.06.016

Hill, R., Boto, E., Rea, M., Holmes, N., Leggett, J., Coles, L., Papastavrou, M., Everton, S., Hunt, B., Sims, D., Bowtell, R., & Brookes, M. (2020). Multi-channel whole-head opm-meg: Helmet design and a comparison with a conventional system. *NeuroImage*, *219*, 116995. https://doi.org/10.1016/j.neuroimage.2020.116995

Honey, C. J., Thesen, T., Donner, T. H., Silbert, L. J., Carlson, C. E., Devinsky, O., Doyle, W. K., Rubin, N., Heeger, D. J., & Hasson, U. (2012). Slow cortical dynamics and the accumulation of information over long timescales. *Neuron*, *76*(2), 423–434. https://doi.org/10.1016/j.neuron.2012.08.011

Iivanainen, J., Carter, T. R., Trumbo, M. C., McKay, J., Taulu, S., Wang, J., Stephen, J. M., Schwindt, P. D., & Borna, A. (2023). Single-trial classification of evoked responses to auditory tones using opm- and squid-meg. *Journal of Neural Engineering*, *20*(5), 056032. https://doi.org/10.1088/1741-2552/acfcd9

Iivanainen, J., Stenroos, M., & Parkkonen, L. (2017). Measuring meg closer to the brain: Performance of on-scalp sensor arrays. *NeuroImage*, *147*, 542–553. https://doi.org/10.1016/J.NEUROIMAGE.2016.12.048

Jääskeläinen, I. P., Koskentalo, K., Balk, M. H., Autti, T., Kauramäki, J., Pomren, C., & Sams, M. (2008). Inter-subject synchronization of prefrontal cortex hemodynamic activity during natural viewing. *The Open Neuroimaging Journal*, *2*(1), 14–19. https://doi.org/10.2174/1874440000802010014

James, O., Park, H., & Kim, S. G. (2019). Impact of sampling rate on statistical significance for single subject fmri connectivity analysis. *Human Brain Mapping*, *40*(11), 3321–3337. https://doi.org/10.1002/hbm.24600

Lankinen, K., Saari, J., Hari, R., & Koskinen, M. (2014). Intersubject consistency of cortical meg signals during movie viewing. *NeuroImage*, *92*, 217–224. https://doi.org/10.1016/j.neuroimage.2014.02.004

Lankinen, K., Saari, J., Hlushchuk, Y., Tikka, P., Parkkonen, L., Hari, R., & Koskinen, M. (2018). Consistency and similarity of meg- and fmri-signal time courses during movie viewing. *NeuroImage*, *173*, 361–369. https://doi.org/10.1016/j.neuroimage.2018.02.045

Lew, B. J., Fitzgerald, E. E., Ott, L. R., Penhale, S. H., & Wilson, T. W. (2021). Three-year reliability of meg resting-state oscillatory power. *NeuroImage*, *243*, 118516. https://doi.org/10.1016/j.neuroimage.2021.118516

Lin, C.-H., Tierney, T. M., Holmes, N., Boto, E., Leggett, J., Bestmann, S., Bowtell, R., Brookes, M. J., Barnes, G. R., & Miall, R. C. (2019). Using optically pumped magnetometers to measure magnetoencephalographic signals in the human cerebellum. *The Journal of Physiology*, *597*(16), 4309–4324. https://doi.org/10.1113/JP277899

Liu, X., Dai, Y., Xie, H., & Zhen, Z. (2022). A studyforrest extension, meg recordings while watching the audio-visual movie forrest gump. *Scientific Data*, *9*(1). https://doi.org/10.1038/s41597-022-01299-1

Lumet, S. (1975). *Dog day afternoon* [Film]. Warner Bros.

MacEvoy, S. P., & Epstein, R. A. (2011). Constructing scenes from objects in human occipitotemporal cortex. *Nature Neuroscience*, *14*(10), 1323–1329. https://doi.org/10.1038/nn.2903

Maffei, A. (2020). Spectrally resolved eeg intersubject correlation reveals distinct cortical oscillatory patterns during free-viewing of affective scenes. *Psychophysiology*, *57*(11). https://doi.org/10.1111/psyp.13652

Marhl, U., Jodko-Wadziska, A., Brühl, R., Sander, T., & Jazbinek, V. (2022). Transforming and comparing data between standard squid and opm-meg systems. *PLoS ONE*, *17*(1), e0262669. https://doi.org/10.1371/journal.pone.0262669

Martín-Buro, M. C., Garcés, P., & Maestú, F. (2016). Test-retest reliability of resting-state magnetoencephalography power in sensor and source space. *Human Brain Mapping*, *37*(1), 179–190. https://doi.org/10.1002/hbm.23027

Martinovic, J., & Busch, N. A. (2011). High frequency oscillations as a correlate of visual perception. *International Journal of Psy-*

chophysiology, 79(1), 32–38. https://doi.org/10.1016/j.ijpsycho.2010.07.004

Mejia, A. F., Nebel, M. B., Shou, H., Crainiceanu, C. M., Pekar, J. J., Mostofsky, S., Caffo, B., & Lindquist, M. A. (2015). Improving reliability of subject-level resting-state fmri parcellation with shrinkage estimators. NeuroImage, 112, 14–29. https://doi.org/10.1016/J.NEUROIMAGE.2015.02.042

Michelmann, S., Bowman, H., & Hanslmayr, S. (2016). The temporal signature of memories: Identification of a general mechanism for dynamic memory replay in humans. PLOS Biology, 14(8), e1002528. https://doi.org/10.1371/journal.pbio.1002528

Mukamel, R., Gelbard, H., Arieli, A., Hasson, U., Fried, I., & Malach, R. (2005). Neuroscience: Coupling between neuronal firing, field potentials, and fmri in human auditory cortex. Science, 309(5736), 951–954. https://doi.org/10.1126/science.1110913

Nakatani, H., Nakamura, Y., & Okanoya, K. (2022). Respective involvement of the right cerebellar crus i and ii in syntactic and semantic processing for comprehension of language. The Cerebellum, 22(4), 739–755. https://doi.org/10.1007/s12311-022-01451-y

Nastase, S. A., Gazzola, V., Hasson, U., & Keysers, C. (2019). Measuring shared responses across subjects using intersubject correlation. Social Cognitive and Affective Neuroscience, 14(6), 667–685. https://doi.org/10.1093/scan/nsz037

Nolte, G. (2003). The magnetic lead field theorem in the quasi-static approximation and its use for magnetoencephalography forward calculation in realistic volume conductors. Physics in Medicine and Biology, 48(22), 3637–3652. https://doi.org/10.1088/0031-9155/48/22/002

Nugent, A. C., Andonegui, A. B., Holroyd, T., & Robinson, S. E. (2022). On-scalp magnetocorticography with optically pumped magnetometers: Simulated performance in resolving simultaneous sources. Neuroimage: Reports, 2(2), 100093. https://doi.org/10.1016/j.ynirp.2022.100093

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2010). Fieldtrip: Open source software for advanced analysis of meg, eeg, and invasive electrophysiological data. Computational Intelligence and Neuroscience, 2011, 1–9. https://doi.org/10.1155/2011/156869

Pajula, J., & Tohka, J. (2014). Effects of spatial smoothing on inter-subject correlation based analysis of fmri. Magnetic Resonance Imaging, 32(9), 1114–1124. https://doi.org/10.1016/J.MRI.2014.06.001

Park, H., Ince, R. A., Schyns, P. G., Thut, G., & Gross, J. (2018). Representational interactions during audiovisual speech entrainment: Redundancy in left posterior superior temporal gyrus and synergy in left motor cortex. PLoS Biology, 16(8), e2006558. https://doi.org/10.1371/journal.pbio.2006558

Proudfoot, M., Woolrich, M. W., Nobre, A. C., & Turner, M. R. (2014). Magnetoencephalography. Practical Neurology, 14(5), 336–343. https://doi.org/10.1136/practneurol-2013-000768

Rea, M., Boto, E., Holmes, N., Hill, R., Osborne, J., Rhodes, N., Leggett, J., Rier, L., Bowtell, R., Shah, V., & Brookes, M. J. (2022). A 90-channel triaxial magnetoencephalography system using optically pumped magnetometers. Annals of the New York Academy of Sciences, 1517(1), 107–124. https://doi.org/10.1111/nyas.14890

Rhodes, N., Rea, M., Boto, E., Rier, L., Shah, V., Hill, R. M., Osborne, J., Doyle, C., Holmes, N., Coleman, S. C., Mullinger, K., Bowtell, R., & Brookes, M. J. (2023). Measurement of frontal midline theta oscillations using opm-meg. NeuroImage, 271, 120024. https://doi.org/10.1016/j.neuroimage.2023.120024

Rier, L., Michelmann, S., Ritz, H., Shah, V., Hill, R. M., Osborne, J., Doyle, C., Holmes, N., Bowtell, R., Brookes, M. J., Norman, K. A., Hasson, U., Cohen, J. D., & Boto, E. (2023). Test-retest reliability of the human connectome: An opm-meg study. Imaging Neuroscience, 1, 1–20. https://doi.org/10.1162/imag_a_00020

Safar, K., Vandewouw, M. M., Sato, J., Devasagayam, J., Hill, R. M., Rea, M., Brookes, M. J., & Taylor, M. J. (2024). Using optically pumped magnetometers to replicate task-related responses in next generation magnetoencephalography. Scientific Reports, 14(1). https://doi.org/10.1038/s41598-024-56878-6

Schaworonkow, N., Blythe, D. A., Kegeles, J., Curio, G., & Nikulin, V. V. (2015). Power-law dynamics in neuronal and behavioral data introduce spurious correlations. Human Brain Mapping, 36(8), 2901–2914. https://doi.org/10.1002/hbm.22816

Schreiber, T., & Schmitz, A. (1996). Improved surrogate data for nonlinearity tests. Physical Review Letters, 77(4), 635–638. https://doi.org/10.1103/PhysRevLett.77.635

Stelzer, J., Chen, Y., & Turner, R. (2013). Statistical inference and multiple testing correction in classification-based multi-voxel

pattern analysis (mvpa): Random permutations and cluster size control. *NeuroImage*, *65*, 69–82. https://doi.org/10.1016/j.neuroimage.2012.09.063

Sun, T., Chi, X., Peng, Y., Zhang, Q., Liu, K., Ma, Y., Ding, M., Ji, N., & Zhang, Y. (2024). A first-in-human application of opm-meg for localizing motor activity area: Compared to functional mri. *NeuroImage*, *304*, 120953. https://doi.org/10.1016/j.neuroimage.2024.120953

Tierney, T. M., Holmes, N., Meyer, S. S., Boto, E., Roberts, G., Leggett, J., Buck, S., Duque-Muñoz, L., Litvak, V., Bestmann, S., Baldeweg, T., Bowtell, R., Brookes, M. J., & Barnes, G. R. (2018). Cognitive neuroscience using wearable magnetometer arrays: Noninvasive assessment of language function. *NeuroImage*, *181*, 513–520. https://doi.org/10.1016/J.NEUROIMAGE.2018.07.035

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B., & Joliot, M. (2002). Automated anatomical labeling of activations in spm using a macroscopic anatomical parcellation of the mni mri single-subject brain. *NeuroImage*, *15*(1), 273–289. https://doi.org/10.1006/nimg.2001.0978

Veen, B. V., Drongelen, W. V., Yuchtman, M., & Suzuki, A. (1997). Localization of brain electrical activity via linearly constrained minimum variance spatial filtering. *IEEE Transactions on Biomedical Engineering*, *44*(9), 867–880. https://doi.org/10.1109/10.623056

Wang, X., Teng, P., Meng, Q., Jiang, Y., Wu, J., Li, T., Wang, M., Guan, Y., Zhou, J., Sheng, J., Gao, J. H., & Luan, G. (2024). Performance of optically pumped magnetometer magnetoencephalography: Validation in large samples and multiple tasks. *Journal of Neural Engineering*, *21*(6). https://doi.org/10.1088/1741-2552/ad9680

Wernicke, C. (1974). *Der aphasische symptomenkomplex*. Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-642-65950-8_1

Woolrich, M. W., Ripley, B. D., Brady, M., & Smith, S. M. (2001). Temporal autocorrelation in univariate linear modeling of fmri data. *NeuroImage*, *14*(6), 1370–1386. https://doi.org/10.1006/nimg.2001.0931

Yang, A. I., Wang, X., Doyle, W. K., Halgren, E., Carlson, C., Belcher, T. L., Cash, S. S., Devinsky, O., & Thesen, T. (2012). Localization of dense intracranial electrode arrays using magnetic resonance imaging. *NeuroImage*, *63*(1), 157–165. https://doi.org/10.1016/j.neuroimage.2012.06.039

Ye, Z., Rüsseler, J., Gerth, I., & Münte, T. F. (2017). Audiovisual speech integration in the superior temporal region is dysfunctional in dyslexia. *Neuroscience*, *356*, 1–10. https://doi.org/10.1016/j.neuroscience.2017.05.017

Zarahn, E., Aguirre, G. K., & D'esposito, M. (1997). Empirical analyses of bold fmri statistics i. spatially unsmoothed data collected under null-hypothesis conditions. *NeuroImage*, *5*(3), 179–197. https://doi.org/10.1006/nimg.1997.0263

Zumer, J. M., Brookes, M. J., Stevenson, C. M., Francis, S. T., & Morris, P. G. (2010). Relating bold fmri and neural oscillations through convolution and optimal linear weighting. *NeuroImage*, *49*(2), 1479–1489. https://doi.org/10.1016/J.NEUROIMAGE.2009.09.020

Supplementary Material

Reliability and signal comparison of OPM-MEG, fMRI & iEEG in a repeated movie viewing paradigm.

Olivia R. Christiano & Sebastian Michelmann

Table S1: Comparison of within-subject maximum correlations after temporally down-sampling and averaging.

Modality	Frequency		Maxima			r (20 Hz)	r (0.67 Hz)	r (0.67 Hz, GA)
			z	MNI Coord.	AAL Label (Left/Right)			
OPM	δ	0.5–4 Hz	12.20	48, –16, 48	Precentral (R)	.06	.13	.33
	θ	4–8 Hz	9.17	–8, –104, 0	Occipital Mid (L)	.04	.14	.35
	α	8–12 Hz	11.98	8, –88, 40	Cuneus (R)	.05	.12	.33
	β	12–28 Hz	8.76	48, –40, 56	Parietal Inf (R)	.03	.14	.44
	γ_1	28–46 Hz	4.65	–48, –16, 56	Postcentral (L)	.02	.06	.14
	γ_2	55–70 Hz	5.04	64, –40, –24	Temporal Inf (R)	.02	.04	.22
	HF	64–116 Hz	4.70	–16, –24, 64	Paracentral Lobule (L)	.02	.06	.23
fMRI	–	–	13.43	–56, –16, 0	Temporal Mid (L)	.32	.28	.67
iEEG	δ	0.5–4 Hz	13.39	–58, –14, 48	Postcentral (L)	.29	.47	–
	θ	4–8 Hz	10.00	–72, –10, 7	SupraMarginal (R)	.22	.54	–
	α	8–12 Hz	10.33	–68, –11, 8	Temporal Sup (L)	.24	.40	–
	β	12–28 Hz	10.94	–67, –21, 16	SupraMarginal (L)	.11	.21	–
	γ_1	28–46 Hz	8.33	38, –82, –7	Occipital Inf (R)	.09	.39	–
	γ_2	55–70 Hz	8.86	–72, –38, 9	Temporal Mid (L)	.12	.40	–
	HF	64–116 Hz	23.16	–67, –21, 16	SupraMarginal (L)	.33	.76	–

Note. For each imaging modality and frequency band, we identify the maximum within-subject z-score (surviving multiple comparisons correction) and report correlation coefficients at that location. First, we report the untransformed correlation at the peak z-score location using data sampled at 20 Hz (r (20 Hz)), where correlations were computed between the first and second movie viewing for each subject and channel and averaged across subjects. Next, we report the corresponding value from the same location using data that were further down-sampled to 0.67 Hz (r (0.67 Hz)). Finally, for OPM and fMRI, we averaged the 0.67 Hz time series across subjects separately for viewing 1 and viewing 2, computed the correlation between the two grand-averaged signals, and report the resulting untransformed coefficient (r (0.67, GA)). For fMRI, analyses were conducted on data projected to the OPM source grid for consistency. r (0.67, GA) is not applicable to the single-subject iEEG data.

Within-viewing correlation analyses

For within-viewing comparisons, we followed the same procedures described in the main Methods section, except we correlated timeseries from the same viewing (e.g., subject i's first viewing with subject j's first viewing) rather than between them. Correlations were computed separately for viewing 1 and viewing 2, then averaged prior to generating surrogate distributions and computing z-scores.

Table S2: Between-subject reliability statistics for OPM and fMRI within movie viewings.

Modality	Frequency Band	% MC	M_z	SD_z	Maxima z	Maxima MNI Coord.	Maxima AAL Label (Left/Right)
OPM	δ 0.5–4 Hz	61.9%	4.0	1.7	10.28	–64, –8, 16	Postcentral (L)
	θ 4–8 Hz	94.6%	5.1	1.5	10.47	32, –80, –48	Crus II (R)
	α 8–12 Hz	87.4%	4.2	1.4	9.63	8, –88, –32	Crus II (R)
	β 12–28 Hz	99.9%	6.0	1.5	11.67	48, –64, 48	Angular (R)
	γ_1 28–46 Hz	9.9%	3.4	0.7	5.60	32, 0, –48	Fusiform (R)
	γ_2 55–70 Hz	2.4%	3.3	0.3	4.58	8, 72, 8	Frontal Sup Medial (R)
	HF 64–116 Hz	27.6%	2.9	0.5	4.48	–16, –72, –40	Cerebellum 7b (L)
fMRI	– –	26.7%	4.6	2.0	12.42	–48, –24, 8	Temporal Sup (L)

Note. All statistics are reported for data surviving correction for MC. For each modality and frequency band, this table reports the percentage of voxels/channels/electrodes surviving FDR correction (% MC), the M and SD of z-scores, and the peak z-score with its MNI coordinate and anatomical label.

Figure S1: Between-subject reliability for OPM and fMRI within movie viewings. FDR-corrected z-scores are shown for: (A) OPM, 3,581 source locations across 10 subjects for 7 frequency bands; (B) fMRI: 3581 voxels, across 11 subjects. (C) Schematic of between-subject correlation analysis. Color bars reflect z-scores (surviving correction for MC).

Table S3: Between-method reliability statistics across frequency bands for OPM, fMRI, and iEEG within movie viewings.

Comparison	Frequency Band		% MC	M_z	SD_z	z	MNI Coord.	AAL Label (Left/Right)
							Maxima	
OPM to fMRI	δ	0.5–4 Hz	0.7%	−3.6	2.3	−5.19	24, −8, 64	Frontal Sup (R)
	θ	4–8 Hz	11.3%	−2.1	3.7	−9.86	−32, −56, 64	Parietal Sup (L)
	α	8–12 Hz	6.3%	−3.9	1.9	−8.64	−40, −88, 8	Occipital Mid (L)
	β	12–28 Hz	11.5%	−3.9	2.5	−9.18	−48, −80, 8	Occipital Mid (L)
	γ_1	28–46 Hz	0.1%	−4.4	0.2	−4.54	−56, 24, 16	Frontal Inf Tri (L)
	γ_2	55–70 Hz	1.7%	3.5	1.9	5.32	−16, −96, 24	Occipital Sup (L)
	HF	64–116 Hz	0.03%	4.2	0.0	4.18	0, −88, 24	Cuneus (L)
OPM to iEEG	δ	0.5–4 Hz	0%	–	–	–	–	–
	θ	4–8 Hz	0.9%	3.7	0.4	4.60	−48, −54, −24	Temporal Inf (L)
	α	8–12 Hz	2.4%	3.3	1.8	4.76	−35, −91, 22	Occipital Mid (L)
	β	12–28 Hz	0%	–	–	–	–	–
	γ_1	28–46 Hz	0%	–	–	–	–	–
	γ_2	55–70 Hz	0%	–	–	–	–	–
	HF	64–116 Hz	0.5%	1.1	4.4	−3.91	−54, 19, −23	Temporal Pole Sup (L)
fMRI to iEEG	δ	0.5–4 Hz	6.3%	−3.5	2.2	−5.83	−11, −64, 75	Precuneus (L)
	θ	4–8 Hz	11.9%	−4.0	1.3	−8.36	−74, −28, 3	Temporal Mid (L)
	α	8–12 Hz	8.2%	−4.6	1.5	−8.79	−66, −8, 3	Temporal Sup (L)
	β	12–28 Hz	7.5%	−4.2	1.1	−7.05	−35, −91, 22	Occipital Mid (L)
	γ_1	28–46 Hz	3.2%	3.9	2.8	7.04	−66, −8, 3	Temporal Sup (L)
	γ_2	55–70 Hz	4.9%	4.3	2.4	8.62	−21, −55, 78	Parietal Sup (L)
	HF	64–116 Hz	9.1%	4.8	1.8	8.58	−74, −28, 3	Temporal Mid (L)

Note. All statistics are reported for data surviving correction for MC. For each comparison and band, we report the percent of voxels/electrodes surviving correction for MC (% MC), *M* and *SD* of *z*-scores, and the peak *z*-score with its MNI coordinate and anatomical label.

Figure S2: Consistent between-method alignment within movie viewings. (A) OPM to fMRI: *z*-scores from OPM subjects (n = 10) and fMRI subjects (n = 11) are shown for 3581 locations. (B) OPM to iEEG: *z*-scores from OPM subjects (n = 10) and single-subject iEEG from 656 electrodes pooled across five subjects. (C) fMRI to iEEG: *z*-scores from fMRI subjects (n = 11) and single-subject iEEG from 656 electrodes pooled across five subjects. Color bars reflect *z*-scores (see above).

Figure S3: Uncorrected cluster maps of SNR differences within and across modalities. All clusters shown exceed the 97.5th or fall below the 2.5th percentile of the empirical SNR distribution; values were not corrected for multiple comparisons and do not reflect statistical significance. (A) Within-modality SNR changes in OPM, computed by subtracting within-subject from between-subject *z*-scores at each channel. Negative values (blue) indicate greater signal consistency within individuals than across individuals. (B) Between-modality SNR changes comparing between-subject OPM to between-method fMRI, where negative values (red) indicate greater signal reliability within OPM than in cross-modal correspondence. SNR changes are shown across frequency bands from delta (0.5- 4 Hz) to high-frequency (64-116 Hz). Diagrams below illustrate the comparison logic for each panel.