From Colonisation to Invasion: Genomic and Phenotypic Comparison of

Faecal and Bloodstream Isolates from the same patients

Aakash Khanijau[1,2], Ellie Allman[1], Ralfh Pulmones[1], Richard N. Goodman[1], Rachel

McGalliard[2,3], Christopher M. Parry[1,3,4], Enitan D. Carrol[2,3], Adam P. Roberts[1]*.

1. Department of Tropical Disease Biology, Liverpool School of Tropical
   Medicine, Pembroke Place, Liverpool, L3 5QA

2. Institute of Infection, Veterinary and Ecological Sciences, University of
   Liverpool

3. Alder Hey Children's NHS Foundation Trust, Liverpool

4. Centre for Tropical Medicine and Global Health, Nuffield Department of
   Medicine, University of Oxford

*Corresponding author, Prof Adam P. Roberts. Liverpool School of Tropical Medicine,

Pembroke Place, Liverpool, L3 5QA. Adam.Roberts@lstmed.ac.uk

Abstract

Gram-negative bloodstream infections (GNBSI) carry a significant global health burden. *Escherichia coli* and *Klebsiella pneumoniae* are the two most common causes of healthcare-associated GNBSI, which may arise from gastrointestinal tract (GIT) colonisation. Understanding genomic and phenotypic adaptations that underpin transition from GIT colonisation to invasive bloodstream infection could improve understanding of pathogenesis.

This study identified 'linked' faecal and blood isolates from children with healthcare-associated GNBSI caused by *E. coli* and *K. pneumoniae*. Linked pairs were compared for antimicrobial resistance, biofilm formation, and underwent comparative genomic analysis via whole-genome sequencing, comparative average nucleotide identity (ANI) and core genome single nucleotide polymorphism (SNP) analysis.

Five isolate pairs (three *E. coli*, two *K. pneumoniae*) showed high relatedness, supporting GIT origin of bloodstream infection. Isolates within pairs had identical virulence genes whereas phenotypic assays revealed changes in antimicrobial susceptibility, with one pair undergoing changes in resistance gene profiles, and increased biofilm formation in 4/5 isolates.

This study provides insight into within-host evolution from gastrointestinal colonisation to bloodstream invasion in Gram-negative pathogens. Convergence on metabolic adaptation and biofilm formation suggests these traits may be advantageous in healthcare-associated GNBSI. Further studies involving larger cohorts alongside functional validation of mutations are needed to better understand GNBSI pathogenesis.

Introduction

Gram-negative bloodstream infections (GNBSI) contribute significantly to the global burden of paediatric sepsis, accounting for 30-50% of all bloodstream infections worldwide [1, 2]. Healthcare-associated bloodstream infections, such as central-line associated bloodstream infections (CLABSI), often affect neonates or other critically ill children, and are associated with significant mortality. [3].

The two most clinically significant organisms implicated in GNBSI are *Klebsiella pneumoniae* and *Escherichia coli* [4, 5]. The emergence of antimicrobial resistance (AMR) in these pathogens, presents a significant challenge. Particularly, extended-spectrum beta-lactamase (ESBL)-producing isolates, which exhibit resistance to third generation cephalosporins and beta-lactam/beta-lactamase inhibitor combinations. As such, the World Health Organization has designated ESBL-producing *K. pneumoniae* and *E. coli* as priority pathogens for which new antibiotics are required [6].

Both *K. pneumoniae* and *E. coli* are persistent colonisers of the human gastrointestinal tract (GIT) which is believed to be an important reservoir for subsequent GNBSI [7]. The transition from GIT colonisation to GNBSI can occur via invasion of other body sites or medical devices, leading to secondary bloodstream infection, or direct translocation across the gut mucosa causing primary bloodstream infection. The latter is implicated in 20-50% of cases of healthcare associated infections [7].

The transition from GIT colonisation to bloodstream invasion depends on host, clinical, and environmental influences, combined with specific pathogen characteristics. Host factors that can predispose to GIT-bloodstream transition

include immunosuppression, critical illness, and antimicrobial exposure [7, 8].

Pathogen factors include advantageous phenotypes such as AMR, bloodstream

fitness and biofilm formation; which may allow invasion, survival and dissemination in

the bloodstream. Biofilm formation is of particular interest in the context of CLABSI,

as growth within a biofilm provides protection against host defence and antimicrobial

agents, and may act as a nidus for ongoing bloodstream dissemination.

Many antimicrobial resistance genes (ARGs) and virulence genes underpinning AMR

and invasive infection respectively are well characterised. However, predisposing

genotypes and specific genetic changes undergone within-host that may underpin

GIT-blood transition are seldom studied, perhaps due to the scarcity of such paired

isolates from the same patient. This includes genomic adaptations like acquisition or

loss of virulence genes and plasmids, and regulatory or metabolic alterations.

Furthermore, the relationship between many genomic adaptations and

advantageous phenotypic changes remains unclear, as well as the relationship

between AMR and invasive infection. Combined phenotypic and genomic

comparison of faecal and bloodstream isolates, obtained from the same patient,

could improve understanding of intra-host evolution involved in the pathogenesis of

GNBSI, and translate to informing intervention strategies.

In this study we aimed to identify genomically 'linked' pairs of faecal and blood Gram-

negative isolates, which were likely to have undergone GIT-blood transition and

evolution within patients to cause GNBSI. For each 'linked' pair, we investigated if

there had been any loss or acquisition of virulence genes, ARGs, and plasmids

between faecal isolates and their blood counterpart. We then investigated if there

were any difference in two phenotypes of interest: AMR and biofilm formation.

Finally, we used a comparative genomic approach to identify further genomic

changes that could explain any observed phenotypic differences or otherwise

contribute to the transition from GIT colonisation to bloodstream invasion.

Materials and Methods

Clinical isolates, clinical data collection and ethics

A library of 81 stored bacterial isolates from Alder Hey Children's Hospital, Liverpool,

UK, was interrogated for pairs of ESBL-producing *E. coli* and *K. pneumoniae* isolates

from the same patient, where the same pathogen had been identified on a routine

faecal surveillance culture and in a subsequent blood culture. Isolates were stored in

LB (Luria-Bertani) Broth (Sigma-Aldrich) and 20% glycerol at -80°C. Anonymised

clinical information was obtained from patient records, including: demographics,

source of GNBSI, and antimicrobial exposure between isolate dates. Ethical

approval for linking anonymised data was granted by the University of Liverpool as

part of a separate study for which the isolate library was created. (REC reference

255669).

Whole Genome Sequencing and comparative genomics

Short-read Whole Genome Sequencing (WGS) and assembly was performed at the

Centre of Genomic Research (University of Liverpool) using the Illumina NovoSeq

SP (2x150 bp). Assembled genomes were analysed for the following characteristics,

which were compared between faecal and blood isolates of each patient pair. Core

genome multi-locus sequence typing (cgMLST) was performed using Kleborate [9]

for *K. pneumoniae* isolates and using Institut de Pasteur database

(https://bigsdb.pasteur.fr/about/) for *E. coli* isolates. Kaptive [10] was used to identify

wzi-type, K-type and O-type of *K. pneumoniae* isolates, and SeroTyper 2.0 and

CHtype to determine the H-type, O-type and FumC- type of *E. coli* isolates.

To assess relatedness further, average Nucleotide Identity (ANI) was then compared

between isolates using OrthoANI [11] Following this, the number of core genome

single nucleotide polymorphisms (SNPs) between faecal and blood isolates of each

pair was calculated. SNPs were predicted using Snippy v.4.3.6

(https://github.com/tseemann/snippy) by aligning raw reads of assemblies to the

reference genomes MG1655 and HS11286, for *E. coli* and *K.*

pneumoniae respectively. Core genome SNPs were extracted with Snippy-core and

intra-species SNPs were plotted and counted in R v.4.3.1 [12]. ResFinder 4.2.2 [13]

was used to determine the presence of ARGs and provide a genotypic prediction of

resistance for each isolate. VF analyzer (http://www.mgc.ac.cn/VFs/) was used to

identify virulence genes in each isolate. PlasmidFinder [14] was used to identify the

presence of plasmids.

Isolate genomes were annotated using RAST annotation tool [20] and pairs

underwent comparative genomic analysis using Breseq version 0.38.3 [21] using

default parameters. Mutations between blood isolates and the putative counterpart

faecal ancestor were categorized as insertions/deletions (indels), non-synonymous

SNPs (nSNPs), synonymous SNPs (sSNPs), or intergenic SNPs (iSNPs). Mutations

within or <100bp upstream of a gene were classified into COG functional categories

using EGGNOG 5.0. Mutations involving hypothetical proteins were classified as

'Function unknown' and mutations without genes within 100bp were 'Unclassified.'

Any mutations relating to virulence genes, ARGs, or previously described

bloodstream survival factors were identified. In addition, mutations in genes

previously reported to affect antimicrobial resistance or biofilm formation were

identified.

Data access.

The bioproject number for this study is PRJNA1295786 which contains the 10

accession numbers for the individual genomes SRX29836134 - SRX29836143.

Antimicrobial Susceptibility Testing (AST):

Six antimicrobials from the contemporaneous local prescribing guidance for Gram-

negative infections were used for AST: Cefotaxime, Gentamicin, Ciprofloxacin,

Amoxicillin-Clavulanic acid (Co-amoxiclav), Piperacillin-Tazobactam and

Meropenem. Kirby-Bauer disc diffusion testing was used for initial AST following

EUCAST guidelines [15]. Zone sizes were interpreted as per the EUCAST clinical

breakpoints, to give a phenotype of Sensitive (S), resistant (R) or in the 'Area of

Technical Uncertainty' (ATU) [16]. Three technical and three biological repeats were

performed, and the mode was recorded as the phenotype for each isolate, and

compared within each pair. In addition, the mean zone size, along with standard

error of the mean (SEM) was recorded for each isolate and Mann-Whitney U test

was used to compare differences between faecal and blood isolates for each patient

pair. Any identified differences between faecal and blood isolates in each patient pair

were confirmed using broth microdilution to quantify the fold-change in minimum

inhibitory concentration (MIC). Broth microdilution was performed following EUCAST

guidelines [17] and with three biological and three technical replicates.

Biofilm formation

Crystal violet assay was performed derived from two previously published protocols

[18, 19]. Isolates were incubated at 37°C on LB agar for 18 h. A single colony of each

isolate was transferred to 10ml Mueller-Hinton broth 2 (MHB2; Sigma-Aldrich) and

incubated at 37°C at 200 rpm for a further 18 h. Cultures were diluted to an OD_{600} of

0.01 in fresh MHB2. 150µl of each isolate was transferred in a U-bottom plate, with

150µl sterile MHB2 used as a negative control. After 24-hour incubation, the biomass

was transferred to a flat-well plate, which was used to measure OD_{600}. The

remaining biofilm in the U-bottom plate wells was stained with 1% crystal violet and

subsequently solubilized with 30% acetic acid. The final OD_{600} was measured in a

flat-well plate. This value was then normalized to the initial OD_{600} of the biomass, to

quantify biofilm formation. The assay was performed as three biological replicates

and four technical replicates. Mean OD_{600} of blood and faecal isolates were

compared using a paired t-test, and SEM calculated.

Statistical analysis and data visualization

All statistical analysis and data visualization was performed using ggplot2 in R

version 4.3.2 [12].

Results

Clinical isolates and clinical context of GNBSI

Five pairs of isolates, two *K. pneumoniae* and three *E. coli,* were identified for use in

this study (Table 1). Time between faecal (1st sample) and blood sample isolation

ranged from 17-144 days. Patients from whom isolates were derived were aged 1

month to 12 years. All five cases of GNBSI were diagnosed as CLABSI, and 4/5

cases occurred on ICU. Patients received between two to six different antimicrobials

in-between faecal and blood sample isolation.

Genomic relatedness of faecal-blood isolate pairs

To initially assess relatedness between faecal and blood isolates, cgMLST was

performed. In 5/5 pairs, faecal and blood isolates shared the same sequence type

(Table 2.) In Pair 1, *K. pneumoniae* isolates were from the sequence type ST922,

and in Pair 2 *K. pneumoniae* isolates were from the sequence type ST461 (Table

2A). In Pair 3 and 4, *E. coli* isolates were from the globally dominant sequence type

ST131, whereas in Pair 5 isolates were from sequence type 12 (Table 2B).

K. pneumoniae isolates were then compared for K-type, O-type and wzi-type, which

were concordant between faecal and blood isolates in both pairs. *E. coli* isolates

were compared or O-type, H-type and FumC -type, and again, all were concordant

within each pair.

Next, relatedness was further assessed using ANI (Figure 1A.) In 4/5 pairs (2, 3, 4,

5) the faecal-blood isolates shared an ANI of 99.99%, whereas isolates in Pair 1

shared an ANI of 99.82%. Within a comparison between all isolates across all patient

pairs, each faecal isolate was most closely related to its corresponding blood isolate

from the same patient. Given these findings, all pairs were at this point considered to

be genomically 'linked' and included for further study.

Isolates were next compared for core genome SNP differences (Figure 1B.) For *K.*

pneumoniae isolates, Pair 2 exhibited the lowest SNP difference (38 SNPs), while

Pair 1 showed the highest (318 SNPs). In *E. coli* isolates, within-pair SNP distances

ranged from 40 SNPs in Pair 4 to 60 SNPs in Pair 3. These findings were in

alignment with the ANI comparison.

Initial genomic comparison: ARGs, virulence genes and plasmids

At this point, all pairs were considered to be 'linked' and an initial genomic

comparison of ARGs, virulence genes and plasmids of the faecal and blood isolates

within each pair was performed (Figure 2). There were between 1-13 ARGs detected

in each isolate, predicted to confer resistance to various drug-classes (Figure 2A).

The presence of at least one ESBL-gene was confirmed in each isolate. ARG

profiles were identical within pairs, except for Pair 4, in which *bla*TEM-1B was

detected in the blood isolate but not the faecal counterpart, suggesting acquisition

between the dates of sampling, possibly during GIT-blood translocation.

There were between 51-86 virulence genes detected in each isolate, encoding

multiple classes of virulence factors (Figure 2B/C). Virulence gene presence was

identical between faecal and blood isolates, indicating no loss or acquisition of

virulence genes in GIT-blood transition in any pair.

There were between 3-8 plasmids detected in each isolate (Figure 2D). Whilst the

majority of pairs demonstrated identical plasmid profiles, indicating stable plasmid

carriage during the likely GIT-blood transition, Pair 1 demonstrated loss of IncFIB(K)

from the faecal isolate and gain of IncN and IncFII(Yp) plasmids in the blood isolate.

Phenotypic comparison: Antimicrobial resistance

Disc diffusion testing was used to screen for a change in antimicrobial resistance

between faecal and blood isolates of each pair. A change in resistance phenotype

was identified for three drug-bug combinations in 2/5 pairs (Supplementary figure 1).

There was also a statistically significant difference in zone size between faecal and

blood isolates that did not cross the EUCAST clinical breakpoint for four drug-isolate

combinations in 2/5 pairs. (Supplementary figure 2, Supplementary table 1).

These within-pair differences were only confirmed by Broth microdilution in two pairs.

Pair 2 demonstrated an increase in MIC for Ciprofloxacin, Piperacillin-Tazobactam,

Cefotaxime and Gentamicin in the blood isolate compared to its faecal counterpart,

indicating increased resistance development to these four antimicrobials between

the two sampling timepoints. (Figure 3A). In Pair 3, MIC was decreased for

Piperacillin-Tazobactam in the blood isolate compared to faecal, indicating increased

susceptibility development in GIT-blood transition. (Figure 3B).

Phenotypic comparison: Biofilm formation

Because all patients were found to be diagnosed with CLABSI, and the importance

of biofilm formation in CLABSI pathogenesis, we investigated for any difference in

biofilm formation between faecal and blood isolates in each pair. In 4/5 pairs, blood

isolates demonstrated significantly increased biofilm formation in comparison to the

faecal counterpart, as determined by a paired t-test. (Figure 4). All p-values are

stated in Supplementary table 2. Overall, the isolates in Pair 2 demonstrated the

strongest biofilm formation.

Comparative genomics: Breseq analysis

Breseq was used to identify mutations in each pair. Most mutations were Indels, and

the majority of these were 'Function unknown' or 'Unclassified.' (Supplementary

figure 3) Mutations occurring within 100bp of genes with a known function were

assigned to COG categories (Figure 5). There were six categories in which

mutations occurred in all pairs. Of these, most occurred in 'L: Replication,

recombination and repair,' 'E: Amino acid transport and metabolism,' and 'G:

Carbohydrate transport and metabolism.'

No mutations were identified in specific ARGs identified by ResFinder, however four

pairs demonstrated mutations in genes encoding efflux pumps and transporter

proteins involved in AMR (Table 3). Four pairs demonstrated mutations in at least

one known virulence factor gene. Two pairs demonstrated mutations in or related to

previously reported bloodstream survival factors. All pairs demonstrated a variety of

mutations in genes related to biofilm formation. In Pair 1, there was evidence of

acquisition of another copy of the *hha* gene, along with modification of the Shine-

Delgarno intergenic region (Figure 6)

Discussion

In this study, we compared blood isolates with the previously acquired faecal isolates

from 5 children with GNBSI, and in each instance demonstrated concordance in

sequence type and species-specific typing methods. This is highly suggestive that

the faecal-blood isolate pairs were genomically 'linked' and that the invasive blood

isolate was likely to have been acquired from the GIT. Average nucleotide identity

has been used as a marker of genomic relatedness in previous studies comparing

faecal-blood isolates in children [22, 23] but there is no defined consensus on a

threshold that indicates isolates as similar enough to be linked. Thresholds of

99.99% have been previously used to indicate isolates being the same strain [24]

Using this threshold, 4/5 pairs (Pairs 2-5) shared ANI values convincing of being the

same strain. The lower ANI in Pair 1 (99.82%) could be somewhat explained by the

identified plasmid changes in this pair. This is consistent with observations that MGE

movement can lead to a lower ANI in isolates that are highly likely to be concordant,

as noted by Vornhagen *et al*. when comparing wzi-type concordant rectal and blood

K. pneumoniae isolates [25]. However, this finding could also be indicative of the

blood isolate having evolved to be a different clade of the same sequence type as

the faecal isolate in this pair, as indicated by the higher number of core genome

SNPs calculated in further analysis.

The number of core genome SNPs between faecal and blood isolates in each pair

ranged from 38 SNPs (Pair 2) to 318 SNPs (Pair 1), indicating varying degrees of

evolutionary distance between faecal and blood isolates in each patient. It is possible

that our BSI isolates, although related, are not immediate descendants of the

cognate faecal isolate and that there could have been closer ancestors in the GIT

which were not picked up during sampling. Similarly, there may have been evolution

within the blood giving rise to multiple descendants, of which we isolated one. A

previous study that identified concordant faecal and blood isolates in patients with *E.*

coli bloodstream infections noted that the number of core genome SNPs between

faecal and blood isolates was 0 [26]. Another study comparing faecal and blood *K.*

pneumoniae isolates found a core genome SNP difference of 6 SNPs between

isolates considered to be concordant strains in one patient and 76 SNPs between

concordant isolates from another patient. [27]. In our study, the number of SNPs we

found is generally higher, and was not proportional to the number of days between

faecal and blood isolation. This may represent varied and convoluted routes taken

from the GIT to the blood e.g. via other body tissues in each case of GNBSI,

requiring adaptation to multiple niches – as opposed to direct GIT-blood translocation

Alternatively, the higher number of core genome SNPs in Pair 1 could be considered

to contradict the hypothesis that this represents a 'linked' pair. However, as with ANI,

there is no SNP threshold that is universally agreed as defining isolates as 'linked'.

For example, whilst one study used a threshold of 17 core genome SNPs for this,

another study looking at persistence of *E. coli* isolates between GI and urinary tracts

used a threshold of ≤500 core genome SNPs [26, 28]. These differences could also

be due to differences in sequencing and bioinformatic technologies, as well as the

reference genomes used in different studies. Such differences in methodology are

recognised as a key barrier to the determination of a defined threshold for strains

similarity [29-31]. In addition, the expected rate of mutation of isolates moving

between different body niches is currently unknown and unpredictable [32], adding to

this challenge. When comparing isolates from the two different patients with *K.*

pneumoniae infection, there are over 30,000 core genome SNPS between each. A

similar number is seen when comparing the *E. coli* isolates of Patient 5 to both

Patient 3 and Patient 4. However, the number of core genome SNPs between

Patient 3 and 4 are more comparable to those seen within the pair of Patient 1. This

raises suspicion for a common source of colonisation of these patients within the

intensive care unit during the time period the isolates were taken. However granular

data of overlapping hospital admission times is unavailable.

ARG profile and subsequently AMR phenotypes were compared within 'linked' pairs.

The transition from GIT colonisation to blood invasion was associated with

acquisition of an ARG in 1/5 pairs, (Pair 4) - and in this case, the acquisition of

*bla*TEM-1B did not lead to increased resistance to any of the beta-lactam/inhibitor

combinations tested. Only in one pair, (Pair 2) did we observe that increases in

resistance to any antimicrobials occurred between the two sample timepoints. This is

despite that in all five cases, the patient from whom isolates were obtained received

multiple antimicrobials during hospital admission. In contrast, for Pair 3, increased

susceptibility to Piperacillin-tazobactam was noted. These findings may reflect

varying selection pressure exerted by dose and duration of antimicrobial therapy, in

balance with environmental adaptation. Unfortunately, the study is limited by lack of

granular data on antimicrobial exposure.

These phenotypic changes in AMR are not explained by acquisition or loss of ARGs

(Figure 2A). However, by using Breseq for whole genome comparison, we identified

deletion of drug-efflux pump repressors that could putatively explain this observation.

Several drug-efflux pumps are conserved across Gram-negative species and

implicated in AMR. These include MFS-pumps, RND-pumps, and ABC-transporters.

The *acrAB-tolC* efflux pump is an RND-pump implicated in resistance to multiple

antimicrobials. [33]. It has multiple regulatory pathways, including the *acrRAB*

operon, in which *acrR* is a repressor, and the *marAB* operon, with *marB* as a

repressor. In Pair 2, *marB* and *acrR* deletions were identified (Table 3), which could

lead to increased efflux pump activity, explaining the changes in susceptibility. The

deletion of *acrR* is particularly interesting because a previous study reported that

acrR knockout in *K. pneumoniae* isolates resulted in increased MIC to the same

antimicrobial classes as seen here [34]. Previous literature also reports an

association between *acrR* disruption with a MDR phenotype [35, 36]. However, it

must be noted that the *acrR* deletion was identified in Pair 5 but without a change in

susceptibility, possibly explained by the variation in other efflux pump changes seen

in each pair.

In 5/5 cases in this study, GIT-blood transition was not associated with the

acquisition of virulence genes when comparing blood isolates to the presumed faecal

ancestor. Because all cases of GNBSI in this study were cases of CLABSI, biofilm

formation was chosen as a virulence phenotype of interest. The finding that 4/5

blood isolates exhibited an increase in biofilm formation compared to faecal

counterparts, suggests in-host adaptation towards increased biofilm formation may

be implicated in the pathogenesis. As noted, the observed phenotypic changes are

not explained by acquisition of virulence genes involved in biofilm formation, but

comparative genomic analysis using Breseq again identified a variety of putative

mechanisms that could potentially underpin our phenotypic observation.

In Pair 2, the *acrR* deletion explored above could also be responsible for increased

biofilm formation, as *acrAB-tolC* has a central role in biofilm formation in *E. coli* and

K. pneumoniae. [37, 38]. Studies also link biofilm formation, AMR and *acrAB-tolC*

activity. In a comparison of *K. pneumoniae* urinary isolates, Vuotto *et al.* observed

that those with the highest biofilm forming-capacity also demonstrated the most

AMR, correlating with increased *acrAB* activity [39]. Although only a sample of three

isolates, this strikingly mirrors the findings of the *K. pneumoniae* blood isolate of Pair

2, raising the possibility of a single intra-host adaptation contributing to both

antimicrobial resistance and biofilm formation. Hennequin *et al* demonstrated this *in*

vitro, reporting that sub-MIC cephalosporin exposure in *E. coli* isolates resulted in

enhanced *acrAB* expression and biofilm formation [40]. This could be an example of

antimicrobial exposure selecting for a resistance-conferring mutation that also

increases virulence and propensity to invasive infection. If seen consistently in a

larger cohort, this could suggest a promising interventional target. For example,

RND-efflux pump inhibitors have been explored for use as antimicrobial-adjuvant

therapy for resistant infections, and for biofilm disruption in invasive infection [41].

Furthermore, in this pair, nsSNPs in two biofilm-related virulence genes, *lcmF* and

Protease III precursor were identified, possibly contributing to the phenotypic

change. Similarly, in Pair 3, two nsSNPs in the biofilm-associated virulence gene

'antigen 43' were noted. Mutations in this gene were noted when Nielson *et al*

compared 'linked' urine and faecal isolates, and hypothesised this related to

increased biofilm formation [42].

Pair 1 exhibited a potential link between biofilm formation and plasmid acquisition.

This pair underwent loss of IncFIB(K) and acquisition of IncFII(Yp) between faecal

and blood isolates (Figure 2D), Breseq analysis revealed a duplication of the *hha*

gene in the blood isolate compared to the faecal isolate (Table 3), with the additional

copy residing on a contig containing multiple IncF-related genes, suggesting

plasmid-mediated acquisition. The *hha* gene encodes haemolysin-expression-

modulating protein and is a regulator of virulence factors involved in cell adhesion

and biofilm formation in Gram-negative species. Studies in *E. coli* have identified *hha*

as a positive regulator of biofilm formation and of virulence [43, 44]. This is less

studied in *K. pneumoniae*, however Bandeira *et al* described *hha* presence to be

associated with *K. pneumoniae* biofilm-forming isolates, particularly on indwelling

central lines [45]. Plasmid-mediated acquisition of *hha* was described by Krall *et al*

who compared 'linked' faecal and blood isolates finding that *hha* acquisition on an

IncFII plasmid increased the invasiveness of *E. coli* isolates in a gut-organoid model

[46]. They concluded *hha* was involved in GIT-blood adaptation but did not explore

biofilm formation specifically. Here, one could similarly hypothesise that IncFII

plasmid-mediated acquisition of *hha* could be an adaptive change leading to

increased biofilm formation and procession to GNBSI.

Other studies report *hha* has a negative influence on biofilm formation [47, 48] which

this hypothesis contradicts. Interestingly, in both copies of the *hha* gene in the blood

isolate, a 3bp GGT insertion was noted 1bp downstream of *hha*. Upon gene

alignment, this appeared to be an AAA duplication, disrupting the Shine-Dalgarno

sequence 7bp before the gene (Figure 6). This could potentially reduce the efficiency

of protein translation. Hence, one could equally hypothesize, assuming a negative-

regulatory role of *hha*, that after plasmid acquisition during GIT colonisation, the

compensatory mutation in the Shine-Dalgarno region led to increased biofilm

formation and aided bloodstream invasion. Further investigation is required to clarify

the role of *hha* in invasion and the combined effect of the observed SNPs, Shine-

Dalgarno mutation, and *hha* copy number on biofilm formation.

The transition from GIT-blood, and subsequent survival in the bloodstream will

involve the ability to adapt to a harsh and dynamic environment, with factors such as

varying nutrient abundance and host immunity to contend with. As noted, this

adaptation did not appear to result from acquisition of virulence genes in these five

cases, but was associated with varied changes across the genome. No single

specific mutation occurred in all pairs, likely reflecting the differing clinical scenarios

and selection pressures driving diverse evolutionary trajectories en-route to

bloodstream infection. However, in all pairs, most mutations occurred in the COG

category 'Replication, recombination and repair', including a mutation in *dam* a

known bloodstream survival factor, in Pair 1 [49]. These genes are involved in

environmental and stress response and selection for mutations could be expected in

adaptation to challenging environments, for example maintaining efficient DNA repair

mechanisms in the presence of reactive oxygen species, host immune defences or

nutrient variability. Another commonality was mutations in 'cell

wall/membrane/envelope biogenesis,' 'cell motility,' and 'signal transduction'

categories. Such mutations might confer altered chemotaxis or adherence, allowing

tissue invasion. Few mutations were observed in 'Cell cycle control and repair' and

'Translation, ribosomal structure and biogenesis.' This demonstrates the strong

selective pressure to conserve key cellular processes integral to bacterial survival

during the transition between GIT and blood environments. Many mutations were

classified as 'Function Unknown' within COG categories but may hold significance.

For example, in Pair 3, a non-synonymous SNP was identified in a phage

endopeptidase gene adjacent to the gene *iss.* The *iss* gene has is associated with

bloodstream survival in *E. coli* through mediating serum and complement resistance,

and increased mRNA copy number has been associated with increased serum

survival [50, 51]. Mutations in the adjacent phage endopeptidase could influence the

activation or expression of *iss*, with effects on bloodstream survival. This represents

a potential avenue for further research.

When comparing blood isolates to their faecal ancestors using Breseq, we noted

mutations related to carbohydrate and amino acid transport in 5/5 pairs. This

included an nsSNP in *ompk36* in Pair 1, a transport protein also previously identified

as a bloodstream survival factor in *Klebsiella pneumoniae* [52]. Further, we identified

mutations related to iron uptake utilisation in 4/5 pairs. Iron availability and chemical

form varies in different tissues and a 'functional hierarchy' whereby iron uptake

systems are utilised variably across environments has been discussed by Garcia *et*

al in the context of UTI [53]. The findings of *fec* operon deletion in Pair 1 and the

transposase mutation near the *fec* operon in Pair 3 may suggest a genomic shift

away from ferric citrate uptake, whilst the *sitC* virulence gene mutation and ferrous

(Fe^{2+}) ABC transporter deletion in Pair 5 may suggest a shift away from Fe^{2+} uptake.

These mutations likely reflect adaptation towards iron-acquisition in the bloodstream,

where ferric citrate is less abundant than the gut and Fe^{2+} is less abundant due to

oxygenated conditions. Further exploration could inform therapeutic strategies. For

example, *tonB* which regulates the *fec* operon has been explored as a potential drug

target against *E. coli* [54]. However, this may not be an appropriate target in the

context of bloodstream infection if functional redundancy of the *fec* operon is a

common adaptation.

The strengths of this study include the combination of genomic with phenotypic

comparison, using isolates from the same patient to gain insight into within-host

evolution and link phenotypic differences to changes at the genomic level. To our

knowledge, there are few studies in the existing literature that have investigated this.

Further, we compared mutations across the whole genome, rather than focusing only

on known ARGs and virulence genes identifying potentially relevant mutations that

would have been missed by a narrower approach. Limitations of our study lie in the

fact that it cannot be definitively concluded that these pairs are 'linked' given the lack

of an existing robustly defined threshold for strain similarity/genomic relatedness -

this in itself highlighting an area for further work. In addition, the retrospective nature

and ability to only include five cases of GNBSI, without granular clinical data,

meaning the significance and generalisability of these findings cannot be assumed.

The use of short-read rather than long-read sequencing makes it harder to discern

plasmid-mediated changes that could contribute to adaptation. Further, phenotypic

assays for biofilm formation demonstrated changes *in vitro*, but likely do not reflect

true clinical conditions.

Whilst this study shows individual instances of mutations of potential clinical

relevance in each pair which may represent avenues for further study; it is clear that

there is a diverse range of possible evolutionary trajectories in GIT-blood transition,

even in this small study, so a common pathway for clinical intervention may be

difficult to elucidate. Future study with a larger sample size that used faecal isolates

from proven 'linked' pairs to colonising isolates that did not achieve bloodstream

invasion, could be a more promising avenue to identify pathogen factors e.g

virulence gene SNPs that may predispose to bloodstream infection. Previous case-

control studies with this approach have been used, but not including WGS data [55].

Analysis of isolates from other body-site isolates would be beneficial, along with

information around antimicrobial exposure, clinical severity, and GNBSI source,

giving a more comprehensive insight into the relevance of adaptations to differing

clinical scenarios, and how findings could translate to clinical intervention.

Overall, this study used genomic and phenotypic comparison of 'linked' pairs of

blood and faecal *K. pneumoniae* and *E. coli* isolates to investigate evolutionary and

adaptive changes relevant to the transition from GIT colonisation to bloodstream

invasion and GNBSI. Phenotypic changes in AMR were variable, however biofilm

formation and metabolic flexibility were common phenotypic adaptations towards

bloodstream infection in this small sample. These changes were not found to be

associated with acquisition of ARGs or virulence genes, but rather, with a diverse

array of genomic changes, including mutation of virulence genes, efflux pumps and

acquisition of plasmid-mediated genes. No single mutation occurred in all pairs,

however changes in functional categories of genes involved in transport and

metabolism were common. These findings demonstrate a snapshot of the vast

evolutionary landscape of intra-host adaptation that is traversed in GNBSI

pathogenesis. However, significant further study is needed to assess whether these

findings could feasibly translate into clinical intervention for GNBSI treatment or

prevention.

Acknowledgements

APR acknowledges funding from the Medical Research Council, Biotechnology and

Biological Sciences Research Council and Natural Environmental Research Council

which are all Councils of UK Research and Innovation (grant no. MR/W030578/1)

under the umbrella of the JPIAMR (Joint Programming Initiative on Antimicrobial

Resistance), and UKRI through the Strength in Places Fund (grant no. SIPF 36348).

EA and RP are by the MRC via the LSTM and Lancaster University PhD Doctoral

Training Program (Grant no. MR/N013514/1)

References

1.   Hallmaier-Wacker, L.K., et al., *Incidence and aetiology of infant Gram-negative*
bacteraemia and meningitis: systematic review and meta-analysis. Arch Dis Child,
2022. **107**(11): p. 988-94.
2.   Rudd, K.E., et al., *Global, regional, and national sepsis incidence and mortality,*
1990-2017: analysis for the Global Burden of Disease Study. Lancet, 2020.
395(10219): p. 200-211.
3.   Wagner, M., et al., *Prospective study on central venous line associated bloodstream*
infections. Arch Dis Child, 2011. **96**(9): p. 827-31.
4.   Kern, W.V. and S. Rieg, *Burden of bacterial bloodstream infection-a brief update on*
epidemiology and significance of multidrug-resistant pathogens. Clin Microbiol Infect,
2020. **26**(2): p. 151-157.
5.   Lipworth, S., et al., *Molecular epidemiology and antimicrobial resistance phenotype*
of paediatric bloodstream infections caused by Gram-negative bacteria. Commun
Med (Lond), 2022. **2**: p. 101.

6.   WHO, *Bacterial pathogens of public health importance to guide research,*
546   *development and strategies to prevent and control antimicrobial resistance.* 2024.
547   7.   Holmes, C.L., et al., *Pathogenesis of Gram-Negative Bacteremia.* Clin Microbiol Rev,
548   2021. **34**(2).
549   8.   Jimenez-Rojas, V., et al., *Gut colonization and subsequent infection of neonates*
550   *caused by extended-spectrum beta-lactamase-producing Escherichia coli and*
551   *Klebsiella pneumoniae.* Front Cell Infect Microbiol, 2023. **13**: p. 1322874.
552   9.   Lam, M.M.C., et al., *A genomic surveillance framework and genotyping tool for*
553   *Klebsiella pneumoniae and its related species complex.* Nat Commun, 2021. **12**(1):
554   p. 4188.
555   10.   Lam, M.M.C., et al., *Kaptive 2.0: updated capsule and lipopolysaccharide locus*
556   *typing for the Klebsiella pneumoniae species complex.* Microb Genom, 2022. **8**(3).
557   11.   Yoon, S.H., et al., *A large-scale evaluation of algorithms to calculate average*
558   *nucleotide identity.* Antonie Van Leeuwenhoek, 2017. **110**(10): p. 1281-1286.
559   12.   R Foundation for Statistical Computing, V., Austria, *R Core Team: R: A language and*
560   *environment for statistical computing.* 2021.
561   13.   Florensa, A.F., et al., *ResFinder - an open online resource for identification of*
562   *antimicrobial resistance genes in next-generation sequencing data and prediction of*
563   *phenotypes from genotypes.* Microb Genom, 2022. **8**(1).
564   14.   Carattoli, A., et al., *In silico detection and typing of plasmids using PlasmidFinder and*
565   *plasmid multilocus sequence typing.* Antimicrob Agents Chemother, 2014. **58**(7): p.
566   3895-903.
567   15.   EUCAST, *Disk diffusion manual version 12.0.* 2024.
568   16.   EUCAST, *Breakpoint tables for interpretation of MICs and zone diameters, version*
569   *14.0* 2024.
570   17.   EUCAST, *Broth microdilution - EUCAST reading guide, version 5.0.* 2024.
571   18.   Merritt, J.H., D.E. Kadouri, and G.A. O'Toole, *Growing and analyzing static biofilms.*
572   Curr Protoc Microbiol, 2005. **Chapter 1**: p. Unit 1B 1.
573   19.   Au - O'Toole, G.A., *Microtiter Dish Biofilm Formation Assay.* JoVE, 2011(47): p.
574   e2437.
575   20.   Aziz, R.K., et al., *The RAST Server: rapid annotations using subsystems technology.*
576   BMC Genomics, 2008. **9**: p. 75.
577   21.   Deatherage, D.E. and J.E. Barrick, *Identification of mutations in laboratory-evolved*
578   *microbes from next-generation sequencing data using breseq.* Methods Mol Biol,
579   2014. **1151**: p. 165-88.
580   22.   Schwartz, D.J., et al., *Gut pathogen colonization precedes bloodstream infection in*
581   *the neonatal intensive care unit.* Science Translational Medicine, 2023. **15**(694).
582   23.   Carl, M.A., et al., *Sepsis From the Gut: The Enteric Habitat of Bacteria That Cause*
583   *Late-Onset Neonatal Bloodstream Infections.* Clinical Infectious Diseases, 2014.
584   **58**(9): p. 1211-1218.
585   24.   Rodriguez-R LM, C.R., Viver T, Feistel DJ, Lindner BG, Venter SN, Orellana LH,
586   Amann R, Rossello-Mora R, Konstantinidis KT., *An ANI gap within bacterial species*
587   *that advances the definitions of intra-species units.* mBio, 2024. **15**: p. e02696-23.
588   25.   Vornhagen, J., et al., *Combined comparative genomics and clinical modeling reveals*
589   *plasmid-encoded genes are independently associated with Klebsiella infection.* Nat
590   Commun, 2022. **13**(1): p. 4459.
591   26.   Ludden, C., et al., *Defining nosocomial transmission of Escherichia coli and*
592   *antimicrobial resistance genes: a genomic surveillance study.* Lancet Microbe, 2021.
593   **2**(9): p. e472-e480.
594   27.   Gorrie, C.L., et al., *Gastrointestinal Carriage Is a Major Reservoir of Klebsiella*
595   *pneumoniae Infection in Intensive Care Patients.* Clinical Infectious Diseases, 2017.
596   **65**(2): p. 208-215.
597   28.   Thanert, R., et al., *Persisting uropathogenic Escherichia coli lineages show*
598   *signatures of niche-specific within-host adaptation mediated by mobile genetic*
599   *elements.* Cell Host Microbe, 2022. **30**(7): p. 1034-1047 e6.

600   29.   Rebelo, A.R., et al., *One day in Denmark: whole-genome sequence-based analysis*
601   *of Escherichia coli isolates from clinical settings.* J Antimicrob Chemother, 2025.
602   30.   Bush, S.J., et al., *Genomic diversity affects the accuracy of bacterial single-*
603   *nucleotide polymorphism-calling pipelines.* Gigascience, 2020. **9**(2).
604   31.   Jamin, C., et al., *Harmonization of whole-genome sequencing for outbreak*
605   *surveillance of Enterobacteriaceae and Enterococci.* Microb Genom, 2021. **7**(7).
606   32.   Wylie, K.M., et al., *Whole-Genome Sequencing of Klebsiella pneumoniae Isolates to*
607   *Track Strain Progression in a Single Patient With Recurrent Urinary Tract Infection.*
608   Front Cell Infect Microbiol, 2019. **9**: p. 14.
609   33.   Okusu, H., *AcrAB Efflux Pump Plays a Major Role in the Antibiotic Resistance*
610   *Phenotype of Escherichia coli MultipleAntibiotic-Resistance (Mar) Mutants.* Journal of
611   Bacteriology, 1996. **178**: p. 306-308.
612   34.   Padilla, E., et al., *Klebsiella pneumoniae AcrAB efflux pump contributes to*
613   *antimicrobial resistance and virulence.* Antimicrob Agents Chemother, 2010. **54**(1): p.
614   177-83.
615   35.   Lv, F., et al., *Overexpression of Efflux Pumps Mediate Pan Resistance of Klebsiella*
616   *pneumoniae Sequence Type 11.* Microbial Drug Resistance, 2021. **27**(10): p. 1405-
617   1411.
618   36.   Chowdhury, N., et al., *Identification of AcrAB-TolC Efflux Pump Genes and Detection*
619   *of Mutation in Efflux Repressor AcrR from Omeprazole Responsive Multidrug-*
620   *Resistant Escherichia coli Isolates Causing Urinary Tract Infections.* Microbiol
621   Insights, 2019. **12**: p. 1178636119889629.
622   37.   Alav, I., J.M. Sutton, and K.M. Rahman, *Role of bacterial efflux pumps in biofilm*
623   *formation.* J Antimicrob Chemother, 2018. **73**(8): p. 2003-2020.
624   38.   Tang, M., et al., *The role and relationship with efflux pump of biofilm formation in*
625   *Klebsiella pneumoniae.* Microbial Pathogenesis, 2020. **147**: p. 104244.
626   39.   Vuotto, C., et al., *Biofilm formation and antibiotic resistance in Klebsiella pneumoniae*
627   *urinary strains.* Journal of applied microbiology, 2017. **123**(4): p. 1003-1018.
628   40.   Hennequin, C., et al., *Antibiotic resistance and plasmid transfer capacity in biofilm*
629   *formed with a CTX-M-15-producing Klebsiella pneumoniae isolate.* Journal of
630   Antimicrobial Chemotherapy, 2012. **67**(9): p. 2123-2130.
631   41.   Reza, A., J.M. Sutton, and K.M. Rahman, *Effectiveness of efflux pump inhibitors as*
632   *biofilm disruptors and resistance breakers in gram-negative (ESKAPEE) bacteria.*
633   Antibiotics, 2019. **8**(4): p. 229.
634   42.   Nielsen, K.L., et al., *Adaptation of Escherichia coli traversing from the faecal*
635   *environment to the urinary tract.* Int J Med Microbiol, 2016. **306**(8): p. 595-603.
636   43.   Prieto, A., et al., *Tracking bacterial virulence: global modulators as indicators.* Sci
637   Rep, 2016. **6**: p. 25973.
638   44.   Barrios, A.F.G., et al., *Hha, YbaJ, and OmpA regulate Escherichia coli K12 biofilm*
639   *formation and conjugation plasmids abolish motility.* Biotechnology and
640   Bioengineering, 2006. **93**(1): p. 188-200.
641   45.   Bandeira, M., et al., *Insights on Klebsiella pneumoniae Biofilms Assembled on*
642   *Different Surfaces Using Phenotypic and Genotypic Approaches.* Microorganisms,
643   2017. **5**(2).
644   46.   Krall, L.J., et al., *Invasiveness of Escherichia coli Is Associated with an IncFII*
645   *Plasmid.* Pathogens, 2021. **10**(12).
646   47.   Ueda, T., et al., *Functions of the Hha and YdgT proteins in transcriptional silencing by*
647   *the nucleoid proteins, H-NS and StpA, in Escherichia coli.* DNA Res, 2013. **20**(3): p.
648   263-71.
649   48.   Sharma, V.K. and B.L. Bearson, *Hha controls Escherichia coli O157:H7 biofilm*
650   *formation by differential regulation of global transcriptional regulators FlhDC and*
651   *CsgD.* Appl Environ Microbiol, 2013. **79**(7): p. 2384-96.
652   49.   Fang, C.T., et al., *DNA adenine methylation modulates pathogenicity of Klebsiella*
653   *pneumoniae genotype K1.* J Microbiol Immunol Infect, 2017. **50**(4): p. 471-477.

50.   Desvaux, M., et al., *Pathogenicity Factors of Genomic Islands in Intestinal and*
Extraintestinal Escherichia coli. Front Microbiol, 2020. **11**: p. 2065.
51.   Xu, W.-y., Y.-j. Li, and C. Fan, *Different loci and mRNA copy number of the increased*
serum survival gene of Escherichia coli. Canadian Journal of Microbiology, 2018.
64(2): p. 147-154.
52.   Chen, J.H., et al., *Contribution of outer membrane protein K36 to antimicrobial*
resistance and virulence in Klebsiella pneumoniae. J Antimicrob Chemother, 2010.
65(5): p. 986-90.
53.   Garcia, E.C., A.R. Brumbaugh, and H.L. Mobley, *Redundancy and specificity of*
Escherichia coli iron acquisition systems during urinary tract infection. Infection and
immunity, 2011. **79**(3): p. 1225-1235.
54.   Mann, R., et al., *Metabolic Adaptations of Uropathogenic E. coli in the Urinary Tract.*
Front Cell Infect Microbiol, 2017. **7**: p. 241.
55.   Krawczyk, B., et al., *Characterisation of Escherichia coli isolates from the blood of*
haematological adult patients with bacteraemia: translocation from gut to blood
requires the cooperation of multiple virulence factors. Eur J Clin Microbiol Infect Dis,
2015. **34**(6): p. 1135-43.

686 **Tables and figures:**

687 **Table 1:** Characteristics of isolate pairs and clinical data of patients with GNBSI

Pair	Species of isolate pair	Days from faeces - blood isolation	Age at GNBSI	CLABSI	ICU patient at time of GNBSI	Antibiotic exposure between faeces and blood isolate
1	*K. pneumoniae*	144	12 years	Y	N	Ciprofloxacin, Clindamycin, Ceftriaxone, Piperacillin-Tazobactam, Co-trimoxazole, Gentamicin
2	*K. pneumoniae*	87	11 months	Y	Y	Ciprofloxacin, Cefalexin, Piperacillin-Tazobactam, Ceftriaxone, Vancomycin, Gentamicin
3	*E. coli*	185	9 months	Y	Y	Co-amoxiclav, Cefalexin
4	*E. coli*	31	12 months	Y	Y	Ciprofloxacin, Meropenem
5	*E. coli*	17	1 month	Y	Y	Ciprofloxacin, Vancomycin

697 **Table 2A:** MLST, K-type, O-antigen type and wzi-type of *K. pneumoniae* isolates

698 within pairs, determined using Kleborate and Kaptive.

Species	Klebsiella pneumoniae			
Pair	Pair 1		Pair 2	
Isolate	Faeces	Blood	Faeces	Blood
MLST	ST922	ST922	ST461	ST461
K type	K28	K28	K25	K25
O type	O1	O1	O1	O1
wzi type	wzi84	wzi84	wzi25	wzi25

701 **Table 2B:** Phylotype and MLST of *E. coli* isolates within pairs using ClermonTyping

702 and Institut de Pasteur database.

Species	Escherichia coli					
Pair	Pair 3		Pair 4		Pair 5	
Isolate	Faeces	Blood	Faeces	Blood	Faeces	Blood
MLST	ST131	ST131	ST131	ST131	ST12	ST12
FumC type	C40	C40	C40	C40	C13	C13
O type	H4	H4	H4	H4	H1	H1
H type	O25	O25	O25	O25	O4	O4

711   **Table 3: Specific mutations identified in relation to phenotypes of interest.**

Pair	Mutation type	Mutation	Annotated gene	Relevance of mutation
1	nsSNP	D47E (GAC→GAA)	ompk36	Bloodstream survival factor
	nsSNP	Q62K (CAA→AAA)	dam	Bloodstream survival factor
	nsSNP	T57A (ACT→GCT)	dam	Bloodstream survival factor
	nsSNP	Q141L (CAG→CTG)	dam	Bloodstream survival factor
	Insertion	+GGT	Haemolysin expression modulating protein/hypothetical protein	Regulator of virulence factors, known to regulate biofilm formation
	iSNP	2 bp→TT	Haemolysin expression modulating protein/hypothetical protein	Regulator of virulence factors, known to regulate biofilm formation
	iSNP	G→T	Haemolysin expression modulating protein/hypothetical protein	Regulator of virulence factors, known to regulate biofilm formation
	iSNP	A→G	Haemolysin expression modulating protein/hypothetical protein	Regulator of virulence factors, known to regulate biofilm formation
	iSNP	A→T	Haemolysin expression modulating protein/hypothetical protein	Regulator of virulence factors, known to regulate biofilm formation
	sSNP	A40A (GCT→GCC)	Haemolysin expression modulating protein	Regulator of virulence factors, known to regulate biofilm formation
	sSNP	R11R (CGC→CGG)	Haemolysin expression modulating protein	Regulator of virulence factors, known to regulate biofilm formation
	Deletion	Δ916 bp	Fec operon	Iron uptake and utilisation
2	nsSNP	T37I (ACT→ATT)	Multiple antibiotic resistance protein MarB	Drug-efflux pump implicated in antibiotic resistance
	Deletion	Δ348 bp	YcAD - MFS transporter	Drug efflux pump implicated in antibiotic resistance
	Deletion	Δ254 bp	permease of drug/metabolite transporter	Drug efflux pump implicated in antibiotic resistance
	Deletion	Δ250 bp	inner membrane protein/drug antiporter	Drug efflux pump implicated in antibiotic resistance
	Deletion	Δ243 bp	LysR	Transcriptional regulator implicated in antibiotic resistance
	Deletion	Δ241 bp	MFS-type transporter	Drug efflux pump implicated in antibiotic resistance
	Deletion	Δ239 bp	RND efflux transporter	Drug efflux pump implicated in antibiotic resistance
	Deletion	Δ236 bp	LysR transcriptional regulator	Drug efflux pump implicated in antibiotic resistance
	Deletion	Δ321 bp	acrR	Transcriptional regulator implicated in antibiotic resistance and biofilm formation
	nsSNP	A434S (GCC→TCC)	Protease III precursor (EC 3.4.24.55)	Biofilm formation
	nsSNP	V701A (GTG→GCG)	T6SS component TssM (IcmF/VasK)	Virulence factor, involved in biofilm formation
3	Deletion	Δ235 bp	Lipopolysaccharide ABC transporter	Transporter protein implicated in antibiotic resistance
	nsSNP	T115A (ACC→GCC)	Phage endopeptidase Rz	Phage protein, adjacent to virulence factor iss (involved in bloodstream survival)
	nsSNP	L237P (CTT→CCT)	Mobile element protein	MGE adjacent to Fec operon (involved in iron uptake and utilisation)
	nsSNP	A238G (GCC→GGC)	Mobile element protein	MGE adjacent to Fec operon (involved in iron uptake and utilisation)
	nsSNP	S932T (AGC→ACC)	Antigen 43	Virulence factor, involved in biofilm formation
	nsSNP	A763V (GCC→GTC)	Antigen 43	Virulence factor, involved in biofilm formation
	sSNP	G765G (GGA→GGT)	Antigen 43	Virulence factor, involved in biofilm formation
4	Deletion	Δ286 bp	MFS type transporter	Drug efflux pump implicated in antibiotic resistance
	sSNP	G977G (GGG→GGT)	Antigen 43	Virulence factor, involved in biofilm formation
	sSNP	G972G (GGT→GGA)	Antigen 43	Virulence factor, involved in biofilm formation
	nsSNP	S932T (AGC→ACC)	Antigen 43	Virulence factor, involved in biofilm formation
	nsSNP	V375A (GTC→GCC)	Antigen 43	Virulence factor, involved in biofilm formation
	Deletion	Δ265 bp	Ferrochelatase	Iron uptake and utilisation
	Deletion	Δ236 bp	Ferredoxin reductase	Iron uptake and utilisation
	Deletion	Δ119 bp	cog3210	Iron uptake and utilisation
	Deletion	Δ95 bp	cog3210	Iron uptake and utilisation
5	Synonymous SNP	L123L (CTG→TTG)	Manganese ABC transporter, inner membrane permease protein SitC	Iron uptake and utilisation
	Deletion	Δ297 bp	Transcriptional regulator, acrR family	Transcriptional regulator implicated in antibiotic resistance and biofilm formation
	Deletion	Δ295 bp	Spermidine/putrescine ABC transporter	Transporter protein implicated in antibiotic resistance
	Deletion	Δ281 bp	Fe2+ ABC transporter, substrate binding protein	Iron uptake/homeostasis
	Non-synonymous SNP	V32E (GTG→GAG)	Transcriptional repressor of aga operon	Transcriptional regulator involved in biofilm formation
	Non-synonymous SNP	M300L (ATG→CTG)	Lipopolysaccharide export system permease protein LptG	Biofilm formation
	Deletion	Δ246 bp	TctA transporter	Drug-efflux pump implicated in antibiotic resistance
	Deletion	Δ245 bp	AaeAB efflux system	Drug efflux pump implicated in antibiotic resistance

713 **Figure 1:** Pairwise comparison of Average Nucleotide Identity and core genome

714 SNPs between isolates

Figure 2A: Antimicrobial resistance gene (ARG) presence in all isolates. Each tile

represents the presence of an ARG as detected by ResFinder 4.2.2. using whole

genome sequencing data. ARGs are categorised by drug class that they are

predicted to confer resistance to.

Figure 2B: Virulence gene presence in *K. pneumoniae* isolates. Each tile

demonstrates the presence of a virulence gene as detected by the VF analyser using

whole genome sequencing data. Virulence genes are categorised by class of

virulence factor encoded by the gene.

Figure 2C: Virulence gene presence in *E. coli* isolates. Each tile demonstrates the

presence of a virulence gene as detected by the VF analyser using whole genome

sequencing data. Virulence genes are categorised by class of virulence factor

encoded by the gene.

Figure 2D: Plasmid presence in all pairs of isolates. Each tile demonstrates the

presence of a Plasmid replicon gene in each isolate as detected by PlasmidFinder,

using whole genome sequencing data.

Figure 3A: Minimum inhibitory concentration (MIC) (μg/ml) as determined by broth

microdilution of Ciprofloxacin, Co-amoxiclav, Piperacillin-Tazobactam, Cefotaxime

and Gentamicin for Pair 2. Each plotted point represents the MIC (μg/mL) of a single

biological replicate in broth microdilution. Dashed line = EUCAST clinical breakpoint

for MIC.

Figure 3B: Minimum inhibitory concentration (MIC) as determined by broth

microdilution of Piperacillin-Tazobactam for Pair 3. Each plotted point represents the

MIC of a single biological replicate in broth microdilution. Dashed line = EUCAST

clinical breakpoint for MIC.

Figure 4: Crystal violet assay for biofilm formation in 96-well plate. Each bar

represents mean OD_{600} (n=12) for faecal and blood isolates after crystal violet

staining, normalised to biomass. This is used as the measure of biofilm formation

capacity. Error bars represent standard error of the mean. * = Statistical significance

as determined by paired t-test, p=0.05.

Figure 5: Heatmap of all mutations assigned to COG categories across isolate pairs.

Each tile represents the total count of mutations within each COG (Cluster of

Orthologous Groups) category for each isolate pair. White tiles represent the

absence of mutations in a specific COG category.

```
Faeces      tattaccgggacacctcaggatat    aaattaatgacgaaac
Blood_1     aactaccgggacacctaaggatataaaaaaccaatgacgaaac
Blood_2     aactaccgggacacctaaggatataaaaaaccaatgacgaaac
```

Figure 6: Alignment of upstream intergenic region and first 10bp (underlined) of *hha*

gene of faecal isolate and two copies in blood isolate. (Yellow = duplication of bp.

Red = intergenic SNP).