Detecting Foldback Artifacts in Long Reads

Jakob M. Heinz[1,2,3], Matthew Meyerson[3,4,5], Heng Li[1,2,*]

[1]Department of Biomedical Informatics, Harvard Medical School, Boston, MA, United States

[2]Department of Data Science, Dana-Farber Cancer Institute, Boston, MA, United States

[3]Cancer Program, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[4]Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA, USA

[5]Department of Genetics, Harvard Medical School, Boston, MA, USA

*Corresponding author: hli@ds.dfci.harvard.edu

Abstract

Long-read sequencing data is useful for detecting large and complex structural variations; however, technical artifacts can lead to false structural variant calls. In our analyses, we became aware of a foldback artifact in long-read data. Therefore, we developed the open-source Breakinator tool to flag putative foldback artifact reads, as well as previously known chimeric artifacts. Through an alignment-based approach, Breakinator can detect artifacts missed by existing quality control tools. We profiled the occurrences of foldbacks and chimeric reads in both nanopore and single-molecule real-time sequences across a range of specimens, library types, sequencing chemistries, sequencing machines, and base-calling software.

Keywords

Long-read sequencing, Nanopore, RNA-Sequencing, Technical artifacts, Quality control

Background

Advances in long-read sequencing technologies, such as those by Pacific Biosciences (PacBio; single-molecule real-time sequencing) and Oxford Nanopore Technologies (ONT; nanopore sequencing), have made the reliable resolution of complex and repetitive regions of the genome possible, which cannot be achieved by short-read sequencing. The ability of long reads to span these complicated regions is useful for detecting large complex structural variations (SV) in the genome [1,2]. Typically, for long-read SV calling, reads are aligned to a reference genome to identify those with a supplementary alignment, a so-called split read, where a portion of the read maps to one genomic region, and the remainder maps to another distinct region causing the aligner to split the read into separate alignment blocks [3–5]. However, systemic technical artifacts in the reads can lead to false SV calls. Chimeric reads have been a pervasive issue throughout the history of long-read sequencing [6–9]. Technical artifacts can arise through issues in library preparation or computational errors in distinguishing independent molecules [10,11]

Here, we have become aware of another type of technical artifact, a foldback (also known as a duplicated inversion) [12]where half the read maps to one strand and the other half maps to the same location, but on the opposite strand, appearing as a near perfect concatemerization of the forward and reverse strands of a read. A previous study by Soneson *et al.* had noted an elevated rate of palindromic reads in ONT direct-cDNA data with supplementary alignments overlapping their primary alignment on the opposite strand. They hypothesized that these may represent un-split reads by the base caller [11]. However, to our knowledge, neither the cause nor the prevalence of these foldback artifacts was explored by any later studies with more recent data, such as the latest ONT nanopores (R10.4) or PacBio HiFi.

Results and Discussion

The Breakinator tool identifies putative foldback and chimeric artifacts by parsing a read

alignment file. Our tool extracts the primary and all supplemental alignments of a read and

classifies them as either a true breakpoint, chimeric, or foldback using the classification scheme in

Fig. 1a. While all metrics can be set by the user, we classify a chimeric artifact as a read that has

segments mapping to different chromosomes, or more than 1 Mb apart within the same

chromosome. We classify a read as a foldback artifact if the supplementary alignment maps to the

same location as the primary alignment (within 200 bps) on the opposite strand. For samples where

true foldback events may be expected, as in cancers, we added a symmetry option to our tool to

only report reads as foldback artifacts if the foldback occurs within a 5% margin on either side of

the middle of the read, to retain true foldback event supporting reads.

Fig. 1 a). Metrics used to define types of breakpoints observed in this study. A "True Break" is a breakpoint that is not found to be chimeric or a foldback, such as the tandem duplication example shown here. A "Foldback" is a read where the second segment maps to approximately the same location (within 200bps) as the first segment, but on the opposite strand. A "Chimeric" is where the second segment maps more than 1 Mb away on the same chromosome (intrachromosomal) or to a different chromosome (interchromosomal). b) IGV schematization of two foldback artifact reads from the HG002_PAW70337_sup sample supporting the same breakpoint at approximately 113,286 kb on chromosome X of the HG002 diploid assembly. Read A (5f80430c-2b83-408b-89b2-63b61b1173cc) is an example of a read with no sequence between the two segments of the

foldback. Read B (115c006d-16a6-47b1-bf5c-c5eb9bb85f69) is an example where there are ~2kb

of sequence between the segments.

To verify that our tool properly identifies foldback events, we used PacBio's samples of

self-complementary adeno-associated viruses (scAAV) CBA-eGFP as a positive control. Since

scAAV-CBA-eGFP constructs were designed to be perfectly palindromic, we expected our tool to

detect them as foldbacks. With the Breakinator, we find that 97.49% CBA-eGFP sample reads are

scAAV compared to the 97.69% reported by Talevich *et al*. [13]. This verification also presented

an interesting additional use case for the Breakinator for the rapid identification of scAAVs in a

sample.

We first noticed a potential technical artifact when analyzing transcriptomic data of the

HCC1395 cell line generated by Cotto *et al*. [14]. In the ONT direct-cDNA samples, we observed

many reads with supplementary alignments overlapping the primary alignment on the opposite

strand, indicative of a foldback SV, that were not present in the ONT direct-RNA sample run on

the same RNA replicate. Manual inspection of the unique breakpoint locations in IGV [15] found

that those detected from direct-RNA had supporting evidence from PacBio HiFi gDNA sequences,

while most of the direct-cDNA breakpoints did not, leading us to suspect that many of these events

were caused by a technical artifact. After developing the Breakinator, we used it to identify all

breakpoints in the sample and merge breakpoints in the same region (see Methods). We found

35,843 unique breakpoint locations with at least five supporting reads in the direct-cDNA samples,

compared to just eight in the direct-RNA. While foldback SVs occur in cancer genomes, they have

not been observed at such a high frequency [12], and are not expected to vary so greatly by library

type.

To explore whether this was a dataset-specific sequencing result or a systemic technical

artifact, we downloaded publicly available datasets representing four library types: direct-RNA,

direct-cDNA, cDNA, and gDNA, from four sample types (HG002 cell line, K562 cell line,

HCC1395 cell line, and mouse tissue). A range of sequencing chemistries, machines, and base-

calling software were evaluated through these samples (Additional file 1). The HG002 samples

offer a useful neutral background to investigate technical artifacts because they have a high-quality

personal diploid reference assembly to which the reads can be aligned [16]Therefore, we would

expect no germline variants, given that this is the reference, only variations between samples of

the cell line, which are likely limited [17]. Each sample was aligned to the corresponding reference

genome with the appropriate Minimap2 [3] parameters for each data type (see Methods).

While an alignment-based artifact detection method, as is implemented by the Breakinator,

may lead to issues in distinguishing true SV reads from artifact reads, the increased specificity

offered by alignment can be useful. Commonly used quality control tools for long reads, such as

fastplong, Porechop, pychopper, and Restrander [18–21], were not able to identify many of the

foldback or chimeric reads flagged by the Breakinator (Additional file 2). We suspect this is due

to their reliance on identifying adapter sequences within the middle of a read sequence to identify

a concatemerized read. We found that typically less than half of the putative artifactual reads

detected by the Breakinator had more than 11 bps of non-mapping sequence between alignments

(Additional file 3), making it unlikely they will be detected with existing methods. This is

highlighted in Fig. 1b, where two foldback reads identified by the Breakinator are shown. The

reads are both nearly perfectly palindromic and map to the same breakpoint location, yet Read A

has no sequence between mappings, while Read B has ~2kb of sequence that could be searched

for retained adaptors. 27 other reads (not shown) cover this foldback location, with no indication

of a breakpoint.

Summary statistics across library types and samples evaluated with the Breakinator are

shown in Fig. 2a. ONT direct-cDNA samples were found to have substantially elevated levels of

foldback artifacts (~10-25% of all reads). Meanwhile, at most one foldback read was observed in

each of the ONT direct-RNA samples. The ONT cDNA samples had the lowest rates of foldback

artifacts (~0.1 – 0.2). We suspect this low rate is due to the cDNA samples being the only datatype

in this study that undergoes PCR amplification, and palindromic sequences such as these foldbacks

are known to be difficult to PCR amplify as they will bind to themselves [22,23]. In ONT gDNA,

we observed foldback rates of ~0.5–0.7%. While only three PacBio samples were investigated,

they did not show any evidence of systemic foldback artifacts. There were virtually no foldback

reads observed in the PacBio HiFi (Revio) or Mas-Seq (Vega) data, while 0.04% of the Iso-Seq

(Sequel II) reads were considered foldbacks. For five of the samples, from K562 and HCC1395

cells, the same RNA replicate was sequenced with different technologies, allowing for direct

comparisons in the rates of foldbacks and chimeras across libraries without replicate bias. It was

observed that for all replicates, the rates of foldbacks and chimeras were much higher in ONT

direct-cDNA library samples, as shown in Fig. 2b.

Fig. 2. a) The observed rates of Foldback, Chimeric, and any split reads out of all reads that passed the filtering criteria for direct-cDNA, direct-RNA, cDNA, and gDNA data from four different sample types: HCC1395, K562, mouse tissue, and HG002. B) Paired comparisons for the five RNA replicates that were sequenced with different technologies for both foldback and chimeric

reads. c) The proportion of Foldback, Chimeric, and True Break reads in relation to the number of split reads. d) Recurrence of breakpoints at 2, 5, and 10 supporting reads for Foldback, Chimeric, and True Break reads.

While the rates of putative foldback and chimeric artifacts across all reads are relatively low for libraries other than ONT direct-cDNA, they represent a far larger proportion of the reads used for SV calling, namely, reads with split alignments. Fig. 2c highlights that these artifacts can cause issues even with ONT cDNA or gDNA data. In cDNA, foldback artifacts account for 10-20% of all split reads, while chimeras are ~30-40%. In gDNA, foldbacks are also 15-20% of split reads, and chimeric reads are 50-60% of all split reads (Fig. 2.c). These artifactual reads were also observed to be recurrent at the same locations, which made it more likely they would be called as a false positive SV. Fig. 2d highlights that many of the locations to which these reads align have more than 2, 5, or 10 reads of support.

Given that the rates of foldback artifacts appear relatively consistent among runs of the same data type, but vary between data types that are prepared with different chemistries, we believe it is most likely that these artifacts arise during library preparation, rather than a failure of the base-caller to recognize a pore-open signal or an adapter sequence, as was hypothesized by Soneson *et al*. [11]. Additionally, we observed both forward and reverse strands of foldback and chimeric reads, which can only arise during library preparation, compared to an informatic chimera, which would only occur in one reading direction. Finally, we investigated the raw signal of eight putative artifactual reads from the SGNEx K562 direct-cDNA replicate 2 run 1 to visualize the electrical signal interpreted by the base-caller. We did not observe any notable difference in signal around

159   the breakpoint region for artifacts compared to true breakpoints (Additional file 4). Thus, we

160   conclude that Breakinator can identify foldback artifacts in sequencing data, which may arise due

161   to an experimental feature in library preparation.

Conclusions

We present evidence of a new technical artifact, a foldback read. If these technical artifacts

are not properly removed, they can lead to tangles in an assembly graph or false-positive SV calls.

We developed the open-source Breakinator tool to flag putative foldback and chimeric technical

artifacts in long-read sequences. Many current quality control tools rely on detecting incorrectly

incorporated adapter sequences, but through a reference alignment-based approach, we can detect

reads that may not have retained any non-sample sequence, but still likely represent a non-real

biological concatenation. In this study, we profiled both ONT and PacBio data across a range of

specimens, library types, sequencing chemistries, sequencing machines, and base-callers. We find

that foldback artifacts occur throughout ONT library types and are particularly high in direct-

cDNA libraries. We believe the Breakinator will offer a useful tool for quality control in the

analysis of long-read data.

Methods

All data used in this study were downloaded from publicly available sources, with the links

available in Additional file 1. The K562 samples were generated by the SGN-Ex consortium [24].

HCC1395 samples were generated by Cotto *et al.* [14]. The mouse tissue samples were generated

by Sessegolo *et al.* [25]. The HG002 gDNA samples were downloaded from ONT's 2025 Genome

in a Bottle Data (GIAB) Release [26]. The HG002 direct-RNA002, PacBio Iso-seq, and Mas-seq

were made available by GIAB. The HG002 dRNA sample SRR30901279 and the HG002 cDNA

data were generated by Zheng *et al*.[27] The HG002 PacBio HiFi gDNA is a public dataset made

available by PacBio [28]. The PacBio scAAV sample was generated by Talevich *et al*. [13]

The samples were aligned to their respective reference genomes using Minimap2 [3] K562

and HCC1395 samples were aligned to T2T-CHM13v2.0 [29,30]. The mouse samples were

aligned to GRCm39 [31]. HG002 samples were aligned to the HG002v1.0.1 diploid genome

[16]To align dRNA ONT, the following parameters were used: "-cx splice -uf -k14 --

secondary=no". For ONT direct-cDNA and cDNA alignments, "-cx splice --secondary=no"

parameters were used. ONT gDNA data were aligned with "-cx map-ont --secondary=no"

parameters. For PacBio Iso-Seq and Mas-Seq data, we used "-cx splice:hq -uf --secondary=no"

parameters. Finally, PacBio HiFi data were aligned with "-cx map-pb-- secondary=no." Aligning

to a diploid genome decreases the mapping quality of many reads, as there will likely be at least

two locations where they map well. To address this issue, HG002 sample reads were aligned to the

HG002v1.0.1 diploid genome by aligning the reads to each haplotype separately using Minimap2

with the flags "--secondary=no" and "--paf-no-hit" specified. We wrote a preprocessing script for

the Breakinator to compare the alignment of every read to the maternal and paternal haplotype

assemblies, choosing to keep the alignment with the highest alignment score. If the alignment

scores were equal, then the maternal haplotype alignment was chosen.

The PAF files were generated using either "paftools.js sam2paf -p" on a SAM file or

Minimap2 with the "--secondary=no" flag. We designed the Breakinator to parse the input PAF

file to detect reads with supplementary alignments. Secondary alignments were ignored.

Alignments with mapping quality less than 10 or lengths less than 200 bps were filtered out. The

primary and all supplementary alignments for a given read were identified and sorted by read start

location. If the first read segment mapped to the forward strand, then the end of the alignment of

the first segment and the start of the second segment were identified as the breakpoint location. If

the first segment mapped to the reverse strand, the reciprocal start and end locations were taken

for the breakpoint. As shown in Fig. 1a, if the breakpoint was to the opposite strand within 200

bps of the end of the alignment of the first segment, it was classified as a foldback. If the breakpoint

was to another chromosome (interchromosomal) or more than 1 Mb away (intrachromosomal), it

was classified as a chimeric read. The 1 Mb default threshold for intrachromosomal chimeras was

chosen because only two introns, out of 177,839, on MANE selected protein-coding genes were

longer than 1Mb [32]. If the breakpoint was not chimeric or a foldback, then it was considered a

true breakpoint. The identified breakpoints and their classification were generated for every

sample. While uncommon, it is possible to have multiple breakpoints on the same read. To classify

at the read level, we used the following approach: if there was any foldback or chimeric breakpoint

on the read, regardless of how many true breakpoints there were, it was classified as an artifact. If

there were multiple foldback or chimeric breakpoints on the read, the read was classified according

to which breakpoint occurred most frequently. If they were equal, then the read was classified as

chimeric. All summary statistics are generated using the read classification.

To compare how existing tools detect the reads we had identified as artifactual, we filtered

all reads with fastplong, porechop, and restrander, respectively [18,19,21]. Fastplong was run with

the following command: "fastplong -i sample.fastq -o filter_pass.fastq --failed_out

failed_reads.fastq." Restrander was run only on our cDNA samples as it didn't support any other

data types, with the command "restrander sample.fastq filter_pass.fastq config.json" Restrander

required a configuration file of adaptor sequences, so for samples that did not have their library

version supported by Restrander, we ran them with the closest version of the library supported by

Restrander. For example, DCS108 samples were run with the SQK-DCS109 config file. Porechop

was also only run on the transcriptomic data using the command: "porechop -i sample.fastq --

discard_middle -o filter_pass.fastq." For each tool, the read IDs that passed their respective filter

were used to subset the reads we found to pass the Breakinator default filter of a minimum of 200

bps of alignment and a mapping quality of 10. The reads classified as foldback, chimeric, or any

breakpoint were similarly subset to include only the reads that passed the respective filter to

calculate the proportions reported in Additional file 2. PyChopper was only run on the three

HCC1395 samples, as we found it to be too slow to run on all samples. The length of the unaligned

sequence between aligning segments of the breakpoint was identified by running the Breakinator

with the "--rcoord" flag to output the read coordinates of the breakpoint along with the reference

coordinates of the breakpoint. The reads were subset to foldback, chimeric, and true breakpoints,

and the proportions of reads with at least 11 bps between the read coordinates of the breakpoint

were calculated for each subset in Additional file 3.

To generate summary statistics for Fig.2, the Breakinator was run on every sample using

the "-i in.paf --tabular -o out.txt" parameters. The PacBio scAAV sample was evaluated with the

Breakinator using the stricter parameters of a symmetry filter and half the default margin (--sym -

-margin 0.025). To determine which breakpoints were recurrent, we extracted the subset of

breakpoints that were flagged as either "Foldback", "Chimeric", or "True Break", respectively,

then for each file we merged any breakpoints that were within 100 bps of each other to account for

inexact alignments around the breakpoint, to determine the number of reads supporting a

breakpoint using our merge_breaks.py script. The merging process was run with the -s flag set to

2, 5, and 10, which sets the minimum number of reads needed to support a breakpoint. The reads

that had support of other reads at the respective support thresholds were summed and divided by

the total number of reads in the subset to determine the proportion of reads with support, as shown

in Fig. 2d.

The SGNEx K562 direct-cDNA replicate 2 run 1 sample was chosen to investigate the raw

electrical signal that the ONT base caller interprets because its fast5 files were publicly available,

and we discovered numerous putative artifacts in the reads. The fast5 files were indexed with

nanopolish using "nanopolish index -d fast5_directory reads.fastq" [33]. Nanopolish eventalign

was used to determine which region of the raw signal corresponds to the breakpoint location in the

read. To explore the signal in read coordinates, we converted the read fastq file to a fasta file to

use as the genome input, then aligned the reads to themselves with "minimap2 -ax map-ont -o

reads2reads.sam reads.fastq reads.fa " and used this as the BAM input for the eventalign command

as follows: "nanopolish eventalign -r reads.fastq -b reads2reads.bam -g reads.fasta --signal-index".

We chose eight reads in total to investigate the raw signal. We investigated foldbacks, chimeras,

and true breakpoints with and without unaligned sequences in the middle, as well as a read with

no breakpoint. The start and end of the breakpoint locations in the read were determined from the

Breakinator output and converted to the signal coordinates using the nanopolish eventalign output.

The fast5 samples were graphed, and the breakpoint coordinates in the signal were highlighted to

manually investigate the signal for any signs of increased noise around the breakpoint.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Additional files

Additional_file_1.xlsx – All public data used in this study and their sources. Includes metadata on datatype, sequencer, pore, chemistry, and base-caller.

Additional_file_2.xlsx – Proportions of reads remaining, including artifact subsets, after filtering with other current tools (fastplong, Porechop, pychopper, and Restrander).

Additional_file_3.xlsx – Proportions of reads that had 11 bps or greater of unmapped read sequence between alignment blocks.

Additional_file_4.html – SGNEx K562 direct-cDNA replicate 2 run 1 raw signal analysis around identified breakpoints for eight reads.

Availability of data and materials

The Breakinator tool and additional software are available on GitHub at https://github.com/jheinz27/breakinator. All data used in this study were previously made publicly available, and their sources are listed in Additional file 1. The alignment files for every sample, as well as all breakpoints identified and their Breakinator classification, are available on Zenodo at: https://zenodo.org/records/15882143.

Competing interests

M.M. holds equity in Bayer, Delve Bio, Isabl, and Karyoverse; consults for Delve Bio; receives research funding from Bayer; and receives patent licensing payments from Bayer and Labcorp.

Funding

J.M.H. is supported by the U.S. National Institutes of Health training grant T32HG002295.

M.M. is supported by NIH grants R35CA197568 and U24CA264029. H.L. is supported by NIH grants U24CA294203 and R01HG010040

Authors' contributions

J.M.H., M.M., and H.L conceived the project. J.M.H. implemented the method, analyzed the data, and drafted the manuscript. M.M. and H.L provided edits. All authors read and approved the final manuscript.

Acknowledgments

We would like to thank Drs. Neng Huang, Owen Hirschi, Alaina Shumate, Maximillian Marin, and Victoria Popic, as well as all members of the Meyerson and Li labs, for their helpful discussions.

References

1. Mahmoud M, Gobet N, Cruz-Dávalos DI, Mounier N, Dessimoz C, Sedlazeck FJ. Structural variant calling: the long and the short of it. Genome Biol. 2019;20.

2. Sedlazeck FJ, Rescheneder P, Smolka M, Fang H, Nattestad M, Von Haeseler A, et al. Accurate detection of complex structural variations using single-molecule sequencing.

Nature Methods 2018 15:6 [Internet]. 2018 [cited 2025 Apr 17];15:461–8. Available

from: https://www.nature.com/articles/s41592-018-0001-7

3. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics [Internet]. 2018

[cited 2025 Apr 15];34:3094–100. Available from:

https://dx.doi.org/10.1093/bioinformatics/bty191

4. Smolka M, Paulin LF, Grochowski CM, Horner DW, Mahmoud M, Behera S, et al. Detection

of mosaic and population-level structural variants with Sniffles2. Nature Biotechnology

2024 42:10 [Internet]. 2024 [cited 2025 Apr 15];42:1571–80. Available from:

https://www.nature.com/articles/s41587-023-02024-y

5. Keskus A, Bryant A, Ahmad T, Yoo B, Aganezov S, Goretsky A, et al. Severus: accurate

detection and characterization of somatic structural variation in tumor genomes using

long reads. medRxiv [Internet]. 2024 [cited 2025 Apr 15];2024.03.22.24304756.

Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC10996739/

6. Buck D, Weirather JL, de Cesare M, Wang Y, Piazza P, Sebastiano V, et al. Comprehensive

comparison of Pacific Biosciences and Oxford Nanopore Technologies and their

applications to transcriptome analysis. F1000Res [Internet]. 2017 [cited 2025 Jun

13];6:100. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC5553090/

7. Eccles D, White R, Pellefigues C, Ronchese F, Lamiable O. Investigation of chimeric reads

using the MinION. F1000Res [Internet]. 2017 [cited 2025 Jun 13];6:631. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/PMC5600009/

8. Yang C, Lo T, Nip KM, Hafezqorani S, Warren RL, Birol I. Characterization and simulation of

metagenomic nanopore sequencing data with Meta-NanoSim. Gigascience [Internet].

2022 [cited 2025 Jun 13];12:1–11. Available from:

https://dx.doi.org/10.1093/gigascience/giad013

9. Stock W, Rousseau C, Dierickx G, D'hondt S, Amadei Martínez L, Dittami SM, et al.

Breaking free from references: a consensus-based approach for community profiling with

long amplicon nanopore data. Brief Bioinform [Internet]. 2024 [cited 2025 Jun 13];26.

Available from: https://dx.doi.org/10.1093/bib/bbae642

10. Li Y, Wang T-Y, Guo Q, Ren Y, Lu X, Cao Q, et al. A Genomic Language Model for Chimera

Artifact Detection in Nanopore Direct RNA Sequencing. bioRxiv [Internet]. 2024 [cited

2025 Jun 13];2024.10.23.619929. Available from:

https://pmc.ncbi.nlm.nih.gov/articles/PMC11526916/

11. Soneson C, Yao Y, Bratus-Neuenschwander A, Patrignani A, Robinson MD, Hussain S. A

comprehensive examination of Nanopore native RNA sequencing for characterization of

complex transcriptomes. Nature Communications 2019 10:1 [Internet]. 2019 [cited 2025

Apr 17];10:1–14. Available from: https://www.nature.com/articles/s41467-019-11272-z

12. Li Y, Roberts ND, Wala JA, Shapira O, Schumacher SE, Kumar K, et al. Patterns of somatic

structural variation in human cancer genomes. Nature 2020 578:7793 [Internet]. 2020

[cited 2025 Apr 17];578:112–21. Available from:

https://www.nature.com/articles/s41586-019-1913-9

13. Talevich E, Tseng E, Diallo A, Sellami N, Elliott A, Cantarel BL, et al. Standardized

Nomenclature and Reporting for PacBio HiFi Sequencing and Analysis of rAAV Gene

Therapy Vectors. bioRxiv [Internet]. 2024 [cited 2025 Jun 11];2024.05.07.592296.

Available from: https://www.biorxiv.org/content/10.1101/2024.05.07.592296v1

14. Cotto KC, Feng YY, Ramu A, Richters M, Freshour SL, Skidmore ZL, et al. Integrated

analysis of genomic and transcriptomic data for the discovery of splice-associated

variants in cancer. Nature Communications 2023 14:1 [Internet]. 2023 [cited 2025 Apr

17];14:1–18. Available from: https://www.nature.com/articles/s41467-023-37266-6

15. Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al.

Integrative Genomics Viewer. Nat Biotechnol [Internet]. 2011 [cited 2025 Apr 18];29:24.

Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC3346182/

16. Rautiainen M, Nurk S, Walenz BP, Logsdon GA, Porubsky D, Rhie A, et al. Telomere-to-

telomere assembly of diploid chromosomes with Verkko. Nat Biotechnol [Internet]. 2023

[cited 2025 Jun 8];41:1474–82. Available from: https://www.nature.com/articles/s41587-

023-01662-6

17. Liu Y, Mi Y, Mueller T, Kreibich S, Williams EG, Van Drogen A, et al. Multi-omic

measurements of heterogeneity in HeLa cells across laboratories. Nat Biotechnol

[Internet]. 2019 [cited 2025 May 26];37:314–22. Available from:

https://www.nature.com/articles/s41587-019-0037-y

18. Chen S. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication

using fastp. iMeta [Internet]. 2023 [cited 2025 May 17];2:e107. Available from:

/doi/pdf/10.1002/imt2.107

19. rrwick/Porechop: adapter trimmer for Oxford Nanopore reads [Internet]. [cited 2025 May

17]. Available from: https://github.com/rrwick/Porechop

20. epi2me-labs/pychopper: cDNA read preprocessing [Internet]. [cited 2025 May 17]. Available

from: https://github.com/epi2me-labs/pychopper

21. Schuster J, Ritchie ME, Gouil Q. Restrander: rapid orientation and artefact removal for long-

read cDNA data. NAR Genom Bioinform [Internet]. 2023 [cited 2025 May

17];5:lqad108. Available from: https://pmc.ncbi.nlm.nih.gov/articles/PMC10748469/

22. Devine SE, Chissoe SL, Eby Y, Wilson RK, Boeke JD. A Transposon-Based Strategy for

Sequencing Repetitive DNA in Eukaryotic Genomes. Genome Res [Internet]. 1997 [cited

2025 Jun 9];7:551–63. Available from: http://genome.cshlp.org/content/7/5/551.full

23. Tanaka H, Tapscott SJ, Trask BJ, Yao MC. Short inverted repeats initiate gene amplification

through the formation of a large DNA palindrome in mammalian cells. Proc Natl Acad

Sci U S A [Internet]. 2002 [cited 2025 Jun 9];99:8772–7. Available from:

/doi/pdf/10.1073/pnas.132275999?download=true

24. Chen Y, Davidson NM, Wan YK, Yao F, Su Y, Gamaarachchi H, et al. A systematic

benchmark of Nanopore long-read RNA sequencing for transcript-level analysis in

human cell lines. Nature Methods 2025 22:4 [Internet]. 2025 [cited 2025 Jun 10];22:801–

12. Available from: https://www.nature.com/articles/s41592-025-02623-4

25. Sessegolo C, Cruaud C, Da Silva C, Cologne A, Dubarry M, Derrien T, et al. Transcriptome

profiling of mouse samples using nanopore sequencing of cDNA and RNA molecules.

Sci Rep [Internet]. 2019 [cited 2025 Jun 2];9:1–12. Available from:

https://www.nature.com/articles/s41598-019-51470-9

26. Genome in a Bottle Data Release 2025.01 | EPI2ME Blog [Internet]. [cited 2025 Jun 10].

Available from: https://epi2me.nanoporetech.com/giab-2025.01/

27. Zheng Z, Yu X, Chen L, Lee Y-L, Xin C, On A, et al. Clair3-RNA: A deep learning-based

small variant caller for long-read RNA sequencing data. [cited 2025 Jun 10]; Available

from: https://doi.org/10.1101/2024.11.17.624050

28. Datasets - PacBio - Highly accurate long-read sequencing [Internet]. [cited 2025 Jun 10].

Available from: https://www.pacb.com/connect/datasets/

29. Nurk S, Koren S, Rhie A, Rautiainen M, Bzikadze A V., Mikheenko A, et al. The complete

sequence of a human genome. Science (1979) [Internet]. 2022 [cited 2025 Apr

17];376:44–53. Available from: https://www.science.org/doi/10.1126/science.abj6987

30. Rhie A, Nurk S, Cechova M, Hoyt SJ, Taylor DJ, Altemose N, et al. The complete sequence

of a human Y chromosome. Nature 2023 621:7978 [Internet]. 2023 [cited 2025 Apr

17];621:344–54. Available from: https://www.nature.com/articles/s41586-023-06457-y

31. Waterston RH, Lindblad-Toh K, Birney E, Rogers J, Abril JF, Agarwal P, et al. Initial

sequencing and comparative analysis of the mouse genome. Nature [Internet]. 2002 [cited

2025 Jul 8];420:520–62. Available from: https://pubmed.ncbi.nlm.nih.gov/12466850/

32. Morales J, Pujar S, Loveland JE, Astashyn A, Bennett R, Berry A, et al. A joint NCBI and

EMBL-EBI transcript set for clinical genomics and research. Nature 2022 604:7905

[Internet]. 2022 [cited 2025 Jun 21];604:310–5. Available from:

https://www.nature.com/articles/s41586-022-04558-8

33. Loman NJ, Quick J, Simpson JT. A complete bacterial genome assembled de novo using only

nanopore sequencing data. Nature Methods 2015 12:8 [Internet]. 2015 [cited 2025 Jun

11];12:733–5. Available from: https://www.nature.com/articles/nmeth.3444