Title: An Autoimmunity-associated allele of *PTPN22* enhances innate antiviral immunity to

protect against acute coronavirus infection

Authors: Alec M. Bevis[1], Kathryn J.L.H. Rosa[1], Nancy Schwarting[1], Tammy R. Cockerham[1],

Catherine M. Kerr[1], Sunil More[2], Anthony R. Fehr[1], Robin C. Orozco[1,*]

Affiliations:

[1] Department of Molecular Biosciences, University of Kansas; Lawrence, Kansas, 66045.

[2] Department of Veterinary Pathobiology, Oklahoma State University; Stillwater, Oklahoma,

74078.

*Corresponding author. Email: orozco.robin@ku.edu. Phone number: (785)-864-2602. Address:

1200 Sunnyside Ave, Room 8035, Lawrence, KS 66045.

Preprint Server:

A preprint of this manuscript was uploaded to BioRxiv on July 18[th], 2025. The link to this

preprint is pasted below:

https://www.biorxiv.org/cgi/content/short/2025.07.15.664967v1

Classifications:

Major – Biological Sciences

Minor – Immunology and Inflammation

Key Words:

PTPN22, coronavirus, Natural Killer (NK) cells

Abstract:

Allelic variation can impact viral clearance and disease severity. Still, our understanding of the effect of the autoimmunity-associated allelic variant of *Ptpn22* (PEP-R619W) on antiviral immunity remains incomplete, as previous reports have only focused on chronic Lymphocytic Choriomeningitis virus (LCMV) infection. This research defines how the loss of *Ptpn22* (PEP-null) and PEP-R619W changes antiviral immunity during an acute coronavirus infection. We address the hypothesis that CRISPR/Cas9-generated PEP-null and PEP-R619W mice have enhanced antiviral immunity over PEP-WT mice during coronavirus infection. Following Mouse Hepatitis Virus (MHV) A59 infection, we interrogated pathology, cytokine production, and cellular responses in the blood, spleen, and liver of PEP-WT, PEP-null, and PEP-R619W mice. Key findings show that PEP-R619W mice have reduced viral titer and weight loss, increased survival, and more mature natural killer (NK) cells in the liver and spleen compared to PEP-WT mice. Interestingly, protection against disease in PEP-null mice was inoculation-dose-dependent, whereas PEP-R619W conferred immunity regardless of infection dose. Further, Rag1-/- PEP-R619W mice had increased survival and reduced viral titer over Rag1-/- PEP-WT mice. PEP-R619W mice also had higher concentrations of IFNγ and enhanced IFNγ production by mature NK cells in the liver at 3 days post-infection. Finally, NK cell depletion elevated PEP-R619W viral titer to similar levels as PEP-WT mice. This is one of the first studies investigating the role of *Ptpn22* within NK cells and demonstrates that the *Ptpn22* allelic variant augments NK cell function and is beneficial during coronavirus infection.

Significance Statement:

Approximately 5-15% of the North American population has the *PTPN22* 1858C>T allele, which has been linked with numerous autoimmune diseases and is considered the highest non-

HLA risk allele for autoimmunity. Due to this link, the *PTPN22* 1858C>T allele is often considered pathologic or detrimental. However, recent studies have demonstrated the benefit of this allele in protection from chronic virus infection and some cancers. Yet, a significant research gap remains in understanding how the *PTPN22* 1858C>T allele impacts the immune response during acute, moribund infections. Using mice that are homozygous for the equivalent, autoimmunity-associated allele in *Ptpn22*, we demonstrate that this allele uniquely augments innate immunity and enhances Natural Killer cell function to protect against coronavirus infection.

Main Text:

INTRODUCTION

Allelic variation in immune-regulatory genes is associated with protective or pathogenic immune responses during disease (1, 2). An example is *PTPN22* and its minor 1858C>T allele (rs2476601) (1-10). This *PTPN22* minor allele is strongly associated with multiple autoimmune diseases, has been cited as the highest non-HLA risk allele for autoimmunity, and is present in 5-15% of North Americans (1-10). The generational persistence of this *PTPN22* allele, despite its strong association with various severe autoimmune diseases, poses an evolutionary paradox. A theory is that the heightened immune responses it drives, though detrimental in autoimmunity, may offer protection against infections and cancer.

PTPN22 encodes for the protein Lyp in humans and its ortholog PEP in mice. In humans and mice, *PTPN22* is expressed constitutively and exclusively within bone marrow-derived cells,

which include all immune cells (3, 4). The *PTPN22* 1858C>T allele results in an amino acid

substitution of an arginine (R) to tryptophan (W) that alters immune-regulatory function in both

human (Lyp-R620W) and murine ortholog (PEP-R619W) models (3-6, 9, 11-21). To better

understand the function of *Ptpn22*, prior studies have frequently employed *Ptpn22* knockout

(PEP-null) mice. The molecular function of PEP has been best characterized as a negative

regulator of T-cell receptor (TCR) and B-cell receptor (BCR) signaling, but its role in other

immune cells is less well understood (3, 4, 11, 13, 16, 22-27). Further, the Lyp-R620W and PEP-

R619W mutations do not directly impact phosphatase activity but lead to a loss of binding

capacity with other proteins, such as Csk, TRAF3, Vav, Grb2, SKAP-HOM, and NLRP3,

resulting in altered immune function (3, 4, 11-13, 17, 20, 21, 24, 28). Studies have defined

several mechanisms in which the *PTPN22* 1858C>T allele potentiates heightened immune

responses (3, 4, 11, 13, 16-21, 23, 24, 29). These same mechanisms that are detrimental during

autoimmunity, such as enhanced T cell function, are protective when combating chronic viral

infection and cancer (18, 19).

Previous studies show that both PEP-null and PEP-R619W mice, unlike wildtype (PEP-WT)

mice, can clear chronic lymphocytic choriomeningitis virus clone 13 (LCMV-cl13) infection,

and this phenotype is attributed to their enhanced antiviral T cell effector function (19, 30-32).

However, the role of PEP and its PEP-R619W variant in virus infection has mostly been studied

in the context of LCMV infection and T cells (19, 27, 30-33). Consequently, much remains

unknown regarding the impact of PEP and its PEP-R619W variant on innate antiviral immunity,

as well as whether PEP-null and PEP-R619W mice will exhibit similar immune responses during

non-LCMV infection models. Despite the comparable immune phenotypes of PEP-null and PEP-

R619W mice in LCMV-cl13 infection, other studies have highlighted several cellular and

molecular differences between these two strains. For example, naïve PEP-null mice have an

increased amount of Tregs, whereas PEP-R619W do not, compared to PEP-WT mice (26).

Additionally, NLRP3-mediated IL-1β production is increased in PEP-R619W myeloid cells, but

decreased in PEP-null myeloid cells, compared to PEP-WT cells (20, 21). These findings

provide additional rationale to investigate differential responses between these strains,

specifically within the context of viral infection. Therefore, we investigated whether PEP-null

mice, PEP-R619W mice, or both can modulate innate antiviral immunity to confer protection

during acute coronavirus infection.

In this study, we investigated the impact of *Ptpn22* (PEP) and its *Ptpn22* 1858C>T allele (PEP-

R619W variant) on the innate immune response following acute coronavirus infection using the

moribund, highly liver-tropic, mouse hepatitis virus (MHV) strain A59. Our findings show that

PEP-R619W enhances natural killer (NK) cell maturation and strengthens NK effector functions,

and that NK cells are necessary to restrict viral replication in the liver during innate time points

in PEP-R619W mice. The results of this research define new mechanisms by which the

autoimmunity-associated allele of *PTPN22* augments innate antiviral immunity and enhances

protection against coronavirus infection.

RESULTS

PEP-R619W reduces disease and viral titer during MHV A59 infection

To determine the impact of *Ptpn22* and its autoimmunity-associated minor allele (PEP-R619W) during acute coronavirus infection, we infected wildtype (PEP-WT) and CRISPR-Cas9 generated *Ptpn22* knockout (PEP-null) and PEP-R619W mice with a common murine strain of coronavirus, mouse hepatitis virus (MHV) strain A59, which primarily infects the liver. In accordance with previous studies in the literature, MHV A59 infection did not cause severe disease following infection in 8-week-old mice, regardless of PEP-genotype (Fig. S1A and 1B) (34-40). Following 5e4 PFU MHV A59 infection, 5-week-old PEP-R619W and PEP-null mice had significantly reduced weight loss at day 5 post-infection (dpi) and 3-5 dpi, respectively, and this was paired with enhanced survival compared to PEP-WT mice (Fig. 1A and 1C). We next examined if the protection observed in PEP-null and PEP-R619W mice would persist at a higher inoculation dose, as early studies utilizing MHV A59 noted that higher viral input doses lead to more severe clinical symptoms (40). Following 1e5 PFU MHV A59 infection, PEP-R619W mice had significantly reduced weight loss at 3-6 dpi and enhanced survival compared to both PEP-WT and PEP-null mice (Fig. 1B and 1D). These data indicate that during 1e5 PFU infection, PEP-null mice are no longer protected from MHV A59 disease and instead phenocopy PEP-WT mice in weight loss and survival (Fig. 1A-D). To study the replication kinetics of MHV A59 *in vivo*, we quantified the viral load of the whole liver following 5e2 PFU infection, as it is the primary site of replication for MHV A59, at 1-, 3-, 5-, and 7- dpi via plaque assay. This dose was chosen as it has similar replication kinetics to higher inoculation doses, but does not induce severe clinical symptoms of weight loss and death, allowing for titer quantification that is not influenced by differences in survival at later timepoints (41, 42). PEP-R619W mice had significantly reduced viral titer at 1-, 3-, and 5- dpi compared to PEP-WT mice (Fig. 1E). However, PEP-null mice only demonstrated reduced titer at 5 dpi (Fig. 1F). Additionally, both

PEP-R619W and PEP-null mice had accelerated clearance compared to PEP-WT mice, as 7/7

PEP-R619W mice and 7/8 PEP-null mice had cleared the infection by 7 dpi, whereas only 2/7

PEP-WT mice had cleared the infection at this time point (Fig. 1E and 1F). Splenic titers showed

a low level of infection at 1-, 3-, and 5- dpi, regardless of genotype (Fig. S1C and S1D). All mice

were able to clear the MHV A59 infection from the spleen by 7 dpi (Fig. S1C and S1D).

To determine if the reduced viral titer observed in the livers of PEP-R619W and PEP-null mice

(Fig. 1E and 1F) was due to PEP-driven myeloid cell-intrinsic changes in these mice, resulting in

decreased viral replication, we differentiated bone-marrow-derived macrophages (BMMs) and

bone-marrow-derived dendritic cells (BMDCs). We infected them with MHV A59 *ex vivo*, as

they are both readily infected *in vivo* by MHV (Fig. S1E) (43-45). We did not detect any

significant differences in viral titer across the three genotypes in either BMMs (Fig. S1F) or

BMDCs (Fig. S1G), indicating that PEP-R619W and PEP-null do not mediate protection by

directly impacting viral replication through cell-intrinsic mechanisms. Histopathologic analysis

of liver sections at 3 dpi following 1e5 PFU infection demonstrated multifocal regions of marked

hepatocellular necrosis in PEP-WT mice, characterized by infiltration of inflammatory cells and

deposition of edema and fibrin (black arrow) (Fig. 1G and Fig. S1H). In comparison, PEP-

R619W and PEP-null mice exhibited markedly reduced hepatic injury, with mild necrosis and

inflammation (Fig. 1G and Fig. S1H). These combined results show that PEP-R619W mice have

improved disease outcomes and reduced viral burden, which includes accelerated clearance,

during MHV A59 infection.

PEP-R619W prolongs the survival of infected mice in the absence of T and B cells

Due to the enhanced protection against disease conferred by PEP-R619W at early days post-infection (Fig. 1), we wanted to determine if a strengthened innate immune response was sufficient to fully protect these mice, or if T and B cells were necessary for survival during MHV A59 infection. To do this, we infected Rag1-/- PEP-WT, Rag1-/- PEP-null, and Rag1-/- PEP-R619W mice with 5e4 PFU or 1e5 PFU MHV A59. Like the immune-competent mice, Rag1-/- PEP-R619W mice had noticeably less weight loss throughout most of the early stages of the infection (Fig. 2A and 2B) and mice survived longer (Fig. 2C and 2D) compared to Rag1-/- PEP-WT mice at both 5e4 and 1e5 PFU MHV A59 infection. Rag1-/- PEP-null mice also have prolonged survival over Rag1-/- PEP-WT at 5e4 PFU infection (Fig. 2C). However, at 1e5 PFU infection, PEP-null mice no longer had prolonged survival (Fig. 2D). These data also show that T and B cells are necessary for overall survival, regardless of PEP- genotype or MHV A59 dose, as all mice on a Rag1-/- background eventually succumb to disease (Fig. 2C and 2D). Next, we measured viral load in the liver at 3- and 5-dpi following 5e2 PFU MHV A59 infection. Rag1-/- PEP-R619W mice had reduced viral burden compared to Rag1-/- PEP-WT mice (Fig. 2E). No difference was detected between Rag1-/- PEP-WT and Rag1-/- PEP-null mice liver titers (Fig. 2F). There was no difference detected in splenic viral titers at 3 and 5 dpi between Rag1-/- PEP-WT and Rag1-/- PEP-R619W mice or Rag1-/- PEP-WT and Rag1-/- PEP-null mice (Fig. S2). Overall, these data suggest that PEP-null enhances the innate immune response during 5e4 PFU infection in both immune competent and Rag1-/- models to improve survival compared to PEP-WT mice, and these results are comparable to PEP-R619W mice (Figs. 1 and 2). However, only PEP-R619W mice showed enhanced protection following 1e5 PFU infection, suggesting that this autoimmune-associated variant modifies the innate response through additional mechanisms

compared to PEP-null, and this confers increased resistance against MHV A59-associated

disease (Figs. 1 and 2).

PEP-R619W mice have increased innate immune presence in the liver and spleen during

MHV A59 infection

We next aimed to determine how the PEP-R619W variant augmented the innate immune

response. We focused on 3 dpi because 1) this is when we can consistently observe a divergence

in weight loss between genotypes, 2) PEP-R619W mice have reduced liver titer and pathology at

this time point, and 3) this is a time point at which the innate response dominates, whereas at 5

dpi, there could be some early T cell activity. We first investigated the innate cellular response in

the spleen after 5e4 PFU MHV A59 infection, as we wanted to determine if the immune

response in PEP-null and PEP-R619W mice would correlate with the improved weight loss and

survival phenotypes observed at this dose. Overall, PEP-R619W and PEP-null mice had

comparable immune phenotypes at 3 dpi in the spleen post-infection (Fig. S3). There was no

difference detected in the bulk number of splenocytes (Fig. S3B) or frequency of CD3e- CD19-

(innate cells) of all live cells (Fig. S3C), regardless of infection or PEP genotype. However, PEP-

R619W and PEP-null mice had significantly increased frequencies of NK cells and DCs

compared to infected PEP-WT mice (Fig. S3D).

Interestingly, PEP-R619W mice had more splenic CD11b+ NK cells than PEP-WT, indicating

PEP-R619W NK cells are significantly more mature than PEP-WT NK cells following infection

(Fig. S3E) (46, 47). Further, there were no differences detected in the frequency of CD8α+

conventional dendritic cells (CD8α+ cDCs), which are likely type 1 conventional dendritic cells

(cDC1s) (Fig. S3F) (48). While the expression of activation markers CD80 and CD86 was

significantly upregulated post-infection in splenic CD8α+ cDCs, no significant difference was

detected between infected genotypes (Fig. S3G and S3H, respectively). Similar results were seen

in splenic CD8α- cDCs, which are likely type 2 conventional dendritic cells (cDC2s) (Fig. S3I-

K), though there was a decrease in CD86 expression in PEP-null and PEP-R619W splenic

CD8α- cDCs compared to PEP-WT post-infection (Fig. S3K) (48).

Next, we wanted to know if PEP-R619W mice had an altered immune response in the liver, the

main site of viral replication, during the 1e5 PFU MHV A59 infection, as they were the only

genotype protected at the inoculation dose (Figs. 1 and 2). At 3 dpi, we did not detect significant

differences in the absolute number of liver cells (Fig. 3A) or in the frequency (%) of CD45+ cells

(Fig. 3B) between our mock mice (uninfected controls) and our infected PEP-WT, PEP-null, and

PEP-R619W mice. However, PEP-R619W mice have a higher proportion of CD3e- CD19- of

CD45+ immune cells, which we define as an immune "innate subset," compared to PEP-WT

mice post-infection (Fig. 3C). Delving further into this innate cell population, PEP-R619W mice

had increased proportions of natural killer (NK) cells, dendritic cells (DCs), and macrophages of

CD45+ cells over both PEP-WT and PEP-null mice post-infection (Fig. 3D). We did not detect

any differences between infected genotypes in plasmacytoid DCs (pDCs), monocytes, or

neutrophils (Fig. 3D). We also did not observe any differences in the proportion of or CD86

expression on Tim4+ macrophages, which are likely Kupffer cells amongst infected PEP-WT,

PEP-null, and PEP-R619W mice (Fig. S4A-C) (49-51). Post-infection, all genotypes increased

CD86 expression on Tim4+ macrophages (Fig. S4C). Within the DC population, there was a

decrease in both the frequency and expression of MHC-II (Fig. S4D and S4E, respectively) in

infected genotypes compared to uninfected controls (mock). Further, we did not identify any

differences in the frequency and expression of CD86 on DCs between infected genotypes (Fig.

S4F and S4G).

MHV A59-infected PEP-R619W mice had more conventional NK cells compared to infected

PEP-WT and PEP-null mice, characterized as CD49b+ CD49a- of NK1.1+ cells, whereas the

inverse is considered tissue-resident type 1 innate lymphoid cells (ILC-1) (Fig. 3E) (52, 53).

Additionally, PEP-R619W conventional NK cells were more mature (CD11b+) compared to

PEP-WT (Fig. 3F), and overall, PEP-R619W mice had more mature conventional NK cells

(CD11b+ CD49b+ CD49a- NK1.1+) compared to both PEP-WT and PEP-null mice during

coronavirus infection (Fig. 3G). These mature NK cells are of particular interest because, as NK

cells mature, they enhance their effector function potential, including increased interferon-

gamma (IFNγ) production and elevated expression of cytotoxic molecules, such as perforin and

granzyme B (46, 47). Collectively, these data demonstrate that the PEP-R619W variant enhances

both splenic and liver innate antiviral immunity, specifically increasing the overall amount and

maturity of NK cells.

PEP-null splenic mature NK cell effector functions are changed depending on the MHV

A59 inoculation dose

Since our data showed that PEP-R619W mice have significantly more mature NK cells in both

the liver and spleen at 3 dpi, we next wanted to determine if the NK cells amongst our mice had

changed functional properties, and if the inoculation dose of MHV A59 would impact these

phenotypes. Again, we observed inoculation dose-dependent differences, where at 5e4 PFU,

PEP-null and PEP-R619W mice had similar NK cell immune phenotypes in the spleen (Fig.

S5A-F). Specifically, following 5e4 PFU infection, PEP-null and PEP-R619W mature splenic

NK cells had less CD107a, a marker of degranulation (Fig. S5A and S5B), increased perforin

(Fig. S5C and S5D), and no detectable difference in IFNγ (Fig. S5E and S5F) compared to PEP-

WT mature splenic NK cells. Yet at 1e5 PFU, PEP-null and PEP-WT phenotypes were more

similar in the spleen (Fig. S5G-L). With 1e5 PFU infection, PEP-R619W mature splenic NK

cells had reduced degranulation marker CD107a (Fig. S5G and S5H), increased perforin (Fig.

S5I and S5J), and an increase in frequency, but not expression, of IFNγ (Fig. S5K and S5L)

compared to PEP-WT and PEP-null mature splenic NK cells. These data provide some insight

into why PEP-null mice have differing phenotypes (survival) during different doses of MHV

A59 infection, as differential immune activation at 1e5 PFU infection may be responsible for the

previously described increase in morbidity (Figs. 1 and 2).

PEP-R619W enhances liver mature NK cell effector function and the overall production of

IFNγ during virus infection

As only PEP-R619W mice, but not PEP-null, had more mature conventional NK (M-cNK) cells

compared to PEP-WT mice at 3 dpi in the liver, we next investigated if PEP-R619W M-cNK

cells have altered effector functions compared to PEP-WT M-cNK cells. Following 1e5 PFU

infection, at 3 dpi in the liver, PEP-R619W M-cNK cells (CD11b+ CD49b+ CD49a- NK1.1+)

have higher expression of (i) IFNγ, (ii) T-bet, (iii) granzyme B, and (iv) perforin, compared to

PEP-WT M-cNK cells (Fig. 4A, i-iv). There was no difference detected in the expression of (v)

TNFα or (vi) CD107a between PEP-WT and PEP-R619W M-cNK cells (Fig. 4A, v-vi). Further,

PEP-R619W M-cNK cells were more poly-functional compared to PEP-WT, as there was an

increased proportion of IFNγ+ Perforin+ and IFNγ+ Perforin+ TNFα+ M-cNK cells in PEP-

R619W mice.

Next, we examined the magnitude of the cytokine/chemokine response both systemically and in

the whole liver at 3 dpi. Strikingly, out of the thirteen different cytokines and chemokines

paneled in the serum (Fig. 4C), only the concentration of IFNγ was significantly increased in

PEP-R619W mice (Fig. 4D), yet all other molecules, including interferon-alpha (IFNα) (Fig.

4E), which is necessary for protection against MHV A59 infection, were unchanged between

infected genotypes (54). Similar results were detected in the whole liver homogenate (Fig. 4F-

H). To corroborate these findings, we measured the levels of IFNβ in the liver at 3 dpi in our

PEP-WT, PEP-null, and PEP-R619W mice. We detected no difference between infected

genotypes (Fig. 4I).

Given the enhanced M-cNK IFNγ expression and the overall increased IFNγ concentration in our

PEP-R619W mice, we next wanted to determine if IFNγ and/or IFNγ-signaling was necessary

for survival and weight retention during MHV A59 infection. To test this, we infected WT, IFNγ

knockout (IFNγ KO), and IFNγ receptor knockout (IFNγR KO) mice with 5e4 PFU of MHV

A59. Both IFNγ KO and IFNγR KO mice had more severe weight loss (Fig. 4J) and nearly 100%

mortality (Fig. 4K) compared to WT mice, demonstrating the importance of IFNγ for protection

during MHV A59 infection.

Taken together, these data show that PEP-R619W liver M-cNK cells produce more IFNγ and are

more poly-functional than PEP-WT M-cNK cells, contributing to the increased levels of IFNγ in

both the serum and liver at 3 dpi. Further, IFNγ and IFNγ signaling are critical for survival. This

suggests that the protection conferred by PEP-R619W during MHV A59 infection may be

driven, in part, by enhanced NK cell IFNγ production.

NK Cell maturation is accelerated in PEP-null and PEP-R619W mice

Due to the increased presence of mature NK cells in the spleens (CD11b+ of NK1.1+) and liver

(CD11b+ of CD49b+ CD49a- NK1.1+) of PEP-R619W mice post-infection (Fig. S3E, Fig. 3E-

G), we examined if PEP-R619W NK cells had differential maturation states compared to PEP-

WT at homeostasis. Previous studies have shown that murine NK cells follow a four-stage

developmental program, defined by the subsequent acquisition and loss of surface receptors

CD27 and CD11b, and these maturation states can be seen in the spleen (46, 47). The four

stages, from least to most mature, are defined as; (i) CD27- CD11b- (Double Negative, DN), (ii)

CD27+ CD11b-, (iii) CD27+ CD11b+ (Double Positive, DP), and (iv) CD27- CD11b+ (46, 47).

This developmental program is associated with the stepwise acquisition of NK cell effector

function, and the accumulation of terminally mature NK cells (CD27- CD11b+) plateaus at 8

weeks of age (46, 47). At 5 weeks of age, PEP-null and PEP-R619W splenic NK cells had

differential maturation states compared to PEP-WT (Fig. 5). Specifically, of the NK cell

repertoire, PEP-WT mice had more DP NK cells, but PEP-null and PEP-R619W had increased

proportions of terminally mature (CD27- CD11b+) NK cells at 5 weeks of age (Fig. 5A, 5D, 5E).

However, at 8 weeks of age, there were no differences detected in the proportion of DP or

terminally mature NK cells between the three genotypes of mice. Collectively, these data

indicate that PEP may negatively impact NK cell maturation at homeostasis, which could play a

role in the observed differences in mature NK cell presence and function in PEP-R619W mice

following MHV A59 infection.

PEP-R619W NK cells restrict viral replication and enhance IFNγ production during

coronavirus infection

Given our observation that NK cells are more mature and functional, and there are more of them

in PEP-R619W mice during infection, we next wanted to determine if NK cells were necessary

for driving disease protection during MHV A59 infection. To test this, we administered an anti-

NK1.1 (α-NK1.1) depleting antibody (clone PK136) or anti-IgG isotype control (α-Iso) two days

before MHV A59 infection. Subsequent injections were administered every four days after the

initial treatment (Fig. 6A). NK depletion was verified at all key time points, both within the

spleen and in the peripheral blood (Fig. S6A-E). Loss of NK cells in PEP-R619W mice resulted

in a decreased capacity to control virus infection, as viral burden in the liver at 3- and 5- dpi rose

to comparable levels of both α-NK1.1 PEP-WT and isotype-treated PEP-WT mice, and were

significantly higher than isotype-treated PEP-R619W mice at these time points (Fig. 6B). In

addition, isotype-treated PEP-R619W mice maintained the reduced levels of viral titer in the

liver compared to isotype-treated PEP-WT mice at these time points (Fig. 6B). These data

indicate that PEP-R619W NK cells effectively reduce liver viral load, but PEP-WT NK cells

have minimal impact on combating viral replication within the liver. Similar results were

detected in the splenic titers (Fig. S6F). These data suggest that in PEP-R619W mice, the

increased presence of NK cells in the liver is one of the key innate cells responsible for

combating viral replication during the early days of infection. Our data support this conclusion,

as all NK-sufficient PEP-R619W mice, including immune-competent (Fig. 1E), Rag1-/- (Fig.

2E), and α-Iso (Fig. 6B) mice, had significantly reduced liver viral titers at days 3 and 5 post-

infection compared to PEP-WT counterpart mice. In contrast, this protection was lost in α-NK1.1

PEP-R619W mice (Fig. 6B). Next, we wanted to determine if NK cells were a critical source of

IFNγ in our PEP-R619W mice. Upon depletion of NK cells, the levels of IFNγ in liver

homogenate at 3 dpi in α-NK1.1 PEP-R619W mice were significantly reduced compared to α-Iso

PEP-R619W mice (Fig. 6C). Further, α-Iso PEP-R619W had a significantly higher concentration

of IFNγ compared to α-Iso PEP-WT, recapitulating the cytokine profiling data previously shown

(Fig. 4G). In conclusion, these data highlight that PEP-R619W enhances NK cell function to

protect against MHV A59 viral replication in the liver through, in part, boosting the production

of NK cell-derived IFNγ.

PEP-R619W NK cells do not mediate weight loss, survival, or liver histopathology during

coronavirus infection

Since PEP-R619W NK cells mediate viral titer at early days post-infection and are a key source

of IFNγ, we next wanted to determine if they were necessary for mediating weight loss and

survival during MHV A59 infection. Following 1e5 PFU MHV A59 infection, NK cell depletion

in either PEP-WT or PEP-R619W mice did not result in changes in weight loss, survival, or in

liver histopathology compared to NK-sufficient PEP-WT and PEP-R619W counterparts (Fig.

6D-F and Fig. S6G).

Overall, these data indicate that NK cells in PEP-R619W mice play a specific role in combating

MHV A59 infection by restricting viral replication and are critical producers of IFNγ within the

primary site of replication, the liver. However, these NK cells are dispensable for protection

against weight loss, survival, and liver damage. While NK cell depletion in PEP-R619W mice

increased viral load and reduced liver IFNγ concentration, the same depletion in PEP-WT mice

did not affect liver viral titer or IFNγ production. This disparity can potentially be attributed to

the reduced number and impaired function of M-cNK cells in the livers of PEP-WT mice at 3 dpi

(Fig. 3D-G, Fig. 4A and 4B), making their absence less impactful on viral replication.

DISCUSSION

Despite the strong genetic link between the minor allele of *PTPN22* and the development of

various autoimmune diseases, this allele persists in the human population. A potential theory lies

in the fact that the mechanisms driving heightened immune responses, while pathogenic in the

context of autoimmunity, are likely protective when defending against infections and cancer.

Previous studies support this reasoning, as PEP-R619W mice exhibit enhanced antiviral

immunity during chronic LCMV-cl13 infection and increased antitumor immunity, resulting in

reduced disease severity compared to PEP-WT mice (18, 19). Similar results in PEP-null mice

have been reported during LCMV-cl13 infection (30-32). However, these diseases are chronic.

Thus, we were interested in determining if PEP-null and/or PEP-R619W mice would also have

improved disease outcomes during acute, moribund coronavirus infection, MHV A59.

Our results show that PEP-R619W and PEP-null do not always phenocopy each other.

Regardless of whether PEP-null mice were fully immune-competent or on a Rag1-/- genetic

background, their phenotypes were comparable to PEP-R619W mice after 5e4 PFU infection,

but demonstrated increased susceptibility following 1e5 PFU infection and mirrored the

outcomes observed in PEP-WT mice (Figs. 1 and 2). Despite similar splenic innate profiles and

NK activation between PEP-null and PEP-R619W mice at 3 dpi following 5e4 PFU infection,

after 1e5 PFU MHV A59 infection, PEP-null splenic NK cell phenotypes were comparable to

PEP-WT (Fig. S3 and S5). These findings add to the current literature, highlighting that there are

differences between PEP-null and PEP-R619W. While PEP-null and PEP-R619W mice may be

similar in certain scenarios, such as in LCMV-cl13 antiviral immunity, there are other examples

where they can differ (19-21, 27, 30-32). This may lie in cellular and/or molecular differences

between these strains. For example, previous studies show that PEP-null mice have more Tregs

than PEP-WT and PEP-R619W mice, and that the PEP-R619W mutation does not directly

impact phosphatase activity (3, 4, 9, 23, 24, 26, 27, 29). Instead, the mutation disrupts molecular

interactions with other proteins, such as TRAF3 and Csk, leading to altered immune function (3,

4, 9, 23, 24, 26, 27, 29). The precise ways in which established molecular interactions and other

potential mechanisms that influence innate immune cell function diverge between PEP-null and

PEP-R619W mice are poorly understood. Further investigation into these mechanisms would

provide crucial insight into the varying outcomes observed between these strains during MHV

A59 infection.

Our data shows that PEP-R619W protects against murine coronavirus, MHV A59, associated

disease (Fig. 1). This is paired with reduced liver viral titer at early days post-infection, and

accelerated clearance compared to PEP-WT mice (Fig. 1). This protection in PEP-R619W mice

was partially mediated by an enhanced innate immune response. This is further strengthened by

the results of our studies using Rag1-/- models, as Rag1-/- PEP-R619W mice had reduced

disease compared to Rag1-/- PEP-WT mice (Fig. 2). These data indicate that PEP-R619W

differentially modulates the innate immune response to defend against acute MHV A59 infection

(Figs. 1 and 2). These data are further supported by the immune phenotyping of both the spleen

and liver at 3 dpi, where PEP-R619W mice had more innate immune cells compared to PEP-WT

mice (Fig. 3, Fig. S3 and S4).

We interrogated the immune response in the liver following 1e5 PFU infection to define how

PEP-R619W uniquely enhanced innate antiviral immunity. Notably, PEP-R619W mice had more

NK cells, DCs, and macrophages compared to PEP-WT and PEP-null mice (Fig. 3D). However,

we detected no differences in type-I interferon (IFN-I) production, which is necessary for

survival following MHV A59 infection (54). There were no changes in the pDC population,

which are critical producers of IFN-I (Fig. 3D), or in the systemic and liver homogenate levels of

IFNα and IFNβ (Fig. 4C, E, F, H, and I) between infected genotypes, indicating that PEP-

R619W protects through alternative mechanisms, such as enhancing NK cell function. NK cells

can directly lyse infected cells through the release of perforin and granzyme B, are the primary

producers of IFNγ during the innate immune response, and these functions are enhanced as NK

cells mature (46, 47, 55, 56). These effector functions are necessary for combating murine

cytomegalovirus (MCMV) disease and viral replication and have been specifically shown to be

vital in driving the innate immune response in the liver post-infection (57-61). As such, we more

closely interrogated the NK cell compartment in PEP-WT and PEP-R619W mice post-MHV

A59 infection.

We investigated the role of NK cells during MHV A59 infection and discovered that: 1) there

were more mature conventional NK (M-cNK) cells in the spleen and liver of PEP-R619W mice

(Fig. 3D-G and Fig. S3D, S3E), 2) PEP-R619W enhanced M-cNK effector function and

polyfunctionality compared to PEP-WT (Fig. 4A and 4B), and 3) PEP-R619W NK cells, but not

PEP-WT NK cells, are necessary for restricting viral replication in the liver at 3 and 5 dpi (Fig.

6B) and are a critical source of IFNγ (Fig. 6C). Taken together, these results indicate that PEP-

R619W augments M-cNK cell antiviral function in the liver during MHV A59 infection, leading

to reduced viral replication, but PEP-R619W M-cNK cells do not directly mediate weight loss,

survival, or liver pathology (Fig. 6D-F). These findings add to recent work and highlight the

protective role of NK cells at the primary site of coronavirus infection, as NK cells are critical

for combating SARS-CoV-2 replication in both human and non-human primate lungs (62-64).

However, our data also demonstrates that viral burden and disease symptoms are not always

directly correlated. This has been observed in previous MHV A59 studies, as well as during

chronic LCMV-cl13 infection, where weight loss is driven by CD8 T-cell immunopathology (34,

65).

We are one of the first groups to investigate the function of PEP in NK cells. The molecular role

of PEP in NK cells remains undefined, despite being the immune cells with the highest Lyp/PEP

expression in humans and mice (66-68). Our results show that at 5 weeks of age, PEP-R619W

mice have more mature NK cells post-MHV A59 infection at 3 dpi in the spleen and liver

compared to PEP-WT (Fig. S3 and Fig. 3). This phenotype persists at homeostasis, where both

PEP-null and PEP-R619W mice had a higher proportion of terminally mature (CD11b+ CD27-)

NK cells in the spleen compared to PEP-WT counterparts (Fig. 5). Despite these findings, the

molecular functions of PEP within NK cells, and how the PEP-R619W variant accelerates

maturity and enhances effector functions during infection, remains unknown. These results

warrant additional studies defining the molecular mechanisms by which PEP and PEP-R619W

regulate NK cell maturity and function at homeostasis and during disease.

In conclusion, our results indicate that the autoimmunity-associated allele of *PTPN22* enhances

innate antiviral immunity against acute murine coronavirus infection, MHV A59, and adds to the

growing literature supporting the protective role of this allele during viral and tumorigenic

challenges (18, 19). We have shown a divergence in protection against MHV A59 infection in

PEP-null mice in an inoculation dose-dependent manner, whereas PEP-R619W was protective

regardless of infection dose. Additionally, we showed that PEP-R619W boosts innate cell

infiltration, enhances NK cell maturation and function, and, overall, reduces disease following

murine coronavirus infection. These data highlight that PEP-R619W uniquely modifies immune

function through distinct mechanisms compared to both PEP-WT and PEP-null, especially in NK

cells. These findings provide further rationale for testing the protective capacity of PEP-R619W

in additional infection models and defining how PEP-R619W augments immunity in a context-

dependent manner.

MATERIALS AND METHODS

Ethics Statement

The University of Kansas (KU) Institutional Animal Care and Use Committee (IACUC)

reviewed and approved all animal studies under protocol number AUS 278-01.

Mice

Both male and female mice at 5 weeks of age were used in all studies unless otherwise noted. We do not observe sex-based differences in these studies. C57BL/6J mice and C57BL/6 Rag1-/- mice were originally purchased from Jackson Labs and then bred and maintained in the University of Kansas Animal Care Unit (ACU) facilities. *Ptpn22* knockout (PEP-null) and *Ptpn22* R619W (PEP-R619W) mice were generated using CRISPR/Cas9 technology using methods previously reported (15, 18, 19, 69). PEP-null Rag1-/- and PEP-R619W Rag1-/- mice were derived as previously described (18). IFNγ KO (RRID: IMSR_JAX:002287) and IFNγR KO (RRID: IMSR_JAX:003288) mice were purchased from Jackson Labs and were housed and maintained in the University of Kansas ACU facilities as detailed above.

Cell Culture

HeLa-MHVR and 17Cl-1 Cells:

HeLa cells expressing the MHV receptor CEACAM1a (HeLa-MHVR) were cultured in Advanced Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% FBS, 1% L-glutamine, and 1% Penicillin/Streptomycin, and 17Cl-1 cells were grown in DMEM supplemented with 5% FBS, 5% Tryptose Phosphate Broth, and 2% Penicillin/Streptomycin (70) (71).

Bone Marrow Dendritic Cells (BMDC) and Macrophages (BMM):

To generate BMDCs, bone marrow cells were isolated from PEP-WT, PEP-null, and PEP-R619W mouse femurs. Following isolation, cells were cultured in Advanced DMEM supplemented with 10% FBS, 1% L-glutamine, 1% Penicillin/Streptomycin, and 100ng/mL FLT3L (Stem Cell Technologies, Vancouver, CA). For BMMs, bone marrow cells were isolated

492 and cultured in Roswell Park Memorial Institute medium (RPMI) supplemented with 10% FBS,

493 1% L-glutamine, and 1% Penicillin/Streptomycin with 50ng/mL M-CSF (Stem Cell

494 Technologies, Vancouver, CA) for 8 days. An additional 10 mL of M-CSF-containing media was

495 added to BMM cultures on day 3, and on day 8, differentiated cells were harvested, counted, and

496 replated for further assays (72).

MHV A59 Production, Titer, and Infections

MHV A59 stocks were grown in 17Cl-1 cells, and viral titers were determined on HeLa cells

expressing the MHV receptor CEACAM1a (HeLa-MHVR), as previously described (42, 70, 71,

73, 74). Briefly, HeLa-MHVRs were seeded in 12-well plates the day before infection. The

following day, once the HeLa-MHVRs were grown to 90-95% confluency, 10-fold dilutions of

titer samples were prepared, and the HeLa-MHVRs were inoculated for a 45-to-60-minute

adsorption phase, followed by 1 mL overlay per well of 1:1 mixture of 1.2% Agarose and 2x

DMEM (10% FBS, 1% L-Gln, 1% Pen/Strep). Once solidified, 500 µL D2 media (DMEM, 2%

FBS, 1% L-Gln, 1% Pen/Strep) was added to the overlay. The plaques were counted and

recorded 48 hours later for viral titer (PFU/sample) calculation.

For *in vivo* infections, mice were inoculated with 5e2 PFU, 5e4 PFU, or 1e5 PFU MHV A59 in a

final volume of 200 µL sterile PBS via intraperitoneal (i.p.) injection (PBS alone for "mock"

mice). Throughout all infections, mice were weighed daily and euthanized if weight loss was

greater than 20% of the original starting weight. At peak infection (days 5-8 post-infection), mice

were checked twice daily to provide more frequent monitoring. To determine viral load, the

whole liver and spleen were harvested and homogenized in 5 mL or 1 mL sterile DMEM,

respectively, centrifuged to remove cellular debris, and stored at -80^0C. The viral titer was

determined on HeLa-MHVR cells as detailed above.

For *ex vivo* Bone Marrow Dendritic Cell (BMDC) and Bone Marrow Macrophage (BMM)

infections, cells were inoculated at a multiplicity of infection (MOI) of 0.1 with a 45-to-60-

minute adsorption phase. The viral inoculum was aspirated, and 1 mL of advanced DMEM

supplemented with 10% FBS, 1% L-glutamine, 1% Penicillin/Streptomycin or RPMI

supplemented with 10% FBS, 1% L-glutamine, and 1% Penicillin/Streptomycin, for BMDC or

BMM, respectively, was added per well and incubated for 24 hours. At 24 hours post-infection,

the cells were scraped and resuspended before being placed in the -80^0C freezer overnight.

Following freeze-thaw, the virus from the supernatant and cells was combined and tittered on

HeLa-MHVRs as described above.

Leukocyte Isolation and Flow Cytometry

Mice were euthanized at indicated time points in line with AVMA and KU IACUC guidelines by

CO2 inhalation followed by cervical dislocation before tissue harvest.

The spleen was excised at indicated days post-infection, smashed over 40 µm filters, rinsed with

DMEM, and centrifuged to remove debris. The cell pellet was resuspended in 2 mL ACK red

cell lysis buffer, incubated at RT for 2 minutes, quenched with 10 mL DMEM, and centrifuged.

The resultant pellet was resuspended in 2 mL advanced DMEM supplemented with 10% FBS,

1% L-glutamine, 1% Penicillin/Streptomycin to generate a single cell suspension.

Whole livers were harvested following cardiac perfusion with 15-20 mL sterile PBS and placed

in 3 mL DMEM supplemented with 10% FBS on ice. Livers were digested and processed using a

murine liver dissociation kit (Miltenyi Biotec, Bergisch Gladbach, North Rhine-Westphalia,

Germany) per the manufacturer's protocol. Following centrifugation, the cell pellet was

resuspended in 1 mL sterile DMEM (1% L-glutamine), gently overlaid onto a 20 mL 33%

percoll gradient in PBS, and centrifuged at 800xg for 25 minutes at room temperature (RT) (75).

After aspiration, the cell pellet was resuspended in 2 mL Advanced DMEM (supplemented with

10% FBS, 1% L-glutamine, 1% Penicillin/Streptomycin) to generate a single cell suspension.

To harvest peripheral blood for NK Depletion verification, 50 µL of blood was acquired via

retro-orbital (r.o.) bleeds using heparinized capillary tubes. 10 µL of blood was plated in a 96-

well plate for each sample. 100 µL of ACK red cell lysis buffer was added and incubated at RT

for 2 minutes, quenched with 100 µL Advanced DMEM (supplemented with 10% FBS, 1% L-

glutamine, 1% Penicillin/Streptomycin), centrifuged, and resuspended in 200 µL of media to

generate a single cell suspension.

All single cell suspensions were counted using Vi-CELL BLU (Beckman Coulter, Brea, CA).

The concentration of each sample was adjusted to plate 2e6 cells/well in round-bottom 96-well

plates for antibody staining and flow cytometric analysis.

All flow cytometry was performed on the Cytek Aurora 5 laser system (355 nm [UV], 405 nm

[Vio], 488 nm [Blue], 561 nm [YG], and 640 nm [Red]) spectral flow cytometer (Fremont, CA,

US). Single stain control Biolegend Compensation Beads were used for unmixing (San Diego,

California, U.S). Unmixed files were analyzed using FlowJo Software (BD Biosciences, San

Diego, California). Antibodies used in varying combinations (depending upon the experiment)

are shown below in the Supplementary Materials (Table S1).

All samples were incubated with diluted Fc block (Biolegend) for 10 minutes and then stained

for 30 minutes at 4^0C in the dark for surface marker staining in serum-free PBS. For intracellular

cytokine staining, cells were incubated with brefeldin A (BFA, Thermo Scientific) for four hours

at 37^0C before surface staining. BD Cytofix and Permwash kit or Tonbo Foxp3 Fix/Perm kit

were used according to manufacturers' protocols for intracellular cytokine or intranuclear

transcription factor staining, respectively.

Cytokine and Chemokine Quantification

500 µL of blood was acquired via retro-orbital (r.o.) bleeds using non-heparinized capillary tubes before euthanasia. After incubating at RT for 30 minutes, the blood was centrifuged at 13,000 RPM for 10 minutes at RT, and the serum supernatant was collected and stored at -80^0C for future analysis. As described above, the whole liver was homogenized in 5 mL DMEM, centrifuged to remove cellular debris, and stored at -80^0C for future analysis. Serum and liver homogenate antiviral cytokines and chemokines were quantified using a LEGENDplex assay (Biolegend), with the serum being diluted 1:2 as described in the manufacturer protocol. In a separate experiment, a portion of the left lateral liver lobe was harvested, weighed, homogenized, and the concentration of IFNβ was determined using the Mouse IFN-Beta ELISA Kit (PBL Assay Science). The levels of IFNγ were determined using LEGEND MAX™ Mouse IFNγ ELISA Kit (Biolegend) in NK cell depletion experiments following whole liver harvest and homogenization at indicated time points.

Natural Killer (NK) cell depletion

Depletion of NK cells was achieved through i.p. injection of 100 µg anti-NK1.1 monoclonal antibodies (clone PK136) two days before infection and followed by 25 µg anti-NK1.1 injections every four days to sustain depletion (Bio-X-Cell). In parallel, separate mice were administered control IgG injections at the same dosage and timings as detailed above (Bio-X-Cell). The experimental outline is shown in Supplementary Figure 6.

Histopathology

Whole livers were harvested following cardiac perfusion with 15-20 mL sterile PBS, followed by 10-15 mL of 4% PFA diluted in PBS, and placed in 20 mL of 10% PFA diluted in PBS

(minimum ratio of 1 gram tissue:10 mL 10% PFA). After 72 hours at 4^0C in the dark, a

representative section from each lateral and medial (right and left) lobe was cut and processed

for routine hematoxylin and eosin (H&E) staining. The liver lesions were blindly scored by an

American College of Veterinary Pathology Board-certified pathologist. The lesions scored were

inflammation, necrosis, and edema/fibrin on a scale of 0–10% (score 1), 10–40% (score 2), 40–

70% (score 3), and >70% (score 4). Cumulative scores were obtained for each mouse and

presented as total pathology score.

Statistical Analysis and Graphing

All graphs and appropriate statistical analysis for data were done in GraphPad Prism (La Jolla,

CA). The type of statistical tests used are listed in each figure legend. Data was considered

statistically significant if the p-value<0.05. Data with a p-value>0.05 was determined non-

significant, and pairwise comparisons were either not labeled or represented with "ns." Figure

legends indicate if data are from pooled experiments or are representative of multiple studies.

Most graphs display mean + SEM, unless otherwise stated.

Supplementary Materials

Figure S1: MHV A59 disease is attenuated in 8-week-old mice, and PEP-R619W does not

impact splenic titer or restrict viral replication in myeloid cells.

Figure S2: Splenic titers are unchanged in Rag1-/- mice regardless of PEP genotype.

Figure S3: PEP-R619W increases the splenic innate immune population.

Figure S4: Representative gating strategy for different liver immune cell populations and

additional quantification of immune cells.

Figure S5: Infection dose-dependent impact on splenic mature NK function.

Figure S6: NK depletion protocol and NK-depleted spleen titer and histopathology scores.

Table S1: Flow Cytometry Antibodies

REFERENCES

1.   P. K. Gregersen, B. Diamond, R. M. Plenge, GWAS implicates a role for quantitative
immune traits and threshold effects in risk for human autoimmune disorders. *Curr Opin
Immunol* **24**, 538-543 (2012).
2.   R. J. Xavier, J. D. Rioux, Genome-wide association studies: a new window into immune-
mediated diseases. *Nat Rev Immunol* **8**, 631-643 (2008).
3.   L. H. Armitage, M. A. Wallet, C. E. Mathews, Influence of PTPN22 Allotypes on Innate
and Adaptive Immune Function in Health and Disease. *Front Immunol* **12**, 636618
(2021).
4.   N. Bottini, E. J. Peterson, Tyrosine phosphatase PTPN22: multifunctional regulator of
immune signaling, development, and disease. *Annu Rev Immunol* **32**, 83-119 (2014).
5.   N. Bottini, T. Vang, F. Cucca, T. Mustelin, Role of PTPN22 in type 1 diabetes and other
autoimmune diseases. *Semin Immunol* **18**, 207-213 (2006).
6.   G. L. Burn, L. Svensson, C. Sanchez-Blanco, M. Saini, A. P. Cope, Why is PTPN22 a
good candidate susceptibility gene for autoimmune disease? *FEBS Lett* **585**, 3689-3698
(2011).
7.   R. C. Chiaroni-Clarke *et al.*, The association of PTPN22 rs2476601 with juvenile
idiopathic arthritis is specific to females. *Genes Immun* **16**, 495-498 (2015).
8.   L. M. Gomez *et al.*, PTPN22 C1858T polymorphism in Colombian patients with
autoimmune diseases. *Genes Immun* **6**, 628-631 (2005).
9.   L. B. Ivashkiv, PTPN22 in autoimmunity: different cell and different way. *Immunity* **39**,
91-93 (2013).
10.   H. Wang *et al.*, Identification of PTPN22, ST6GAL1 and JAZF1 as psoriasis risk genes
demonstrates shared pathogenesis between psoriasis and diabetes. *Exp Dermatol* **26**,
1112-1117 (2017).
11.   H. H. Chang *et al.*, PTPN22 modulates macrophage polarization and susceptibility to
dextran sulfate sodium-induced colitis. *J Immunol* **191**, 2134-2143 (2013).
12.   X. Dai *et al.*, A disease-associated PTPN22 variant promotes systemic autoimmunity in
murine models. *J Clin Invest* **123**, 2024-2036 (2013).
13.   Y. Wang *et al.*, The autoimmunity-associated gene PTPN22 potentiates toll-like receptor-
driven, type 1 interferon-dependent immunity. *Immunity* **39**, 111-122 (2013).
14.   Y. Wang *et al.*, PTPN22 Variant R620W Is Associated With Reduced Toll-like Receptor
7-Induced Type I Interferon in Systemic Lupus Erythematosus. *Arthritis Rheumatol* **67**,
2403-2414 (2015).
15.   X. Lin *et al.*, CRISPR-Cas9-Mediated Modification of the NOD Mouse Genome With
Ptpn22R619W Mutation Increases Autoimmune Diabetes. *Diabetes* **65**, 2134-2138
(2016).
16.   A. M. Wallis *et al.*, TRAF3 enhances TCR signaling by regulating the inhibitors Csk and
PTPN22. *Sci Rep* **7**, 2081 (2017).

17. E. Fiorillo *et al.*, Autoimmune-associated PTPN22 R620W variation reduces
phosphorylation of lymphoid phosphatase on an inhibitory tyrosine residue. *J Biol Chem*
285, 26506-26518 (2010).
18. R. C. Orozco, K. Marquardt, K. Mowen, L. A. Sherman, Proautoimmune Allele of
Tyrosine Phosphatase, PTPN22, Enhances Tumor Immunity. *J Immunol* **207**, 1662-1671
(2021).
19. R. C. Orozco *et al.*, Autoimmunity-associated allele of tyrosine phosphatase gene
PTPN22 enhances anti-viral immunity. *PLoS Pathog* **20**, e1012095 (2024).
20. M. R. Spalinger *et al.*, NLRP3 tyrosine phosphorylation is controlled by protein tyrosine
phosphatase PTPN22. *J Clin Invest* **126**, 1783-1800 (2016).
21. M. R. Spalinger *et al.*, PTPN22 regulates NLRP3-mediated IL1B secretion in an
autophagy-dependent manner. *Autophagy* **13**, 1590-1601 (2017).
22. K. Hasegawa *et al.*, PEST domain-enriched tyrosine phosphatase (PEP) regulation of
effector/memory T cells. *Science* **303**, 685-689 (2004).
23. T. Vang *et al.*, LYP inhibits T-cell activation when dissociated from CSK. *Nat Chem Biol*
8, 437-446 (2012).
24. E. L. Hornick, A. M. Wallis, G. A. Bishop, TRAF3 enhances type I interferon receptor
signaling in T cells by modulating the phosphatase PTPN22. *Sci Signal* **15**, eabn5507
(2022).
25. A. Gjorloff-Wingren, M. Saxena, S. Williams, D. Hammi, T. Mustelin, Characterization
of TCR-induced receptor-proximal signaling events negatively regulated by the protein
tyrosine phosphatase PEP. *Eur J Immunol* **29**, 3845-3854 (1999).
26. C. J. Maine *et al.*, PTPN22 alters the development of regulatory T cells in the thymus. *J*
Immunol **188**, 5267-5275 (2012).
27. C. J. Maine, K. Marquardt, J. Cheung, L. A. Sherman, PTPN22 controls the germinal
center by influencing the numbers and activity of T follicular helper cells. *J Immunol*
192, 1415-1424 (2014).
28. H. H. Chang, N. Dwivedi, A. P. Nicholas, I. C. Ho, The W620 Polymorphism in PTPN22
Disrupts Its Interaction With Peptidylarginine Deiminase Type 4 and Enhances
Citrullination and NETosis. *Arthritis Rheumatol* **67**, 2323-2334 (2015).
29. T. Vang *et al.*, Autoimmune-associated lymphoid tyrosine phosphatase is a gain-of-
function variant. *Nat Genet* **37**, 1317-1319 (2005).
30. C. J. Maine, J. R. Teijaro, K. Marquardt, L. A. Sherman, PTPN22 contributes to
exhaustion of T lymphocytes during chronic viral infection. *Proc Natl Acad Sci U S A*
113, E7231-E7239 (2016).
31. T. Jofra *et al.*, Protein tyrosine phosphatase PTPN22 has dual roles in promoting
pathogen versus homeostatic-driven CD8 T-cell responses. *Immunol Cell Biol* **95**, 121-
128 (2017).
32. T. Jofra *et al.*, Extrinsic Protein Tyrosine Phosphatase Non-Receptor 22 Signals
Contribute to CD8 T Cell Exhaustion and Promote Persistence of Chronic Lymphocytic
Choriomeningitis Virus Infection. *Front Immunol* **8**, 811 (2017).
33. X. Zhou, S. Ramachandran, M. Mann, D. L. Popkin, Role of lymphocytic
choriomeningitis virus (LCMV) in understanding viral immunology: past, present and
future. *Viruses* **4**, 2650-2669 (2012).
34. Z. B. Zalinger, R. Elliott, S. R. Weiss, Role of the inflammasome-related cytokines Il-1
and Il-18 during infection with murine coronavirus. *J Neurovirol* **23**, 845-854 (2017).

35.   L. Zhao *et al.*, Antagonism of the interferon-induced OAS-RNase L pathway by murine
695   coronavirus ns2 protein is required for virus replication and liver pathology. *Cell Host*
696   *Microbe* **11**, 607-616 (2012).
697   36.   J. K. Roth-Cross *et al.*, Organ-specific attenuation of murine hepatitis virus strain A59 by
698   replacement of catalytic residues in the putative viral cyclic phosphodiesterase ns2. *J*
699   *Virol* **83**, 3743-3753 (2009).
700   37.   J. Das Sarma *et al.*, Demyelinating and nondemyelinating strains of mouse hepatitis virus
701   differ in their neural cell tropism. *J Virol* **82**, 5519-5526 (2008).
702   38.   K. C. MacNamara, M. M. Chua, P. T. Nelson, H. Shen, S. R. Weiss, Increased epitope-
703   specific CD8+ T cells prevent murine coronavirus spread to the spinal cord and
704   subsequent demyelination. *J Virol* **79**, 3370-3381 (2005).
705   39.   E. Lavi, D. H. Gilden, M. K. Highkin, S. R. Weiss, Persistence of mouse hepatitis virus
706   A59 RNA in a slow virus demyelinating infection in mice as detected by in situ
707   hybridization. *J Virol* **51**, 563-566 (1984).
708   40.   E. Lavi, D. H. Gilden, M. K. Highkin, S. R. Weiss, MHV-A59 pathogenesis in mice. *Adv*
709   *Exp Med Biol* **173**, 237-245 (1984).
710   41.   K. K. Eriksson, L. Cervantes-Barragan, B. Ludewig, V. Thiel, Mouse hepatitis virus liver
711   pathology is dependent on ADP-ribose-1"-phosphatase, a viral function conserved in the
712   alpha-like supergroup. *J Virol* **82**, 12325-12334 (2008).
713   42.   C. M. Kerr *et al.*, PARP12 is required to repress the replication of a Mac1 mutant
714   coronavirus in a cell- and tissue-specific manner. *J Virol* **97**, e0088523 (2023).
715   43.   B. C. Turner, E. M. Hemmila, N. Beauchemin, K. V. Holmes, Receptor-dependent
716   coronavirus infection of dendritic cells. *J Virol* **78**, 5486-5490 (2004).
717   44.   O. L. Wijburg, M. H. Heemskerk, C. J. Boog, N. Van Rooijen, Role of spleen
718   macrophages in innate and acquired immune responses against mouse hepatitis virus
719   strain A59. *Immunology* **92**, 252-258 (1997).
720   45.   H. Zhou, S. Perlman, Preferential infection of mature dendritic cells by the JHM strain of
721   mouse hepatitis virus. *Adv Exp Med Biol* **581**, 411-414 (2006).
722   46.   S. Kim *et al.*, In vivo developmental stages in murine natural killer cell maturation. *Nat*
723   *Immunol* **3**, 523-528 (2002).
724   47.   L. Chiossone *et al.*, Maturation of mouse NK cells is a 4-stage developmental program.
725   *Blood* **113**, 5488-5496 (2009).
726   48.   D. Vremec, J. Pooley, H. Hochrein, L. Wu, K. Shortman, CD4 and CD8 expression by
727   dendritic cell subtypes in mouse thymus and spleen. *J Immunol* **164**, 2978-2986 (2000).
728   49.   H. Wu *et al.*, TIM-4 interference in Kupffer cells against CCL4-induced liver fibrosis by
729   mediating Akt1/Mitophagy signalling pathway. *Cell Prolif* **53**, e12731 (2020).
730   50.   M. Ni *et al.*, T-Cell Immunoglobulin and Mucin Domain-Containing Protein-4 Is Critical
731   for Kupffer Cell Homeostatic Function in the Activation and Resolution of Liver
732   Ischemia Reperfusion Injury. *Hepatology* **74**, 2118-2132 (2021).
733   51.   W. Li, Y. Yang, L. Yang, N. Chang, L. Li, Monocyte-derived Kupffer cells dominate in
734   the Kupffer cell pool during liver injury. *Cell Rep* **42**, 113164 (2023).
735   52.   P. D. Krueger *et al.*, Murine liver-resident group 1 innate lymphoid cells regulate optimal
736   priming of anti-viral CD8+ T cells. *J Leukoc Biol* **101**, 329-338 (2017).
737   53.   D. K. Sojka *et al.*, Tissue-resident natural killer (NK) cells are cell lineages distinct from
738   thymic and conventional splenic NK cells. *Elife* **3**, e01659 (2014).

54. L. Cervantes-Barragan *et al.*, Control of coronavirus infection through plasmacytoid
dendritic-cell-derived type I interferon. *Blood* **109**, 1131-1137 (2007).
55. N. K. Bjorkstrom, B. Strunz, H. G. Ljunggren, Natural killer cells in antiviral immunity.
Nat Rev Immunol **22**, 112-123 (2022).
56. S. N. Waggoner *et al.*, Roles of natural killer cells in antiviral immunity. *Curr Opin Virol*
16, 15-23 (2016).
57. T. P. Salazar-Mather, J. S. Orange, C. A. Biron, Early murine cytomegalovirus (MCMV)
infection induces liver natural killer (NK) cell inflammation and protection through
macrophage inflammatory protein 1alpha (MIP-1alpha)-dependent pathways. *J Exp Med*
187, 1-14 (1998).
58. T. P. Salazar-Mather, T. A. Hamilton, C. A. Biron, A chemokine-to-cytokine-to-
chemokine cascade critical in antiviral defense. *J Clin Invest* **105**, 985-993 (2000).
59. K. A. Daniels *et al.*, Murine cytomegalovirus is regulated by a discrete subset of natural
killer cells reactive with monoclonal antibody to Ly49H. *J Exp Med* **194**, 29-44 (2001).
60. A. O. Dokun *et al.*, Specific and nonspecific NK cell activation during virus infection.
Nat Immunol **2**, 951-956 (2001).
61. J. Loh, D. T. Chu, A. K. O'Guin, W. M. Yokoyama, H. W. t. Virgin, Natural killer cells
utilize both perforin and gamma interferon to regulate murine cytomegalovirus infection
in the spleen and liver. *J Virol* **79**, 661-667 (2005).
62. H. Balachandran *et al.*, NK cells modulate in vivo control of SARS-CoV-2 replication
and suppression of lung damage. *PLoS Pathog* **20**, e1012439 (2024).
63. C. A. Fielding *et al.*, SARS-CoV-2 host-shutoff impacts innate NK cell functions, but
antibody-dependent NK activity is strongly activated through non-spike antibodies. *Elife*
11 (2022).
64. C. van Eeden, L. Khan, M. S. Osman, J. W. Cohen Tervaert, Natural Killer Cell
Dysfunction and Its Role in COVID-19. *Int J Mol Sci* **21** (2020).
65. H. Baazim *et al.*, CD8(+) T cells induce cachexia during chronic viral infection. *Nat*
Immunol **20**, 701-710 (2019).
66. A. B. Begovich *et al.*, A missense single-nucleotide polymorphism in a gene encoding a
protein tyrosine phosphatase (PTPN22) is associated with rheumatoid arthritis. *Am J Hum*
Genet **75**, 330-337 (2004).
67. BioGPS (Ptpn22 (protein tyrosine phosphatase, non-receptor type 22).
68. K. Douroudis, A. Shcherbakova, H. Everaus, A. Aints, PTPN22 gene regulates natural
killer cell proliferation during in vitro expansion. *Tissue Antigens* **76**, 315-318 (2010).
69. S. Pelletier, S. Gingras, D. R. Green, Mouse genome engineering via CRISPR-Cas9 for
study of immune function. *Immunity* **42**, 18-27 (2015).
70. E. Hemmila *et al.*, Ceacam1a-/- mice are completely resistant to infection by murine
coronavirus mouse hepatitis virus A59. *J Virol* **78**, 10156-10165 (2004).
71. H. Zhou, S. Perlman, Mouse hepatitis virus does not induce Beta interferon synthesis and
does not inhibit its induction by double-stranded RNA. *J Virol* **81**, 568-574 (2007).
72. X. Zhang, R. Goncalves, D. M. Mosser, The isolation and characterization of murine
macrophages. *Curr Protoc Immunol* **Chapter 14**, 14 11 11-14 11 14 (2008).
73. A. R. Fehr *et al.*, The nsp3 macrodomain promotes virulence in mice with coronavirus-
induced encephalitis. *J Virol* **89**, 1523-1536 (2015).

74. L. S. Voth *et al.*, Unique Mutations in the Murine Hepatitis Virus Macrodomain
Differentially Attenuate Virus Replication, Indicating Multiple Roles for the
Macrodomain in Coronavirus Replication. *J Virol* **95**, e0076621 (2021).
75. W. Shi *et al.*, Isolation and purification of immune cells from the liver. *Int
Immunopharmacol* **85**, 106632 (2020).

ACKNOWLEDGMENTS:

We thank the KU Flow Cytometry Core, KU Animal Care Unit, and all members of the Orozco
lab for their discussions and assistance. We also thank Dr. Hans Dalton for his edits on this
manuscript.

Funding Sources:

NIH T32 Chemical Biology T32 Training Grant GM13206 (AB)
KU Chemical Biology of Infectious Disease (COBRE) National Institute of General Medical
Sciences (NIGMS) of the National Institutes of Health under award number P20GM113117
(RCO, ARF)
KU Center for Genomics Seed Grant (RCO)
Funds from the University of Kansas (RCO)
NIGMS R35GM138029 (ARF)
NIH T34 GM062232 (KJLHR)

Author contributions:

Conceptualization: AMB, ARF, RCO

Methodology: AMB, KJLHR, TC, CMK, SM, ARF, RCO

Investigation: AMB, KJLHR, NS, CMK, SM, RCO

Visualization: AMB, KJLHR, SM, ARF, RCO

Funding acquisition: AMB, KJLHR, RCO

Project administration: RCO

Supervision: SM, ARF, RCO

Writing – original draft: AMB, SM, ARF, RCO

Writing – review & editing: AMB, KJLHR, NS, TRC, CMK, SM, ARF, RCO

Competing interests:

Authors declare that they have no competing interests.

Data and materials availability:

All data are available in the main text or the supplementary materials. Raw data is available upon
request.

 Main Text Figure Legends:

 Figure 1: PEP-R619W mice have reduced disease during MHV A59 Infection.

 5-week-old C57BL/6J mice with *Ptpn22* wild type (PEP-WT, black circle), *Ptpn22* knockout

 (PEP-null, red square), or the *Ptpn22* autoimmune-associated allele (PEP-R619W, blue triangle),

 were infected with 5e4 PFU (A, C) or 1e5 PFU (B, D) MHV A59 (i.p). Mice are tracked by

 weight (A, B) and survival (C, D). Respective spider plots for individual mouse weights are

 shown below (A, B). To measure viral burden, liver homogenates from infected (5e2 PFU) PEP-

 WT, PEP-null, and PEP-R619W mice were used for plaque assay (E, F). Histopathology of the

 livers of mice at 3 dpi following PBS injection (Mock) or 1e5 PFU MHV A59 infection, with

 representative photomicrographs of liver sections exhibiting areas of necrosis, inflammation, and

 edema/fibrin accumulation (black arrow) and cumulative pathology scores represented as mean +

 SEM. Scale bar = 500 μm (G). Weight loss and survival studies pooled from 3 individual

 experiments for each dose: Group sizes for 5e4 PFU infection (A, C) were PEP-WT n = 28, PEP-

 null n = 14, PEP-R619W n = 17. Group sizes for 1e5 PFU infection (B, D) were PEP-WT n = 31,

 PEP-null n = 27, PEP-R619W n = 31. For weight loss statistical analysis, significant p-values

 from One-way ANOVA with Tukey's Multiple Comparisons at each dpi are represented by #,

 comparing PEP-WT vs. PEP-R619W, * comparing PEP-null vs. PEP-R619W, or ∇ comparing

 PEP-WT vs. PEP-null. Log-rank Mantel-Cox Test (survival), *p<0.05, ***p<0.001. For whole

 liver titer (E, F), each symbol represents an individual mouse. The dashed line indicates the limit

 of detection (L.O.D.). Certain PEP-WT mice are represented in both graphs, as some

 experiments had all three genotypes, whereas others contained only PEP-WT vs PEP-null or

 PEP-R619W. Combined data from 2 independent experiments, multiple unpaired T-Tests, PEP-

 WT vs PEP-R619W, PEP-WT vs PEP-null, at each time point, *p<0.01, **p<0.01. 7 dpi was

analyzed using Fisher's Exact Test, Cleared vs Not Cleared, #p<0.05. For histopathological

analysis (G), mock mice received sterile PBS injection only and are a mixture of all three

genotypes. Each dot represents an individual mouse, with p-values from One-way ANOVA with

Tukey's Multiple Comparisons shown, *p<0.05.

Figure 2: PEP-R619W innate cells provide some protection against disease, but T and B

cells are necessary for survival.

Rag1-/- PEP-WT (black circle), Rag1-/- PEP-null (red square), or Rag1-/- PEP-R619W (blue

triangle), mice were infected with 5e4 PFU (A, C) or 1e5 PFU (B, D) MHV A59 (i.p), and were

tracked by weight (A, B) and survival (C, D). Respective spider plots for individual mouse

weights are shown below (A, B). To measure viral burden, liver homogenates from infected (5e2

PFU) Rag1-/- PEP-WT, Rag1-/- PEP-null, and Rag1-/- PEP-R619W mice were used for plaque

assay (E, F). Weight loss and survival studies pooled from 2 individual experiments for each

dose: Group sizes for 5e4 PFU (A, C) were Rag1-/- PEP-WT n = 16, Rag1-/- PEP-null n = 6,

Rag1-/- PEP-R619W n = 16. Group sizes for 1e5 PFU (B, D) were Rag1-/- PEP-WT n = 17,

Rag1-/- PEP-null n = 13, Rag1-/- PEP-R619W n = 15. For weight loss statistical analysis,

significant p-values from One-way ANOVA with Tukey's Multiple Comparisons at each dpi are

represented by # comparing Rag1-/- PEP-WT vs. Rag1-/- PEP-R619W and * comparing Rag1-/-

PEP-null vs. Rag1-/- PEP-R619W. Log-rank Mantel-Cox Test (survival), *p<0.05, **p<0.01. For

whole liver titers (E, F), each symbol represents an individual mouse. The dashed line indicates

the limit of detection (L.O.D.). Certain Rag1-/- PEP-WT mice are represented in both graphs, as

some experiments had all three genotypes, whereas others contained only Rag1-/- PEP-WT vs

Rag1-/- PEP-null or Rag1-/- PEP-R619W. Combined data from 2 independent experiments,

multiple unpaired T-Tests, Rag1-/- PEP-WT vs Rag1-/- PEP-R619W, Rag1-/- PEP-WT vs

Rag1-/- PEP-null, at each time point. *p<0.01, **p<0.01.

Figure 3: PEP-R619W mice have more innate immune and mature NK cells in the Liver.

Following PBS injection (Mock, white bar, circle) or 1e5 PFU MHV A59 infection (i.p) in PEP-

WT (black bar, circle), PEP-null (red bar, square), and PEP-R619W (blue bar, triangle) mice, at 3

dpi the liver was perfused, harvested, and antibody stained for flow cytometric analysis. The

absolute count of liver cells for all genotypes (A). Frequencies (%) of CD45+ cells of live cells

(B) and of CD3e- CD19- of CD45+ Live (C). Frequency (%) of several innate cells of CD45+

cells is shown (D). Gating strategy for all innate subsets is: Lymphocytes> Single cell x2>

autofluorescent-> Live> CD45+> CD3e- CD19-> Ly6G- Ly6C+/- (non-neutrophils). NK Cells:

innate subsets> NK1.1+. DCs: innate subsets> NK1.1-> CD11c+ PDCA-1-. pDCs: innate

subsets> NK1.1-> PDCA-1+ CD11c+/-. Monocytes: innate subsets> NK1.1-> Ly6C^{hi} CD11b+.

Macrophage: innate subsets> NK1.1-> Ly6C- CD11b+/-> F4/80+ CD11b+. Neutrophils:

Lymphocytes> Single cell x2> autofluorescent-> Live> CD3e- CD19-> Ly6G+ Ly6C+.

Frequency (%, i) of CD45+ and absolute count (#, ii) of conventional NK cells (E). Frequency

(%) mature conventional NK cells (F). Frequency (%, i) of CD45+ and absolute count (#, ii)

mature conventional NK cells (G). Gating strategy for NK subsets: Conventional NK Cells:

innate subsets> NK1.1+> CD49b+ CD49a-. Mature Conventional NK Cells: Conventional NK

Cells> CD11b+. Quantification of flow cytometric data was pooled from 6 independent

experiments. Each symbol represents an individual mouse. Mock mice are a mixture of all 3

genotypes. p-values from One-way ANOVA with Tukey's Multiple Comparisons are shown for

each panel. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Figure 4: PEP-R619W augments mature conventional NK antiviral functions during coronavirus infection.

3 dpi liver M-cNK analysis following 1e5 PFU MHV A59 infection (i.p) in PEP-WT (black bar, circle) and PEP-R619W (blue bar, triangle) mice. Expression (gMFI) of IFNγ (i), T-bet (ii), Granzyme B (iii), Perforin (iv), TNFα (v), and CD107a (vi) in Mature Conventional NK cells, with representative histograms, with fluorescence minus one (FMO) controls included for each marker, below (A). Proportion of IFNγ- (grey), IFNγ+ only (pink), IFNγ+ Perforin+ (teal), and IFNγ+ Perforin+ TNFα+ (purple) of Mature Conventional NK Cells (B). Expression of 13 different cytokines and chemokines were quantified using LEGENDplex. Radar plots showing the 13 different cytokines from the serum (C) and liver (F), with select cytokines, IFNγ (D, G) and IFNα (E, H), quantified to the right. IFNβ concentration in the liver at 3 dpi [ELISA] (I). C57BL/6 mice with wild type (PEP-WT, black circle), IFNγ knockout (IFNγ KO, purple triangle), or IFNγ Receptor knockout (IFNγR KO, orange square) were infected with 5e4 PFU MHV A59 (i.p). Mice are tracked by weight (J) and survival (K). Quantification of flow cytometric data is representative of 3 independent experiments. p-values from Welch's unpaired t test are shown, $*p<0.05$, $**p<0.01$, $***p<0.001$. LEGENDplex assays are from 1 experiment. Each symbol represents an individual mouse. Mock mice received PBS injections only and are a mixture of all 3 genotypes. p-values from One-way ANOVA with Tukey's Multiple Comparisons are shown for each panel. $*p<0.05$, $**p<0.01$, $***p<0.001$. Weight loss and survival studies pooled from 2 individual experiments. Group sizes were PEP-WT n = 19, IFNγ KO n = 4, and IFNγR KO n = 15. For weight loss statistical analysis, significant p-values from One-way ANOVA with Tukey's Multiple Comparisons at each dpi are represented by # comparing PEP-

WT vs. IFNγ KO, or * PEP-WT vs. IFNγR KO. Log-rank Mantel-Cox Test (survival),

****p<0.0001.

Figure 5: PEP-null and PEP-R619W splenic NK cell repertoires have higher proportions of

terminally mature NK cells at 5 weeks of age.

Whole spleens were harvested and stained for flow cytometric analysis from 5-week-old or 8-

week-old naïve mice as indicated. Representative flow plots from 5-week-old mice are shown

from CD3e- CD19- populations, pie charts represent the proportion of the NK1.1+ CD49b+

population that are: CD27- CD11b- (double negative, DN), CD27+ CD11b-, CD27+ CD11b+

(double positive, DP), or CD11b+ CD27- (A). Quantification for frequency (%) of CD27-

CD11b- (double negative, DN) (B), CD27+ CD11b- (C), CD27+ CD11b+ (double positive, DP)

(D), or CD11b+ CD27- (E) NK cells (NK1.1+ CD49b+). Gating strategy for examined NK cells:

Lymphocytes> Single cell x2> autofluorescent-> Live> CD3e- CD19-> NK1.1+ CD49b+>

CD27+/- CD11b+/-. Frequencies of flow cytometric data are combined from 5 independent

experiments. Each symbol represents an individual mouse. p-values from Two-way ANOVA

with Tukey's Multiple Comparisons are shown for each panel. *p<0.05, **p<0.01, ***p<0.001,

****p<0.0001.

Figure 6: PEP-R619W NK Cells control viral burden and are key producers of IFNγ

during MHV A59 infection.

PEP-WT and PEP-R619W mice received a natural killer cell depleting antibody (α-NK1.1,

Clone PK136) or non-depleting isotype control antibody (α-Iso) as indicated prior- and post-

MHV A59 infection (i.p) (A). Following 5e2 PFU MHV A59 infection in α-Iso PEP-WT (black,

black circle), α-Iso PEP-R619W (blue, blue triangle), α-NK1.1 PEP-WT (clear, black circle) and

α-NK1.1 PEP-R619W (clear, blue triangle) mice, the whole liver was harvested at the indicated

dpi, homogenized, and viral titer was quantified via Plaque Assay (B). Mice were injected with

PBS (Mock) or infected with 1e5 PFU MHV A59 (i.p), and at 3 dpi, the whole liver was

harvested, homogenized, and the levels of IFNγ were determined by ELISA (C). Following 1e5

PFU MHV A59 infection, mice were tracked by weight (D) and survival (E). Respective spider

plots for individual mouse weights are shown below (D). Histopathology of the livers of mice at

3 dpi following 1e5 PFU MHV A59 infection, with representative photomicrographs of liver

sections exhibiting areas of necrosis, inflammation, and edema/fibrin accumulation (black arrow)

and cumulative pathology scores represented as mean + SEM. Scale bar = 500 μm (F). Each

symbol represents an individual mouse for whole liver titer (B) and IFNγ ELISA (C). Viral titers

are pooled data from 2 independent experiments at each time point, comparisons are shown from

multiple unpaired T-Tests, α-Iso PEP-WT vs α-NK1.1 PEP-WT , α-Iso PEP-R619W vs α-NK1.1

PEP-R619W, *p<0.01. The dashed line indicates the limit of detection (L.O.D.). IFNγ ELISA is

representative of two independent experiments, p-values from One-way ANOVA with Tukey's

Multiple Comparisons are shown, *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. Mock mice

are a mixture of both genotypes. Weight loss and survival studies were pooled from 2 individual

experiments. Group sizes were α-Iso PEP-WT n = 18, α-Iso PEP-R619W n = 17, α-NK1.1 PEP-

WT n = 19, and α-NK1.1 PEP-R619W n = 17. For weight loss statistical analysis (D), significant

p-values from One-way ANOVA with Tukey's Multiple Comparisons at each dpi are represented

by # comparing α-NK1.1 PEP-WT vs α-NK1.1 PEP-R619W, or * comparing α-Iso PEP-WT vs.

α-Iso PEP-R619W. Log-rank Mantel-Cox Test (survival) (E), ***p<0.001. For histopathological

analysis, each dot represents an individual mouse, with p-values from One-way ANOVA with

Tukey's Multiple Comparisons shown, *p<0.05.