PIKfyve governs endoplasmic reticulum-lysosome contacts to modulate endoplasmic

reticulum dynamics

Nicala Jenkins[1,2], Zainab Adamji[1,2], Nourhan R. Almasri[1], Sharifa Lomiyev[1], and Roberto J.

Botelho[1,2#]

[1]Department of Chemistry and Biology and [2]Molecular Science Graduate Program, Toronto

Metropolitan University, Toronto, Ontario, M5B2K3, Canada

[#]To whom correspondence should be sent: rbotelho@torontomu.ca

Running title: PIKfyve and the endoplasmic reticulum

Keywords: Lipid kinases, organelles, membrane, ER, morphology, membrane contact sites

Abbreviations

Arl8b: ADP-ribosylation factor like 8b; **Atg18**: autophagy-related protein 18; **ClC7:** chlorine

exchange transporter 7; **CLIMP63:** 63 kDa cytoskeleton linking protein; **EB1:** end-binding

protein 1; **ER**: endoplasmic reticulum; **FFAT**: two phenylalanine's (FF) in an acidic tract; **Fig4:**

factor-Induced gene 4; **FYCO1:** FYVE and coiled-coil domain autophagy adaptor 1; **FYVE:** fab1,

YOTB, Vac1 and EEA1 binding domain; **FYVE4A:** fab1, YOTB, Vac1 and EEA1 binding domain

mutant; **GFP:** green fluorescent protein; **GST:** glutathione s-transferase; **GTPase**: guanine

triphosphatase; **KIF5B:** kinesin-1 heavy chain isoform 5B; **LAMP1:** lysosomal associate

membrane protein 1; **LCR:** low complexing region; **mCherry/mCh:** monomeric cherry, red

fluorescent protein; **MCS:** membrane contact site; **PI(3,5)P$_2$**:phosphatidylinositol 3,5-

bisphosphate; **PI(3)P**: phosphatidylinositol 3-phosphate; **PIKfyve**: phosphoinositide FYVE-type

zinc finger containing; **PIP**: phosphoinositide; **PtdIns:** phosphatidylinositol; **PLA:** proximity

ligation assay; **PX Domain:** phox homology domain; **Rab5:** ras-related protein-5; **Rab7:** ras-

related protein-7; **RILP:** rab interacting lysosomal protein; **RILPC33**: rab interacting lysosomal

protein, c terminal 33aa fragment; **Rtn4a:** reticulon 4A; **RUFY3:** RUN and FYVE domain

containing 3; **RUFY4:** RUN and FYVE domain containing 4; **siRNA:** small interfering RNA; **SnxA:**

sorting nexin A; **Snx19:** sorting nexin 19; **TAC:** tip attachment complex; **TRPML1:** transient

receptor potential channel for mucolipin-1; **VAPA:** vesicle associated membrane protein-

associated protein A; **VAPB:** vesicle associated membrane protein-associated protein B; **VPS34:**

vacuolar sorting protein 34; **VPS34 In1:** vacuolar sorting protein 34 inhibitor-1

Summary

Lysosomes are essential to clear unwanted cellular material delivered by constant membrane

fusion. Membrane fission is thus required to balance lysosome size, number, and composition.

PIKfyve is a lipid kinase that converts phosphatidylinositol-3-phosphate [PtdIns(3)P] to

phosphatidylinositol-3,5-bisphosphate [PtdIns(3,5)P_2] and promotes lysosome fission since

lysosomes coalesce into larger, but fewer organelles in PIKfyve defective cells. Here, we reveal a

role for PIKfyve in regulating endoplasmic reticulum (ER) dynamics. We show the ER is less

reticulated and motile in cells acutely inhibited for PIKfyve. Partly, this arises because

lysosomes cluster perinuclearly and are less motile, which appears to arrest ER hitchhiking, a

process in which lysosomes pull and form ER tubules via microtubule-based movement.

Secondly, the ER morphology is distorted because of hyper-tethering of protrudin, an ER

transmembrane protein, to lysosomes via excess PtdIns(3)P and protrudin's FYVE domain. Thus,

our findings reveal that PIKfyve activity balances phosphoinositides at ER-lysosome contact

sites to govern ER properties and has significant implications to our understanding of disease

linked to PIKfyve dysfunction such as neurodegeneration.

Introduction

Lysosomes are acidic and degradative organelles that receive and digest a plethora of molecular

cargo from the endocytic, autophagic and phagocytic pathways – they are thus responsible for

maintaining cellular homeostasis by clearing a wide variety of unwanted, damaged, and

potentially dangerous material (Ballabio and Bonifacino, 2020; Inpanathan and Botelho, 2019).

Lysosomes are highly dynamic because they continuously undergo fusion-fission cycles with

maturing late endosomes, transport vesicles from the *trans*-Golgi network, and between

themselves (Saffi and Botelho, 2019; Luzio et al., 2007). This incessant intermixing is dependent

on bidirectional movement of lysosomes on microtubules driven by kinesin and dynein motors

(Cabukusta and Neefjes, 2018). Perhaps due to this dynamic nature, lysosomes are a

heterogeneous collection of organelles that differ in shape, size, position, and function(Saffi

and Botelho, 2019; Inpanathan and Botelho, 2019). Due to this, and unless otherwise indicated,

we use the term *lysosomes* to encapsulate this full spectrum including late endosomes,

terminal lysosomes, and their hybrids, endolysosomes.

Given this complexity, lysosomes are subject to various regulatory inputs including by

the Rab7 and Arl8b GTPases and the PIKfyve lipid kinase (Ballabio and Bonifacino, 2020; Bissig

et al., 2017; de Lartigue et al., 2009; Cantalupo et al., 2001; Khatter et al., 2015). PIKfyve is a

conserved lipid kinase that localizes to endosomes and lysosomes via its FYVE domain and

phosphatidylinositol-3-phosphate [PtdIns(3)P], which is then converted by PIKfyve into

phosphatidylinositol-3,5-bisphosphate [PtdIns(3,5)P_2] (Ikonomov et al., 2001; Sbrissa et al.,

2002, 2007). While deletion of PIKfyve is embryonic lethal, tissue specific knockouts or

mutations that abate PtdIns(3,5)P_2 is associated with immunity, inflammation, metabolic

defects, neurological conditions and more (Ikonomov et al., 2011; Min et al., 2014; Chow et al.,

2007; Tsuruta et al., 2009; Cheng et al., 2025; Min et al., 2019; Prado et al., 2025). At the

molecular level, there remain few known PtdIns(3,5)P_2-binding proteins, which include Ca^{2+} and

Cl^- channels like TRPML1 and ClC7 (Leray et al., 2022; Dong et al., 2010), the actin regulator

cortactin (Hong et al., 2015), membrane curvature factors like the yeast Atg18 (Gopaldass et al.,

2017), and the *Dictyostelium* SnxA, which is proposed as a reporter for PtdIns(3,5)P_2 (Vines et

al., 2023). Through these and other unknown effectors, PIKfyve and PtdIns(3,5)P_2 collectively

control osmotic regulation, lysosome fusion-fission cycling, the degradative capacity of

lysosomes, and nutrient and stress sensing (Bissig et al., 2017; Choy et al., 2018; Leray et al.,

2022; Dong et al., 2010; Bridges et al., 2012; Sharma et al., 2019). The most glaring defect

caused by PIKfyve loss of function is lysosome enlargement and the concomitant decline in

lysosome number (Bissig et al., 2017; Choy et al., 2018; Sharma et al., 2019). This suggests that

PIKfyve regulates lysosome fusion-fission cycling with the evidence pointing to a specific role in

fission (Bissig et al., 2017; Choy et al., 2018; Sharma et al., 2019). However, the molecular

mechanism impacted by PIKfyve loss of function resulting in defective fission remains unclear.

Interestingly, endosomes and mitochondrial sites of fission are demarcated by endoplasmic

reticulum (ER)-dependent contacts (Friedman et al., 2013; Abrisch et al., 2020; Friedman et al.,

2011), but how this relates to PIKfyve and lysosomes in general is unresolved.

The ER is a reticulated membrane network composed of membrane sheets and tubules

extending throughout the cell (Westrate et al., 2015). The ER is organized into distinct

structural and functional domains, including the rough ER (ribosome-associated) and smooth

ER, that can undergo remodelling in response to stresses such as such as changes to protein

quality control systems involved in ER-associated degradation, protein chaperones and ER-

phagy (Chen et al., 2023; Goyal and Blackstone, 2013). There are several known proteins that

shape the ER. For example, ER sheets are formed by integral membrane proteins such as

CLIMP63, which forms bridge-like structures connecting opposing ER bilayers (Shibata et al.,

2010). In comparison, ER tubules are dynamic, continually forming, branching, and rearranging

through the action of proteins such as Reticulons and Atlastins (Wang et al., 2016; Hu et al.,

2009; Voeltz et al., 2006; Goyal and Blackstone, 2013).

ER remodeling can also occur through motor and cytoskeleton-driven processes. This

includes microtubule tip attachment complex (TAC)-mediated movement and ER-sliding, which

rely on the interaction between the ER membrane and microtubules to facilitate the formation

of new ER tubules (Waterman-Storer and Salmon, 1998; Rodríguez-García et al., 2020; Spits et

al., 2021), and ER hitchhiking. During ER hitchhiking, the ER forms a contact with an endosome

or lysosome, and the latter then moves via a microtubule and motor-dependent process

(Guimaraes et al., 2015; Spits et al., 2021). The attached ER then co-migrates with the motile

endosome/lysosome to grow a new ER tubule (Guimaraes et al., 2015; Spits et al., 2021). For ER

hitchhiking to occur, stable and physical contact sites between the ER and

endosomes/lysosomes are required (Voeltz et al., 2024). Moreover, and unlike ER-sliding and

TAC mediated mechanisms, which relies on direct motor attachment to the ER, hitchhiking uses

endosome/lysosome-motor complexes to displace the attached ER-tubules (Spits et al., 2021;

Raiborg et al., 2015b; a). These complexes are thought to be maintained by tethering proteins,

such as protrudin that bridge the two organelles together (Raiborg et al., 2015b; a; Pedersen et

al., 2020; Spits et al., 2021).

Protrudin, which was originally identified as a protein that promotes neurite formation

(Shirane and Nakayama, 2006), is an ER-transmembrane protein that together with VAPA,

another ER transmembrane protein, and the lysosomal Rab7 GTPase, form an ER-lysosome

tethering complex (Raiborg et al., 2015a; Pedersen et al., 2020; Matsuzaki et al., 2011; Saita et

al., 2009). Protrudin has a FYVE domain and a protein-binding low complexing region (LCR) that

respectively are thought to bind to PtdIns(3)P and GTP-bound Rab7 on endosomes/lysosomes

(Raiborg et al., 2015a; Pedersen et al., 2020). Protrudin then extends its cytoplasmic region to

loop around and interact with VAPA using the FFAT motif. Through this contact, protrudin is

thought to load kinesin-1 onto lysosome-bound FYCO1, a Rab7 effector that promotes

lysosome anterograde movement and binds PtdIns(3)P (Pedersen et al., 2020; Raiborg et al.,

2015a; Jongsma et al., 2024; Matsuzaki et al., 2011; Pankiv et al., 2010). Collectively, protrudin

not only tethers ER-lysosomes together, but also modulates their relative dynamics (Cabukusta

and Neefjes, 2018; Jongsma et al., 2016, 2024).

Overall, there is bidirectional relationship between the ER and organelles of the endo-

lysosomal pathway through contact sites. ER contacts appear to define areas of endosome, and

possibly lysosome fission, while endosomes and lysosomes help reshape ER via hitchhiking.

Given this, we postulated that ER-driven fission may malfunction in PIKfyve inhibition resulting

in lysosome coalescence. Additionally, we hypothesized that lysosome coalescence during

PIKfyve loss may alter ER morphology and dynamics. Here, we provide evidence that PIKfyve

inhibition potently distorts the ER morphology and dynamics by hindering ER hitchhiking and

stabilizing protrudin interaction with lysosomes via excessive PtdIns(3)P levels.

Results

Lysosome morphology is relatively intact in cells manipulated for ER morphology

PIKfyve inhibition enlarges lysosomes via coalescence, ostensibly by perturbing fission (Choy et

al., 2018; Bissig et al., 2017; Sharma et al., 2019). On the other hand, sites of tubular endosome

fission are demarcated by ER tubules that wrap around endosomes (Friedman et al., 2013;

Rowland et al., 2014). Conceivably then, lysosome coalescence and fission defects in PIKfyve-

inhibited cells may be the corollary of perturbed contacts between ER and endo-lysosomal

structures. To test this hypothesis, we over-expressed CLIMP63 or Rtn4A, which respectively

bias ER towards sheets and reticulated tubules (Shibata et al., 2010; Wang et al., 2016; Wu and

Voeltz, 2021; Voeltz et al., 2006), and examined their effect on lysosome morphology in resting

and apilimod-treated cells.

As expected, Cos7A cells overexpressing mCh-CLIMP63 displayed an overwhelmingly

sheet-like ER network with a dramatic decline of ER tubules **(Fig. 1A).** Conversely, cells

expressing eGFP-Rtn4 exhibited an extensive and exaggerated tubular ER network **(Fig 1B).** We

then quantified the average lysosome number and the average volume of individual lysosomes

in cells over-expressing mCherry or GFP to control for transfection, or over-expressing CLIMP63

or Rtn4 after incubation with apilimod to block PIKfyve. Overexpression of these proteins on

their own did not alter lysosome size and total lysosome number **(Fig. 1C-F).** This suggests that

distortion of ER morphology by these approaches does not readily change the fission-fusion

dynamics of dextran-loaded lysosomes. As expected, apilimod-treatment reduced lysosome

number and increased their size in GFP/mCherry-expressing cells **(Fig. 1C-F)**. Relative to these

conditions, the overexpression of CLIMP63 or Rtn4 did not significantly alter the total lysosome

number **(Fig. 1C-F)**, but they respectively enhanced and reduced apilimod-driven changes in the

volume of individual lysosomes **(Fig. 1D and 1F).** These results indicate that manipulating ER

morphology does not readily mimic the effects of PIKfyve on lysosome dynamics, and thus it is

unlikely that PIKfyve-regulation of lysosome fusion-fission cycling depends on ER-lysosome

contacts.

PIKfyve inhibition distorts ER morphology

While ER morphology had relatively minor effects on lysosome size, we next considered the

converse: whether lysosome distortion in PIKfyve-inhibited cells would alter the ER spatial

organization. To test this, we expressed ER-mNeonGreen in Cos7A cells to label the ER and

labelled lysosomes with a fluorescent dextran. Moreover, we opted to use a pharmacological

approach to block PIKfyve for these experiments because this permitted acute disturbance of

lysosomes, while genetic or epigenetic methods like siRNA require at least a day, which might

cause a host of indirect effects.

In vehicle-treated cells, the ER network exhibited a series of sheets and tubules throughout

the cell volume **(Fig 2A)**, some of which were juxtaposed to dextran-labelled lysosomes **(Fig.**

2B). Strikingly, the ER network was visibly distorted in PIKfyve-inhibited cells, exhibiting

numerous "black holes" **(Fig. 2A).** Many of these voids, but not all, were occupied with

Alexa647-dextran **(Fig. 2B).** To determine if some of these voids were also cytosolic in nature or

contoured by other swollen endosomal structures, we expressed cytosolic mCherry or mCherry-

Rab5. We found that none of the apparent ER-decorated voids labelled with cytosolic mCherry

(Sup. Fig 1A), but a subpopulation was lined with mCherry-Rab5 **(Sup. Fig 1B)**. Overall, the ER

"black holes" were occupied with either dextran-labelled lysosomes or mCherry-Rab5

endosomes **(Sup. Fig 1C-D)**, indicating that separate populations of enlarged endosomal and

lysosomal organelles can alter the ER architecture. Similar observations were made in cells

inhibited with YM201636 **(Sup. Fig 2A-B).** To complement our visual inspection and buttress

these assertions, we quantified the ER morphology by extracting total ER skeletal length,

number of branches and junctions per cell, and average branch length. Apilimod and YM201636

both caused a significant decline in ER total skeletal length, loss of ER junctions and branch

numbers, though not for ER branch length **(Fig. 2C-F** and **Sup. Fig 2C-F).** Collectively, these data

respectively indicate a reduction in the total ER network size and loss of the ER

interconnectivity in cells negated for PIKfyve activity.

Loss of PIKfyve activity impairs ER dynamics

We next examined if ER motility was disturbed in PIKfyve-inhibited cells. To assess this, we

employed a previously described method that measures membrane displacement by tracking

pixel intensity changes over time (Spits et al., 2021). To achieve this, we tracked ER-

mNeonGreen pixels over time in control and PIKfyve-hindered cells, binning pixel displacement

into three categories: low (displacement of 0-3 pixels), intermediate (displacement of 4-7

pixels), and rapid ER movement (8 or more pixels). Relative to vehicle-treated cells, we

observed that cells exposed to apilimod or YM201636 displayed a greater number of pixels with

low membrane displacement and a corresponding drop in pixels with rapid membrane

displacement (**Fig. 3**; **Sup. Fig. 2A, 2G-I**; **Sup. Videos 1-3**). These results reveal that the ER is less

motile in cells inhibited for PIKfyve activity further reinforcing the idea that endo-lysosomal

coalescence distorts ER architecture and dynamics.

ER hitchhiking is blunted in PIKfyve-perturbed cells

Conceivably, the distortion of the ER spatio-temporal properties in PIKfyve-deficient cells could

simply arise from physical hindrance of swollen endo-lysosomes compressing the intricate ER

network of tubules and sheets. While physical hindrance of swollen lysosomes may very well

help disfigure the ER network, we noted that there were few swollen lysosomes in the

periphery of PIKfyve-inhibited cells (quantified below), which nevertheless sustained loss of ER

complexity. We thus postulated that a more sophisticated mechanism was at play.

The ER is partly sculpted by forming and extending new ER tubules, which can then fuse

with existing tubules to promote ER interconnectivity (Westrate et al., 2015; Guo et al., 2018;

Shibata et al., 2010; Spits et al., 2021). There are several processes that drive ER shaping that

depend on microtubules: ER sliding, TAC-dependent ER extension, and ER hitchhiking whereby

a moving endo-lysosome drags and grows a nascent ER tubule (Spits et al., 2021; Grigoriev et

al., 2008; Rodríguez-García et al., 2020; Guo et al., 2018; Shibata et al., 2010). We thus

investigated if ER spatial-temporal changes sustained in PIKfyve-inhibited cells resulted because

these processes were perturbed, especially in regions of the ER not physically occupied by

swollen endo-lysosomes.

First, using immunofluorescence against α-tubulin to analyze microtubule morphology,

we did not observe a significant difference in microtubule organization between vehicle and

apilimod-treated cells **(Fig. 4A)** when quantifying parameters such as number of microtubule

branches and junctions per cell, and average microtubule branch length **(Fig. 4B-D).**   This was

also true for the number and kinetics of EB1-GFP comets, which associate with microtubule

plus-ends **(Fig. 4E-H; Sup. Videos 4-6).** We then turned to measuring ER hitchhiking events by

tracking ER and lysosome movement over one-minute movies at 0.3 Hz. Timelapses were then

analysed for lysosome-ER co-migration in control and PIKfyve-inhibited cells. We observed an

average of 4-6 hitchhiking events/min/100 μm^2 area in control cells, which substantially

dropped to 0-1 events/min/100 μm^2 in apilimod exposed cells **(Fig. 4I, J; Sup. Videos 7-8).**

These observations support a model whereby the botched ER morphology and dynamics in

PIKfyve-defective cells are partly underpinned by failed ER hitchhiking.

Lysosome distribution and movement are altered in PIKfyve-inhibited cells

Hitchhiking requires the ER to attach to a lysosome and for the lysosome to move (Spits et al.,

2021; Pedersen et al., 2020; Voeltz et al., 2024). We thus reasoned that abrogation of ER

hitchhiking in PIKfyve-hindered cells may ensue from a defect in attachment and/or lysosome

motility. We first tested lysosome motility and distribution in control and PIKfyve-impaired

cells. Using shell analysis, we reveal that lysosomes collapse into the perinuclear region and

become less frequent in the cell periphery **(Fig. 5A-C).** This was accompanied by a significant

decline in lysosome speed and total distance travelled in apilimod-treated Cos7A cells relative

to vehicle-treated cells (**Fig. 5E-G),** made visible through kymographs that revealed a marked

decrease in the motility of lysosomes upon PIKfyve inhibition (**Fig. 5D)**. Collectively, loss of

PIKfyve activity is associated with hindered lysosome motility and displacement to the

perinuclear region. This implies that in the absence of PIKfyve activity, lysosomes likely sustain a

loss of kinesin and/or hyperactivation of dynein activities. As a corollary, this could impair ER

hitchhiking and disrupt the ER network.

PIKfyve inhibition alters ER-lysosome contact sites

There are likely several types of ER-lysosome contact sites (Rudnik et al., 2024; Raiborg et al.,

2015a; Allison et al., 2017; Du et al., 2023; Bonet-Ponce and Cookson, 2022; Cai et al., 2022;

Gao et al., 2022; Saric et al., 2021). However, contact sites formed by the ER transmembrane

proteins, protrudin and VapA/B, and the lysosomal Rab7 GTPase, are linked to lysosome and ER

position and motility via kinesin (Pedersen et al., 2020; Raiborg et al., 2015a; Matsuzaki et al.,

2011). Therefore, we tested if PIKfyve blockade altered the protrudin-VapA-Rab7 contact site to

understand our earlier observations.

We found that protrudin and VapA appeared proximal in equal measures between

PIKfyve-enabled and PIKfyve-blocked cells (**Fig. 6A, E**), which is perhaps not surprising since

both are ER transmembrane proteins. However, we observed an inverse relationship between

Rab7 and these proteins: there were fewer proximal events between VapA and Rab7 (**Fig. 6B,**

F), but higher between protrudin and Rab7 (**Fig. 6C, G).** This did not readily translate to altered

Rab7 and KIF5B though (**FIG. 6D, H).** To explore this further, we tested the status of Rab7

GTPase. We did not observe a change in total Rab7 associated with LAMP1 by

immunofluorescence (**Sup Fig. 3A-C**) in these cells and experimental conditions. Similarly, we

did not observe a change in the cellular levels of GTP-loaded Rab7 as reported by affinity

precipitation of GTP-bound Rab7 using GST-RILP-C33 in HeLa cells **(Sup Fig. 3D, E)**, though it is

possible that specifics pools of GTP-Rab7 may be altered in these conditions.

These data suggest that the Rab7-VapA-protrudin ER-lysosome contact site is

undergoing a reorganization rather than a complete loss, whereby protrudin-Rab7 interaction is

promoted at the expense of Rab7-VapA proximity. Since PIKfyve inhibition concomitantly

increases PtdIns(3)P while reducing PtdIns(3,5)P$_2$ levels (Choy et al., 2018; Zolov et al., 2012)

and protrudin has a FYVE domain, we postulated that this causes the hyper-stabilization of

protrudin-Rab7 interactions potentially locking ER-lysosome contact sites in a given

conformation.

Lipid tethering mechanisms are altered upon PIKfyve inhibition

To test the above hypothesis, we expressed GFP-chimeras of wild-type (protrudin^{wt}) and a

FYVE4A-mutant protrudin (protrudin^{FYVE4A}; Raiborg et al., 2015a) in cells expressing and/or

silenced for endogenous protrudin (**Fig. 7A**). Protrudin^{wt}-GFP and protrudin^{FYVE4A}-GFP both

exhibited ER-like distribution in the presence of (non-targeting) or the absence of endogenous

protrudin (siRNA; **Fig. 7A**). Remarkably, cells expressing protrudin^{wt}-GFP but inhibited for

PIKfyve displayed intense protrudin^{wt}-GFP ring structures that decorated both dextran-labelled

and dextran-negative vacuoles (**Fig. 7A**). Since protrudin is an ER-transmembrane protein, this

means that the ER membrane or regions of the ER membrane wrap around vacuoles induced in

PIKfyve-disabled cells. This concept is supported by 3D reconstruction of protrudin[wt]-GFP which

extends and at least partially encloses dextran-loaded vacuoles (**Sup Fig. 4A**). Importantly, the

protrudin[FYVE4A]-GFP mutant failed to form these rings around dextran-loaded lysosomes in

PIKfyve-inhibited cells (**Fig. 7A**), indicating that the protrudin-labelled ER membranes tightly

encircle the enlarged lysosomes likely in a PtdIns(3)P-dependent manner. Qualitative

observations suggests that overexpression of protrudin[wt]-GFP, but not of the protrudin[FYVE4A]-

GFP protein reduces the size of PIKfyve-induced vacuoles, reminiscent of observations when

cells overexpressed Rtn4a (**Fig. 1F**).

We then determined if excess protrudin wrapping around vacuoles was at least partially

responsible for the altered ER morphometrics in PIKfyve-arrested cells by skeletonizing the

protrudin-GFP fluorescence. First, protrudin-GFP-based ER morphometrics was

indistinguishable when cells were mock-silenced or silenced for endogenous protrudin (**Fig. 7B-**

D, first two groups). Second, apilimod elicited dramatic changes to ER morphometrics in cells

expressing wild-type protrudin-GFP whether mock or silenced for endogenous protrudin (**Fig.**

7B-D, compare groups 1 and 3, and groups 2 and 4). We then compared the ER morphometrics

by expressing protrudin[FYVE4A]-GFP in cells silenced for endogenous protrudin. In vehicle-treated

cells, the ER morphology was similar to cells expressing protrudin[wt]-GFP (**Fig. 7B-D**, compare

groups 2 and 5). Importantly, the ER changes driven by apilimod were annulled if cells

expressed protrudin[FYVE4A]-GFP relative to protrudin[wt]-GFP (**Fig. 7B-D**, compare groups 4 and 6).

Thus, a functional FYVE domain on protrudin was necessary for the apilimod-induced changes

to ER morphometrics.

We then explored if the absence of PtdIns(3)P or expression of protrudin[FYVE4A]-GFP

rescued the relative proximity of protrudin-Rab7 and Rab7-VapA in PIKfyve-defective cells. First,

as a control, we note that protrudin silencing eliminated the number of proximity events

between Rab7-protrudin PLA (**Fig. 7E**) and Rab7-VapA PLA **(Fig.7F),** suggesting protrudin's role

as a contact site tether is important. Second, VPS34-IN1 which inhibits PtdIns(3)P production by

Vps34, reversed the excess Rab7-protrudin interactions (**Fig. 7G)** and recovered Rab7-VapA

interactions elicited by PIKfyve suppression (**Fig. 7H)**. This intimates that excess PtdIns(3)P in

PIKfyve-arrested cells reorganizes the Rab7-VapA-protrudin interactions. Third, we compared

the proximity of these proteins in cells expressing protrudin[wt]-GFP or protrudin[FYVE4A]-GFP upon

endogenous protrudin silencing **(Fig. 7I)**. As before, apilimod promoted Rab7-protrudin

interactions in cells expressing protrudin[wt]-GFP (**Fig. 7I)**. Importantly, there was no difference

between vehicle and apilimod treated cells expressing protrudin[FYVE4A]-GFP (**Fig. 7I)**. Even

expressing protrudin[FYVE4A]-GFP alone reduced Rab7-protrudin interactions indicating that

protrudin-lipid interaction is important for the organization of this contact site **(Fig. 7I)**.

Overall, we propose that PIKfyve inhibition leads to excess PtdIns(3)P levels, which

inappropriately recruits and locks protrudin to lysosomes via its FYVE domain. This hyper-

recruitment distorts the balance of ER-lysosome contact sites by shifting interactions toward

Rab7-protrudin tethering and away from a conformation wherein Rab7-VapA are more

proximal. The disproportionate lysosomal localization of protrudin also disrupts ER morphology.

Discussion

PIKfyve is a well-established regulator of the endolysosomal pathway, governing the membrane

fusion-fission cycle of endosomes and lysosomes (Choy et al., 2018; Bissig et al., 2017; Sbrissa

et al., 2002; Ikonomov et al., 2001; Sharma et al., 2019). Here, we extend PIKfyve's sphere of

influence by revealing that PIKfyve modulates ER morphology and dynamics – cells acutely

inhibited for PIKfyve present a less reticulate and dynamic ER. Ostensibly, these changes to the

ER are driven partly by two mechanisms. First, the enlarged lysosomes become less motile,

collapsing to the cell centre – as a corollary, we observed fewer ER hitchhiking events, which

promotes ER remodelling and motility (Spits et al., 2021; Guimaraes et al., 2015). Second, the

ER membrane appears "glued" to enlarged lysosomes by extensive protrudin-dependent

binding, likely due to a surfeit of PtdIns(3)P (Choy et al., 2018; Zolov et al., 2012). Thus, we

identified an important role for PIKfyve in modulating ER properties beyond those known for

the endolysosomal system.

PIKfyve modulation of lysosome and ER architecture

Inhibition of PIKfyve and PtdIns(3,5)P_2 hinders lysosome fission eliciting their coalescence (Choy

et al., 2018; Bissig et al., 2017; Gopaldass et al., 2017; Sharma et al., 2019). Currently, we do not

know the fission factors controlled by PIKfyve and PtdIns(3,5)P_2 in mammalian cells, though

Atg18 may be such a factor in yeast (Gopaldass et al., 2017; Dove et al., 2004). On the other

hand, ER tubules contact and demarcate sites of mitochondrion and endosome fission

(Rowland et al., 2014; Friedman et al., 2013, 2011). To test if PIKfyve and ER-dependent fission

were linked, we overexpressed CLIMP63 and Rtn4a to respectively abate and promote ER

tubule formation. We reasoned that CLIMP63 overexpression would lead to enlarged

lysosomes, mimicking PIKfyve arrest, while excess Rtn4a might rescue lysosome coalescence in

PIKfyve-hindered cells by promoting ER tubules; however, we did not discern these predicted

behaviors. While we observed effects on individual lysosome volume in CLIMP63/Rnt4a-

modified cells, this was not accompanied by altered lysosome number, which intimates the

effects on lysosome size by CLIMP63 and Rtn4a overexpression were not because of altered

fusion-fission cycling, as is the case for PIKfyve (Choy et al., 2018; Sharma et al., 2019). Instead,

we anticipate that processes such as modified biogenesis and/or lipid transfer may drive these

changes to lysosome size in this context. It remains possible that CLIMP63 and Rtn4a

overexpression is not ideal approach to test the relationship between PIKfyve-driven processes

and ER.

Conversely, disturbing PIKfyve greatly perturbed ER morphometrics, which became less

reticulated and more sheet-like, decorated with various spherical voids filled with enlarged

lysosomes or endosomes. While some of these ER morphometric changes reflect steric

hindrance by the enlarged endosomes and lysosomes, our data point to a direct role for PIKfyve

in modulating ER properties by calibrating protrudin-dependent ER-lysosome contact sites by

balancing the types and levels of phosphoinositides. Supporting this model, ER morphometrics

were rescued in PIKfyve-arrested cells expressing the FYVE4A-protrudin mutant, but not when

expressing wild-type protrudin, in cells silenced for endogenous protrudin – we discuss the

nature of this contact further below. While physical "glueing" of the ER membrane to

lysosomes likely helps transform the ER morphology in PIKfyve-defective cells, we anticipate

that this may also directly affect ER morphogenic factors like CLIMP63, Atlastin, and reticulons

since protrudin interacts with these factors (Chang et al., 2013; Hashimoto et al., 2014).

PIKfyve modulates lysosome and ER dynamics

A striking consequence of PIKfyve inhibition is the concurrent loss of lysosome and ER motility, which we propose are related processes. While microtubule morphometrics and dynamics appeared normal in PIKfyve-inhibited cells, the enlarged lysosomes were clearly perinuclear and less motile in our experimental models. While we do not precisely know how PIKfyve modulates lysosome dynamics, an attractive mechanism is that PIKfyve tightly balances PtdIns(3)P and PtdIns(3,5)P_2 levels in space and time to govern various PtdIns(3)P-interacting proteins such as FYCO1, RUFY3, RUFY4, protrudin, and SNX19, most of which modulate motor function (Pankiv et al., 2010; Saric et al., 2021; Raiborg et al., 2015a; Jongsma et al., 2024; Kumar et al., 2022; Keren-Kaplan et al., 2022; Jongsma et al., 2016, 2024). Remarkably, our observations on protrudin behavior align well with previous work by Saric *et al. on* SNX19, another ER transmembrane protein that carries a PX domain that binds to PtdIns(3)P (Mas et al., 2014; Saric et al., 2021). These authors observed that SNX19 forms an ER-lysosome contact site that restricts endo-lysosome motility and that enlarged lysosomes in PIKfyve inhibited cells were extensively decorated with SNX19 and were less motile (Saric et al., 2021). Collectively, we propose that PIKfyve activity tightly modulates the relative levels of PtdIns(3)P and PtdIns(3,5)P_2 to coordinate factors like SNX19, protrudin, FYCO1, and RUFY3/4 at ER-lysosome contacts with consequences on lysosome motility. Interestingly, PIKfyve forms a conserved protein complex with its antagonistic phosphatase Fig4 that back-converts PtdIns(3,5)P_2 to PtdIns(3)P; it is tempting to suggest that such an antagonistic complex evolved to enable exquisite control of PtdIns(3)P and PtdIns(3,5)P_2 at spatially-restricted membrane contact sites

to switch motor activity during bidirectional movement (Botelho et al., 2008; Jin et al., 2008;

Sbrissa et al., 2007; Lees et al., 2020; Botelho, 2009). This phosphoinositide-based switch may

interface with the previously observed role of repeated ER-endosome contacts and

microtubule-associated proteins in switching motor activity on endosomes (Raiborg et al.,

2015a; Jongsma et al., 2024)

As for loss of ER dynamism, we propose that this is partly due to declined lysosome

motility, and consequently, reduced ER hitchhiking, which is a major driver of ER remodeling

(Spits et al., 2021). However, ER hitchhiking also depends on the formation of lysosome-ER

contact sites. Based on our observations, ER-lysosome contact sites not only form, but may in

fact undergo hyper-tethering during PIKfyve arrest as visualized for protrudin by us and for

SNX19 by *Saric et al*, both of which indicate that ER membrane subdomains envelope enlarged

lysosomes and related organelles. This implies that bidirectional transport of lysosomes and ER

may depend on regulated conformational or compositional changes to these contacts which

consequently alter kinesin and/or dynein-mediated activity. Conceivably, PIKfyve inhibition

traps specific contact sites labeled with SNX19 or protrudin in a specific state arresting the

putative cycling necessary to elicit ER (and lysosome) dynamics.

PIKfyve and regulation of ER-lysosome contact sites

Protrudin is thought to load kinesin-1 onto FYCO1 on lysosomes (Raiborg et al., 2015a). To do

this, protrudin binds to PtdIns(3)P and Rab7 on lysosomes and VapA on the ER membrane

(Raiborg et al., 2015a). Nonetheless, there is little known about how the structural organization

of these proteins at these contact sites – whether all interactions are concurrent or if they are

subject to a regulated conformational change that enables ER and/or lysosome dynamics. Since

we observed an increase in protrudin-Rab7 proximity and a converse decrease in VapA-Rab7

proximity in PIKfyve arrested cells, this suggests that PIKfyve aids in such a

conformational/compositional change at this ER-lysosome contact site. Since VPS34-IN1 can

reverse these effects, we posit that the surplus PtdIns(3)P caused by PIKfyve inhibition locks

protrudin in a specific conformation that favours Rab7 and sacrifices VapA interaction. This in

turn may occlude Rab7 from interacting with other factors. Overall, our preferred model is that

PIKfyve converts PtdIns(3)P to PtdIns(3,5)P_2 to transiently promote protrudin dissociation from

lysosomal membrane to elicit a conformational change that promotes ER and lysosome

dynamics. Of course, PtdIns(3,5)P_2 itself may have a regulatory role in this process, but we do

not know the targets of this lipid in this context, if any. Together with effects on SNX19-

mediated (Saric et al., 2021) and protrudin-mediated contact sites, our results suggest that

PIKfyve coordinates the molecular machinery at the ER-lysosome interface, influencing both

organelle tethering and downstream functional outcomes such as remodeling, fission, and

motility.

Conclusion

Collectively, our results suggest that PIKfyve supports ER network integrity by governing

lysosome dynamics, distribution, and ER-lysosome contact sites. In its absence, swollen and

immobile lysosomes accumulate that are hyper-tethered to ER membranes in a specific

conformation; consequently, ER hitchhiking fails, thereby contributing to the compromised ER

dynamics. These results illuminate a previously underappreciated aspect of PIKfyve function

and provide mechanistic insight into how lysosomal dysfunction can drive architectural and

dynamic changes of neighboring organelles. Moreover, the PIKfyve pathway is linked to

neurological diseases and is a potential therapeutic target against certain cancers and viral

infections (Chow et al., 2007; Cai et al., 2013; Babu et al., 2024; Cheng et al., 2025; Bao et al.,

2023; Kang et al., 2020). The physiological link between the PIKfyve pathway and these diseases

is generally thought to occur through changes to the endolysosomal pathway. However, given

our findings, mutations in the PIKfyve pathway that lead to disease may reflect altered ER

function as well, particularly in highly secretory or polarized cells where proper ER architecture

and dynamics are critical.

Methods and Materials

Cell culture

COS-7 and HeLa cell strains were cultured in Dulbecco's Modified Eagle Medium (DMEM;

Wisent, St. Bruno, QC) supplemented with 10% heat-inactivated fetal bovine serum (FBS;

Wisent) and 1% penicillin-streptomycin (10,000 U/mL) (P/S; Gibco, Burlington, ON). All cells

were authenticated, checked for contamination and cultured at 37°C with 5% CO_2.

Plasmids, siRNA gene silencing, and transfection

Plasmids were obtained from Addgene as follows: ER-mNeonGreen (#137804), mCherry-

CLIMP63 (#136293), pHAGE2 GFP-Rtn4a (#86684), pcDNA3.1-mCherry (#128744), pcDNA3-

eGFP (#13031), human EB1 in pEGFP N1 to express EB1-GFP (#39299). Plasmids encoding

protrudin[WT]-GFP, protrudin[FYVE4A]-GFP were a kind gift from Dr. Camilla Raiborg (Oslo University

Hospital) and was previously described (Raiborg et al., 2015a). Transient transfections were

performed with FuGENE HD (Promega, Madison, WI) following the manufacturer's instructions,

using a 3:1 DNA to transfection reagent ratio. The transfection mixture was replaced with fresh

medium after 4-6 h and cells were used 24 h post-transfection.

Silencing RNA oligonucleotides were transfected into HeLa cells with Lipofectamine

RNAiMAX Transfection Reagent (Invitrogen, Carlsbad, CA) following the manufacturer's

instructions. One round of knockdown was performed using 25 nM ON-TARGETplus ZFYVE27

(the standard gene name for protrudin) individual siRNA (J-016349-12-0010) or ON-TARGETplus

non-targeting control (Sigma-Aldrich) over 24 h.

Organelle labelling

Lysosomes in COS-7A were labelled with 320 µg/mL Alexa-Fluor 488-, 546- or 647-conjugated

dextran (ThermoFisher), while lysosomes in HeLa cells were labelled with 150 µg/mL by

incubating cells in cell-specific complete medium for 1 h at 37°C with 5% CO_2, followed by

washing in phosphate-buffered saline (PBS) and replenishing with fresh complete medium for 1

h before live-cell imaging. Since this method likely labels a spectrum of late endosomes,

lysosomes and endolysosomes, the term *lysosomes* reflects this range of organelles in this

work. ER-NeonGreen expression was used to label bulk ER.

Pharmacological treatments

PIKfyve inhibition was accomplished by treating cells with 80-240 nM apilimod (Selleck

Chemicals LLC, Houston, TX) or 1 μM YM-201636 (Cayman Chemical Company, Ann Arbor, MI)

for 1-2 h – specific concentrations and times are indicated in each figure legend. Vps34 Class III

PI 3-kinase was arrested by treating cells with 1 μM VPS34-IN1 (Chemical Company, Ann Arbor,

MI) for 30 min.

Immunofluorescence

For immunofluorescence, following treatments, cells were fixed with 4% paraformaldehyde

(PFA; v/v) for 15 min. After washing with PBS, residual PFA was quenched with 100 mM glycine

for 10 min, then cells were blocked and permeabilized with 0.1% (v/v) Triton X-100 for 5 min.

Cells were then subjected to mouse monoclonal (DM1A) antibody against ▯-tubulin (1:100;

#T9026; Sigma-Aldrich), rabbit monoclonal (D9F52) antibody against Rab7 (1:100; #9367),

mouse monoclonal (E9O7E) antibody against Rab7 (1:100; #95746), mouse monoclonal (D4O1S)

antibody against LAMP1 (1:100; #15665) and rabbit monoclonal (D2D11) antibody against

LAMP1 (1:200; #9091; all from Cell Signaling Technology). Dylight-conjugated secondary donkey

antibodies against rabbit and mouse (1:1000; Novus Biologicals Canada) were used.

Proximity ligation assays

PLA was performed with Duolink In Situ Red Starter Kit Mouse/Rabbit (Sigma Aldrich,

DUO92101) according to the manufacturer's instructions. Briefly, HeLa cells were seeded, fixed

with 4% PFA for 15 min, quenched with 100 mM glycine for 10 min and permeabilized with

0.1% Triton X-100 for 5 min – all at RT respectively. Cells were co-incubated with one mouse

and one rabbit antibody from the following; rabbit monoclonal (D9F52) antibody against Rab7

(1:100; #9367; Cell Signaling Technology), mouse monoclonal (E9O7E) antibody against Rab7

(1:100; #95746; Cell Signaling Technology), mouse monoclonal (4C12) antibody against VAPA

(1:100; #293278; Santa Cruz Biotechnology), rabbit polyclonal antibody against ZFYVE27 (1:100;

12680-1-AP; Proteintech), rabbit monoclonal (E3H4J) antibody against KIF5B (1:250; #47610;

Cell Signaling Technology), followed by incubation with PLA probes. After probe incubation,

cells were washed and incubated with ligation mixtures to promote DNA ligation. Next,

amplification mixtures were added to cells for Rolling Circle Amplification (RCA) and

hybridization of fluorescently labelled oligonucleotides to occur. A PLA mounting media

containing DAPI was added to washed cells when mounted. PLA signals from cells were

detected using confocal microscopy.

Fluorescence microscopy

Confocal images were obtained using a Quorum Diskovery spinning disc confocal system

(Quorum Technologies, Inc., Guelph, ON) and a Quorum ReSCAN confocal system (Quorum

Technologies, In., Guelph, ON). The Diskovery microscope consisted of an inverted fluorescence

microscope (DMi8; Leica) equipped with an Andor Zyla 4.2 Megapixel sCMOS camera (Oxford

Instruments, Belfast, UK), a 63x oil immersion objective (1.4 NA), and standard excitation and

emission filter sets and lasers were used for all fluorophores. The microscope system was

controlled by the MetaMorph acquisition software (Molecular Devices, LLC, San Jose, CA, USA).

The ReSCAN microscope consisted of an inverted fluorescence microscope (DMi8 Automated;

Leica) equipped with a ORCA-Flash V3 digital sCMOS camera (Hamamatsu, Shizuoka Prefecture,

Japan) , a 63x oil immersion objective (0.6 to 1.4 NA) , and standard excitation and emission

filter sets and lasers were used for all fluorophores. The microscope system was controlled by

the Volocity acquisition software (Quorum Technologies, In., Guelph, ON). For all live-cell

imaging, cells were maintained in DMEM free of phenol red in a Chamlide microscope-mounted

chamber at 37°C and 5% CO_2. For z-stack imaging, 20-50 images were acquired with an

interplanar distance of 0.2 to 0.6 µm, where origin was set to the centre of the cell. For

timelapse obtained for ER and lysosome dynamics and EB1-GFP dynamics, frames were

acquired every 0.25 s for 1 min. For timelapse obtained for ER-hitchhiking, frames were

acquired every 0.3 s for 1 min.

ER and lysosome morphology analysis

Image analysis was conducted using Volocity and Fiji software. ER morphology analysis was

conducted on Volocity using an ER protocol designed to provide skeletal length measurements

through the thinning (or skeletonizing) of the ER until only a single-voxel-wide-line remains. The

protocol then counts the number of voxels along the path and provides information such as

skeletal length, curvature, branching and fragmentation. Further ER morphology analysis, along

with protrudin-GFP morphology analysis was completed using a semi-automated junction

analysis plug-in for Fiji provided by Spits *et al (2021)* from the Leiden University Medical Centre

in the Netherlands. This plug-in was based on the analyze skeleton feature on Fiji where an ER

mask is initially created and the skeleton representation is analyzed for the number of junctions

and branches (Spits et al., 2021). Lysosome morphology analysis was done on Volocity using a

"Measure Objects" protocol designed to provide voxel/pixel count, volume and area, and total

number measurements.

ER and lysosome dynamics analysis

ER dynamics were analyzed using a membrane displacement analysis macro for Fiji also

provided by *Spits et al (2021)* that analyzed mobile ER fractions based on #pixels/frame ranging

from 0-9. This method is largely based on optical flow analysis which compares corresponding

areas in two frames and generates a map that depicts the frame-to-frame movement of specific

image areas. The extent of movement is reflected by the change in pixel color intensity

between each frame (Spits et al., 2021). Seven separate 60-pixel x 60-pixel regions of interest

around the ER for one single cell per field of view were selected in a clockwise manner for

seven cells and the mobile ER fractions were determined. We then binned movement into

three categories of change (0-3 #pixels/frame), (4-7 #pixels/frame) and (8-9 #pixels/frame) to

define low-, intermediate-, and rapid-movement.

Lysosome dynamics were analyzed using Single Particle Tracking analysis for Fiji that

analyzed lysosome dynamics upon PIKfyve inhibition using time-lapse images acquired every

0.25 s for 1 min. Default settings were kept with the LoG detector (Laplacian of Gaussian). In

vehicle treated cells, the diameter of the lysosome was set to 1.5 μm with a quality threshold of

500. A uniform color was selected for color spots. The simple LAP Tracker was selected with the

linking max distance set to 7 μm, the gap closing distance set to 5 μm, and the gap closing max

frame set to 2 μm. A track index was generated, and the following parameters were extracted:

total distance travelled, mean speed, and overall track displacement. In PIKfyve inhibited cells,

the same process was followed with the exception that the diameter of the lysosome was set to

2 μm with a quality threshold of 250; and both the linking max distance and gap closing max

distance were set to 5 μm. Lysosome motility was also shown using a "Reslice" protocol in Fiji

to create Kymographs between conditions, where a segmented line was drawn along the x-axis

of the cell randomly, with a line width of 2 pixels; generating a space-time projection displaying

spatial position (x-axis) versus time (y-axis).

The frequency of ER-hitchhiking events upon PIKfyve inhibition was analyzed using

Time-lapse analysis acquired every 0.25 s for 1 min. Cells post-treatment were analyzed to

determine the number of ER-hitchhiking events for multiple, randomized (100 μm^2) regions of

interest (ROI). To quantify the number of events, Fiji software was utilized to select ROIs based

on available ER periphery with co-migrating lysosomes. A positive hitchhiking event was

identified if the lysosome localized with the tip of a growing ER tubule, initiated movement, co-

travelled with the ER tubule, and "dropped it off" at another independent ER tubule.

Hitchhiking events were tracked and identified qualitatively by eye.

Lysosome distribution analysis

To determine the change in lysosome distribution, Shell analysis on Fiji software was utilized.

Using the Freehand Selection Tool on Fiji, the cell was outlined to create a "total cell" ROI. This

was followed by using the Enlarge Tool on Fiji to create a shell by shrinking the "total cell" ROI

by 30-50 pixels to create a second layer of shell. This process was repeated to create a third and

fourth layer of shell. The raw integrated density was calculated for each shell independently to

determine the sum of the values of pixel intensities in the respective shell. The raw integrated

density of the inner most shell as a fraction of the "total cell" ROI was calculated and compared

to the outer most shell as a fraction of the "total cell" ROI.

3D Image Reconstruction

Three-dimensional (3D) image reconstruction was performed using Imaris Software (V9.5,

Bitplane, Oxford Instruments). Raw z-stack images (plane thickness: 0.2 μm) were acquired

using a ReSCAN Confocal Microscope (previously explained) and imported into Imaris for

analysis. Prior to reconstruction, image stacks were subjected to background subtraction

preprocessing, with a rolling ball background subtraction of 50 pixels applied across all images

and conditions using Fiji software. Within Imaris, 3D volumes were generated using the Surpass

Mode, and surfaces were rendered using the Surface or Volume modules, with thresholds set

manually based on signal intensity. Reconstruction parameters, including smoothing and

thresholding method were consistently applied across all samples to ensure comparability. For

visualization, coloring was applied to distinguished channels representing lysosomes (magenta)

and protrudin (gray).

Recombinant protein expression and purification

GST-fused RILP-C33 (truncated mutant) and GST-fused empty vectors (ampicillin resistant) were

expressed in BL21 competent cells, grown at 37 ºC and shaking at 200 rpm over-night . Cells

were cultured until the OD_{600} reached 0.6. After treatment with 1 μM isopropyl-ß-D-

thiogalactoside (IPTG) for 4 h at 30 ºC and shaking at 200 rpm, the induced recombinant GST-

RILP-C33 and GST proteins were centrifuged at 4 ºC for 30 minutes at 5000 xg, with the

supernatant discarded and pellet resuspended in 1X bacterial lysis buffer (50 mM Tris, 150 mM

NaCl, 10 mM β-mercaptoethanol, pH 7.5 with protease inhibitor cocktail (Sigma-Aldrich). The

bacterial lysates were then homogenized using a EmulsiFlex-C3 pressure homogenizer

(AVESTIN, Canada). Once homogenized, the recombinant GST-RILP-C33 and GST- lysates were

centrifuged at 4 ºC for 30 minutes at 5000 xg, pelleted, and the supernatant was incubated with

50% glutathione-Sepharose 4B bead resuspension made following manufacturer's instructions

(Millipore Sigma, G317-0756-01) at a dilution of 1:50 (beads to bacterial lysate supernatant,

respectively) for 2 h at 4 ºC with gentle shaking. The samples were then centrifuged at 4 ºC for

5 minutes at 5000 xg, pelleted, supernatant discarded, and pellet resuspended in 1X bacterial

lysis buffer. The samples were analyzed by western blotting to confirm protein expression and

purification.

Affinity precipitation of GTP-bound Rab7

HeLa cells were pre-treated with either vehicle control (DMSO), apilimod, or YM-201636 and

lysed in 700 µL modified- 1X RIPA Buffer (25 mM Tris HCl, 150 mM NaCl, 1% TritonX-100, 1% Na

deoxychlorate, pH 7.4 with protease inhibitor cocktail (Sigma-Aldrich)). Cells were then

incubated on ice for 30 minutes, collected into clean microcentrifuge tubes and centrifuged at

4°C for 5 minutes at 3,000 xg. Prior to centrifugation, 10% of each lysate sample was collected

reserved as an input control. These input samples were also centrifuged at 4°C for 5 minutes at

3,000 xg and subsequently mixed with a 1:1 ratio of 2X Laemmli Sample Buffer (0.5 M Tris, pH

6.8, glycerol and 10% sodium dodecyl sulfate (SDS), 1:10 β-mercaptoethanol). Recombinant

RILP-C33 and GST bound to glutathione-Sepharose 4B beads previously explained were mixed

with HeLa cell lysates at a dilution of 1:14 (beads to cell lysate, respectively). These mixtures

were incubated for 2 h at 4 ºC with gentle shaking. The samples were then centrifuged at 4°C

for 5 minutes at 500 xg, with remaining supernatant discarded. The sample pellets were eluted

using modified-1X RIPA Buffer (1:2 ratio of sample to buffer, respectively). The supernatant

from each wash was moved to a clean microcentrifuge tube and a 1:1 ratio of 2X Laemmli

Sample was added to the samples, boiled at 96°C for 10 minutes and resolved via SDS-PAGE.

SDS-PAGE and Western blotting

Following treatment, whole cell lysates were prepared in 200 µL 2x Laemmli Sample Buffer (0.5

M Tris, pH 6.8, glycerol and 10% SDS) supplemented with a protease and phosphatase inhibitor

cocktail (Complete 1x protease inhibitor (Sigma-Aldrich), Halt phosphatase inhibitor cocktail

(ThermoFisher Scientific). Lysates were heated at 95°C for 10 min, and 10% β-mercaptoethanol

and 5% bromophenol blue were added to the cell lysates. Proteins were resolved by Tris-

Glycine SDS-PAGE and transferred on to a polyvinylidene difluoride (PVDF) membrane. The

PVDF membrane was blocked for 1 h at RT in blocking buffer composed of 5% BSA in 1x final

Tris-buffered saline-Tween (TBS-T; 20 nM Tris, 150 mM NaCl, and 0.1% Tween-20) After

blocking, membranes were washed and incubated with primary antibody (at designated

dilution) overnight at 4°C. After primary incubation, membranes were washed and then

subjected to secondary antibody (at designated dilution) incubation for 1 h at RT. After

incubation with secondary antibody, membranes were washed and imaged using the Chemi-

Doc Imaging System (Bio-Rad). Western blot signals were analyzed and quantified using

ImageLab 6.1 (Bio-Rad). Band intensity was quantified by signal integration in an area

corresponding to the appropriate band.

Statistical Analysis

All experiments were performed at least three independent times with the specific total

number indicated in the respective figure legends along with sample size. The mean and

measure of variation such as standard deviation (STD) or standard error of the mean (SEM) are

indicated. When shown, violin plots indicate range, followed dotted lines to display 25-75%

percentile, the dashed line indicates median of the sampled populations. Sample means of

independent experiments are shown and colour coded to match the independent experiment

across conditions. All experimental conditions were statistically analyzed with the actual tests

disclosed in the respective figure legend. Unless data was normalized, sampling was assumed to

be normally distributed. As such, data was typically assessed with paired Student's t-test when

comparing two populations, with a repeated-measures one-way ANOVA for three or more

samples with one parameter, or repeated-measures two-way ANOVA when two parameters

were considered. The Geisser-Greenhouse correction was applied. Post-hoc tests were

employed based on assumptions and comparisons as recommended by GraphPad Prism and

disclosed in the respective figure legends when comparing multiple conditions in non-

normalized controls. ANOVA tests were coupled to Dunnett's Multiple Comparison Test for

one-way analysis and Tukey's multiple comparison test for two-way analysis, to analyze

pairwise conditions. Here, we accept P<0.05 as indication of statistical significance, but we

disclose all p values for greater transparency.

References

Abrisch, R.G., S.C. Gumbin, B.T. Wisniewski, L.L. Lackner, and G.K. Voeltz. 2020. Fission and
fusion machineries converge at ER contact sites to regulate mitochondrial morphology.
Journal of Cell Biology. 219. doi:10.1083/JCB.201911122.

Allison, R., J.R. Edgar, G. Pearson, T. Rizo, T. Newton, S. Günther, F. Berner, J. Hague, J.W.
Connell, J. Winkler, J. Lippincott-Schwartz, C. Beetz, B. Winner, and E. Reid. 2017.
Defects in ER–endosome contacts impact lysosome function in hereditary spastic
paraplegia. *The Journal of Cell Biology*. jcb.201609033. doi:10.1083/jcb.201609033.

Babu, S., K.A. Nicholson, J.D. Rothstein, A. Swenson, P.J. Sampognaro, P. Pant, E.A. Macklin, S.
Spruill, S. Paganoni, T.F. Gendron, M. Prudencio, L. Petrucelli, D. Nix, S. Landrette, E.
Nkrumah, K. Fandrick, J. Edwards, and P.R. Young. 2024. Apilimod dimesylate in C9orf72
amyotrophic lateral sclerosis: a randomized phase 2a clinical trial. *Brain*. 147:2998–
3008. doi:10.1093/brain/awae109.

Ballabio, A., and J.S. Bonifacino. 2020. Lysosomes as dynamic regulators of cell and organismal
homeostasis. *Nat Rev Mol Cell Biol*. 21:101–118. doi:10.1038/s41580-019-0185-4.

Bao, Y., Y. Qiao, J.E. Choi, Y. Zhang, R. Mannan, C. Cheng, T. He, Y. Zheng, J. Yu, M. Gondal, G.
Cruz, S. Grove, X. Cao, F. Su, R. Wang, Y. Chang, I. Kryczek, M. Cieslik, M.D. Green, W.
Zou, and A.M. Chinnaiyan. 2023. Targeting the lipid kinase PIKfyve upregulates surface
expression of MHC class I to augment cancer immunotherapy. *Proc Natl Acad Sci U S A*.
120:e2314416120. doi:10.1073/pnas.2314416120.

Bissig, C., I. Hurbain, G. Raposo, and G. van Niel. 2017. PIKfyve activity regulates reformation of
terminal storage lysosomes from endolysosomes. *Traffic*. 18:747–757.
doi:10.1111/tra.12525.

Bonet-Ponce, L., and M.R. Cookson. 2022. The endoplasmic reticulum contributes to lysosomal
tubulation/sorting driven by LRRK2. *Mol Biol Cell*. 33:ar124. doi:10.1091/mbc.E22-04-
0139.

Botelho, R.J. 2009. Changing phosphoinositides "on the fly": How trafficking vesicles avoid an
identity crisis. *BioEssays*. 31:1127–1136. doi:10.1002/bies.200900060.

Botelho, R.J., J.A. Efe, D. Teis, and S.D. Emr. 2008. Assembly of a Fab1 phosphoinositide kinase
signaling complex requires the Fig4 phosphoinositide phosphatase. *Mol Biol Cell*.
19:4273–4286. doi:10.1091/mbc.e08-04-0405.

Bridges, D., J.-T. Ma, S. Park, K. Inoki, L.S. Weisman, and A.R. Saltiel. 2012. Phosphatidylinositol
3,5-bisphosphate plays a role in the activation and subcellular localization of

mechanistic target of rapamycin 1. *Molecular biology of the cell*. 23:2955–62.
doi:10.1091/mbc.E11-12-1034.

Cabukusta, B., and J. Neefjes. 2018. Mechanisms of lysosomal positioning and movement.
Traffic. 19:761–769. doi:10.1111/tra.12587.

Cai, S., Y. Wu, A. Guillén-Samander, W. Hancock-Cerutti, J. Liu, and P. De Camilli. 2022. In situ
architecture of the lipid transport protein VPS13C at ER-lysosome membrane contacts.
Proc Natl Acad Sci U S A. 119:e2203769119. doi:10.1073/pnas.2203769119.

Cai, X., Y. Xu, A.K. Cheung, R.C. Tomlinson, A. Alcázar-Román, L. Murphy, A. Billich, B. Zhang, Y.
Feng, M. Klumpp, J.-M. Rondeau, A.N. Fazal, C.J. Wilson, V. Myer, G. Joberty, T.
Bouwmeester, M.A. Labow, P.M. Finan, J.A. Porter, H.L. Ploegh, D. Baird, P. De Camilli,
J.A. Tallarico, and Q. Huang. 2013. PIKfyve, a class III PI kinase, is the target of the small
molecular IL-12/IL-23 inhibitor apilimod and a player in Toll-like receptor signaling.
Chemistry & biology. 20:912–21. doi:10.1016/j.chembiol.2013.05.010.

Cantalupo, G., P. Alifano, V. Roberti, C.B. Bruni, and C. Bucci. 2001. Rab-interacting lysosomal
protein (RILP): The Rab7 effector required for transport to lysosomes. *EMBO Journal*.
20:683–693. doi:10.1093/emboj/20.4.683.

Chang, J., S. Lee, and C. Blackstone. 2013. Protrudin binds atlastins and endoplasmic reticulum-
shaping proteins and regulates network formation. *Proceedings of the National
Academy of Sciences*. 110:14954–14959. doi:10.1073/pnas.1307391110.

Chen, X., C. Shi, M. He, S. Xiong, and X. Xia. 2023. Endoplasmic reticulum stress: molecular
mechanism and therapeutic targets. *Sig Transduct Target Ther*. 8:352.
doi:10.1038/s41392-023-01570-w.

Cheng, C., J. Hu, R. Mannan, T. He, R. Bhattacharyya, B. Magnuson, J.P. Wisniewski, S. Peters,
S.A. Karim, D.J. MacLean, H. Karabürk, L. Zhang, N.J. Rossiter, Y. Zheng, L. Xiao, C. Li, D.
Awad, S. Mahapatra, Y. Bao, Y. Zhang, X. Cao, Z. Wang, R. Mehra, P. Morlacchi, V. Sahai,
M. Pasca di Magliano, Y.M. Shah, L.S. Weisman, J.P. Morton, K. Ding, Y. Qiao, C.A.
Lyssiotis, and A.M. Chinnaiyan. 2025. Targeting PIKfyve-driven lipid metabolism in
pancreatic cancer. *Nature*. doi:10.1038/s41586-025-08917-z.

Chow, C.Y., Y. Zhang, J.J. Dowling, N. Jin, M. Adamska, K. Shiga, K. Szigeti, M.E. Shy, J. Li, X.
Zhang, J.R. Lupski, L.S. Weisman, and M.H. Meisler. 2007. Mutation of FIG4 causes
neurodegeneration in the pale tremor mouse and patients with CMT4J. *Nature*. 448:68–
72. doi:10.1038/nature05876.

Choy, C.H., G. Saffi, M.A. Gray, C. Wallace, R.M. Dayam, Z.Y.A. Ou, G. Lenk, R. Puertollano, S.C.
Watkins, and R.J. Botelho. 2018. Lysosome enlargement during inhibition of the lipid
kinase PIKfyve proceeds through lysosome coalescence. *Journal of Cell Science*.
131:jcs213587. doi:10.1242/JCS.213587.

Dong, X., D. Shen, X. Wang, T. Dawson, X. Li, Q. Zhang, X. Cheng, Y. Zhang, L.S. Weisman, M.
Delling, and H. Xu. 2010. PI(3,5)P(2) controls membrane trafficking by direct activation
of mucolipin Ca(2+) release channels in the endolysosome. *Nature communications*.
1:38. doi:10.1038/ncomms1037.

Dove, S.K., R.C. Piper, R.K. McEwen, J.W. Yu, M.C. King, D.C. Hughes, J. Thuring, A.B. Holmes,
F.T. Cooke, R.H. Michell, P.J. Parker, and M.A. Lemmon. 2004. Svp1p defines a family of
phosphatidylinositol 3,5-bisphosphate effectors. *EMBO J.* 23:1922–1933.
doi:10.1038/sj.emboj.7600203.

Du, Y., W. Chang, L. Gao, L. Deng, and W.-K. Ji. 2023. Tex2 is required for lysosomal functions at
TMEM55-dependent ER membrane contact sites. *J Cell Biol*. 222:e202205133.
doi:10.1083/jcb.202205133.

Friedman, J.R., J.R. DiBenedetto, M. West, A.A. Rowland, and G.K. Voeltz. 2013. Endoplasmic
reticulum-endosome contact increases as endosomes traffic and mature. *Molecular*
Biology of the Cell. 24:1030–1040. doi:10.1091/mbc.E12-10-0733.

Friedman, J.R., L.L. Lackner, M. West, J.R. DiBenedetto, J. Nunnari, and G.K. Voeltz. 2011. ER
tubules mark sites of mitochondrial division. *Science*. 334:358–362.
doi:10.1126/science.1207385.

Gao, Y., J. Xiong, Q.-Z. Chu, and W.-K. Ji. 2022. PDZD8-mediated lipid transfer at contacts
between the ER and late endosomes/lysosomes is required for neurite outgrowth. *J Cell*
Sci. 135:jcs255026. doi:10.1242/jcs.255026.

Gopaldass, N., B. Fauvet, H. Lashuel, A. Roux, and A. Mayer. 2017. Membrane scission driven by
the PROPPIN Atg18. *The EMBO Journal*. 36:e201796859. doi:10.15252/embj.201796859.

Goyal, U., and C. Blackstone. 2013. Untangling the web: mechanisms underlying ER network
formation. *Biochim Biophys Acta*. 1833:2492–2498. doi:10.1016/j.bbamcr.2013.04.009.

Grigoriev, I., S.M. Gouveia, B. van der Vaart, J. Demmers, J.T. Smyth, S. Honnappa, D. Splinter,
M.O. Steinmetz, J.W. Putney, C.C. Hoogenraad, and A. Akhmanova. 2008. STIM1 is a MT-
plus-end-tracking protein involved in remodeling of the ER. *Curr Biol*. 18:177–182.
doi:10.1016/j.cub.2007.12.050.

Guimaraes, S.C., M. Schuster, E. Bielska, G. Dagdas, S. Kilaru, B.R.A. Meadows, M. Schrader, and
G. Steinberg. 2015. Peroxisomes, lipid droplets, and endoplasmic reticulum "hitchhike"
on motile early endosomes. *J Cell Biol*. 211:945–954. doi:10.1083/jcb.201505086.

Guo, Y., D. Li, S. Zhang, Y. Yang, J.-J. Liu, X. Wang, C. Liu, D.E. Milkie, R.P. Moore, U.S. Tulu, D.P.
Kiehart, J. Hu, J. Lippincott-Schwartz, E. Betzig, and D. Li. 2018. Visualizing Intracellular
Organelle and Cytoskeletal Interactions at Nanoscale Resolution on Millisecond
Timescales. *Cell*. 175:1430-1442.e17. doi:10.1016/j.cell.2018.09.057.

Hashimoto, Y., M. Shirane, F. Matsuzaki, S. Saita, T. Ohnishi, and K.I. Nakayama. 2014. Protrudin
regulates endoplasmic reticulum morphology and function associated with the
pathogenesis of hereditary spastic paraplegia. *J Biol Chem*. 289:12946–12961.
doi:10.1074/jbc.M113.528687.

Hong, N.H., A. Qi, and A.M. Weaver. 2015. PI(3,5)P2 controls endosomal branched actin
dynamics by regulating cortactin-Actin interactions. *Journal of Cell Biology*. 210:753–
769. doi:10.1083/jcb.201412127.

Hu, J., Y. Shibata, P.-P. Zhu, C. Voss, N. Rismanchi, W.A. Prinz, T.A. Rapoport, and C. Blackstone.
2009. A class of dynamin-like GTPases involved in the generation of the tubular ER
network. *Cell*. 138:549–561. doi:10.1016/j.cell.2009.05.025.

Ikonomov, O.C., D. Sbrissa, K. Delvecchio, Y. Xie, J.-P.P.J.-P. Jin, D. Rappolee, and A. Shisheva.
2011. The phosphoinositide kinase PIKfyve is vital in early embryonic development:
preimplantation lethality of PIKfyve-/- embryos but normality of PIKfyve+/- mice.
Journal of Biological Chemistry. 286:13404–13413. doi:10.1074/jbc.M111.222364.

Ikonomov, O.C., D. Sbrissa, and A. Shisheva. 2001. Mammalian cell morphology and endocytic
membrane homeostasis require enzymatically active phosphoinositide 5-kinase PIKfyve.
The Journal of biological chemistry. 276:26141–7. doi:10.1074/jbc.M101722200.

Inpanathan, S., and R.J. Botelho. 2019. The Lysosome Signaling Platform: Adapting With the
Times. *Frontiers in Cell and Developmental Biology*. 7:113. doi:10.3389/fcell.2019.00113.

Jin, N., C.Y. Chow, L. Liu, S.N. Zolov, R. Bronson, M. Davisson, J.L. Petersen, Y. Zhang, S. Park, J.E.
Duex, D. Goldowitz, M.H. Meisler, and L.S. Weisman. 2008. VAC14 nucleates a protein
complex essential for the acute interconversion of PI3P and PI(3,5)P(2) in yeast and
mouse. *The EMBO journal*. 27:3221–34. doi:10.1038/emboj.2008.248.

Jongsma, M.L.M., N. Bakker, L.M. Voortman, R.I. Koning, E. Bos, J.J.L.L. Akkermans, L. Janssen,
and J. Neefjes. 2024. Systems mapping of bidirectional endosomal transport through the
crowded cell. *Current Biology*. 34:4476-4494.e11. doi:10.1016/j.cub.2024.08.026.

Jongsma, M.L.M., I. Berlin, R.H.M. Wijdeven, L. Janssen, G.M.C. Janssen, M.A. Garstka, H.
Janssen, M. Mensink, P.A. van Veelen, R.M. Spaapen, and J. Neefjes. 2016. An ER-
Associated Pathway Defines Endosomal Architecture for Controlled Cargo Transport.
Cell. 166:152–166. doi:10.1016/j.cell.2016.05.078.

Kang, Y.-L., Y. Chou, P.W. Rothlauf, Z. Liu, T.K. Soh, D. Cureton, J.B. Case, R.E. Chen, M.S.
Diamond, S.P.J. Whelan, and T. Kirchhausen. 2020. Inhibition of PIKfyve kinase prevents
infection by Zaire ebolavirus and SARS-CoV-2. *Proceedings of the National Academy of
Sciences*. 202007837. doi:10.1073/pnas.2007837117.

Keren-Kaplan, T., A. Sarić, S. Ghosh, C.D. Williamson, R. Jia, Y. Li, and J.S. Bonifacino. 2022.
RUFY3 and RUFY4 are ARL8 effectors that promote coupling of endolysosomes to
dynein-dynactin. *Nature Communications*. 13:1506. doi:10.1038/s41467-022-28952-y.

Khatter, D., V.B. Raina, D. Dwivedi, A. Sindhwani, S. Bahl, and M. Sharma. 2015. The small
GTPase Arl8b regulates assembly of the mammalian HOPS complex on lysosomes.
Journal of cell science. 128:1746–61. doi:10.1242/jcs.162651.

Kumar, G., P. Chawla, N. Dhiman, S. Chadha, S. Sharma, K. Sethi, M. Sharma, and A. Tuli. 2022.
RUFY3 links Arl8b and JIP4-Dynein complex to regulate lysosome size and positioning.
Nature Communications. 13:1540. doi:10.1038/s41467-022-29077-y.

de Lartigue, J., H. Polson, M. Feldman, K. Shokat, S.A. Tooze, S. Urbé, and M.J. Clague. 2009.
PIKfyve regulation of endosome-linked pathways. *Traffic*. 10:883–893.
doi:10.1111/j.1600-0854.2009.00915.x.

Lees, J.A., P. Li, N. Kumar, L.S. Weisman, and K.M. Reinisch. 2020. Insights into Lysosomal
PI(3,5)P2 Homeostasis from a Structural-Biochemical Analysis of the PIKfyve Lipid Kinase
Complex. *Mol Cell*. 80:736-743.e4. doi:10.1016/j.molcel.2020.10.003.

Leray, X., J.K. Hilton, K. Nwangwu, A. Becerril, V. Mikusevic, G. Fitzgerald, A. Amin, M.R. Weston,
and J.A. Mindell. 2022. Tonic inhibition of the chloride/proton antiporter ClC-7 by
PI(3,5)P2 is crucial for lysosomal pH maintenance. *eLife*. 11:e74136.
doi:10.7554/eLife.74136.

Luzio, J.P., P.R. Pryor, and N.A. Bright. 2007. Lysosomes: fusion and function. *Nature reviews.*
Molecular cell biology. 8:622–632. doi:10.1038/nrm2217.

Mas, C., S.J. Norwood, A. Bugarcic, G. Kinna, N. Leneva, O. Kovtun, R. Ghai, L.E. Ona Yanez, J.L.
Davis, R.D. Teasdale, and B.M. Collins. 2014. Structural basis for different
phosphoinositide specificities of the PX domains of sorting nexins regulating G-protein
signaling. *J Biol Chem*. 289:28554–28568. doi:10.1074/jbc.M114.595959.

Matsuzaki, F., M. Shirane, M. Matsumoto, and K.I. Nakayama. 2011. Protrudin serves as an
adaptor molecule that connects KIF5 and its cargoes in vesicular transport during
process formation. *Mol Biol Cell*. 22:4602–4620. doi:10.1091/mbc.E11-01-0068.

Min, S.H., A. Suzuki, T.J. Stalker, L. Zhao, Y. Wang, C. McKennan, M.J. Riese, J.F. Guzman, S.
Zhang, L. Lian, R. Joshi, R. Meng, S.H. Seeholzer, J.K. Choi, G. Koretzky, M.S. Marks, and
C.S. Abrams. 2014. Loss of PIKfyve in platelets causes a lysosomal disease leading to
inflammation and thrombosis in mice. *Nature communications*. 5:4691.
doi:10.1038/ncomms5691.

Min, S.H., A. Suzuki, L. Weaver, J. Guzman, Y. Chung, H. Jin, F. Gonzalez, C. Trasorras, L. Zhao,
L.A. Spruce, S.H. Seeholzer, E.M. Behrens, and C.S. Abrams. 2019. PIKfyve Deficiency in

Myeloid Cells Impairs Lysosomal Homeostasis in Macrophages and Promotes Systemic
Inflammation in Mice. *Molecular and Cellular Biology*. 39. doi:10.1128/mcb.00158-19.

Pankiv, S., E. a. Alemu, A. Brech, J.-A.A. Bruun, T. Lamark, A. Øvervatn, G. Bjørkøy, T. Johansen,
A. Overvatn, G. Bjørkøy, and T. Johansen. 2010. FYCO1 is a Rab7 effector that binds to
LC3 and PI3P to mediate microtubule plus end - Directed vesicle transport. *Journal of
Cell Biology*. 188:253–269. doi:10.1083/jcb.200907015.

Pedersen, N.M., E.M. Wenzel, L. Wang, S. Antoine, P. Chavrier, H. Stenmark, and C. Raiborg.
2020. Protrudin-mediated ER–endosome contact sites promote MT1-MMP exocytosis
and cell invasion. *Journal of Cell Biology*. 219:e202003063. doi:10.1083/jcb.202003063.

Prado, D.S., R.T. Cattley, A.B. Sonego, P. Sutariya, S. Wu, M. Lee, W.C. Boggess, M.J. Shlomchik,
and W.F. Hawse. 2025. The phospholipid kinase PIKFYVE is essential for Th17
differentiation. *J Exp Med*. 222:e20240625. doi:10.1084/jem.20240625.

Raiborg, C., E.M. Wenzel, N.M. Pedersen, H. Olsvik, K.O. Schink, S.W. Schultz, M. Vietri, V. Nisi,
C. Bucci, A. Brech, T. Johansen, and H. Stenmark. 2015a. Repeated ER-endosome
contacts promote endosome translocation and neurite outgrowth. *Nature*. 520:234–8.
doi:10.1038/nature14359.

Raiborg, C., E.M. Wenzel, and H. Stenmark. 2015b. ER –endosome contact sites: molecular
compositions and functions. *The EMBO Journal*. 34:1848–1858.
doi:10.15252/embj.201591481.

Rodríguez-García, R., V.A. Volkov, C.-Y. Chen, E.A. Katrukha, N. Olieric, A. Aher, I. Grigoriev, M.P.
López, M.O. Steinmetz, L.C. Kapitein, G. Koenderink, M. Dogterom, and A. Akhmanova.
2020. Mechanisms of Motor-Independent Membrane Remodeling Driven by Dynamic
Microtubules. *Current Biology*. 30:972-987.e12. doi:10.1016/j.cub.2020.01.036.

Rowland, A.A., P.J. Chitwood, M.J. Phillips, and G.K. Voeltz. 2014. ER contact sites define the
position and timing of endosome fission. *Cell*. 159:1027–1041.
doi:10.1016/j.cell.2014.10.023.

Rudnik, S., S. Heybrock, E. Coyaud, Z. Xu, D. Neculai, B. Raught, V. Oorschot, C. Heus, J.
Klumperman, and P. Saftig. 2024. The lysosomal lipid transporter LIMP-2 is part of
lysosome-ER STARD3-VAPB-dependent contact sites. *J Cell Sci*. 137:jcs261810.
doi:10.1242/jcs.261810.

Saffi, G.T., and R.J. Botelho. 2019. Lysosome Fission: Planning for an Exit. *Trends in Cell Biology*.
29:635–646. doi:10.1016/j.tcb.2019.05.003.

Saita, S., M. Shirane, T. Natume, S.-I. Iemura, and K.I. Nakayama. 2009. Promotion of neurite
extension by protrudin requires its interaction with vesicle-associated membrane
protein-associated protein. *J Biol Chem*. 284:13766–13777.
doi:10.1074/jbc.M807938200.

Saric, A., S.A. Freeman, C.D. Williamson, M. Jarnik, C.M. Guardia, M.S. Fernandopulle, D.C.
Gershlick, and J.S. Bonifacino. 2021. SNX19 restricts endolysosome motility through
contacts with the endoplasmic reticulum. *Nat Commun*. 12:4552. doi:10.1038/s41467-
021-24709-1.

Sbrissa, D., O.C. Ikonomov, Z. Fu, T. Ijuin, J. Gruenberg, T. Takenawa, and A. Shisheva. 2007.
Core protein machinery for mammalian phosphatidylinositol 3,5-bisphosphate synthesis
and turnover that regulates the progression of endosomal transport: Novel Sac
phosphatase joins the ArPIKfyve-PIKfyve complex. *Journal of Biological Chemistry*.
282:23878–23891. doi:10.1074/jbc.M611678200.

Sbrissa, D., O.C. Ikonomov, and A. Shisheva. 2002. Phosphatidylinositol 3-phosphate-interacting
domains in PIKfyve. Binding specificity and role in PIKfyve. Endomenbrane localization.
The Journal of biological chemistry. 277:6073–9. doi:10.1074/jbc.M110194200.

Sharma, G., C.M. Guardia, A. Roy, A. Vassilev, A. Saric, L.N. Griner, J. Marugan, M. Ferrer, J.S.
Bonifacino, and M.L. DePamphilis. 2019. A family of PIKFYVE inhibitors with therapeutic
potential against autophagy-dependent cancer cells disrupt multiple events in lysosome
homeostasis. *Autophagy*. 15:1694–1718. doi:10.1080/15548627.2019.1586257.

Shibata, Y., T. Shemesh, W.A. Prinz, A.F. Palazzo, M.M. Kozlov, and T.A. Rapoport. 2010.
Mechanisms Determining the Morphology of the Peripheral ER. *Cell*. 143:774–788.
doi:10.1016/j.cell.2010.11.007.

Shirane, M., and K.I. Nakayama. 2006. Protrudin induces neurite formation by directional
membrane trafficking. *Science*. 314:818–821. doi:10.1126/science.1134027.

Spits, M., I.T. Heesterbeek, L.M. Voortman, J.J. Akkermans, R.H. Wijdeven, B. Cabukusta, and J.
Neefjes. 2021. Mobile late endosomes modulate peripheral endoplasmic reticulum
network architecture. *EMBO reports*. 22. doi:10.15252/embr.202050815.

Tsuruta, F., E.M. Green, M. Rousset, and R.E. Dolmetsch. 2009. PIKfyve regulates CaV1.2
degradation and prevents excitotoxic cell death. *Journal of Cell Biology*. 187:279–294.
doi:10.1083/jcb.200903028.

Vines, J.H., H. Maib, C.M. Buckley, A. Gueho, Z. Zhu, T. Soldati, D.H. Murray, and J.S. King. 2023.
A PI(3,5)P2 reporter reveals PIKfyve activity and dynamics on macropinosomes and
phagosomes. *Journal of Cell Biology*. 222:e202209077. doi:10.1083/jcb.202209077.

Voeltz, G.K., W.A. Prinz, Y. Shibata, J.M. Rist, and T.A. Rapoport. 2006. A class of membrane
proteins shaping the tubular endoplasmic reticulum. *Cell*. 124:573–586.
doi:10.1016/j.cell.2005.11.047.

Voeltz, G.K., E.M. Sawyer, G. Hajnóczky, and W.A. Prinz. 2024. Making the connection: How
membrane contact sites have changed our view of organelle biology. *Cell*. 187:257–270.
doi:10.1016/j.cell.2023.11.040.

Wang, S., H. Tukachinsky, F.B. Romano, and T.A. Rapoport. 2016. Cooperation of the ER-shaping
proteins atlastin, lunapark, and reticulons to generate a tubular membrane network.
eLife. 5:e18605. doi:10.7554/eLife.18605.

Waterman-Storer, C.M., and E.D. Salmon. 1998. Endoplasmic reticulum membrane tubules are
distributed by microtubules in living cells using three distinct mechanisms. *Current*
Biology. 8:798–807. doi:10.1016/S0960-9822(98)70321-5.

Westrate, L.M., J.E. Lee, W.A. Prinz, and G.K. Voeltz. 2015. Form follows function: the
importance of endoplasmic reticulum shape. *Annu Rev Biochem*. 84:791–811.
doi:10.1146/annurev-biochem-072711-163501.

Wu, H., and G.K. Voeltz. 2021. Reticulon-3 Promotes Endosome Maturation at ER Membrane
Contact Sites. *Developmental Cell*. 56:52-66.e7. doi:10.1016/j.devcel.2020.12.014.

Zolov, S.N., D. Bridges, Y. Zhang, W.-W. Lee, E. Riehle, R. Verma, G.M. Lenk, K. Converso-Baran,
T. Weide, R.L. Albin, A.R. Saltiel, M.H. Meisler, M.W. Russell, and L.S. Weisman. 2012. In
vivo, Pikfyve generates PI(3,5)P2, which serves as both a signaling lipid and the major
precursor for PI5P. *Proceedings of the National Academy of Sciences of the United States*
of America. 109:17472–7. doi:10.1073/pnas.1203106109.

964 **Funding**

965 RJB was a recipient of a Canada Research Chair (#950-232333) with contributions from Toronto

966 Metropolitan University and Faculty of Science. The Natural Sciences and Engineering Research

967 Council (NSERC) of Canada funded this research with Discovery research grant to RJB

968 (RGPIN-2020-04343). NRA was partly funded by an NSERC Undergraduate Student Research

973   **Acknowledgements**

974   We would like to thank Dr. Camilla Raiborg (Oslo University Hospital) for kindly providing the

975   plasmids encoding the GFP fusion of the wild-type and of the FYVE4A mutant of protrudin. We

976   also thank Kimberly Lau at the Sick Kids Imaging Facility for advise on Imaris and 3D

977   reconstruction.

Figure Legends

Figure 1: Overexpression of ER morphogenic proteins does not readily mimic PIKfyve

inhibition in COS-7A cells. (A, B) Confocal images of COS-7A cells labeled with Dextran, Alexa

Fluor-488 (magenta) and transiently expressing mCh-CLIMP63/mCherry (gray) (A) or Alexa

Fluor-546 expressing eGFP-Rtn4a/eGFP (gray) (B). Cells were exposed to either 0.01% vehicle

(DMSO) or 80 nM apilimod for 2 h. **(C-D)** Quantification of lysosome number and volume upon

expression of CLIMP63 to induce ER sheets. **(E-F)** Quantification of lysosome number and

volume upon expression of Rtn4a to promote ER tubulation. Scale bar = 20 μm . All experiments

were repeated four independent times. Data points from matching independent experiments

are color coded. Shown is the mean ± SEM. Data are based on 30-50 transfected cells per

condition per experiment. Data was analyzed using repeated measures two-way ANOVA and

Tukey's multiple comparisons test and p values are shown.

Figure 2: PIKfyve loss of function alters the ER morphology. (A, B) Confocal images of COS-7A

cells transiently transfected with ER-mNeonGreen (gray), labeled with Dextran, Alexa Fluor 647

(magenta), and treated with 0.01% vehicle (DMSO) or 80 apilimod for 2 h. Skeletonize function

was applied respectively to trace ER morphology (A). **(C-F)** Quantification of ER morphology

including ER skeletal length (C), number of ER junctions per cell (D), number of branches per cell

(E), and average branch length (μm) (F). Scale bar: full size = 20 μm, zoom insert = 5 μm. All

experiments were repeated four independent times. Data points from matching independent

experiments are colour coded. Shown is the mean ± SEM. Data are based on 15-20 transfected

cells per condition per experiment. Data was analyzed using a two-tailed paired Student's t-test

and p values are shown.

1008

Figure 3: ER motility is reduced in PIKfyve inhibited cells. (A) Representative time-lapse

confocal images of COS-7A cells transiently transfected with ER-mNeonGreen (gray). Zoom

inserts show ER redistribution as indicated by the white arrow. Cells were exposed to either

0.02% vehicle (DMSO) or 240 nM apilimod for 2 h. **(B-D)** Quantification of membrane

displacement analysis categorized by low movement (B), intermediate movement (C), and rapid

movement (D). Scale bar: full size = 20 μm, zoom insert = 5 μm. All experiments were repeated

five independent times. Data points from matching independent experiments are colour coded.

Shown is the mean ± SEM. Data are based on 7 randomized ROIs for 7-10 transfected cells, per

condition per experiment. Data was analyzed using a two-tailed paired Student's t-test and p

values are shown.

1019

Figure 4. ER-remodeling via hitchhiking is reduced upon PIKfyve inhibition. (A) Confocal

images of HeLa cells immunostained for ⬚-tubulin (gray) treated with 0.02% vehicle (DMSO),

240 nM apilimod or 1 μM YM-201636 for 2 h. **(B-D)** Quantification of microtubule morphology

including number of branches per cell (B), number of junctions per cell (C), and average branch

length (D). For B through D, all experiments were repeated four independent times. **(E)**

Confocal images of COS-7A cells transiently transfected with EB1-GFP displayed as Rainbow

RBG LUT and treated as in A. **(F-H)** Semi-quantification of microtubule dynamics including

percentage of cells with >10 EB1-puncta (F), percentage of cells with mobile EB1-puncta (G),

and percentage of cells with EB1-puncta moving away from the nucleus (H). For F-H, all

experiments were repeated three independent times. For B-D and F-H, data was analyzed using

an Ordinary one-way ANOVA test and Dunnett's multiple comparisons test. **(I)** Quantification of

the number of ER-hitchhiking events that occurred in a 100 μm^2 ROI over one minute, repeated

three independent times. **(J)** Confocal images of COS-7A cells transiently transfected with ER-

mNeonGreen (gray) and labeled with Dextran, Alexa Fluor 647 (magenta) and exposed to either

0.02% vehicle (DMSO) or 240 nM apilimod. Zoom inserts show an ER-hitchhiking event

occurring over 2.5 seconds as indicated by the white arrow. Data were analyzed using a two-

tailed paired Student's t-test and p values are shown. Data points from matching independent

experiments are color coded. Shown is the mean ∓ SEM. Scale bar: full size = 20 μm, zoom

insert = 5 μm. P values are shown.

1039

Figure 5. PIKfyve activity supports lysosome mobility and spatial distribution. (A) Confocal

images of COS-7A cells labeled with Dextra, Alexa Fluor 647 (magenta) and exposed to either

0.02% vehicle (DMSO) or 240 nM apilimod for 2 h. Scale bar = 20 μm. **(B)** Schematic

representation of the changes to lysosome spatial distribution between control and PIKfyve

inhibited conditions. **(C)** Quantification of the distribution of lysosomes in the perinuclear

region versus the peripheral region of the cell. A ratio of the number of lysosomes in the

perinuclear region to total number of lysosomes in the cell was determined – the same was

done for the peripheral region respectively. **(D)** Kymograph of lysosome movement along a

linear ROI. The vertical axis represents time (s), and the horizontal axis represents distance

along a selected path. Diagonal or segmented lines indicate motility; vertical lines represent

stationary organelles. **(E-G)** Quantification of lysosome motility including track mean speed (E),

total distance travelled (F), and track displacement (G). All experiments were repeated four

independent times. Data points from matching independent experiments are colour coded.

Shown is the mean ± SEM. Data were analyzed using a two-way ANOVA and Tukey's multiple

comparison test for (C) and two-tailed paired Student's t-test for (E-G) and p values are shown.

1055

Figure 6. Loss of PIKfyve activity alters Protrudin-VapA-Rab7 ER-lysosome contact site. (A-D)

Confocal images of HeLa cells treated with 0.02% vehicle (DMSO), 240 nM apilimod, or 1 μM

YM-201636 for 2 h followed by Proximity Ligation Assay between protrudin and VapA (A), or

Rab7 and VapA (B), or Rab7 and protrudin (C), or Rab7 and Kinesin-1 (KIF5B; D). Images were

generated by maximum projection of z-stacks of the PLA signal (magenta); DAPI-stained nucleus

is shown in gray. Negative controls where one of the primary antibodies was missing are

shown. Scale bar = 15 μm. **(E-H)** Quantification of the number of PLA dots per cell for each

protein pair tested in A-D, as indicated in image. All experiments were repeated four

independent times with matching independent experiments colour coded. Shown is the mean

± SEM. Data were analyzed using an Ordinary one-way ANOVA and Dunnett's multiple

comparisons test and p values are shown.

1067

Figure 7. Protrudin hyper-tethers ER and lysosomes during PIKfyve inhibition to alter ER

morphology. (A) Confocal images of HeLa cells co-transfected with non-targeting siRNA control

or siRNA oligonucleotides against protrudin, and protrudin^{wt}-GFP or protrudin^{FYVE4A}-GFP (gray).

Cells were labeled with Dextran, Alexa 546 (magenta) and then exposed to either 0.02% vehicle

(DMSO) or 240 nM apilimod for 2 h. Scale bar: full size = 20 μm, zoom insert = 5 μm. **(B-D)**

Quantitative analysis of ER morphology based on protrudin^{wt}-GFP or protrudin^{FYVE4A}-GFP

fluorescence in conditions shown in A: number of protrudin branches per cell (B), junctions per

cell (C), and average branch length (D). **(E, F)** Quantification of number of PLA dots between

Rab7 and protrudin (E) and Rab7 and VapA (F) in control silenced and protrudin-silenced cells

with and without apilimod. **(G, H)** Quantification of PLA dots between Rab7 and protrudin (G)

and Rab7-VapA (H) in cells treated with vehicle, apilimod, or Vps34-IN to abate PtdIns(3)P

levels. **(I)** Quantification of PLA dots between Rab7 and protrudin in mock-silenced or

protrudin-silenced cells expressing protrudin^{wt}-GFP or protrudin^{FYVE4A}-GFP exposed to either

vehicle or apilimod. All experiments were repeated three independent times and colour-code

matched. Shown is the mean $\pm$ SEM. Data for (B-F) and (I) were analyzed using two-way

ANOVA and Tukey's multiple comparison test; Data for (G-H) were analyzed using a one-way

ANOVA and Tukey's multiple comparison test. P values are shown.

1085

Supplementary Figure Legend

Supplementary Figure 1. Voids in ER architecture are filled with vacuolated endosomes and

lysosomes upon PIKfyve inhibition. (A-B) Confocal images of COS-7A cells labeled with Dextran,

Alexa Fluor 647 (magenta) and transiently expressing ER-mNeonGreen (gray) or mCherry (cyan)

(A) or mRFP-Rab5 (cyan) (B). Cells were exposed to either 0.01% DMSO or 80 nM apilimod for 2

h. Scale bar: full size = 20 μm, zoom insert = 5 μm. **(C-D)** Quantification of Mander's Analysis

done on the colocalization between Dextra, Alexa 647 and mRFP-Rab5. All experiments were

repeated three independent times. Data points from matching independent experiments are

colour coded and shown as mean $\pm$ SEM from 30-50 transfected cells per condition per

experiment. Data were analyzed using a two-tailed Students t-test and p values are shown.

1096

Supplementary Figure 2. ER morphology and dynamics are altered upon PIKfyve inhibition

using YM-201636. (A) Representative time-lapse confocal images of COS-7A cells transiently

transfected with ER-mNeonGreen (gray). Zoom inserts show ER redistribution as indicated by

the white arrow and Skeletonize function was applied respectively to trace ER morphology. **(B)**

Confocal images of COS-7A cells transiently transfected with ER-mNeonGreen (gray) and

labelled with Dextran, Alexa-647 (magenta). Cells were exposed to either 0.02% vehicle (DMSO)

or 1 µM YM-201636 for 2 h. Scale bar: full size = 20 µm, zoom insert = 5 µm. **(C-F)**

Quantification of ER morphology including ER skeletal length (C), number of ER junctions per

cell (D), number of branches per cell (E), and average branch length (F). **(G-I)** Quantification of

membrane displacement analysis categorized by low movement (G), intermediate movement

(H), and rapid movement (I). Experiments (C-F) were repeated four independent times;

Experiments (G-I) were repeated six independent times. Data points from matching

independent experiments are colour coded. Shown in the mean $\pm$ SEM. Data from C through F

are based on 15-20 transfected cells per condition per experiment. Data from G through I are

based on 7 randomized ROIs for 7-10 transfected cells, per condition per experiment. Data

were analyzed using a two-tailed Student's t-test and p values are shown.

1113

Supplementary Figure 3. Upon PIKfyve inhibition Rab7 GTPase status appears unaltered. (A)

HeLa cells immuno-stained for LAMP1 (magenta) and Rab7 (gray) following treatment with

0.02% vehicle (DMSO) and 240 nM apilimod for 2 h. **(B-C)** Quantitative analysis displaying the

average Mander's Coefficient for the fraction of Rab7 colocalized with LAMP1 (B), and fraction

of LAMP1 colocalized with Rab7 (C). **(D)** Representative Western blots of affinity purification of

Rab7-GTP using GST-RILPC33, displaying total Rab7 levels from input samples and GTP-Rab7

levels recovered with the GST negative control and GST-RILPC33 from cells exposed to 0.02%

vehicle (DMSO), 240 nM apilimod or 1 µM YM-201636 for 2 h. **(E)** Quantitative analysis of

affinity precipitated GTP-Rab7 from GST negative control and GST-RILPC33. GST-control and

GST-RILPC33 were normalized first to input levels of β-actin and then Rab7. All experiments

were repeated three independent times. Data points from matching independent experiments

are colour coded and shown as the mean ± STD. Data from B through C was analyzed using a

two-tailed Student's t-test. Data from E was analyzed using a two-way ANOVA and Tukey's

multiple comparisons test and p values are shown.

1128

Supplementary Figure 4. Three-dimensional reconstruction of protrudin-positive ER. (A)

Three-dimensional reconstruction of HeLa cells co-transfected with non-targeting siRNA control

or siRNA oligonucleotides against protrudin, and expressing protrudin^wt-GFP or protrudin^FYVE4A-

GFP (gray). Cells were labeled with Dextran, Alexa 546 (magenta) and then exposed to either

0.02% vehicle (DMSO) or 240 nM apilimod for 2 h. Scale bar: full frame = 20 μm, zoom insert =

0.7 μm. **(B)** Representative Western blots of non-targeting siRNA control or siRNA

oligonucleotides against protrudin displaying protrudin and clathrin levels. **(C)** Quantitative

analysis displaying protrudin protein levels upon transfection with siNT or siProtrudin after

normalized to loading control, clathrin. All experiments were repeated three independent

times. Data points from matching independent experiments are colour coded. Shown in the

mean ± STD. Data was analyzed using a two-tailed Student's t-test and p values are shown.

1140

Supplementary Video Legend

Supplemental Video 1. ER motility is retained upon treatment with vehicle control (DMSO).

COS-7A cells were transiently transfected with ER-mNeonGreen (gray) and treated with 0.02%

vehicle (DMSO) for 2 h. The ER network undergoes continuous remodeling through tubule

extension, retraction, and branching, reflecting its highly dynamic and interconnected nature.

Time-lapse images were acquired at a rate of 4 frame/s for 1 minute. Still frames from Video 1

are displayed in Figure 3A. Scale bar: 20 µm.

1148

1149   **Supplemental Video 2. ER motility is reduced upon acute treatment with PIKfyve inhibitor**

1150   **Apilimod.** COS-7A cells were transiently transfected with ER-mNeonGreen (gray) and treated

1151   with 240 nM apilimod for 2 h. Time-lapse images were acquired at a rate of 4 frame/s for 1

1152   minute. Still frames from Video 2 are displayed in Figure 3A. Scale bar: 20 µm.

1153

1154   **Supplemental Video 3. ER motility is reduced upon acute treatment with PIKfyve inhibitor**

1155   **YM-201636.** COS-7A cells were transiently transfected with ER-mNeonGreen (gray) and treated

1156   with 1 µM YM-201636 for 2 h. Time-lapse images were acquired at a rate of 4 frame/s for 1

1157   minute. Still frames from Video 3 are displayed in Supplemental Figure 2A. Scale bar: 20 µm.

1158

Supplemental Video 4. EB1-GFP dynamics under vehicle condition conditions. COS-7A cells

were transiently transfected with EB1-GFP displayed as a rainbow LUT scale, where red is the

highest intensity level and blue is the lowest intensity level. Cells were treated with 0.02%

vehicle (DMSO) for 2 h. EB1-GFP comets mark the plus-ends of growing microtubules, enabling

visualization of dynamic microtubules. Time-lapse images were acquired at a rate of 4 frame/s

for 1 minute. Still frames from Video 4 are displayed in Figure 4E. Scale bar: 20 µm.

1165

1166 **Supplemental Video 5. EB1-GFP dynamics appear unaltered upon PIKfyve inhibition using**

1167 **apilimod.** COS-7A cells were transiently transfected with EB1-GFP displayed as a rainbow LUT

1168 scale, where red is the highest intensity level and blue is the lowest intensity level. Time-lapse

1169 images were acquired at a rate of 4 frame/s for 1 minute. Still frames from Video 5 are

1170 displayed in Figure 4E. Scale bar: 20 μm.

1171

1172 **Supplemental Video 6. Microtubule EB1 dynamics remain unaffected by YM-201636**

1173 **treatment.** COS-7A cells were transiently transfected with EB1-GFP displayed as a rainbow LUT

1174 scale, where red is the highest intensity level and blue is the lowest intensity level. Time-lapse

1175 images were acquired at a rate of 4 frame/s for 1 minute. Still frames from Video 6 are

1176 displayed in Figure 4E. Scale bar: 20 μm.

1177

Supplemental Video 7. Normal ER-movement via organelle hitchhiking in vehicle-treated

cells. COS-7A cells were transiently transfected with ER-mNeonGreen (gray) and labeled with

Dextran, Alexa 647 (magenta) followed by treatment with 0.02% vehicle (DMSO) for 2 h.

Positive hitchhiking events were scored based on ER movement driven by physical association

with a motile dextran labelled lysosome. Timelapse images were acquired at a rate of 3.3

frames/s for 1 minute. Still frames from Video 7 are displayed in Figure 4J. Scale bar: 20 μm.

1184

Supplemental Video 8. ER-hitchhiking frequency drops upon PIKfyve inhibition. COS-7A cells

were transiently transfected with ER-mNeonGreen (gray) and labeled with Dextran, Alexa 647

(magenta) followed by treatment with 240 nM apilimod for 2 h. Positive hitchhiking events

were scored based on ER movement driven by physical association with a motile dextran

labelled lysosome. Timelapse images were acquired at a rate of 3.3 frames/s for 1 minute. Still

frames from Video 7 are displayed in Figure 4J. Scale bar: 20 µm.

1191

FIGURE 1.

FIGURE 2.

FIGURE 3.

FIGURE 4.

FIGURE 6.

FIGURE 7.