Whole blood transcriptional signatures of age and survival identified in Long Life Family and Integrative Longevity Omics Studies

Mengze Li [1,2], Zeyuan Song [3], Eric Reed [4], Tanya T. Karagiannis [3,5], Stacy Andersen [6], Michael Brent [7], Chase Mateusiak [7], Sandeep Acharya [7], Wooseok J. Jung [7], Shu Liao [7], Mary K. Wojczynski [8], Mary F. Feitosa [8], Jeffrey R. O'Connell [9], May E. Montasser [9], Roland J. Thorpe Jr [10], Konstantin Arbeev [11], Sofiya Milman [4,12], Albert Tai [5], Thomas T. Perls [6], Paola Sebastiani [3,5]*, Stefano Monti [1,2,13]*

Bioinformatics Program, Boston University, Boston 02215, MA, USA.

Section of Computational Biomedicine, School of Medicine, Boston University, Boston 02118, MA, USA.

Institute for Clinical Research and Health Policy Studies, Tufts Medical Center, Boston, MA 02111, USA.

Department of Medicine, Institute for Aging Research, Albert Einstein College of Medicine, Bronx, NY 10461, USA.

Department of Medicine, School of Medicine, Tufts University, Boston, MA 02111, USA.

Department of Medicine, Chobanian & Avedisian School of Medicine, Boston University, Boston 02118, MA, USA.

Department of Computer Science and Engineering, Washington University, St Louis, MO 63130, USA.

Department of Genetics, Washington University School of Medicine, St. Louis, MO 63130, USA.

Department of Medicine, University of Maryland School of Medicine, Baltimore, MD 21201, USA.

Program for Research on Men's Health, Hopkins Center for Health Disparities Solutions, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD 21205, USA.

Social Science Research Institute, Duke University, Durham, NC 27708, USA.

Department of Genetics, Institute for Aging Research, Albert Einstein College of Medicine, Bronx, NY 10461, USA.

Department of Biostatistics, Boston University School of Public Health, Boston, 02118, MA

* Corresponding author

Author's notes:

This manuscript has been peer-reviewed and accepted by GeroScience (Springer). This bioRxiv article
reflects the published version, incorporating revisions made in response to reviewers' comments. The
main content, results, and conclusions remain unchanged from the previous version, while the
Discussion section has been expanded to further address the functional annotation and clinical
relevance of the identified markers.

Copy of the acceptance letter:

The editors are pleased to inform you that your manuscript, JAAA-D-25-01946R1 entitled "Whole blood
transcriptional signatures of age and survival identified in Long Life Family and Integrative Longevity
Omics Studies" has been accepted for publication in GeroScience as an Original Article. The editors
commend you on your outstanding contribution to the journal. Your manuscript can be published online
ahead of print within approximately two weeks.

Abstract

Although aging is a universal event, some individuals are able to achieve extreme longevity. The Long-Life Family Study (LLFS) enrolls participants from families enriched with long-lived individuals, serves as a valuable dataset for studying ageing phenotypes and identify potential intervention targets. We analyzed the association between age at blood draw and 16,284 RNAseq-based blood transcriptomic data from 2,167 LLFS participants with ages ranging from 18 to 107, replicated the results in the Integrative Longevity Omics Study (ILO) dataset of 20,884 RNAseq-based blood transcriptomic data from 419 participants, with ages ranging from 60 to 108, and further compared our findings to a published reference aging signature.

We identified 4,227 transcripts increasing and 4,044 transcripts decreasing with age, and enrichment analysis revealed age-related upregulation of inflammatory and senescence-related pathways, and downregulation of MYC and Wnt/β-catenin targets, among others. Further, a subset of transcripts showed age associations unique to the longevity-enriched cohorts (LLFS and ILO). We also identified 314 transcripts significantly associated with mortality risk and found that pro-survival gene sets included NK cell-mediated cytotoxicity and GPCR signaling. Finally, increased transcriptomic age predicted using transcriptomic clock was strongly associated with increased mortality. In summary, this study identified robust transcriptomic signatures of aging and mortality in a longevity-enriched population, highlighting key biological pathways such as immune modulation, inflammation, and senescence.

Keywords

Aging, whole blood, transcriptomics, aging clock, mortality risk, longevity-enriched population.

Introduction

The biology of aging is complex and multifaceted [1] [2], with age-related changes observed across various hallmarks and omics-based profiles [1] [2] [3]. Given the growing elderly population and the increased risk for morbidity at older ages [4], significant efforts have been devoted to better understand human aging at the molecular level and to identify interventions that can maintain health [5]. Blood is a minimally invasive and reliable source of biomarkers for research and clinical diagnostics and many markers have been identified in plasma or serum [6], [7] [8]. Additionally, single-cell analyses have shown that the blood cell-type composition changes substantially during aging [9].

Aging is associated with increased risk for morbidity and mortality. However, some individuals can postpone or delay age-related diseases and extend their health span to extreme old age [10] [11], [12], [13]. Evidence is growing that these individuals harbor a combination of genetic and genomic features

that either promote or reflect the biological mechanisms underlying this phenomenon. For example,
we discovered unique protein signatures in centenarians [14], and distinctive metabolic profiles in
long lived individuals [8]. Whole-genome sequencing analysis of semi-supercentenarians and
supercentenarians has uncovered unique somatic genetic mutations in these individuals [15]. These
findings underscore the value of studying long-lived individuals to uncover protective mechanisms to
achieve longevity. As a critical omics layer, transcriptomic profiling holds significant potential to
provide valuable insights into the biological mechanisms of aging and to identify potential
therapeutic targets.

A previous study of the blood transcriptome of 14,983 individuals of European ancestry by Peters et
al., identified 1,497 genes differentially expressed with chronological age, and found these genes to be
enriched for the presence of potentially functional CpG-methylation sites in enhancer and insulator
regions [7]. A second study of blood specimens of 3,388 adult individuals showed the ability of the
transcriptome to identify molecular sub-types distinguished by their sex, age, and immune and
metabolic status [16]. In all of these studies, the analysis was conducted on age ranges typical of the
general population, thus excluding extreme old age individuals. Expanding these analyses to include
long-lived subjects could offer valuable insights into the protective mechanisms underlying healthy
aging and extreme longevity. The Long Life Family Study (LLFS) has collected whole blood
transcriptomic data from individuals in families enriched with long-lived members, offering a unique
opportunity to study the transcriptional changes associated with extreme old age.

An emerging and related area of research is the evaluation of biological age as distinct from
chronological age, recognizing that the rate of aging varies among individuals and that chronological
age is not always a reliable indicator of health status. To address this question, omics-based aging
clocks have been developed. Further exploration of the potential of transcriptomic clocks to measure
various aspects of aging and extreme longevity in humans could provide valuable insights into the
biological mechanisms associated with and/or contributing to these processes.

In this study, we analyzed the whole blood transcriptomic profile of 2,167 individuals from longevity-
enriched families, with the aim of identifying new gene markers and leading biological pathways
regarding human aging, extreme longevity, and mortality risk. Additionally, a transcriptomic aging
clock was constructed using elastic net and was bias-corrected [17], and the predicted age
acceleration was analyzed in relation to mortality risk.

Methods

Study participants.

The Long Life Family Study (LLFS) enrolled families with evidence of exceptional longevity across the
United States and Denmark between 2006 and 2009. Study participants are followed longitudinally
until death. We described study design, genetic, genomic and phenotypic data previously [18]. The
Integrative Longevity Omics (ILO) study enrolled ~1400 centenarians, their biological offspring, and

spouses of the offspring between 2019 and 2024 from North America. Study participants completed a
comprehensive assessment of physical and cognitive functions, and answered questionnaires on
medical history and medication use, which are harmonized to LLFS. LLFS was approved by the
Washington University in St. Louis IRB. ILO was approved by Albert Einstein College of Medicine IRB.
All participants provided informed consent.

RNA-seq data generation and QC.

The RNA-seq data in the LLFS was generated from March 2021 to November 2023 at Washington
University in St. Louis. Details are provided in [19]. Briefly, total RNA was extracted from PAXgene™
Blood RNA tubes using the Qiagen PreAnalytiX PAXgene Blood miRNA Kit (Qiagen, Valencia, CA). The
Qiagen QIAcube extraction robot performed the extraction according to the company's protocol. The
RNA-Seq data were processed with the nf-core/rnaseq pipeline version 3.3 using STAR/RSEM [20].
Samples with high intergenic read percentage (> 8%), suspicious sex chromosome gene expression, or
without sufficient genotype or phenotype information were excluded. Transcripts with less than 10
counts per million in at least 3% of samples across the original 4189-sample series were excluded,
yielding a final set of 16,284 transcripts from 2,167 subjects for the downstream analysis.

The RNA-seq profiles of 434 ILO participant blood samples were generated from December 2023 to
February 2024 in 10 batches, prepared on separate days, at the Tufts University Core Facility
Genomics Core in Boston, MA. Total RNA was extracted from PAXgene™ Blood RNA tubes using the
QIASymphony PAXgene Blood RNA Kit (Qiagen, Valencia, CA). The Qiagen QIASymphony SP/AS nucleic
acid isolation robotic system performed the extraction according to the company's protocol. Quality
of the isolated RNA samples was checked by measuring their RNA quality numbers and concentrations
using the Agilent Fragment Analyzer (Agilent, Santa Clara, CA). RNA-seq libraries were prepared using
the Illumina Stranded Total RNA with RiboZero Plus preparation kit (Illumina, San Diego, CA). RNA-seq
libraries were sequenced using the Illumina NovaSeq X Plus (Illumina, San Diego, CA), generating 150-
base paired-end reads. Raw RNA-seq reads were processed using the nf-core/rnaseq pipeline (version
3.14) [20]. Main steps included adapter trimming, quality control, and alignment steps. Nextera
transposase adapter sequences were removed using TrimGalore (version 0.6.7) [21] [22]. Trimmed
reads were aligned to the human reference genome, GRCh38, using STAR version 2.7.9a [23], followed
by gene expression quantification using Salmon version 1.10.1 [24]. In addition, the genomic origins of
all reads, which could be one of exonic, intronic, or intergenic, were quantified using Qualimap
version 2.3 [25]. Following read processing, low-quality samples were flagged based on various quality
control metrics, including total reads, total aligned reads, total reads counted in genes, percentage of
total reads that were aligned, percentage of aligned reads that were counted, percentage of intergenic
reads, and the total number of genes with non-zero counts. For each metric, samples were flagged
based on interquartile range outlier detection of all samples, and samples that flagged as outliers in
multiple metrics were excluded from further analysis. As a result, 419 of the 434 profiles were retained.
Following sample filtering, transcripts were excluded if they were poorly quantified in any of the 10
batches, defined as having zero counts in at least 20% of samples within at least one batch. As a
result, 20,884 of the 57,890 transcripts were retained.

For both LLFS and ILO transcriptomic data, raw transcript counts were normalized using the R
package *DESeq2* [26] and log2-transformed.

Genotype data.

To control for population structure and familial relatedness among the study participants, we
calculated genome-wide principal components (PCs) and genetic relationship matrix (GRM) from the
LLFS whole genome sequencing data released in December 2021 using the pipeline described in [27].
The genetic data were described in [28].

Transcriptomic signature of chronological age.

In LLFS data, we applied a linear mixed-effects model to regress each transcript's expression level
against age at blood draw using the *fitNullModel* function in the R package *GENESIS* [29]. The model
included a random intercept and specified the variance-covariance structure of the random effects
using the GRM to account for participant relatedness. The initial dataset comprised 2,167 participants;
after excluding individuals with incomplete covariate data, we retained 2,101 participants from 183
families for the final analysis (Table 1). Each gene-level model incorporated covariate adjustments for
sex, education level, recruitment center, RNA-seq instrument batch, percentage of intergenic reads,
and the first ten genotype-derived principal components.

1,786 participants had medication information available, indicating whether they were taking
medications for hypertension, heart disease, high cholesterol, and diabetes (Figure S1 A). To assess
the impact of medication on gene expression levels, we repeated the analysis, including these four
medication groups as additional covariates to assess their effects on the association between gene
expression and age. The results showed no significant effect of medication on gene expression (Figure
S1B, C). Therefore, medication was not included as a covariate in the subsequent analyses.

In ILO data, we applied a linear model to regress each transcript's expression levels against age at
blood draw using the *glm()* base R function. Each gene-level model incorporated covariate
adjustments for sex, education level, profile preparation date batch, and percentage of intergenic
reads. For each gene, profiles with outlier expression levels were excluded from modeling, which was
defined as exceeding four standard deviations from the 20% trimmed mean expression of all profiles.

For both analyses of LLFS and ILO transcriptomic data, we corrected nominal p-values for multiple
testing using the Benjamini-Hochberg false discovery rate (FDR) procedure (q-values).

Replication of the transcriptomic signature of chronological age.

We compared our results with the transcriptomic signature of age reported by Peters et.al. [7] and the
dataset in the ILO study. Genes were then categorized according to whether they increased or
decreased with age, and Fisher's exact tests were performed separately for the two groups to evaluate
the significance of the overlap between the marker genes in the two datasets. The background for

both tests was defined as the intersection of genes available in both studies. When comparing our
study to Peters et.al., this overlap included 9,463 genes; for the comparison with ILO, the overlap was
14,728 genes. The null hypothesis of Fisher's exact test is that a gene's association with age in one
study is independent of its association in the other study. Therefore, a significant deviation from this
null hypothesis would indicate a successful replication of age-associated gene markers between the
two studies.

Transcriptomic signature of mortality risk.

We conducted a survival analysis to identify gene transcripts that predict mortality risk. This analysis
included 2,057 participants, 698 of whom were deceased (Table 1). A Cox-proportional hazards (Cox-
PH) model was fitted for each gene transcript data using the *coxph* function from the R package
survival [30]. Pedigree ID was included as the cluster term to account for family membership. In this
analysis, we used time to death as the outcome of interest, and batch-corrected and standardized
gene expression data as the predictor. We adjusted the model by age at enrollment, sex, education
level, enrollment center, and percentage of intergenic reads in the RNA-seq experiment. To obtain the
batch-corrected gene expression values, we first fitted a linear regression model for each gene, using
the batch indicator as a predictor. The residuals from this model were then extracted and
standardized to have a mean of 0 and a standard deviation of 1.

Gene set enrichment analysis.

We mapped Ensembl IDs of the mRNAs to gene symbols using biomaRt [31] and retrieved gene set
collections from MSigDB [32], including HALLMARK, KEGG, and REACTOME. Gene set enrichment
analysis was performed with hypeR [33] using a hypergeometric test. Gene sets with q ≤ 0.05 after FDR
correction were considered significant and retained. We applied fgsea [34] to perform the rank-based
Kolmogorov-Smirnov (KS) test, where the gene sets were ranked using their Z-score. Additionally, to
specifically assess cellular senescence, we tested enrichment of the SenMayo gene set [35] with the KS
test using fgsea. There were 125 genes included in the SenMayo gene set, and 67 of them passed the
QC filter in the RNA-seq preprocessing and used for enrichment analysis.

Co-expression analysis.

We used Weighted Gene Co-expression Network Analysis (WGCNA) [36] to detect gene co-expression
modules among the 2,550 genes most largely affected by age (absolute Z-score > 6, q ≤ 0.01). We first
obtained residuals by regressing each transcript on age, sex, education level, recruitment center, RNA-
seq batch number, and percentage of intergenic reads in the RNA-seq experiment. We then performed
module detection using these residuals. The clustering method was selected to be "average", and the
correlation type was set to be "signed". We defined the hub gene in each module as the gene that has
the highest correlation with the module's eigengene.

Construction of a transcriptomic aging clock.

We generated an aging clock using standardized (mean = 0, SD = 1) RNA expression levels, with
chronological age as the response variable. We trained the model on data from 2,167 participants,
using the elastic net method implemented in the R package *glmnet* [37]. We used cross-validation to
determine optimal α and λ values, with alpha values ranging between 0 and 1 in increments of 0.05.
The model yielding the lowest RMSE was chosen, corresponding to α = 0.05 and λ = 2.215. The
importance of each of the genes included in the model was estimated using the permutation method,
as implemented in the R package *vip* [38]. Predicted ages by the model showed a systematic bias, with
younger individuals being overestimated in predicted age and older individuals being
underestimated. To address this bias, we employed the correction proposed by Cole et al. [17], [39].
Survival analysis based on a Cox-PH model was performed using delta-age, defined as the difference
between corrected predicted age and chronological age, as the regression variable, with a positive
(negative) value predicting the subject to be biologically older (younger). The same set of covariates
used in the survival signature analysis was included in the model.

235 Results

Participants' characteristics.

Pre-processing and quality-control procedures yielded robust quantification of 16,284 transcripts
across 2,167 individuals from 184 families in the LLFS cohort, with participant ages ranging from 18 to
107 years (Table 1). The ILO cohort comprised 419 individuals with ages ranging from 60 to 108 years
(Table 1).

Chronological age impacts the expression of multiple genes.

In the LLFS cohort, we identified 4,227 age-associated mRNA transcripts mapping to 3,777 genes
whose expression increased with age, and 4,044 transcripts mapping to 3,710 genes whose expression
decreased with age (1% FDR, corrected using Benjamini-Hochberg false discovery rate correction,
Figure 1). The top 20 genes associated with age are shown in Table 2 and the results for all genes are
available in Table S1.

The expression level changes with age of several notable example genes are visualized in Figure 2A,
and the volcano plot illustrating all effect sizes and significance of the analyzed genes is shown in
Figure 2B. These include genes decreasing with age such as CD248 (Est = -0.0476, q = 4.81e-179),
implicated in extracellular matrix binding activity and cell migration; 5'-Nucleotidase Ecto (NT5E, Est =
-0.027, q = 3.55e-95), pivotal in lymphocyte differentiation and the calcification process in joints and
arteries; and the MYC proto-oncogene (MYC, Est = -0.012, q = 3.28e-62), acting as a proto-oncogene
and contributing to cell cycle progression and apoptosis [40].

Many of the most significant gene markers that decreased with age are involved in immune response
and immune cell development. Examples include the NEDD4 (neural precursor cell expressed
developmentally down-regulated protein 4) binding protein 3 (N4BP3, Est = -0.0219, q = 1.05e-92),

involved upstream of the innate immune response and shown to regulate antiviral innate immune

signaling pathways [40] [41]; the chemokine receptor 6 (CCR6, Est = -0.0202, q = 2.03e-76), expressed

in various immune cells and shown to guide cell migration during immune responses [42]; the T-Cell

specific transcription factor 7 (TCF7, Est = -0.0146, q = 1.82e-74), crucial for the development of natural

killer cells and innate lymphoid cells; CD27 (Est = -0.0146, q = 2.01e-62), required for T cell

maintenance and B cell activation; and CD28 (Est = -0.0141, q = 2.74e-60), critical for T-cell

proliferation and survival [40].

In contrast, zinc finger e-box-binding homeobox 2 (ZEB2) was among the most significantly

upregulated genes with age (Est = 0.0087, q = 7.12e-65). As a transcription factor, ZEB2 plays a critical

role in epithelial-to-mesenchymal transition [43], is essential for the differentiation of age-associated

B cells, and has been implicated in autoimmune pathogenesis [44]. Notably, studies in mice have

shown that silencing Zeb2-NAT enhances the reprogramming efficiency of aged fibroblasts,

underscoring its potential as a therapeutic target for preserving stem cell pluripotency [45]. Another

gene, leucine rich repeat containing G protein-coupled receptor 6 (LGR6, Est = 0.0198, q = 1.53e-34),

was associated with chronic activation of the Wnt/β-catenin pathway, a process that promotes

cellular senescence, and exacerbates the progression of chronic obstructive pulmonary disease

(COPD) [46].

Successful replication of genes associated with age.

Comparison of the aging signature we identified in LLFS with a previously published human blood

transcriptomic signature of age by Peters et al. [7] showed significant replication, both in terms of

correlation of the Z-scores (R = 0.78, p < 2.2e-16, Figure 2C) and in terms of signature overlap (Fisher

test p < 2.2e-16, Figure 2D), confirming the robustness of our results (Table S1). The systematic bias in

Z-score values can be attributed to the difference in sample size (n = 14,983 in Peters et al. and n =

2,101 in LLFS).

We performed a similar comparison with the ILO cohort and observed a strong correlation in age-

related gene expression estimates across the genes shared by the two studies (R = 0.82; Figure 2F).

This correlation increased to R = 0.94 when considering only the genes that were significantly

associated with age (q ≤ 0.01) in either LLFS or ILO (Figure 2G). The overlap of significant gene markers

identified in the two studies is also illustrated in Figure 2E.

Functional annotation shows multiple biological processes affected by age.

All analyzed gene markers were ranked by Z-score, with those showing the greatest age-related

increases placed at the top and those with the greatest decreases at the bottom. Enrichment of gene

sets across multiple compendia was then assessed using KS tests applying a significance threshold of

q ≤ 0.05. Figure 3A displays the top enrichment results in the HALLMARK compendium, with the full

results available in Table S2.

The set of genes increasing with age was found to be significantly enriched in inflammation and
cancer-related pathways, such as interferon gamma response (q = 3.87e-32, including PSMB10, IFI30,
and VAMP5), and TNF-α signaling via NF-κB (q = 1.01e-10, including KLF10, PLEK, and CCL4). Given the
reported relationship between inflammation and cell senescence [47], we further tested the
enrichment of the SenMayo gene set [35] in the age signature, and showed a significant association of
the gene set with markers that increase with age (enrichment score = 0.528, p = 4.39e-7, Figure 3B).
The set of genes decreasing with age was enriched in pathways related to cell growth, such as MYC
targets (q = 1.83e-12), E2F targets (q = 0.0038), and Wnt/β-catenin signaling (q = 0.0129).

WGCNA-based module detection [36] of the 2,550 aging markers with absolute Z-score greater than 6
identified 12 modules of highly correlated genes, each containing between 26 and 474 genes, and with
the majority of the genes in each module changing with age in the same direction (Figure 3C). The
modules were further annotated by their hub genes, defined as the most 'central' genes within each
module as measured by their correlation with the module's first eigen value, and by gene set over-
representation enrichment analysis (Figure 3D, E). This dimensionality reduction approach further
highlighted dominant patterns in the aging signature.

For instance, the turquoise module, which is the largest module and primarily consists of genes that
increase with age, was found to be enriched in immune-, inflammation-, and stress-related pathways,
such as inflammatory response, interferon gamma response, and hypoxia (Figure 3D), indicating
stress and activation of the immune system. The hub gene, interferon gamma receptor 2 (IFNGR2), is
representative of the interferon pathway.

Conversely, the blue, brown, and yellow modules, which are primarily composed of genes that
decrease with age, were found to be enriched in MYC targets as well as IL-2 and STAT5 signaling
pathways. The hub genes for these modules were T-cell specific factor 7 (TCF7), Ena/Vasodilator-
stimulated phosphoprotein-like gene (EVL), which is associated with cytoskeleton remodeling and cell
migration, and Ribosomal Protein L10A (RPL10A), which encodes a ribosomal protein [40]. These
findings are consistent with increased inflammation and reduced protein synthesis and cell
proliferation in older individuals (Figure 3D; Table 3). Interestingly, the estrogen response early
pathway was enriched in the blue module, which comprises genes that significantly decrease with
age. In contrast, the estrogen response late pathway was enriched in the red module, which primarily
includes genes that increase with age.

Several blood transcripts predict mortality risk.

We found 122 mRNA transcripts, encoding 122 genes, whose increased expression was associated
with higher mortality risk (HR > 1), and 192 mRNA transcripts, encoding 175 genes, whose increased
expression was associated with lower mortality risk (HR < 1, Figure 1). The full results are provided in
Table S1.

KS-based enrichment analysis identified several gene sets significantly associated with mortality risk.
Gene sets associated with increased mortality risk (HR > 1) included MYC targets (q = 8.67e-20), E2F

targets (q = 2.74e-3), and heme metabolism (q = 1.35e-12). In contrast, gene sets associated with
decreased mortality risk (HR < 1) included natural killer cell-mediated cytotoxicity (q = 2.47e-5) and
signaling by GPCR (q = 1.88e-5). The full list of results is provided in Table S2.

A subset of aging markers is uniquely associated with age in long-lived individuals.

Since both LLFS and ILO cohorts are enriched for individuals who have achieved exceptional longevity
(EL), we compared our findings with previously published aging signatures by Peters et al. [7], whose
study did not include EL individuals, to identify aging markers potentially unique to this distinct
population. Across the three studies, we observed an overlap of 9,358 gene markers. To define
markers specific to the long-lived population, we selected genes whose expression levels were
significantly associated with age (q ≤ 0.01) in both LLFS and ILO but not associated with age (p > 0.25)
in the Peters et al. dataset. Based on these criteria, we identified 176 and 110 gene markers that
uniquely increased and decreased with age, respectively, in LLFS and ILO. Notably, 14 of these genes
were also significantly associated with mortality risk in LLFS (q ≤ 0.05), with 8 of them meeting a more
stringent threshold of q ≤ 0.01 (Figure 4A, Figure S2). Z-score normalized expression levels of these
genes in LLFS, stratified by age groups in ten-year intervals, are visualized as a heatmap in Figure S2,
highlighting age-related expression changes.

Gene set enrichment analysis revealed pathways significantly associated with markers uniquely
linked to age in longevity-enriched families (Figure 4B). Several pathways, such as IL6-Jak-Stat3
signaling (q = 7.8e-4) and the p53 pathway (q = 2.5e-3), overlapped with those enriched among the
broader set of age-associated markers described earlier. However, additional pathways emerged in
this subset, including Mitotic Spindle (q = 0.018), which was not identified in the original aging
signature analysis. Other pathways, such as UV Response (q = 8.3e-3) and Glycolysis (q = 0.018), were
identified in both analyses; notably, their enrichment significance remained relatively stable despite
the smaller number of age-associated genes specific to longevity-enriched families.

Age Acceleration derived from a transcriptomic clock predicts increased mortality risk.

We fitted a transcriptomic clock using an elastic net model on all transcripts that passed quality
control, resulting in a final model that included 1,432 transcripts with non-zero coefficients, with their
distribution visualized in Figure 5A. The top genes ranked by permutation-based variable importance
are shown in Figure 5B.

Upon applying the model to the dataset, a systematic bias was observed in the predicted ages (Figure
5C), which was also observed in previous studies [6], [17]. The bias was corrected using Cole's formula
[17], [39], as shown in Figure 5D. Delta-age was defined as the difference between the adjusted
predicted age and chronological age, and its impact on mortality risk was evaluated both as a
continuous variable and as categorical groups. Cox-PH models were applied with adjustments for
relevant covariates. As a continuous variable, delta-age was significantly positively associated with
mortality risk (HR = 1.108, p = 3.33e-18, Figure 5E). When recoded as a three-value categorical variable,
survival analysis revealed that individuals with a negative delta-age of more than three years had a

significantly lower mortality risk (HR = 0.6403, p = 3.43e-5, Figure 5E), whereas those with a positive
delta-age of more than three years showed a significantly higher mortality risk (HR = 1.561, p = 7.81e-
7, Figure 5E). These trends are also reflected in the age-adjusted Kaplan-Meier curve of Figure 5E.

Discussion

In this study, we performed a large-scale transcriptomics profiling of subjects from families enriched
for long-lived individuals and spanning a wide age range, between 18 and 107 years. The large sample
size (n = 2,167) and the wide age range allowed us to robustly identify signatures associated with
chronological age, mortality risk, and extreme longevity, and to integrate these results toward the
prioritization of candidate unique aging markers.

Our analysis detected 8,271 genes exhibiting significant associations with age (q ≤ 0.01), with roughly
equal numbers increasing and decreasing over the lifespan. These age-related expression changes
highlight major transcriptional remodeling with age, reflecting coordinated shifts in immune function,
inflammation, cellular stress responses, and cell cycle regulation. Notably, genes involved in immune
cell activation and differentiation, such as CD28, CD27, TCF7, and CCR6, exhibited marked declines
with age, consistent with the well-documented phenomenon of immunosenescence. Conversely,
genes such as ZEB2 and LGR6 were significantly upregulated with age, suggesting increased activity in
pathways related to cellular senescence, epithelial-to-mesenchymal transition, and Wnt/β-catenin
signaling, which are hallmarks of aging and chronic disease progression. This finding is consistent
with a previous genome-wide association study in LLFS, which also identified several genes involved
in the Wnt/β-catenin signaling pathway as being associated with age-related traits [48].

Our results strongly corroborate previous findings, particularly the transcriptomic signature of aging
reported by Peters et al. Despite differences in sample size and demographic composition, we
observed strong replication of gene-level associations and pathway enrichments, confirming the
robustness and generalizability of our aging signature. Replication across the ILO cohort further
strengthens our findings and demonstrates that transcriptomic signatures of aging are largely
conserved across different populations and family-based study designs.

Enriched gene sets revealed biological processes known to play a role in aging.

Functional enrichment analyses revealed that genes upregulated with age were consistently enriched
in inflammatory and immune response pathways, including TNF-α signaling and interferon gamma
response. This aligns with the concept of "inflammaging," a state of chronic, low-grade inflammation
that contributes to age-related morbidity [1]. The enrichment of the SenMayo gene set among age-
upregulated transcripts further emphasizes the role of cellular senescence in the aging process [35],
as well as its link to inflammation and its established role as a hallmark of aging [1]. In contrast, genes
downregulated with age were enriched for MYC and E2F targets and Wnt/β-catenin signaling,
suggesting declining proliferative capacity and regenerative potential with age. Prior studies have
shown that MYC also plays a role in nutrient sensing [2], and its downregulation can promote

longevity and health span in mice [49]. Additionally, some of the key biological functions regulated by
MYC, such as ribosome biogenesis and protein translation, are essential for cell proliferation and
survival [50]. These findings highlight the interconnected roles of cell cycle regulation, nutrient
sensing, and MYC signaling in the aging process.

To distill these complex changes into interpretable networks, we employed WGCNA to identify co-
expression modules among highly age-associated genes. These modules revealed distinct
transcriptional programs reflecting core aging processes, including immune activation, translational
suppression, and cytoskeletal remodeling. For example, the turquoise module, enriched for
inflammatory and interferon pathways, was anchored by the hub gene *IFNGR2*. Conversely, modules
composed of downregulated genes were enriched for MYC target genes and IL-2/STAT5 signaling, with
hub genes such as TCF7, EVL, and RPL10A pointing to declining immune signaling and ribosomal
function.

Interestingly, our data also revealed differential enrichment of early and late estrogen response
pathways, with early response genes declining and late response genes increasing with age. Though
both pathways have been implicated in breast cancer biology [51] [52], their distinct age-related
regulation may reflect complex hormonal changes in aging and could have implications for sex-
specific disease risk.

Mortality-associated genes offer insight into underlying biological mechanisms.

Many of the identified transcripts were also predictive of mortality. Gene sets associated with
increased mortality risk included MYC and E2F targets, as well as pathways related to heme
metabolism, whereas protective gene sets included those involved in chemokine signaling, natural
killer (NK) cell-mediated cytotoxicity, and G protein–coupled receptor (GPCR) signaling, the latter
previously shown to influence cellular senescence and cell cycle control [53]. These findings suggest
that a robust immune response may support the detection and clearance of abnormal or senescent
cells, thereby reduce the risk of malignancy and contribute to improved survival. Conversely, our
observation that elevated expression of *MYC* targets is associated with increased mortality aligns with
experimental evidence that heterozygous Myc ± mice exhibit extended lifespan compared to wild type
(Myc +/+) mice [54].

Elevated expression of several ribosomal proteins, including RPL32, RPL19, and RPS8, was
significantly associated with increased mortality risk. These genes contribute to the activation of
several pathways linked to nutrient sensing and cellular biogenesis, including the EIF2AK4 (GCN2)
response to amino acid deficiency, seleno-amino acid metabolism, and cellular responses to
starvation. In addition, they were enriched in the *regulation of SLITs and ROBOs expression* and *ROBO*
receptor signaling, pathways previously implicated in inflammation and tissue remodeling [55].
Interestingly, these same ribosomal proteins were found to decrease in expression with chronological
age. This may reflect the well-documented decline in efficient protein synthesis capacity with age [1].
However, in the context of disease or compensatory responses, dysregulation of these pathways may

contribute to maladaptive cellular states, thereby increasing mortality risk. Further investigation is
warranted to clarify the context-specific roles of these ribosomal genes in aging and survival.

Taken together, these findings support the potential utility of blood transcriptomic markers as
prognostic indicators of biological age and mortality risk. They also underscore the importance of
specific transcriptional programs – particularly those related to immune surveillance, cellular
metabolism, and translational control – in mediating resilience or susceptibility to age-related
decline.

Functional insights into aging markers unique to individuals from families enriched in
exceptional longevity.

A particularly novel aspect of our study was the identification of a subset of aging markers that are
associated with age in the LLFS and ILO cohorts, but not in the general population. These unique
markers were enriched in pathways such as IL6-Jak-Stat3 signaling and p53 signaling, as well as
mitotic spindle and glycolysis pathways not observed in the broader age signature. The emergence of
these unique signatures in long-lived families suggests that distinct regulatory programs may underlie
healthy aging and exceptional longevity.

Some of these markers may play a protective role against age-related decline. For example, regulator
of G protein signaling 3 (RGS3), which increases with age only in longevity-enriched cohorts, was
previously shown to confer protection against pathological changes in fibroblasts [56]. Although not
yet extensively studied, it has demonstrated potential for clinical application. Members of the RGS
family have been proposed as biomarkers in cancer [57], highlighting their growing relevance in
clinical settings. In addition, while no studies have directly examined RGS3 in this context, it
modulates the KRAS pathway [58] which is involved in cancer progression. A clinical trial for non-small
cell lung cancer is currently underway (ClinicalTrials.gov ID: NCT06582771) that targets KRAS and
includes RGS3 as one of its evaluation points. Taken together, a higher expression level of RGS3 seem
to be beneficial as it could suppress oncogenic KRAS activity, which aligns with our hypothesis, and it
has already showcased its potential application in clinical settings. Similarly, CD200, which is known
to be elevated in pancreatic ductal adenocarcinoma [59], exhibited a unique age-related decrease in
the longevity enriched families, suggesting a potentially beneficial regulatory role in aging within this
cohort. CD200R1, the receptor for CD200, has already been identified as a promising therapeutic
target in a phase 1 clinical trial [60], supporting the potential utility of CD200 in clinical applications.

In contrast, however, the functional roles of some other markers are less clear. For instance, NR1D1, a
nuclear receptor with reported anti-inflammatory role through inhibition of M1 macrophage
polarization via suppression of the NF-κB pathway [61], showed a decline with age. This may indicate
a deleterious shift in immune regulation. However, given NR1D1's involvement in circadian rhythm
regulation, light exposure, and physical activity [62], its downregulation may also reflect
environmental or behavioral changes with age, rather than intrinsic biological decline.

Similarly, PIK3R3 (est = 8.961e-3, q = 1.397e-14) and RNF114 (est = 2.871e-3, q = 3.703e-14), both of
which demonstrated age-associated increases in expression, have been implicated in promoting
colorectal cancer cell proliferation [63] [64], with PIK3R3 also upregulated in liver cancer cells [65].
Although these associations might initially suggest harmful effects, it is important to consider the
context of these findings. Prior studies were conducted in cancer cell lines, whereas our observations
are derived from whole blood transcriptomic data in non-diseased individuals. While these genes are
linked to proliferative signaling in cancer, their expression in healthy aging may reflect preserved
proliferative or regenerative capacity. Given that aging is typically associated with diminished cell
proliferation and tissue maintenance, it is possible that increased expression of such genes may,
under non-pathological conditions, support tissue repair and homeostasis, underscoring their
potentially beneficial context-dependent roles.

Pathway analysis further reveals unique patterns in those families enriched with exceptional
longevity. Several immune and inflammation-related pathways, including IL6-Jak-Stat3 signaling, the
p53 pathway, and TNF-α signaling via NF-κB, were enriched both in this subset and in the broader
aging marker analysis (Figure 3A). However, additional pathways emerged as unique to the long-lived
population. Notably, the UV response up pathway, associated with external stress, was enriched
among genes that uniquely increased with age. Additionally, enrichment of glycolysis and the mitotic
spindle pathways, which are involved in energy metabolism and cell division, suggests heightened
cellular activity and proliferative potential in individuals from families with exceptional longevity.

Delta-age predicted using a transcriptomics aging clock is associated with mortality risk.

Our findings demonstrate that delta-age, defined as the difference between the biological age
predicted by the clock and the chronological age, is a strong predictor of all-cause mortality. Positive
delta-age values were associated with increased mortality risk, while negative values were predictive
of reduced risk. These associations remained robust whether we analyzed delta-age as a continuous
or categorical variable. These results align with previous reports of biological clocks capturing aging-
related functional decline [6], [17], and they further support the utility of transcriptomic clocks as
meaningful biomarkers of aging. The effect sizes observed here (HR = 1.108 per unit delta-age; HR =
1.561 for those >3 years older) underscore the relevance of transcriptome-derived age acceleration
not only as a reflection of biological aging but also as a clinically informative metric for mortality risk
stratification, assessment of individual health status, and monitoring intervention effects.

Limitations and Future Directions

Blood cell type composition changes.

The composition of blood cell types changes with age, thereby influencing gene expression levels in
whole blood [66]. For instance, a single-cell study in centenarians observed a decrease in naive CD4 T-
cells, memory CD4 T-cells, and naive B-cells, alongside an increase in CD8 T-cells, natural killer cells,
and monocytes proportions among the extreme old individuals compared to younger age groups [9].

In this study, we did not incorporate blood cell type information, as our primary objective was to
identify easily accessible aging markers in blood, the rationale being that even if a marker's
significance is induced by changes in blood cell type composition, it remains a valid aging marker.
Current work is ongoing to further investigate the association between cell type composition with age.

Analysis of longitudinal changes.

All findings discussed in this study are the results of cross-sectional analyses. A previous longitudinal
multi-omics study identified diverse aging patterns among individuals, highlighting that longitudinal
changes within individuals can diverge from the cross-sectional trends observed in molecular features
[67]. Currently, LLFS is actively collecting and profiling blood transcriptome data at various time
points. At the time this manuscript is being prepared, LLFS has collected transcriptomic data from
over 1,100 participants across at least two time points, with time intervals ranging from 5 to 17 years.
Among them, more than 300 participants have measurements at three time points, providing a
valuable dataset for evaluating individual longitudinal changes. As a subsequent step, we will explore
longitudinal changes in transcriptomic profiles across participants of different ages, which will offer
deeper insight into the dynamics of molecular features over time.

Integration of multi-omics, single-cell, and tissue-specific analysis.

This study focused exclusively on whole blood transcriptomic analysis. However, aging and mortality
risk are complex processes that involve changes across multiple molecular and tissue-specific levels.
Future research that integrates multi-omics datasets may yield a more comprehensive understanding
of these processes. Moreover, while whole blood offers advantages such as minimally invasive sample
collection and relevance in systemic biological studies, it is a heterogeneous tissue and lacks cellular
specificity. Analyzing specific tissue types, and further diving into the single-cell space could provide a
more detailed and systematic view of the aging process.

Grant Funding

This research was supported by NIH-NIA U19 AG063893 and UH2/UH3AG064704 (ILO).

Competing Interest Declaration

539   The authors declare no competing interests.

Ethics Declaration

This study was conducted in accordance with the relevant local and international ethical guidelines. All enrolled participants (or their legal guardian or representative, when applicable) provided informed consent. All study personnel and data analysts have completed and maintained current human-subjects research training under IRB oversight. All participant data used, including phenotypical data, genetic data, and RNA-seq data, were de-identified and are stored on secure servers with restricted access.

Author Contribution Declaration

Analysis, methods and manuscript writing: Mengze Li.

Analysis, methods and manuscript editing: Zeyuan Song, Eric Reed, Tanya T. Karagiannis, Stacy Andersen.

RNA-seq data acquiring and processing: Michael Brent, Chase Mateusiak, Sandeep Acharya, Wooseok J. Jung, Jeffrey R. O'Connell, Sofiya Milman.

Participant recruiting, administration, phenotype data collection and processing: Shu Liao, Mary K. Wojczynski, Mary F. Feitosa, Konstantin Arbeev, May E. Montasser, Albert Tai, Roland J. Thorpe.

Analysis guidance, ideation, manuscript editing, and funding acquiring: Stefano Monti, Paola Sebastiani, Thomas T. Perls.

References

[1] C. López-Otín, M. A. Blasco, L. Partridge, M. Serrano, and G. Kroemer, "Hallmarks of aging: An expanding universe," *Cell*, vol. 186, no. 2, pp. 243–278, Jan. 2023, doi: 10.1016/j.cell.2022.11.001.

[2] P. P. Singh, B. A. Demmitt, R. D. Nath, and A. Brunet, "The Genetics of Aging: A Vertebrate Perspective," *Cell*, vol. 177, no. 1, pp. 200–220, Mar. 2019, doi: 10.1016/j.cell.2019.02.038.

[3] J. Rutledge, H. Oh, and T. Wyss-Coray, "Measuring biological age using omics data," *Nat Rev Genet*, vol. 23, no. 12, pp. 715–727, Dec. 2022, doi: 10.1038/s41576-022-00511-7.

[4] B. K. Kennedy *et al.*, "Geroscience: Linking Aging to Chronic Disease," *Cell*, vol. 159, no. 4, pp. 709–713, Nov. 2014, doi: 10.1016/j.cell.2014.10.039.

[5] M. A. Espeland *et al.*, "Clinical Trials Targeting Aging and Age-Related Multimorbidity," *GERONA*, p.
glw220, Oct. 2016, doi: 10.1093/gerona/glw220.

[6] T. Tanaka *et al.*, "Plasma proteomic signature of age in healthy humans," *Aging Cell*, vol. 17, no. 5, p.
e12799, Oct. 2018, doi: 10.1111/acel.12799.

[7] NABEC/UKBEC Consortium *et al.*, "The transcriptional landscape of age in human peripheral blood,"
Nat Commun, vol. 6, no. 1, p. 8570, Dec. 2015, doi: 10.1038/ncomms9570.

[8] P. Sebastiani *et al.*, "Metabolite signatures of chronological age, aging, survival, and longevity," *Cell*
Reports, vol. 43, no. 11, p. 114913, Nov. 2024, doi: 10.1016/j.celrep.2024.114913.

[9] T. T. Karagiannis *et al.*, "Multi-modal profiling of peripheral blood cells across the human lifespan
reveals distinct immune cell signatures of aging and longevity," *eBioMedicine*, vol. 90, p. 104514, Apr.
2023, doi: 10.1016/j.ebiom.2023.104514.

[10] P. Sebastiani and T. T. Perls, "The Genetics of Extreme Longevity: Lessons from the New England
Centenarian Study," *Front. Gene.*, vol. 3, 2012, doi: 10.3389/fgene.2012.00277.

[11] P. Sebastiani *et al.*, "Families Enriched for Exceptional Longevity also have Increased Health-Span:
Findings from the Long Life Family Study," *Front. Public Health*, vol. 1, 2013, doi:
10.3389/fpubh.2013.00038.

[12] S. L. Andersen, P. Sebastiani, D. A. Dworkis, L. Feldman, and T. T. Perls, "Health Span Approximates
Life Span Among Many Supercentenarians: Compression of Morbidity at the Approximate Limit of Life
Span," *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, vol. 67A, no. 4,
pp. 395–405, Apr. 2012, doi: 10.1093/gerona/glr223.

[13] K. Ismail *et al.*, "Compression of Morbidity Is Observed Across Cohorts with Exceptional Longevity," *J*
American Geriatrics Society, vol. 64, no. 8, pp. 1583–1591, Aug. 2016, doi: 10.1111/jgs.14222.

[14] P. Sebastiani *et al.*, "Protein signatures of centenarians and their offspring suggest centenarians age
slower than other humans," *Aging Cell*, vol. 20, no. 2, Feb. 2021, doi: 10.1111/acel.13290.

[15] P. Garagnani *et al.*, "Whole-genome sequencing analysis of semi-supercentenarians," *eLife*, vol. 10, p.
e57849, May 2021, doi: 10.7554/eLife.57849.

[16] M. Schmidt *et al.*, "The Human Blood Transcriptome in a Large Population Cohort and Its Relation to
Aging and Health," *Front. Big Data*, vol. 3, p. 548873, Oct. 2020, doi: 10.3389/fdata.2020.548873.

[17] A.-M. G. De Lange and J. H. Cole, "Commentary: Correction procedures in brain-age prediction,"
NeuroImage: Clinical, vol. 26, p. 102229, 2020, doi: 10.1016/j.nicl.2020.102229.

[18] M. K. Wojczynski *et al.*, "NIA Long Life Family Study: Objectives, Design, and Heritability of Cross-
Sectional and Longitudinal Phenotypes," *The Journals of Gerontology: Series A*, vol. 77, no. 4, pp. 717–
727, Apr. 2022, doi: 10.1093/gerona/glab333.

[19] S. Acharya *et al.*, "A methodology for gene level omics-WAS integration identifies genes influencing
traits associated with cardiovascular risks: the Long Life Family Study," *Hum. Genet.*, vol. 143, no. 9–
10, pp. 1241–1252, Oct. 2024, doi: 10.1007/s00439-024-02701-1.

[20] Harshil Patel *et al.*, *nf-core/rnaseq: nf-core/rnaseq v3.18.0 - Lithium Lynx*. (Dec. 20, 2024). Zenodo. doi:
10.5281/ZENODO.14537300.

[21] F. Krueger, F. James, P. Ewels, E. Afyounian, and B. Schuster-Boeckler, *FelixKrueger/TrimGalore:*
v0.6.7 - DOI via Zenodo. (July 23, 2021). Zenodo. doi: 10.5281/ZENODO.5127899.

[22] M. Martin, "Cutadapt removes adapter sequences from high-throughput sequencing reads," *EMBnet*
j., vol. 17, no. 1, p. 10, May 2011, doi: 10.14806/ej.17.1.200.

[23] A. Dobin *et al.*, "STAR: ultrafast universal RNA-seq aligner," *Bioinformatics*, vol. 29, no. 1, pp. 15–21,
Jan. 2013, doi: 10.1093/bioinformatics/bts635.

[24] R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, and C. Kingsford, "Salmon provides fast and bias-aware
quantification of transcript expression," *Nat Methods*, vol. 14, no. 4, pp. 417–419, Apr. 2017, doi:
10.1038/nmeth.4197.

[25] K. Okonechnikov, A. Conesa, and F. García-Alcalde, "Qualimap 2: advanced multi-sample quality
control for high-throughput sequencing data," *Bioinformatics*, vol. 32, no. 2, pp. 292–294, Jan. 2016,
doi: 10.1093/bioinformatics/btv566.

[26] M. I. Love, W. Huber, and S. Anders, "Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2," *Genome Biol*, vol. 15, no. 12, p. 550, Dec. 2014, doi: 10.1186/s13059-014-0550-8.

[27] Z. Song, A. Gurinovich, A. Federico, S. Monti, and P. Sebastiani, "nf-gwas-pipeline: A Nextflow
Genome-Wide Association Study Pipeline," *JOSS*, vol. 6, no. 59, p. 2957, Mar. 2021, doi:
10.21105/joss.02957.

[28] E. W. Daw *et al.*, "A Paradigm For Calling Sequence In Families: The Long Life Family Study," May 24,
2024. doi: 10.1101/2024.05.23.595584.

[29] S. M. Gogarten *et al.*, "Genetic association testing using the GENESIS R/Bioconductor package,"
Bioinformatics, vol. 35, no. 24, pp. 5346–5348, Dec. 2019, doi: 10.1093/bioinformatics/btz567.

[30] T. M. Therneau, "survival: Survival Analysis." p. 3.8-3, June 22, 2001. doi:
10.32614/CRAN.package.survival.

[31] S. Durinck *et al.*, "BioMart and Bioconductor: a powerful link between biological databases and
microarray data analysis," *Bioinformatics*, vol. 21, no. 16, pp. 3439–3440, Aug. 2005, doi:
10.1093/bioinformatics/bti525.

[32]   A. Liberzon, A. Subramanian, R. Pinchback, H. Thorvaldsdóttir, P. Tamayo, and J. P. Mesirov,
634   "Molecular signatures database (MSigDB) 3.0," *Bioinformatics*, vol. 27, no. 12, pp. 1739–1740, June
635   2011, doi: 10.1093/bioinformatics/btr260.

636   [33]   A. Federico and S. Monti, "hypeR: an R package for geneset enrichment workflows," *Bioinformatics*,
637   vol. 36, no. 4, pp. 1307–1308, Feb. 2020, doi: 10.1093/bioinformatics/btz700.

638   [34]   C. Alexey Sergushichev [Aut, *fgsea*. (2017). Bioconductor. doi: 10.18129/B9.BIOC.FGSEA.

639   [35]   D. Saul *et al.*, "A new gene set identifies senescent cells and predicts senescence-associated
640   pathways across tissues," *Nat Commun*, vol. 13, no. 1, p. 4827, Aug. 2022, doi: 10.1038/s41467-022-
641   32552-1.

642   [36]   P. Langfelder and S. Horvath, "WGCNA: an R package for weighted correlation network analysis," *BMC
643   Bioinformatics*, vol. 9, no. 1, p. 559, Dec. 2008, doi: 10.1186/1471-2105-9-559.

644   [37]   J. Friedman, T. Hastie, and R. Tibshirani, "Regularization Paths for Generalized Linear Models via
645   Coordinate Descent," *J Stat Softw*, vol. 33, no. 1, pp. 1–22, 2010.

646   [38]   B. Greenwell M. and B. Boehmke C., "Variable Importance Plots—An Introduction to the vip Package,"
647   *The R Journal*, vol. 12, no. 1, p. 343, 2020, doi: 10.32614/RJ-2020-013.

648   [39]   J. H. Cole *et al.*, "Brain age predicts mortality," *Mol Psychiatry*, vol. 23, no. 5, pp. 1385–1392, May 2018,
649   doi: 10.1038/mp.2017.62.

650   [40]   G. Stelzer *et al.*, "The GeneCards Suite: From Gene Data Mining to Disease Genome Sequence
651   Analyses," *CP in Bioinformatics*, vol. 54, no. 1, June 2016, doi: 10.1002/cpbi.5.

652   [41]   C. Wang, T. Ling, N. Zhong, and L.-G. Xu, "N4BP3 Regulates RIG-I-Like Receptor Antiviral Signaling
653   Positively by Targeting Mitochondrial Antiviral Signaling Protein," *Front. Microbiol.*, vol. 12, p. 770600,
654   Nov. 2021, doi: 10.3389/fmicb.2021.770600.

655   [42]   H. T. Meitei, N. Jadhav, and G. Lal, "CCR6-CCL20 axis as a therapeutic target for autoimmune
656   diseases," *Autoimmunity Reviews*, vol. 20, no. 7, p. 102846, July 2021, doi:
657   10.1016/j.autrev.2021.102846.

658   [43]   C. Vandewalle, "SIP1/ZEB2 induces EMT by repressing genes of different epithelial cell-cell junctions,"
659   *Nucleic Acids Research*, vol. 33, no. 20, pp. 6566–6578, Nov. 2005, doi: 10.1093/nar/gki965.

660   [44]   D. Dai *et al.*, "The transcription factor ZEB2 drives the formation of age-associated B cells," *Science*,
661   vol. 383, no. 6681, pp. 413–421, Jan. 2024, doi: 10.1126/science.adf8531.

662   [45]   B. Bernardes De Jesus *et al.*, "Silencing of the lncRNA Zeb2-NAT facilitates reprogramming of aged
663   fibroblasts and safeguards stem cell pluripotency," *Nat Commun*, vol. 9, no. 1, p. 94, Jan. 2018, doi:
664   10.1038/s41467-017-01921-6.

[46] E. E. Cortesi *et al.*, "Increased LGR6 Expression Sustains Long-Term Wnt Activation and Acquisition of
Senescence in Epithelial Progenitors in Chronic Lung Diseases," *Cells*, vol. 10, no. 12, p. 3437, Dec.
2021, doi: 10.3390/cells10123437.

[47] J.-L. Ren, J.-S. Pan, Y.-P. Lu, P. Sun, and J. Han, "Inflammatory signaling and cellular senescence,"
Cellular Signalling, vol. 21, no. 3, pp. 378–383, Mar. 2009, doi: 10.1016/j.cellsig.2008.10.011.

[48] M. F. Feitosa *et al.*, "Genetic Pleiotropy Between Pulmonary Function and Age-Related Traits: The
Long Life Family Study," *The Journals of Gerontology: Series A*, vol. 79, no. 3, Mar. 2024, doi:
10.1093/gerona/glac046.

[49] J. W. Hofmann *et al.*, "Reduced Expression of MYC Increases Longevity and Enhances Healthspan,"
Cell, vol. 160, no. 3, pp. 477–488, Jan. 2015, doi: 10.1016/j.cell.2014.12.016.

[50] C. V. Dang, K. A. O'Donnell, K. I. Zeller, T. Nguyen, R. C. Osthus, and F. Li, "The c-Myc target gene
network," *Seminars in Cancer Biology*, vol. 16, no. 4, pp. 253–264, Aug. 2006, doi:
10.1016/j.semcancer.2006.07.014.

[51] M. Oshi *et al.*, "Degree of Early Estrogen Response Predict Survival after Endocrine Therapy in Primary
and Metastatic ER-Positive Breast Cancer," *Cancers*, vol. 12, no. 12, p. 3557, Nov. 2020, doi:
10.3390/cancers12123557.

[52] T. Takeshita *et al.*, "Clinical Relevance of Estrogen Reactivity in the Breast Cancer Microenvironment,"
Front. Oncol., vol. 12, p. 865024, May 2022, doi: 10.3389/fonc.2022.865024.

[53] P. Santos-Otte, H. Leysen, J. Van Gastel, J. O. Hendrickx, B. Martin, and S. Maudsley, "G Protein-
Coupled Receptor Systems and Their Role in Cellular Senescence," *Computational and Structural
Biotechnology Journal*, vol. 17, pp. 1265–1277, 2019, doi: 10.1016/j.csbj.2019.08.005.

[54] E. V. Prochownik and H. Wang, "Lessons in aging from Myc knockout mouse models," *Front. Cell Dev.
Biol.*, vol. 11, p. 1244321, Aug. 2023, doi: 10.3389/fcell.2023.1244321.

[55] M. Tong, T. Jun, Y. Nie, J. Hao, and D. Fan, "The Role of the Slit/Robo Signaling Pathway," *J. Cancer*,
vol. 10, no. 12, pp. 2694–2705, 2019, doi: 10.7150/jca.31877.

[56] F. Xu *et al.*, "RGS3 inhibits TGF-β1/Smad signalling in adventitial fibroblasts," *Cell Biochemistry &
Function*, vol. 35, no. 6, pp. 334–338, Aug. 2017, doi: 10.1002/cbf.3280.

[57] C. Yang *et al.*, "Function and regulation of RGS family members in solid tumours: a comprehensive
review," *Cell Commun Signal*, vol. 21, no. 1, p. 316, Nov. 2023, doi: 10.1186/s12964-023-01334-7.

[58] C. Li, A. Vides, D. Kim, J. Y. Xue, Y. Zhao, and P. Lito, "The G protein signaling regulator RGS3 enhances
the GTPase activity of KRAS," *Science*, vol. 374, no. 6564, pp. 197–201, Oct. 2021, doi:
10.1126/science.abf1730.

[59]   F. Choueiry *et al.*, "CD200 promotes immunosuppression in the pancreatic tumor
698   microenvironment," *J Immunother Cancer*, vol. 8, no. 1, p. e000189, June 2020, doi: 10.1136/jitc-2019-
699   000189.

700   [60]   S. Kummar *et al.*, "First-in-Human Study of 23ME-00610, an Antagonistic Antibody for Genetically
701   Validated CD200R1 Immune Checkpoint, in Participants with Advanced Solid Malignancies," *Cancer
702   Research Communications*, vol. 5, no. 1, pp. 94–105, Jan. 2025, doi: 10.1158/2767-9764.CRC-24-0568.

703   [61]   Z.-Y. Zhang-sun *et al.*, "Targeting NR1D1 in organ injury: challenges and prospects," *Military Med Res*,
704   vol. 10, no. 1, p. 62, Dec. 2023, doi: 10.1186/s40779-023-00495-3.

705   [62]   D. Gubin *et al.*, "Light Exposure, Physical Activity, and Indigeneity Modulate Seasonal Variation in
706   NR1D1 (REV-ERBα) Expression," *Biology*, vol. 14, no. 3, p. 231, Feb. 2025, doi:
707   10.3390/biology14030231.

708   [63]   Q. Chen, X. Sun, X. Luo, J. Wang, J. Hu, and Y. Feng, "PIK3R3 inhibits cell senescence through p53/p21
709   signaling," *Cell Death Dis*, vol. 11, no. 9, p. 798, Sept. 2020, doi: 10.1038/s41419-020-02921-z.

710   [64]   S. Liu *et al.*, "RNF114 facilitates the proliferation, stemness, and metastasis of colorectal cancer,"
711   *Pathology - Research and Practice*, vol. 248, p. 154716, Aug. 2023, doi: 10.1016/j.prp.2023.154716.

712   [65]   W. Lin *et al.*, "PIK3R3 is upregulated in liver cancer and activates Akt signaling to control cancer
713   growth by regulation of CDKN1C and SMC1A," *Cancer Medicine*, vol. 12, no. 13, pp. 14413–14425, July
714   2023, doi: 10.1002/cam4.6068.

715   [66]   BIOS Consortium *et al.*, "Correction for both common and rare cell types in blood is important to
716   identify genes that correlate with age," *BMC Genomics*, vol. 22, no. 1, p. 184, Dec. 2021, doi:
717   10.1186/s12864-020-07344-w.

718   [67]   S. Ahadi *et al.*, "Personal aging markers and ageotypes revealed by deep longitudinal profiling," *Nat
719   Med*, vol. 26, no. 1, pp. 83–90, Jan. 2020, doi: 10.1038/s41591-019-0719-5.

Figures

Figure 1 Results summary

Figure 1. Summary of sample size and identified gene markers.

Figure 2 Identified age gene markers and replication

Figure 2. Identified age gene markers and replication.

A. Scatter plots of the expression levels of selected top aging markers. Blue line represents the fitted regression line. Z-scores and q-values (FDR correction) are shown in the caption.

B. Volcano plot of the analyzed gene markers. A few top markers are labeled.

C. Scatter plot of Z-scores from regression analysis of age for 9,463 shared genes across LLFS and Peters.et.al. X-axis: Z-scores from LLFS. Y-axis: Z-scores from Peters et al. Blue line represents the fitted regression line, with R denoting the Pearson correlation coefficient.

D. Venn diagrams of the number of significant gene markers that increase and decrease with age in LLFS and Peters et. al. Significant markers are defined by q ≤ 0.01 (FDR correction). Fisher's exact test was conducted separately for markers increasing (up) and decreasing (down) with age, with each background set to be 9,463, which is the shared number of the genes analyzed in both studies.

E. Venn diagram of the number of significant gene markers that increase and decrease with age in LLFS and ILO. Significant markers are defined by q ≤ 0.01 (FDR correction).

F. Scatter plot of estimates of the effects of all the genes analyzed in both LLFS and ILO, totaling 14,728 genes. X-axis: estimates from LLFS. Y-axis: estimates from ILO. Blue line represents the fitted regression line, with R denoting the Pearson correlation coefficient.

G. Scatter plot of estimates of the effects of the genes that are significantly associated with age (q ≤ 0.01) in either LLFS or ILO, totaling 4,924 genes. X-axis: estimates from LLFS. Y-axis: estimates from ILO. Blue line represents the fitted regression line, with R denoting the Pearson correlation coefficient.

Figure 3 Functional annotation of gene markers and co-expression module detection of age associated genes

Figure 3. Functional annotation and co-expression module detection of age associated genes

A. Gene sets from the HALLMARK compendium that significantly enriched (q ≤ 0.05) in gene markers increasing (Age_up) and decreasing (Age_dn) with age using Kolmogorov-Smirnov (KS) test. Gene markers are ranked by absolute Z score.

B. Enrichment analysis of the SenMayo gene set using Kolmogorov-Smirnov (KS) test, demonstrating cellular senescence activity is enriched in markers increasing with age. Gene markers are ranked left to right based on their high-to-low age Z-scores.

C to E represents the co-expression modules detected among the top aging markers (absolute Z-score > 6, q ≤ 0.01 using FDR correction) using WGCNA.

C. Module sizes, with color representing the number of markers increasing and decreasing with age within the module.

D. Gene sets from the HALLMARK compendium that are significantly enriched (q ≤ 0.05) in each module. Modules with no enriched gene sets are not included.

E. Volcano plot of genes with correlation above 0.75 with the eigenvalue of their respective modules, with color denoting the associated module. Hub genes of each module are labeled.

Figure 4 Unique markers associated with age in longevity enriched families (LLFS and ILO)

Figure 4. Unique markers associated with age in longevity enriched families (LLFS and ILO)

A. Venn diagram illustrating the number of gene markers that are exclusively significantly associated with age in the LLFS and ILO cohorts (q ≤ 0.01), and with mortality risk (q ≤ 0.05), but not associated with age in the Peters et al. study (p > 0.25).

B. Enriched gene sets among the gene markers that are exclusively associated with age in the LLFS and ILO cohorts.

Figure 5 Construction of the transcriptomics aging clock and the association between age deviation (delta-age) and mortality risk.

Figure 5. Construction of the transcriptomics aging clock and the association between age deviation (delta-age) and mortality risk.

A. Histogram showing the distribution of feature coefficients. Only the 1,432 transcripts with non-zero coefficients are included. The darker blue line illustrates the density.

B. Top important features in the aging clock, with importance scores derived by the permutation method. Red and green dots denote features positively (est > 0) and negatively (est < 0) associated with age (q ≤ 0.01) in the aging signature analysis, respectively.

C. Comparison of predicted age versus chronological age, which shows a systematic bias in the predicted age before correction.

D. Corrected predicted age (using the Cole method) versus chronological age, demonstrating the removal of the bias.

E. Kaplan-Meier curve of mortality risk based on delta-age (defined by corrected predicted age minus chronological age) groups, with coefficients and significance indicated below. Normal: individuals whose corrected predicted age differs from their chronological age by no more than 3 years. Decel: individuals predicted to be over 3 years younger than their chronological age. Accel: individuals predicted to be over 3 years older than their chronological age. Age Accel Years: age acceleration as a continuous variable. CA: Chronological age. Female: Effect of female compared to male.

Figure S1. Medication has minimal changes in association between gene expression level and age.

A

B

green line: p = 0.05

C

blue line: x=y

Supplement Figure 1. Medication has minimal changes in association between gene expression level and age.

A. Bar plots showing the number of participants taking each medication in each age groups.

B. Volcano plots showing effect size estimates versus p-values for gene expression associations with four medications, analyzed using the same model as the age association test. Genes significantly associated with medication (FDR q ≤ 0.01) are labeled. The green horizontal dashed line indicates p = 0.05.

C. Scatter plots comparing the estimates and the negative log 10 nominal p-values of the analyses with and without medication added to the model. The systematic decrease in p-value significance when including medications is explained by the reduced sample size since only a fraction of the participants have medication information available. The blue dash line represents the identity line x=y.

Figure S2 Heatmap of the expression level of the markers exclusively in LLFS and ILO.

Age Group

*: mortality q <= 0.05
**:mortality q <= 0.01

Supplement Figure 2. Heatmap of the expression level of the markers exclusively in LLFS and ILO. Heatmap showing the normalized expression levels of 286 gene markers exclusively associated with age in LLFS and ILO cohorts, across different age groups. Genes significantly associated with mortality risk (q ≤ 0.05 and q ≤ 0.01) are highlighted. Genes are clustered using hierarchical clustering based on their expression patterns.

Tables

Table 1. Characteristics of the study participants.

Aging Analysis (LLFS)				
	N	*Age Min*	*Age Median*	*Age Max*
Male	943	18	64	101
Female	1,158	19	63	107
Field Center*	*Total N*	*BU*	*DK*	*NY*	*PT*
	2,101	487	935	152	527

Aging Analysis (ILO)				
	N	*Age Min*	*Age Median*	*Age Max*
Male	135	62	88	107
Female	284	60	87	108

Survival Analysis (LLFS)		
	N	*Alive*	*Deceased*
Male	919	577	342
Female	1,138	782	356
Age	*Min*	*Median*	*Max*
Male	18	64	101
Female	19	63	107
Age Follow up	*Min*	*Median*	*Max*
Male	0	12	18
Female	0	12	18

*Field center: BU: Boston, USA. DK: Denmark. NY: New York, USA. PT: Pittsburgh, USA.

Table 2. Top 20 age associated genes, ranked by p-value.

Ensembl_ID	symbol	Age_Est	Age_SE	Age_Z	Age_p	Age_q
ENSG00000174807.4	CD248	-0.0476	0.00165	-28.9	2.95E-183	4.81E-179
ENSG00000173114.13	LRRN3	-0.0393	0.0016	-24.5	5.92E-133	4.82E-129
ENSG00000135318.12	NT5E	-0.027	0.00128	-21.1	6.53E-99	3.55E-95
ENSG00000145911.6	N4BP3	-0.0219	0.00105	-20.8	2.57E-96	1.05E-92
ENSG00000183691.6	NOG	-0.0347	0.00178	-19.5	9.01E-85	2.93E-81
ENSG00000112486.17	CCR6	-0.0202	0.00107	-18.9	7.48E-80	2.03E-76
ENSG00000081059.20	TCF7	-0.0146	0.000783	-18.7	7.83E-78	1.82E-74
ENSG00000099204.20	ABLIM1	-0.0142	0.000791	-18	1.71E-72	3.47E-69
ENSG00000184613.11	NELL2	-0.0207	0.00116	-17.8	7.48E-71	1.35E-67
ENSG00000167106.12	EEIG1	-0.015	0.000843	-17.7	2.03E-70	3.31E-67
ENSG00000144290.17	SLC4A10	-0.0397	0.00225	-17.6	1.21E-69	1.78E-66
ENSG00000232021.7	LEF1-AS1	-0.0261	0.00148	-17.6	1.66E-69	2.25E-66
ENSG00000169554.22	ZEB2	0.0087	0.000499	17.4	5.68E-68	7.12E-65
ENSG00000139193.4	CD27	-0.0146	0.000853	-17.1	1.73E-65	2.01E-62
ENSG00000136997.21	MYC	-0.012	0.000704	-17.1	3.02E-65	3.28E-62
ENSG00000215788.11	TNFRSF25	-0.0139	0.000821	-17	1.70E-64	1.73E-61
ENSG00000228956.9	SATB1-AS1	-0.0157	0.000929	-16.9	4.96E-64	4.75E-61
ENSG00000089012.14	SIRPG	-0.0143	0.00085	-16.9	8.61E-64	7.79E-61
ENSG00000178562.18	CD28	-0.0141	0.000842	-16.8	3.19E-63	2.74E-60
ENSG00000126353.3	CCR7	-0.019	0.00114	-16.6	5.56E-62	4.52E-59

Table 3. Table of modules detected from 2,550 significant transcriptomic markers (absolute Z-score > 6, q ≤ 0.01) using WGCNA, annotated by the top enriched gene sets from each compendium (HALLMARK, KEGG, and REACTOME). Only gene sets enriched with significance q < 0.05 are included. Size: the total number of markers within the module. Age_up and Age_dn: number of markers increase and decrease with age.

module	size	age_up	age_dn	hub_gene	annot_HALLMARK_label	annot_KEGG_label	annot_REACTOME_label
turquoise	474	469	5	IFNGR2	Inflammatory Response	Leishmania Infection	Neutrophil Degranulation
blue	395	0	395	TCF7	Estrogen Response Early	NA	Repression Of Wnt Target Genes
brown	383	1	382	EVL	Myc Targets V2	NA	Generation Of Second Messenger Molecules
yellow	289	1	288	RPL10A	Myc Targets V1	Ribosome	Eukaryotic Translation Elongation
green	263	256	7	PTPRE	NA	Lysosome	Neutrophil Degranulation
red	256	252	4	TBX21	Allograft Rejection	Natural Killer Cell Mediated Cytotoxicity	NA
black	192	0	192	TNFRSF13C	NA	Primary Immunodeficiency	Antigen Activates B Cell Receptor Bcr Leading To Generation Of Second Messengers
pink	83	83	0	CD68	Apical Junction	Lysosome	Neutrophil Degranulation
magenta	77	0	77	PLCG1	NA	NA	NA
purple	66	66	0	STAT1	Interferon Gamma Response	Proteasome	Interferon Signaling
greenyellow	46	0	46	DCAF1	Myc Targets V2	NA	NA
tan	26	26	0	SAT1	Complement	NA	Innate Immune System

Supplement Table 1. Comprehensive table of all the mRNA associations with age in LLFS and ILO, and mortality risk in LLFS. age_sig are defined by if the gene marker has a q value (FDR corrected) ≤ 0.01. age_dir refers to if the expression of the gene marker increase (Inc) or decrease (Dec) with age.

Supplement Table 2. Significantly enriched gene sets and pathways (q ≤ 0.05) identified from gene markers associated with age (Age_up and Age_dn) and mortality risk (HR > 1 and HR < 1). Enrichment was assessed using a Kolmogorov–Smirnov (KS) test with genes ranked by Z score. label: gene set or pathway names. gene set: total number of genes in this gene set. overlap: the number of genes overlapped between the gene marker list and the gene set. pval and fdr: p-value and q-value (FDR correction) of the enrichment significance. score: enrichment score calculated from fgsea package. hits: the overlapped gene symbols between the gene set and the analyzed data set.