Proteasome Inhibition Reduces NBR1 Protein Levels in mATG8-Deficient Cells Independently of Lysosomal Degradation and RAB27A

Cristóbal Cerda-Troncoso (1)⚓, Eloísa Arias-Muñoz (1), Gabriela Vargas (1), Belén Gaete-Ramírez (1,2), Karina Cereceda (1), Viviana A. Cavieres (1)§ and Patricia V. Burgos (1,2)§

1. Centro de Biología Celular y Biomedicina CEBICEM, Facultad de Ciencias, Universidad San Sebastián, Avenida del Valle Norte 725, Santiago 8580704, Chile.
2. Centro Científico y Tecnológico de Excelencia Ciencia & Vida, Facultad de Medicina y Ciencia, Universidad San Sebastián, Avenida del Valle Norte 725, Santiago 8580704, Chile.

⚓ **Current address**
Laboratory for extracellular vesicle research, Department of Human Genetics, KU Leuven. O&N6, Herestraat 49 box 610, 3000 Leuven, Belgium.

§ **Corresponding authors**
Cavieres VA, email: viviana.cavieres@uss.cl
Burgos PV, email: patricia.burgos@uss.cl

48 **Graphical abstract**

Graphical Abstract
Cerda-Troncoso C, et al., 2025

49
50 *Created with BioRender https://BioRender.com/geoh3sz*
51
52 *In the absence of mATG8s, NBR1 may be recruited to multivesicular bodies (MVBs) via*
53 *HRS, for later degradation in lysosomes (left panel). Under proteasomal inhibition (right*
54 *panel), NBR1 associated with MVBs (with ubiquitinated cargos) is downregulated,*
55 *independent of lysosomes. Possibly, this downregulation is due to a secretion process,*
56 *through a RAB27A-independent mechanism.*
57

Abstract

Autophagy is a lysosome-dependent degradation process that involves autophagosome
formation, typically mediated by mammalian ATG8 proteins (mATG8s).
Autophagosomes can still form in their absence, suggesting alternative mechanisms.
NBR1, a selective autophagy receptor, has been shown to compensate for autophagy
defects. Here, we examined NBR1 regulation under proteotoxic stress in mATG8s-
deficient HeLa cells. NBR1 levels were elevated in mATG8s knockout (KO) cells under
basal conditions but decreased significantly after treatment with the proteasome
inhibitor MG132. This reduction was not prevented by lysosomal inhibition with BafA1,
indicating a non-lysosomal mechanism. Silencing of RAB27A reduced basal NBR1 levels,
and the effects of MG132 were no longer observed, likely due to already diminished
NBR1. Remaining NBR1 localized to puncta positive for ubiquitin and the ESCRT-0
component HRS, suggesting involvement of a ubiquitin-dependent endosomal pathway.
Overall, our results suggest that under proteotoxic stress and impaired autophagy, cells
activate alternative routes, potentially involving unconventional secretion, to regulate
NBR1 levels.

Introduction

(Macro)Autophagy is a catabolic process that delivers cytoplasmic components to the
lysosome for degradation through double-membrane autophagosomes. Its biogenesis
requires the coordinated action of autophagy-related (ATG) proteins. In yeast, ATG8 is
essential for autophagosome formation (1), while in mammals, this function is fulfilled
by ATG8 orthologs (mATG8s), including LC3 and GABARAP protein families. Notably,
unlike yeast, mammals can form autophagosomes even in the absence of mATG8s (2).
Selective cargo recognition is mediated by autophagy receptors such as NBR1 and
SQSTM1/p62 (3). In the absence of mATG8s, cargo can instead be secreted via
autophagy-dependent secretion (4). Notably, NBR1 compensates for the loss of ATG7,
an essential protein involved in mATG8s lipidation by promoting autophagosome
formation through an alternative mechanism (5).

Given the central role of NBR1 in coordinating selective autophagy and potential
compensatory pathways, we explored how NBR1 behaves under conditions of
proteasomal inhibition in the absence of mATG8s. For this, HeLa cells, both WT and
mATG8s KO, were compared, revealing that mATG8s KO cells displayed elevated levels
of NBR1 relative to WT. Surprisingly, proteasomal inhibition with MG132 led to a
significant decrease in NBR1 levels in mATG8s KO cells. Inhibiting lysosomal degradation
using BafA1 failed to prevent this reduction, suggesting a non-lysosomal mechanism.
Moreover, silencing of RAB27A, a GTPase involved in endo-lysosomal vesicle trafficking
and secretion, reduced basal NBR1 levels and abolished its further decrease by MG132,
likely due to already diminished levels rather than a requirement for RAB27A. Residual
NBR1 localized in punctate structures positive for ubiquitin and the ESCRT-0 component
HRS, suggesting a ubiquitin-dependent endosomal mechanism. Altogether, these
findings suggest that under proteotoxic stress and impaired autophagy, cells engage
alternative pathways, potentially involving unconventional secretion of NBR1.
100
101

102
103
Methods

Reagents and Antibodies
MG132 (cat#47479), Bafilomycin A1 (BafA1, cat#B1793), anti-NBR1 (cat#sc-130380,
NBP171703), anti-EEA1 (cat#61045), anti-CD63 (cat#ab8219), anti-Cathepsin D
(cat#AF1014), anti-HRS (cat#ab15539), anti-ubiquitin (cat#AUB01), and anti-RAB27A
(cat#168013) were used.

110
Cell Culture, Treatments, and Western Blot
HeLa WT and mATG8s KO cells (courtesy of Dr. Maho Hamasaki, Osaka University), and
HEK293T cells (cat#632180), were cultured in DMEM High Glucose (cat#12800-017)
supplemented with 10% heat-inactivated FBS (cat#sv30160.03), 100 U/mL penicillin,
and 100 µg/mL streptomycin (cat#15070-063), at 37°C and 5% CO_2. Cells were seeded
after trypsinization with 0.05% trypsin and 1 mM EDTA (cat#25300-54).
For treatments, cells were incubated with 10 µM MG132 for 8 h or 100 nM BafA1 for 4
h. DMSO was used as vehicle control.
Cells were washed with cold PBS (cat#46-013) and lysed in RIPA buffer [50 mM Tris-HCl,
150 mM NaCl, 5 mM EDTA, 1% NP-40, 1% sodium deoxycholate, 0.1% SDS, pH 7.4]
supplemented with protease inhibitors (cat#P8340). Lysates were sonicated on ice (3
pulses, 2–3 s, 40 mA) and processed for SDS-PAGE, Western blotting, and membrane
detection following previously described protocols (6).

124
125   *Immunofluorescence and Confocal Microscopy*
126   Cells grown on coverslips were fixed, permeabilized, and stained as previously described
127   (6). Images were acquired using a Leica TCS SP8 confocal microscope (6).

128
129   *Lentiviral Transduction and Stable Knockdown*
130   RAB27A knockdown was achieved by lentiviral transduction with shRNA as previously
131   described (6). mATG8s KO cells were selected with 6 µg/mL puromycin (cat#P8833).
132   Knockdown efficiency was confirmed by Western blot.

133
134   *Data Analysis*
135   Densitometry and image quantification were performed using FIJI (ImageJ), Excel 2022,
136   and Prism 10.0 (macOS). Statistical significance was determined using a one-tailed t-
137   test.

138
Results

<u>Proteasomal inhibition with MG132 reduces NBR1 levels in HeLa cells lacking mATG8s</u>
<u>proteins, independently of lysosomal degradation</u>
The absence of mATG8s in HeLa cells perturbs autophagosome trafficking, alters their
morphology, and slows down their biogenesis rate (2), ultimately leading to the
accumulation of selective autophagy receptors. Consistent with this, we confirmed the
accumulation of NBR1 in mATG8s-KO HeLa cells compared to wild-type (WT) by Western
blot analysis (Fig. 1A, lanes 1 and 3; Fig. 1B). Considering that in mATG8s-KO cells, cells
may rely on the proteasomal system to maintain proteostasis, where NBR1 can promote

substrate delivery to the proteasome, we explore whether proteasome inhibition
increases NBR1 levels. Contrary to our expectations, treatment with 10 µM MG132 for
8 h led to a significant decrease in NBR1 protein levels in mATG8s-KO cells, as shown by
Western blot (Fig. 1A line 3 and line 4; Fig. 1B) and immunofluorescence (Fig. 1C and
1D), a phenotype that was not observed in WT cells. Proteasome inhibition is known to
activate lysosomal biogenesis and autophagosome degradation (7). We tested whether
the reduction in NBR1 levels was due to lysosomal degradation. Cells were treated with
10 µM MG132 for 8 h, with 100 nM BafA1 added during the final 2 h to inhibit lysosomal
activity. Lysosomal inhibition did not restore NBR1 levels in mATG8s-KO cells (Fig. 1E
and 1F), indicating that the MG132-induced decrease in NBR1 is not mediated by
lysosomal degradation.

Residual NBR1 levels in response to catalytic proteasome inhibition co-localize with
ESCRT-0 component HRS and Ubiquitin aggregates.
To gain insight into the trafficking pathways involved, we next assessed the subcellular
distribution of NBR1 following MG132 treatment. Given that NBR1 can be incorporated
into endosomes under basal conditions in yeast (8), we evaluated its co-localization with
markers of the endolysosomal pathway in mATG8s-KO cells treated with MG132,
including early endosomes (EEA1), the ESCRT-0 component HRS (involved in endosomal
sorting (9)), late endosomes (CD63), and lysosomes (Cathepsin D, CathD). Confocal
microscopy analysis revealed a significant co-localization of NRB1 with HRS, compared
to the other markers (Fig. 2A and 2B), predominantly within large circular structures (Fig.
2A, arrowheads). Since NBR1 recognizes ubiquitinated cargos for degradation, we next
examined the co-localization of NBR1 and HRS with ubiquitin following proteasome
inhibition in mATG8s-KO cells. Confocal analysis showed that both NBR1 and HRS co-
localize with ubiquitin after MG132 treatment (Fig. 2C and 2D), within similar structures.

Proteasomal inhibition with MG132 reduces NBR1 levels in mATG8-KO cells, via a
RAB27A-independent mechanism.
Given the observed association between HRS and NBR1 after proteasome inhibition,
and that HRS mediates earlier steps in intraluminal vesicle formation at endosomes,
later released as exosomes via MVB-plasma membrane fusion regulated by the GTPase
RAB27A (9), we investigated whether NBR1 reduction could be mediated by RAB27A-
dependent secretion. To test this, we generated stable RAB27A knockdown in mATG8s-
KO cells via lentiviral transduction (shRAB27A), using cells expressing a shRNA against
luciferase as a control (shLuc). Western blot analysis revealed reduced basal NBR1 levels
in shRAB27A compared to shLuc cells (Fig. 2E, lanes 1 and 3; Fig. 2F). We next assessed
the effect of proteasome inhibition in these lines. As expected, MG132 treatment
reduced NBR1 levels in shLuc cells (Fig. 2E, lanes 1 and 2; Fig. 2F). In contrast, MG132
treatment did not further decrease NBR1 levels in shRab27A cells (Fig. 2E, lanes 3 and
4; Fig. 2F), suggesting that the reduction of NBR1 upon proteasomal inhibition occurs
through a RAB27A-independent mechanism.

Discussion

Here, we show that NBR1 protein levels, elevated in mATG8s-KO cells, decrease under
two distinct stress conditions: proteasomal inhibition and RAB27A silencing. The

remaining NBR1 signal localizes to dot-like structures that co-localize with the ESCRT-0
component HRS and ubiquitin, suggesting involvement of the ubiquitin-dependent
endosomal pathway. Consistent with this, lysosomal inhibition using BafA1 did not
restore NBR1 levels, and no intracellular accumulation was observed, supporting the
notion that its reduction is not due to lysosomal degradation. RAB27A silencing in
mATG8s-KO cells also reduced NBR1 levels, suggesting that in the absence of this
GTPase, NBR1 may be rerouted toward degradation or secretion via yet unidentified
mechanisms. Although we did not directly assess secretion, our findings align with the
possibility that NBR1 is released via a non-canonical pathway under cellular stress. The
absence of MG132-induced NBR1 reduction in RAB27A-silenced cells likely reflects
already diminished basal levels due to RAB27A depletion, rather than indicating its
requirement in the MG132 response. Thus, NBR1 downregulation upon proteasome
inhibition likely proceeds through a RAB27A-independent mechanism.

These results suggest that when canonical autophagy is compromised, cells become
more vulnerable to additional stressors such as proteasome inhibition or vesicular
trafficking disruption. In compensation, they may activate alternative pathways
involving selective autophagy receptors that bind ubiquitinated cargo, with HRS
potentially contributing. HRS has been implicated in autophagy-dependent secretory
autophagy, typically requiring LC3 (10). However, our data point to an LC3-independent
mechanism, consistent with the absence of mATG8s, where NBR1 may be redirected
into ubiquitin-enriched vesicles. Whether these vesicles are ultimately secreted, and the
nature of the underlying pathway, remain to be determined.

Given that aging and chronic proteotoxic stress impair canonical autophagy, identifying
alternative routes for protein handling may offer new insights into cellular resilience and
protein quality control.

221
Acknowledgments

We kindly thank to Drs. Manuel Varas-Godoy (Universidad San Sebastián) and Matías
Ostrowski (Universidad de Buenos Aires) for providing the plasmids used in lentiviral
particle production. We are also grateful to Dr. Nicolás Albornoz for his contribution to
the generation of raw ~~~+~data not included in this work.

Fundings

This research was funded by Fondo Nacional de Desarrollo Científico y Tecnológico de
Chile (FONDECYT) grants numbers 1211261 (Patricia V. Burgos), and 3220485 (Viviana
A. Cavieres); and Vicerrectoria de Investigación y Doctorados, Universidad San Sebastián
for PhD scholarship (Cristóbal Cerda-Troncoso).

Author contributions

Conceptualization, **C.C-T.**, **V.A.C** and **P.V.B.**; Methodology, **C.C.T.**, **A.E-M**., **B.G-R.**, **K.C.**,
and **V.A.C**; Formal analysis, **C.C-T.**, **A.E-M.**, **V.A.C.**, and **P.V.B.**; Resources, **C.C-T.**, **V.A.C.**,
and **P.V.B.**; writing—original draft preparation, **C.C-T.**, **V.A.C.**, and **P.V.B.** writing—
review and editing, **C.C-T.**, **V.A.C.**, and **P.V.B.**; funding acquisition, **V.A.C.**, and **P.V.B.**

239
240
241 **Conflict of interest statement**

242 The authors declare no conflict of interest.
243
References

1. Nakatogawa H, Ichimura Y, Ohsumi Y. Atg8 controls phagophore expansion during
autophagosome formation. Mol Biol Cell. 2008 Jul;19(7):3290-8. doi:
10.1091/mbc.E07-12-1292.

2. Nguyen TN, Padman BS, Usher J, Oorschot V, Ramm G, Lazarou M. Atg8 family
LC3/GABARAP proteins are crucial for autophagosome-lysosome fusion but not
autophagosome formation during PINK1/Parkin mitophagy and starvation. J Cell
Biol. 2016 Dec 19;215(6):857-874. doi: 10.1083/jcb.201607039.

3. Vargas JNS, Hamasaki M, Kawabata T, Youle RJ, Yoshimori T. The mechanisms and
roles of selective autophagy in mammals. Nat Rev Mol Cell Biol. 2023
Mar;24(3):167–85. doi:10.1038/s41580-022-00542.

4. Tan HWS, Luo S, Wang Z, et al. A degradative to secretory autophagy switch
mediates mitochondria clearance in the absence of the mATG8-conjugation
machinery. Nat Commun. 2022 Jun 28;13:3720. doi: 10.1038/s41467-022-31213-7.

5. Shoemaker CJ, Huang H, Weir NR, et al. Receptor-mediated clustering of FIP200
bypasses the role of LC3 lipidation in autophagy. EMBO J. 2020 Nov
23;39(23):e104948. doi: 10.15252/embj.2020104948.

6. Cerda-Troncoso C, et al. Chemo-small extracellular vesicles released in cisplatin-
resistance ovarian cancer cells are regulated by the lysosomal function. J Extracell
Vesicles. 2024 May 30; 3(6):e157.. doi: 10.1002/jex2.157

7. Zhang X, Garstka MA, Wang Z, et al. Proteasome Inhibition Activates Autophagy-
Lysosome Pathway and the Formation of Aggresome in Human Mature Dendritic
Cells. Front Cell Dev Biol. 2019;7:282. doi: 10.3389/fcell.2019.00282.

8. Liu XM, Sun LL, Hu W, Ding YH, Dong MQ, Du LL. ESCRTs cooperate with a selective
autophagy receptor to mediate vacuolar targeting of soluble cargos. Mol Cell. 2015
Sep 17;59(6):1035–42. doi:10.1016/j.molcel.2015.07.034.

9. Hessvik NP, Llorente A. Current knowledge on exosome biogenesis and release. Cell
Mol Life Sci. 2018 Jan;75(2):193-208. doi: 10.1007/s00018-017-2595-9.

10. Leidal AM, Huang HH, Marsh T, Solvik T, Zhang D, Ye J, Kai F, Goldsmith J, Liu JY,
Huang YH, Monkkonen T, Vlahakis A, Huang EJ, Goodarzi H, Yu L, Wiita AP, Debnath
J. LC3-dependent EV loading and secretion (LDELS) promotes TFRC secretion and
microenvironment remodeling. Nat Cell Biol. 2023 May;25(5):741-754. doi:
10.1038/s41556-023-01154-3.

284 **Figures and legends**

Figure 1
Cerda-Troncoso C, et al., 2025

291
292
293
Figure 1. Catalytic proteasome inhibition reduces NBR1 levels in mATG8-deficient cells.
(A) Western blot of endogenous NBR1 and β-actin in HeLa WT and HeLa mATG8s KO
cells treated with 10 µM MG132 for 0 or 8 h. Cells were lysed with RIPA buffer. (B)
Densitometric quantification of NRB1 levels shown in (A). NS: No-Significative **P<0.01,
***P<0,001; unpaired-parametric t-Test. (C) Immunofluorescence analysis of NBR1
distribution in HeLa WT and HeLa mATG8s-KO cells treated with 10 µM MG132 for 0 or
8 h; Scale bar 10 µm. (D) Quantification of NBR1 positive puncta per cell from images in
(C); ***P<0,001; unpaired-parametric t-Test. (E) Western blot of endogenous NBR1 and
β-actin in HeLa mATG8s KO cells treated with 10 µM MG132 for 0 or 8 h, with or without
100 nM BafA1 added during the final 2 h. (F) Densitometric quantification of NBR1 levels
from (E); NS: No-Significative, *P<0.05, ***P<0.001; unpaired-parametric t-Test.

338
339
340

Figure 2
Cerda-Troncoso C, et al., 2025

349
350
Figure 2. Residual NBR1 localizes to HRS and ubiquitin structures following catalytic
proteasome inhibition. (A) Confocal analysis of NBR1 with endolysosomal markers in
mATG8s KO cells treated with 10 μM MG132 for 0 or 8 h. Markers include EEA1 (early
endosomes), HRS (MVBs), CD63 (secretory MVBs), and CathD (lysosomes); arrowheads
indicate areas of co-localization. (B) Representative quantification of NBR1 co-
localization with HRS from panel (A). (C) Mander's coefficient quantification of NBR1 co-
localization with the indicated endolysosomal markers after MG132 treatment;
***P<0.001; unpaired-parametric t-Test. (D) Confocal analysis of NBR1 or HRS co-
localization with ubiquitin after 0 or 8 h of 10 μM MG132 treatment in mATG8s KO cells;
arrowheads show co-localization. (D) Mander's coefficient quantification of ubiquitin
co-localization with NBR1 or HRS following MG132 treatment as shown in (C);
***P<0.001; unpaired-parametric t-Test. (F) Western blot analysis of NBR1 and β-actin
protein levels in mATG8s KO either shLuc and shRAB27A cells, treated with 10 μM
MG132 for 0 or 8 h. (G) Densitometric quantification of Western blot shown in (F); NS:
No-Significative **P<0.01; unpaired-parametric t-Test.
366