In vitro **anti-cancer properties of a commercially available polyherbal nutraceutical**
(Vernolac) capsule on cancer stem cell-like (NTERA-2 cl.D1) cells.

Poorni Chanuka Rathnayake[1], Duvinika Chalani Senavirathna[1], Matheen Muhammadh Milhan[1],

Sandani De Vass Gunawaradane[1], Nirwani Natasha Seneviratne[1], Dipun Nirmal Perera[1],

Umapriyatharshini Rajagopalan[1], Kanishka Sithira Senathilake[1], Sameera Ranganath

Samarakoon[1*]

[1]Institute of Biochemistry, Molecular Biology and Biotechnology, University of Colombo

No. 90, Cumaratunga Munidasa Mawatha, Colombo 03, Sri Lanka.

*Corresponding author:

Sameera R Samarakoon

Institute of Biochemistry, Molecular Biology and Biotechnology, University of Colombo

No. 90, CumaratungaMunidasa Mawatha, Colombo 03, Sri Lanka.

Telephone: +94 11 2552528 (Ext- 318)

Fax: +94-11-2553683

Email: sam@ibmbb.cmb.ac.lk

Abstract

Vernolac is a commercially available polyherbal nutraceutical formulation comprising *Vernonia zeylanica* aerial parts, *Nigella sativa* seeds, *Hemidesmus indica* roots, *Smilax glabra* rhizome, and *Leucas zeylanica* aerial parts. Previous studies have demonstrated anti-cancer activities of phytochemicals derived from these individual plant components. However, the anti-cancer properties of the supercritical CO_2 extract of Vernolac remain unexplored against cancer stem-like cell populations. The current study is focused on the anti-cancer potential of Vernolac extract on NTERA-2 cl.D1 cancer stem-like model, a human embryonal carcinoma-derived pluripotent cell line. Gas Chromatography-Mass Spectrometry (GC-MS) was performed for phytochemical analysis. Several in vitro assays evaluated the anti-cancer properties of the Vernolac extract. Cytotoxicity was assessed using the Sulforhodamine B assay, and apoptosis induction was determined by Acridine Orange/Ethidium Bromide staining and the caspase 3/7 activity assay. Scratch assay was used to evaluate cell migration, and Reverse Transcriptase quantitative Polymerase Chain Reaction (RT-qPCR) was performed to analyze the expression levels of apoptosis-related genes (*TP53, BIRC5*) and the autophagy-related gene *mTOR*. Further, the free-radical scavenging activity of Vernolac extract was assessed using 2,2-diphenyl-1-picrylhydrazyl and 2,2′-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) assay. The reactive oxygen species levels in NTERA-2 cl.D1 cells were quantified using the nitroblue tetrazolium assay. GC-MS analysis revealed 20 phytochemical constituents in the supercritical CO_2 extract. *In vitro* assay results demonstrated that Vernolac extract exhibits significant anti-proliferative activity in NTERA-2 cl.D1 cells, with an IC_{50} of 41.12 µg/mL at 48 h, while exerting minimal effects on non-cancerous MCF-10A cells ($IC_{50} > 1000$ µg/mL). Fluorescence microscopy and caspase 3/7 assay showed that Vernolac extract leads to early apoptosis in NTERA-2 cl.D1 cells. RT-qPCR revealed the upregulation of tumor suppressor protein P53 while downregulating *BIRC5* and *mTOR*. Additionally, Vernolac extract inhibited the migration rate of NTERA-2 cl.D1 cells and elevated intracellular reactive oxygen species levels. These findings suggest that the supercritical CO_2 extract of Vernolac exerts potent anticancer properties against NTERA-2 cl.D1 cancer stem-like model, highlighting its therapeutic potential for targeting cancer stem cells.

Key words: Anti-cancer, Vernolac, Supercritical CO_2 extraction, Nutraceutical, Cancer Stem Cell like models

Introduction

Cancer is one of the most common causes of mortality worldwide [1]. Despite various cancer therapies being available, conventional treatments often show limited success due to the challenges, including treatment resistance and tumor recurrence, and adverse side effects [1,2]. These limitations are mainly due to a subpopulation of tumor-initiating cells known as cancer stem cells (CSCs), which possess self-renewal and multi-lineage differentiation capabilities, making them resistant to many conventional therapies, thereby sustaining tumor heterogeneity and recurrence [3–6]. Different treatment strategies are being investigated to overcome the limitations of conventional treatment options. Natural pharmaceuticals are a treatment option currently being investigated for their favorable properties, such as fewer side effects, diverse modes of action, and reduced potential of treatment resistance [7–12]. Moreover, polyherbal extracts are emerging as potential cancer treatment options, and numerous phytochemicals in traditional polyherbal formulations have been shown to exert anticancer effects, with several constituents targeting CSCs through various molecular pathways [13–15]. The NTERA-2 cl.D1 cell line, a well-characterized human pluripotent embryonal carcinoma cell line, is frequently used as a model to study CSC-like model phenotype behavior and evaluate therapeutic responses, serving as a validated *in vitro* platform for evaluating novel anticancer agents targeting stemness-associated pathways [14,16].

Commercially available nutraceutical 'Vernolac' is coined after the main constituent *Vernonia zeylanica*. Vernolac is formulated using five different herbal plants, including *Vernonia zeylanica* aerial parts, *Nigella sativa* seeds, *Hemidesmus indica* roots, *Smilax glabra* rhizome, and *Leucas zeylanica* aerial parts [8]. Natural phytochemicals in nutraceuticals reveal multiple advantages through their antioxidant, anti-inflammatory, and cancer prevention potentials [17–23]. Alpha-hedrin, Thymoquinone, and Vernolactone are some of the phytochemicals present in Vernolac formulation, which possess anticancer, antioxidant, and anti-inflammatory properties [15,24–30]. An *in vitro* study conducted by Abeysinghe et al. [15] and Mendis et al. [22] demonstrated that Vernolactone in *V. zeylanica* triggers the antiproliferative activity and apoptosis in NTERA-2 cl.D1 cells, as well as the triads of breast cancer cells (MDAMB-231, MCF-7, SKBR-3) in a time and dose-dependent manner. In addition, Samarakoon et al. [23], Galhena et al. [25], and Pathiranage et al. [26] have reported that a polyherbal mixture comprised of *N. sativa seeds, H. indicus* roots and *S. glabra* rhizome contain possible anticancer and anti-inflammatory effects.

Although the anticancer effects of individual phytochemicals present in the constituent plants of
the Vernolac have been demonstrated in various studies [13,15,22,23,31–34], the integrative
anticancer effect of the complete polyherbal formulation extracted using the supercritical CO_2
method has not yet been evaluated, particularly on CSC-like cell populations. Therefore, the
present study was designed to investigate the anticancer potential of the supercritical CO_2 extract
of Vernolac against NTERA-2 cl.D1 cells, to elucidate its efficacy as a potential CSC-targeted
anticancer therapeutic.

Materials and methods

Preparation of Vernolac extract

A polyherbal formulation comprised of *V. zeylanica* aerial parts, *N. sativa* seeds, *H. indica* roots,
S. glabra rhizome, and *L. zeylanica* aerial parts was extracted using the supercritical fluid
extraction method. Plant materials were collected in September 2024 (Batch No. V24085). The
extraction was performed using supercritical carbon dioxide (CO_2) at 42°C, at 30 MPa pressure
and the extracts were collected at a CO_2 flow rate of 4 mL/min for 1 h.

Cell culture

All the cell lines were cultured and maintained according to American Type Culture Collection
(ATCC) guidelines. NTERA-2 cl.D1 (ATCC CRL-1973) cells were cultured in Dulbecco's
Modified Eagle medium (DMEM), and MCF-10A cells were cultured in Mammary Epithelial Cell
Basal Medium (MEBM). Both the media types were supplemented with 10%, fetal bovine serum
(FBS), 50 U/mL penicillin, and 50 µg/mL streptomycin. Cells were maintained in a humidified
incubator with 5% CO_2 at 37°C.

Determination of cell viability by SRB assay

Briefly, cells were cultured in 96-well cell culture plates at a density of 7.5×10^3 cells/well in 200
µL of medium for 24h. Cells were treated with varying concentrations of Vernolac extract (6.25,
12.5, 25, 50, 100, 200, and 400 µg/mL) with < 0.1 % DMSO and incubated for 24 and 48h. After
incubation, sulforhodamine B (SRB) assay was performed as described in Rajagopalan et al. [35]
with minor modifications. Absorbance was measured at 540 nm using a Synergy HT microplate
reader, BioTek Instruments, USA.

Fluorescence microscopic analysis

As described by Abeysinghe et al. [12], NTERA-2 cl.D1 cells (2×10^5 cells/mL) were cultured on a cell culture-treated coverslip and incubated for 24h. The cells were treated with different doses of Vernolac (5.0, 10, and 20 µg/mL) and incubated for 24h. Paclitaxel (1.0, 2.0, and 4.0 µg/mL) was used as a positive control. At 24h post-treatment, cells were washed with ice-cold PBS and fixed with 4% formaldehyde solution in the dark for 10 minutes. Cells were stained using Acridine orange and Ethidium bromide (AO\EB) solution (20 µL). Excess stains were removed with PBS (1 mL); cells were air-dried prior to mounting with anti-fading solution. The stained cells were observed under the fluorescence microscope (BX51 TRF, Olympus Corporation, Tokyo, Japan).

Caspase 3/7 luminescence assay

NTERA-2 cl.D1 cells were cultured in a cell culture-treated 96-well plate (20,000 cells/ well in 100 µL of media) and incubated for 24h. Cells were treated with different concentrations of Vernolac extract (2.5, 5.0, 10, and 20 µg/mL) and incubated for 24h. Caspase 3/7 Glo$^{®}$ reagent (100 µL) was added to each treated well and incubated for 1h in the dark condition at 37°C as described by Ediriweera et al. [36]. After incubation and 15 minutes in the shaker, Luminescence was measured using a microplate reader (Synergy HT, Biotek Instruments, USA).

Nitrobluetetrazolium Reactive Oxygen Species (NBT-ROS) assay

NTERA-2 cl.D1 cancer cells (5×10^4 cells/well) were cultured in a 96-well plate using DMEM growth medium with 10% FBS and incubated for 24h. The extract was treated with Vernolac extract 2.5, 5.0, 10, and 20 µg/mL and incubated for 24h. At 24h post-incubation, the NBT-ROS assay was performed according to Ediriweera et al. [37]. Using the Synergy HT microplate reader, absorbance was measured at 560 nm, and ROS expression levels were analyzed as percentage relative values of the untreated control.

2,2-diphenyl-1-picrylhydrazyl (DPPH) assay

Free-radical scavenging capacity was measured using 2,2-diphenyl-1-picrylhydrazyl (DPPH) reagent according to the method described by Ediriweera et al. [36]. Volume of 50 µL from each concentration of Vernolac extract (0.01, 0.023, 0.046, 0.093, 0.1875, 0.375, 0.75, 1.5, and 3 µg/mL) prepared with DMSO by the double dilution method. was added in 96-well plates. Samples were further diluted with methanol (90 µL/well) and mixed with 60 µL of DPPH solution (2 mg/ mL) in each well in a dark environment. Ascorbic acid was used as a positive control. The plates were incubated at room temperature in the dark for 10 min, and the absorbance was measured at 517 nm using Synergy HT microplate reader. Free-radical scavenging capacity of each concentration was calculated in percentage using the formula (Absorbance $_{control}$ – Absorbance $_{sample}$)/Absorbance $_{control}$ × 100%, where control contains all the reagents used except Vernolac extract.

2,2′-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) assay

The 2,2′-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) radical scavenging activity of Vernolac extract was measured described by num A dilution series of Vernolac extract (7.8125, 15.625, 31.25, 62.5, 125, 250, 500, and 1000 µg/mL) was prepared in PBS prior to the assay. Vernolac (160 µL) and ABTS (40 µL) were added to each well, and the plate was incubated for 10 min at room temperature. Ascorbic acid was used as a positive control. Absorbance of each well was measured at 734nm using a microplate reader. The percentage radical scavenging activity was calculated as (Abcontrol – Absample) /(Abcontrol) × 100.

Cell migration assay

NTERA-2 cl.D1 cells were seeded in 96-well plates at a density of $2x10^4/200$ µL and incubated up to 100% confluence until a fully confluent cell monolayer was observed. As described by Abeysinghe et al. [12], a vertical scratch was made on the monolayer formed in each well and washed twice with PBS to remove detached cells and debris. Then, the cells were treated with

different concentrations of Vernolac extract (5, 10, 20, and 40 µg/mL) and paclitaxel as the positive

control in separate wells. Untreated cells were used as controls. Closure of the scratch made was

observed by a phase-contrast microscope at every 12h of incubation, and the area of closures was

calculated using the recorded images. The rate of migration affected by respective treatments was

calculated using the change in the cell-free area (wound) over time.

Reverse Transcriptase quantitative polymerase chain reaction (RT-qPCR)

NTERA-2 cl.D1 cells (2.5×10^5 cells/mL) were seeded into 25 cm^2 cell culture flasks and incubated

for 24h in 37 C and 5 % CO2 : 95 % air environment. Then the cells were exposed to different

concentrations (10 and 20 µg/mL) of Vernolac extract under the previously mentioned conditions.

Total RNA was isolated using PromegaSV Total RNA isolation system, USA (Cat. No: Z3100SV)

according to the manufacturer's instructions. Complementary DNA (cDNA) from isolated total

RNA was synthesized using Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase

system (Cat.no: M1701, Promega Corporation, Madisons, USA). Gene expression profiling of

TP53, *BIRC5,* and *mTOR* was carried out with *GADPH* gene as the internal control using

QuantiNova® SYBR® Green PCR Kit (Cat. No. 208052, Qiagen) and the qPCR reaction was

carried out with QuantStudio® 5dx Real-Time PCR Systems (Applied Biosystems, Massachusetts,

USA) on 96-well PCR plates. *GADPH* was used as the internal control. The relative gene

expression analysis was performed using the Livak method, where fold changes were calculated

using $2^{-\Delta\Delta Ct}$ equation [38]. Primers used for gene expression profiling are given in Table 1, and the

PCR cycle conditions were as follows: initial denaturation at 95°C for 10 min, and amplification

in three steps for 40 cycles (denaturation at 95˚C for 30 s; annealing at 58˚C for 1 min, and

extension at 60˚C for 30 s). Annealing temperature for gene *mTOR* was maintained at 52˚C.

Table 01: Primers used for expression of *P53, BIRC5, mTOR, GAPDH*

Gene	Forward & reverse primers
P53:	Forward: 5′-TCTGGCCCCTCCTCAGCATCTT
	Reverse: 5′-TTGGGCAGTGCTCGCTTAGTGC
BIRC5:	Forward:5′-TGGCCGCTCCTCCCTCAGAAAA
	Reverse:5′- GCTGCTGCCTCCAAAGAAAGCG
mTOR:	Forward: 5′- TGGGACAGCATGGAAGAATA
	Reverse: 5′- TGTTGTGCCAAGGAGAAGAG
GAPDH:	Forward: 5′- GGCATTGCCCTCAACGACCAC
	Reverse: 5′- ACATGACAAGGTGCGGCTCCCTA

Gas chromatography and mass spectroscopic analysis (GC-MS)

GC-MS analysis was conducted on an Agilent 7890 GC system coupled with a 5975 MS detector (Agilent Technologies, Inc., Santa Clara, CA, USA). Separation was achieved on a J&W DB-5 MS capillary column (30.0 m × 250 μm, film thickness 0.25 μm; 5% phenyl methyl siloxane). Ultra-high pure helium (99.999%) served as the carrier gas at a constant flow rate of 1 mL/min. Injections were performed in split mode with a 1:10 ratio using a 1 μL sample volume. Electron ionization was performed at 70 eV ionization energy, while injector and detector temperatures were maintained at 260 °C and 320 °C, respectively. The oven temperature program started at 110 °C (held for 5 min), increased to 280 °C at 20 °C/min (held for 1 min), and then further ramped to 320 °C at 20 °C/min (held for 5 min). Data acquisition and instrument control were managed using MS Solution software. Compound identification was achieved by comparing mass spectra with those in the NIST (National Institute of Standards and Technology) library.

Statistical analysis

Data analysis was performed using GraphPad Prism 8.0.1 (GraphPad Software Inc., San Diego, CA, USA). Each experiment was triplicated (n=3) for biological triplicates. One-way analysis of

variance (ANOVA) was used to compare means across multiple groups in the caspase 3/7 activity
assay and NBT-ROS assay, followed by Bonferroni's post hoc test for multiple pairwise
comparisons. For gene expression analysis, one-way ANOVA followed by Dunnett's post hoc test
was performed to compare treatment groups against the control group. Statistical significance was
set at $P < 0.05$.

Results

Vernolac extract Characterization

The Vernolac extract yielded 3% (Batch No. V24085). The GC–MS analysis revealed 20 distinct
compound peaks. Table 02 summarizes the principal phytochemicals identified in the extract.

Table 02: GC/MS analysis of Vernolac Extract

	Compounds
GC/MS analysis	Alpha -Tocopherol acetate Alpha-pinene Beta-Myrcene Beta-pinene D-Limonene Dodecanoic acid 1,6-Octadien-3-ol, 3,7-dimethyl- Carvacrol Cycloartenol Thymoquinone 1,3,5-Trimethyl benzene Alpha terpinyl acetate Hexadecanoic acid Linalyl acetate Stigmasterol Octadecanoic acid Oleic acid Palmitic acid Lauric acid Acetic acid,phenylmethyl ester

Notably, several bioactive phytochemicals were identified, including α-Tocopherol acetate, Thymoquinone, Carvacrol, Cycloartenol, Stigmasterol, and Oleic acid, which are known for their antioxidant, anti-inflammatory, and therapeutic properties. Other detected compounds included monoterpenes (α-Pinene, β-Myrcene, β-Pinene, D-Limonene, Linalyl acetate, α-Terpinyl acetate), fatty acids (Lauric acid, Palmitic acid, Hexadecanoic acid, Octadecanoic acid), and additional aromatic or ester derivatives (1,3,5-Trimethyl benzene, Acetic acid phenylmethyl ester).

Cytotoxicity and cell proliferation assay

As indicated by the results of the SRB assay (Table 03), Vernolac extract exerted a potential to mediate dose- and time-dependent suppression of NTERA-2 cl.D1 cell proliferation. Both Vernolac extract, and Paclitaxel showed a dose and time-dependent anti-proliferative activity against NTERA-2 cl.D1 cells.

Table 03: IC$_{50}$ values (µg/mL) of Vernolac extract and Paclitaxel in NTERA-2 cl.D1 and MCF-10 cells.

Cell type	Vernolac extract (µg/mL)		Paclitaxel (µg/mL)	
	24h	48h	24h	48h
NTERA-2 cl.D1 cells	64.5	41.12	4.96	1.79
MCF-10A cells	1075	803.5	11.90	6.70

Fluorescence microscopic observations

Acridine Orange (AO)/Ethidium Bromide (EB) stained NTERA-2 cl.D1 cells were incubated 24h after the treatment with different concentrations of Vernolac extract and Paclitaxel. By employing Acridine orange/Ethidium Bromide staining, clear identification of apoptosis-associated membrane alterations was achieved. The apoptotic changes of the cells were distinguished and visualized by the fluorescence staining, yellow/orange fluorescence indicates early apoptotic cells while red fluorescence indicates late apoptosis, demonstrating disruption of cellular membrane. The induction of apoptosis was increased in Vernolac extract (5, 10 and 20 µg/mL) - and Paclitaxel (1, 2, and 4 µg/mL) treated NTERA-2 cl.D1 cells in a dose-dependent manner (Figure 01).

Figure 01: AO/EB stained Fluorescence microscopic observation of NTERA-2 cl.D1 cells under 200X magnification 24h post-treatment incubation. A – Untreated control, B – Paclitaxel (1 µg/mL), C – Paclitaxel (2 µg/mL), D – Paclitaxel (4 µg/mL), E –Vernolac extract (15 µg/mL).F– Vernolac extract - (30 µg/mL), G –Vernolac extract (60 µg/mL).

Expression of caspase 3 and Caspase 7 activity in NTERA-2 cl.D1 cells

Activities of caspase 3 and 7 were assessed using the Caspase Glo®3/7 assay kit (Promega, USA) according to the manufacturer's guidelines. Results are presented as the mean $\pm$ standard deviation values. Vernolac extract significantly increased the activities of caspase 3 and caspase 7 in NTERA-2 cl.D1 cells at the concentrations of 10 µg/mL (P < 0.001) and 20 µg/mL (P < 0.0001) as shown in Figure 02.

Caspase 3/7 activity of Extract

Figure 02: Expression of Caspase 3 and 7 in NTERA-2 cl.D1 cells after 24h post-treatment with

Vernolac extract. (Results are indicated as mean $\pm$ standard deviation of duplicate determination.

*** P < 0.001 and ****P < 0.0001 compared to the control).

Antioxidant activity of Vernolac Extract

The 2,2-diphenyl-1-picrylhydrazyl **(**DPPH) assay and 2,2′-Azino-bis(3-ethylbenzothiazoline-6-

sulfonic acid) (ABTS) were performed to evaluate the antioxidant activity of Vernolac extract.

Ascorbic acid was used as a positive control in both assays. Vernolac extract showed a low free

radical scavenging activity (EC_{50} > 1000 µg/mL) in both ABTS and DPPH assays. The obtained

EC_{50} values of free radical scavenging activity of Vernolac extract and Ascorbic are summarized

in Table 04.

Table 04: EC_{50} values of free radical scavenging activity for Vernolac extract and Ascorbic acid.

Sample	EC_{50} values (µg/mL)
Vernolac extract	> 1000
Ascorbic acid	53.678

Effect of the Vernolac extract on ROS levels

NBT-ROS assay was performed to analyze the ROS production in NTERA-2 cl.D1 cells after 24h of Vernolac extract. A $P < 0.001$ was considered to indicate a strong statistically significant result. As seen in the bar graph (Figure 03), treatment with 10 µg/mL or 20 µg/mL of Vernolac extract caused a major increase in ROS activity compared to the non-treated control group ($P < 0.001$).

Figure 03: Levels of reactive oxygen species (ROS) detected in NTERA-2 cl.D1 cells after 24h incubation with Vernolac extract treatment under different concentrations (Results are indicated as mean $\pm$ Standard deviation. ***$P < 0.001$ compared to the control).

Analysis of the wound healing rate of NTERA-2

The NTERA-2 cl.D1 cells were incubated until they created a confluent monolayer at the bottom of the well. An artificial gap was made on the confluent monolayer, and both Vernolac extract and Paclitaxel were treated separately according to a concentration gradient. After 24h of drug treatment, untreated control cells have migrated at a rate of 62.126 µm/h. The cell migration rate of Vernolac extract and Paclitaxel is graphically represented as shown in Figure 04. The morphological representation of cell migration at 0h and after the gap of control closure is shown in Figure 05.

Figure 04: Cell migration rates of Vernolac extract with the concentrations of 5, 10, 20, and 40 µg/mL and Paclitaxel 0.25, 0.5, and 1 µg/mL treated NTERA-2 cl.D1 cells compared to untreated control under different concentrations.

Figure 05: Phase contrast light microscopic images under 10× magnification captured after 0h of creating the gap in NTERA cl.D1 cell monolayer, followed by Vernolac extract with 5, 10, 20, and 40 µg/mL concentrations (A-D), paclitaxel with 0.25, 0.5, and 1 µg/mL concentrations (I–K), and the Untreated control (O). Cell migration after 22h of creating the gap, followed by Vernolac extract (E-H), paclitaxel (L–N), and Untreated control (P).

Effect of Vernolac extract on the expression of *P53*, *BIRC5,* and *mTOR*

Results of the RT- qPCR (Figure 06) demonstrated that the Vernolac extract enhanced *TP53* mRNA expression in human NTERA-2 cl.D1 cells in a significant (P < 0.001) dose-dependent manner. Furthermore, treatment with all concentrations of supercritical Vernolac extract and Paclitaxel resulted in a significant downregulation (P < 0.05) of the expression of *mTOR* and *BIRC5*.

Figure 06: mRNA expression of selected genes on Vernolac extract and Paclitaxel treatment. (A-Vernolac extract on *TP53* expression, B - Vernolac extract on *BIRC5* expression, C - Vernolac extract on *mTOR* expression, D- Paclitaxel on *TP53* expression, E - Vernolac extract on *BIRC5* expression, F - Vernolac extract on *mTOR* expression). (Results are indicated as mean ± standard deviation, duplicate determination. *P < 0.05, **P < 0.01, ***P < 0.001, and ****P < 0.0001. compared to the control).

Discussion

Previous studies have shown that traditional polyherbal formulation demonstrates significant anti-proliferative activities against several cancer cell lines, including MCF-7, HepG2, and NCI-H292 [23,26,32,39,40]. Sailu et al. [13] and Abeysinghe et al. [15] have elucidated that phytochemicals such as alpha-hederin and Vernolactone present in the constituents of Vernolac have proven anticancer activity against cancer stem-like cells. Chemical analysis using GC/MS identified 20 compounds in all the biological Triplicates of Vernolac extract (Table 02). The present *in vitro* investigation employed supercritical fluid extraction with liquid CO_2, a superior extraction methodology that preserves thermolabile compounds while eliminating organic solvent residues, to evaluate the anticancer efficacy of Vernolac extract against the pluripotent human testicular embryonal carcinoma (NTERA-2 cl.D1) cell line.

Consistent with these earlier findings, as shown by Abeysinghe et al. [12], the profiles obtained through SRB assay of this study demonstrated dose and time-dependent cytotoxicity against NTERA-2 cl.D1 cells. Vernolac extract showed an anti-proliferative effect against the cells with an IC_{50} value of 41.12 µg/mL after 48h treatment, while demonstrating significantly lower cytotoxicity against non-tumorigenic epithelial MCF-10A cells (IC_{50} value of 1075 µg/mL), suggests selective cytotoxicity towards cancer stem cells (Table 03). 'Vernolactone', which is an active compound of the polyherbal isolated from *V. Zeylanica*, has been shown a significant cytotoxicity against NTERA-2 cl.D1 (0.4808 µg/mL at 72h) cells with Paclitaxel [15]. Similarly, Mendis et al. [22], Samarakoon et al. [23] and Pathiranage et al. [26] have also reported that a polyherbal mixture, including several constituents of Vernolac (*N. sativa, H. indicus* and *S. glabra,*) displayed a dose-dependent inhibition of cell proliferation in treated cancer cell lines (MCF-7, MDA-MB-231, HepG2), aligning with the findings.

The fluorescent and phase-contrast microscopic data revealed that Vernolac extract induces dose-dependent apoptosis-related cytomorphological changes in NTERA-2 cl.D1 cells (Figure 01), including cell membrane blebbing and cell shrinkage. Furthermore, compared to the untreated cells (control), Vernolac extract significantly ($P < 0.0001$) enhanced the enzymatic activities of executioner caspases, caspase 3, and caspase 7 at a 20 µg/mL concentration, indicating activation of the terminal phase of programmed cell death (Figure 01). Concentration gradient for the Caspase 3/7 assay was obtained after several dose optimizations, and depending on the Caspase 3/7 activity

results, the concentration of Vernolac extract for the other core assays, including NBT-ROS and
RT-qPCR has been decided. These findings align with previous research done by Abeysinghe et
al. [15], who reported that Vernolactone significantly (P < 0.0001) upregulated caspase 3/7 activity
at both 2 μg/mL and 4 μg/mL concentrations when compared to the untreated NTERA-2 cl.D1
cells (untreated control). A traditional polyherbal mixture formulated using *N. sativa*, *H. indicus*
and *S. glabra* has also demonstrated an elevated caspase 3/7 activation against several cancer cell
lines (HepG2, MCF-7, MDA-MB-231, SKBR-3), suggesting a conserved apoptotic mechanism
among these phytotherapeutic agents [22,26,32]. Further, extracts of *N. sativa* seed extract
demonstrated an elevated caspase 3 activity on the MDA-MB-231 breast cancer cell line.

In this study, the DPPH and ABTS free radical scavenging assays revealed that Vernolac extract
contains a low antioxidant activity (EC_{50} value > 1000 μg/mL) compared to the positive control:
Ascorbic acid (EC_{50} = 53.678 μg/mL) (Table 03). The obtained result aligns with a previous study
demonstrating that Vernolactone possesses low antioxidant activity (free radical scavenging
capacity) through the DPPH and ABTS assays [15]. However, multiple investigations have
indicated that *N. sativa, H. indicus,* and *S. glabra* extracts have shown free radical scavenging
activity through multiple analyses [26,31]. Comprehensive antioxidant profiling using
complementary analysis is recommended to fully characterize the antioxidant capacity of Vernolac
extract. Despite, extract of Vernolac having a minimal antioxidant activity, a significantly (P <
0.001) elevated ROS level was determined in NTERA-2 cl.D1 at the concentrations of 10 and 20
μg/mL (Figure 03). This pro-oxidant effect in cancer cells, contrasting with limited antioxidant
capacity in cell-free systems, suggests a selective modulation of redox homeostasis within the
cellular microenvironment, potentially contributing to the observed cytotoxicity [41].

The effect of Vernolac extract on the migration of NTERA-2 cl.D1 cells was evaluated using a
wound healing assay. A dose-dependent migration rate (< 3 μm/h across all tested concentrations)
(Figure 04), demonstrated minor or no reduction in the wound width throughout the experiment in
Vernolac-treated cells compared to the untreated NTERA-2 cl.D1 cells (Figure 05), indicates
significantly reduced migratory capacity. Similarly, paclitaxel also restricted cellular migration to
rates below 1.5μm/h for all tested concentrations. Low serum concentrations in the cell medium
were maintained to ensure the suppression of cell proliferation [42]. This pronounced inhibition

of cellular motility indicates impairment of cytoskeletal reorganization and extracellular matrix
interaction, critical processes in tumor cell invasiveness and metastatic potential [43].

Moreover, RT-qPCR was performed to evaluate the mRNA expression of several autophagy and
apoptosis-related genes (*TP53*, *BIRC5,* and *mTOR*). The tumor suppressor *TP53* is a key gene that
is essential for DNA repair, apoptosis, and triggering of cell cycle arrest, and plays a role in
autophagy [44]. Moreover, *p53* involves suppression of *BIRC5* at both mRNA and protein
expression levels to promote apoptosis [43]. Lower doses of Vernolac extract and paclitaxel were
used to maintain the cells in viable conditions until the cells were under the early apoptotic stage
at the time of RNA extraction. Vernolac extract led to a significant (P < 0.001) upregulated tumor
suppressor *TP53* expression, with a 9.67-fold change at 10 $\mu g/mL$ and 13.29 fold change at 20
μg/mL compared to the untreated control (Figure 06). Concurrently, Vernolac extract-treated
NTERA-2 cells resulted in significant (P < 0.05) downregulation of mammalian target of
rapamycin (*mTOR*) and *BIRC5* mRNA expression at both 10 and 20 μg/mL concentrations(Figure
06). According to Abeysinghe et al. [15], it was shown that Vernolactone in *V. zeylanica*
significantly induced apoptosis and autophagy in NTERA-2 cl.D1 cells through modulation of the
PI3K/Akt/mTOR pathways. Key phytochemicals of *Nigella sativa*; Beta catenin, Alpha-hederin
and Thymoquinone, have been reported to inhibit DNA synthesis, induce cell cycle arrest, and
promote apoptosis via both p53-dependent and p53-independent mechanisms [13,33,34,45].
Collectively, the above-mentioned studies indicate that active compounds in the constituents of
Vernolac extract exert their anticancer effects through coordinated modulation of apoptotic and
autophagy-related pathways.

Vernolac extract comprises *V. zeylanica* aerial parts, *N. sativa* seeds, *H. indica* roots, *S. glabra*
rhizome, and *L. zeylanica* aerial parts investigated in the present study, exerting anti-proliferative
and anti-cancer effects via apoptosis, autophagy, and cell cycle arrest through upregulation of
mRNA expression of *TP53*and downregulation of *BIRC5* and *mTOR* mRNA expression. The
observed cytotoxic and apoptotic effects likely result from the action of a single active compound
and/or synergistic interactions among various phytochemicals within the extract, potentially
offering advantages over single-compound therapies [13,22,35,45]. Furthermore, in vitro
investigations are necessary to demonstrate and confirm the precise molecular mechanisms
underlying the anticancer activity, and subsequent *in vivo* studies are essential to evaluate its

efficacy and the biological effects of Vernolac extract within living organisms (physiologically

relevant systems), particularly its safety profile.

Conclusion

This study demonstrates that Vernolac extract showed selective antiproliferative effects on

NTERA-2 cl.D1, cancer stem-like cells by inducing apoptosis and autophagy, while exhibiting

minimal cytotoxicity toward noncancerous epithelial cells (MCF-10A). These findings support the

potential of Vernolac extract as a promising therapeutic candidate targeting cancer stem cells.

Limitations of this study include the absence of assays such as CSC markers, autophagy flux

measurements, apoptosis validation, and migration controls. These limitations will be addressed

in future investigations on Vernolac.

Acknowledgments

The authors gratefully acknowledge the support provided by the Institute of Biochemistry, Molecular Biology and Biotechnology (IBMBB), University of Colombo. The authors gratefully acknowledge Fadna Life Science Pvt. Ltd for kindly providing the Vernolac extract and supporting its chemical characterization, which greatly contributed to the advancement of this research.

Disclosure statement.

The authors report there are no competing interests to declare.

Source of Funding

This work was funded by the Master of Science project, Institute of Biochemistry Molecular Biology and Biotechnology, University of Colombo, Sri Lanka.

CRediT author statement

PCR: Investigation, Formal analysis, Writing – original draft, Writing – review and editing ; **DCS, MMM**: Writing – original draft, Writing – review and editing ; **SDVG, NNS, DNP**: Writing – review and editing ; **UR,KSS**: Supervision, Writing – review and editing ; **SRS**: Conceptualization, Writing – review and editing, Supervision, Project administration.

References

[1]   Siegel RL, Kratzer TB, Giaquinto AN, Sung H, Jemal A. Cancer statistics, 2025. CA Cancer J Clin 2025. https://doi.org/10.3322/caac.21871.

[2]   Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin 2024;74(3):229–63. https://doi.org/10.3322/caac.21834.

[3]   Pal R, Ravindran G. Assessment of pluripotency and multilineage differentiation potential of NTERA-2 cells as a model for studying human embryonic stem cells. vol. 39. 2006.

[4]   Khan SU, Fatima K, Aisha S, Malik F. Unveiling the mechanisms and challenges of cancer drug resistance. Cell Communication and Signaling 2024;22(109):1-25. https://doi.org/10.1186/s12964-023-01302-1.

[5]   Chu X, Tian W, Ning J, Xiao G, Zhou Y, Wang Z, et al. Cancer stem cells: advances in knowledge and implications for cancer therapy. Signal Transduct Target Ther 2024;9:1-70. https://doi.org/10.1038/s41392-024-01851-y.

[6]   Roszkowska M. Multilevel Mechanisms of Cancer Drug Resistance. Int J Mol Sci 2024;25. https://doi.org/10.3390/ijms252212402.

[7]   Garg P, Malhotra J, Kulkarni P, Horne D, Salgia R, Singhal SS. Emerging Therapeutic
Strategies to Overcome Drug Resistance in Cancer Cells. Cancers (Basel) 2024;16(13): 1-
25. https://doi.org/10.3390/cancers16132478.

[8]   Afkar A, Afolabi BO, Piyathilake PC, Perera DN, Senathilake K, Wijerathne SK, et al.
Anticancer Therapeutic Potential of Natural Products. Journal of Desk Research Review
and Analysis 2024;2(1):67–91. https://doi.org/10.4038/jdrra.v2i1.29.

[9]   Safe S. Natural products as anticancer agents and enhancing their efficacy by a mechanism-
based precision approach. Exploration of Drug Science 2024:408–27.
https://doi.org/10.37349/eds.2024.00054.

[10] Naeem A, Hu P, Yang M, Zhang J, Liu Y, Zhu W, et al. Natural Products as Anticancer
Agents: Current Status and Future Perspectives. Molecules 2022;27.
https://doi.org/10.3390/molecules27238367.

[11] Shaik BB, Katari NK, Jonnalagadda SB. Role of Natural Products in Developing Novel
Anticancer Agents: A Perspective. Chem Biodivers 2022;19(11):1-13.
https://doi.org/10.1002/cbdv.202200535.

[12] Asma ST, Acaroz U, Imre K, Morar A, Shah SRA, Hussain SZ, et al. Natural
Products/Bioactive Compounds as a Source of Anticancer Drugs. Cancers (Basel) 2022;14.
https://doi.org/10.3390/cancers14246203.

[13] Saliu TP, Seneviratne NN, Faizan M, Rajagopalan U, Perera DC, Adhikari A, et al. In silico
identification and in vitro validation of alpha-hederin as a potent inhibitor of Wnt/β-catenin
signaling pathway in breast cancer stem cells. In Silico Pharmacol 2024;12(31):1-15.
https://doi.org/10.1007/s40203-024-00199-z.

[14] Tharmarajah L, Samarakoon SR, Ediriweera MK, Piyathilaka P, Tennekoon KH,
Senathilake KS, et al. In Vitro Anticancer Effect of Gedunin on Human Teratocarcinomal
(NTERA-2) Cancer Stem-Like Cells. Biomed Res Int 2017;2017(1):1-9.
https://doi.org/10.1155/2017/2413197.

[15] Abeysinghe NK, Thabrew I, Samarakoon SR, Ediriweera MK, Tennekoon KH, Pathiranage
VPC, et al. Vernolactone Promotes Apoptosis and Autophagy in Human Teratocarcinomal
(NTERA-2) Cancer Stem-Like Cells. Stem Cells Int 2019;2019.
https://doi.org/10.1155/2019/6907893.

[16] Borlongan MC, Rodriguez T, Putthanbut N, Wang H, Lee JY. Modeling of cancer stem cells
and the tumor microenvironment Via NT2/D1 cells to probe pathology and treatment for
cancer and beyond. Discover Oncology 2025;16. https://doi.org/10.1007/s12672-025-
02158-2.

[17] Anand U, Dey A, Chandel AKS, Sanyal R, Mishra A, Pandey DK, et al. Cancer
chemotherapy and beyond: Current status, drug candidates, associated risks and progress in
targeted therapeutics. Genes Dis 2023;10:1367–401.
https://doi.org/10.1016/j.gendis.2022.02.007.

[18] Ali Abdalla YO, Subramaniam B, Nyamathulla S, Shamsuddin N, Arshad NM, Mun KS, et
al. Natural Products for Cancer Therapy: A Review of Their Mechanism of Actions and
Toxicity in the Past Decade. J Trop Med 2022;2022. https://doi.org/10.1155/2022/5794350.

[19] Guo M, Jin J, Zhao D, Rong Z, Cao LQ, Li AH, et al. Research Advances on Anti-Cancer
Natural Products. Front Oncol 2022;12. https://doi.org/10.3389/fonc.2022.866154.

[20] Atanasov AG, Zotchev SB, Dirsch VM, Orhan IE, Banach M, Rollinger JM, et al. Natural
products in drug discovery: advances and opportunities. Nat Rev Drug Discov
2021;20:200–16. https://doi.org/10.1038/s41573-020-00114-z.

[21] Yun BW, Yan Z, Amir R, Hong S, Jin YW, Lee EK, et al. Plant natural products: History,
limitations and the potential of cambial meristematic cells. Biotechnol Genet Eng Rev
2012;28:47–60. https://doi.org/10.5661/bger-28-47.

[22] Mendis AS, Thabrew I, Ediriweera MK, Samarakoon SR, Tennekoon KH, Adhikari A, et
al. Isolation of a New Sesquiterpene Lactone From Vernonia Zeylanica (L) Less and its
Anti-Proliferative Effects in Breast Cancer Cell Lines. Anticancer Agents Med Chem
2018;19:410–24. https://doi.org/10.2174/1871520619666181128163359.

[23] Samarakoon SR, Thabrew I, Galhena P, De Silva D, Tennekoon K. A comparison of the
cytotoxic potential of standardized aqueous and ethanolic extracts of a polyherbal mixture
comprised of Nigella sativa (seeds), Hemidesmus indicus (roots) and Smilax glabra
(rhizome). Pharmacognosy Res 2010;2:335–42. https://doi.org/10.4103/0974-8490.75451.

[24] Samarakoon SR, Thabrew I, Galhena P, De Silva D, Tennekoon K. A comparison of the
cytotoxic potential of standardized aqueous and ethanolic extracts of a polyherbal mixture
comprised of Nigella sativa (seeds), Hemidesmus indicus (roots) and Smilax glabra
(rhizome). Pharmacognosy Res 2010;2:335–42. https://doi.org/10.4103/0974-8490.75451.

[25] Galhena PB, Samarakoon SR, Thabrew MI, Weerasinghe GAK, Thammitiyagodage MG,
Ratnasooriya WD, et al. Anti-inflammatory activity is a possible mechanism by which the
polyherbal formulation comprised of Nigella sativa (seeds), Hemidesmus indicus (root), and
Smilax glabra (rhizome) mediates its antihepatocarcinogenic effects. Evidence-Based
Complementary and Alternative Medicine 2012;2012.
https://doi.org/10.1155/2012/108626.

[26] Pathiranage VC, Thabrew I, Samarakoon SR, Tennekoon KH, Rajagopalan U, Ediriweera
MK. Evaluation of anticancer effects of a pharmaceutically viable extract of a traditional
polyherbal mixture against non-small-cell lung cancer cells. J Integr Med 2020;18:242–52.
https://doi.org/10.1016/j.joim.2020.02.007.

[27] Daulagala PWHKP. The Journal of the Society for Tropical Plant Research Medicinally
important Leucas zeylanica (L.) R. Br.: A review 2021.
https://doi.org/10.22271/tpr.2021.v8.i2.015.

[28] Wu H, Wang Y, Zhang B, Li YL, Ren ZX, Huang JJ, et al. Smilax glabra Roxb.: A Review
of Its Traditional Usages, Phytochemical Constituents, Pharmacological Properties, and

Clinical Applications. Drug Des Devel Ther 2022;16:3621–43.
https://doi.org/10.2147/DDDT.S374439.

[29] Ratnasooriya WD, Deraniyagala SA, Peiris SKJS. Antinociceptive potential of the Sri
Lankan endemic plant Vernonia zeylanica. Pharm Biol 2007;45:525–32.
https://doi.org/10.1080/13880200701215042.

[30] Rukshala D, de Silva ED, Ranaweera BVLR, Fernando N, Handunnetti SM. Anti-
inflammatory effect of leaves of Vernonia zeylanica in lipopolysaccharide-stimulated RAW
264.7 macrophages and carrageenan-induced rat paw-edema model. J Ethnopharmacol
2021;274. https://doi.org/10.1016/j.jep.2021.114030.

[31] Galhena PB, Samarakoon SR, Thabrew MI, Weerasinghe GAK, Thammitiyagodage MG,
Ratnasooriya WD, et al. Anti-inflammatory activity is a possible mechanism by which the
polyherbal formulation comprised of Nigella sativa (seeds), Hemidesmus indicus (root), and
Smilax glabra (rhizome) mediates its antihepatocarcinogenic effects. Evidence-Based
Complementary and Alternative Medicine 2012;2012.
https://doi.org/10.1155/2012/108626.

[32] Samarakoon SR, Thabrew I, Galhena PB, Tennekoon KH. Modulation of apoptosis in
human hepatocellular carcinoma (HepG2 cells) by a standardized herbal decoction of
Nigella sativa seeds, Hemidesmus indicus roots and Smilax glabra rhizomes with anti-
hepatocarcinogenic effects. BMC Complement Altern Med 2012;12.
https://doi.org/10.1186/1472-6882-12-25.

[33] Pathiranage VC, Thabrew I, Samarakoon SR, Tennekoon KH, Rajagopalan U, Ediriweera
MK. Evaluation of anticancer effects of a pharmaceutically viable extract of a traditional
polyherbal mixture against non-small-cell lung cancer cells. J Integr Med 2020;18:242–52.
https://doi.org/10.1016/j.joim.2020.02.007.

[34] Samarakoon SR, Thabrew I, Galhena PB, Tennekoon KH. Effect of standardized decoction
of Nigella sativa seed, Hemidesmus indicus root and smilax glabra rhizome on the
expression of p53 and p21 genes in human hepatoma cells (HepG2) and mouse liver with
chemically-induced hepatocarcinogenesis. Tropical Journal of Pharmaceutical Research
2012;11:51–61. https://doi.org/10.4314/tjpr.v11i1.7.

[35] Rajagopalan U, Samarakoon SR, Tennekoon KH, Malavige N, Silva ED. Screening of five
sri lankan endemic plants for anti-cancer effects on breast cancer stem cells isolated from
mcf-7 and mda-mb-231 cell lines. Tropical Journal of Pharmaceutical Research
2018;17:1825–32. https://doi.org/10.4314/tjpr.v17i9.21.

[36] Ediriweera MK, Tennekoon KH, Adhikari A, Samarakoon SR, Thabrew I, De Silva ED.
New halogenated constituents from Mangifera zeylanica Hook.f. and their potential anti-
cancer effects in breast and ovarian cancer cells. J Ethnopharmacol 2016;189:165–74.
https://doi.org/10.1016/j.jep.2016.05.047.

[37] Ediriweera MK, Tennekoon KH, Samarakoon SR, Thabrew I, De Silva ED. Induction of
Apoptosis in MCF-7 Breast Cancer Cells by Sri Lankan Endemic Mango (Mangifera
zeylanica) Fruit Peel through Oxidative Stress and Analysis of its Phytochemical
Constituents. J Food Biochem 2017;41. https://doi.org/10.1111/jfbc.12294.

[38] Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time
quantitative PCR and the 2-ΔΔCT method. Methods 2001;25:402–8.
https://doi.org/10.1006/meth.2001.1262.

[39] Sakthi Saravanan V, Selvam P, Parthiban P, Chandra Bose Ps, Paul Raj S. Anticancer activity
of Polyherbal formulation anti cancer activity of polyherbal formulation. International
journal of pharmacy and pharmaceutical analysis 2015;02(03):01-05.

[40] Bhatt D, Vaghela K, Jethva K, Patel S, Zaveri M. Development and standardization of
Polyherbal formulation for the management of breast cancer. International Journal of
Pharmaceutical Science and Research 2017;2:2455–4685.

[41] Petronek MS, Stolwijk JM, Murray SD, Steinbach EJ, Zakharia Y, Buettner GR, et al.
Utilization of redox modulating small molecules that selectively act as pro-oxidants in
cancer cells to open a therapeutic window for improving cancer therapy. Redox Biol
2021;42. https://doi.org/10.1016/j.redox.2021.101864.

[42] Grada A, Otero-Vinas M, Prieto-Castrillo F, Obagi Z, Falanga V. Research Techniques Made
Simple: Analysis of Collective Cell Migration Using the Wound Healing Assay. Journal of
Investigative Dermatology 2017;137:11–6. https://doi.org/10.1016/j.jid.2016.11.020.

[43] Brábek J, Mierke CT, Rösel D, Veselý P, Fabry B. The role of the tissue microenvironment
in the regulation of cancer cell motility and invasion. 2010.

[44] Abdollahzadeh N, Tafrihi M. Exploring in vitro anticancer potential of Ferula gummosa gum
on PC-3 cells: insights from cellular and molecular studies. Egyptian Journal of Basic and
Applied Sciences. 2025;12(1): 86-93. doi: 10.1080/2314808X.2024.2448403.

[45] Ndreshkjana B, Çapci A, Klein V, Chanvorachote P, Muenzner JK, Huebner K, et al.
Combination of 5-fluorouracil and thymoquinone targets stem cell gene signature in
colorectal cancer cells. Cell Death Dis 2019;2019;379(10): https://doi.org/10.1038/s41419-
019-1611-4.