1

European ash pangenome reveals widespread structural variation and genetic basis of low ash dieback susceptibility

Authors: Daniel P. Wood[1], Mohammad Vatanparast[1], Dario Galanti[1], Catherine Gudgeon[1], Katherine Wheeler[1,2], Emma Curran[3], Levi Yant[3,4], Richard Whittet[5], Richard A. Nichols[6], Richard J. A. Buggs[1,6], Laura J. Kelly[1]

1. Royal Botanic Gardens, Kew, Richmond, UK
2. Centre for Novel Agricultural Products, University of York, York, UK
3. School of Life Sciences, University of Nottingham, Nottingham, UK
4. Linnean Centre for Plant Biology, Department of Plant Biology, Uppsala BioCenter, Swedish University of Agricultural Sciences, Uppsala, Sweden
5. Forest Research, Northern Research Station, Roslin, Midlothian, UK
6. School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK

Email: d.wood@kew.org; l.kelly@kew.org

Abstract

European Ash (*Fraxinus excelsior*) is a keystone forest tree species, whose populations are being decimated by ash dieback disease (ADB). Uncovering the genetic basis of low susceptibility to this devastating disease relies on a comprehensive understanding of genetic variation present in *F. excelsior.* A linear reference genome from a single individual cannot contain the total sequence variability within a species, including its genic regions; a pangenome more fully captures total sequence content. In this study, we developed a *F. excelsior* pangenome reference using a new chromosomal-level

phased linear reference genome, and *de novo* assemblies and long-read data from a

geographically diverse set of fifty *F. excelsior* samples, with particular focus on

individuals showing low ADB susceptibility. We identified 362,965 structural variants

(SVs) present in more than three individuals, including 174Mb of sequence absent from

the linear reference genome (22% of the linear reference size). We demonstrated that

failing to explicitly link SVs with gene sequences can lead to substantial overestimation

of dispensable genes (those that vary in their presence/absence between individuals)

due to variability in the annotation process. Controlling for this reduced the fraction of

the genome estimated as dispensable from 35.9% to 8.7%, identifying 3,412 high-

confidence dispensable genes, including 141 annotated with gene ontology terms

associated with defence response. We used the pangenome to analyse existing

genomic data from over 1,200 individuals to identify loci associated with reduced

susceptibility to ADB. This revealed 220 single nucleotide polymorphisms (SNPs)

showing allele frequency shifts between healthy and highly damaged pools of

individuals that are broadly consistent across the mainly UK seed sources sampled,

explicitly demonstrating the existence of a shared genetic component to low ADB

susceptibility.

Introduction

High quality reference genomes constructed from single individuals have been used to

great effect to assess genetic diversity within species, allowing mapping of short reads

from many individuals and discovery of genetic variants[1]. However, this approach

provides limited insights into larger structural variants (SVs) between individuals and

can lead to some miscalling of single nucleotide polymorphisms (SNPs)[2]. Pangenomes

– reference genome data structures incorporating highly contiguous sequences from a

representative set of individuals – have revealed the total sequence contained in SVs

can exceed the average genome size of a species[3–6]. These SVs frequently encompass

genes, with some studies identifying >50% of genes within a species as dispensable

(i.e. not present in all individuals[5,7–9]). Pangenomes thus often contain genes that would

be entirely missed if using a single reference genome approach[10,11]. Indeed, novel

disease resistance genes are frequently identified in pangenomes[12–14] including in

trees[15–17], suggesting pangenome approaches may be particularly valuable when

seeking to detect loci contributing to pathogen resistance. Analysis of existing data

using pangenome references has uncovered the basis of missing heritability in some

systems[18] and has enhanced our understanding of the genetic basis of a variety of

traits, often adding complementary information to that uncovered by analyses using a

linear reference genome[5,19–21]. Thus, to better understand genetic variability within a

species and its impact on traits, pangenome references incorporating diverse samples

are required[10,22].

Fraxinus excelsior (European ash), a keystone tree species within Europe, is under

severe threat from ash dieback disease (ADB) caused by the invasive non-native

fungus *Hymenoscyphus fraxineus*[23,24]. Populations of ash are undergoing substantial

and rapid mortality[25–27], with high associated management costs and biodiversity loss[28–

30], attracting substantial public concern[31–34]. A small fraction of trees (as low as 0.5%[35])

remain healthy under *H. fraxineus* inoculum pressure, with quantitative resistance to

ADB having a narrow-sense heritability of approximately 0.4[35–39] . Understanding the

genetic basis of this trait is important for ascertaining the feasibility of producing new

individuals with lower susceptibility[40–44]. Stocks et al. (2019)[42] identified 192 SNPs

significantly associated with low ADB susceptibility when using a p-value threshold of 1

x 10^{-13} in mainly UK populations by mapping short reads to a single reference genome.

Studies in other European populations did not find SNPs strongly associated with low

ADB susceptibility[43,44], with one study indicating many of the most significant from

Stocks et al.[42] were at very low frequencies in Austrian populations[45], raising the

possibility of diverse mechanisms of low susceptibility between populations in different

parts of the species' range[42–47]. A pangenome reference for *F. excelsior* that captures

variation from different genetic backgrounds would represent an invaluable resource to

enable improved understanding of this critically important trait. A recently published

"superpangenome" of the *Fraxinus* genus identifies interspecific SVs associated with

salt tolerance, suggesting a role for these variants in adaptation to abiotic stressors over

millions of years[48]. However, this study does not capture intraspecific SVs in *F.*

excelsior that may contribute to low ADB susceptibility.

Here we construct a chromosome level, phased genome assembly for European ash

and a graph pangenome using long-read sequencing from a diverse sample of 50

individuals across Europe. This revealed 174Mb of additional sequence captured in

SVs, an increase of 22% from the new linear reference genome size from a single

individual. By explicitly associating putative dispensable genes with SVs, we identified

3,412 high confidence dispensable genes in *F. excelsior*, including 141 with gene

ontology terms associated with defense response. Reanalysis of poolseq data (i.e.

sequence data from a mixture of approximately equal amounts of DNA from multiple

individuals) from over 1,200 individuals[42] revealed 220 SNPs with broadly consistent

allele frequency shifts between unhealthy and unhealthy pairs of pools across a set of

mainly UK seed sources, providing explicit evidence for a shared genetic component to

low ADB susceptibility.

100   Results

101   A high quality linear reference genome for European ash

A phased linear reference genome was produced for a single *F. excelsior* individual,

with total sizes of 793 and 787 Mb, N50s of 32.9 and 32.2 Mb, and L90s of 22 for the

haplotype 1 (henceforth referred to as "BATG-1.0") and haplotype 2 assemblies

respectively (Supplementary Table 1). *Fraxinus excelsior* has 23 pairs of

chromosomes[49] – the top 23 scaffolds of BATG-1.0 and haplotype 2 contain 93.7% and

94.5% of the total sequence in each assembly, with the 24th scaffold for each of the

haplotypes being 2.1% and 1.5% of the size of the 23rd scaffold. Three short scaffolds

were identified as potential contaminants (with lengths between 14,626 and 35,091bp),

representing 0.009% of the total assembly size - these were removed from downstream

analysis. Telomeres were detected at both ends of 19 of the top 23 scaffolds in BATG-

1.0 and 17 of the top 23 in haplotype 2**.** Together these results indicate the assemblies

are pseudochromosomal. Unlike other recently published pseudochromosomal

assemblies of *F. excelsior*[48,50], this assembly is not scaffolded using the *F.*

pennsylvanica genome assembly[51]. We also assembled and annotated the *F. excelsior*

chloroplast genome; this was 155.6kb in length with 130 annotated genes.

98.2% and 98.3% of eudicot BUSCOs were completely assembled in the BATG-1.0 and

haplotype 2 genomes; 21% of BUSCOs were duplicated, a similar proportion as in other

F. excelsior assemblies (Supplementary Table 1) and consistent with the fact that there

are several whole genome duplication events in the Oleaceae, the family to which *F.*

excelsior belongs[48,52]. 98.3% of PacBio HiFi reads mapped to one or both of the haploid

assemblies. The total assembly sizes of 787Mb and 793Mb are smaller than the 1C

genome sizes of 877Mb and 840Mb estimated by flow cytometry in Sollars et al.[53]

(2017) and Siljak-Yakovlev et al. (2014)[54] respectively - different estimates could result

from methodological variation or inter-individual genome size differences. Analysing

27bp kmers from the PacBio reads using GenomeScope[55] gave an estimated genome

size range of 823-825.4Mb for this individual (Supplementary Figure 1) - other kmer

lengths tested (see Methods) gave very similar size estimates. The coverage of read

mapping to both phased assemblies did not show clear peaks of higher depth beyond

the main homozygous peak, which would be indicative of collapsed sequences

(Supplementary Figure 2). This suggests the remaining difference between estimated

genome sizes and assembly length may be due to highly repetitive regions such as

centromeres and ribosomal DNA that are challenging to assemble. The assembly sizes

are similar to the size of unpublished pseudochromosomal assemblies of *F. excelsior*

that do not use *F. pennsylvanica* for scaffolding (i.e. FRAX_001_PL and

daFraExce3.hap1.1; see Supplementary Table 1). The LTR Assembly Index (LAI) was

14.85 and 14.62 for BATG-1.0 and haplotype 2 respectively, with 5.35-5.44 intact

LTRs/Mb (comparable to other recent *Fraxinus* assemblies), with the top 23 scaffolds in

each having average LAIs of 20.74 and 20.62 respectively (Supplementary Table 1) - a

score of >10 indicates a reference standard genome[56].

65.6% of BATG-1.0 consists of repeats. LTR elements occupy 33.0% of the sequence,

most of which comprises Ty1/Copia (15.3% of genome sequence) or Gypsy/DIRS1

(15.9% of genome sequence) families, with DNA transposons covering 6.2% of the

sequence. Unclassified repeats covered 19.9% of the sequence. The total repetitive

fraction is similar to the estimated repeat content of the assembly of another *F. excelsior*

individual (69.12%) and assemblies for some other *Fraxinus* species such as *F.*

pennsylvanica from the *Fraxinus* superpangenome[48], but is substantially higher than a

separate estimate for *F. pennsylvanica* (48.4%[51]). Olive, also a member of the

Oleaceae, has a lower repeat content than *F. excelsior* (56.2%)[57], although tomato (a

fellow asterid with a similar 1C genome size) is more similar (65%-72%)[58,59].

Annotation of BATG-1.0 was performed using BRAKER3[60]. Ten runs of BRAKER3 were

performed with short-read RNA-seq data and protein orthologue information, and 10

runs with long-read RNA-seq data and protein orthologues. These identified 31,926

genes annotated in every run, with a further 1,736 genes annotated in every run of the

short-read annotation only and 1,246 genes annotated in every run of the long-read

annotation only. The median gene length of genes exclusively identified by long-reads

was 43% longer than those identified only by short reads (Supplementary Figure 3),

although different underlying tissue types were used to generate the long-read and

short-read RNA-seq data, making these differences challenging to explicitly link to the

length of RNA-seq input data. A further 3,075 genes were inconsistently annotated, in

either the short- or long-read runs. To reduce the chances of erroneously identifying

genes as dispensable in the pangenome, we initially retained all these genes, giving a

total of 37,983. This is a similar number to that reported in the BATG-0.5 assembly

(38,949)[53] and the *F. pennsylvanica* assembly (35,470)[51] but lower than the 43,392

genes reported in the recent Fexc_ONT_UoY *F. excelsior* assembly[50] and substantially

lower than the *Fraxinus* assemblies used in the *Fraxinus* superpangenome (48,389-

53,538)[48] including their *F. excelsior* assembly (50,327). These latter two *F. excelsior*

assemblies have more genes annotated than are plausibly missing from BATG-1.0 due

to being dispensable (see pangenome annotation below), suggesting differences are

likely due to different annotation approaches rather than intraspecific variation in gene

content. Using OrthoFinder[61] to identify orthologues from BATG-1.0 and available

annotations for *F. excelsior* assemblies (BATG-0.5, Fexc_ONT_UoY) as well as *F.*

pennsylvanica, *Olea europaea*, *Solanum lycopersicum*, *Erythranthe guttata*, *Sesamum*

indicum and *Salvia miltiorrhiza* (Supplementary Table 2), 0.6% of BATG-1.0 genes

lacked an orthologue in another assembly, compared with 1.1% in BATG-0.5 and 7.3%

in Fexc_ONT_UoY. 98.7% of eudicot BUSCO proteins were found in the total BATG-1.0

protein set, suggesting the annotation captures the vast majority of genes in the

assembly.

Structural variant calling and pangenome construction

Low susceptibility to ADB is likely a highly polygenic trait[42,47], so we prioritised sampling

a larger number of individuals to construct the pangenome instead of focusing on

generating fewer highly contiguous assemblies, even if this restricts the types of SV that

can be analysed. Oxford Nanopore Technologies (ONT) data was generated for 49

individuals from a variety of European provenances, as well as for the reference

individual (Figure 1A, Supplementary Table 3). PCA of SNPs called from the 49

individuals revealed little geographic structure (Figure 1B), as was also found to be the

case when a similar set of provenances were analysed in earlier work[53].

Heterozygous structural variants called between the two high quality haploid genomes

of the reference individual were used to assess the accuracy of calling SVs using lower

coverage ONT data from the same individual with a variety of methods. These included

mapping the ONT reads to the reference haploid genome and calling SVs using

Sniffles2[62] and cuteSV[63], *de novo* assembling the ONT reads with Flye[64], Shasta[65] or

nextDenovo[66] and calling SVs from contigs mapped against the reference using swim-

asm[67], and the combination of read mapping and assembly mapping approaches (see

Methods). This analysis indicated that the read mapping approaches had higher recall

of SVs detected using the high quality haploid genomes, while assembly mapping

approaches had higher precision, and a combination of read mapping and mapping of

the assembly produced by Shasta gave the highest recall (84.4%) and F1 score

(77.9%), with a precision of 72.2% (Extended Data Figure 1). For insertions, deletions

and tandem duplications, this method also gave the highest F1; for inversions, using

swim-asm from a Flye *de novo* assembly gave a higher F1, although the overall

precision/recall for inversion and tandem repeats was low (Supplementary Figure

4).The mapping and/or Shasta assembly based approach was therefore used to call

SVs for all 50 samples. Within each sample, a median of 41.6% of SVs called with

cuteSV and Sniffles2 were also called with swim-asm, and a median of 72.6% of SV

called with svim-asm were also called with cuteSV and Sniffles2 (Supplementary Figure

5).

Merging SVs across all individuals identified a total of 737,995 SVs (Figure 2A).

Merging samples sequentially resulted in fewer new SVs discovered in the fiftieth

sample compared with the first, but the number newly detected did not plateau,

indicating that more samples would be required to exhaustively sample SVs in *F.*

excelsior (Figure 2B). It is also possible that sample-specific SVs represent sample-

specific assembly or mapping errors - to attempt to control for this, only SVs present in

more than three individuals were retained for further analysis (Figure 2A), although this

may result in rare SVs being missed. This resulted in 362,965 SVs, of which 170,202

were insertions, 189,724 were deletions, 295 were inversions and 2,744 were tandem

duplications (Figure 2A; Supplementary Figure 6). The total number of SVs represented

396Mb of variable sequence across the *F. excelsior* individuals, including 174Mb of

insertion sequences not present in the linear reference genome (Figure 2C). 99.9% of

SVs were on the 23 pseudochromosomal scaffolds. Insertions and deletions were

generally more frequent towards the ends of chromosome arms (Figure 2D;

Supplementary File 1). 69% of SVs overlapped with an annotated repeat in the

reference genome; the distribution of base pairs overlapping different SV types closely

reflected the makeup of the genome as a whole (i.e. mostly LTRs and DNA

transposons), although rRNAs and tRNAs were relatively unlikely to overlap segregating

SVs, likely due to their core role in the cell machinery (Supplementary Figure 7) .The set

of 362,965 SVs was used to construct a pangenome using vg[68] which had 30,710,268

nodes and 31,069,469 edges.

Pangenome annotation: explicitly linking dispensable genes with SVs

To identify dispensable genes with an explicit link to SVs in the pangenome, a genome sequence was generated for each individual by using the SV genotype calls to modify the BATG-1.0 reference fasta (i.e. deleting deletions, inserting insertion sequence, etc). Genes were *de novo* annotated for each individual genome sequence and consolidated across individuals using OrthoFinder to identify a set of unique gene models. Of the 48,356 genes, 30,992 (64.1%) were present in all individuals including the reference (Figure 3). Of the 10,411 genes absent from the BATG-1.0 genome (i.e. putatively undiscovered dispensable genes), 6,009 did not overlap with an SV in any individual. For those that did overlap with SVs, only 1,474 did so with an F1 score (a measure of how consistently a gene presence or absence was associated with a SV, see Methods) of 0.8 or above (Figure 3, Extended Data Figure 2). As in each case the non-SV sequence is identical to the reference sequence, these putatively dispensable genes may be the result of stochasticity in the annotation process, or the result of SVs elsewhere in the genome altering the mapping of RNA-seq data and thereby reducing evidence to support the gene annotation. Similarly, of the 6,953 genes present in the reference assembly but absent from one or more of the other assemblies, 3,190 did not overlap with a SV in any individual. For those that did overlap with a SV, only 1,938 did so with an F1 score of 0.8 or above (Figure 3, Extended Data Figure 2). Putatively dispensable genes with an F1 score of less than 0.8 were classified as invariant if present in the reference genome, and if absent were removed from the analysis. This

brought the total number of genes in the pangenome to 39,419, of which 3,412 were

dispensable (8.7%% of the total). This is a substantial reduction in dispensable genes

compared with the estimated 35.9% of genes without filtering for those consistently

overlapping with SVs (Figure 3). Putatively dispensable genes without a consistent SV

association (i.e. those excluded from our final annotation) could be affected by the non-

comprehensive RNA-seq dataset (only 72% of individuals in the pangenome were

represented in the RNA-seq data) or the particular annotation pipeline used. The final

pangenome gene set included 99.0% complete eudicot BUSCOs – slightly more than

the 98.7% captured in the BATG-1.0 protein sequences alone.

Functional annotation indicated that high confidence dispensable genes overall had

fewer annotations than indispensable genes; putatively dispensable genes that were

filtered out also had a similar reduction in functional annotations compared with

indispensable genes (Figure 4A). This may be a result of these gene subsets being less

conserved across species and/or less well annotated. High-confidence dispensable

genes were enriched for eight GO terms including "RNA-templated DNA biosynthetic

process", "retrotransposition" and "floral meristem growth" (Supplementary Table 4).

Putatively dispensable genes that were filtered out were enriched for 28 terms including

"defense response" and "regulation of auxin polar transport" that were not significant in

the high-confidence dispensable genes (Supplementary Table 4). However, a direct

comparison of the high confidence dispensable genes with the filtered genes provided

little evidence of GO terms only enriched in the dispensable genes, or the filtered

genes, differing meaningfully in frequency between these two sets (Figure 4B). The

3,412 dispensable genes include those with GO terms indicating a role in response to

stress (355), developmental process (341) and defence response (141), with 26 genes

linked to a function in "defense response to fungus" specifically (Supplementary Table

4). Dispensable genes with a "defense response" GO term belonged to a variety of

protein superfamilies (Supplementary Table 5), including cytochrome P450s (three

genes) and flagellin-sensing 2 (FLS2)-related serine/threonine protein kinases (two),

with Nucleotide Binding Site, Leucine Rich Repeat (NBS-LLR) superfamily being the

most frequently annotated (eight genes). Genes annotated for functions in leaf

development (42) and leaf senescence (16) were also included within the dispensable

set – variation in Spring leaf flushing and Autumn leaf senescence has been linked to

low ADB susceptibility in some studies[38,44,69,70].

Read mapping, SNP calling and SV calling using the pangenome

We next tested whether using our pangenome reference improved read mapping for

short-read sequenced individuals, compared to the BATG-1.0 linear reference

sequence. While overall mapping to our pangenome produced little effect on the total

number of short reads mapped in proper pairs (Extended Data Figure 3), we observed

general improvements based on other metrics: this includes increases in mean mapping

quality and alignment score (1.4% and 1.2% on average, respectively Extended Data

Figure 3), a reduction in the total number of substitutions (3.0% on average, Extended

Data Figure 3), and a large reduction in soft clipped sequence (33.5% on average,

Extended Data Figure 3). Overall, this indicates the pangenome improves the accuracy

of short-read read mapping. Mapped reads in each case were used to call SNPs -

overall numbers of SNPs and quality metrics were similar using both approaches; only

5.6% and 3.4% of SNPs called in the reference-based and pangenome approaches

respectively were not called with the other approach. The quality of SNPs that were not

detected with both approaches appeared similar to the general background set of SNPs

(Supplementary Figures 8,9; Supplementary Table 6). The median distance of unique

SNPs from one approach to SNPs from the other approach was <30bp for both callsets,

although the reference-based callset included some SNPs >1kb from the nearest SNP

in the pangenome-based callsets (Supplementary Figure 9). This indicates that while

the improved overall mapping to the pangenome results in some SNP-calling

differences, the overall impact on SNP-based analysis is likely to be minimal compared

to that associated with using a high-quality linear reference genome.

Examination of allelic depths of homozygous alternate, reference and heterozygous

sites showed SVs and SNPs exhibit similar distributions of read support for reference

and alternate alleles (Extended Data Figure 4). The minor allele frequency (MAF)

distributions and deviation from Hardy-Weinburg equilibrium (HWE) for the two variant

types were also similar (Extended Data Figure 5A, B). Patterns of linkage decay with

distance were broadly similar in SNP-SNP, SV-SV and SNP-SV comparisons, with

slightly faster linkage decay between SVs, and between SVs and SNPs, compared with

between SNPs (Extended Data Figure 5C). Together, these suggest SNP and SV calls

behave in a similar way for individually sequenced samples.

To investigate the accuracy of allele frequency estimations from short-read poolseq

data, we constructed an *in silico* pool with an equivalent number of individuals and

depth to the average *in vitro* pool from Stocks et al.[42]. Allele frequencies for SNPs were

estimated using Popoolation2[71]. Similar software does not exist for estimating SV allele

frequencies in poolseq data using a pangenome. Whilst vg[68] does not explicitly support

poolseq allele frequency estimation, we called SVs for the pools using vg and used the

depths of reference vs. alternate alleles to estimate SV frequency in the pools.

Correlation between the alternate allele frequencies from the individual based calling

and estimates from the *in silico* pool was high for SNPs (slope = 0.94, R^2 = 0.93). For

SVs, whilst the overall correlation was also high (slope = 0.96, R^2 = 0.85), there were a

minority of SVs where the pool-based estimates differed substantially from the individual

based calling (Extended Data Figure 6A, B). Poolseq data from Stocks et al.[42] was

mapped to the pangenome, and SNP and SV frequencies analysed. PCA recovered

similar population structures from SV and SNP frequencies estimated using these pools

(Extended Data Figure 6C, D) as has been found in other studies[68,72]. Together, these

results suggest that allele frequencies of most SVs can be captured accurately by

mapping poolseq data to a pangenome.

GWAS with SVs and SNPs

We reanalysed poolseq data from Stocks et al. (2019)[42] using the pangenome. In that

study, for 13 seed sources, a pool of healthy trees (no visible signs of ADB damage)

and a pool of unhealthy trees (for which the majority of individuals had a dead/partially

dead main stem; 42 trees per pool on average) from field trials heavily affected by ADB

were sequenced (see Methods for details). We performed a GWAS for low ADB

susceptibility using SV allele frequencies estimated from this data, with a Cochran-

Mantel Haenszel (CMH) test[73,74]. This test identifies associations of treatments and

outcomes while taking into account stratification - it is a commonly used statistic for

identifying associations from paired poolseq data[42,71,75–79]. This identified 531 significant

SVs with a p-value cutoff of 1 x 10^{-13} (the same significance threshold as used for the

CMH test in Stocks et al.[42]; Extended Data Figure 7A). However, many of these sites

showed inconsistent allele frequency estimates from the *in silico* pools vs. individual

genotyping as assessed in the above section (261 had a difference in estimated

frequency of > 0.2; see Extended Data Figure 7B vs. Extended Data Figure 6B). Whilst

some of the 531 significant SVs may be truly associated with low ADB susceptibility,

many are likely false positives resulting from errors in the estimation of allele

frequencies. These results indicate SV allele frequency estimates from poolseq data are

insufficiently accurate for a reliable GWAS.

We next considered the estimated allele frequencies for SNPs called from the Stocks et

al.[42] poolseq data using our pangenome, and used these to perform a separate GWAS

for low susceptibility to ADB using the CMH test. This displayed high p-value inflation

(Supplementary Figure 10A). 866 SNPs were identified as significant after Bonferroni

correction with □ = 0.05 (uncorrected p-value threshold = 1.52 x 10^{-9}), and 404,503 with

5% false discovery rate using the Benjamini-Hochberg procedure. 92 SNPs had a p-

value of less than 1 x 10^{-13} (the significance threshold used for the CMH test in Stocks

et al.). Wiberg et al. (2017)[75] demonstrated that heterogeneity in SNP allele frequency

shifts among pairs of pools can be identified as a main effect, causing p-value inflation

in the CMH test. Deviation from homogeneity, as measured either by a significant Woolf

test[80] p-value (p < 0.05: 53.4% of significant SNPs), or by an I^2 statistic > 0.25 (73.2% of

SNPs)[81], was common, and was particularly pronounced for SNPs with the lowest CMH

test p-values (Supplementary Figure 11). Indeed, many of the SNPs with the lowest p-

values (less than 1×10^{-13}) had high read depth variation between pools (Extended Data

Figure 8), consistent with the presence of rare SVs distorting estimates of allele

frequency (see Supplementary Note 1 for further discussion).

A quasibinomial generalized linear model (qGLM) test - a GLM for proportion/binomial

data including an overdispersion parameter - has been shown to be better able to

identify consistent allele frequency shifts across heterogeneous pairs of pools[75]. This

analysis did not identify any significant sites, although substantial p-value deflation

suggests this test may be overly conservative (Supplementary Figure 12A). Indeed, the

lowest p-values displayed consistent allele frequency shifts, but with shifts in allele

frequency between healthy and unhealthy pools similar to random subsets of SNPs

(Supplementary Note 2, Supplementary Figure 12B-F). Different approaches to

modelling the effect of provenance using GLMs either led to a similar lack of significant

SNPs, or substantial p-value inflation (Supplementary Note 2, Supplementary Figure

13). This indicates limited evidence for SNPs behaving in a highly consistent manner

across pools. We attempted to explicitly assess heterogeneity in allele frequency

changes across pairs of pools using the Fixation index (F_{ST}) or Fisher's exact test, and

considering sites where multiple pairs of pools showed a strongly supported allele

frequency shift. However, this proved ineffective for identifying sites shared by subsets

of pools (with minimal overlap found even between biological replicates from Stocks et

al.[42]), likely due to the inherent noise in individual allele frequency estimates from

poolseq data (see Supplementary Note 3).

Despite inflated p-values, the 866 SNPs passing CMH test Bonferroni correction

displayed large allele frequency shifts compared to the genome-wide average, also

displaying a consistent direction of change in most pairs of pools (Figure 5), suggesting

they are likely enriched for SNPs associated with ADB resistance in the majority of seed

sources analysed. 220 of these SNPs did not violate the CMH test assumption of

homogeneity, or display high standard deviation in read depth potentially resulting from

SVs not captured in the pangenome (see Methods), suggesting their low p-values are

more likely to represent a true association with tree health status. 34 genes contained

one or more of these SNPs, including two encoding pentatricopeptide-repeat containing

proteins, ribosomal subunit proteins, and transcription and translation factors

(Supplementary Table 5). A further 177 genes were within 10kb of these SNPs

(Supplementary Table 5), of which 16 were dispensable. No GO terms were

significantly enriched in either the 34 genes overlapping the SNPs, or the 211 genes

within 10kb, after correction for multiple testing (Bonferroni correction with $\square$ = 0.05;

uncorrected p-value threshold of 1.53 x 10^{-5}; Supplementary Table 4), with "immune

response-activating cell surface receptor signaling pathway" and "brassinosteroid

metabolic process" among those in the 10 most enriched terms. The set of 211 genes

included nine with a "defence response" GO term annotation. These included i) a gene

belonged to the NBS-LRR superfamily, ii) a dispensable gene homologous to an

Arabidopsis globulin family protein, and iii) a phosphatidylinositol-specific phospholipase

C homologue that directly overlapped the significant SNPs.

Discussion

Here we present a new chromosome-level linear reference genome (BATG-1.0) and a

pangenome for European ash from a geographically diverse sample of individuals,

enriched for those with low susceptibility to ADB. The pangenome of 50 individuals

captures 362,965 SVs, representing 396Mb of variable sequence, including 174Mb that

is absent from the linear reference genome (equivalent to approximately 22% of its

length). This is broadly similar to other recent pangenome studies, although there is

substantial variability across these[3,15,82–95] (Extended Data Figure 9A, 9B;

Supplementary Table 7). Our analyses likely underestimate the total SV diversity in *F.*

excelsior, as each additional individual contributed novel SVs to the collection. A

comparison of SVs called between the chromosome-level haploid assemblies, and ONT

data for the same individual, indicates our SV calling approach has a precision of 72.2%

(although a recall of 84.4%) compared with using chromosome-level phased

assemblies, suggesting some SVs included in the pangenome may be false positives.

However, it should be noted that i) as heterozygous sites only have half the read

support of homozygous alternate sites it is likely the latter can be detected with higher

recall and precision, and ii) the fact that we only included SVs called in more than three

samples in the pangenome potentially reduces the number of false positives, although

at the cost of excluding rare SVs. Our results highlight the substantial diversity of SVs in

European ash, consistent with high SNP genetic diversity[53], but also underline that more

extensive sampling is required to fully capture segregating SVs.

We identified 8.7% of genes in the pangenome as dispensable; this is substantially

lower than most other plant pangenomes, which typically report 20-60% of genes or

gene families as being variable[3,15,82–93,95] (Extended Data Figure 9C, Supplementary

Table 7). As noted above, our pangenome only contains SVs called in more than three

individuals, so potentially misses the rarest dispensable genes. However, most of the

difference in the proportion of genes that are dispensable is explained by our filtering of

genes not reliably associated with SVs; without this filter, a more typical 35.9% of genes

in the pangenome are classified as dispensable (Extended Data Figure 9C,

Supplementary Table 7). Despite leading to a dramatic reduction in dispensable genes,

we believe this filter is justified as variability in the annotation process has previously

been found to underlie putatively dispensable gene sets[92,95]. Indeed, previous

pangenome studies have identified "core" genes as being more highly expressed

[7,84,87,90,96,97]. This relationship could be due to these genes performing critical biological

functions, but also could be the result of more highly expressed genes having more

RNA-seq data evidence and being more accurately annotated as a result. Functional

groups typically found as enriched among dispensable genes in other studies, such as

defense and auxin response[3,7,84,92,94,98] were enriched in our initial set of putatively

dispensable genes, but not following our filtering process. It should however be stressed

that there was no clear functional distinction between the final set of dispensable genes

and those removed during filtering - a smaller set of genes will inherently have lower

power to reliably detect enrichment in GO terms. More comprehensive RNA-seq

sampling from tissues/conditions across individuals, and manual curation of repetitive

elements, may lead to more genes being annotated - including as dispensable - in

future studies. We should also note that we aimed to identify genes specifically within

our characterised SVs; in studies where *de novo* assemblies are explicitly incorporated

(e.g. all-against-all pangenomes and comparative genomics more broadly) specific

testing of DNA sequence differences underlying presence/absence of genes should be

performed, as has been suggested elsewhere[99]. Our results emphasise the importance

of confirming that putatively dispensable genes are driven by underlying sequence

differences, to distinguish them from possible non-biological annotation variability. The

genes we identified as dispensable with high confidence include 141 with homology to

defence-related genes, one of which was within 10kb of the 220 SNPs associated with

low ADB susceptibility, emphasising the importance of pangenomes in understanding

the mechanistic basis of disease resistance traits.

Using a pangenome reference allows improved short-read mapping for *F. excelsior*, as

has been found for other species[11], and we were able to successfully call individual

SNPs and SVs from individually sequenced samples. However, there was not a clear

improvement in SNP calling compared to using the BATG-1.0 linear reference genome.

SV genotyping from short-reads is generally more challenging than SNPs[100–102], and we

are not aware of previous studies calling allele frequencies of poolseq data mapped to a

pangenome graph. Our results indicate poolseq data can be used to estimate allele

frequencies accurately for most SVs (Extended Data Figure 6), but the approach used

generates more discrepancies with individual based genotyping than is the case for

SNPs. SVs showing these discrepancies were particularly likely to be identified as

significant in a GWAS using a CMH test, resulting in many likely false positives. We

were unable to identify the cause of these discrepancies (see

https://github.com/vgteam/vg/issues/4443). We should emphasise that vg is not

designed for analysing poolseq data; the development of dedicated software for

estimating SV allele frequencies from poolseq data, equivalent to those available for

SNPs, is required in order to make full use of poolseq data in the pangenome era.

Nevertheless, some of the SVs identified as significant from the CMH test are potentially

involved in low ADB susceptibility; future studies using individually genotyped data

could be used to confirm this.

Despite broadly accurate SNP allele frequency estimates from poolseq mapped to a

pangenome, our reanalysis of data from Stocks et al (2019)[42] highlighted known

challenges of analysing poolseq data[103]. The commonly used CMH test can assign very

low p-values to heterogenous allele frequency shifts among pairs of pools[75], and

uncharacterised SVs can distort allele frequency estimations. A combination of these

sources of error meant that despite mapping to a pangenome and using relatively

stringent depth filtering, a likely rare, large/high copy number SV produced the lowest p-

values in our dataset using this test (Supplementary Note 1). We find a subset of 220

SNPs with significant p-values from the CMH test that do not violate its assumptions of

heterogeneity, or display unusual patterns of depth, and shift in frequency between

unhealthy and healthy pools in a broadly consistent manner across most seed source

zones (Figure 5). We would note the inflation of p-values from the CMH test makes the

threshold of inclusion in this group (passing Bonferroni correction with □ = 0.05)

somewhat arbitrary. Nevertheless, these results indicate a shared genetic component to

low susceptibility exists across a broad range of provenances. It should be noted that

more densely sampled regions in the Stocks et al.[42] dataset (e.g. Great Britain) may

make SNPs consistently shifting in these regions more likely to be identified. Whilst no

GO terms were significantly enriched in genes near these SNPs, the "brassinosteroid

metabolic process" term was highly ranked - brassinosteroid biosynthesis genes were

enriched near SNPs predictive of autumn leaf yellowing in a previous study, a trait

which has been associated with low ADB susceptibility[44]. Variation in the frequency of

trees with low susceptibility exists between provenances[35,36], suggesting there may be

further geographically restricted genetic variation associated with low ADB

susceptibility, although we are not able to accurately characterise this in our dataset.

Higher coverage per pool, or individual based sequencing, would allow more accurate

characterisation of variants restricted to subsets of provenances.

In Stocks et al (2019)[42], the consistency of allele frequency shifts, and the potentially

confounding effect of SVs, were not explicitly examined. Indeed, the maximum per-pool

depth for sites considered was 37.5X the average coverage, rather than the more

stringent 2.5x in the present study, increasing the potential for SVs to impact allele

frequency estimates. It is likely therefore that the 192 SNPs with p-values lower than 10^{-13}

in Stocks et al. are similarly, or more severely, enriched for problematic sites, as

those in our study. This may explain the lack of correlation of CMH p-values with

estimated effect sizes in genomic prediction (GP) of ADB damage, and the fact that

around two thirds of their top 192 GWAS hits are ranked outside the top 2,500 SNPs by

GP effect size (Stocks et al. 2019). It may also explain a lack of shared genes between

Doonan et al. (2025) and Stocks et al.[44], although we should note that ADB health

scores represent a complex phenotype underpinned by several mechanisms, which

may reduce the chances of identifying similar underpinning regions across studies.

Nevertheless, a GP model trained on the top 10,000 GWAS loci from Stocks et al. was

able to predict tree health status with an accuracy of 67%[42], despite likely including

many loci violating the assumptions of the CMH test. This is consistent with our results

which suggest that despite containing false positives, SNPs with significant CMH test p-

values after Bonferroni correction are nevertheless enriched with loci that display

broadly consistent allele frequency shifts between healthy and unhealthy pools.

Overall, our newly developed pangenome captures a diverse set of SVs from across the

geographic range of European ash, including a large quantity of previously

uncharacterised sequence and dispensable genes, making it an essential resource for

full investigation of the role of both SNPs and SVs in a breadth of ecological traits for

this important tree species. After carefully accounting for spurious signals in our GWAS,

we detect signals of a shared component of low susceptibility to ADB across UK

provenances. Explicitly identifying variants with consistent effects across a range of

seed sources may be useful in developing widely-applicable genomic prediction models

for low ADB susceptibility, and ultimately informing breeding programs to restore this

threatened species.

Methods

Reference assembly

To construct a phased, chromosome-level assembly, we sampled a potted graft growing at Royal Botanic Gardens, Kew. The graft was created by Forest Research in 2017 using a rootstock and a scion from *F. excelsior* individuals of unknown provenance. Leaf tissue from this graft was flash frozen in liquid nitrogen in May 2022 and sent to Cantata Bio, CA, USA (formerly Dovetail Genomics) on dry ice, where DNA extractions, library preparation, sequencing and genome assembly was performed as outlined below.

DNA was extracted by Cantata Bio using an in-house CTAB protocol and quantified using Qubit 2.0 Fluorometer (Life Technologies, Carlsbad, CA, USA). The SMRTbell Express Template Prep Kit 2.0 (PacBio, Menlo Park, CA, USA) using the manufacturer recommended protocol was used to construct a PacBio SMRTbell library (~20kb) for PacBio Sequel. The library was bound to polymerase using the Sequel II Binding Kit 2.0 (PacBio) and loaded onto PacBio Sequel II). Sequencing was performed on PacBio Sequel II 8M SMRT cells, resulting in 66Gb data (approximately 79X coverage based on the prime 1C genome size estimate for *F. excelsior* of 840Mb[54,104]. For the Omni-C libraries, chromatin was fixed in place with formaldehyde in the nucleus and then extracted. Fixed chromatin was digested with DNAse I, chromatin ends were repaired and ligated to a biotinylated bridge adapter followed by proximity ligation of adapter containing ends. After proximity ligation, crosslinks were reversed and the DNA purified. Purified DNA was treated to remove biotin that was not internal to ligated fragments.

Sequencing libraries were generated using NEBNext Ultra enzymes and Illumina-

compatible adapters. Biotin-containing fragments were isolated using streptavidin beads

before PCR enrichment of each library. The library was sequenced on an Illumina

HiSeqX platform to produce approximately 30X sequence coverage in 150bp paired-

end reads. The combined Illumina and PacBio data was used to produce an initial

assembly with hifiasm v0.15.4-r347[105] using default parameters. The *de novo* assembly

and Dovetail OmniC library reads were used as input data for HiRise[106] to produce a

draft assembly. OmniC library reads were then aligned to the draft input assembly using

BWA[107] and reads with MQ < 50 removed. The mapping information was analyzed

using HiRise to produce a likelihood model for genomic distance between read pairs,

and the model was used to identify and break putative misjoins, and to score and make

joins.

After receiving the assemblies and raw data from Cantata Bio, the haploid genome

assemblies were assessed using QUAST v5.2[108] and BUSCO v5.5.4 using the

eudicots_odb10 database[109]. To assess potential contaminants, scaffolds from the most

contiguous of the two haploid genomes (BATG-1.0) were searched against the NCBI nt

database (downloaded 21/02/23) using blast+ v2.11.0[110] with settings -max_target_seqs

10 -max_hsps 1 -evalue 1e-25. To assess coverage, the PacBio data was mapped to

the haploid genomes using minimap2 v2.5[111]. The combined data was analysed using

blobtools v4.0.7[112]. . To assess LTR completeness, LTRharvest from genome-tools

v1.6.6 [113] and LTR_finder v1.07[114] were run with default parameters, other than

specifying a 85% similarity threshold, length range 100-5000bp, starting/ending motif

TG/CA and maximum distance of LTR starting positions of 20,000. LAI was calculated

using LTR_retreiver v3.0.4[56]. To estimate genome size using kmers, LVgs v1.00[115] was

run on the PacBio fastq files, with kmer sizes between 17 and 77bp in increments of 10,

and results analysed using Genomescope2 v2.1.0[55].

The PacBio data was used to *de novo* assemble a chloroplast genome for *F. excelsior*

using Oatk v1.0[116] using HMMER v.3.1b2 (hmmer.org) with parameter settings -c 30 -k

1001 and a chloroplast gene profile database provided by OatK (embryophyta_pltd).

Annotation was performed using GeSeq v2.03[117] using default settings.

Long-read RNA extraction and sequencing

Long-read RNA seq data from leaf and cambium tissue was generated to annotate the

new genome assembly from the reference individual. Leaf material was collected from

the same grafted tree as sampled for the reference assembly (see above) in May 2022

and flash frozen in liquid nitrogen. For cambium tissue, twigs were sampled in January

2023 and the layer of tissue immediately below the bark was isolated using a fresh

razor blade before flash freezing in liquid nitrogen. Tissue was stored in a -80 freezer

until RNA extraction.

For long-read RNA extraction, a modified version of the Qiagen RNeasy Plant Mini Kit

standard protocol was used (Supplementary Note 2). In brief, 25-30mg tissue was used

with the standard kit protocol (including the optional DNase step) and eluted in 50µl

RNase-free water. 450µl buffer RLT was added to the RNA, and the RNeasy kit protocol

was repeated (without the DNase step). RNA was quantified using a Nanodrop 2000

and a Promega QuantiFluor® RNA System Kit using a Quantus fluorometer. Samples

were shipped to the Deepseq facility at the University of Nottingham and fragment

length determined with an Agilent Tapestation using an RNA Screentape kit.

cDNA libraries were prepared using an Oxford Nanopore PCR-cDNA Sequencing Kit

(SQK-PCS109), loaded onto R9.4.1 flowcells and sequenced on a Gridion X5 Mk1 –

this was performed once for the cambium library and twice for the leaf library. Runs

were performed with high-accuracy basecalling (running on MinKNOW software version

22.12.5). All data reported relates to reads that passed a base calling quality filter

(PHRED score >9).

Sampling for pangenome construction

In sampling individuals for the ash pangenome, we aimed to i) sample a broad range of

European populations, with ii) a particular focus on UK diversity by sampling individuals

representing different NSZ in the UK[118] (see

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachmen

t_data/file/797909/Regions_of_provenance_and_seed_zone_map.pdf). With this

approach we aimed to capture a global set of variants from across the species range,

whilst increasing our chances of capturing SVs segregating in the mainly UK

populations for which short-read data is available from Stocks et al. [42].

In total, 50 samples were used for long-read sequencing. To maximise the chances of

identifying SVs segregating in the poolseq data from Stocks et al. [42] one individual from

each of the 13 NSZ used in the Forest Research (FR) mass screening field trial at North

Burlingham, Norfolk, UK (Site 16)[42] was sequenced, with the exception of NSZ-204

where two were sequenced, as individually sequenced data for this provenance were

available from Stocks et al.[42] The North Burlingham site was planted in 2013 with two-

year-old saplings that were grown for non-research purposes in commercial nurseries

from seed sources from 15 different provenances. The site contained four complete

replicate plots for each seed source, with 5,000 trees per ha (a spacing of

1 m × 2 m). At the time of sampling, the site had high *H. fraxineus* inoculum pressure

from naturally established infections, and ADB damage was severe throughout. In 2022

(when trees were eleven years old) individuals were sampled that appeared relatively

healthy within each plot, and scored between 3 and 5 on the damage scale of Pliura et

al.[37] (where 3 is a tree with moderate damage from ADB, 4 a tree with slight damage

and 5 a tree with no visible signs of damage) to increase the chances of capturing

alleles for low ADB susceptibility. Three wild trees of unknown age from seed zones in

the UK not covered by the FR trial populations (NSZ-305, NSZ-406, NSZ-402) were

also included to further capture UK diversity. To increase the likelihood of including

more of the range-wide diversity of structural variation in *F. excelsior*, one individual

from each of 29 mostly mainland European provenances was sampled from the

Realising Ash's Potential (RAP) trial (established in 2005), as well as two from

provenances in the FRAXIGEN trial (established in 2004), both at Paradise Wood, Earth

Trust, Oxfordshire. Finally, leaf material from the individual used for the linear reference

genome was also included. Sample information is outlined in Supplementary Table 3,

including approximate coordinates of origin where available. A map based on these

coordinates is shown in Figure 1A , overlaid with a shapefile of the *F. excelsior* native

range from Caudullo et al.[119]. We note our sampling does not comprehensively cover

the geographic range of *F. excelsior* - in particular, the far south western, south eastern

European, west Asian, and far northern areas of the range are unrepresented in our

samples (See Fig. 1A).

Sampling for long-read DNA sequencing was performed in May 2022. For samples

collected in the field, intact stems with leaves were excised from individual trees using

secateurs or a pole pruner, wrapped in damp tissue paper and sealed in airtight plastic

bags before returning to RBG Kew for storage at 4°C. Leaf tissue was removed from the

stems and frozen within 48 hours of collection by immersing in dry ice, and then stored

at -80°C prior to DNA extraction.

High molecular weight DNA extraction, and RNA extraction, from

pangenome samples

For high molecular weight DNA extraction, a modified CTAB protocol was used[50] (see

Supplementary Note 6 for full protocol). In brief, 300-400mg tissue was disrupted in

liquid nitrogen and added to 8ml Carlson buffer with 20µl beta-mercaptoethanol and

40µl Qiagen Proteinase K. This was incubated at 65^{o}C for 1 hour, before adding 16µl

Qiagen Rnase A and incubated for a further 15 minutes. 1 volume of SEVAG solution

was added, the tube inverted, and centrifuged for 10mins at 4^{o}C before the aqueous

phase was removed. This step was then repeated. 0.7 volumes of isopropanol was

added and the tube incubated at 4^{o}C for 15mins. The sample was centrifuged at 4^{o}C for

30mins to precipitate the DNA and resuspended in Qiagen buffer AE before being

redissolved in 5ml Qiagen Buffer G2 for 15 minutes. A Qiagen Genomic Tip 100/G Kit

was used to purify the DNA; the "Isolation of Genomic DNA from Blood, Cultured Cells,

Tissue, Yeast or Bacteria using Genomic-tips" protocol from the kit handbook was used,

with the modification of using 10ml rather than 7.5ml buffer QC. DNA was eluted in

100µl buffer 1x TE and left to resuspend at room temperature and then at 4^{o}C for up to

3 and 4 days respectively. DNA was quantified using a Quantus Flurometer with

QuantiFlour dsDNA kit, and a Nanodrop 2000. Integrity was measured using an Agilent

Tapestation with a Genomic DNA kit, and the DNA shipped to the University of York on

dry ice to prevent fragmentation during transit, where library preparation was performed

as described below.

Short fragment elimination was performed using a Circulomics SRE Standard Kit.

Library preparation was performed using the Oxford Nanopore Technologies (ONT)

SQKNBD114.24 (Protocol version NBE_9169_v114_revD_15Sep2022-promethion.pdf)

library preparation kit, with the following modifications: i) Reaction volumes were

proportionately scaled up to 60µl total for end repair and A-tailing, and incubation times

increased to 30 minutes each at 20°C and 65°C, ii) Reaction volumes were doubled for

the barcode adapter ligation step, which was performed at room temperature for 1 hour.

Barcode-ligated samples were bead cleaned separately before pooling at appropriate

volumes prior to adapter ligation. Native barcodes were ligated and reads demultiplexed

by barcode following the run. Libraries were loaded onto PromethION flowcells, runs

commenced on a Gridion with live super accuracy base calling (running on MinKNOW

software version 22.10.5) with barcode demultiplexing and adapter trimming. All data

reported relates to reads that passed a base calling quality filter (PHRED score >10).

For 35/50 samples with remaining leaf tissue after DNA extraction, RNA from frozen

material was extracted for short read sequencing (outlined in Supplementary Table 3).

For one additional sample (PG5), fresh leaf tissue was collected in May 2024 and

stored in Thermofisher RNALater, and stored at 4°C until RNA extraction in November

2025. The same RNA extraction protocol was used as previously described

(Supplementary Note 5). Samples were sent to the Centre for Genomic Research

(CGR), Liverpool. Analysis using an Agilent Tapestation did not reveal any evidence of

genomic DNA contamination. cDNA libraries were prepared using a New England

Biolabs NEBNext polyA selection and Ultra II Directional RNA library preparation kit,

and sequenced on half a lane on an Illumina NovaSeq X Plus using 25B chemistry, with

paired-end 15-bp reads. Raw data was trimmed by CGR using Cutadapt[120] v4.5 with

option -O 3. Further trimming was performed using Trimmomatic[121] with settings

LEADING:7 TRAILING:7 ILLUMINACLIP:2:30:7 MINLEN:70, resulting in an average of

63 million read pairs per sample after trimming (Supplementary Table 3).

707 Long-read DNA data analysis

The simplex ONT DNA sequence data generated was trimmed, firstly using

Trimmomatic [121] with settings LEADING:7 TRAILING:7, followed by Nanofilt v2.8.0[122]

with minimum average read quality of 7 and length 1000. Quality was inspected and

summary statistics generated using Nanoplot v1.4.10[123], FastQC v0.11.9[124] and

multiQC v1.14[125].

Reads were mapped to BATG-1.0 using minimap2 v2.5[111], with PCR duplicates and

reads with mapping quality (MAPQ) <20 removed using samtools v1.9[107]. Mean depth

after filtering was 22.7X (Supplementary Table 3).

To inspect population structure of these samples, SNPs were called individually for all

samples other than the reference genome individual using Clair3 v.1.0.5[126] with model

r941_prom_sup_g5014. Gvcfs were merged using glnexus v1.4.1-0-g68e25e5n[127].

Genotypes with DP < 8 or > 60 were set to missing using bcftools v1.16[128], and sites

with >10% missing data or MAF < 0.05 were excluded. Plink2 v2.00a3.3LM[129] was used

to identify unlinked SNPs (using setting –indep-pairwise 50 10 0.1) and calculate a PCA

using default settings.

Read-based SV calling was performed using the long-read data generated for all 50

samples using Sniffles2 v2.0.7[62] and cuteSV v2.0.3[63], two tools that rank highly across

several comparisons of read-based SV calling software[130–137]. Both tools were run with

a minimum read support of 5, and cuteSV with the additional parameters –

max_cluster_bias_INS 100 –max_cluster_bias_DEL 100 –diff_ratio_merging_INS 0.3 –

diff_ratio_merging_DEL 0.3. Breakends/translocations (BND) or sites which were

genotyped as homozygous reference were removed.

To complement the read-based SV calling, *de novo* assemblies were produced from the

ONT read data. *De novo* assembly from trimmed reads was performed using shasta

v0.11.1[65] with the Nanopore-May2022 config file, Flye v2.9.2-b1786[64] with default

settings, and nextDenovo v.2.5.0[66] with an estimated genome size of 850Mb. Assembly

contiguity was assessed using QUAST. SVs were called using these assemblies by

mapping them to BATG-1.0 using minimap2 in asm10 mode. SVs were called with

svim-asm v.1.02[67] using alignments with a minimum MAPQ of 20 and minimum SV size

of 50. Haplotype 2 of the linear reference genome was also mapped to BATG-1.0, and

SVs called using the above method. SV calls between the two pseudochromosomal

haploid genomes were compared to SV calls from the ONT data from the same

individual using the various approaches, with SV sets merged using SURVIVOR

v1.0.7[138] using settings 200 2 1 1 1 50 for the intersection of sets and 200 1 1 1 1 50 for

the union. A 200bp merge distance yielded similar numbers of SVs called with Sniffles2

and cuteSV to values between 10 and 300bp yielded similar numbers of SVs, with

larger or smaller values resulting in fewer SVs. Assuming the SV calls between the two

haploid linear reference genomes are a "truth" set, precision, recall and F1 were

calculated for the various SV calling methods from the lower coverage ONT data. Based

on these comparisons (see Results), SVs for each individual were included if they were

called in at least one of i) cuteSV and Sniffles2 ii) svim-asm from mapping the shasta

assembly to BATG-1.0.

These SVs were merged across all 50 individuals using SURVIVOR with settings 200 2

1 1 1 50. SVs that overlapped with Ns in the reference were removed. SVs that were

called in more than 3/50 individuals were identified using bcftools v1.16 and used for

downstream analysis. The overlap between these SVs and the repeat annotation was

assessed using the bedtools v2.28.0[139] window function with parameter -w 1.

Consensus sequences for insertions and deletions were searched against the ncbi nt

database using blast+ with parameters -max_hsps 1 -evalute $1e^{-25}$, with the resulting

data loaded into blobtools. Based on analysis with blobtools, four sequences were

identified as putative contaminants and the associated SVs removed from the vcf. The

resulting vcf, as well as the BATG-1.0 fasta was used to construct a pangenome using

the vg v1.64.0[68] construct command with option -v -S -a.

761 Linear reference and pangenome annotation

Repeats were identified using Repeatmodeler v2.0.1[140] in LTRStruct mode. The outputs

of this were combined with the custom repeat library used previously to mask the

BATG-0.5 assembly[53], and used to mask the genome with Repeatmasker v4.1.2[141] with

the parameters -wublast -nolow. Telomere sequences were identified using Teloscope

v0.0.9[142] with default settings using the canonical plant telomeric repeat CCCTAAA. To

annotate the linear reference genome, short-read RNA-seq data from leaf, root,

cambium and flower tissue that had been used to annotate the BATG-0.5 assembly[53],

was downloaded from the European Nucleotide Archive (ENA), as well as RNA seq

data from Emerald Ash Borer infested and control phloem tissue[143], and ADB infested

and control petiole tissue[144] (Supplementary Table 8).Together with the short-read leaf

RNA-seq data for the BATG-1.0 individual generated as part of this study, these data

were mapped mapped to BATG-1.0 using STAR v2.7.10b with default parameters[145].

The long-read RNA-seq data generated in this study was trimmed firstly using

Trimmomatic v0.39[121] with settings LEADING:7 TRAILING:7, followed by Nanofilt

v2.8.0[122] with minimum average read quality of 7 and minimum length 100. The trimmed

data was then mapped to the BATG-1.0 reference genome using minimap2 v2.5[111]. We

acknowledge that this RNA-seq data does not provide comprehensive coverage of

tissues/timepoints/conditions in the species, and that some genes may therefore be

missed. Variability in the gene annotation process could lead to false positives during

the identification of variable genes[16,146], so we repeated the annotation process multiple

times and combined the results. BRAKER3[60] was run 10 times on the softmasked

genome using the bam files from the short read data, and protein orthologue data from

Viridiplantae from orthodb v11[147]. A version of BRAKER3 capable of using long-reads

(singularity build braker3_lr.sif docker://teambraker/braker3:devel) was also run 10

times using the bam files from the long read data and the same protein orthologue set

as used with the short read data. To combine the outputs of the runs, the amino acid

sequences of the longest transcript per gene (minimum length 30 amino acids) for each

run were clustered into orthogroups using Orthofinder v2.5.5[61]. Within each orthogroup,

groups of sequences with non-overlapping positions within the BATG-1.0 assembly

(likely representing recent duplicates) were separated using the bedtools intersect

utility. For each of these groups, the gene model with the longest transcript from the set

of 20 BRAKER3 runs was chosen. In some cases, OrthoFinder split genes with largely

overlapping coding sequences into separate groups. In cases where two genes with a

coding sequence on the same strand overlapped by >50% for one of the genes, only

the gene producing the longest protein was retained (for further discussion on clustering

see Supplementary Note 7). The features for the selected gene models were combined

into a single bed file. The amino acid sequences from the final set of transcripts was

used as the input for Orthofinder, along with protein sequences for BATG-0.5[53],

Fraxinus pennsylvanica[51], *Olea europaea* subsp. *europaea*[148], *Solanum*

lycopersicum[149], *Erythranthe guttata*[150], *Sesamum indicum*[151] and *Salvia miltiorrhiza*[152]

(Supplementary Table 2).

To annotate the pangenome, sequences and locations of structural variants were used

to modify the sequence of the BATG-1.0 fasta, which was then annotated. As most SVs

were defined from read mapping to BATG-1.0 rather than from the individual *de novo*

assemblies, this approach maximises the opportunity for novel genes/gene deletions to

be explicitly linked to underlying SVs. The steps were as follows (see Extended Data

Figure 10 for an explanatory diagram): for each individual, a non-overlapping set of SVs

detected in that individual was extracted from the combined vcf of SVs used to build the

pangenome. Where SVs overlapped, insertions were retained in priority to other types

of SVs, and longer SVs were retained in priority to shorter ones, as these may be more

likely to capture coding sequences absent in BATG-1.0 This process resulted in 2,394

deletions, 96 insertions and 21 inversions being left unrepresented in any of the

samples (0.7% of SVs). Tandem duplicates (0.5% of SVs), which were not included in

the graph pangenome, were also excluded. Based on these sets of SVs, sequence was

then inserted, deleted, or inverted in the reference fasta (the SV sequence in the

combined vcf of SVs used to build the pangenome was used, so the same sequence

was used to represent a given SV in different individuals) to produce a transformed

fasta for each individual in turn. The transformed fasta was then annotated for each

individual. Repeats were annotated and softmasked as described above. Long and

short read RNA-seq data from the reference individual were mapped to each

transformed fasta using the same methods as for the reference sequence. For the 35

non-reference individuals where short-read RNA-seq data was generated as part of this

study, leaf RNA-seq data from each individual was also mapped to the corresponding

transformed fasta and used in downstream annotation. A single run of the short-read

BRAKER3 protocol, and a single run of the long-read BRAKER3 protocol was

performed as detailed above. For these two runs for each individual, the Orthofinder

approach as outlined previously was used to consolidate the outputs of BRAKER3 into

a single set of proteins. Orthofinder was then run again for each set of proteins across

the 50 individuals, as well as the proteins from the consensus annotation of BATG-1.0.

Non-overlapping groups of sequences were separated using bedtools as before. For

each gene, bedtools intersect was used to identify SVs that overlapped the gene

sequence. Novel genes not present in the BATG-1.0 genome should overlap with an

SV, as should deletions of genes present in the reference genome. For each novel gene

that overlapped a SV at least once (i.e. in at least one sample, including the reference

individual sequenced with long reads), an F1 score was calculated to identify the

strength of the association between the new gene and the SV: $2 \times N\_Gene\_SV / (2 \times$

$N\_Gene\_SV + N\_Gene\_NSV + N\_NGene\_SV)$ where N_Gene_SV is the number of

samples with the gene and the SV, N_Gene_NSV is the number of samples with the

gene but without the SV, and N_NGene_SV is the number of samples without the gene

but with the SV. For novel genes overlapping more than one SV, the SV with the

highest F1 score was retained as potentially causal for the novel gene. An equivalent

calculation was performed for genes present in the BATG-1.0 genome but absent in at

least one sample, where an SV overlapped with the annotation coordinates for the gene

in the reference. To account for genes potentially being associated with multiple SVs,

we also calculated an F1 score of gene presence/absence with all overlapping SVs - if

this scored higher than any individual SV, this score was taken forward. Genes with an

F1 score of 0.8 or above were retained in the final pangenome annotation as being

dispensable; for those with an F1 score of < 0.8, those absent in the BATG-1.0 genome

were removed, and those present in the BATG-1.0 genome were classified as invariant.

Genes that did not overlap an SV in any sample were classified as invariant if they were

annotated in BATG-1.0 and were absent in some other samples, and discarded if they

were absent in BATG-1.0 but present in some other samples. To assess the impact of

OrthoFinder potentially oversplitting genes, in a separate analysis, orthogroups

containing at least one pair of genes with a coding sequence on the same strand

overlapping by more than 50% were merged, and SV/Gene F1 scores recalculated. The

results for this were qualitatively similar to the unmerged dataset, so the latter was used

subsequently - for more discussion see Supplementary Note 7.

The complete set of putatively dispensable genes (i.e. before removing those with an F1

score <0.8) was functionally annotated. Protein sequences were searched against

Araport 11 representative gene peptides (downloaded 14/09/2022), the NCBI SR

database (downloaded 20/08/24), and UniProt/SwissProt database (downloaded

24/07/24) using DIAMOND v2.1.9.163[153] in blastp mode. Interproscan v5.69.101.0[154]

with default settings was used to search for protein domains. GO terms were assigned

using eggNOG-mapper v2.1.12[155] by using DIAMOND blastp to search the eggNOG

database. GO enrichment was performed using topGO v2.62.0[156] using annot =

annFun.gene2GO, nodeSize = 10 and the exact Fisher test with Bonferroni correction.

OrthoFinder was run with the complete set of potential pangenome protein sequences

using the same species and parameters described above.

SNP and SV calling from short read data from individuals and

pools

To test the impact of using the pangenome on read mapping accuracy, short-read data

from 42 individuals separately whole genome sequenced in Stocks et al.[42] (this number

selected to reflect the average number of individuals in a pool in that study, see

generation of *in silico* pools below) was mapped using vg giraffe[68] to either i) the full

pangenome or ii) the BATG-1.0 genome converted into pangenome format using vg

construct. The resulting gam files were converted to bam files in the BATG-1.0

coordinate system using vg surject. Mapping statistics were calculated using samtools.

For each mapping approach, SNPs were called using bcftools v1.19 call and calls

merged across individuals. Genotypes with DP <5 or >50 were set to missing, then sites

with >10% missing individuals or a MAF of <0.05 were removed.

For SV calling from these individuals, vg pack was used with parameter -Q 5, and SVs

called using vg call with the vcf used to construct the pangenome as a reference. For

SNP calling, bams were produced using vg surject. SNPs were then called using

bcftools v1.19 call. SNPs and SVs were merged across individuals, and calls with DP <

5 or >50 were set to missing, then sites with >10% missing individuals or a MAF of

<0.05 were removed. To produce similar numbers of SNPs and SVs for linkage

analysis, 1% of SNPs were randomly sampled using vcflib v1.0.3[157] function

vcfrandomsample. SNP and SV data were combined using bcftools v1.19 concat option.

Pairwise linkage between markers was calculated using plink v1.9.-170906[158], with

parameters –ld-window-kb 1000 and –ld-window 99999. Linkage decay with distance

between SVs alone, between SNPs and SVs, and between SNPs alone were analysed

separately.

To identify the accuracy of SNP and SV calling from poolseq data using the

pangenome, read files from the 42 individually sequenced samples (which each

contained approximately the same number of reads, equating to approximately 22X 1C

genome coverage) were combined, and 10% of read pairs randomly subsampled with

seqkit v2.8.0[159]. This corresponds to approximately 2.2X coverage per individual, the

average per-individual coverage of the poolseq data in Stocks et al.[42]. SNPs and SVs

were called from this *in silico* pool using the approaches above, but with sites being set

to missing if they had a DP < 50 or DP > 200. Allele frequencies for SNPs/SVs were

calculated as the number of reads supporting the alternate allele divided by the total

number of reads. For the merged individually called SNPs/SVs, the correlation between

the alternate allele frequency in the individually called samples and the estimated allele

frequency for the *in silico* pool was tested with a linear model in R v4.3.1[160].

GWAS for low susceptibility to ash dieback from poolseq data

A poolseq dataset from Stocks et al.[42] was used to perform a GWAS for low

susceptibility to ADB. In brief, this data was generated as follows: leaf samples were

taken from four FR mass screening trial sites in England (including the North

Burlingham site where individuals were sampled for constructing the pangenome), from

six year old ash trees from a total of 12 different seed sources (10 from UK NSZ, one

from Ireland [IRL-DON] and one from Germany [DEU]), as well as a group from a

breeding seedling orchard (sourced in the UK) planted by the Future Trees Trust (FTT-

SO). For each of the 13 sources, a pool of healthy trees (no visible damage) and a pool

of unhealthy trees (main stem dead, or partially dead) - 42 trees per pool on average -

were sequenced with the Illumina HiSeq X platform in 150bp paired-end reads. Tree

health was assessed based on a tree health score designed to assess the overall level

of damage from ADB - it should be noted that low susceptibility to ADB is a complex

trait, which may not be associated with long-term tree health, and may be underpinned

by multiple underlying mechanisms. Two seed sources had a second pair of pools,

produced as biological replicates, for a total of 30 pools (excluding one technical

replicate). See Stocks et al.[42] for full details. The trimmed, paired short read data from

ENA (accession ERA1732352, Submitted Files) was used.

Here, these reads were mapped to the pangenome and read depths for the reference or

alternate paths were extracted from the vcf for each pool to produce estimates of SV

allele frequencies, as described in the previous section. The SV allele frequencies were

then merged across pools. For SNPs, the output of vg giraffe was converted to linear

reference coordinates using vg surject, with PCR duplicates and reads with MAPQ < 20

removed using samtools v1.9[161]. SNPs were called from the resulting bam files using

samtools mpileup with -d 8000 and the mpileup2sync function from Popoolation2[71]

v1.201 with –min-qual 20 to produce a .sync file. Sites were removed if there was a third

most frequent allele with a total depth higher than 15. The frequencies of major alleles

for i) SVs and ii) a random 1% of SNPs were used to perform PCA using the R prcomp

function.

To identify SV and SNP allele frequency changes associated with ADB damage,

Cochran-Mantel Haenszel (CMH) tests were performed using the cmh-test.pl function

from Popoolation2, with parameters min_count 15, min_coverage 40 and

max_coverage 200. For sites with a significant result ($p < 1 \times 10^{-13}$), the Woolf test was

used from the vcd R package[162] to test for significant deviation from homogeneity

across groups using the alternate/reference read counts for each pool. To quantify

heterogeneity in allele frequency shifts, Cochran's Q[73] and I^2 [81] were calculated.

Cochran's Q was calculated as $Q = \sum_{i=1}^{k} w_i (x_i - x_w)^2$ from Cochran (1954)[73], with

x_i being the estimated effect size from pool i, x_w the weighted mean effect size estimate

across all k pairs of pools, and w_i being the inverse of the variance estimated for each

pair. For the effect sizes, we used natural logarithm transformed of individual odds

ratios [i.e. ln(ad/bc)], where a = alt reads in healthy, b = ref reads in healthy, c = alt

reads in unhealthy, d = ref reads in unhealthy) as recommended in Woolf (1955)[80] and

performed in other studies[163]. The variance in this measure was estimated as 1/a + 1/b

+ 1/c + 1/d, as specified in Woolf (1955)[80]. The natural logarithm transformed overall

CMH test odds ratio is used as x_w. These values were used to calculate Q. I^2 was

calculated from these as I^2 = (Q - df) / Q, where df is the degrees of freedom; in this

case, the number of pairs of pools minus one. For SNPs with p < 0.05 from the CMH

test after Bonferroni correction, SNPs with a standard deviation of depth of >15,

indicative of potentially uncaptured SVs that may interfere with allele frequency

estimation (see Results, Supplementary Note 1), were removed, along with SNPs that

showed deviation from homogenous odds ratios (violating an assumption of the CMH

test) - namely, SNPs with either an I^2 value of <0.25, or a Woolf test p-value of <0.05.

220 of the SNPs classified as significant after Bonferroni correction remained after this

filtering. Genes from the pangenome that overlapped these SNPs, or were within 10kb,

were identified using bedtools. For those with annotated GO terms, enrichment was

performed as described above for dispensable vs. indispensable genes.

An alternative statistical test for GWAS, a quasibinomial generalized linear model

(qGLM), was run using the poolFreqDiffTest_QGLM.py script from the PoolFreqDiff

repository[75], using parameters -mincnt 15 -minc 40 -maxc 200 -n 45. Different methods

of accounting for the effect of provenance were investigated, see Supplementary Note 2

for further details. Fixation index (F_{st}) and Fisher's exact test were also calculated in

sliding windows for pairs of pools - see Supplementary Note 3 for further details.

Acknowledgements

We would like to thank i) Genomics Laboratory, Bioscience Technology Facility, University of

York, ii) Deep Seq: Next Generation Sequencing Facility, University of Nottingham, iii) Cantata

Bio LLC, California for their contribution to sequencing and genome assembly, iv) Earth Trust

and other landowners for facilitating access for sampling. We would also like to thank Mike

Charters and Owen Blake at RBG Kew, Domen Finzgar and Liz Richardson at Forest Research,

Jo Clark at Future Trees Trust, and Paul Hill at the Earth Trust for assistance with sample

acquisition, and Laszlo Csiba and the Plant Health and Adaptation team at Kew for valuable

advice on this work. We are grateful to Andrea Harper and Sara Franco Ortega for early access

to their DNA extraction protocol. This research utilised Queen Mary's Apocrita HPC facility,

supported by QMUL Research-IT (http://doi.org/10.5281/zenodo.438045). This work was funded

by the UK Government Department for Environment Food & Rural Affairs through a Centre for

Forest Protection grant (no. CFP2208) to LJK, DPW and RW. The results of this project LL2317

were obtained with the financial contribution of the Ministry of Education, Youth and Sports as

part of the targeted support of the ERC CZ program, through the contribution of LY. DG and LJK

received support from the Natural Environment Research Council [grant number NE/X015351/1

to LJK]. Figure 1 "contains, or is based on, information supplied by the Forestry Commission. ©

Crown copyright and database right [2009] Ordnance Survey [100021242]"

Data availability

988 Source Data are provided with this paper. Sequencing data is available on NCBI under

989 BioProjects PRJNA906474 (long-read DNA sequence data), PRJNA942609 (long-read RNA

990 sequencing data) and PRJNA1390666 (short-read RNA sequencing data). Annotation files for

991 the BATG-1.0 genome and pangenome will be made available on Zenodo.

Code availability

993 Code needed to reproduce the analysis will be made available at

994 https://github.com/danielwood1992/Ash_pangenome.

1000

1001

1002

1003

1004

1005

1006

1007

1008 # Figures

1009

A

B

1010
1011

Figure 1 - Geographic origins and genetic structure of samples

A) Map showing location of origin for individuals used to construct the pangenome; see Methods

and Supplementary Table 1 for full details. Points represent individuals sampled or approximate

sample locations for seedlots from which individuals were grown. Light yellow areas represent

the native range of *F. excelsior* using shapefiles from Caudullo et al.[119]. B) Principal components

analysis of single nucleotide polymorphism (SNP) calls from long-read sequencing data from

individuals sampled. First two principal components are plotted, with the percentage of variation

explained by each principal component shown on the axis labels. In Figures A and B, individuals

are coloured by country of origin (see Supplementary Table 1).

1021

1022

1023

1024

1025

1026
1027

1028

1029

1030

1031

1032

1033

1034

Figure 2 - Structural variant properties

A), C): The total number of SVs (A) and total quantity of sequence (C) of different SV classes, at different minimum numbers of individuals called from long-read data. Colour-coding indicates different types of SVs: insertions (green), deletions (orange), tandem duplications (light blue) and inversions (dark blue). A vertical red line at n = 3 indicates the cutoff for SVs being included in the pangenome. Note: to aid visualisation, separate plots for inversions and tandem repeats are available in Supplementary Figure 6. B) Saturation curves of sequentially merging SVs from different individuals. Each sequential series of SVs was generated by combining individuals in a random order, one at a time, with the total number of SVs in the group being displayed; separate series of random combinations are designated by colour, with points in the same series joined by a line. The X axis represents the number of individuals merged, the Y axis the total number of SVs in those individuals. D) Circular plot of feature densities per chromosome in 100kb windows. The 23 largest scaffolds are labelled 1-23 in descending order of size.Tracks, from outer to inner correspond to density of i) genes (blue), ii) insertions (orange), iii) deletions (red) and iv) inversions (purple). Larger plots for each individual chromosome, also including repeat and tandem duplication density, are presented in Supplementary File 1.

1053

1054

Figure 3 - Consistency of association of putatively dispensable genes with

structural variants

A) Illustration of classification of genes in the pangenome based on their presence/absence,

and how this corresponds with structural variants (SVs). Genome sequence is depicted as a

blue solid line; SVs by light blue dashed lines (representing deleted sequence relative to the

reference in this illustration). Genes are depicted by black arrows. For illustration, each gene in

the reference sequence (top line) is homologous to the gene immediately below them in the

example sample (bottom line). For those present in the BATG-1.0 reference, genes can be

present in every other sample (1) or sometimes absent (2, 3). The absence of a gene can either

coincide with an SV overlapping the gene in at least one sample (2), or with no overlap with an

SV (3). For genes absent from the reference but present in at least one sample, this presence

can either coincide with an overlapping SV in at least one sample (4), or never coincide with an

overlapping SV (5). B) Barplot of the number of genes in each category in the pangenome,

where numbers correspond to those in part A. Following reclassification of genes in each

category as invariant (orange), dispensable (blue), or discarded from the analysis (grey)

according to how consistently they overlap with SVs based on an F1 score. Genes in category 2

with an F1 score > 0.8 are classified as dispensable (blue); those with a score below that

threshold are categorised as invariant. Genes in category 3 are never associated with an SV so

are all categorised as invariant. Genes in category 4 with an F1 score > 0.8 are classified as

dispensable, those with a score below that are discarded from the analysis. Genes in category 5

are never associated with an SV so are discarded from the analysis. In an approach where

genes absent in some individuals are treated as dispensable, all the genes in categories 2-5

would be treated as dispensable genes.

1078

1091 **Figure 4 - Annotation of putatively dispensable genes**

1092

A) Barplots showing the percentage of genes annotated in the eggNOG, InterPro and UniProt

databases, for genes included in the pangenome after filtering (black), genes removed during

filtering (grey), included genes classified as dispensable (i.e. the high-confidence dispensable

genes; light blue) and included genes classified as invariant (orange). B) Results of Gene

Ontology (GO) enrichment analysis of the included dispensable genes, against a background of

the included dispensable genes plus the putatively dispensable genes removed during filtering

for not being consistently associated with an SV, with points coloured by significance in other

GO enrichment analyses. Values along the x-axis represent the ratio of the observed number of

genes in the included dispensable genes to the number of expected genes in the total

background set. Y-axis values represent the $-\log_{10}$ transformed p-values from the GO

enrichment test. The red dashed line indicates the significance threshold. Points are coloured

according to whether those GO terms were enriched in GO enrichment tests for other sets of

genes. Gray points represent GO terms identified as significantly enriched in the genes

removed during filtering, against a background of excluded plus included genes in the

pangenome. Blue points represent GO terms identified as significantly enriched in genes

classified as variable after filtering, compared to a background of all genes included after

filtering. Purple points represent GO terms identified as significantly enriched in both the

previous comparisons described. The remaining GO terms are coloured in yellow.

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120 **Figure 5 - Shifts in direction between healthy and unhealthy pools in candidate**

1121 **SNPs compared to background SNPs**

1122

1123 For a random 1% of SNPs estimated from poolseq data (panels A and B), and SNPs with

1124 significant p-values from a Cochran-Mantel Haenszel CMH) test, with Bonferroni correction, ☐ =

1125 0.05, (panels C and D), and SNPs with a CMH p-value $< 1 \times 10^{-13}$ (panels E and F), i) the mean

1126 absolute value of the allele frequency shift between healthy and unhealthy pools (A, C and E),

1127 and ii) the proportion of all pairs of pools shifting in the most common direction (B, D and F).

1128

1129

1130

1131

1132

1133

1134

1135

1136 # Extended Data

1137

1138

Extended Data Figure 1 - Precision, recall and F1 scores for different SV calling

methods

Values of the precision (orange), recall (blue) and F1 score (green) of SV calling for different

methods. Values were based on comparing SVs called using the long-read sequencing data

from the reference individual, mapped to BATG-1.0 either as individual reads or assembled

contigs, with the "truth" set comprising variants called by mapping the second haplotype of the

reference genome to BATG-1.0 and calling SVs with svim-asm. The methods considered were:

A) SVs called by both cuteSV and Sniffles2; B) SVs called by svim-asm with a mapping of the

de novo assembly produced by shasta; C) SVs called by svim-asm with a mapping the *de novo*

assembly produced by Flye; D) SVs called by svim-asm with a mapping of *de novo* assembly

produced by nextDeNovo; E) SVs called by methods A and/or B; F) SVs called by methods A

and/or C; G) SVs called by methods A and/or D.

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

Extended Data Figure 2 - F1 scores for putatively variable genes association with SVs

Histograms showing A) for genes absent from the linear reference annotation but present in at

least one other individual, but not overlapping with an SV, the number of individuals where this

gene was present; B) for genes present in the linear reference annotation but absent from at

least one other individual, but not overlapping with an SV the number of individuals where the

gene was absent, C) for genes absent from the linear reference annotation but present in at

least one individual with an SV overlapping the gene location, the best F1 score of the SVs

overlapping the gene, D) for genes present in the linear reference individual but absent from at

least one individual where an SV overlaps with the gene sequence, the best F1 score of the

SVs overlapping the gene.

1175

1176

Extended Data Figure 3 - Read mapping statistics for the linear reference vs. the

pangenome

Read mapping statistics for short reads from 42 individuals sequenced in Stocks et al. (2019),

mapped using giraffe either to the linear reference genome in pangenome format (BATG-1.0),

or to the full pangenome (Pangenome). Each individual is represented by a different colour -

1182   lines connect values in the reference-only "pangenome" to the full pangenome in the individual

1183   panels. The panels outline i) the total number of aligned reads, ii) the total number of paired

1184   reads, iii) the total number of properly paired reads, iv) the number of megabases of softclipped

1185   sequences, v) the mean alignment score, vi) the mean mapping quality and vii) the total number

1186   of substitutions, as estimated by the vg stats function.

1187

1188

1189

1190

1191

Extended Data Figure 4 - Allele depths for SNP and SV calls from data for an example

individual

Histograms outlining allelic depths for A) SNPs and B) SVs called from an example individual -

Individual 164 from Stocks et al. (2019). The panels indicate - for homozygous reference calls

(top row), heterozygote calls (middle row) and homozygous alternate calls (bottom row) - the

1197 alternate allele depth (first column), reference allele depth (second column) and the alternate

1198 allele depth as a proportion of the total allelic depth (last column).

1199

1200

1201

1202

1203

1204

1205

Extended Data Figure 5 - Statistics of SNP and SV calls combined across individuals

A) Histograms of minor allele frequencies for SNPs (top panel) and SVs (bottom panel) called

from a set of individually sequenced ash trees from Stocks et al. (2019). Red vertical dotted line

indicates the median value of minor allele frequency for both datasets. B) For the same dataset,

estimates of the probability of deviation from Hardy-Weinburg equilibrium (HWE) calculated by

bcftools. C) Estimates of linkage disequilibrium decay by distance between SNPs (orange line),

SVs (purple line) and between SNPs and SVs (blue line). Pairwise linkage disequilibrium was

estimated using plink2, with mean R^2 values in predefined bins (red points) estimated in R.

1214

1215

Extended Data Figure 6 - SV and SNP calling from Poolseq data

A), B) Heatmaps of alternate allele frequencies for SNPs (A) and SVs (B) estimated from

individual calls (X axis) from a subset of individually sequenced ash trees from Stocks et al.

(2019) vs. an allele frequency estimate from *in silico* pools generated by randomly

downsampling reads for these individuals, and measuring the allelic depth (AD; Y axis). Colour

scale represents the number of sites in a square. C), D) Plots of the first two principal

components (PCs) from principal components analysis from allele frequencies estimated for

SNPs (C) and SVs (D) from the poolseq data from Stocks et al. (2019)[42]; axis labels represent

the percentage of total variance explained by the PCs. Point labels indicate pool provenances

from Stocks et al. (2019) for a subset of pools (i.e. those outside the main dense cluster). ".1"

and ".2" represent biological replicate pools. Colours correspond to pool health scores - healthy

(green) and unhealthy (black).

1228

1229

1230

1231

1232

1233

Extended Data Figure 7 - GWAS using SV allele frequencies estimated from poolseq

data

A) Manhattan plot for SVs called from poolseq data from Stocks et al. 2019, identifying

association with ash dieback disease damage. Y axis values represent the $-\log_{10}$ p-values of a

Cochran-Mantel Haenszel (CMH) test (red horizontal line corresponds to $p = 1 \times 10^{-13}$). Sites are

ordered by position along scaffolds in the BATG-1.0 genome assembly, with alternating black

and blue colours representing the different scaffolds. Scaffolds are listed in order of size, with

the largest on the left and smallest on the right. B) For SVs identified as significant in the CMH

1242   test, heatmap of alternate allele frequencies estimated from individual calls (X axis) from a

1243   subset of individually sequenced ash trees from Stocks et al. vs. an allele frequency estimate

1244   from *in silico* pools generated by randomly down sampling reads for these individuals and

1245   measuring the allelic depth (AD; Y axis). Colour scale represents the number of sites in a

1246   square.

1247

Extended Data Figure 8 - Depth statistics for SNPs with significant p-values from

Cochran-Mantel-Haenszel tests.

Histograms showing, for different sets of SNP sites called from the Stocks et al. 2019 poolseq

data mapped to the pangenome, i) the ratio of the mean depth of healthy pools vs. unhealthy

pools (panels A, C, E), and ii) the standard deviation of read depths across all the pools. Panels

indicate sets of SNPs - i) a random 1% of total SNPs (A, B), ii) 92 SNPs with a p-value $< 10^{-13}$

(C, D) and iii) 866 SNPs with a p-value passing Bonferroni correction with ☐ = 0.05 (E,F).

1256
1257

1258

1259

1260

1261

1262

1263

Extended Data Figure 9 - Comparison of key statistics with other recent pangenome studies.

Results from the present study compared with a selection of plant pangenome studies published since 2022; pangenome studies using primarily short-read data, or incorporating multiple species, were excluded. Labels correspond to study organism and publication date, detailed in Supplementary Table 7. Number of samples included in the pangenome is on the X axis. Y axis

1264

1265

1266

1267

1268

1269

1270

numbers correspond to total number of SVs discovered (A), additional sequence included in the

pangenome, as a percentage of the reference genome size of that species (B), or the number

dispensable genes as a proportion of the total number of genes in the pangenome. Results from

the present study are in blue; in panel C, Ash_2026 (unfiltered) refers to the proportion of

dispensable genes prior to filtering, Ash_2026 (filtered) refers to the proportion after filtering for

consistent association with SVs - see Results for further details.

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

Extended Data Figure 10 - Explanatory diagram for the pangenome annotation

Explanatory diagram for annotation of the pangenome - see Methods section "Linear reference and pangenome annotation" for full description. For the pangenome annotation, the BATG-1.0 linear reference was transformed using the vcf file of SV calls from each of the 50 ONT samples (1 - S1, S2 and S3 correspond to 3 example samples). For each sample, calls from the vcf used to transform the BATG-1.0 fasta; where an insertion was called, the consensus insertion sequence was inserted at the appropriate position, for deletions the relevant sequence from the BATG-1.0 fasta was deleted. This resulted in a transformed fasta for each sample (2). Using the RNA-seq data from the reference individual, each of the transformed fasta files was annotated (3). Based on sequence homology and position in the genome, genes were clustered across each of the transformed fasta files (4). For each gene cluster, the strength of association of genes with each SV was assessed (5): genes that were consistently associated with an SV were kept as variable genes, whereas those not showing a consistent pattern with an overlapping SV were not.

References

1. Ellegren, H. Genome sequencing and population genomics in non-model organisms. *Trends Ecol. Evol.* **29**, 51–63 (2014).

2. Field, M. A. *et al.* Recurrent miscalling of missense variation from short-read genome sequence data. *BMC Genomics* **20**, 546 (2019).

3. Tang, D. *et al.* Genome evolution and diversity of wild and cultivated potatoes. *Nature* **606**, 535–541 (2022).

4. Zhang, F. *et al.* Long-read sequencing of 111 rice genomes reveals significantly larger pan-genomes. *Genome Res.* **32**, 853–863 (2022).

5. Chen, S. *et al.* Gene mining and genomics-assisted breeding empowered by the pangenome of tea plant *Camellia sinensis*. *Nat. Plants* **9**, 1986–1999 (2023).

6. Liu, Z. *et al.* Grapevine pangenome facilitates trait genetics and genomic breeding. *Nat. Genet.* **56**, 2804–2814 (2024).

7. Liang, Y.-Y. *et al.* Pan-genome analysis reveals local adaptation to climate driven by introgression in oak species. *Mol. Biol. Evol.* **42**, msaf088 (2025).

8. Lynch, R. C. *et al.* Domesticated cannabinoid synthases amid a wild mosaic cannabis pangenome. *Nature* 1–10 (2025).

9. He, F. *et al.* Pan-genomic analysis highlights genes associated with agronomic traits and enhances genomics-assisted breeding in alfalfa. *Nat. Genet.* **57**, 1262–1273 (2025).

10. Bayer, P. E., Golicz, A. A., Scheben, A., Batley, J. & Edwards, D. Plant pan-genomes are the new reference. *Nat Plants* **6**, 914–920 (2020).

11. Crysnanto, D. & Pausch, H. Bovine breed-specific augmented reference graphs facilitate accurate sequence read mapping and unbiased variant discovery. *Genome Biol.* **21**, 184 (2020).

12. Bayer, P. E. *et al.* Modelling of gene loss propensity in the pangenomes of three *Brassica*

species suggests different mechanisms between polyploids and diploids. *Plant Biotechnol.*

J. **19**, 2488–2500 (2021).

13. Mackenzie, A. *et al.* Wheat stripe rust resistance locus YR63 is a hot spot for evolution of

defence genes - a pangenome discovery. *BMC Plant Biol.* **23**, 590 (2023).

14. Zhou, H. *et al.* Pan-genome and transcriptome analyses provide insights into genomic

variation and differential gene expression profiles related to disease resistance and fatty

acid biosynthesis in eastern black walnut (*Juglans nigra*). *Hortic. Res.* **10**, uhad015 (2023).

15. Su, Y. *et al.* Phased telomere-to-telomere reference genome and pangenome reveal an

expansion of resistance genes during apple domestication. *Plant Physiol.* **195**, 2799–2814

(2024).

16. Lovell, J. T. *et al.* Four chromosome scale genomes and a pan-genome annotation to

accelerate pecan tree breeding. *Nat. Commun.* **12**, 4125 (2021).

17. Hämälä, T. *et al.* Genomic structural variants constrain and facilitate adaptation in natural

populations of *Theobroma cacao*, the chocolate tree. *Proc. Natl. Acad. Sci. U. S. A.* **118**,

(2021).

18. Zhou, Y. *et al.* Graph pangenome captures missing heritability and empowers tomato

breeding. *Nature* **606**, 527–534 (2022).

19. Khan, A. W. *et al. Cicer* super-pangenome provides insights into species evolution and

agronomic trait loci for crop improvement in chickpea. *Nat. Genet.* **56**, 1225–1234 (2024).

20. Huang, Y. *et al.* Pangenome analysis provides insight into the evolution of the orange

subfamily and a key gene for citric acid accumulation in citrus fruits. *Nat. Genet.* **55**, 1964–

1975 (2023).

21. Yan, H. *et al.* Pangenomic analysis identifies structural variation associated with heat

tolerance in pearl millet. *Nat. Genet.* **55**, 507–518 (2023).

22. Lei, L. *et al.* Plant Pan-Genomics Comes of Age. *Annu. Rev. Plant Biol.* **72**, 411–435

(2021).

23. Gross, A., Holdenrieder, O., Pautasso, M., Queloz, V. & Sieber, T. N. *Hymenoscyphus*
pseudoalbidus, the causal agent of European ash dieback. *Mol. Plant Pathol.* **15**, 5–21
(2014).

24. Pautasso, M., Aas, G., Queloz, V. & Holdenrieder, O. European ash (*Fraxinus excelsior*)
dieback – A conservation biology challenge. *Biol. Conserv.* **158**, 37–49 (2013).

25. Coker, T. L. R. *et al.* Estimating mortality rates of European ash (*Fraxinus excelsior*) under
the ash dieback (*Hymenoscyphus fraxineus*) epidemic. *Plants People Planet* **1**, 48–58
(2019).

26. Pacia, A. *et al.* Ash dieback in forests and rural areas—history and predictions. *Forests* **14**,
2151 (2023).

27. George, J.-P., Sanders, T. G. M., Timmermann, V., Potočić, N. & Lang, M. European-wide
forest monitoring substantiate the neccessity for a joint conservation strategy to rescue
European ash species (*Fraxinus spp.*). *Sci. Rep.* **12**, 4764 (2022).

28. Hill, L. *et al.* The £15 billion cost of ash dieback in Britain. *Curr. Biol.* **29**, R315–R316
(2019).

29. Mitchell, R. J. *et al.* Cumulative impact assessments of multiple host species loss from plant
diseases show disproportionate reductions in associated biodiversity. *J. Ecol.* **110**, 221–231
(2022).

30. Combes, M., Webber, J. & Boddy, L. Current understanding and future prospects for ash
dieback disease with a focus on Britain. *Forestry (Lond.)* **97**, 678–691 (2024).

31. Fellenor, J. *et al.* 'I'd like to report a suspicious looking tree': Public concern, public
attention and the nature of reporting about ash dieback in the United Kingdom. *Public*
Underst. Sci. **28**, 339–356 (2019).

32. Jepson, P. R. & Arakelyan, I. Developing publicly acceptable tree health policy: public
perceptions of tree-breeding solutions to ash dieback among interested publics in the UK.
For. Policy Econ. **80**, 167–177 (2017).

33. Rallapalli, G. *et al.* Lessons from Fraxinus, a crowd-sourced citizen science game in
genomics. *Elife* **4**, e07460 (2015).

34. Tomlinson, I. The discovery of ash dieback in the UK: the making of a focusing event. *Env.*
Polit. **25**, 709–728 (2016).

35. Clark, J. Clark 2024. The Living Ash Project - 10 years on. *ResearchGate* (2024).

36. Stocks, J. J., Buggs, R. J. A. & Lee, S. J. A first assessment of *Fraxinus excelsior* (common
ash) susceptibility to *Hymenoscyphus fraxineus* (ash dieback) throughout the British Isles.
Sci. Rep. **7**, 16546 (2017).

37. Pliūra, A., Lygis, V., Suchockas, V. & Bartkevičius, E. Performance of twenty four European
Fraxinus excelsior populations in three Lithuanian progeny trials with a special emphasis on
resistance to *Chalara fraxinea*. *Baltic Forestry* **17**, 17–163 (2011).

38. McKinney, L. V., Nielsen, L. R., Hansen, J. K. & Kjær, E. D. Presence of natural genetic
resistance in *Fraxinus excelsior* (Oleraceae) to *Chalara fraxinea* (Ascomycota): an
emerging infectious disease. *Heredity (Edinb.)* **106**, 788–797 (2011).

39. Seidel, H., Šeho, M. & Fussi, B. Hope for ash conservation and propagation—single
individuals can be highly resistant to an invasive pathogen. *J. Plant Dis. Prot. (2006)* **132**,
1–15 (2025).

40. Plumb, W. J. *et al.* The viability of a breeding programme for ash in the British Isles in the
face of ash dieback. *Plants People Planet* **2**, 29–40 (2020).

41. Sniezko, R. A. & Koch, J. Breeding trees resistant to insects and diseases: putting theory
into application. *Biol. Invasions* **19**, 3377–3400 (2017).

42. Stocks, J. J. *et al.* Genomic basis of European ash tree resistance to ash dieback fungus.
Nat Ecol Evol **3**, 1686–1696 (2019).

43. Meger, J., Ulaszewski, B., Pałucka, M., Kozioł, C. & Burczyk, J. Genomic prediction of
resistance to *Hymenoscyphus fraxineus* in common ash (*Fraxinus excelsior* L.) populations.
Evol. Appl. **17**, e13694 (2024).

44. Doonan, J. M. *et al.* Multiple, single trait GWAS and supervised machine learning reveal the

genetic architecture of *Fraxinus excelsior* tolerance to ash dieback in Europe. *Plant Cell*

Environ. **48**, 3793–3809 (2025).

45. Szukala, A. *et al.* Limitations of ad hoc genotyping in detecting ash dieback tolerance in

Fraxinus excelsior. Tree Genet. Genomes **21**, 2024.10.11.617393 (2025).

46. Metheringham, C. L. *et al.* Rapid polygenic adaptation in a wild population of ash trees

under a novel fungal epidemic. *Science* **388**, 1422–1425 (2025).

47. McKinney, L. V. *et al.* The ash dieback crisis: genetic variation in resistance can prove a

long□term solution. *Plant Pathol.* **63**, 485–499 (2014).

48. Liu, J. N. *et al.* Pan-genome analyses of 11 *Fraxinus* species provide insights into salt

adaptation in ash trees. *Plant Commun.* **0**, 101137 (2024).

49. Sax, K. & Abbe, E. C. Chromosome numbers and the anatomy of the secondary xylem in

the oleaceae. *J. Arnold Arbor.* (1932) doi:10.5962/p.185244.

50. Franco Ortega, S. *et al. Fraxinus excelsior* updated long-read genome reveals the

importance of MADS-box genes in tolerance mechanisms against ash dieback. *G3*

(Bethesda) **15**, 2024.12.20.629733 (2025).

51. Huff, M. *et al.* A high-quality reference genome for *Fraxinus pennsylvanica* for ash species

restoration and research. *Mol. Ecol. Resour.* **22**, 1284–1302 (2022).

52. Julca, I., Marcet-Houben, M., Vargas, P. & Gabaldón, T. Phylogenomics of the olive tree

(*Olea europaea*) reveals the relative contribution of ancient allo- and autopolyploidization

events. *BMC Biol.* **16**, 15 (2018).

53. Sollars, E. S. A. *et al.* Genome sequence and genetic diversity of European ash trees.

Nature **541**, 212–216 (2017).

54. Siljak-Yakovlev, S., Temunović, M., Robin, O., Raquin, C. & Frascaria-Lacoste, N.

Molecular–cytogenetic studies of ribosomal RNA genes and heterochromatin in three

European *Fraxinus* species. *Tree Genet. Genomes* **10**, 231–239 (2014).

55. Ranallo-Benavidez, T. R., Jaron, K. S. & Schatz, M. C. GenomeScope 2.0 and Smudgeplot

for reference-free profiling of polyploid genomes. *Nat. Commun.* **11**, 1432 (2020).

56. Ou, S. & Jiang, N. LTR_retriever: A highly accurate and sensitive program for identification

of long terminal repeat retrotransposons. *Plant Physiol.* **176**, 1410–1422 (2018).

57. Rao, G. *et al.* De novo assembly of a new *Olea europaea* genome accession using

nanopore sequencing. *Hortic Res* **8**, 64 (2021).

58. Jouffroy, O., Saha, S., Mueller, L., Quesneville, H. & Maumus, F. Comprehensive

repeatome annotation reveals strong potential impact of repetitive elements on tomato

ripening. *BMC Genomics* **17**, 624 (2016).

59. Su, X. *et al.* A high-continuity and annotated tomato reference genome. *BMC Genomics* **22**,

898 (2021).

60. Gabriel, L. *et al.* BRAKER3: Fully automated genome annotation using RNA-Seq and

protein evidence with GeneMark-ETP, AUGUSTUS and TSEBRA. *bioRxiv*

doi:10.1101/2023.06.10.544449.

61. Emms, D. M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biol.* **20**, 238 (2019).

62. Smolka, M. *et al.* Detection of mosaic and population-level structural variants with Sniffles2.

Nat. Biotechnol. (2024) doi:10.1038/s41587-023-02024-y.

63. Jiang, T. *et al.* Long-read-based human genomic structural variation detection with cuteSV.

Genome Biol. **21**, 189 (2020).

64. Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads

using repeat graphs. *Nat. Biotechnol.* **37**, 540–546 (2019).

65. Shafin, K. *et al.* Nanopore sequencing and the Shasta toolkit enable efficient de novo

assembly of eleven human genomes. *Nat. Biotechnol.* **38**, 1044–1053 (2020).

66. Hu, J. *et al.* NextDenovo: an efficient error correction and accurate assembly tool for noisy

long reads. *Genome Biol.* **25**, 107 (2024).

67. Heller, D. & Vingron, M. SVIM-asm: structural variant detection from haploid and diploid

genome assemblies. *Bioinformatics* **36**, 5519–5521 (2021).

68. Sirén, J. *et al.* Pangenomics enables genotyping of known structural variants in 5202

diverse genomes. *Science* **374**, abg8871 (2021).

69. Stener, L.-G. Clonal differences in susceptibility to the dieback of *Fraxinus excelsior* in

southern Sweden. *Scand. J. For. Res.* **28**, 205–216 (2013).

70. Landolt, J., Gross, A., Holdenrieder, O. & Pautasso, M. Ash dieback due to

Hymenoscyphus fraxineus: what can be learnt from evolutionary ecology? *Plant Pathol.* **65**,

1056–1070 (2016).

71. Kofler, R., Pandey, R. V. & Schlötterer, C. PoPoolation2: identifying differentiation between

populations using sequencing of pooled DNA samples (Pool-Seq). *Bioinformatics* **27**,

3435–3436 (2011).

72. Mérot, C. *et al.* Genome assembly, structural variants, and genetic differentiation between

lake whitefish young species pairs (*Coregonus* sp.) with long and short reads. *Mol. Ecol.*

32, 1458–1477 (2023).

73. Cochran, W. G. Some methods for strengthening the common χ 2 tests. *Biometrics* **10**, 417

(1954).

74. Mantel, N. & Haenszel, W. Statistical aspects of the analysis of data from retrospective

studies of disease. *J. Natl. Cancer Inst.* **22**, 719–748 (1959).

75. Wiberg, R. A. W., Gaggiotti, O. E., Morrissey, M. B. & Ritchie, M. G. Identifying consistent

allele frequency differences in studies of stratified populations. *Methods Ecol. Evol.* **8**,

1899–1909 (2017).

76. Micheletti, S. J. & Narum, S. R. Utility of pooled sequencing for association mapping in

nonmodel organisms. *Mol. Ecol. Resour.* **18**, 825–837 (2018).

77. Gazda, M. A. *et al.* A genetic mechanism for sexual dichromatism in birds. *Science* **368**,

1270–1274 (2020).

78. Brennan, R. S. *et al.* Experimental evolution reveals the synergistic genomic mechanisms

of adaptation to ocean warming and acidification in a marine copepod. *Proc. Natl. Acad.*

Sci. U. S. A. **119**, e2201521119 (2022).

79. Audet, T., Krol, J., Pelletier, K., Stewart, A. D. & Dworkin, I. Sexually discordant selection is

associated with trait-specific morphological changes and a complex genomic response.

Evolution **78**, 1426–1440 (2024).

80. Woolf, B. On estimating the relation between blood group and disease. *Ann. Hum. Genet.*

19, 251–253 (1955).

81. Higgins, J. P. T., Thompson, S. G., Deeks, J. J. & Altman, D. G. Measuring inconsistency in

meta-analyses. *BMJ* **327**, 557–560 (2003).

82. Ou, L. *et al.* Pan-genome of cultivated pepper (*Capsicum*) and its use in gene presence-

absence variation analyses. *New Phytol.* **220**, 360–363 (2018).

83. Barchi, L. *et al.* Improved genome assembly and pan-genome provide key insights into

eggplant domestication and breeding. *Plant J.* **107**, 579–596 (2021).

84. Li, H. *et al.* Graph-based pan-genome reveals structural and sequence variations related to

agronomic traits and domestication in cucumber. *Nat. Commun.* **13**, 682 (2022).

85. Liang, L. *et al.* Genome and pan-genome assembly of asparagus bean (*Vigna unguiculata*

ssp. sesquipedialis) reveal the genetic basis of cold adaptation. *Front. Plant Sci.* **13**,

1059804 (2022).

86. He, Q. *et al.* A graph-based genome and pan-genome variation of the model plant *Setaria*.

Nat. Genet. **55**, 1232–1242 (2023).

87. Chen, J. *et al.* Pangenome analysis reveals genomic variations associated with

domestication traits in broomcorn millet. *Nat. Genet.* **55**, 2243–2254 (2023).

88. Lyu, X. *et al.* Pan-genome analysis sheds light on structural variation-based dissection of

agronomic traits in melon crops. *Plant Physiol.* **193**, 1330–1348 (2023).

89. He, X. *et al.* Pangenome analysis reveals transposon-driven genome evolution in cotton.

BMC Biol. **22**, 92 (2024).

90. Guo, M. *et al.* Analyzes of pan-genome and resequencing atlas unveil the genetic basis of

jujube domestication. *Nat. Commun.* **15**, 9320 (2024).

91. Fang, Y. *et al.* Pan-genome and phylogenomic analyses highlight *Hevea* species

delineation and rubber trait evolution. *Nat. Commun.* **15**, 7232 (2024).

92. Lian, Q. *et al.* A pan-genome of 69 *Arabidopsis thaliana* accessions reveals a conserved

genome structure throughout the global species range. *Nat. Genet.* **56**, 982–991 (2024).

93. Guo, N. *et al.* A graph-based pan-genome of *Brassica oleracea* provides new insights into

its domestication and morphotype diversification. *Plant Commun.* **5**, 100791 (2024).

94. Liang, Q. *et al.* A view of the pan-genome of domesticated Cowpea (V*igna unguiculata* [L.]

Walp.). *Plant Genome* **17**, e20319 (2024).

95. Glick, L. & Mayrose, I. The effect of methodological considerations on the construction of

gene-based plant pan-genomes. *Genome Biol. Evol.* **15**, evad121 (2023).

96. Zhao, K. *et al.* Pangenome analysis reveals structural variation associated with seed size

and weight traits in peanut. *Nat. Genet.* **57**, 1250–1261 (2025).

97. Hou, Y. *et al.* Haplotype-based pangenomes reveal genetic variations and climate

adaptations in moso bamboo populations. *Nat. Commun.* **15**, 8085 (2024).

98. Meng, Q. *et al.* Pangenome analysis reveals yield- and fiber-related diversity and

interspecific gene flow in *Gossypium barbadense* L. *Nat. Commun.* **16**, 4995 (2025).

99. Brůna, T., Sreedasyam, A., Harder, A. M. & Lovell, J. T. Evolutionary and methodological

considerations when interpreting gene presence-absence variation in pangenomes. *NAR*

Genom. Bioinform. **8**, lqag011 (2026).

100.Ebler, J. *et al.* Pangenome-based genome inference allows efficient and accurate

genotyping across a wide spectrum of variant classes. *Nat. Genet.* **54**, 518–525 (2022).

101.Eggertsson, H. P. *et al.* GraphTyper2 enables population-scale genotyping of structural

variation using pangenome graphs. *Nat. Commun.* **10**, 5402 (2019).

102. Mahmoud, M. *et al.* Structural variant calling: the long and the short of it. *Genome Biol.* **20**,

246 (2019).

103. Schlötterer, C., Tobler, R., Kofler, R. & Nolte, V. Sequencing pools of individuals - mining

genome-wide polymorphism data without big funding. *Nat. Rev. Genet.* **15**, 749–763

(2014).

104. Pellicer, J. & Leitch, I. J. The Plant DNA C-values database (release 7.1): an updated

online repository of plant genome size data for comparative studies. *New Phytol.* **226**, 301–

305 (2020).

105. Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo

assembly using phased assembly graphs with hifiasm. *Nat. Methods* **18**, 170–175 (2021).

106. Putnam, N. H. *et al.* Chromosome-scale shotgun assembly using an *in vitro* method for

long-range linkage. *Genome Res.* **26**, 342–350 (2016).

107. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**, 1754–1760 (2009).

108. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for

genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

109. Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A. & Zdobnov, E. M. BUSCO update:

Novel and streamlined workflows along with broader and deeper phylogenetic coverage for

scoring of eukaryotic, prokaryotic, and viral genomes. *Mol. Biol. Evol.* **38**, 4647–4654

(2021).

110. Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421

(2009).

111. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–

3100 (2018).

112. Challis, R., Richards, E., Rajan, J., Cochrane, G. & Blaxter, M. BlobToolKit - interactive

quality assessment of genome assemblies. *G3 (Bethesda)* **10**, 1361–1374 (2020).

113. Ellinghaus, D., Kurtz, S. & Willhoeft, U. LTRharvest, an efficient and flexible software for de novo detection of LTR retrotransposons. *BMC Bioinformatics* **9**, 18 (2008).

114. Xu, Z. & Wang, H. LTR_FINDER: an efficient tool for the prediction of full-length LTR retrotransposons. *Nucleic Acids Res.* **35**, W265–8 (2007).

115. Xing, J., Hao, J., Tang, C., Xie, S. & Liu, K. A closed-loop method for precise genome size estimation using HiFi reads. *BMC Genomics* **26**, 878 (2025).

116. Zhou, C. *et al.* Oatk: a de novo assembly tool for complex plant organelle genomes. *bioRxiv* 2024.10.23.619857 (2024) doi:10.1101/2024.10.23.619857.

117. Tillich, M. *et al.* GeSeq - versatile and accurate annotation of organelle genomes. *Nucleic Acids Res.* **45**, W6–W11 (2017).

118. Using local stock for planting native trees and shrubs. *Forest Research* https://www.forestresearch.gov.uk/publications/using-local-stock-for-planting-native-trees-and-shrubs/ (1999).

119. Caudullo, G., Welk, E. & San-Miguel-Ayanz, J. Chorological maps for the main European woody species. *Data Brief* **12**, 662–666 (2017).

120. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* **17**, 10 (2011).

121. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

122. De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M. & Van Broeckhoven, C. NanoPack: visualizing and processing long-read sequencing data. *Bioinformatics* **34**, 2666–2669 (2018).

123. De Coster, W. & Rademakers, R. NanoPack2: population-scale evaluation of long-read sequencing data. *Bioinformatics* **39**, btad311 (2023).

124. Andrews, S. Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (2010).

125. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

126. Zheng, Z. *et al.* Symphonizing pileup and full-alignment for deep learning-based long-read
variant calling. *Nat Comput Sci* **2**, 797–803 (2022).

127. Lin, M. F. *et al.* GLnexus: joint variant calling for large cohort sequencing. *bioRxiv* 343970
(2018) doi:10.1101/343970.

128. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

129. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer
datasets. *Gigascience* **4**, 7 (2015).

130. Zhou, A., Lin, T. & Xing, J. Evaluating nanopore sequencing data processing pipelines for
structural variation identification. *Genome Biol.* **20**, 237 (2019).

131. De Coster, W. *et al.* Structural variants identified by Oxford Nanopore PromethION
sequencing of the human genome. *Genome Res.* **29**, 1178–1187 (2019).

132. Bolognini, D. & Magi, A. Evaluation of Germline Structural Variant Calling Methods for
Nanopore Sequencing Data. *Front. Genet.* **12**, 761791 (2021).

133. Yildiz, G. *et al.* Benchmarking Oxford Nanopore Read Alignment-Based Structural Variant
Detection Tools in Crop Plant Genomes. *bioRxiv* 2022.09.23.508909 (2022)
doi:10.1101/2022.09.23.508909.

134. Jiang, T. *et al.* Long-read sequencing settings for efficient structural variation detection
based on comprehensive evaluation. *BMC Bioinformatics* **22**, 552 (2021).

135. Duan, X., Pan, M. & Fan, S. Comprehensive evaluation of structural variant genotyping
methods based on long-read sequencing data. *BMC Genomics* **23**, 324 (2022).

136. Liu, Y. *et al.* Comparison of structural variants detected by PacBio-CLR and ONT
sequencing in pear. *BMC Genomics* **23**, 830 (2022).

137. Dierckxsens, N., Li, T., Vermeesch, J. R. & Xie, Z. A benchmark of structural variation
detection by long reads through a realistic simulated model. *Genome Biol.* **22**, 342 (2021).

138. Jeffares, D. C. *et al.* Transient structural variations have strong effects on quantitative traits

and reproductive isolation in fission yeast. *Nat. Commun.* **8**, 14061 (2017).

139. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* **26**, 841–842 (2010).

140. Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable

element families. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9451–9457 (2020).

141. Tarailo-Graovac, M. & Chen, N. Using RepeatMasker to identify repetitive elements in

genomic sequences. *Curr. Protoc. Bioinformatics* **Chapter 4**, Unit 4.10 (2009).

142. Formenti, G. *et al.* Gfastats: conversion, evaluation and manipulation of genome sequences

using assembly graphs. *Bioinformatics* **38**, 4214–4216 (2022).

143. Doonan, J. M. *et al.* Transcriptome profiling of *Fraxinus excelsior* genotypes infested by

emerald ash borer. *Sci. Data* **10**, 680 (2023).

144. Chano, V. *et al.* Transcriptional time-course analysis during ash dieback infection revealed

different responses in tolerant and susceptible *Fraxinus excelsior genotypes. BMC Plant*

Biol. **25**, 107 (2025).

145. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

146. Lovell, J. T. *et al.* The genomic landscape of molecular responses to natural drought stress

in *Panicum hallii. Nat. Commun.* **9**, 5213 (2018).

147. Kuznetsov, D. *et al.* OrthoDB v11: annotation of orthologs in the widest sampling of

organismal diversity. *Nucleic Acids Res.* **51**, D445–D451 (2023).

148. Unver, T. *et al.* Genome of wild olive and the evolution of oil biosynthesis. *Proc. Natl. Acad.*

Sci. U. S. A. **114**, E9413–E9422 (2017).

149. Tomato Genome Consortium. The tomato genome sequence provides insights into fleshy

fruit evolution. *Nature* **485**, 635–641 (2012).

150. Hellsten, U. *et al.* Fine-scale variation in meiotic recombination in *Mimulus* inferred from

population shotgun sequencing. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 19478–19482 (2013).

151. Wang, L. *et al.* Genome sequencing of the high oil crop sesame provides insight into oil

biosynthesis. *Genome Biol.* **15**, R39 (2014).

152. Pan, X. *et al.* Chromosome-level genome assembly of *Salvia miltiorrhiza* with orange roots

uncovers the role of Sm2OGD3 in catalyzing 15,16-dehydrogenation of tanshinones. *Hortic.*

Res. **10**, uhad069 (2023).

153. Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life scale

using DIAMOND. *Nat. Methods* **18**, 366–368 (2021).

154. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics*

30, 1236–1240 (2014).

155. Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.

EggNOG-mapper v2: Functional annotation, orthology assignments, and domain prediction

at the metagenomic scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).

156. Alexa A, R. J. *topGO: Enrichment Analysis for Gene Ontology*.

http://bioconductor.org/packages/topGO/ (2024).

157. Garrison, E., Kronenberg, Z. N., Dawson, E. T., Pedersen, B. S. & Prins, P. A spectrum of

free software tools for processing the VCF variant call format: vcflib, bio-vcf, cyvcf2, hts-nim

and slivar. *PLoS Comput. Biol.* **18**, e1009123 (2022).

158. Purcell, S. *et al.* PLINK: a tool set for whole-genome association and population-based

linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

159. Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A cross-platform and ultrafast toolkit for FASTA/Q

file manipulation. *PLoS One* **11**, e0163962 (2016).

160. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing, Vienna, Austria. (2023).

161. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–

2079 (2009).

162. vcd: Visualizing Categorical Data. https://cran.r-universe.dev/vcd.

163.Kossowsky, J. *et al.* The separation anxiety hypothesis of panic disorder revisited: a meta-

analysis. *Am. J. Psychiatry* **170**, 768–781 (2013).

1666