Conservation and divergence of UVR8-COP1/SPA-HY5 signaling in UV-B responses of

Marchantia polymorpha

Yuanke Liang[1,‖,#], Roman Podolec[1,¶,#], Richard Chappuis[1], Emmanuel Defossez[2], Gaétan Glauser[3],

Johannes Rötzer[4,5], Sara Christina Stolze[6], Liam Dolan[4], Hirofumi Nakagami[6], Emilie Demarsy[1],

Roman Ulm[1,7,*]

[1]Department of Plant Sciences, Section of Biology, Faculty of Sciences, University of Geneva,

1211 Geneva 4, Switzerland

[2]Institute of Biology, University of Neuchâtel, 2000 Neuchâtel, Switzerland

[3]Neuchâtel Platform of Analytical Chemistry, University of Neuchâtel, 2000 Neuchâtel,

Switzerland

[4]Gregor Mendel Institute of Molecular Plant Biology (GMI), Austrian Academy of Sciences, 1030

Vienna, Austria

[5]Vienna BioCenter PhD Program, Doctoral School of the University of Vienna and Medical

University of Vienna, 1030 Vienna, Austria

[6]Max-Planck Institute for Plant Breeding Research, D-50829, Cologne, Germany

[7]Institute of Genetics and Genomics of Geneva (iGE3), University of Geneva, 1211 Geneva 4,

Switzerland

[‖]Present address: Max-Planck-Institute of Molecular Plant Physiology, Potsdam-Golm, Germany

[¶]Present address: Whitehead Institute for Biomedical Research, Cambridge, MA, USA

[#]Y.L. and R.P. contributed equally to this work.

*Author for correspondence: roman.ulm@unige.ch (R.U.)

ABSTRACT

Ultraviolet-B radiation (UV-B) poses a major challenge to all forms of plant life. The liverwort

Marchantia polymorpha (Marchantia) serves as a key model organism to study signaling pathways

and to infer their evolution throughout the green lineage. Marchantia expresses key components of

UV-B signaling, including the photoreceptor UV RESISTANCE LOCUS 8 (MpUVR8), the

WD40-repeat protein REPRESSOR OF UV-B PHOTOMORPHOGENESIS (MpRUP), the E3 ubiquitin ligase complex CONSTITUTIVELY PHOTOMORPHOGENIC 1 / SUPPRESSOR OF phyA-105 (MpCOP1/MpSPA), and the transcriptional regulator ELONGATED HYPOCOTYL 5 (MpHY5). Here, we show that MpUVR8 exists as a homodimer in its ground-state in vivo, then monomerizes and accumulates in the nucleus upon UV-B activation. Activated MpUVR8 interacts with MpCOP1, triggering growth inhibition, gene expression changes, biosynthesis of UV-absorbing metabolites, and photoprotection, which overall contributes to UV-B stress tolerance. MpRUP facilitates redimerization of MpUVR8 and Mp*rup* null mutants show enhanced UV-B photomorphogenesis, indicating that MpRUP efficiently represses MpUVR8 signaling. Unlike the case in Arabidopsis and in contrast to the strong Mp*cop1* mutant phenotype, Mp*spa* mutants develop only a weak constitutive photomorphogenesis phenotype, indicating that COP1 function is more independent of SPA in Marchantia than in Arabidopsis. Interestingly, however, Mp*spa* is linked with a hyper-responsive UV-B phenotype, suggesting that MpSPA is a negative regulator of MpUVR8 signaling. Our findings demonstrate that core components of UV-B signaling existed in the last common ancestor of extant land plants; however, regulatory interactions have diversified in different lineages since their divergence more than 400 million years ago.

INTRODUCTION

Photoautotrophic plants have a "love/hate relationship" with light. Whereas plant life is unable to survive with too little light, excessive sunlight also poses a major challenge, in part because it contains potentially damaging ultraviolet-B radiation (UV-B; 280–315 nm) (Demarsy et al., 2018). Plants have a unique and conserved UV-B photoreceptor UV RESISTANCE LOCUS 8 (UVR8) that is crucially important for UV-B acclimation, tolerance, and survival (Kliebenstein et al., 2002; Rizzini et al., 2011; Tilbrook et al., 2016; Li et al., 2018; Soriano et al., 2018; Rai et al., 2019; Podolec et al., 2021; Hu et al., 2024; Stockenhuber et al., 2024). The UVR8 signaling pathway has mainly been outlined in Arabidopsis, with UVR8 being a homodimer in its ground state, which exhibits monomerization after absorption of UV-B photons by intrinsic tryptophan (Trp) residues, followed by accumulation in the nucleus (Kaiserli and Jenkins, 2007; Rizzini et al., 2011; Christie et al., 2012; Wu et al., 2012; Yin et al., 2016; Li et al., 2020; Qian et al., 2020; Podolec et al., 2021; Fang et al., 2022). Activated UVR8 then interacts with CONSTITUTIVELY PHOTOMORPHOGENIC 1 (COP1) (Favory et al., 2009; Yin et al., 2015; Wang et al., 2022).

COP1 forms an E3 ubiquitin ligase complex together with SUPPRESSOR OF phyA-105 proteins

(SPA1–4 in Arabidopsis) (Laubinger et al., 2004; Chen et al., 2010). The COP1/SPA complex

represses photomorphogenesis by ubiquitinating photomorphogenesis-promoting factors, directing

them to degradation by the 26S proteasome (Podolec and Ulm, 2018; Ponnu and Hoecker, 2021).

UV-B-activated UVR8 inhibits the activity of the COP1-SPA complex through a cooperative, high-

affinity binding mechanism that displaces substrate proteins from COP1, thus allowing the

stabilization of key transcription factors, including the basic leucine zipper (bZIP) transcriptional

regulators ELONGATED HYPOCOTYL 5 (HY5) and HY5 HOMOLOG (HYH) (Holm et al.,

2002; Ulm et al., 2004; Oravecz et al., 2006; Binkert et al., 2014; Lau et al., 2019). This ultimately

leads to broad changes in gene expression upon UV-B exposure (Brown et al., 2005; Favory et al.,

2009; Tavridou et al., 2020). COP1/SPA together with HY5/HYH thus represents a major

regulatory hub in light signaling in Arabidopsis (Holm et al., 2002; Ulm et al., 2004; Ponnu et al.,

2019). In Arabidopsis, the two WD-40 repeat proteins REPRESSOR OF UV-B

PHOTOMORPHOGENESIS 1 (RUP1) and RUP2 act as negative feedback regulators of the UVR8

photocycle by facilitating UVR8 redimerization, thereby repressing UV-B signaling (Gruber et al.,

2010; Heijde and Ulm, 2013; Podolec et al., 2021; Wang et al., 2023).

The core of UVR8 signaling seems to be evolutionarily conserved in the green lineage,

since it is present in the alga *Chlamydomonas reinhardtii* (hereafter Chlamydomonas) (Allorent et

al., 2016; Tilbrook et al., 2016; Tokutsu et al., 2019; Dannay et al., 2025) and the liverwort

Marchantia polymorpha (hereafter Marchantia) (Clayton et al., 2018; Soriano et al., 2018; Kondou

et al., 2019). Notably, Marchantia has minimal gene redundancy in many signaling pathways, and

many studies in the past decade have contributed to our understanding of the origins and evolution

of major plant regulatory pathways (Bowman et al., 2017; Liang et al., 2022; Frangedakis et al.,

2024). The Marchantia genome indeed possesses single homologs for genes encoding the known

UVR8 signaling-related components, such as MpUVR8, MpRUP, MpCOP1, MpSPA, and MpHY5.

Although a few studies have explored the role of some of these components in UV-B (Clayton et

al., 2018; Soriano et al., 2018; Kondou et al., 2019) and visible light signaling (Zhang et al., 2021;

Yelina et al., 2024), a deeper understanding of their activities and roles in UV-acclimation is

warranted because it remains largely unknown to what extent the precise molecular mechanisms

by which these components mediate UV acclimation and tolerance are conserved or divergent

throughout the plant kingdom, thus limiting our ability to fully explain how plants sense, respond

to, and survive in environments with fluctuating UV levels.

We investigated if the major UVR8 signaling components have functionally diverged

during plant evolution, and how mechanisms underlying UV-B tolerance may have evolved during

the colonization of land by plants. We extend our understanding of the crucial role of MpUVR8 in

UV-B acclimation and survival in Marchantia, and provide unequivocal evidence clarifying

conservation of the UVR8 photocycle in Marchantia: MpUVR8 forms UV-B-responsive

homodimers, then monomerizes, interacts with MpCOP1, and accumulates in nuclei in response to

UV-B. MpRUP facilitates MpUVR8 redimerization to repress signaling, with Mp*rup* mutants

showing extreme dwarfism specifically under UV-B. Although Mp*cop1* mutants exhibited

enhanced constitutive photomorphogenesis and extreme dwarfism comparable to UV-B-grown

Mp*rup* mutants, Mp*spa* mutants showed only a weak constitutive photomorphogenesis phenotype

in visible light; however, Mp*spa* mutants were surprisingly hyper-responsive to UV-B, suggesting

that MpSPA plays a divergent role from that in Arabidopsis. Notably, our transcriptome data

revealed that only a few UV-B-regulated genes were entirely MpHY5-dependent. This is different

from Arabidopsis, where HY5/HYH serves as a major regulator in the UV-B signaling pathway.

Moreover, we provide a genome-wide view of MpUVR8-mediated transcriptional responses in

Marchantia and identify UV-B-regulated gene clusters and metabolites key to UV-B protection.

RESULTS

MpUVR8 is homodimeric in its ground state, and monomerizes and accumulates in nuclei in response to UV-B

In Marchantia, UVR8 is encoded by a single locus, but alternative splicing of the 11th intron

produces two transcripts, namely Mp*UVR8.1* and Mp*UVR8.2*, resulting in two different proteins

whereby residues 414 to 426 present in MpUVR8.1 are absent in MpUVR8.2 (Supplementary Fig.

S1) (Soriano et al., 2018). Whether MpUVR8 forms homodimers *in vivo* or is primarily monomeric

with a divergent inactivation mechanism is unclear (Soriano et al., 2018). We thus first tested if

MpUVR8.1 and MpUVR8.2 form dimers using a luciferase complementation imaging (LCI) assay

in transiently transformed *Nicotiana benthamiana* leaves (Zhou et al., 2018). Resulting LCI signals

showed interactions of MpUVR8.1 and MpUVR8.2 both with and between themselves, indicating

that MpUVR8 forms homodimers *in vivo* (Fig. 1A to C). In AtUVR8, the aspartate-96 (D96) and

aspartate-107 (D107) residues are key to homodimer formation and mutating each to asparagine impairs UVR8^{D96N,D107N} dimer formation (Christie et al., 2012; Wu et al., 2012; Heilmann et al., 2016; Podolec et al., 2021). Consistent with the conservation of these residues in MpUVR8 (Supplementary Fig. S1), no LCI signal was detected with the mutated MpUVR8.1^{D95N,D106N} and MpUVR8.2^{D95N,D106N} variants (Fig. 1A and B), indicating that these residues are also key for MpUVR8 homodimerization.

To further test if endogenous MpUVR8 forms homodimers, we crosslinked protein extracts from Marchantia with dithiobis (succinimidyl propionate) (DSP), as described previously for Arabidopsis (Rizzini et al., 2011; Podolec et al., 2021). Using a specific anti-MpUVR8 antibody, we detected a band at the size of the predicted MpUVR8 dimer at about 100 kDa in the absence of UV-B, and this band was strongly reduced in extracts from UV-B-treated plants (Fig. 1D). These data indicate that UV-B exposure induces MpUVR8 monomerization. Of note, UV-B-activated monomeric MpUVR8 was not well detected in crosslinked extracts, as in Arabidopsis (Rizzini et al., 2011; Podolec et al., 2021). However, we detected MpUVR8 at its monomeric size of about 48 kDa (predicted sizes for MpUVR8.1 and MpUVR8.2 are 48.6 and 47.1 kDa, respectively) both following the reversal of DSP crosslinking with the reducing agent β-mercaptoethanol (β-ME) and in the absence of crosslinking (Fig. 1D). This indicates that MpUVR8 levels are not affected by UV-B treatment or crosslinking.

To define MpUVR8 subcellular localization in Marchantia, we transformed wild-type Tak-1 with the full genomic Mp*UVR8* locus containing an N-terminal green fluorescent protein (GFP) tag and imaged the resulting fluorescence. GFP-MpUVR8 accumulated in the nucleus only when plants were exposed to UV-B (Fig. 1E), similar to AtUVR8 (Kaiserli and Jenkins, 2007; Yin et al., 2016) and the 35S promoter–driven expression of Citrine-MpUVR8 in Tak-1 (Kondou et al., 2019). This contrasts with the apparent constitutive nuclear localization of GFP-MpUVR8.1 and GFP-MpUVR8.2 when expressed with the 35S promoter in Arabidopsis and *N. benthamiana* (Soriano et al., 2018).

Taken together, we conclude that endogenous MpUVR8.1 and MpUVR8.2 form homodimers in Marchantia and possess conserved aspartate residues necessary for MpUVR8 homodimer formation. Moreover, upon UV-B exposure, MpUVR8 monomerizes and accumulates in nuclei.

MpRUP facilitates MpUVR8 redimerization

We tested if MpUVR8 interacts with MpRUP, the putative ortholog of AtRUP1 and AtRUP2 (Gruber et al., 2010; Clayton et al., 2018). MpUVR8 and MpRUP exhibited interaction in yeast two-hybrid (Fig. 2A and B), LCI assays in *N. benthamiana* (Fig. 2C and D), and MpRUP co-purified with GFP-MpUVR8 in affinity purification coupled with mass spectrometry (AP-MS) (Supplementary Fig. S2). Using immunoblotting of crosslinked protein extracts, we assayed if MpRUP facilitates MpUVR8 redimerization in Marchantia. In wild-type Tak-1 plants, MpUVR8 dimer amount was strongly reduced upon exposure to UV-B, then reverted to initial levels within approximately 1 h once under -UV-B conditions (Fig. 2E). However, in a Mp*rup* mutant generated using CRISPR/Cas9 mutagenesis in the Tak-1 background (Supplementary Fig. S3), MpUVR8 dimer levels recovered slowly and did not reach "-UV" ground state levels even after 6 h (Fig. 2E). We thus conclude that MpRUP facilitates MpUVR8 redimerization, indicating that there is a conserved mechanism for negative feedback regulation of UVR8 activity.

MpUVR8 and MpRUP antagonistically regulate UV-B-dependent growth, pigment accumulation, and acclimation

To explore the phenotypic consequences of MpUVR8 and MpRUP action, we established robust UV-B conditions for phenotypic and molecular analysis. Under continuous white light, no striking differences were detectable between Tak-1, Mp*uvr8*, and Mp*rup* (Fig. 3A, and Supplementary Fig. S4A). However, when exposed to white light supplemented with narrowband UV-B (1.5 μmol m^{-2} s^{-1}), Tak-1 exhibited a slight growth inhibition and slight accumulation of pigments (Fig. 3A, and Supplementary Fig. S4A). These responses were reduced, if not absent, in Mp*uvr8* mutant and Mp*RUP^OE* overexpression lines (Fig. 3A, and Supplementary Fig. S4A and F). By sharp contrast, Mp*rup* mutants exhibited a striking UV-B-hyperresponsive photomorphogenic phenotype, with extreme growth inhibition and strong pigmentation of the thallus specifically under UV-B (Fig. 3A, and Supplementary Fig. S4A) (Clayton et al., 2018), similar to transgenic lines overexpressing Mp*UVR8.1* or Mp*UVR8.2* photoreceptors (Supplementary Fig. S4A and E). We conclude that MpRUP plays a major role in MpUVR8 inactivation through redimerization and that Mp*rup* mutants show an extreme, UV-B-conditional growth phenotype.

To directly assess the effect of MpUVR8-mediated acclimation on photoprotection, we measured F_v/F_m representing the maximum quantum yield of photosystem II (PSII) (Murchie and

Lawson, 2013). Tak-1, Mp*uvr8*, and Mp*rup* plants were grown for 7 days under white light supplemented with narrowband UV-B (acclimated; +/-) or not (non-acclimated; -/-) (Fig. 3B). Acclimated and non-acclimated plants were then exposed to 1 h of broadband UV stress (+/+ and -/+, respectively). We measured F_v/F_m before and after UV stress in both acclimated and non-acclimated plants (Fig. 3B). Non-stressed Tak-1, Mp*uvr8*, and Mp*rup* displayed an F_v/F_m value close to 0.8 (Fig. 3C), which is characteristic of healthy Marchantia plants (Messant et al., 2023). When exposed to UV stress, F_v/F_m decreased significantly and to the same extent in non-acclimated plants of all genotypes (Fig. 3C), suggesting photoinhibition resulting from PSII damage. However, applying the same UV stress treatment on acclimated plants revealed a UVR8-dependent photoprotective effect: in Tak-1 and even more so in Mp*rup*, F_v/F_m following UV-B stress was higher in acclimated compared to non-acclimated plants (Fig. 3C). This effect was MpUVR8-dependent, as it was absent in Mp*uvr8* (Fig. 3C). Mp*UVR8.1*[OE] and Mp*UVR8.2*[OE] overexpression lines (Supplementary Fig. S4E) and Mp*uvr8-1*/*Pro*[MpUVR8]:*GFP*-Mp*UVR8* complementation lines were similar to Mp*rup*, whereas Mp*RUP*[OE] overexpression lines phenocopied Mp*uvr8* (Supplementary Fig. S4B and F).

Next, we tested the survival rates of Tak-1, Mp*uvr8*, and Mp*rup* under UV-B stress. Plants were acclimated or not under narrowband UV-B for 7 days, followed by a 7- or 9-h broadband UV-B stress treatment, after which all plants were returned to white light for 7 days of recovery (Fig. 3D). Without acclimation, Tak-1, Mp*uvr8*, and Mp*rup* did not survive subsequent UV-B stress (Fig. 3E). Acclimated wild-type plants did not survive the 9-h UV-B stress treatment but some gemmalings resumed growth after 7 h of UV-B stress. Mp*rup* mutants recovered more than Tak-1 under both stress treatments (Fig. 3E, Supplementary Fig. S4C and D), indicating enhanced UV-B protection. By contrast, the Mp*uvr8* mutant did not survive the UV-B stress treatment even with prior UV-B acclimation, demonstrating that UV-B photoprotection was dependent on MpUVR8 (Fig. 3E). Complementing the Mp*uvr8-1* mutant with a *Pro*[MpUVR8]:Mp*UVR8* construct restored UV-B stress survival in acclimated plants (Supplementary Fig. S4C). By contrast, overexpression of MpRUP driven by the strong Mp*EF1α* promoter (Tak-1/Mp*RUP*[OE] and Mp*rup-2*/Mp*RUP*[OE]) phenocopied the Mp*uvr8* mutant phenotype (Supplementary Fig. S4D).

We prepared methanolic extracts from gemmalings and measured the absorption spectra between 280–400 nm to detect UV-absorbing compounds. There was a broad peak in UV-absorbance at ~340 nm for UV-B-exposed Tak-1 plants, which was absent for the Mp*uvr8* mutant

(Fig. 3F), indicating Marchantia plants accumulate UV-absorbing compounds in a UV-B- and UVR8-dependent manner. Consistent with their enhanced UV-B acclimatory response and resulting UV-B tolerance, extracts from UV-B-exposed Mp*rup* mutants had an even greater peak in UV absorption (Fig. 3F). Notably, the Mp*uvr8-1* Mp*rup-2* double mutant exhibited photoprotection and UV-absorbing capacity comparable to the Mp*uvr8-1* single mutant (Supplementary Fig. S5), demonstrating that the enhanced photoprotection and UV-B-absorbing capacity of Mp*rup* is dependent on MpUVR8.

We conclude that MpUVR8 is key in mediating protection of the photosynthetic apparatus and survival under UV-B stress, with MpRUP functioning as a key negative regulator of MpUVR8-dependent responses. Moreover, enhanced photoprotection under UV-B is associated with the accumulation of UV-absorbing compounds.

Mp*cop1* mutants exhibit constitutive photomorphogenesis and UV-B tolerance

Interaction of UVR8 with the COP1/SPA E3 ligase complex and its subsequent inactivation is a key step in UV-B signal transduction (Favory et al., 2009; Yin et al., 2015; Lau et al., 2019). MpCOP1 and MpSPA interacted in yeast two-hybrid assays, both in the absence and presence of UV-B (Fig. 4A) (Zhang et al., 2021). By contrast, MpUVR8 interacted with MpCOP1 specifically under UV-B in yeast two-hybrid (Fig. 4A), LCI (Fig. 4B), and AP-MS assays (Supplementary Fig. S2).

Arabidopsis *cop1* null mutants have a seedling-lethal phenotype (McNellis et al., 1994) and no Marchantia Mp*cop1* mutant has been recovered and described until now (Zhang et al., 2021). We generated two viable Mp*cop1* null mutant alleles using CRISPR/Cas9 mutagenesis (Supplementary Fig. S3). These mutants displayed stunted growth, delayed development, and reduced gemmae production compared to that in wild-type Tak-1 when grown under standard conditions (Fig. 5A). Whereas the Mp*cop1* mutants developed gemma cups and gemmae and thus could complete their asexual life cycle,, they failed to complete their sexual life cycle because antheridiophores did not develop when induced under continuous white light supplemented with far-red light.

Mp*cop1* mutants exhibited constitutively enhanced photoprotection, as evidenced by their elevated F_v/F_m ratios after UV-B stress (Fig. 5B). The strong constitutively photomorphogenic phenotype and enhanced photoprotection of Mp*cop1* mutants agreed with how Mp*cop1* mutants

survived prolonged broadband UV-B stress exposure, both with or without acclimation (Fig. 5C;
note that the plants are viable as determined by non-stressed levels of F_v/F_m after recovery).
Absorption spectra of methanolic extracts also revealed constitutive hyperaccumulation of UV-
absorbing compounds in Mp*cop1* (Fig. 5D). Thus, MpCOP1 activity as key repressor of
photomorphogenesis is evident in Marchantia.

Mp*spa* mutants show only a minor constitutively photomorphogenic growth phenotype, but
strongly enhanced UV-B tolerance
To investigate the role of the single MpSPA protein, we generated two independent CRISPR/Cas9
knockout alleles (Supplementary Fig. S3). Surprisingly, both Mp*spa* alleles exhibited a phenotype
comparable to wild-type Tak-1 under both white light and white light supplemented with UV-B
(Fig. 5A). This was in stark contrast to the strong constitutively photomorphogenic phenotype
observed for the Mp*cop1* mutant (Fig. 5A) and Arabidopsis *spa* quadruple mutants (Ordonez-
Herrera et al., 2015).

Furthermore, we tested the UV-B acclimation response of Mp*spa* mutants. Mp*spa* showed
enhanced basal UV-B tolerance when compared to wild type ("-/+", Fig. 5B), as well as after UV-
B acclimation ("+/+", Fig. 5B). Mp*spa* mutants also showed an increased survival rate after
prolonged UV-B stress if plants were previously acclimated to low UV-B (Fig. 5C). In accordance
with their improved UV-B stress tolerance, Mp*spa* mutants constitutively accumulated more UV-
absorbing compounds which became further pronounced upon UV-B-mediated acclimation (Fig.
5D).

As enhanced UV-B-acclimation of Mp*spa* resembled Mp*rup* mutants, we investigated
whether MpSPA function was also associated with MpUVR8 redimerization. However, in contrast
to Mp*rup*, Mp*spa* showed efficient MpUVR8 redimerization (Supplementary Fig. S6A).
Additionally, overexpression of Mp*SPA* and Mp*COP1* reduced basal UV-B tolerance but did not
abolish UV-B-induced photoprotection, in contrast to Mp*RUP* overexpression (Supplementary Fig.
S6B to D). We conclude that MpCOP1 activity as a repressor of photomorphogenesis is not fully
dependent on MpSPA, and that the absence of MpSPA enhances UV-B tolerance.

MpUVR8 signaling components mediate transcriptional changes in response to UV-B

We analyzed the UVR8-mediated transcriptome in 7-day-old Marchantia gemmalings in response to 3 h of narrowband UV-B (Supplementary Tables S1 and S2). In Tak-1, we identified 480 and 114 genes up- and downregulated by UV-B, respectively ($|$log$_2$ fold-change$|$ > 1; p-adj < 0.05), which all exhibited MpUVR8- and MpCOP1-dependent expression changes (Supplementary Tables S3A, B and S4), thus representing the transcriptional MpUVR8 signaling output.

In wild-type Tak-1, Mp*rup*, and Mp*spa*, we identified a combined total of 756 UV-B-induced genes and 298 UV-B-repressed genes ($|$log$_2$ fold-change$|$ > 1; p-adj < 0.05), with substantial overlaps between the genotypes (Fig. 6A to C, Supplementary Tables S3A, B and S4). Interestingly, induction or repression of transcripts after 3 h of UV-B was enhanced in Mp*rup* and Mp*spa* compared to Tak-1 (Fig. 6C), consistent with their UV-B hyper-responsive phenotypes (Fig. 3 and Fig. 5). Consistent with this, over two-thirds of genes that were up- or downregulated ($|$log$_2$ fold-change$|$ > 1) specifically in Mp*rup* and Mp*spa* ("unique Mp*rup*" and "unique Mp*spa*") were up- or downregulated in Tak-1 as well when considering a lower fold-change threshold ($|$log$_2$ fold-change$|$ > 0 but <1; p-adj < 0.05) (Fig. 6A and B). Notably, whereas the transcriptomic profile of the Mp*cop1-1* mutant was unresponsive to UV-B, it resembled the profile of UV-B-responsive genotypes after UV-B exposure, with a large portion of genes constitutively up- or downregulated (Fig. 6C and D). These data support the role of MpCOP1 as a repressor of UV-B photomorphogenesis and COP1-inactivation by UVR8 as a key mechanism for UV-B-responsive gene expression changes.

Using hierarchical clustering, we defined seven major clusters of gene expression changes upon UV-B exposure (Fig. 6C and D, and Supplementary Table S4). Clusters 1 and 5 contain UV-B-downregulated genes, with gene expression in Mp*cop1* that is constitutively down- or upregulated, respectively. Cluster 7 consists of UV-B-downregulated genes that are more highly expressed in Mp*spa*. Clusters 4 and 6 contain genes that are induced after 3 h of UV-B in wild type, Mp*rup*, and Mp*spa*, but not in Mp*uvr8* and Mp*cop1*, and their constitutive expression in Mp*cop1* is either elevated or comparable in comparison to that in wild type, respectively. Clusters 2 and 3 contain genes induced after 3 h of UV-B exposure in wild type, with hyper-induced expression in Mp*spa* and Mp*rup*, respectively. We profiled the expression of components of the UVR8 pathway in the data: Mp*HY5*, Mp*RUP*, and Mp*COP1* (cluster 2) were upregulated upon UV-B in wild type (Supplementary Table S4), whereas Mp*UVR8* was constitutively expressed.

MpSPA plays a minor role in repressing visible light photomorphogenesis compared to MpCOP1

In Arabidopsis, *spa1234* (*spaQn*) quadruple mutants undergo constitutive photomorphogenesis with a phenotype similar to strong *cop1* mutants (such as *cop1-1*) and stronger than the weak *cop1-4* allele (Fig. 7A) (McNellis et al., 1994; Laubinger et al., 2004; Ordonez-Herrera et al., 2015). However, Marchantia Mp*spa* mutants exhibited only a minor growth phenotype in white light compared to the strongly affected Mp*cop1* mutants (Fig. 7A, Fig. 5A). Thus, we investigated the transcriptome profile Mp*spa* and Mp*cop1* mutants in more detail.

By comparing white light–grown Mp*cop1* and Mp*spa* versus Tak-1, we found that in Mp*cop1*, 1,256 and 1,898 genes were significantly up- or downregulated ($\left|\log_2 \text{fold-change}\right| > 1$; p-adj < 0.05), respectively (Fig. 7B and C). By stark contrast, only 60 genes were upregulated and 15 genes were downregulated in Mp*spa* using the same threshold criteria (Fig. 7B and C, and Supplementary Table S3C and D), which is significantly different from Arabidopsis, where over 7,000 differentially expressed genes (DEGs) were observed between the *spaQ* mutant and the wild type (Pham et al., 2020). Of those genes dysregulated in Mp*spa*, 40 out of 60 and 12 out of 15 genes were also up- and downregulated, respectively, in Mp*cop1*. We further checked if genes dysregulated in Mp*cop1* but not in Mp*spa* were in fact dysregulated in Mp*spa* but at a lower fold-change threshold. Indeed, out of the 1,256 upregulated genes in Mp*cop1*, 486 were shared with Mp*spa* when considering a looser threshold for genes upregulated in Mp*spa* ($\left|\log_2 \text{fold-change}\right| > 0$; p-adj < 0.05; Fig. 7B). Similarly, out of the 1,898 downregulated genes in Mp*cop1*, 349 were also downregulated in Mp*spa* (Fig. 7C). This indicates a larger overlap between genes dysregulated in Mp*cop1* and Mp*spa* mutants in comparison to wild type, but also identifies clear differences between MpCOP1- and MpSPA-dependent genes at the genome-wide level.

A broader examination of gene expression levels across wild type, Mp*cop1*, and Mp*spa* genotypes confirmed that, although the overall profile of Mp*spa* was similar to wild-type, many transcripts had levels that were intermediate between Tak-1 and Mp*cop1* (Fig. 7D, Supplementary Table S5). Using hierarchical clustering, we defined 11 main clusters of gene expression (Fig. 7E), specifically four for Mp*cop1*-upregulated genes (clusters 1, 2, 7 and 10), six for Mp*cop1*-downregulated genes (clusters 3, 4, 5, 6, 8 and 9), and a small cluster of genes specifically upregulated in Mp*spa* (cluster 11). Interestingly, the cluster groups defined by Mp*cop1*-up- and

downregulated patterns of gene expression could be discriminated by the extent to which expression in Mp*spa* aligned with the expression in Mp*cop1* (Fig. 7E).

Finally, since we observed that the regulation of many genes was dependent on MpCOP1 under white light, and we established that MpCOP1 has a conserved role as a repressor of photomorphogenesis, we tested if genes dysregulated in Mp*cop1* were induced by UV-B in UV-B-responsive genotypes. We observed that most genes upregulated under white light in Mp*cop1* were induced by UV-B in Tak-1, Mp*rup*, or Mp*spa* (Fig. 7D), whereas many genes constitutively downregulated in Mp*cop1* were repressed by UV-B. However, some of the Mp*cop1*-downregulated genes were induced after 3 hours of UV-B, suggesting more complex dynamics between induction under UV-B and steady-state levels in Mp*cop1*.

Overall, our analysis shows that MpCOP1 plays a major role in repressing the light-inducible transcriptome of Marchantia, whereas MpSPA plays a minor yet detectable role, likely as part of the repressive COP1-SPA complex (Zhang et al., 2021).

MpHY5 promotes MpUVR8 signaling

The induction of Mp*HY5* expression is UV-B- and MpUVR8-dependent, repressed by both MpRUP and MpSPA, and Mp*HY5* is constitutively upregulated in the Mp*cop1-1* mutant (Supplementary Table S4). To test the role of MpHY5 in Marchantia, we generated Mp*hy5* knockout mutants by CRISPR/Cas9 (Supplementary Fig. S3). We further investigated the photoprotection and UV-B-absorbing capacity of Mp*hy5* mutants. Mp*hy5* mutants showed slightly reduced basal photoprotection, as determined by measuring the F_v/F_m ratio after UV-B stress exposure (-/+; Fig. 8A). Interestingly, and in contrast to Arabidopsis *hy5 hyh* double mutants (Leonardelli et al., 2024), Mp*hy5* mutants were able to acclimate to UV-B (+/+ versus -/+; Fig. 8A), but Mp*hy5* mutants nevertheless showed a significantly reduced capacity to protect their photosynthetic machinery from subsequent UV-B-induced damage compared to UV-B-acclimated wild type (+/+; Fig. 8A). In agreement, in contrast to wild type, UV-B-acclimated Mp*hy5* mutants did not survive 7 h of UV-B stress (Fig. 8B). Finally, we measured the UV-B absorbance profiles of methanolic extracts from different genotypes and found that, consistent with the reduced photoprotection observed in UV-B-exposed Mp*hy5*, Mp*hy5* accumulated slightly less UV-B-absorbing compounds compared to Tak-1 wild type (Fig. 8C).

To further understand how MpHY5 promotes MpUVR8 signaling, we performed another RNA-seq experiment with 7-day-old Tak-1 and Mp*hy5-1* Marchantia gemmalings exposed or not to 3 h of narrowband UV-B (Supplementary Tables S1, S2, S6 and S7). Using heatmap analysis and hierarchical clustering, we examined the 501 genes regulated upon UV-B exposure in Tak-1 and/or Mp*hy5-1* (│log2 fold-change│ > 1; p-adj < 0.05) (Supplementary Table S7). This analysis revealed 10 major clusters of gene expression changes in response to UV-B exposure (Fig. 8D, and Supplementary Fig. S7). Notably, few genes were entirely dependent on Mp*HY5*, though certain trends were observed where some genes showed reduced or enhanced UV-B-induced expression compared to the wild-type Tak-1. Mp*HY5-1* (one base deletion, see Supplementary Fig. S3) and Mp*COP1* were hyperinduced in Mp*hy5-1* (cluster 3), while the induction of Mp*RUP* was unaffected by the Mp*hy5-1* mutation (cluster 1). Most UV-B-induced genes were constitutively upregulated in Mp*cop1-1* (Fig. 8D, and Supplementary Table S3), and we noticed that the most MpHY5-dependent genes (clusters 2 and 7) were associated with the highest levels of constitutive upregulation in Mp*cop1*, while the genes that showed the most hyperinduction in Mp*hy5* were associated with the lowest constitutive expression levels in Mp*cop1* (Fig. 8D, Supplementary Fig. S8 and Supplementary Table S6). This is in agreement with an evolutionary conservation of the COP1-HY5 regulatory module in Marchantia photomorphogenesis. Taken together, our data indicate that MpHY5 acts antagonistically to MpCOP1 and is involved in mediating UVR8-responsive gene expression in Marchantia, although to a much lesser extent than that facilitated by HY5 and HYH in Arabidopsis.

Aureusidin synthase MpAUS contributes to the synthesis of UV-absorbing pigments that support Marchantia UV-B tolerance

Several genes involved in the phenylpropanoid biosynthetic pathway, including *PHENYLALANINE AMMONIA LYASE 1* (Mp*PAL1*), *CHALCONE SYNTHASE 2* (Mp*CHS2*), *CHALCONE ISOMERASE-LIKE* (Mp*CHIL2*), and *AUREUSIDIN SYNTHASE* (Mp*AUS*, also named Mp*PPO* (Berland et al., 2019) and Mp*AS1* (Furudate et al., 2023)), were MpUVR8-dependently induced upon UV-B exposure (Fig. 9A, Supplementary Fig. S9). Notably, the transcript level of Mp*AUS*, encoding a key enzyme in biosynthesis of the auronidin flavonoid pigments (Berland et al., 2019; Furudate et al., 2023), exhibited strong upregulation in response to UV-B and was constitutively upregulated in Mp*cop1* (Fig. 9A). Additionally, RT-qPCR analysis

confirmed that the induction of Mp*AUS* following 3 h of UV-B exposure was absent or significantly reduced in Mp*uvr8* and Mp*hy5* mutants, respectively, compared to that in Tak-1 wild type (Fig. 9B). To test the role of Mp*AUS* in UV tolerance, we generated frameshift knockout mutants of Mp*AUS* in Tak-1 using CRISPR/Cas9 mutagenesis (Supplementary Fig. S3). The F_v/F_m ratio in UV-B-acclimated plants after UV-B stress was slightly reduced in Mp*aus-2* compared to that in Tak-1, but not in Mp*rup* Mp*aus* and Mp*spa* Mp*aus* compared to that in Mp*rup* and Mp*spa*, respectively (Fig. 9C, and Supplementary Fig. S10A). In addition, fewer UV-B-acclimated Mp*aus-2* mutants survived subsequent UV-B stress compared to Tak-1 wild type (Fig. 9D and Supplementary Fig. S10B). The Mp*aus* mutation decreased the survival rate of Mp*spa* Mp*aus* double mutants (Supplementary Fig. S10B) but did not affect the survival rate of Mp*rup* Mp*aus* double mutants (Fig. 9D). Altogether, this suggests that MpAUS plays a role in UV stress tolerance, and compensatory effects can occur upon elevated UVR8 activity. Mutation of Mp*aus* affected thallus growth slightly under UV-B in Tak-1 wild type but not in Mp*rup*; however, UV-B-grown Mp*rup* Mp*aus* mutants failed to accumulate characteristic red pigments (Fig. 9E). The UV-B-induced accumulation of UV-absorbing compounds was reduced by the Mp*aus* mutation in a wild-type background, although to a variable extent (Fig. 9F, and Supplementary Fig. S10C). A similar reduction of UV-absorbing compounds was detected in Mp*rup* Mp*aus* and Mp*spa* Mp*aus* versus that in Mp*rup* and Mp*spa* single mutants, respectively (Fig. 9F, and Supplementary Fig. S10C). However, the UV absorbance of Mp*rup* Mp*aus* extracts was much higher than that of wild-type extracts, suggesting that the overaccumulation of UV-absorbing compounds in Mp*rup* occurs in the absence of MpAUS. Next, we performed untargeted metabolomics of 9-day-old wild type Tak-1, Mp*uvr8*, Mp*rup*, and Mp*aus* grown in white light for 7 days and exposed or not to supplemental UV-B for 2 days, following which we detected a total of 609 metabolites that we annotated using a workflow based on SIRIUS 6.2, including CSI:FingerID and CANOPUS, coupled with a taxonomically informed reannotation using the Tima R package (Duhrkop et al., 2019; Rutz and Allard, 2025) (Supplementary Table S8). Fifty-eight metabolites accumulated in response to UV-B in wild-type Tak-1 that were not observed in Mp*uvr8* (Fig. 9G, Supplementary Fig. S11 and Supplementary Table S9). Over half of these (30) were assigned to compounds from the shikimate-phenylpropanoid biosynthetic pathway, whereas metabolites from that pathway represented only about a quarter of all detected compounds (151/609) (Fig. 9H and Supplementary Table S9). This indicates an overrepresentation of shikimate-phenylpropanoid compounds among those UVR8-

regulated metabolites. Similar to Arabidopsis, flavones and flavonol glycosides accumulated in an MpUVR8-dependent manner upon UV-B exposure, which is consistent with previous reports (Supplementary Fig. S11, and Supplementary Table S9, (Clayton et al., 2018)). However, unlike in Arabidopsis whereby sinapate esters play a major role in UV-B photoprotection (Leonardelli et al., 2024), no sinapate esters were detected in Marchantia (Supplementary Fig. S11A, B). Notwithstanding this, 30 of the 58 compounds that accumulated in response to UV-B in wild-type Tak-1 also accumulated in Mp*rup* (Fig. 9G). In addition, 59 compounds were specifically induced under UV-B in Mp*rup*, including auronidin-4-neohesperidoside, a known product of MpAUS (Berland et al., 2019). However, we did not detect the aglycon form of auronidin, possibly because this compound may be present in the cell wall as an insoluble polymer (Jibran et al., 2024). Importantly, we found that the Mp*aus* metabolome was almost non-responsive to UV-B in our conditions (only a total of 17 metabolites accumulated, of which 6 overlapped with wild-type Tak-1), suggesting that, in addition to impairing the synthesis of auronidins, Mp*aus* mutation also affects the accumulation of many other flavonoids.

Taken together, our data suggest that MpAUS contributes to UV-B acclimation in wild type through its direct or indirect effect on the synthesis of diverse phenylpropanoids. However, Mp*rup* and Mp*rup* Mp*aus* absorption spectra reveal the potential contribution of further unidentified, MpAUS-independent UV-absorbing compounds that function in UV-B acclimation in Marchantia.

DISCUSSION

UV-B exposure elicits UVR8-dependent transcriptional changes underlying UV-B acclimation phenotypes in plants (Kliebenstein et al., 2002; Brown et al., 2005; Favory et al., 2009; Tilbrook et al., 2016; Tavridou et al., 2020; Hu et al., 2024; Dannay et al., 2025). The present study provides compelling evidence demonstrating that the core UVR8 photocycle is conserved among land plants, defines the MpUVR8-mediated genome-wide transcriptional response, and provides physiological evidence for how MpUVR8 regulates acclimation and UV-stress tolerance in Marchantia using a broad panel of knock-out mutants in the UVR8 signaling pathway and its output.

In contrast to that in Arabidopsis and the moss *Physcomitrium patens* (hereafter Physcomitrium), MpUVR8 has been reported to form weaker dimers and increased levels of monomer *in vivo* in the absence of UV-B, which is observed when analyzing MpUVR8 dimer/monomer ratio by SDS-PAGE and immunoblotting using samples that are not heat-

denatured before electrophoresis (Soriano et al., 2018). Although this type of assay is accurate for

the dimer/monomer status of wild-type AtUVR8 (Rizzini et al., 2011; Heijde and Ulm, 2013;

Heilmann and Jenkins, 2013; Findlay and Jenkins, 2016; Moriconi et al., 2018), it can make weak

dimeric UVR8 mutant variants appear monomeric, whereas *in vitro* assays (Christie et al., 2012;

Wu et al., 2012; Heijde et al., 2013) as well as more stringent assays using protein-protein

crosslinking identify these variants as homodimeric *in vivo* (Rizzini et al., 2011; Podolec et al.,

2021). Thus, we employed crosslinking to show that MpUVR8 is indeed dimeric *in vivo* in the

absence of UV-B, consistent with the UV-B-dependent dimer-to-monomer transition being a

conserved feature of the UVR8 photocycle (Rizzini et al., 2011; Tilbrook et al., 2016; Podolec et

al., 2021). Furthermore, we provide evidence that negative feedback regulation through MpRUP-

facilitated redimerization of MpUVR8 is conserved in Marchantia (Heijde et al., 2013; Podolec et

al., 2021; Zhang et al., 2021). Consistent with this, the Mp*rup* mutant displayed a strong UV-B

photomorphogenic phenotype, characterized by both growth inhibition resulting in a small size and

a strong pigmentation of the thallus (this study and (Clayton et al., 2018)). Interestingly, this

conditional phenotype of Mp*rup* specifically under UV-B phenocopies the constitutive Mp*cop1*

phenotype, in agreement with active UVR8-mediated inactivation of COP1 underlying UV-B

signaling (Yin et al., 2015; Lau et al., 2019). We further show that GFP-MpUVR8 expressed under

its own promoter localizes mainly to the cytosol and displays UV-B-dependent nuclear

accumulation in Marchantia, confirming findings using a *Citrine*-Mp*UVR8* overexpression line

(Kondou et al., 2019). Thus, we conclude that the homodimeric ground state of UVR8, its UV-B

dependent monomerization, nuclear accumulation, and COP1 inactivation, as well as negative

feedback regulation of UVR8 through RUP-facilitated redimerization are well-conserved in

Marchantia.

In flowering plants, COP1 forms an E3 ligase complex together with SPA proteins, which

together function as suppressors of photomorphogenesis; i.e. COP1 requires SPA proteins to

function (Hoecker, 2017; Podolec and Ulm, 2018). Since non-plant COP1 orthologs are not

regulated by light, it is hypothesized that the evolution of SPA proteins was a crucial step in

enabling photoreceptor-mediated regulation of COP1/SPA complex activity (Ranjan et al., 2014;

Balcerowicz et al., 2017; Podolec and Ulm, 2018). Active phytochrome and cryptochrome

photoreceptors directly interact with SPA proteins, which represses the E3 ligase activity of the

COP1/SPA complex (Hoecker, 2017; Podolec and Ulm, 2018). In agreement, Arabidopsis

quadruple *spa* mutants exhibit strong constitutive photomorphogenesis (Ordonez-Herrera et al., 2015). However, it is of note that *spa* quadruple null mutants can complete their life cycle whereas *cop1* null mutants cannot (McNellis et al., 1994; Ordonez-Herrera et al., 2015), suggesting that COP1 can function partially in a SPA-independent manner. COP1, but not SPA proteins, are also necessary for transient gene expression changes upon exposure to UV-B (Favory et al., 2009). Recently, the role of COP1/SPA has started to be explored in more phylogenetically distant land plant lineages such as bryophytes, including Physcomitrium (Kreiss et al., 2023). A Physcomitrium Pp*cop1_9x* mutant lacking all nine PpCOP1(a-i) orthologs was viable and displayed a rather minor growth phenotype, certainly much less severe than the Arabidopsis *cop1* mutant phenotypes (Kreiss et al., 2023). In this study, we analyzed mutants of the single Mp*COP1* gene in light-grown Marchantia plants and found that, although they completed their asexual life cycle, they exhibited profound transcriptome changes and a severe constitutively photomorphogenic phenotype. Additionally, consistent with the described role of COP1 in UV-B signaling in Arabidopsis and Chlamydomonas (Oravecz et al., 2006; Favory et al., 2009; Allorent et al., 2016; Tilbrook et al., 2016), transient gene expression changes upon exposure to UV-B were dependent on MpCOP1. The Marchantia genome contains a single gene encoding MpSPA that interacts with MpCOP1 and MpCRY (Zhang et al., 2021). Surprisingly, Mp*spa* null mutants exhibited at most a mild constitutively photomorphogenic phenotype in white light, unlike Mp*cop1*. Interestingly, in Physcomitrium, a Pp*spa_ab* double mutant displayed an intermediate phenotype between that of the wild type and Pp*cop1_9x*. This suggests that SPA proteins indeed play a lesser role in repressing light signaling in bryophytes compared to in angiosperms (Huang et al., 2013; Ordonez-Herrera et al., 2015; Artz et al., 2019; Kreiss et al., 2023) and that high dependence of COP1 on the accessory SPA proteins evolved later, meaning that SPA protein evolution is separate to the evolution of COP1 activity regulation by light. This is further supported by the discovery of the UVR8- and cryptochrome photoreceptor–mediated COP1 inactivation mechanism of high-affinity competitive binding and displacement of COP1 substrates (Lau et al., 2019; Ponnu et al., 2019; Trimborn et al., 2025). Moreover, complementation of corresponding Arabidopsis mutant phenotypes was possible with a Physcomitrium COP1 ortholog but not a SPA ortholog (Ranjan et al., 2014), further indicating the divergence of SPA function between evolutionary lineages.

Enhanced UV-B-induced changes in gene expression, increased accumulation of UV-absorbing compounds, and enhanced UV-B acclimation and photoprotection against UV stress in

Mp*spa* mutants compared to that in wild type revealed a repressor role for SPA in the UV-B signaling pathway of Marchantia. This is similar to Mp*rup* mutants, although long-term growth under UV-B was not as strongly inhibited in Mp*spa*. MpSPA also did not facilitate MpUVR8 redimerization like observed for MpRUP. Several hypotheses can be proposed to explain the Mp*spa* phenotype. It has been reported that MpSPA can promote the ubiquitination of MpHY5 by MpCOP1 (Zhang et al., 2021). Consequently, the lack of MpSPA could result in reduced ubiquitination of MpHY5 by MpCOP1, leading to higher activation of downstream genes. Also, UV-B-activated MpUVR8 may inactivate MpCOP1 more strongly in the absence of MpSPA, leading to an enhanced UV-B response. MpSPA might function by "shielding" the MpCOP1/SPA complex from being inhibited by MpUVR8 under UV-B by providing an additional UVR8-irrepressible activity. Moreover, it has been suggested that SPA proteins promote the phosphorylation of several proteins involved in light signaling in Arabidopsis, including UVR8 and HY5 (Wang et al., 2021; Liu et al., 2024). Therefore, in the event such activity is conserved, the absence of MpSPA may affect the phosphorylation status and activity of light signaling components, enabling the Mp*spa* mutant to respond more strongly to UV-B and to produce more "sunscreen" metabolites. Future work is required to elucidate the precise mechanism by which MpSPA exerts its repressive effect on UV-B signaling in Marchantia, and to determine if a similar SPA protein function exists in Arabidopsis or whether this represents a species-specific UV-B response regulatory mechanism. Overall, genetic analysis of COP1/SPA function suggests that, whereas the role of COP1 is widely conserved within the green lineage, the importance of SPA proteins in regulating COP1 activity and repressing photomorphogenesis increased during plant evolution.

It is of note that the role and importance of the transcriptional regulators HY5/HYH in UVR8-signaling has also changed during plant evolution. HY5 is a major player in UVR8-signaling in Arabidopsis (Ulm et al., 2004; Brown et al., 2005; Oravecz et al., 2006; Binkert et al., 2014), whereas the closest Chlamydomonas homolog of HY5, CrBLZ3, plays only a minor role in the algal UV-B response (Dannay et al., 2025). Instead, the B-box family transcription factor and CrCOP1 substrate CrCONSTANS is the main player in the CrUVR8 signaling pathway (Gabilly et al., 2019; Tokutsu et al., 2019; Dannay et al., 2025). In Marchantia, MpHY5 controls a subset of MpUVR8-induced genes but has a less prominent role than its counterpart in Arabidopsis. Other transcription factor(s) mediating MpUVR8 signaling remain to be identified.

In land plants, UVR8 activates the phenylpropanoid pathway, increasing the accumulation

of visible anthocyanins and colorless, UV-absorbing phenylpropanoids, such as flavonoids and

hydroxycinnamic acids, which overall contribute to photoprotection. Sinapate esters play a major

UV-sunscreen role in Arabidopsis (Stracke et al., 2010; Ferreyra et al., 2021; Leonardelli et al.,

2024), but are not present in Marchantia. Anthocyanins are also absent in Marchantia, but a red

pigment derived from chalcones, auronidin, is strongly accumulated upon different stresses (Albert

et al., 2018; Berland et al., 2019). Auronidin biosynthesis is catalyzed by MpAUS, a polyphenol

oxidase (PPO) enzyme found exclusively in liverworts and angiosperms, but not in hornworts and

mosses (Fig. 9A) (Davies et al., 2022; Furudate et al., 2023). Expression of Mp*AUS* is strongly

upregulated by UV-B in a UVR8-dependent manner. As Mp*AUS* overaccumulation following

Mp*MYB14* overexpression was associated with increased UV-B tolerance (Clayton et al., 2018),

we further characterized the role of MpAUS in UV-B photoprotection. A combination of genetics,

physiological analysis, and metabolomics revealed that MpAUS contributes to photoprotection

against UV-B. However, this contribution may not solely be attributed to the synthesis of auronidin,

as the accumulation of many other UV-B absorbing compounds was affected in the Mp*aus* mutant.

This indicates that MpAUS has specificity for more than just auronidin biosynthesis as previously

proposed (Berland et al., 2019) and may accept different substrates (Furudate et al., 2023). Future

experiments will be necessary to more precisely characterize how different classes of

phenylpropanoids contribute to photoprotection, and to investigate other types of photoprotective

mechanisms in Marchantia.

In conclusion, our study provides a comprehensive understanding of the UV-B perception

and signaling mechanisms in the liverwort model organism Marchantia. Our findings further

confirm and strengthen the notion that the UVR8 photoreceptor and its associated signaling

components are conserved in Marchantia, highlighting the evolutionary importance of this pathway

in mediating UV-B responses across land plants. Our analysis of the COP1/SPA E3 ligase complex

and its HY5 substrate in Marchantia also revealed divergences in the functions of this

photomorphogenesis-repressing complex. Overall, our work further validates Marchantia as a

useful system to investigate light signaling and its evolution in plants.

MATERIALS AND METHODS

Plant materials

The *Marchantia polymorpha* accessions Takaragaike-1 (Tak-1) and Takaragaike-2 (Tak-2) were used as male and female wild type, respectively (Ishizaki et al., 2008). The Mp*uvr8-1* mutant was described previously (Kondou et al., 2019), whereas the following mutants were generated using CRISPR–Cas9-mediated mutagenesis (Sugano et al., 2018; Sauret-Güeto et al., 2020): Mp*rup-2,* Mp*rup-3,* Mp*cop1-1,* Mp*cop1-2,* Mp*spa-9,* Mp*spa-10,* Mp*hy5-1,* Mp*hy5-3,* Mp*hy5-7,* Mp*aus-2,* and Mp*aus-3*. Two different gene-specific guide RNAs (gRNAs, see Supplemental Table S10) were selected using the CRISPRdirect (https://crispr.dbcls.jp/) or CasFinder (https://marchantia.info/tools/casfinder/) tools (Aach et al., 2014). Corresponding complementary DNA oligos were annealed and ligated into the BsaI sites of pMpGE_En03, then subcloned to the binary vector pMpGE010 (Sugano et al., 2018) using Gateway LR Clonase II (Thermo Fisher Scientific). To generate the Mp*hy5* and Mp*aus* mutants, complementary oligos were inserted into BbsI sites of the L1_lacZgRNA-Ck2 and L1_lacZgRNA-Ck3 vectors, respectively, after which these were recombined by SapI restriction enzymes with the L1_HyR-Ck1 and L1_Cas9-Ck4 vector into the pCsA acceptor plasmid using Golden Gate Loop Assembly (Sauret-Güeto et al., 2020). Following transformation, frame-shift and early stop codon mutants were identified using PCR genotyping and Sanger sequencing (Supplemental Table S10, and Supplemental Fig. S3 with actual mutations).

To generate Pro_{MpUVR8}:*GFP*-Mp*UVR8* reporter lines, we modified the previously described pMpGWB301-MpUVR8 genomic construct (Kondou et al., 2019). The vector was digested with MluI and HpaI restriction enzymes and the GFP coding sequence was ligated immediately upstream of the Mp*UVR8* start codon. To create Pro_{MpCHS2}:*LUC* reporter lines, the predicted Mp*CHS2* promoter including 5′UTR sequence (3,000 bp region upstream of the Mp*CHS2* start codon) was amplified from Tak-1 genomic DNA and the PCR product was inserted into pDONR207 via Gateway BP reaction. The Pro_{MpCHS2} sequence was then transferred into the pMpGWB331 vector by LR reaction, followed by transformation into the Tak-1 wild-type background. To create Pro_{MpCHS2}:*LUC* lines in the Mp*uvr8-1* and Mp*rup-1* backgrounds, Tak-1/Pro_{MpCHS2}:*LUC* was first crossed with Tak-2 to obtain Tak-2/Pro_{MpCHS2}:*LUC* in female background, and then the male Mp*uvr8-1* or Mp*rup-2* mutants were crossed with the female Tak-2/Pro_{MpCHS2}:*LUC* line. To generate overexpression lines in Tak-1, the coding sequences including

stop codons of Mp*UVR8.1*, Mp*UVR8.2*, Mp*RUP*, Mp*COP1*, and Mp*SPA* were first cloned into the pDONR207 entry vector via BP reaction, then transferred via LR reaction into the pMpGWB336 destination vector containing the *ELONGATION FACTOR 1α* (Mp*EF1α*, Mp3g23400) promoter and an N-terminal Red Fluorescent Protein (RFP) tag. Agrobacterium-mediated Marchantia thallus transformation was performed as previously described (Kubota et al., 2013). Marchantia spore transformation was used to generate Mp*hy5* and Mp*aus* mutants as previously described (Chiyoda et al., 2008).

Mp*rup-2* Mp*aus-2*, Mp*spa-10* Mp*aus-2*, and Mp*uvr8-1* Mp*rup-2* Cas9-free double mutants were generated by crossing: first Mp*aus-2* (male) and Mp*rup-2* (male) were crossed with Tak-2 to select for female lines, followed by crossing Mp*rup-2* (male) or Mp*spa-10* (male) with Mp*aus-2* (female), and Mp*uvr8-1* (male) with Mp*rup-2* (female), as previously described (Mulvey and Dolan, 2023). The inheritance of the parental Mp*rup-2* Mp*aus-2* and Mp*spa-10* Mp*aus-2* alleles was evaluated by Sanger sequencing for hygromycin-sensitive T1 plants. For those plants where the inheritance of the parental alleles was confirmed, replica plating on plates with and without hygromycin was repeated to further verify the absence of the Cas9 transgene.

Growth conditions and light treatments

For maintenance, Marchantia plants were cultured aseptically on half-strength Gamborg's B5 medium with 1.2% (w/v) agar at 22°C under continuous white light (50 μmol m^{-2} s^{-1}) in a CLF GroBank growth chamber. For UV-B acclimation experiments, continuous white light was supplemented with narrowband UV-B tubes (Philips TL20W/01RS, 1.5 μmol m^{-2} s^{-1}). For UV-B stress treatments, plates were irradiated under broadband UV-B tubes (Philips TL20W/12RS; 25 μmol m^{-2} s^{-1}).

Crosslinking and immunoblotting

Crosslinking of proteins and immunoblot analysis of 7-day-old Marchantia gemmalings was performed following a modified version of a previously described method (Rizzini et al., 2011), except that the extraction buffer did not contain leupeptine, dichloroisocumarin, MG115, and PSI. Polyclonal rabbit anti-MpUVR8[(3-17)] antibodies raised and purified against a synthetic peptide (amino acids C + 3–17: CDSKMSSNERPRRRVL) were used as the primary antibodies (Eurogentec), with horseradish peroxidase (HRP)-conjugated anti-rabbit immunoglobulins (Dako

A/S) as secondary antibodies. The signal was revealed using ECL Select Western Blotting
Detection Reagent (GE Healthcare) and analyzed using an Amersham Imager 680 camera system
(GE Healthcare).

Quantitative reverse transcription PCR (RT-qPCR) analysis
RNA was isolated with the RNeasy Plant Mini kit (Qiagen) according to the manufacturer's
instructions and was treated on-column with the RQ1 RNase-Free DNase (Promega), before
serving as template for cDNA synthesis using the TaqMan Reverse Transcription Reagents kit
(Thermo Fisher Scientific) with a 1:1 mix of oligo-dT and random hexamer primers. RT-qPCR
was performed using the PowerUp SYBR Green Master Mix (Thermo Fisher Scientific) on a
QuantStudio^{TM} 5 Real-Time PCR System (Thermo Fisher Scientific). Mp*EF1α* was used as
reference gene for normalization (Saint-Marcoux et al., 2015), and expression values were
calculated using the $2^{(-\Delta\Delta Ct)}$ method (Livak and Schmittgen, 2001). Gene-specific primers are listed
in Supplemental Table S10. Every experiment was performed using three biological replicates, and
all expression values were normalized against the average of untreated wild type, which was set to
1.

Yeast two-hybrid
Yeast two-hybrid assays were performed as described previously (Rizzini et al., 2011). Gateway-
based cloning was used to insert the Mp*UVR8.1*, Mp*UVR8.2*, Mp*RUP*, Mp*COP1*, and Mp*SPA*
coding sequences downstream of the LexA DNA binding domain (BD) in pBTM116-D9 (Stelzl et
al., 2005) or GAL4 activation domain (AD) in pGADT7-GW (Marrocco et al., 2006). The empty
vectors were used as negative controls. All the constructs were transformed into the yeast strain
L40 using the lithium-acetate protocol (Gietz et al., 2014). For the interaction assays, serial
dilutions of transformed yeast colonies were grown under narrowband UV-B (1.5 μmol m^{-2} s^{-1}) at
30°C for three days either under a WG305 (+UV-B) or a WG368 (-UV-B control) cut-off filter
(Schott Glaswerke). For quantification, transformed yeast colonies were grown under +UV-B or -
UV-B control conditions for 16 h. Ten colonies were then combined and resuspended in YPD
media, OD$_{600}$ was measured, and cells were spun down and washed. The quantitative interaction
assay was carried out with chlorophenol red-β-D-galactopyranoside (CPRG; Roche Applied
Science) as substrate, according to the Yeast Protocols Handbook (Clontech, Version PR973283).

Affinity purification – mass spectrometry (AP–MS)

AP–MS was performed using a Marchantia transgenic line expressing Pro_{MpUVR8}:GFP-Mp$UVR8$, with wild-type Tak-1 and a Pro_{MpUBQ2}:$mVenus$ line as negative controls. Plants were grown for 7 days under continuous white light (50 µmol m^{-2} s^{-1}) without UV-B, or under continuous white light (50 µmol m^{-2} s^{-1}) supplemented with narrowband UV-B (1.5 µmol m^{-2} s^{-1}) followed by a 20 min broadband UV-B exposure (25 µmol m^{-2} s^{-1}) immediately prior to harvest. Total protein was extracted and GFP-Trap agarose beads (Chromotek) were used for affinity purifications. Eluted proteins were subjected to LC-MS/MS analysis.

Sample preparation and LC-MS/MS data acquisition: enriched proteins were submitted to an on-bead digestion using trypsin. In brief, dry beads were re-dissolved in 25 µL digestion buffer 1 (50 mM Tris, pH 7.5, 2M urea, 1mM DTT, 5 ng/µL trypsin) and incubated for 30 min at 30 °C in a Thermomixer with 400 rpm. Next, beads were pelleted and the supernatant was transferred to a fresh tube. Digestion buffer 2 (50 mM Tris, pH 7.5, 2M urea, 5 mM CAA) was added to the beads, after mixing the beads were pelleted, the supernatant was collected and combined with the previous one. The combined supernatants were then incubated o/n at 32°C in a Thermomixer with 400 rpm; samples were protected from light during incubation. The digestion was stopped by adding 1 µL TFA and desalted with C18 Empore disk membranes according to the StageTip protocol (Rappsilber et al., 2003).

Dried peptides were re-dissolved in 2% ACN, 0.1% TFA (10 µL) for analysis. Samples were analyzed using an EASY-nLC 1200 (Thermo Fisher) coupled to a Q Exactive Plus mass spectrometer (Thermo Fisher). Peptides were separated on 16 cm frit-less silica emitters (New Objective, 75 µm inner diameter), packed in-house with reversed-phase ReproSil-Pur C18 AQ 1.9 µm resin (Dr. Maisch). Peptides were loaded on the column and eluted for 115 min using a segmented linear gradient of 5% to 95% solvent B (0 min : 5%B; 0-5 min -> 5%B; 5-65 min -> 20%B; 65-90 min ->35%B; 90-100 min -> 55%; 100-105 min ->95%, 105-115 min ->95%) (solvent A 0% ACN, 0.1% FA; solvent B 80% ACN, 0.1%FA) at a flow rate of 300 nL/min. Mass spectra were acquired in data-dependent acquisition mode with a TOP15 method. MS spectra were acquired in the Orbitrap analyzer with a mass range of 300–1750 m/z at a resolution of 70,000 FWHM and a target value of 3×10^6 ions. Precursors were selected with an isolation window of 1.3 m/z (Q Exactive Plus). HCD fragmentation was performed at a normalized collision energy of 25.

MS/MS spectra were acquired with a target value of 10^5 ions at a resolution of 17,500 FWHM, a

maximum injection time (max.) of 55 ms and a fixed first mass of m/z 100. Peptides with a charge

of +1, greater than 6, or with unassigned charge state were excluded from fragmentation for MS^2,

dynamic exclusion for 30s prevented repeated selection of precursors.

Data analysis: raw data were processed using MaxQuant software (version 1.6.3.4,

http://www.maxquant.org/) (Cox and Mann, 2008) with label-free quantification (LFQ) and iBAQ

enabled (Tyanova et al., 2016).

MS/MS spectra were searched by the Andromeda search engine against a combined

database containing the sequences from *M. polymorpha* (MpTak_v6.1r2.protein.fasta;

https://marchantia.info/download/MpTak_v6.1r2/) and sequences of 248 common contaminant

proteins and decoy sequences. Trypsin specificity was required and a maximum of two missed

cleavages allowed. Minimal peptide length was set to seven amino acids. Carbamidomethylation

of cysteine residues was set as fixed, oxidation of methionine and protein N-terminal acetylation

as variable modifications. The match between runs option was enabled. Peptide-spectrum-matches

and proteins were retained if they were below a false discovery rate of 1% in both cases. The MS

data have been deposited in the ProteomeXchange Consortium via the PRIDE partner repository

with the dataset identifier PXD065691.

Statistical analysis of the MaxLFQ values was carried out using Perseus (version 1.5.8.5,

http://www.maxquant.org/). Quantified proteins were filtered for reverse hits and hits "identified

by site" and MaxLFQ values were log2 transformed. After grouping samples by condition only

those proteins were retained for the subsequent analysis that had three valid values in one of the

conditions. Missing values were imputed from a normal distribution (1.8 downshift, separately for

each column). Significantly enriched proteins were identified using a p-value cutoff of < 0.01 and

a log10 fold change threshold of ±1.0 (Supplementary Table S11).

F_v/F_m measurements

A PSI (Photon Systems Instruments) FluorCam 800 MF was used to measure the maximum

quantum efficiency of PSII (F_v/F_m). Plates containing 15 gemmalings per genotype were incubated

in darkness for at least 5 minutes, which was sufficient to ensure a steady state under the plant

growth conditions. Maximum fluorescence in the dark (F_m) was measured using saturating pulses

of 2000 µmol m^{-2} s^{-1} white light (400–720 nm) for 960 ms, followed by detection pulses of orange-

red light (620 nm) lasting 10 µs. The F_v/F_m ratio was calculated using the formula $F_v/F_m = (F_m -$

$F_o)$ / F_m, where F_m represents the maximal fluorescence, and F_o represents the minimal

fluorescence in the dark-adapted state.

Luciferase complementation imaging (LCI) and reporter assays

The split-luciferase assays were performed as previously described (Zhou et al., 2018). Site-

directed mutagenesis was performed as described previously (Sawano and Miyawaki, 2000) to

generate MpUVR8.1^{D95N,D106N} and MpUVR8.2^{D95N,D106N}. The pCambia1300–nLUC vector was

first digested with BamHI and SalI, while the pCambia1300–cLUC vector was digested with KpnI

and SalI. Subsequently, the cDNA fragments of Mp*UVR8.1*, Mp*UVR8.2*, Mp*UVR8.1^{D95N,D106N}*,

Mp*UVR8.2^{D95N,D106N}*, Mp*RUP*, and Mp*COP1* were cloned into the two digested vectors using

Gibson assembly. *Agrobacterium tumefaciens* strain GV3101 harboring the indicated

combinations of constructs expressing nLUC and cLUC fusion proteins were mixed in a 1:1 ratio

and infiltrated into *N. benthamiana* leaves. The plants were kept in the growth chamber for 2 days

before being incubated with 1 mM luciferin (D-luciferin sodium salt, Yeasen Biotechnology) for

10 min (in darkness). Luciferase activity was detected using a NightSHADE LB985 In Vivo Plant

Imaging System (Berthold).

For *Pro$_{MpCHS2}$:LUC* luciferase reporter assays, gemmae were transferred into 96-well plates

containing ½-strength Gamborg's B5 medium with 1.2% (w/v) agar and grown under at 22°C

continuous white light (50 µmol m^{-2} s^{-1}) in a CLF GroBank growth chamber for two days. Four-

µL of 1 mM luciferin (Yeasen Biotechnology) were added to each well and the following day plants

were exposed to UV-B for 20 min and luminescence was recorded using a Centro XS LB 960

luminometer.

Extraction of phenolic compounds, measurements of absorption spectra, and high-

performance thin-layer chromatography (HPTLC)

Phenolic compounds were extracted in 100 µL of 80% (v/v) methanol from 50 mg of fresh plant

material homogenized using a Silamat S6 (Ivodar Vivadent). Extracts were incubated for 10 min

at 70°C on a shaker and centrifuged for 10 min at 12,000 *g* at room temperature (RT). A 40-µL

aliquot of clear supernatant from each extract was transferred to a transparent 96-well plate and

absorption spectra (200–1000 nm wavelengths) were measured using a TECAN Spark 10M

Multimode Microplate Reader. In case absorbance values exceeded the linear range, aliquots were
uniformly diluted with extraction buffer.

HPTLC was used to analyze the phenolic compound profile using silica-60 HPTLC glass
plates (Sigma-Aldrich), as described previously (Stracke et al., 2010; Leonardelli et al., 2024).

Microscopy

Pro_{MpUVR8}:*GFP*-Mp*UVR8* transgenic gemmalings were grown for 7 days under
50 µmol m^{-2}s^{-1} white light and irradiated for 20 min with 25 µmol m^{-2} s^{-1} broadband UV-B.
Imaging was performed using a Leica Thunder Widefield microscope in both GFP and bright-field
modes. Z-stack images were acquired at 2-µm intervals. Image processing and Z-projection were
conducted using ImageJ software.

Transcriptome analysis

Gemmalings were grown for 7 days in 50 µmol m^{-2} s^{-1} white light and exposed or not to
supplemental UV-B (1.5 µmol m^{-2} s^{-1}) for 3 h. Total RNA was extracted using the RNeasy Plant
Mini kit (Qiagen) and Illumina Tru-Seq libraries were generated from cDNA and sequenced using
Illumina NovaSeq 100-bp single-end at the iGE3 genomics platform of the University of Geneva
(Supplementary Table S1). Raw reads were trimmed for quality using Trim Galore (v0.6.7;
github.com/FelixKrueger/TrimGalore) with "-q 25 --stringency 5" as parameters. Trimmed reads
were mapped to the MpTak.v7.1 genome reference using STAR (v2.7.4; (Dobin et al., 2013),
followed by read quantification using featureCounts (Subread v2.0.1; (Liao et al., 2019). The raw
counts were normalized and differential expression analysis was performed using DESeq2 (Love
et al., 2014). Genes with a log$_2$ fold change greater than 1 and a p-adj value of less than 0.05 were
considered as significantly differentially expressed. Genes with lower log2 fold change values and
p-adj < 0.05 were considered when comparing different genotypes in which gene expression
changes may occur at varying magnitude (Supplementary Table S2). Euler plots were generated
using the eulerr package in R. Heatmaps were generated using the heatmap.2 function in the gplots
package in R. The raw RNA-sequencing data has been deposited in the NCBI Gene Expression
Omnibus (www.ncbi.nlm.nih.gov/geo) and is accessible through GEO Series accession number
GSE296364.

Untargeted metabolomics

7-day-old Marchantia gemmalings were ground in liquid nitrogen using a mortar and pestle. 50 mg of tissue powder were transferred into a 1.5-mL tube and compounds were extracted with 5 volumes of 80% (v/v) methanol containing 0.1% (v/v) formic acid. After centrifugation, the supernatant was transferred to an HPLC glass vial fitted with a conical insert. A 0.75-µL injection of the extract was performed into ultrahigh performance liquid chromatography coupled to high-resolution mass spectrometry (UHPLC-HRMS) equipped with an electrospray ionization (ESI) source operating in negative (ESI-) ionization mode. Specifically, the system used was an Acquity UPLC I-Class (Waters) interfaced with a Synapt XS QTOF (Waters). The parameters used for untargeted metabolomics were based on (Defossez et al., 2023) with minor adjustments. Briefly, chromatography was done on a Waters Acquity UPLC™ HSS T3 column (100 × 2.1 mm, 1.8 µm) with mobile phases consisting of $H_2O + 0.05\%$ (v/v) formic acid (solvent A) and acetonitrile + 0.05% (v/v) formic acid (solvent B). A linear gradient from 0 to 100% B in 10.0 min at a flow rate of 0.5 mL/min was applied. MS detection was performed in negative electrospray ionisation using data-dependent acquisition (DDA). The mass range for both MS and MS/MS was 50–1200 Da. Scan times for MS and MS/MS were 0.1 and 0.05 s, respectively. The top 7 precursor ions were selected for MS/MS with an intensity threshold of 15,000 counts per second. An exclusion list of the 150 most intense background ions was generated from a blank sample run just before the plant samples, and a dynamic exclusion of 1.5 s was applied after each MS/MS acquisition. For MS/MS acquisition, a ramped collision energy of 5–40 V (at *m/z* 50) and 20–70 V (at *m/z* 1200) was set. The entire system and method were controlled by MassLynx 4.2 (Waters). Raw profile data were first centroided in MassLynx and further converted to MzML using MSConvert (ProteoWizard). We used MZmine 4.5 for feature detection, deconvolution, and alignment of DDA (data-dependent acquisition) data, with noise thresholds set to 2000 for MS1 and 50 for MS2 (Schmid et al., 2023). The resulting peak table and MS/MS spectral data were exported and annotated using SIRIUS 6.2 (Duhrkop et al., 2019), including CSI:FingerID for structural identification and CANOPUS for the prediction of compound classes, superclasses, and biosynthetic pathways. Taxonomically informed reannotation of molecular features was performed using the Tima R package (Rutz and Allard, 2025).

Statistical analyses

Statistical analyses were performed using GraphPad Software Prism 10 software (San Diego, California). Statistical significance of the differences between means was determined using one- or two-way analysis of variance (ANOVAs) followed by Tukey's post hoc honestly significant difference (HSD) test for multiple comparisons. For metabolomics, adjusted p-values were calculated and corrected using the Benjamini-Hochberg in R.

Accession numbers

Mp2g11590 (Mp*UVR8*), Mp2g18040 (Mp*RUP*), Mp5g12010 (Mp*COP1*), Mp3g25460 (Mp*SPA*), Mp1g16800 (Mp*HY5*), Mp2g05850 (Mp*AUS*), Mp2g07060 (Mp*CHS2*), Mp1g10150 (Mp*PAL1*), Mp6g00020 (Mp*C4H1*), Mp8g05070 (Mp*C4H2*), Mp8g05080 (Mp*C4H3*), Mp1g27780 (Mp*4CL1*), Mp1g05060 (Mp*4CL2*), Mp2g22700 (Mp*CHS6*), Mp5g01410 (Mp*CHIL2*), Mp6g05290 (Mp*CHI*), Mp1g26540 (Mp*F3H*), Mp3g23400 (Mp*EF1α*).

Supplementary data

Supplementary Figure S1. Alignment of Marchantia MpUVR8.1 and MpUVR8.2 with Arabidopsis AtUVR8.

Supplementary Figure S2. Volcano plot of GFP-MpUVR8 AP-MS under +UV versus -UV conditions.

Supplementary Figure S3. Gene structure and indication of the mutated sites in the corresponding mutants generated by CRISPR/Cas9.

Supplementary Figure S4. UV-B photomorphogenesis and photoprotection in MpUVR8 and MpRUP transgenic lines.

Supplementary Figure S5. Enhanced photoprotection and UV-B-absorbing capacity of Mp*rup* depends on MpUVR8.

Supplementary Figure S6. MpSPA does not affect MpUVR8 redimerization, but MpCOP1 and MpSPA overexpression reduce basal UV-B tolerance.

Supplementary Figure S7. MpHY5 plays a minor role in the regulation of UV-B-induced gene expression changes.

Supplementary Figure S8. MpHY5 and MpCOP1 regulate UV-B-induced gene expression in an antagonist manner.

Supplementary Figure S9. LUC activity of a *Pro*_{MpCHS2}*:LUC* reporter in Tak-1, Mp*uvr8-1* and Mp*rup-2* backgrounds upon UV-B exposure.

Supplementary Figure S10. MpAUS contributes to protection against UV-B stress.

Supplementary Figure S11. MpUVR8- and MpAUS-dependent regulation of metabolite accumulation upon UV-B exposure.

Supplementary Table S1. List of RNA-seq libraries.

Supplementary Table S2. Raw read counts, normalized read counts and differential expression analysis from RNA-seq experiments.

Supplementary Table S3. Overlaps between UV-B regulated genes in Tak-1, Mp*rup* and Mp*spa* (lists corresponding to Figure 6A,B) and overlaps between genes misregulated in Mp*cop1* and Mp*spa* in white light (lists corresponding to Figure 7B,C).

Supplementary Table S4. Hierarchical clustering of UV-B-regulated gene expression.

Supplementary Table S5. Hierarchical clustering of MpCOP1/SPA-regulated gene expression.

Supplementary Table S6. Overlaps between genes misregulated in Mp*cop1* in white light and genes regulated by MpHY5 after UV-B exposure (lists corresponding to Supplementary Figure S8).

Supplementary Table S7. Hierarchical clustering of UV-B-regulated gene expression in Tak-1 and Mp*hy5*.

Supplementary Table S8. Methanol-soluble metabolites of Marchantia in presence and absence of UV-B.

Supplementary Table S9. Overlaps of methanol-soluble metabolites in response to UV-B between Tak-1, Mp*uvr8*, Mp*rup* and Mp*aus*. Metabolite lists corresponding to Figure 9G.

Supplementary Table S10. Primer list.

Supplementary Table S11. AP-MS data.

ACKNOWLEDGEMENTS

We would like to thank Youichi Kondou (Kanto Gakuin University College of Science and Engineering, Yokohama, Japan) for kindly providing the Mp*uvr8-1* mutant and the pMpGWB301-MpUVR8 construct, Félix Rico-Reséndiz (University of Geneva) for sharing protocols and tips regarding Marchantia cultivation and transformation, Yanfei Zhou (The New Zealand Institute for Plant and Food Research Limited, Palmerston North, New Zealand) for precious suggestions, and Anne Harzen (Max-Planck Institute for Plant Breeding Research) for MS sample preparation. The

RNA-seq experiments were performed at the iGE3 Genomics Platform of the University of Geneva (https://ige3.genomics.unige.ch/). This work was supported by the University of Geneva, and, in part, the Swiss National Science Foundation (grant 310030_207716 to R.U.).

AUTHOR CONTRIBUTIONS

Y.L., R.P., and R.U. conceived and designed the research; Y.L., R.P., and R.C. performed the experiments; G.G. and E.Def. performed and contributed metabolomic data; S.C.S and H.N. performed and analyzed LC-MS/MS data; J.R. and L.D. contributed Mp*hy5* mutants; Y.L., R.P., E.Dem., and R.U. analyzed the data and wrote the manuscript. All authors reviewed and approved the submitted manuscript.

Competing interests: The authors have declared that no competing interests exist.

REFERENCES

Aach J, Mali P, Church GM (2014) CasFinder: Flexible algorithm for identifying specific Cas9 targets in genomes. bioRxiv: 005074

Albert NW, Thrimawithana AH, McGhie TK, Clayton WA, Deroles SC, Schwinn KE, Bowman JL, Jordan BR, Davies KM (2018) Genetic analysis of the liverwort *Marchantia polymorpha* reveals that R2R3MYB activation of flavonoid production in response to abiotic stress is an ancient character in land plants. New Phytol 218: 554-566

Allorent G, Lefebvre-Legendre L, Chappuis R, Kuntz M, Truong TB, Niyogi KK, Ulm R, Goldschmidt-Clermont M (2016) UV-B photoreceptor-mediated protection of the photosynthetic machinery in *Chlamydomonas reinhardtii*. Proc Natl Acad Sci USA 113: 14864-14869

Artz O, Dickopf S, Ranjan A, Kreiss M, Abraham ET, Boll V, Rensing SA, Hoecker U (2019) Characterization of spa mutants in the moss Physcomitrella provides evidence for functional divergence of SPA genes during the evolution of land plants. New Phytol 224: 1613-1626

Balcerowicz M, Kerner K, Schenkel C, Hoecker U (2017) SPA proteins affect the subcellular localization of COP1 in the COP1/SPA ubiquitin ligase complex during photomorphogenesis. Plant Physiol 174: 1314-1321

Berland H, Albert NW, Stavland A, Jordheim M, McGhie TK, Zhou Y, Zhang H, Deroles SC, Schwinn KE, Jordan BR, Davies KM, Andersen OM (2019) Auronidins are a previously unreported class of flavonoid pigments that challenges when anthocyanin biosynthesis evolved in plants. Proc Natl Acad Sci USA 116: 20232-20239

Binkert M, Kozma-Bognar L, Terecskei K, De Veylder L, Nagy F, Ulm R (2014) UV-B-responsive association of the Arabidopsis bZIP transcription factor ELONGATED HYPOCOTYL5 with target genes, including its own promoter. Plant Cell 26: 4200-4213

Bowman JL, Kohchi T, Yamato KT, Jenkins J, Shu S, Ishizaki K, Yamaoka S, Nishihama R, Nakamura Y, Berger F, Adam C, Aki SS, Althoff F, Araki T, Arteaga-Vazquez MA, Balasubrmanian S, Barry K, Bauer D, Boehm CR, Briginshaw L, Caballero-Perez J, Catarino B, Chen F, Chiyoda S, Chovatia M, Davies KM, Delmans M, Demura T, Dierschke T, Dolan L, Dorantes-Acosta AE, Eklund DM, Florent SN, Flores-Sandoval E, Fujiyama A, Fukuzawa H, Galik B, Grimanelli D, Grimwood J, Grossniklaus U, Hamada

T, Haseloff J, Hetherington AJ, Higo A, Hirakawa Y, Hundley HN, Ikeda Y, Inoue K, Inoue SI, Ishida S, Jia Q, Kakita M, Kanazawa T, Kawai Y, Kawashima T, Kennedy M, Kinose K, Kinoshita T, Kohara Y, Koide E, Komatsu K, Kopischke S, Kubo M, Kyozuka J, Lagercrantz U, Lin SS, Lindquist E, Lipzen AM, Lu CW, De Luna E, Martienssen RA, Minamino N, Mizutani M, Mizutani M, Mochizuki N, Monte I, Mosher R, Nagasaki H, Nakagami H, Naramoto S, Nishitani K, Ohtani M, Okamoto T, Okumura M, Phillips J, Pollak B, Reinders A, Rovekamp M, Sano R, Sawa S, Schmid MW, Shirakawa M, Solano R, Spunde A, Suetsugu N, Sugano S, Sugiyama A, Sun R, Suzuki Y, Takenaka M, Takezawa D, Tomogane H, Tsuzuki M, Ueda T, Umeda M, Ward JM, Watanabe Y, Yazaki K, Yokoyama R, Yoshitake Y, Yotsui I, Zachgo S, Schmutz J (2017) Insights into land plant evolution garnered from the *Marchantia polymorpha* genome. Cell 171: 287-304 e215

Brown BA, Cloix C, Jiang GH, Kaiserli E, Herzyk P, Kliebenstein DJ, Jenkins GI (2005) A UV-B-specific signaling component orchestrates plant UV protection. Proc Natl Acad Sci USA 102: 18225-18230

Chen H, Huang X, Gusmaroli G, Terzaghi W, Lau OS, Yanagawa Y, Zhang Y, Li J, Lee JH, Zhu D, Deng XW (2010) Arabidopsis CULLIN4-damaged DNA binding protein 1 interacts with CONSTITUTIVELY PHOTOMORPHOGENIC1-SUPPRESSOR OF PHYA complexes to regulate photomorphogenesis and flowering time. Plant Cell 22: 108-123

Chiyoda S, Ishizaki K, Kataoka H, Yamato KT, Kohchi T (2008) Direct transformation of the liverwort *Marchantia polymorpha* L. by particle bombardment using immature thalli developing from spores. Plant Cell Rep 27: 1467-1473

Christie JM, Arvai AS, Baxter KJ, Heilmann M, Pratt AJ, O'Hara A, Kelly SM, Hothorn M, Smith BO, Hitomi K, Jenkins GI, Getzoff ED (2012) Plant UVR8 photoreceptor senses UV-B by tryptophan-mediated disruption of cross-dimer salt bridges. Science 335: 1492-1496

Clayton WA, Albert NW, Thrimawithana AH, McGhie TK, Deroles SC, Schwinn KE, Warren BA, McLachlan ARG, Bowman JL, Jordan BR, Davies KM (2018) UVR8-mediated induction of flavonoid biosynthesis for UVB tolerance is conserved between the liverwort *Marchantia polymorpha* and flowering plants. Plant J 96: 503-517

Cox J, Mann M (2008) MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol 26: 1367-1372

Dannay M, Bertin C, Cavallari E, Albanese P, Tolleter D, Giustini C, Menneteau M, Brugiere S, Coute Y, Finazzi G, Demarsy E, Ulm R, Allorent G (2025) Photoreceptor-induced LHL4 protects the photosystem II monomer in *Chlamydomonas reinhardtii*. Proc Natl Acad Sci USA 122: e2418687122

Davies KM, Landi M, van Klink JW, Schwinn KE, Brummell DA, Albert NW, Chagne D, Jibran R, Kulshrestha S, Zhou Y, Bowman JL (2022) Evolution and function of red pigmentation in land plants. Ann Bot 130: 613-636

Defossez E, Bourquin J, von Reuss S, Rasmann S, Glauser G (2023) Eight key rules for successful data-dependent acquisition in mass spectrometry-based metabolomics. Mass Spectrom Rev 42: 131-143

Demarsy E, Goldschmidt-Clermont M, Ulm R (2018) Coping with 'dark sides of the sun' through photoreceptor signaling. Trends Plant Sci 23: 260-271

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29: 15-21

Duhrkop K, Fleischauer M, Ludwig M, Aksenov AA, Melnik AV, Meusel M, Dorrestein PC, Rousu J, Bocker S (2019) SIRIUS 4: a rapid tool for turning tandem mass spectra into metabolite structure information. Nat Methods 16: 299-302

Fang F, Lin L, Zhang Q, Lu M, Skvortsova MY, Podolec R, Zhang Q, Pi J, Zhang C, Ulm R, Yin R (2022) Mechanisms of UV-B light-induced photoreceptor UVR8 nuclear localization dynamics. New Phytol 236: 1824-1837

Favory JJ, Stec A, Gruber H, Rizzini L, Oravecz A, Funk M, Albert A, Cloix C, Jenkins GI, Oakeley EJ, Seidlitz HK, Nagy F, Ulm R (2009) Interaction of COP1 and UVR8 regulates UV-B-induced photomorphogenesis and stress acclimation in Arabidopsis. EMBO J 28: 591-601

Ferreyra MLF, Serra P, Casati P (2021) Recent advances on the roles of flavonoids as plant protective molecules after UV and high light exposure. Physiol Plant 173: 736-749

Findlay KM, Jenkins GI (2016) Regulation of UVR8 photoreceptor dimer/monomer photo-equilibrium in Arabidopsis plants grown under photoperiodic conditions. Plant Cell Environ 39: 1706-1714

Frangedakis E, Yelina NE, Billakurthi K, Hua L, Schreier T, Dickinson PJ, Tomaselli M, Haseloff J, Hibberd JM (2024) MYB-related transcription factors control chloroplast biogenesis. Cell 187: 4859-4876 e4822

Furudate H, Manabe M, Oshikiri H, Matsushita A, Watanabe B, Waki T, Nakayama T, Kubo H, Takanashi K (2023) A polyphenol oxidase catalyzes aurone synthesis in *Marchantia polymorpha*. Plant Cell Physiol 64: 637-645

Gabilly ST, Baker CR, Wakao S, Crisanto T, Guan K, Bi K, Guiet E, Guadagno CR, Niyogi KK (2019) Regulation of photoprotection gene expression in Chlamydomonas by a putative E3 ubiquitin ligase complex and a homolog of CONSTANS. Proc Natl Acad Sci USA 116: 17556-17562

Gietz R, Smith J, Burke D (2014) Yeast genetics: methods and protocols.

Gruber H, Heijde M, Heller W, Albert A, Seidlitz HK, Ulm R (2010) Negative feedback regulation of UV-B-induced photomorphogenesis and stress acclimation in Arabidopsis. Proc Natl Acad Sci USA 107: 20132-20137

Heijde M, Binkert M, Yin R, Ares-Orpel F, Rizzini L, Van De Slijke E, Persiau G, Nolf J, Gevaert K, De Jaeger G, Ulm R (2013) Constitutively active UVR8 photoreceptor variant in Arabidopsis. Proc Natl Acad Sci USA 110: 20326-20331

Heijde M, Ulm R (2013) Reversion of the Arabidopsis UV-B photoreceptor UVR8 to the homodimeric ground state. Proc Natl Acad Sci USA 110: 1113-1118

Heilmann M, Jenkins GI (2013) Rapid reversion from monomer to dimer regenerates the ultraviolet-B photoreceptor UV RESISTANCE LOCUS8 in intact Arabidopsis plants. Plant Physiol 161: 547-555

Heilmann M, Velanis CN, Cloix C, Smith BO, Christie JM, Jenkins GI (2016) Dimer/monomer status and in vivo function of salt-bridge mutants of the plant UV-B photoreceptor UVR8. Plant J 88: 71-81

Hoecker U (2017) The activities of the E3 ubiquitin ligase COP1/SPA, a key repressor in light signaling. Curr Opin Plant Biol 37: 63-69

Holm M, Ma LG, Qu LJ, Deng XW (2002) Two interacting bZIP proteins are direct targets of COP1-mediated control of light-dependent gene expression in Arabidopsis. Genes Dev 16: 1247-1259

Hu S, Chen Y, Qian C, Ren H, Liang X, Tao W, Chen Y, Wang J, Dong Y, Han J, Ouyang X, Huang X (2024) Nuclear accumulation of rice UV-B photoreceptors is UV-B- and OsCOP1-independent for UV-B responses. Nat Commun 15: 6396

Huang X, Ouyang X, Yang P, Lau OS, Chen L, Wei N, Deng XW (2013) Conversion from CUL4-based COP1-SPA E3 apparatus to UVR8-COP1-SPA complexes underlies a distinct biochemical function of COP1 under UV-B. Proc Natl Acad Sci USA 110: 16669-16674

Ishizaki K, Chiyoda S, Yamato KT, Kohchi T (2008) Agrobacterium-mediated transformation of the haploid liverwort *Marchantia polymorpha* L., an emerging model for plant biology. Plant Cell Physiol 49: 1084-1091

Jibran R, Hill SJ, Lampugnani ER, Hao P, Doblin MS, Bacic A, Vaidya AA, O'Donoghue EM, McGhie TK, Albert NW, Zhou Y, Raymond LG, Schwinn KE, Jordan BR, Bowman JL, Davies KM, Brummell DA (2024) The auronidin flavonoid pigments of the liverwort *Marchantia polymorpha* form polymers that modify cell wall properties. Plant J 120: 1159-1175

Kaiserli E, Jenkins GI (2007) UV-B promotes rapid nuclear translocation of the Arabidopsis UV-B specific signaling component UVR8 and activates its function in the nucleus. Plant Cell 19: 2662-2673

Kliebenstein DJ, Lim JE, Landry LG, Last RL (2002) Arabidopsis UVR8 regulates ultraviolet-B signal
transduction and tolerance and contains sequence similarity to human regulator of chromatin
condensation 1. Plant Physiol 130: 234-243
Kondou Y, Miyagi Y, Morito T, Fujihira K, Miyauchi W, Moriyama A, Terasawa T, Ishida S, Iwabuchi K, Kubo
H, Nishihama R, Ishizaki K, Kohchi T (2019) Physiological function of photoreceptor UVR8 in UV-B
tolerance in the liverwort *Marchantia polymorpha*. Planta 249: 1349-1364
Kreiss M, Haas FB, Hansen M, Rensing SA, Hoecker U (2023) Co-action of COP1, SPA and cryptochrome in
light signal transduction and photomorphogenesis of the moss *Physcomitrium patens*. Plant J 114:
159-175
Kubota A, Ishizaki K, Hosaka M, Kohchi T (2013) Efficient Agrobacterium-mediated transformation of the
liverwort *Marchantia polymorpha* using regenerating thalli. Biosci Biotechnol Biochem 77: 167-
172
Lau K, Podolec R, Chappuis R, Ulm R, Hothorn M (2019) Plant photoreceptors and their signaling
components compete for COP1 binding via VP peptide motifs. EMBO J 38: e102140
Laubinger S, Fittinghoff K, Hoecker U (2004) The SPA quartet: a family of WD-repeat proteins with a central
role in suppression of photomorphogenesis in arabidopsis. Plant Cell 16: 2293-2306
Leonardelli M, Tissot N, Podolec R, Ares-Orpel F, Glauser G, Ulm R, Demarsy E (2024) Photoreceptor-
induced sinapate synthesis contributes to photoprotection in Arabidopsis. Plant Physiol 196: 1518-
1533
Li H, Li Y, Deng H, Sun X, Wang A, Tang X, Gao Y, Zhang N, Wang L, Yang S, Liu Y, Wang S (2018) Tomato
UV-B receptor SlUVR8 mediates plant acclimation to UV-B radiation and enhances fruit chloroplast
development via regulating SlGLK2. Sci Rep 8: 6097
Li X, Ren H, Kundu M, Liu Z, Zhong FW, Wang L, Gao J, Zhong D (2020) A leap in quantum efficiency through
light harvesting in photoreceptor UVR8. Nat Commun 11: 4316
Liang Y, Heyman J, Xiang Y, Vandendriessche W, Canher B, Goeminne G, De Veylder L (2022) The wound-
activated ERF15 transcription factor drives *Marchantia polymorpha* regeneration by activating an
oxylipin biosynthesis feedback loop. Sci Adv 8: eabo7737
Liao Y, Smyth GK, Shi W (2019) The R package Rsubread is easier, faster, cheaper and better for alignment
and quantification of RNA sequencing reads. Nucleic Acids Res 47: e47
Liu W, Giuriani G, Havlikova A, Li D, Lamont DJ, Neugart S, Velanis CN, Petersen J, Hoecker U, Christie JM,
Jenkins GI (2024) Phosphorylation of Arabidopsis UVR8 photoreceptor modulates protein
interactions and responses to UV-B radiation. Nat Commun 15: 1221
Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR
and the 2(-Delta Delta C(T)) Method. Methods 25: 402-408
Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. Genome Biol 15: 550
Marrocco K, Zhou Y, Bury E, Dieterle M, Funk M, Genschik P, Krenz M, Stolpe T, Kretsch T (2006) Functional
analysis of EID1, an F-box protein involved in phytochrome A-dependent light signal transduction.
Plant J 45: 423-438
McNellis TW, von Arnim AG, Araki T, Komeda Y, Misera S, Deng XW (1994) Genetic and molecular analysis
of an allelic series of *cop1* mutants suggests functional roles for the multiple protein domains.
Plant Cell 6: 487-500
Messant M, Hani U, Hennebelle T, Guerard F, Gakiere B, Gall A, Thomine S, Krieger-Liszkay A (2023)
Manganese concentration affects chloroplast structure and the photosynthetic apparatus in
Marchantia polymorpha. Plant Physiol 192: 356-369
Moriconi V, Binkert M, Costigliolo C, Sellaro R, Ulm R, Casal JJ (2018) Perception of sunflecks by the UV-B
photoreceptor UV RESISTANCE LOCUS8. Plant Physiol 177: 75-81

Mulvey H, Dolan L (2023) RHO GTPase of plants regulates polarized cell growth and cell division orientation
during morphogenesis. Curr Biol 33: 2897-2911 e2896
Murchie EH, Lawson T (2013) Chlorophyll fluorescence analysis: a guide to good practice and
understanding some new applications. J Exp Bot 64: 3983-3998
Oravecz A, Baumann A, Mate Z, Brzezinska A, Molinier J, Oakeley EJ, Adam E, Schafer E, Nagy F, Ulm R
(2006) CONSTITUTIVELY PHOTOMORPHOGENIC1 is required for the UV-B response in Arabidopsis.
Plant Cell 18: 1975-1990
Ordonez-Herrera N, Fackendahl P, Yu X, Schaefer S, Koncz C, Hoecker U (2015) A *cop1 spa* mutant deficient
in COP1 and SPA proteins reveals partial co-action of COP1 and SPA during Arabidopsis post-
embryonic development and photomorphogenesis. Mol Plant 8: 479-481
Pham VN, Paik I, Hoecker U, Huq E (2020) Genomic evidence reveals SPA-regulated developmental and
metabolic pathways in dark-grown Arabidopsis seedlings. Physiol Plant 169: 380-396
Podolec R, Demarsy E, Ulm R (2021) Perception and signaling of ultraviolet-B radiation in plants. Annu Rev
Plant Biol 72: 793-822
Podolec R, Lau K, Wagnon TB, Hothorn M, Ulm R (2021) A constitutively monomeric UVR8 photoreceptor
confers enhanced UV-B photomorphogenesis. Proc Natl Acad Sci USA 118: e2017284118
Podolec R, Ulm R (2018) Photoreceptor-mediated regulation of the COP1/SPA E3 ubiquitin ligase. Curr
Opin Plant Biol 45: 18-25
Ponnu J, Hoecker U (2021) Illuminating the COP1/SPA ubiquitin ligase: fresh insights into its structure and
functions during plant photomorphogenesis. Front Plant Sci 12: 662793
Ponnu J, Riedel T, Penner E, Schrader A, Hoecker U (2019) Cryptochrome 2 competes with COP1 substrates
to repress COP1 ubiquitin ligase activity during Arabidopsis photomorphogenesis. Proc Natl Acad
Sci USA 116: 27133-27141
Qian C, Chen Z, Liu Q, Mao W, Chen Y, Tian W, Liu Y, Han J, Ouyang X, Huang X (2020) Coordinated
transcriptional regulation by the UV-B photoreceptor and multiple transcription factors for plant
UV-B responses. Mol Plant 13: 777-792
Rai N, Neugart S, Yan Y, Wang F, Siipola SM, Lindfors AV, Winkler JB, Albert A, Brosche M, Lehto T, Morales
LO, Aphalo PJ (2019) How do cryptochromes and UVR8 interact in natural and simulated sunlight?
J Exp Bot 70: 4975-4990
Ranjan A, Dickopf S, Ullrich KK, Rensing SA, Hoecker U (2014) Functional analysis of COP1 and SPA
orthologs from Physcomitrella and rice during photomorphogenesis of transgenic Arabidopsis
reveals distinct evolutionary conservation. BMC Plant Biol 14: 178
Rappsilber J, Ishihama Y, Mann M (2003) Stop and go extraction tips for matrix-assisted laser
desorption/ionization, nanoelectrospray, and LC/MS sample pretreatment in proteomics. Anal
Chem 75: 663-670
Rizzini L, Favory JJ, Cloix C, Faggionato D, O'Hara A, Kaiserli E, Baumeister R, Schafer E, Nagy F, Jenkins GI,
Ulm R (2011) Perception of UV-B by the Arabidopsis UVR8 protein. Science 332: 103-106
Rutz A, Allard P (2025) tima: taxonomically informed metabolite annotation (2.11.0) [R package]. Zenodo:
https://doi.org/10.5281/zenodo.5797920
Saint-Marcoux D, Proust H, Dolan L, Langdale JA (2015) Identification of reference genes for real-time
quantitative PCR experiments in the liverwort *Marchantia polymorpha*. PLoS One 10: e0118678
Sauret-Güeto S, Frangedakis E, Silvestri L, Rebmann M, Tomaselli M, Markel K, Delmans M, West A, Patron
NJ, Haseloff J (2020) Systematic tools for reprogramming plant gene expression in a simple model,
Marchantia polymorpha. ACS Synth Biol 9: 864-882
Sawano A, Miyawaki A (2000) Directed evolution of green fluorescent protein by a new versatile PCR
strategy for site-directed and semi-random mutagenesis. Nucleic Acids Res 28: E78
Schmid R, Heuckeroth S, Korf A, Smirnov A, Myers O, Dyrlund TS, Bushuiev R, Murray KJ, Hoffmann N, Lu
M, Sarvepalli A, Zhang Z, Fleischauer M, Duhrkop K, Wesner M, Hoogstra SJ, Rudt E, Mokshyna O,

Brungs C, Ponomarov K, Mutabdzija L, Damiani T, Pudney CJ, Earll M, Helmer PO, Fallon TR, Schulze
T, Rivas-Ubach A, Bilbao A, Richter H, Nothias LF, Wang M, Oresic M, Weng JK, Bocker S, Jeibmann
A, Hayen H, Karst U, Dorrestein PC, Petras D, Du X, Pluskal T (2023) Integrative analysis of
multimodal mass spectrometry data in MZmine 3. Nat Biotechnol 41: 447-449
Soriano G, Cloix C, Heilmann M, Nunez-Olivera E, Martinez-Abaigar J, Jenkins GI (2018) Evolutionary
conservation of structure and function of the UVR8 photoreceptor from the liverwort *Marchantia*
polymorpha and the moss *Physcomitrella patens*. New Phytol 217: 151-162
Stelzl U, Worm U, Lalowski M, Haenig C, Brembeck FH, Goehler H, Stroedicke M, Zenkner M, Schoenherr
A, Koeppen S, Timm J, Mintzlaff S, Abraham C, Bock N, Kietzmann S, Goedde A, Toksoz E, Droege
A, Krobitsch S, Korn B, Birchmeier W, Lehrach H, Wanker EE (2005) A human protein-protein
interaction network: a resource for annotating the proteome. Cell 122: 957-968
Stockenhuber R, Akiyama R, Tissot N, Milosavljevic S, Yamazaki M, Wyler M, Arongaus AB, Podolec R, Sato
Y, Widmer A, Ulm R, Shimizu KK (2024) UV RESISTANCE LOCUS 8-mediated UV-B response is
required alongside CRYPTOCHROME 1 for plant survival in sunlight under field conditions. Plant
Cell Physiol 65: 35-48
Stracke R, Favory JJ, Gruber H, Bartelniewoehner L, Bartels S, Binkert M, Funk M, Weisshaar B, Ulm R (2010)
The Arabidopsis bZIP transcription factor HY5 regulates expression of the *PFG1*/*MYB12* gene in
response to light and ultraviolet-B radiation. Plant Cell Environ 33: 88-103
Sugano SS, Nishihama R, Shirakawa M, Takagi J, Matsuda Y, Ishida S, Shimada T, Hara-Nishimura I, Osakabe
K, Kohchi T (2018) Efficient CRISPR/Cas9-based genome editing and its application to conditional
genetic analysis in *Marchantia polymorpha*. PLoS One 13: e0205117
Tavridou E, Pireyre M, Ulm R (2020) Degradation of the transcription factors PIF4 and PIF5 under UV-B
promotes UVR8-mediated inhibition of hypocotyl growth in Arabidopsis. Plant J 101: 507-517
Tavridou E, Schmid-Siegert E, Fankhauser C, Ulm R (2020) UVR8-mediated inhibition of shade avoidance
involves HFR1 stabilization in Arabidopsis. PLoS Genet 16: e1008797
Tilbrook K, Dubois M, Crocco CD, Yin R, Chappuis R, Allorent G, Schmid-Siegert E, Goldschmidt-Clermont
M, Ulm R (2016) UV-B perception and acclimation in *Chlamydomonas reinhardtii*. Plant Cell 28:
966-983
Tokutsu R, Fujimura-Kamada K, Matsuo T, Yamasaki T, Minagawa J (2019) The CONSTANS flowering
complex controls the protective response of photosynthesis in the green alga Chlamydomonas.
Nat Commun 10: 4099
Trimborn L, Kuttig F, Ponnu J, Yu P, Korsching KR, Lederer P, Urquiza-Garcia U, Zurbriggen MD, Hoecker U
(2025) Cryptochrome 1 promotes photomorphogenesis in Arabidopsis by displacing substrates
from the COP1 ubiquitin ligase. Plant J 121: e70071
Tyanova S, Temu T, Cox J (2016) The MaxQuant computational platform for mass spectrometry-based
shotgun proteomics. Nat Protoc 11: 2301-2319
Ulm R, Baumann A, Oravecz A, Mate Z, Adam E, Oakeley EJ, Schafer E, Nagy F (2004) Genome-wide analysis
of gene expression reveals function of the bZIP transcription factor HY5 in the UV-B response of
Arabidopsis. Proc Natl Acad Sci USA 101: 1397-1402
Wang L, Wang Y, Chang H, Ren H, Wu X, Wen J, Guan Z, Ma L, Qiu L, Yan J, Zhang D, Huang X, Yin P (2023)
RUP2 facilitates UVR8 redimerization via two interfaces. Plant Commun 4: 100428
Wang W, Paik I, Kim J, Hou X, Sung S, Huq E (2021) Direct phosphorylation of HY5 by SPA kinases to regulate
photomorphogenesis in Arabidopsis. New Phytol 230: 2311-2326
Wang Y, Wang L, Guan Z, Chang H, Ma L, Shen C, Qiu L, Yan J, Zhang D, Li J, Deng XW, Yin P (2022) Structural
insight into UV-B-activated UVR8 bound to COP1. Sci Adv 8: eabn3337
Wu D, Hu Q, Yan Z, Chen W, Yan C, Huang X, Zhang J, Yang P, Deng H, Wang J, Deng X, Shi Y (2012) Structural
basis of ultraviolet-B perception by UVR8. Nature 484: 214-219

Yelina NE, Frangedakis E, Wang Z, Schreier TB, Rever J, Tomaselli M, Forestier ECF, Billakurthi K, Ren S, Bai
Y, Stewart-Wood J, Haseloff J, Zhong S, Hibberd JM (2024) Streamlined regulation of chloroplast
development in the liverwort *Marchantia polymorpha*. Cell Rep 43: 114696
Yin R, Arongaus AB, Binkert M, Ulm R (2015) Two distinct domains of the UVR8 photoreceptor interact
with COP1 to initiate UV-B signaling in Arabidopsis. Plant Cell 27: 202-213
Yin R, Skvortsova MY, Loubery S, Ulm R (2016) COP1 is required for UV-B-induced nuclear accumulation of
the UVR8 photoreceptor. Proc Natl Acad Sci USA 113: E4415-4422
Zhang C, Zhang Q, Guo H, Yu X, Liang W, Chen Y, Yin R, Lin L (2021) Tomato SlRUP is a negative regulator
of UV-B photomorphogenesis. Mol Hortic 1: 8
Zhang L, Li T, Su S, Peng H, Li S, Li K, Ji L, Xing Y, Zhang J, Du X, Bian M, Liao Y, Yang Z, Zuo Z (2021) Functions
of COP1/SPA E3 ubiquitin ligase mediated by MpCRY in the liverwort *Marchantia polymorpha*
under blue light. Int J Mol Sci 23: 158
Zhou Z, Bi G, Zhou JM (2018) Luciferase complementation assay for protein-protein interactions in plants.
Curr Protoc Plant Biol 3: 42-50

1184

Figure 1. MpUVR8.1 and MpUVR8.2 form dimers in their ground state, monomerize upon UV-B exposure, and accumulate in the nucleus.

(A-C) Luciferase complementation imaging (LCI) assays assessing monomer interaction for MpUVR8.1 (A) and MpUVR8.2 (B), as well as their D95N,D106N variants, and MpUVR8.1 and MpUVR8.2 (C). MpUVR8 versions were fused to N-terminal (nLUC) or C-terminal (cLUC) fragments of luciferase, expressed in *N. benthamiana* leaves and imaged 2 days after infiltration. An image of a representative leaf is shown. Cps, counts per second.

(D) Immunoblot analysis of MpUVR8 dimer/monomer levels in 7-day-old wild type (Tak-1) gemmalings irradiated (+UV) or not (-UV) for 20 min with broadband UV-B. Protein extracts were either crosslinked with DSP, DSP crosslinked then reversed with the reducing agent β-mercaptoethanol (DSP + β-ME), or non-treated.

(E) Subcellular localization of GFP-MpUVR8 in 7-day-old gemmalings of a Tak-1/*Pro*$_{MpUVR8}$:*GFP*-Mp*UVR8* line, grown in white light and irradiated or not for 20 min with 25 μmol m^{-2} s^{-1} broadband UV-B (+UV and -UV, respectively). Bars = 200 μm.

Figure 2. MpRUP facilitates MpUVR8 redimerization.

(A) Yeast two-hybrid assays of MpUVR8.1 and MpUVR8.2 interactions with MpRUP. Representative images of serial dilutions spotted on non-selective SD/-Trp/-Leu double dropout (DDO) and selective SD/-Trp/-Leu/-His triple dropout (TDO) media are shown. AD, transcriptional activation domain; BD, DNA-binding domain; EV, empty vector.

(B) Quantitative yeast two-hybrid assays of MpUVR8.1 and MpUVR8.2 interactions with MpRUP. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Different letters indicate statistically significant differences between the means (P < 0.01) as determined by one-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test. β-Gal., β-galactosidase; MU, Miller units.

(C,D) LCI assays of MpRUP interactions with MpUVR8.1 (C) and MpUVR8.2 (D) fused to N-terminal (nLUC) or C-terminal (cLUC) luciferase fragments, expressed in *N. benthamiana* leaves and imaged 2 days after infiltration. Images of representative leaves are shown. Cps, counts per second.

(E) Immunoblot analysis of MpUVR8 levels in 7-day-old wild type (Tak-1) and Mp*rup-2* gemmalings irradiated for 20 min with broadband UV-B (+UV) before recovery in white light (WL) for the indicated times (1–6 h). Non-irradiated samples (-UV) are included as MpUVR8 ground state controls alongside a negative control -UV Mp*uvr8-1* sample. Top: MpUVR8 dimer levels in DSP-crosslinked protein extracts; bottom: total MpUVR8 levels in non-crosslinked extracts.

Figure 3. UV-B acclimation and survival is impaired in Mp*uvr8* and enhanced in Mp*rup*.

Figure 3. UV-B acclimation and survival is impaired in Mp*uvr8* and enhanced in Mp*rup*.

(A) Growth phenotype of 4-week-old wild-type (Tak-1), Mp*uvr8-1*, and Mp*rup-2* grown in white light supplemented (WL+UV) or not (WL-UV) with UV-B (1.5 µmol m^{-2} s^{-1}). Bars = 1 cm.

(B) Scheme for UV-B treatment and sampling for F_v/F_m measurements. Light yellow bar: white light (WL, 50 µmol m^{-2} s^{-1}), non-acclimated condition; light blue bar: white light supplemented with narrowband UV-B (WL+UVB, 1.5 µmol m^{-2} s^{-1}), acclimated condition; light/dark purple bars: broadband UV stress (UV BB, 25 µmol m^{-2} s^{-1}). Red stars indicate timepoints of measurements: -/-, non-acclimated, non-stressed; -/+, non-acclimated, stressed; +/-, acclimated, non-stressed; +/+, acclimated, stressed.

(C) F_v/F_m in 7-day-old wild-type (Tak-1), Mp*uvr8-1*, and Mp*rup-2* gemmalings grown and sampled as indicated in (B). Individual data points of biological replicates and means ± SD are shown (*n* = 3). Different letters indicate statistically significant differences between the means (P < 0.01) as determined by two-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

(D) Scheme for UV-B stress survival experiments. Plants were grown in WL or WL+UVB for 7 days, then exposed to 7 or 9 h of broadband UV-B stress, followed by recovery for 7 days in WL, after which they were imaged.

(E) UV-B stress survival in Tak-1, Mp*uvr8-1*, and Mp*rup-2* gemmalings grown as indicated in (D). Representative images are shown; bars = 1 cm.

(F) Absorbance spectra (280–400 nm) of methanolic extracts from 9-day-old Tak-1, Mp*uvr8-1*, and Mp*rup-2* gemmalings grown for 7 days in white light and exposed (WL+UV-B, purple lines) or not (WL, blue lines) to supplemental UV-B (1.5 µmol m^{-2} s^{-1}) for 2 days. The gray line shows the absorbance of the buffer (blank). Values of independent measurements and means ± SEM are shown (*n* = 3).

Figure 4. MpUVR8 and MpCOP1 interact in a UV-B-dependent manner.

(**A**) Yeast two-hybrid assay of the interactions of MpCOP1 with MpSPA and MpUVR8, as well as AtCOP1 with MpUVR8, in the presence (+UV) or absence (-UV) of UV-B. Left, growth assay; right, quantitative assay. Yeasts were grown on non-selective SD/-Trp/-Leu (DDO) or selective SD /-Trp/-Leu/-His (TDO) media. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Different letters indicate statistically significant differences between the means (P < 0.01) as determined by one-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test. AD, Gal4 activation domain; BD, LexA DNA-binding domain; β-Gal., β-galactosidase; MU, Miller units; EV, empty vector.

(**B**) LCI assays of MpCOP1 with MpUVR8.1 and MpUVR8.2. N-terminal (nLUC) or C-terminal (cLUC) luciferase fusion proteins were expressed in *N. benthamiana* leaves and imaged 2 days after infiltration, with (+UV) or without (-UV) exposure to 20 min of broadband UV-B. An image of a representative leaf is shown.

Figure 5. Mp*cop1* shows constitutive photomorphogenesis and UV-B tolerance, whereas Mp*spa* shows enhanced UV-B-induced UV-B tolerance.
(**A**) Growth phenotype of 4-week-old wild-type (Tak-1), Mp*cop1-1*, Mp*cop1-2*, Mp*spa-9*, and Mp*spa-10* plants grown in white light supplemented (WL +UV) or not (WL -UV) with UV-B. Scale bars = 1 cm.
(**B**) F_v/F_m measurements in 7-day-old wild-type (Tak-1), Mp*cop1-1*, *Mpcop1-2*, Mp*spa-9*, and Mp*spa-10* gemmalings grown in white light (non-acclimated) or white light supplemented with UV-B (acclimated), without (non-stressed) or with (stressed) additional exposure to broadband UV-B stress for 60 min. -/-, non-acclimated, non-stressed; -/+, non-acclimated, stressed; +/-, acclimated, non-stressed; +/+, acclimated, stressed. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Different letters indicate statistically significant differences between the means (P < 0.01) as determined by two-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.
(**C**) UV-B stress survival in wild-type (Tak-1), Mp*cop1-1*, Mp*cop1-2*, Mp*spa-9*, and Mp*spa-10* plants grown for 7 days in white light supplemented (acclimated) or not (non-acclimated) with UV-B, then exposed or not to 9 h of broadband UV-B stress. Pictures were taken after a further 7-day recovery period in white light.
(**D**) Absorbance spectra (280–400 nm) of methanolic extracts from 9-day-old wild-type (Tak-1), Mp*cop1-1,* Mp*cop1-2*, Mp*spa-9*, and Mp*spa-10* gemmalings grown for 7 days in white light supplemented (WL+UV-B, purple lines) or not (WL, blue lines) with UV-B. The gray line shows the absorbance of the buffer (blank). Values of independent measurements and means ± SEM are shown (*n* = 3).

Figure 6. Transcriptome analysis reveals MpUVR8- and MpCOP1-dependent, UV-B-responsive gene regulation that is repressed by MpRUP and MpSPA

(**A**) Euler plots showing overlaps between upregulated differentially expressed genes (DEGs) (p-adj < 0.05) upon exposure to 3 h UV-B in 7-day-old wild-type (Tak-1), Mp*rup*, and Mp*spa* gemmalings. Left: overlap between genes upregulated at least two-fold ($\log_2$FC > 1) in either Tak-1, Mp*rup*, or Mp*spa*. Right: overlap between upregulated ($\log_2$FC > 1) genes that are unique to Mp*rup* and/or Mp*spa*, and Tak-1 upregulated genes without a fold-change threshold ($\log_2$FC > 0). Corresponding gene lists: Supplementary Table S3a.

(**B**) Euler plots showing overlaps between downregulated DEGs (p-adj < 0.05) upon exposure to 3 h UV-B in 7-day-old Tak-1, Mp*rup*, and Mp*spa* gemmalings. Left: overlap between genes downregulated at least two-fold ($\log_2$FC < -1) in either Tak-1, Mp*rup*, or Mp*spa*. Right: overlap between downregulated ($\log_2$FC < -1) genes that are unique to Mp*rup* and/or Mp*spa*, and Tak-1 downregulated genes without a fold-change threshold ($\log_2$FC < 0). Corresponding gene lists: Supplementary Table S3b.

(**C**) Hierarchical clustering and heatmap of UV-B-regulated gene expression in Tak-1, Mp*uvr8-1,* Mp*rup-2,* Mp*cop1-1*, and Mp*spa-9* plants grown for 7 days under white light and exposed or not to supplemental UV-B for 3 h. All genes with a greater than two-fold change in expression in response to UV-B and a p-adj < 0.05 in at least one genotype are shown (total = 1005 genes). Red and blue colors indicate normalized read count Z-scores. The color bar on the left side indicates 7 major profiles (clusters) of gene expression.

(**D**) Gene expression profiles in clusters identified by hierarchical clustering. Z-scores for normalized read count values for each gene (gray lines) and for the cluster average (colored line) are shown.

Figure 7. MpSPA plays a minor role in repressing Marchantia photomorphogenesis under visible light.

(**A**) Representative images of 4-day-old Arabidopsis seedlings of wild-type (Col-0), *cop1-4*, and *spaQn* mutants, and 28-day-old Marchantia thalli of wild-type (Tak-1), Mp*cop1-1*, and Mp*spa-10* grown in white light. Scale bars: 4 mm or 1 cm, as indicated.

(**B**) Euler plots showing overlaps between upregulated DEGs (p-adj < 0.05) in Mp*cop1* and Mp*spa* mutants compared to Tak-1 wild type. Gemmalings were grown for 7 days in white light. Top: overlap between genes upregulated at least two-fold ($log_2FC > 1$) in either Mp*cop1*/Tak-1 or Mp*spa*/Tak-1. Bottom: overlap between genes upregulated at least two-fold ($log_2FC > 1$) in Mp*cop1*/Tak-1 and genes upregulated without a fold-change threshold ($log_2FC > 0$) in Mp*spa*/Tak-1. Corresponding gene lists: Supplementary Table S3c.

(**C**) Euler plots showing overlaps between downregulated DEGs (p-adj < 0.05) in Mp*cop1* and Mp*spa* mutants compared to Tak-1 wild type. Gemmalings were grown for 7 days in white light. Top: overlap between genes downregulated at least two-fold ($log_2FC < -1$) in either Mp*cop1*/Tak-1 or Mp*spa*/Tak-1. Bottom: overlap between genes downregulated at least two-fold ($log_2FC < -1$) in Mp*cop1*/Tak-1 and genes downregulated without a fold-change threshold ($log_2FC < 0$) in Mp*spa*/Tak-1. Corresponding gene lists: Supplementary Table S3d.

(**D**) Hierarchical clustering and heatmap of gene expression in Tak-1, Mp*cop1-1*, and Mp*spa-9* plants grown for 7 days in white light. All genes with a greater than two-fold change in expression between mutants and wild type and a p-adj < 0.05 in at least one comparison are shown (total = 2975 genes). Red and blue colors indicate normalized read count Z-scores. The color bar on the left side indicates 11 major profiles (clusters) of gene expression. The color bars on the right indicate if genes are significantly regulated after UV-B exposure in Tak-1, Mp*rup*, and Mp*spa*.

(**E**) Gene expression profiles in clusters identified by hierarchical clustering. Z-scores for normalized read count values for each gene are shown as gray points; their distribution is represented as a colored box plot.

Figure 8. MpHY5 is involved in UV-B and visible light photomorphogenesis.

(**A**) F_v/F_m measurements in wild-type (Tak-1), Mp*hy5-1*, Mp*hy5-3*, and Mp*hy5-7* plants grown for 7 days in white light (non-acclimated) or white light supplemented with UV-B (acclimated), without (not stressed) or with (stressed) additional exposure to broadband UV-B stress for 60 min. -/-, not acclimated, not stressed; -/+, not acclimated, stressed; +/-, acclimated, not stressed; +/+, acclimated, stressed. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Letters indicate statistically significant differences between the means (P < 0.01) as determined by two-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

(**B**) UV-B stress survival of Tak-1, Mp*hy5-1*, Mp*hy5-3*, and Mp*hy5-7* plants grown for 7 days in white light supplemented (Accl.) or not (Non-accl.) with UV-B, then exposed to 7 h of broadband UV-B stress (UV stress 7 h) or not (Control). Pictures were taken after recovery for another 7 days in white light. Bars = 1 cm.

(**C**) Absorbance spectra (280–400 nm) of methanolic extracts from 9-day-old Tak-1, Mp*hy5-1*, Mp*hy5-3*, and Mp*hy5-7* gemmalings grown for 7 days in white light and exposed (WL+UV-B, purple lines) or not (WL, blue lines) to supplemental UV-B (1.5 µmol m⁻² s⁻¹) for 2 days. The gray lines show absorbance of the buffer (blank). Values of independent measurements and means ± SEM are shown (*n* = 3).

(**D**) Hierarchical clustering and heatmap of UV-B-regulated gene expression in Tak-1 and Mp*hy5-1* plants grown for 7 days and exposed, or not, to supplemental UV-B for 3h. All genes with a greater than two-fold change in expression under UV-B and a p-adj < 0.05 in at least one genotype are shown (total = 501 genes). Red and blue colors indicate normalized read count Z-scores. The color bar on the left side indicates 10 major profiles (clusters) of gene expression. The color bars on the right indicate if genes are significantly misregulated in white light in Mp*cop1* compared to Tak-1, at varying fold-change thresholds.

Figure 9. Auronidins and related flavonoid pigments may play an important role in protecting Marchantia from UV-B stress.

Figure 9. Auronidins and related flavonoid pigments may play an important role in protecting Marchantia from UV-B stress.

(**A**) Left: Schematic of the phenylpropanoid pathway leading to flavonoid pigments and the occurrence of biosynthetic enzymes in different land plant lineages (according to Davies et al., 2022). A colored circle next to an enzyme step indicates that the activity is present in the corresponding plant lineage, as indicated by the color key. An empty white circle indicates its absence in that lineage. A black circle indicates that there is some support for the presence of the enzyme, provided from either sequence analysis or presence of the biosynthetic product, but that definitive evidence is lacking. Right: Heat map showing gene expression levels as determined by RNA-seq of phenylpropanoid biosynthetic enzymes in Tak-1, Mp*uvr8-1*, Mp*rup-2*, Mp*cop1-1*, and Mp*spa-10* plants grown for 7 days in white light exposed (+) or not (-) to supplemental UV-B for 3 h. The $\log_2$ fold change of normalized read counts relative to Tak-1 without UV-B exposure (/Tak-1-UV $\log_2$FC) was calculated. Enzyme abbreviations: PAL, PHENYLALANINE AMMONIA LYASE; C4H, CINNAMATE 4-HYDROXYLASE; 4CL, COUMAROYL CoA LIGASE; CHS, CHALCONE SYNTHASE; CHI, CHALCONE ISOMERASE; CHIL, CHALCONE ISOMERASE-LIKE; AUS, AUREUSIDIN SYNTHASE; FGT, FLAVONOID O-GLYCOSYLTRANSFERASES; FNS, FLAVONE SYNTHASE.

(**B**) RT-qPCR analysis of Mp*AUS* in 7-day-old Tak-1, Mp*uvr8-1*, Mp*hy5-1*, and Mp*hy5-3* plants grown for 7 days in white light and exposed to supplemental UV-B for 3 h (WL+UV-B) or not (WL). Individual data points of biological replicates and means ± SD are shown (*n* = 3). Letters indicate statistically significant differences between the means (P < 0.01) as determined by two-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

(**C**) F_v/F_m measurements in Tak-1, Mp*rup-2*, Mp*aus-2*, and Mp*rup-2* Mp*aus-2* gemmalings grown for 7 days in white light (non-acclimated) or white light supplemented with UV-B (acclimated), without (not stressed) or with (stressed) additional exposure to broadband UV-B stress for 60 min. -/-, not acclimated, not stressed; -/+, not acclimated, stressed; +/-, acclimated, not stressed; +/+, acclimated, stressed. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Letters indicate statistically significant differences between the means (P < 0.01) as determined by two-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

(**D**) UV-B stress survival of Tak-1, Mp*rup-2*, Mp*aus-2*, and Mp*rup-2* Mp*aus-2* plants grown for 7 days in white light supplemented (Accl.) or not (Non-accl.) with UV-B, then exposed to 7 h of broadband UV-B (UV stress 7 h) or not (Control). Pictures were taken after recovery for another 7 days in white light. Bars = 1 cm.

(**E**) Growth phenotype of 4-week-old wild-type (Tak-1), Mp*rup-2*, Mp*aus-2*, and Mp*rup-2* Mp*aus-2* plants grown in white light supplemented with UV-B (WL +UV) or not (WL). Bar = 1 cm.

(**F**) Absorbance spectra (280–400 nm) of methanolic extracts from 9-day-old Tak-1, Mp*rup-2*, Mp*aus-2*, and Mp*rup-2* Mp*aus-2* gemmalings grown for 7 days in white light and exposed (WL+UV-B, purple lines) or not (WL, blue lines) to supplemental UV-B for 2 days. The gray lines show absorbance of the buffer (blank). Values of independent measurements and means ± SEM are shown (*n* = 3).

(**G**) and (**H**) Metabolomic analysis of 9-day-old plants of Tak-1, Mp*uvr8-1*, Mp*rup-2*, and Mp*aus-2* grown for 7 days in white light and exposed (+) or not (-) to supplemental UV-B (1.5 µmol m^{-2} s^{-1}) for 2 days (*n* = 5, 12 plants per replicate). (G) Venn diagram showing overlaps between upregulated metabolites (FC> or =1.5, p-adj < or =0.05) upon exposure to 2 days UV-B in 9-day-old wild-type (Tak-1), Mp*uvr8-1* (Mp*uvr8*), Mp*rup-2* (Mp*rup*), and Mp*aus-2* (Mp*aus)* gemmalings. (H) Distribution of the number of compounds induced under UV-B in wild type (FC> or =1.5, p-adj < or =0.05), according to their chemical classes. Corresponding metabolite lists: Supplementary Table S9.

```
AtUVR8     1 MAEDMAADEVTAPPRKVLIISAGASHSVALLSGDIVCSWGRGEDGQLGHGDAEDRPSPTQ   60
MpUVR8.1   1 -MEDSKMSSNERPRRRVLFVSAGASHSVALLTGDSVCSWGRGEDGQLGHGDADERHLPTI   59
MpUVR8.2   1 -MEDSKMSSNERPRRRVLFVSAGASHSVALLTGDSVCSWGRGEDGQLGHGDADERHLPTI   59
                                                      *

AtUVR8    61 LSALDGHQIVSVTCGADHTVAYSQSGMEVYSWGWGDFGRLGHGNSSDLFTPLPIKALHGI  120
MpUVR8.1  60 ISTLGECEISAITCGADHTTAYADSTRTVYSWGWGDFGRLGHGNSSDLFIPHPIKALRGI  119
MpUVR8.2  60 ISTLGECEISAITCGADHTTAYADSTRTVYSWGWGDFGRLGHGNSSDLFIPHPIKALRGI  119
                                           *  * * *              *

AtUVR8   121 RIKQIACGDSHCLAVTMEGEVQSWGRNQNGQLGLGDTEDSLVPQKIQAFEGIRIKMVAAG  180
MpUVR8.1 120 EISQIACGDSHCLAITTQGDVHSWGRNQNGQLGLGHNEDSLVPKKVVAFEGIAVKMLAAG  179
MpUVR8.2 120 EISQIACGDSHCLAITTQGDVHSWGRNQNGQLGLGHNEDSLVPKKVVAFEGIAVKMLAAG  179
                              *

AtUVR8   181 AEHTAAVTEDGDLYGWGWGRYGNLGLGDRTDRLVPERVTSTGGEKMSMVACGWRHTISVS  240
MpUVR8.1 180 AEHTAAVTESGKLYGWGWGRYGNLGLGDRNDRLVPAEVAGVNNEKFTLVACGWRHTIAVA  239
MpUVR8.2 180 AEHTAAVTESGKLYGWGWGRYGNLGLGDRNDRLVPAEVAGVNNEKFTLVACGWRHTIAVA  239
                            *  *                                    *

AtUVR8   241 YSGALYTYGWSKYGQLGHGDLEDHLIPHKLEALSNSFISQISGGWRHTMALTSDGKLYGW  300
MpUVR8.1 240 ESGNLYTFGWSKYGQLGHGDYLDHLVPHQVAALKNNPIVQISGGWRHTVALDSSGRLYGW  299
MpUVR8.2 240 ESGNLYTFGWSKYGQLGHGDYLDHLVPHQVAALKNNPIVQISGGWRHTVALDSSGRLYGW  299
                        *                            *                 *

AtUVR8   301 GWNKFGQVGVGNNLDQCSPVQVRFPDDQKVVQVSCGWRHTLAVTERNNVFAWGRGTNGQL  360
MpUVR8.1 300 GWNKFGQVGVGNTDDQTSPQLVKNLAEEKVKQVSCGWRHTVCITERGNVFSWGRGTSGQL  359
MpUVR8.2 300 GWNKFGQVGVGNTDDQTSPQLVKNLAEEKVKQVSCGWRHTVCITERGNVFSWGRGTSGQL  359
                    *                              *             *

AtUVR8   361 GIGESVDRNFPKIIEALSVDGASGQHIESSNIDPSSGKSWVSPAERYAVVPDET------  414
MpUVR8.1 360 GHGDAVDRSLPKRIEALSADGLACQELESSRTTTSTGSHWISPSERYAVVPVETVRSLSR  419
MpUVR8.2 360 GHGDAVDRSLPKRIEALSADGLACQELESSRTTTSTGSHWISPSERYAVVPVET------  413
                                                            *

AtUVR8   415 ----GLTDGSSKGNGGDISVPQTDVKRVRI--                              440
MpUVR8.1 420 NMHQNANVGDNEVPSSDASVPDTDVKRLRTGF                             451
MpUVR8.2 414 -------VGDNEVPSSDASVPDTDVKRLRTGF                             438
```

Supplementary Figure S1. Alignment of Marchantia MpUVR8.1 and MpUVR8.2 with Arabidopsis AtUVR8.
Red asterisks (*) highlight conserved tryptophan (W) amino acid residues, including the AtUVR8 key triad tryptophans W233 (W232 in MpUVR8), W285 (W284 in MpUVR8), and W337 (W336 in MpUVR8), within their conserved GWRHT sequence motif. Green asterisks (*) highlight AtUVR8 aspartates D96 (D95 in MpUVR8) and D107 (D106 in MpUVR8).

Supplementary Figure S2. Volcano plot of GFP-MpUVR8 AP-MS under +UV versus -UV conditions.
AP–MS was performed using a Marchantia Pro_{MpUVR8}:GFP-Mp$UVR8$ (genomic) transgenic line grown for 7 days under continuous white light (50 µmol m⁻² s⁻¹) without UV-B, or under continuous white light supplemented with narrowband UV-B (1.5 µmol m⁻² s⁻¹) followed by a 20 min broadband UV-B exposure (25 µmol m⁻² s⁻²). Significantly different proteins were identified using a p-value cutoff of < 0.01 and a log10 fold change threshold of ±1.0. Grey dots represent non-significant proteins; red dots indicate proteins significantly enriched in +UV compared to –UV; blue dots indicate proteins significantly enriched in –UV compared to +UV. Arrows highlight proteins of interest.

Supplementary Figure S3. Gene structure and indication of the mutated sites in the corresponding mutants generated by CRISPR/Cas9.

Supplementary Figure S3. Gene structure and indication of the mutated sites in the corresponding mutants generated by CRISPR/Cas9.

Untranslated (5' and 3' UTRs) and coding sequences are highlighted as white and colored boxes, respectively, with specific domains indicated. Introns are represented by lines connecting exons. The positions of the guide RNA (gRNA) (red bar, sequence in blue), protospacer adjacent motif (PAM) sequences (sequence in red), different mutations including resulting amino acid changes (in yellow), translated amino acids of wild-type (in purple) and the number of translated amino acids are also highlighted. The Mp*rup-2* mutant exhibits an insertion of one adenine (A) nucleotide, while Mp*rup-3* displays 27-bp insertion with a 15-bp deletion of endogenous sequence. Mp*cop1-1* and Mp*cop1-2* mutants show an 11-bp deletion and a 2-bp deletion with one cytosine (C) nucleotide substituted by thymine (T), respectively. Mp*spa-9* has an insertion of one guanine (G) nucleotide for gRNA1, whereas Mp*spa-10* exhibits an insertion of one cytosine (C) nucleotide for gRNA2. The Mp*hy5-1* mutant displays a deletion of one adenine (A) nucleotide, whereas Mp*hy5-3* and Mp*hy5-7* mutants show a 5-bp deletion and 11-bp deletion, respectively. Mp*aus-2* shows a 2-bp deletion and Mp*aus-3* shows an insertion of one thymine (T) nucleotide. All these mutations, except for Mp*rup-3*, result in a frameshift and early STOP codon in the coding sequence.

Supplementary Figure S4. UV-B photomorphogenesis and photoprotection in MpUVR8 and MpRUP transgenic lines.

Supplementary Figure S4. UV-B photomorphogenesis and photoprotection in MpUVR8 and MpRUP transgenic lines.

(**A**) Growth phenotype of 4-week-old wild-type (Tak-1), Mp*rup-2*, Mp*uvr8-1*, Mp*uvr8-1/Pro*$_{MpUVR8}$:GFP-Mp*UVR8* line #5, Mp*RUP*^{OE} #1, Mp*rup-2*/Mp*RUP*^{OE} #4, Mp*UVR8.1*^{OE} #1, and Mp*UVR8.2*^{OE} #1 plants grown in white light supplemented (WL +UV) or not (WL -UV) with UV-B. Bar = 1 cm.

(**B**) F_v/F_m measurements in wild-type (Tak-1), Mp*uvr8-1*, Mp*UVR8.1*^{OE} lines #1 and #2, Mp*UVR8.2*^{OE} #1 and #2, Mp*uvr8-1/Pro*$_{MpUVR8}$:GFP-Mp*UVR8* #4 and #5, Mp*rup-2*, Mp*RUP*^{OE} #1 and #3, and Mp*rup-2*/Mp*RUP*^{OE} #4 grown for 7 days in white light (non-acclimated) or white light supplemented with UV-B (acclimated), without (non-stressed) or with (stressed) additional exposure to broadband UV-B stress for 60 min. -/-, non-acclimated, non-stressed; -/+, non-acclimated, stressed; +/-, acclimated, non-stressed; +/+, acclimated, stressed. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Dashed horizontal lines indicate levels corresponding to acclimated and stressed Tak-1 (blue), Mp*uvr8-1* (green), and Mp*rup-2* (orange) plants. Different letters indicate statistically significant differences between the means (P < 0.01) as determined by two-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

(**C,D**) UV-B stress tolerance of indicated plant genotypes grown for 7 days in white light supplemented (acclimated) or not (non-acclimated) with UV-B, then exposed or not to 6 hours of broadband UV-B. Pictures were taken after recovery for another 7 days in white light. Bar = 1 cm. (C) #1, #2 and #3: three independent Mp*uvr8-1/Pro*$_{MpUVR8}$:Mp*UVR8* lines.

(**E**) Immunoblot analysis of MpUVR8 and RFP-MpUVR8 levels in 14-day-old Tak-1, Mp*UVR8.1*^{OE}#1, Mp*UVR8.1*^{OE}#2, Mp*UVR8.2*^{OE}#1, Mp*UVR8.2*^{OE}#2, and Mp*uvr8-1* thallus. Ponceau staining is shown as loading control. Mp*UVR8.1*^{OE} = Tak-1/*Pro*$_{MpEF1α}$:RFP-UVR8.1; Mp*UVR8.2*^{OE} = Tak-1/*Pro*$_{MpEF1α}$:RFP-UVR8.2.

(**F**) RT-qPCR analysis of Mp*RUP* expression level in 14-day-old Tak-1 and two Mp*RUP*^{OE} lines. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Different letters indicate statistically significant differences between the means (P < 0.05) as determined by one-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

Supplementary Figure S5. Enhanced photoprotection and UV-B-absorbing capacity of Mp*rup* depends on MpUVR8.

(**A**) F_v/F_m measurements in wild-type (Tak-1), Mp*uvr8-1*, Mp*rup-2*, and Mp*uvr8-1* Mp*rup-2* grown for 7 days in white light (non-acclimated) or white light supplemented with UV-B (acclimated), without (non-stressed) or with (stressed) additional exposure to broadband UV-B stress for 60 min. -/-, non-acclimated, non-stressed; -/+, non-acclimated, stressed; +/-, acclimated, non-stressed; +/+, acclimated, stressed. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Different letters indicate statistically significant differences between the means (P < 0.01) as determined by two-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

(**B**) UV-B stress resistance in Tak-1, Mp*uvr8-1*, Mp*rup-2*, and Mp*uvr8-1* Mp*rup-2* plants grown for 7 days in white light supplemented (acclimated) or not (non-acclimated) with UV-B, then exposed or not to 7 hours of broadband UV-B. Pictures were taken after recovery for another 7 days in white light. Bars = 1 cm.

(**C-F**) Absorbance spectra (280–400 nm) of methanolic extracts from 9-day-old Tak-1 (A), Mp*uvr8-1* (B), Mp*rup-2* (C), and Mp*uvr8-1* Mp*rup-2* (D) gemmalings grown for 7 days in white light and exposed (WL+UV-B, purple lines) or not (WL, blue lines) to supplemental UV-B (1.5 µmol m^{-2} s^{-1}) for 2 days. The gray line shows the absorbance of the buffer (blank). Values of independent measurements and means ± SEM are shown (*n* = 3).

Supplementary Figure S6. MpSPA does not affect MpUVR8 redimerization, but MpCOP1 and MpSPA overexpression reduces basal UV-B tolerance.

(**A**) Immunoblot analysis of MpUVR8 levels in 7-day-old Mp*spa-10* gemmalings irradiated for 20 min with broadband UV-B before recovery in white light (WL) for the indicated times. Top: MpUVR8 dimer levels in DSP-crosslinked protein extracts. Bottom: total MpUVR8 levels in non-crosslinked extracts.

(**B**) RT-qPCR analysis of Mp*SPA* expression level in 14-day-old plants of Tak-1 and Mp*SPA*^*OE*^ lines *#1* and *#4*. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Letters indicate statistically significant differences between the means (P < 0.05) as determined by one-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

(**C**) RT-qPCR analysis of Mp*COP1* expression level in 14-day-old plants of Tak-1, Mp*COP1*^*OE*^ lines *#4* and *#6*. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Letters indicate statistically significant differences between the means (P < 0.05) as determined by one-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

(**D**) F_v/F_m measurements in wild-type (Tak-1), Mp*RUP*^*OE*^ lines *#1* and *#3*, Mp*SPA*^*OE*^ *#1* and *#4*, and Mp*COP1*^*OE*^ *#4* and *#6* grown for 7 days in white light (non-acclimated) or white light supplemented with UV-B (acclimated), without (not stressed) or with (stressed) additional exposure to broadband UV-B stress for 60 min. -/-, not acclimated, not stressed; -/+, not acclimated, stressed; +/-, acclimated, not stressed; +/+, acclimated, stressed. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Letters indicate statistically significant differences between the means (P < 0.01) as determined by two-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

Supplementary Figure S7. MpHY5 plays a minor role in the regulation of UV-B-induced gene expression changes.
Gene expression profiles in clusters identified by hierarchical clustering. Z-scores for normalized read count values for each gene (gray lines) and for the cluster average (colored line) are shown.

Supplementary Figure S8. MpHY5 and MpCOP1 regulate UV-B-induced gene expression in an antagonist manner.
(**A**) Euler plots showing overlaps between upregulated DEGs (p-adj < 0.05) in Mp*cop1*/Tak-1 in white light (left: log$_2$FC > 1; right: log$_2$FC > 2) and genes that are induced after 3h of UV-B. UV-B-upregulated genes are separated into 3 categories: genes that are induced in a MpHY5-independent manner ("HY5-indep.", Fig. S6 cluster 1), genes that are dependent on MpHY5 for their full induction or hypoinduced in Mp*hy5* ("HY5-dep.", Fig. S6 clusters 2 and 7) and genes that are hyperinduced in Mp*hy5* or whose induction is repressed by HY5 ("HY5-repr.", Fig. S6 clusters 3 and 9). Corresponding gene lists: Supplementary Table S6a.
(**B**) Euler plots showing overlaps between downregulated DEGs (p-adj < 0.05) in Mp*cop1*/Tak-1 in white light (left: log$_2$FC < -1; right: log$_2$FC < -2) and genes that are repressed after 3h of UV-B. UV-B-downregulated genes are separated into 3 categories: genes that are repressed in a MpHY5-independent manner ("HY5-indep.", Fig. S6 clusters 4, 8 and 10), genes that are somewhat constitutively repressed in Mp*hy5* ("hy5-repr.", Fig. S6 cluster 6) and genes that are derepressed in white light-grown Mp*hy5* ("hy5-derepr.", Fig. S6 cluster 5). Corresponding gene lists: Supplementary Table S6b.
(**C**) Heatmap of all genes upregulated (log$_2$FC > 1 and p-adj < 0.05) by UV-B in Tak-1 and/or Mp*hy5*. Red and blue colors indicate normalized read count Z-scores in RNA-seq experiments including Tak-1 and Mp*hy5* (left) and Tak-1, Mp*cop1* and Mp*spa* (right). Z-scores were calculated separately for the two experiments and genes were ordered in a decreasing order according to the Z-score in Mp*hy5*+UV (arrow).
(**D**) Heatmap of all genes downregulated (log$_2$FC < -1 and p-adj < 0.05) by UV-B in Tak-1 and/or Mp*hy5*. Red and blue colors indicate normalized read count Z-scores in RNA-seq experiments including Tak-1 and Mp*hy5* (left) and Tak-1, Mp*cop1* and Mp*spa* (right). Z-scores were calculated separately for the two experiments and genes were ordered in a decreasing order according to the Z-score in Mp*hy5*+UV (arrow).

Supplementary Figure S9. LUC activity of a Pro_{MpCHS2}:LUC reporter in Tak-1, Mp*uvr8-1*, and Mp*rup-2* backgrounds upon UV-B exposure.

(**A**) The counts per second (cps) of luciferase (LUC) activity were normalized to average expression level measured in 3-day-old gemmalings of the following lines: Tak-1/Pro_{MpCHS2}:LUC, Mp*uvr8-1*/Pro_{MpCHS2}:LUC, and Mp*rup-2*/Pro_{MpCHS2}:LUC. These gemmalings were grown in 96-well plates under white light, and pre-sprayed with luciferin 24 h before a 20-min exposure to broadband UV-B. The LUC activity was monitored for up to 360 min after the UV-B induction. Values of independent measurements and means ± SD are shown (n = 8).

(**B**) Analysis of LUC activity in 14-day-old thalli of Tak-1/Pro_{MpCHS2}:LUC, Mp*uvr8-1*/Pro_{MpCHS2}:LUC, and Mp*rup-2*/Pro_{MpCHS2}:LUC lines followed by 20 min of broadband UV-B exposure at 4 h (left) and 8 h (right).

Supplementary Figure S10. MpAUS contributes to protection against UV-B stress.

(**A**) F_v/F_m measurements in wild-type (Tak-1), Mp*aus-2,* Mp*spa-10*, Mp*spa-10* Mp*aus-2*, and Mp*uvr8-1* gemmalings grown for 7 days in white light (non-acclimated) or white light supplemented with UV-B (acclimated), without (not stressed) or with (stressed) additional exposure to broadband UV-B stress for 60 min. -/-, not acclimated, not stressed; -/+, not acclimated, stressed; +/-, acclimated, not stressed; +/+, acclimated, stressed. Individual data points of biological replicates and means ± SD are shown (*n* = 3). Dashed horizontal lines indicate levels corresponding to Tak-1 not acclimated stressed (green) and acclimated stressed (blue) plants. Letters indicate statistically significant differences between the means (P < 0.01) as determined by two-way ANOVA followed by Tukey's post hoc honestly significant difference (HSD) test.

(**B**) UV-B stress survival of Tak-1, Mp*aus-2*, Mp*spa-10*, Mp*spa-10* Mp*aus-2*, and Mp*uvr8-1* plants grown for 7 days in white light supplemented (acclimated) or not (non-acclimated) with UV-B, then exposed to 7 h or 9 h of broadband UV-B or not. Pictures were taken after recovery for another 7 days in white light. Bars = 1 cm.

(**C**) Absorbance spectra (280–400 nm) of methanolic extracts from 9-day-old Tak-1, Mp*spa-10*, Mp*aus-2,* Mp*spa-10* Mp*aus-2*, and Mp*uvr8-1* gemmalings grown for 7 days in white light and exposed (WL+UV-B, purple lines) or not (WL, blue lines) to supplemental UV-B for 2 days. The gray lines show absorbance of the buffer (blank). Values of independent measurements and means ± SEM are shown (*n* = 3).

A Tak-1   Mp*uvr8-1*   Mp*rup-2*
    −   +   −   +   −   +   UV-B

DPBA staining

B Tak-1 −UV

C

feature 266, 3.693_445.0776
Apigenin-7-glucoside

feature300_3.81_461.0725
Luteolin 7 O gluc

feature64_5.358_257.0816
Lunularic acid

feature228_6.823_423.1601
Isoriccardin C

feature261_6.952_439.1551
MarchantinA

feature431_2.52_591.1354
auronidin-4-neo

Supplementary Figure S11. MpUVR8- and MpAUS-dependent regulation of metabolite accumulation upon UV-B exposure.
(**A**) High-performance thin-layer chromatography (HPTLC) analysis of flavonol glycoside levels in 7-d-old gemmalings of wild type (Tak-1), Mp*uvr8-1*, and Mp*rup-2* grown for 7 days in white light and exposed, or not, to supplemental UV-B for 2 days. DPBA, diphenylboric acid 2-aminoethylester.
(**B-C**) Metabolomic analysis of 9-day-old plants of Tak-1, Mp*uvr8-1*, Mp*rup-2*, and Mp*aus-2* grown for 7 days in white light and exposed (+) or not (−) to supplemental UV-B (1.5 μmol m^{-2} s^{-1}) for 2 days (n = 5, 12 plants per replicate). (B) UHPLC-MS chromatogram from -UV-B Tak-1 sample. Metabolites corresponding to major peaks are annotated based on their accurate mass and MS2 spectra. (C) Relative quantifications of compounds in Tak-1, Mp*uvr8-1* (*uvr8*), Mp*rup-2* (*rup*), and Mp*aus-2* (*aus*) methanolic extracts from plants exposed to UV-B (purple, right bar, UV) or not (yellow, left bar, no UV). Individual data points of biological replicates and means ± SD are shown (n = 4).