Deficiency in POLE exonuclease causes synthetic lethality in

highly aneuploid cancer cells

Eun Jung Kim[1], Alejandro Chibly[2], Kristel Dorighi[3], Jack Kamm[2], Szymon Juszkiewicz[3], Kevin Cabrera[4],

Dexter Jin[4], Rohit Reja[2], Yangmeng Wang[1], Rachel Nakagawa[1], Minyi Shi[3], Yuxin Liang[3], Anwesha Dey[3],

Ethan S. Sokol[4,*], Steven Gendreau[1,*], Meng He[1,5,*]

[1]Translational Medicine, Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA

[2]Computational Sciences, Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA

[3]Research Biology, Genentech, Inc., 1 DNA Way, South San Francisco, CA 94080, USA

[4]Foundation Medicine, Inc., 400 Summer Street, Boston, MA 02210, USA

[*]Correspondence: hem8@gene.com, esokol@foundationmedicine.com and gendreas@gene.com

[5]Lead contact

12 Summary

Aneuploidy is a hallmark of cancer and is associated with drug resistance and poor clinical

outcomes across diverse cancer types. However, no therapies have been clinically established

to target highly aneuploid tumors. By analyzing a real-world dataset comprising nearly half a

million tumor samples subjected to comprehensive genomic profiling, we identified a striking

mutual exclusivity between POLE exonuclease domain mutations and high aneuploidy burden.

This observation was independently validated using data from The Cancer Genome Atlas

(TCGA) and the Cancer Cell Line Encyclopedia (CCLE). Probabilistic modeling revealed that

the elevated quantity and unique spectrum of mutations induced by POLE exonuclease

deficiency increase the likelihood of inactivating essential genes on chromosome arms

harboring losses, leading to a synthetic lethal phenotype in highly aneuploid cells. Functional

experiments demonstrated that POLE exonuclease activity is essential for the viability of highly

aneuploid cancer cell lines but is dispensable in diploid cells. These findings suggest that

selective inhibition of POLE exonuclease may represent a promising therapeutic strategy for

targeting highly aneuploid tumors.

27 Keywords

28 aneuploidy, exonuclease, POLE, synthetic lethality, essential genes, chromosome, tumor

29 mutational burden, adenine base editor, microsatellite instability

Introduction

Aneuploidy is a cellular state characterized by the acquisition of an abnormal number of chromosomes or chromosome arms [1]. These chromosomal abnormalities are observed in over 90% of solid tumors and approximately 50% of hematological malignancies [2-4]. Aneuploidy has been recognized as both a consequence and a driver of tumorigenesis [5-7]. While aneuploidy often reduces fitness in normal cells, cancer cells tolerate and even exploit chromosomal imbalances for increased fitness [8-10]. Though the molecular mechanisms underpinning the initiation of aneuploidy is largely unknown, recent advances in single-cell genomics, karyotype-resolved sequencing, and functional genetics have illuminated the paradoxical role of aneuploidy in tumorigenesis [11]. Aneuploid genomes in cancer cells offer a substrate for clonal diversification under selective pressures, enabling tumor cells to adapt to microenvironmental stressors, evade immune detection, and resist therapeutic interventions [12,13]. Additionally, while chromosomal alterations can amplify oncogenes or eliminate tumor suppressors, the effects of chromosomal abnormalities can extend to global transcriptomic and proteomic disruption [14]. Previous work has implicated aneuploidy in the induction of proteotoxic and metabolic stress, disruption of mitotic fidelity, and activation of tumor-suppressive checkpoints, such as p53 [15-17].

Clinically, high aneuploidy burden, as well as specific recurrent aneuploidies—such as gain of chromosome 1q or 8q, or loss of 17p or 8p—are associated with poor prognosis, therapeutic resistance, and aggressive disease behavior across numerous cancer types, including breast, colorectal, ovarian, and lung cancers [10,18,19]. Mechanistically, aneuploidy promotes resistance through multiple routes, including increased intratumoral heterogeneity, reprogramming of apoptotic and stress response pathways, altered drug metabolism, and modulation of the tumor immune microenvironment [20]. For instance, tumors with high levels of aneuploidy often exhibit reduced antigen presentation and increased immunosuppressive signaling, features that correlate with diminished response to immunotherapies [21].

Therapeutic strategies targeting aneuploidy and consequential stresses have gained increased attention. Early preclinical efforts focused on amplifying the intrinsic vulnerabilities of aneuploid cells, such as proteotoxic and mitotic stress, through the use of proteasome inhibitors, heat shock protein (HSP) inhibitors, or spindle assembly checkpoint modulators [22-24]. More recently, synthetic lethality screens have identified aneuploidy-specific dependencies, including elevated reliance on autophagy, DNA repair pathways, and the unfolded protein response

[25,26]. For example, cells with whole-chromosome gains are more sensitive to centrosome

clustering inhibitors or agents targeting oxidative phosphorylation and energy homeostasis [27].

Despite promising preclinical results, clinical translation remains challenging. Many stress

pathway inhibitors lack tumor specificity or induce systemic toxicity [28]. Nonetheless, early-

phase trials have evaluated compounds such as the HSP90 inhibitor ganetespib and

proteasome inhibitor bortezomib, with mixed results [22,23,29].

In this study, we analyzed a real-world database comprising half a million patient samples

across over 400 cancer types and subtypes to identify potential synthetic lethal targets for high

aneuploidy burden. We identified and validated the exonuclease domain of POLE as a potential

therapeutic target for tumors exhibiting high aneuploidy burden.

Results

Characterization of chromosomal arm-level aneuploidy in a real-world comprehensive genomic profiling database

We examined a real-world database of comprehensive genomic profiling results derived from

499,320 patient samples. This database encompasses over 90 different primary cancer types,

including non-small cell lung cancer, colorectal cancer, breast cancer, pancreatic cancer, ovary

cancer, prostate cancer, etc. (Figure 1A). Patient and tumor characteristics, including

chromosomal sex, histological cancer type, local/metastasis biopsy site and genetic ancestry,

are well represented in this database (Figure 1B).

A computational algorithm was run to call chromosomal arm level gains and losses for 40

autosomal chromosomal arms for each sample [30,31]. The arm-level aneuploidy caller

revealed unique chromosome alterations across different histological cancer types (Figure 1C-

1F). Consistent with previous reports, 8p loss, 17p loss, 1q gain and 8q gain are particularly

prevalent in carcinomas, as compared to sarcomas, gliomas and leukemias [3,18,19], whereas

whole chromosome 7 gain and whole chromosome 10 loss were observed as unique features in

gliomas [32]. On the other hand, leukemia exhibited significantly less arm-level aneuploidy [33].

Upon examining the arm-level aneuploidy across different primary tumor types (Supplemental

Figure 1), we also observed distinct distributions of specific chromosomal alterations. For

instance, while 1q gain is common across various carcinoma cancer types, it is particularly

common in breast cancer with a near 70% incidence rate. 3p loss is particularly enriched in

small cell, head and neck, cholangiocarcinoma and kidney cancers with an incidence rate over
50%. Another notable example is 13q gain which is detected in approximately 70% of colorectal
cancer but is rare in most other cancer types.

To estimate the aneuploidy burden in each tumor type, we calculated an aneuploidy score by
counting the total number of chromosome arm losses or gains in a tumor sample [3,34,35]. Not
surprisingly, we observed pronounced differences in aneuploidy scores across tumor types
(Figure 1G). Hematological malignancies such as leukemia, non-Hodgkin lymphoma, and
diffuse large B-cell lymphoma exhibit significantly less aneuploidy. Even within solid tumor
types, gliomas, mesotheliomas, and prostate cancers tend to have significantly lower
aneuploidy scores compared to breast cancer, non-small cell lung cancer and small cell cancer.

Additionally, we observed statistically significant differences in aneuploidy scores among
patients with different genetic ancestries (Supplemental Figure 2A). Patients of Admixed
American ancestry generally display lower aneuploidy compared to patients of African ancestry.
Higher aneuploidy burden has been associated with metastasis [36,37], which was consistent
with our observation of significantly higher aneuploidy scores in tumor samples collected from
metastatic sites (Supplemental Figure 2B). A strong correlation between aneuploidy scores and
patient age was also noted (Supplemental Figure 2C). Chromosomal arm losses and gains
contribute equally to aneuploidy scores, and no correlation was observed between these two
types of alterations (Supplemental Figure 2D), suggesting these events are likely selected
independently during cancer evolution.

Elastic net regression model identifies genetic mutation of POLE as a top
negative predictor of aneuploidy score

To systematically evaluate the impact of patient and tumor characteristics, including sex, tumor
subtype, genetic ancestry, metastasis status, and genomic alterations on aneuploidy, we
deployed an elastic net regression model using aneuploidy score as the dependent variable
(see Method for details). The comprehensive results from the elastic net regression model can
be found in Supplemental Table S1, with an optimal alpha value of 0.25 (Supplemental Figure
3A). Genomic alterations with non-zero coefficients are summarized in Figure 2A. The model
identified *TP53* alteration as the top positive predictor of aneuploidy score, a well-established
driver of cancer genomic instability and aneuploidy [4,6,38]. In non-transformed primary cells,
TP53 inactivation is often required for cells to tolerate aneuploidy [8,39,40]. Thus, *TP53*

alteration serves as a positive control, demonstrating the robustness of our model. Further

examination showed that tumors with homozygous loss (CN) of *TP53* exhibit significantly higher

aneuploidy scores than those with *TP53* short variants (SV) or structural rearrangements (RE)

(Figure 2B and Supplemental Figure 3B). Apart from *TP53*, our model identified several other

genomic alterations as top positive predictors for aneuploidy scores, such as *AR, GID4, AKT3,*

and FGF10. Although the functional link between these genomic alterations and aneuploidy

remains unclear, it is interesting to note that alterations of *GID4, AKT3,* and *FGF10* are

predominantly through amplification (Supplemental Figure 3C), and the association between *AR*

and aneuploidy score is driven by *AR* amplification in prostate cancer (Supplemental Figure

3D).

The model also identified several negative predictors for aneuploidy scores, such as *GNAQ,*

BCL2A1, CD74, FOXL2, EWSR1, and notably *POLE*. Although their functional links with

aneuploidy are unclear, the strong signal that alterations in *POLE* is associated with lower

aneuploidy scores warranted further investigation. *POLE* encodes the catalytic subunit of the

DNA polymerase epsilon complex and is most frequently mutated in endometrial (1.5%) and

colorectal cancers (0.5%) (Figure 2C). *POLE* encodes a protein with multiple functional

domains, including the exonuclease domain, the DNA polymerase domain, and the less

characterized DUF1744 domain [41,42] (Figure 2D). Hotspot mutations in the *POLE*

exonuclease domain have been reported [41,43] and separating tumors with mutations within

this domain versus outside of the domain revealed that the negative association with high

aneuploidy burden is driven predominantly by mutations within the exonuclease domain (Figure

2E and Supplemental Figure 3E). This association was validated in The Cancer Genome Atlas

(TCGA) (Figure 2F) and the Cancer Cell Line Encyclopedia (CCLE) (Figure 2G). Across all

three datasets, tumors with exonuclease domain mutations of *POLE* are near diploid compared

to those with mutations outside the exonuclease domain or *POLE* wild-type tumors.

We also looked to see if there was an association between aneuploidy score and *POLD1*

alterations. *POLD1* encodes the catalytic subunit of DNA polymerase delta. Both *POLE* and

POLD1 are essential B-family DNA polymerases in eukaryotic cells, playing crucial roles in DNA

replication and repair [44-47]. Both enzymes possess 5′ to 3′ polymerase and 3′ to 5′

exonuclease proofreading activities, ensuring high-fidelity DNA synthesis. However, they differ

in their primary functions during replication: POLE is primarily responsible for leading strand

synthesis, while POLD1 mainly synthesizes the lagging strand [44]. Mutations in the

exonuclease domains of POLE and POLD1 can compromise their proofreading abilities, leading

to increased mutation rates [41,42]. Interestingly, our elastic net regression model identified

POLE alteration as a top negative predictor to aneuploidy scores, but not *POLD1*. We therefore

examined the association between aneuploidy scores and *POLD1* mutation status. While there

was a statistically significant reduction in aneuploidy score in *POLD1* exonuclease (a.a.304-

a.a.533) mutated tumors, the effect size was minimal and less consistent across cancer types

compared to *POLE* (Figure 2H and Supplemental Figure 3F).

Deficiency in POLE exonuclease causes synthetic lethality in highly-aneuploid

cells by inactivating essential genes on chromosome arms with losses

We were interested in understanding the underlying mechanism that drives the mutual

exclusivity between POLE exonuclease mutations and high aneuploidy burden in tumor cells.

Mutual exclusivity in biology has been proposed to stem from either functional redundancy or

synthetic lethality [48]. POLE exonuclease mutations are often associated with ultra mutated

tumors, characterized by extremely high tumor mutation burdens (TMB), in many cases

exceeding 500,000 mutations per haploid genome (Figure 3A and Supplemental Figure 4A)

[45]. Although there is no clear association between TMB and aneuploidy scores when

examined across all tumors (Supplemental Figure 4C) [3], we observed a strong negative

correlation between TMB and aneuploidy scores in *POLE*-mutated tumors (Figure 3B). POLE

exonuclease mutations are not only associated with ultrahigh mutations but also associated with

a unique mutational signature [49,50], leading us to the hypothesis that the ultrahigh TMB and

unique mutation spectrum caused by POLE exonuclease mutations lead to synthetic lethality in

tumor cells with high aneuploidy burden (Figure 3C). We proposed that in highly aneuploid

tumor cells with numerous chromosome arm losses and gains, the inactivation of POLE's

exonuclease domain would lead to ultrahigh mutations across the genome. The unique

spectrum associated with these mutations may have a high probability of inactivating essential

genes located on chromosome arms with losses, leading to cancer cell death (Figure 3C).

Consistently, in POLD1 exonuclease mutated tumors where we did not observe strong mutual

exclusivity with high aneuploidy burden (Figure 2H), the TMB is substantially lower in

comparison to POLE exonuclease mutated tumors (Supplemental Figure 4B).

Consistent with our hypothesis, we observed in the real-world database that chromosomal arm

losses and random mutations (limited to variants with unknown significance) are less likely to

co-occur (odds ratio < 1) (Figure 3D). Next, we developed a probabilistic model to estimate the

likelihood of mutations caused by POLE exonuclease deficiency causing lethality by inactivating

essential genes (see Method for details).

$$Pr_{cell\ death\ per\ cell\ cycle} \approx 1 - exp\ (-\ n \times P_{arm} \times \alpha \times m)$$

In this model, n stands for the number of chromosome arms with losses in a tumor cell, P_{arm}

stands for the average probability of a mutation caused by POLE exonuclease deficiency hitting

an essential gene per chromosome arm, α stands for the portion of deleterious mutations hitting

essential genes, and it remains a challenge to determine the true value of α. Lastly, m stands

for the number of mutations POLE exonuclease deficiency can cause in a cell per cell cycle per

haploid genome. Based on this model, we determined that for a typical solid tumor cell with an

average of 6 chromosome arm losses (based on the real-world database), if 10% (α) of

mutations caused by POLE exonuclease deficiency hitting an essential gene are deleterious,

the probability of inactivating at least one essential gene, leading to cell death per cell cycle,

approaches 100%. If we assume α = 0.01, the probability of cell death per cell cycle under the

same condition is approximately 25% (Figure 3E). As the value of α becomes smaller, the

probability of cell death per cell cycle is less likely to induce a synthetic lethality or mutual

exclusivity phenotype.

The elastic net regression model also identified microsatellite instability as another top negative

predictor for aneuploidy score (Supplemental Table 1, Supplemental Figure 4F). Mutational

signatures related to mismatch repair deficiency also negatively correlate with low aneuploidy

burden (Supplemental Figure 4D). In previous reports, a synthetic lethal interaction between

microsatellite instability and high aneuploidy burden was also proposed, however, the

underlying mechanisms are unknown [51-53]. Interestingly, even though microsatellite unstable

tumors exhibit low aneuploidy burden, almost comparable to POLE exonuclease mutated

tumors (Supplemental figure 4F), these tumors showed significantly lower TMB compared to

POLE exonuclease-mutated tumors (Supplemental Figures 4E and 4G), but more comparable

to POLD1 exonuclease-mutated tumors. However, the TMB in POLD1 exonuclease-mutated

tumors are likely not sufficient to drive the mutual exclusivity with high aneuploidy burden

(Figure 2H). Therefore, it remains to be determined if the mechanism we proposed here for

POLE can be applied to microsatellite unstable tumors.

POLE exonuclease activity is essential for highly-aneuploid cancer cells but not

diploid cells

While our probabilistic modeling provided evidence that the mutual exclusivity between POLE

exonuclease mutations and high aneuploidy burden in tumor cells are likely driven by synthetic

lethality, the unknown value of the key variable α undermines the model's validity. Next, in order

to experimentally validate that POLE exonuclease activity is required for survival of highly-

aneuploid cancer cells, but not diploid cells, we utilized an inducible adenine base editor to

perform a mutagenesis screen using guide RNAs targeting POLE's exonuclease domain (Figure

4A) [54]. We selected four cancer cell lines with high aneuploidy scores (EFE184, MDST8,

SW480, SW837) and four diploid cell lines (DLD1, HCT116, HEC265, LS180) (Figure 4B). We

first validated the inducible expression of the adenine base editor across a few cell lines,

including a non-cancerous cell line 293, an aneuploidy cell line SW837 and a diploid cell line

DLD1(Supplemental Figure 5A). Then, we performed a mutagenesis screen on the eight cancer

cell lines (Figure 4C, see Method for details). Seven of the eight cell lines propagated post-

lentiviral transduction, except for MDST8 (Figure 4D). Expression of the adenine base editor

was maintained during the screen (Figure 4E), and guide RNAs were amplified at passage 8,

sequenced, and quantified. Guide RNAs targeting POLE's exonuclease domain were

consistently underrepresented in the three highly aneuploid cell lines (EFE184, SW480,

SW837), but not in the diploid cell lines (DLD1, HCT116, HEC265, LS180) (Figure 4F,

Supplemental Figure S5). This supports our hypothesis that POLE exonuclease domain

inactivation induces synthetic lethality in highly aneuploid cells but not in diploid cells.

Consistent with the base editor screen, we showed that, in the highly aneuploid cell line SW480,

ectopic expression of wild type POLE rescued the fitness defect caused by *POLE* knockdown,

while the two hot-spot mutations in the POLE exonuclease domain (P286R and V411L) failed to

rescue the defect (Figure 4G, H). Together, our experimental data across multiple cancer cell

line models demonstrated the essentiality of POLE exonuclease activity in highly aneuploid

cells.

243   **Discussion**

Developing therapies for cancers with high aneuploidy burden is of high medical importance.

Synthetic lethality is an attractive strategy for identifying therapeutic targets, due to the unknown

molecular mechanisms causing aneuploidy and the complex biological aberrations associated

with it. In this study, we leveraged a real-world cancer genomics database and identified that

mutations in the POLE exonuclease domain are mutually exclusive with high aneuploidy

burden. We hypothesized and validated that POLE exonuclease inactivation causes synthetic

lethality in highly aneuploid cells, likely by inactivating essential genes with copy number losses.

This discovery leads to a therapeutic hypothesis that selective inhibition of POLE exonuclease

may benefit patients with highly-aneuploid tumors.

Previous reports indicate that endometrial or colorectal cancer patients with POLE-mutated

tumors often have favorable clinical outcomes, even without immune checkpoint inhibitors

[55,56]. The mechanisms underlying this prognosis are not fully understood. One hypothesis

suggests that these tumors may exhibit high tumor antigens, enhancing immune system

recognition [57]. We speculate that the near-diploidy nature of *POLE*-mutated tumors could also

contribute to these favorable clinical outcomes. Additionally, selective inhibition of POLE

exonuclease might potentiate immune checkpoint inhibitors by enhancing tumor antigens

through hypermutation.

However, inhibiting POLE's exonuclease domain carries risks. This domain plays a critical role

in genome duplication proofreading for both cancer and normal cells. Individuals with germline

mutations in POLE's exonuclease domain often develop Polymerase Proofreading-Associated

Polyposis (PPAP) and have a higher risk of colorectal and uterine cancers later in life [45,46].

Interestingly, these patients do not exhibit strong acute disease symptoms. Murine models show

that heterozygous inactivation of POLE exonuclease presents no discernible phenotypes

compared to wild-type animals, while homozygous inactivation correlates with higher cancer

incidence, similar to patients with heterozygous mutations [58]. The hotspot somatic mutation

P286R exhibits a pronounced tumorigenesis phenotype in animal models that is not observed

with germline mutations [59]. This aligns with yeast studies where analogous substitutions

induced strong mutator phenotypes exceeding those of conventional proofreading mutations

[60]. The introduction of a charged residue within the exonuclease site appears to sterically

prevent channeling of the nascent DNA strand containing an erroneous base towards the

exonuclease active site residues [61], instead likely facilitating continuous DNA synthesis

(Supplemental Figure 6A). Similarly, V411L substitution lies within the 'wedge helix' directly

involved in the separation of erroneous base-containing nascent DNA strand from the template

during the transfer from polymerase to exonuclease site (Supplemental Figure 6B) [62]. A

longer side chain of leucine compared to valine is likely to push forward the wedge helix,

creating a stronger physical barrier preventing efficient transfer of the leading strand from the

polymerase to the exonuclease site. This might explain the high prevalence and hypermutator

phenotype of the V411L mutation in cancers.

Although there is precedent for developing small molecule inhibitors of RNase H-like

exonucleases [63], it remains to be determined whether similar pharmacological interventions

could induce hypermutator activity in POLE and what the therapeutic implications of such an

approach might be. Despite the safety and druggability challenges associated with POLE

exonuclease, our work proposes a novel concept of introducing synthetic lethality in highly

aneuploid tumor cells. Aneuploidy might expose many essential genes as vulnerabilities due to

large-scale chromosome losses and gains. Strategies to inhibit these essential genes might

offer therapeutic opportunities.

305 **Methods**

306 **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
POLE	Abcam	Cat. #Ab226848
Flag	Cell Signaling Technology	Cat. #14793s; RRID: AB_2572291
CAS9	Cell Signal Technology	Cat. #14697; RRID: AB_2750916
Tubulin	Cell Signal Technology	Cat. #3873s; RRID: AB_1904178
α-Actin	Cell Signal Technology	Cat. #3700s; RRID: AB_2242334
IRDye® 800CW	LiCorBio	Cat. #926-32211
IRDye® 680LT	LiCorBio	Cat. #926-68022
Critical commercial assays		
BCA assay Kit	Thermo Fisher Scientific	23228
DNA extraction Kit	QIAGEN	69506
Q5 High fidelity 2x MM Kit	New England BioLabs	M0492s
Qubit™ 1X dsDNA High Sensitivity (HS) and Broad Range (BR) Assay Kit	Thermo Fisher Scientific	Q33230
PCR purification Kit	QIAGEN	28106
SE Cell Line 4D-Nucleofector® X Kit	Lonza	V4XC-1032
SF Cell Line 4D-Nucleofector® X Kit	Lonza	V4XC-2032
Lipofectamine 3000 Kit	Thermo Fisher Scientific	L3000-015
Lipofectamine RNAiMAX	Invitrogen	13778-075
Oligonucleotides		
Primer gRNA_ABE Forward: TCTTGTGGAAAGGACGACTCGA	Integrated DNA Technologies	N/A
Primer gRNA_ABE Reverse: CGTACTCACTATAGGGCGAATTG	Integrated DNA Technologies	N/A
On-TargetPlus Human POLE siRNA GCGAGGAACAGGCGAAAUA	Dharmacon	L-020132-00-0005
Recombinant DNA		
POLE WT	Genentech PARS	DNA1AI68314
POLE. P286R	Genentech PARS	DNA1AI68315
POLE. V411L	Genentech PARS	DNA1AI68316
ABE Lentiviral Vector	Genscript	N/A
Software and algorithms		
Graphpad Prism 8.0 software	GraphPad Software, Inc.	http://www.graphpad.com/scientific software/prism/
R	Version 4.4	https://www.r-project.org/
glmnet	version 4.1-8	https://cran.r-project.org/web/packages/glmnet/index.html

Comprehensive genomic profiling in a real-world database

Comprehensive genomic profiling of formalin-fixed, paraffin-embedded (FFPE) tissue sections of 499,320 tumor samples during routine clinical care was performed using the FDA-approved FoundationOne®CDx or the laboratory developed tests, FoundationOne® or FoundationOne®Heme in a Clinical Laboratory Improvement Amendments–certified, College of American Pathologists–accredited laboratory (Foundation Medicine Inc.). Hybrid capture was carried out on at least 300 cancer-related genes and select introns from 28-34 genes frequently rearranged in cancer. Arm level gains and losses were called in a research use only capacity based on statistically lower or higher log2 CN for at least 50% of the arm, as previously described [31]. Approval for this study, including a waiver of informed consent and Health Insurance Portability and Accountability Act (HIPAA) waiver of authorization, was obtained from the Western Institutional Review Board (protocol no. 20152817).

Elastic net regression

The R package glmnet version 4.1-8 (https://cran.r-project.org/web/packages/glmnet/index.html) was used to fit an elastic net regression model using aneuploidy score (0-40) as a dependent variable. The following features were selected as predictor variables: disease ontology terms (n=487), metastasis status (n=2), chromosome sex (n=2), genetic ancestry (n=5), microsatellite instability (n=2) and genetic alterations of evaluable genes (n=349). The outputs of the predictor variables were converted to binary (0,1). The real-world dataset was split into a training set and a test set with a ratio of 2:1. To identify the optimal model, a total number of 21 different alpha values (0-1) were tested with an interval of 0.05. Mean Squared Error (MSE) was calculated for each alpha value to determine the performance of each model. Alpha value with the smallest MSE was chosen for the final model.

Probabilistic modeling of process of inactivating essential genes

Deficiency in POLE exonuclease leads to a unique mutational signature [49,50], with approximately 30% of the mutations on trinucleotide motif TCT and 20% of the mutations on TCG. This unique mutation spectrum allows us to estimate the probability of a mutation caused by POLE exonuclease deficiency hitting an essential gene. The list of essential genes in mammalian cells were extracted from a previous study [64]. First, we compute the probability of a mutation caused by POLE exonuclease deficiency falling on the genomic coding region for essential genes on each autosomal chromosome arm, through each trinucleotide motif:

339   P_{TCT-1p} = probability of a mutation hitting an essential gene on chr1p through motif TCT

340                          $P_{TCT-1p} = M_{TCT-1p} \div N_{TCT}$

341   N_{TCT} = total number of the trinucleotide motif TCT on human genome

342   M_{TCT-1p} = total number of the trinucleotide motif TCT on coding regions for all essential genes
343   located on chr1p

344   P_{TCG-1p} = probability of a mutation hitting an essential gene on chr1p through motif TCG

345                          $P_{TCG-1p} = M_{TCG-1p} \div N_{TCG}$

346   N_{TCG} = total number of the trinucleotide motif TCG on human genome

347   M_{TCG-1p} = total number of the trinucleotide motif TCG on coding regions for all essential genes
348   located on chr1p

349   Similar calculations can be done for all trinucleotide motifs.

350   P_{1p} = probability of a mutation hitting an essential gene on chr1p through any motif

351                    $P_{1p} = \Sigma\, w_i \times P_{i-1p}$, i: all trinucleotide motifs;

352   w_i: probability of a mutation through motif i [49]

353
354   TCT and TCG are the dominant motifs [49], we therefore can simplify the equation above

355                    $P_{1p} \approx 0.6 \times P_{TCT-1p} + 0.4 \times P_{TCG-1p}$

356   Similarly, the probabilities for all the 40 autosomal chromosome arms can be calculated as:

357                    $P_{1q} \approx 0.6 \times P_{TCT-1q} + 0.4 \times P_{TCG-1q}$

358   ...

359                    $P_{22q} \approx 0.6 \times P_{TCT-22q} + 0.4 \times P_{TCG-22q}$

360   The values of P_{1p} - P_{22q} can be computed from the trinucleotide count table in supplementary

361 table 2. In order to evaluate the probability as a function of chromosome arm loss and other

362 variables, we computed an average P_{arm} that will be used for all chromosomal arms in diploid

363 cells.

$$P_{arm} = (P_{1p} + P_{1q} + P_{2p} + P_{2q} + \ldots + P_{22q}) \div 40 = 0.00057$$

365 Even though we can compute the probability of a mutation hitting essential genes for each

366 chromosome arm, it remains unknown what portion of those mutations, when hitting an

367 essential gene, will lead to loss of function. We therefore introduced a variable $\boldsymbol{\alpha}$, proportion of

368 mutations on essential genes that are deleterious, to control the impact.

$$P_{arm} \times \alpha$$

Lastly, we need to determine the number of mutations (**m**) tumor cells can accumulate per cell

cycle per haploid genome when POLE exonuclease is deficient. This number was estimated

using the same approach as previously reported [65]. Briefly, the number of cell cycles (**cc**)

required for a single POLE exonuclease mutated cancer cell to give rise to a tumor were

calculated based on the average tumor volume for stage IV colorectal cancer (45cm^3) [66], and

the average volume of a mammalian cell ($1.8 \times 10^{-6} \text{ mm}^3$) [65]. The variable **m** is then calculated

as

$$m = total\ number\ of\ mutations \div cc \div 2$$

The average total number of mutations in POLE exonuclease mutated tumors is approximately

540,000 (TMB × 3,000) per genome based on the real-world data and TCGA, and the number

of cell cycles (**cc**) is approximately 36, based on which we computed **m** = 7500. However, the

TMB is estimated from a bulk of cancer cells, this estimation assumes that tumor cells are not

evolutionarily divergent, therefore this could be an overestimation of **m**. In a POLE exonuclease

deficient tumor cell with **n** chromosome arm losses, the probability of inactivating at least one

essential gene with copy number loss per cell cycle is:

$$Pr_{cell\ death\ per\ cell\ cycle} \approx 1 - (1 - n \times P_{arm} \times \alpha)^m$$

386 As **m** is large and **n** × P_{arm} × $\boldsymbol{\alpha}$ is small, this equation can be simplified as

$$Pr_{cell\ death\ per\ cell\ cycle} \approx 1 - \exp(-n \times P_{arm} \times \alpha \times m)$$

which follows from the Poisson approximation to the Binomial distribution, or equivalently the

fact that $(1 - p)^m \to \exp(-\lambda)$ when $m \to \infty$, $pm \to \lambda$. This formula could alternatively be derived by

assuming that mutations occur as a Poisson process with total rate **m** per cell cycle, of which **n**

× P_{arm} × α are lethal.

We note that the approximation above contains several simplifications that affect its accuracy.

First, there is a wide variance in the observed TMB across POLE-exonuclease deficient

samples, and thus in **m**. Secondly, different samples with the same **n** may have different

numbers of haploid bases depending on which chromosome arms are lost. Therefore, it would

improve accuracy to account for the specific TMB and aneuploidy pattern for any given sample.

Additionally, both **m** and **n** are estimated from bulk rather than single-cell data. Different cells

within a sample may have different values of **n**, causing overdispersion; meanwhile, the

estimated TMB includes mutations that are not present in all cells, which may inflate **m**, since it

is based on the number of mutations per cell divided by the number of cell cycles.

Western blots

Whole cell lysates were extracted using RIPA buffer (50 mM Tris HCl, 150 mM NaCl, 1.0% (v/v)

IGEPAL CA-630, 0.5% (w/v) Sodium Deoxycholate, 1.0 mM EDTA, 0.1% (w/v) SDS and 0.01%

(w/v) sodium azide at a pH of 7.4) supplemented with protease and phosphatase inhibitors

(Thermo Fisher Scientific, 78427,1861279). Protein concentrations were determined using the

BCA assay. Equal amounts of protein (20 µg per lane) were separated on 4-12% acrylamide

gels (Invitrogen, wg1403bx10) and transferred to membranes (Bio-Rad 1704159) using a

transfer system (trans-blot turbo, Bio-Rad). Membranes were blocked in 5% BSA (Gibco,15260-

037) in TBST (Tris-buffered saline with 0.1% Tween-20) for 30 minutes at room temperature,

followed by an overnight incubation at 4°C with primary antibodies diluted in blocking buffer at

1:500. After washing three times with TBST, membranes were incubated with secondary

antibodies for 1 hour at room temperature. Protein bands were visualized using Odyssey CLx

Infrared Imaging System.

Inducible adenine base editor screen

Tad-8e(V106W) was cloned downstream of a TRE3G doxycycline-inducible promoter in a

lentiviral expression vector with the EF1a promoter driving a puromycin resistance gene linked

to Tet-ON 3G. The vector also contains a human U6 promoter driving a single guide RNA

cassette into which the guide RNA library containing 89 POLE exonuclease tiling guide RNAs,

85 non-targeting control guide RNAs and 46 putative positive control guide RNAs were cloned.

All cloning was performed by Genscript. To perform the screen, 10 million cells were seeded in

15 cm plates and infected with the adenine base editor lentiviral library at a multiplicity of

infection (MOI) of 0.2. After 48 hours, the medium was replaced with a fresh growth medium

containing 3 µg/ml puromycin for 3–5 days. Following antibiotic selection, cells were trypsinized

and re-seeded at a density of 2 million cells per 10 cm dish in triplicates, with doxycycline added

to induce base editor expression. Cells were maintained for 8 passages, cell pellets containing 2

million cells were collected at the end of the experiment and before doxycycline treatment (as

baseline).

DNA Isolation and gRNA amplification

Genomic DNA was isolated from 2 million cells collected from the adenine base editor screen

using the QIAGEN DNA extraction kit, following the manufacturer's protocol. gRNAs were

amplified by PCR using the C1000 Touch Thermocycler (Bio-Rad). The PCR amplification was

performed using a standard three-step thermocycling protocol for 25 cycles: denaturation at

95 °C, primer annealing at 55 °C, and extension at 72 °C, employing the Q5 High-Fidelity 2X

Master Mix (New England BioLabs, M0492S). The resulting amplicons were purified using a

PCR purification kit (Qiagen, 28106) for downstream sequencing.

gRNA amplicon library preparation and sequencing

For generation of gRNA amplicon libraries, the KAPA HyperPrep Kit (Roche) that has

incorporated custom adapters and library amplification PCR primers from Integrated DNA

Technologies was used with an input of 40ng of amplicon DNA. Amplicon libraries were

quantified, and the average library size was determined using the D1000 ScreenTape on 4200

TapeStation (Agilent Technologies). Libraries were pooled and the Qubit dsDNA HS Assay Kit

(Thermo Fisher Scientific) was used to quantify the pool. Library pools were sequenced on

NextSeq 2000 (Illumina) to generate 3 million paired-end 51-base pair reads for each sample.

Analyses of base editor screen

Gene-level significance for differential barcode abundance was assessed using the FRY test

implemented in the limma R package

(https://bioconductor.org/packages/release/bioc/html/limma.html). Each cell line was evaluated

individually comparing the gRNA abundance targeting POLE exonuclease versus a set of non-

targeting control (NTC) gRNAs. For each gene, multiple gRNAs targeting that gene were

considered as a predefined set, and the FRY method was used to estimate a log2-Fold change

in barcode abundance with a corresponding p value.

POLE knockdown and rescue experiments

To establish cell lines stably expressing different *POLE* variants (WT, P286R, and V411L) in

SW480 cells, we transfected the cells with plasmids encoding human *POLE* variants using

electroporation. Briefly, 10^5 cells were suspended with 2 μg plasmids in the electroporation

buffer (SE and SF cell line 4D X kits, Lonza Bioscience). Electroporation was carried out using

optimized settings per the manufacturer's protocol (Lonza, 4D Nucleofector X unit). Post-

transfection, cells were plated in complete growth medium for recovery. Stable expression of

POLE was verified by western blot following puromycin selection for three weeks. For rescue

experiments, endogenous POLE expression was silenced using siRNA (5'-

GCGAGGAACAGGCGAAAUA). Cells were transfected with a siRNA mixture containing 20 nM

POLE-targeting siRNA and RNAiMAX transfection reagent (Invitrogen, Cat# 13778-075)

according to the manufacturer's protocol. Twenty-four hours post-transfection, 2,000 cells were

seeded into a 96-well plate and monitored by the Incucyte system (Essen BioScience) every 12

hours over a period of seven days. Incucyte's built-in software tools were employed to evaluate

cell confluence.

Resource availability

Lead contact

Further information or requests for data and resources should be directed to, and will be fulfilled

by the lead contact, Dr. Meng He (hem8@gene.com)

Materials availability

The all-in-one inducible adenine base editor system generated for this study can be requested

through the standard Genentech MTA process. Please reach out to the lead contact for details.

Data and code availability

Public datasets used in this study include TCGA and CCLE datasets, which are available for

download from cBioPortal (https://www.cbioportal.org/) and DepMap

(https://depmap.org/portal/), respectively. The sequencing data for the real-world database are

derived from clinical samples. All the relevant data supporting the findings of this study are

provided within the main manuscript and its supplementary files. Due to HIPAA requirements,

we are not authorized to share individualized patient genomic data, which contains potentially

identifying or sensitive patient information. Foundation Medicine is committed to collaborative

data analysis, with well-established, and widely utilized mechanisms by which investigators can

query our core genomic database of >700,000 de-identified sequenced cancers to obtain

aggregated datasets. For more information and mechanisms of access, please contact the

corresponding author(s) or the Foundation Medicine, Inc. Data Governance Council at

data.governance.council@foundationmedicine.com. All computational codes used in the

analysis are available for research purposes and can be provided upon request.

Acknowledgments

489   This study is funded by F. Hoffmann-La Roche AG

Author contributions

M.H. and S.G. contributed to the conception of the study. E.S., K.C., D.J. and S.M. contributed

to the curation of the real-world database. M.H. performed analysis. E.J.K. performed the wet

lab experiments. A.C. analyzed the base editor screen. K.D. designed the base editor system

and contributed to the interpretation of the data. S.J. performed the structural analysis of POLE.

R.R. and J. K. contributed to the statistical modeling. Y.W. and R. N. contributed to the

interpretation of the data and the writing and revision of the manuscript. M.S. and Y.L.

generated library from the gRNA library and performed next generation sequencing. A.D.

contributed to the interpretation of the data. E.J.K. and M.H. wrote the manuscript with input

from all authors. M.H., E.S., and S.G. provided supervision of this study.

Declaration of interests

501   M.H., S.G., E.J.K., A.C., K.D., S.J., R.R., J. K., Y.W., R. N., M.S., Y.L., and A.D. are Genentech Inc.

502   employees and may own stocks from F. Hoffmann-La Roche AG; E.S., K.C., and D.J. are

503   Foundation Medicine Inc. employees and may own stocks from F. Hoffmann-La Roche AG.

References

References

1.   Gordon, D. J., Resio, B., & Pellman, D. (2012). Causes and consequences of aneuploidy in
cancer. Nature Reviews Genetics, 13(3), 189–203. https://doi.org/10.1038/nrg3123

2.   Duijf, P. H. G., & Benezra, R. (2013). The cancer biology of whole-chromosome instability.
Oncogene, 32(40), 4727–4736. https://doi.org/10.1038/onc.2012.616

3.   Taylor, A. M., Shih, J., Ha, G., Gao, G. F., Zhang, X. Y., Berger, A. C., Schumacher, S. E., Wang,
C., Hu, H., Liu, J. F., Lazar, A. J., Cherniack, A. D., Beroukhim, R., Meyerson, M., & Network, C. G. A. R.
(2018). Genomic and Functional Approaches to Understanding Cancer Aneuploidy. Cancer Cell, 33(4),
676–+. https://doi.org/10.1016/j.ccell.2018.03.007

4.   Dewhurst, S. M., McGranahan, N., Burrell, R. A., Rowan, A. J., Grönroos, E., Endesfelder, D.,
Joshi, T., Mouradov, D., Gibbs, P., Ward, R. L., Hawkins, N. J., Szallasi, Z., Sieber, O. M., & Swanton, C.
(2014). Tolerance of Whole-Genome Doubling Propagates Chromosomal Instability and Accelerates
Cancer Genome Evolution. Cancer Discovery, 4(2), 175–185. https://doi.org/10.1158/2159-8290.Cd-13-
0285

5.   Sansregret, L., Vanhaesebroeck, B., & Swanton, C. (2018). Determinants and clinical implications
of chromosomal instability in cancer. Nature Reviews Clinical Oncology, 15(3), 139–+.
https://doi.org/10.1038/nrclinonc.2017.198

6.   Davoli, T., Xu, A. W., Mengwasser, K. E., Sack, L. M., Yoon, J. C., Park, P. J., & Elledge, S. J.
(2013). Cumulative Haploinsufficiency and Triplosensitivity Drive Aneuploidy Patterns and Shape the
Cancer Genome. Cell, 155(4), 948–962. https://doi.org/10.1016/j.cell.2013.10.011

7.   Vasudevan, A., Schukken, K. M., Sausville, E. L., Girish, V., Adebambo, O. A., & Sheltzer, J. M.
(2021). Aneuploidy as a promoter and suppressor of malignant growth. Nature Reviews Cancer, 21(2),
89–103. https://doi.org/10.1038/s41568-020-00321-1

8.   Santaguida, S., & Amon, A. (2015). Short- and long-term effects of chromosome mis-segregation
and aneuploidy. Nature Reviews Molecular Cell Biology, 16(8), 473–485. https://doi.org/10.1038/nrm4025

9.   Zhu, J., Tsai, H. J., Gordon, M. R., & Li, R. (2018). Cellular Stress Associated with Aneuploidy.
Developmental Cell, 44(4), 420–431. https://doi.org/10.1016/j.devcel.2018.02.002

10.   Ben-David, U., & Amon, A. (2020). Context is everything: aneuploidy in cancer. Nat Rev Genet,
21(1), 44–62. https://doi.org/10.1038/s41576-019-0171-x

11.   Leibowitz, M. L., Zhang, C. Z., & Pellman, D. (2015). Chromothripsis: A New Mechanism for

Rapid Karyotype Evolution. Annual Review of Genetics, Vol 49, 49, 183–211.

https://doi.org/10.1146/annurev-genet-120213-092228

12.   McGranahan, N., & Swanton, C. (2017). Clonal Heterogeneity and Tumor Evolution: Past,

Present, and the Future. Cell, 168(4), 613–628. https://doi.org/10.1016/j.cell.2017.01.018

13.   Bakhoum, S. F., & Cantley, L. C. (2018). The Multifaceted Role of Chromosomal Instability in

Cancer and Its Microenvironment. Cell, 174(6), 1347–1360. https://doi.org/10.1016/j.cell.2018.08.027

14.   Stingele, S., Stoehr, G., Peplowska, K., Cox, J., Mann, M., & Storchova, Z. (2012). Global

analysis of genome, transcriptome and proteome reveals the response to aneuploidy in human cells.

Molecular Systems Biology, 8. 608. https://doi.org/10.1038/msb.2012.40

15.   Sheltzer, J. M., & Amon, A. (2011). The aneuploidy paradox: costs and benefits of an incorrect

karyotype. Trends in Genetics, 27(11), 446–453. https://doi.org/10.1016/j.tig.2011.07.003

16.   Santaguida, S., Vasile, E., White, E., & Amon, A. (2015). Aneuploidy-induced cellular stresses

limit autophagic degradation. Genes & Development, 29(19), 2010–2021.

https://doi.org/10.1101/gad.269118.115

17.   Carter, S. L., Eklund, A. C., Kohane, I. S., Harris, L. N., & Szallasi, Z. (2006). A signature of

chromosomal instability inferred from gene expression profiles predicts clinical outcome in multiple human

cancers. Nature Genetics, 38(9), 1043–1048. https://doi.org/10.1038/ng1861

18.   Beroukhim, R., Mermel, C. H., Porter, D., Wei, G., Raychaudhuri, S., Donovan, J., Barretina, J.,

Boehm, J. S., Dobson, J., Urashima, M., Mc Henry, K. T., Pinchback, R. M., Ligon, A. H., Cho, Y. J.,

Haery, L., Greulich, H., Reich, M., Winckler, W., Lawrence, M. S.,…Meyerson, M. (2010). The landscape

of somatic copy-number alteration across human cancers. Nature, 463(7283), 899–905.

https://doi.org/10.1038/nature08822

19.   Zack, T. I., Schumacher, S. E., Carter, S. L., Cherniack, A. D., Saksena, G., Tabak, B., Lawrence,

M. S., Zhang, C. Z., Wala, J., Mermel, C. H., Sougnez, C., Gabriel, S. B., Hernandez, B., Shen, H., Laird,

P. W., Getz, G., Meyerson, M., & Beroukhim, R. (2013). Pan-cancer patterns of somatic copy number

alteration. Nature Genetics, 45(10), 1134–U1257. https://doi.org/10.1038/ng.2760

20.   Davoli, T., Uno, H., Wooten, E. C., & Elledge, S. J. (2017). Tumor aneuploidy correlates with

markers of immune evasion and with reduced response to immunotherapy. Science, 355(6322).

https://doi.org/10.1126/science.aaf8399

21.   Ly, P., Brunner, S. F., Shoshani, O., Kim, D. H., Lan, W. J., Pyntikova, T., Flanagan, A. M.,
Behjati, S., Page, D. C., Campbell, P. J., & Cleveland, D. W. (2019). Chromosome segregation errors
generate a diverse spectrum of simple and complex genomic rearrangements. Nature Genetics, 51(4),
705–+. https://doi.org/10.1038/s41588-019-0360-8

22.   Richardson, P. G., Barlogie, B., Berenson, J., Singhal, S., Jagannath, S., Irwin, D., Rajkumar, S.
V., Srkalovic, G., Alsina, M., Alexanian, R., Siegel, D., Orlowski, R. Z., Kuter, D., Limentani, S. A., Lee, S.,
Hideshima, T., Esseltine, D. L., Kauffman, M., Adams, J.,…Anderson, K. C. (2003). A phase 2 study of
bortezomib in relapsed, refractory myeloma. New England Journal of Medicine, 348(26), 2609–2617.
https://doi.org/DOI 10.1056/NEJMoa030288

23.   Jhaveri, K., Taldone, T., Modi, S., & Chiosis, G. (2012). Advances in the clinical development of
heat shock protein 90 (Hsp90) inhibitors in cancers. Biochimica Et Biophysica Acta-Molecular Cell
Research, 1823(3), 742–755. https://doi.org/10.1016/j.bbamcr.2011.10.008

24.   Tang, Y. C., Williams, B. R., Siegel, J. J., & Amon, A. (2011). Identification of Aneuploidy-
Selective Antiproliferation Compounds. Cell, 144(4), 499–512. https://doi.org/10.1016/j.cell.2011.01.017

25.   Oromendia, A. B., Dodgson, S. E., & Amon, A. (2012). Aneuploidy causes proteotoxic stress in
yeast. Genes & Development, 26(24), 2696–2708. https://doi.org/10.1101/gad.207407.112

26.   Dürrbaum, M., Kuznetsova, A. Y., Passerini, V., Stingele, S., Stoehr, G., & Storchová, Z. (2014).
Unique features of the transcriptional response to model aneuploidy in human cells. Bmc Genomics, 15.
https://doi.org/10.1186/1471-2164-15-139

27.   Cohen-Sharir, Y., McFarland, J. M., Abdusamad, M., Marquis, C., Bernhard, S. V., Kazachkova,
M., Tang, H., Ippolito, M. R., Laue, K., Zerbib, J., Malaby, H. L. H., Jones, A., Stautmeister, L. M., Bockaj,
I., Wardenaar, R., Lyons, N., Nagaraja, A., Bass, A. J., Spierings, D. C. J.,…Ben-David, U. (2021).
Aneuploidy renders cancer cells vulnerable to mitotic checkpoint inhibition. Nature, 590(7846), 486–+.
https://doi.org/10.1038/s41586-020-03114-6

28.   McConkey, D. J. (2017). The integrated stress response and proteotoxicity in cancer therapy.
Biochemical and Biophysical Research Communications, 482(3), 450–453.
https://doi.org/10.1016/j.bbrc.2016.11.047

29.   Jhaveri, K., Miller, K., Rosen, L., Schneider, B., Chap, L., Hannah, A., Zhong, Z. Y., Ma, W. N.,
Hudis, C., & Modi, S. (2012). A Phase I Dose-Escalation Trial of Trastuzumab and Alvespimycin
Hydrochloride (KOS-1022; 17 DMAG) in the Treatment of Advanced Solid Tumors. Clinical Cancer
Research, 18(18), 5090–5098. https://doi.org/10.1158/1078-0432.Ccr-11-3200

30.   Pavlick, D., Montesion, M., Heerema, N. A., Severson, E. A., Brennan, T., Frampton, G. M., He,
J., Byrd, J. C., Levine, R. L., Burd, A., Vergilio, J. A., & Albacker, L. A. (2018). A Next-Generation
Sequencing-Based Karyotyping Algorithm Reveals the Genomic Structure of Acute Myeloid Leukemia.
Blood, 132. https://doi.org/10.1182/blood-2018-99-116195

31.   Sharaf, R., Pavlick, D. C., Frampton, G. M., Cooper, M., Jenkins, J., Danziger, N., Haberberger,
J., Alexander, B. M., Cloughesy, T., Yong, W. H., Liau, L. M., Nghiemphu, P. L., Ji, M., Lai, A.,
Ramkissoon, S. H., & Albacker, L. A. (2021). FoundationOne CDx testing accurately determines whole
arm 1p19q codeletion status in gliomas. Neuro-Oncology Advances, 3(1).
https://doi.org/10.1093/noajnl/vdab017

32.   Brennan, C. W., Verhaak, R. G. W., McKenna, A., Campos, B., Noushmehr, H., Salama, S. R.,
Zheng, S. Y., Chakravarty, D., Sanborn, J. Z., Berman, S. H., Beroukhim, R., Bernard, B., Wu, C. J.,
Genovese, G., Shmulevich, I., Barnholtz-Sloan, J., Zou, L. H., Vegesna, R., Shukla, S. A.,…Network, T.
R. (2013). The Somatic Genomic Landscape of Glioblastoma. Cell, 155(2), 462–477.
https://doi.org/10.1016/j.cell.2013.09.034

33.   Papaemmanuil, E., Gerstung, M., Bullinger, L., Gaidzik, V. I., Paschka, P., Roberts, N. D., Potter,
N. E., Heuser, M., Thol, F., Bolli, N., Gundem, G., Van Loo, P., Martincorena, I., Ganly, P., Mudie, L.,
McLaren, S., O'Meara, S., Raine, K., Jones, D. R.,…Campbell, P. J. (2016). Genomic Classification and
Prognosis in Acute Myeloid Leukemia. New England Journal of Medicine, 374(23), 2209–2221.
https://doi.org/10.1056/NEJMoa1516192

34.   Zerbib, J., Bloomberg, A., & Ben-David, U. (2025). Targeting vulnerabilities of aneuploid cells for
cancer therapy. Trends in Cancer, 11(7), 642–664. https://doi.org/10.1016/j.trecan.2025.04.005

35.   Zheng, Y., Ahmad, K., & Henikoff, S. (2025). Total whole-arm chromosome losses predict
malignancy in human cancer. Proceedings of the National Academy of Sciences, 122(18).
https://doi.org/10.1073/pnas.2505385122

36.   Bloomfield, M., & Duesberg, P. (2016). Inherent variability of cancer-specific aneuploidy
generates metastases. Molecular Cytogenetics, 9. https://doi.org/10.1186/s13039-016-0297-x

37.   Hieronymus, H., Schultz, N., Gopalan, A., Carver, B. S., Chang, M. T., Xiao, Y. H., Heguy, A.,
Huberman, K., Bernstein, M., Assel, M., Murali, R., Vickers, A., Scardino, P. T., Sander, C., Reuter, V.,
Taylor, B. S., & Sawyers, C. L. (2014). Copy number alteration burden predicts prostate cancer relapse.
Proceedings of the National Academy of Sciences of the United States of America, 111(30), 11139–
11144. https://doi.org/10.1073/pnas.1411446111

38.   Zhang, C. Z., Spektor, A., Cornils, H., Francis, J. M., Jackson, E. K., Liu, S. W., Meyerson, M., &
Pellman, D. (2015). Chromothripsis from DNA damage in micronuclei. Nature, 522(7555), 179–+.
https://doi.org/10.1038/nature14493

39.   Thompson, S. L., & Compton, D. A. (2010). Proliferation of aneuploid human cells is limited by a
p53-dependent mechanism. Journal of Cell Biology, 188(3), 369–381.
https://doi.org/10.1083/jcb.200905057

40.   Fukasawa, K., Choi, T., Kuriyama, R., Rulong, S., & VandeVoude, G. F. (1996). Abnormal
centrosome amplification in the absence of p53. Science, 271(5256), 1744–1747. https://doi.org/DOI
10.1126/science.271.5256.1744

41.   Rayner, E., van Gool, I. C., Palles, C., Kearsey, S. E., Bosse, T., Tomlinson, I., & Church, D. N.
(2016). A panoply of errors: polymerase proofreading domain mutations in cancer. Nature Reviews
Cancer, 16(2), 71–81. https://doi.org/10.1038/nrc.2015.12

42.   Yuan, Z. N., Georgescu, R., Schauer, G. D., O'Donnell, M. E., & Li, H. L. (2020). Structure of the
polymerase ε holoenzyme and atomic model of the leading strand replisome. Nature Communications,
11(1). https://doi.org/10.1038/s41467-020-16910-5

43.   Henninger, E. E., & Pursell, Z. F. (2014). DNA Polymerase ε and Its Roles in Genome Stability.
Iubmb Life, 66(5), 339–351. https://doi.org/10.1002/iub.1276

44.   Hübscher, U., Nasheuer, H. P., & Syväoja, J. E. (2000). Eukaryotic DNA polymerases, a growing
family. Trends in Biochemical Sciences, 25(3), 143–147. https://doi.org/Doi 10.1016/S0968-
0004(99)01523-6

45.   Mosalem, O., Coston, T. W., Imperial, R., Mauer, E., Thompson, C., Yilma, B., Bekaii-Saab, T. S.,
Stoppler, M. C., & Starr, J. S. (2024). A comprehensive analysis of POLE/POLD1 genomic alterations in
colorectal cancer. Oncologist, 29(9), e1224–e1227. https://doi.org/10.1093/oncolo/oyae098

46.   Gola, M., Stefaniak, P., Godlewski, J., Jereczek-Fossa, B. A., & Starzynska, A. (2023). Prospects
of POLD1 in Human Cancers: A Review. Cancers, 15(6). https://doi.org/10.3390/cancers15061905

47.   Palles, C., Cazier, J. B., Howarth, K. M., Domingo, E., Jones, A. M., Broderick, P., Kemp, Z.,
Spain, S. L., Almeida, E. G., Salguero, I., Sherborne, A., Chubb, D., Carvajal-Carmona, L. G., Ma, Y.,
Kaur, K., Dobbins, S., Barclay, E., Gorman, M., Martin, L.,…Consortium, W. (2013). Germline mutations
affecting the proofreading domains of POLE and POLD1 predispose to colorectal adenomas and
carcinomas. Nature Genetics, 45(2), 136–144. https://doi.org/10.1038/ng.2503

48.   Deng, Y. L., Luo, S. Y., Deng, C. Y., Luo, T., Yin, W. K., Zhang, H. Y., Zhang, Y., Zhang, X. X.,

Lan, Y. J., Ping, Y. Y., Xiao, Y., & Li, X. (2019). Identifying mutual exclusivity across cancer genomes:

computational approaches to discover genetic interaction and reveal tumor vulnerability. Briefings in

Bioinformatics, 20(1), 254–266. https://doi.org/10.1093/bib/bbx109

49.   Alexandrov, L. B., Kim, J., Haradhvala, N. J., Huang, M. N., Ng, A. W. T., Wu, Y., Boot, A.,

Covington, K. R., Gordenin, D. A., Bergstrom, E. N., Islam, S. M. A., Lopez-Bigas, N., Klimczak, L. J.,

McPherson, J. R., Morganella, S., Sabarinathan, R., Wheeler, D. A., Mustonen, V., Getz,

G.,…Consortium, P. (2020). The repertoire of mutational signatures in human cancer. Nature, 578(7793),

94–+. https://doi.org/10.1038/s41586-020-1943-3

50.   Campbell, B. B., Light, N., Fabrizio, D., Zatzman, M., Fuligni, F., de Borja, R., Davidson, S.,

Edwards, M., Elvin, J. A., Hodel, K. P., Zahurancik, W. J., Suo, Z. C., Lipman, T., Wimmer, K., Kratz, C.

P., Bowers, D. C., Laetsch, T. W., Dunn, G. P., Johanns, T. M.,…Shlien, A. (2017). Comprehensive

Analysis of Hypermutation in Human Cancer. Cell, 171(5), 1042–+.

https://doi.org/10.1016/j.cell.2017.09.048

51.   Pino, M. S., & Chung, D. C. (2010). The Chromosomal Instability Pathway in Colon Cancer.

Gastroenterology, 138(6), 2059–2072. https://doi.org/10.1053/j.gastro.2009.12.065

52.   Setton, J., Zinda, M., Riaz, N., Durocher, D., Zimmermann, M., Koehler, M., Reis-Filho, J. S., &

Powell, S. N. (2021). Synthetic Lethality in Cancer Therapeutics: The Next Generation. Cancer Discovery,

11(7), 1626–1635. https://doi.org/10.1158/2159-8290.Cd-20-1503

53.   Zhou, Y. M., Chen, X. F., Chen, J., Kendrick, C. D., Ramanathan, R. K., Graham, R. P., Kossick,

K. F., Boardman, L. A., & Barrett, M. T. (2024). Genomic landscape of diploid and aneuploid microsatellite

stable early onset colorectal cancer. Scientific Reports, 14(1). https://doi.org/10.1038/s41598-024-59398-

5

54.   Richter, M. F., Zhao, K. V. T., Eton, E., Lapinaite, A., Newby, G. A., Thuronyi, B. W., Wilson, C.,

Koblan, L. W., Zeng, J., Bauer, D. E., Doudna, J. A., & Liu, D. V. R. (2020). Phage-assisted evolution of

an adenine base editor with improved Cas domain compatibility and activity. Nature Biotechnology, 38(7),

883–U884. https://doi.org/10.1038/s41587-020-0453-z

55.   Yao, X. H., Feng, M., & Wang, W. (2024). The Clinical and Pathological Characteristics of POLE-

Mutated Endometrial Cancer: A Comprehensive Review. Cancer Management and Research, 16, 117–

125. https://doi.org/10.2147/Cmar.S445055

56.   Yu, S., Sun, Z., Zong, L., Yan, J., Yu, M., Chen, J., & Lu, Z. (2022). Clinicopathological and
molecular characterization of high-grade endometrial carcinoma with POLE mutation: a single center
study. Journal of Gynecologic Oncology, 33(3). https://doi.org/10.3802/jgo.2022.33.e38

57.   Ma, X. X., Riaz, N., Samstein, R. M., Lee, M., Makarov, V., Valero, C., Chowell, D., Kuo, F. S.,
Hoen, D., Fitzgerald, C. W. R., Jiang, H., Alektiar, J., Alban, T. J., Juric, I., Parthasarathy, P. B., Zhao, Y.,
Sabio, E. Y., Verma, R., Srivastava, R. M.,…Chan, T. A. (2022). Functional landscapes of and mutations
in checkpoint blockade-dependent antitumor immunity. Nature Genetics, 54(7), 996–+.
https://doi.org/10.1038/s41588-022-01108-w

58.   Albertson, T. M., Ogawa, M., Bugni, J. M., Hays, L. E., Chen, Y., Wang, Y. P., Treuting, P. M.,
Heddle, J. A., Goldsby, R. E., & Preston, B. D. (2009). DNA polymerase ε and δ proofreading suppress
discrete mutator and cancer phenotypes in mice. Proceedings of the National Academy of Sciences of
the United States of America, 106(40), 17101–17104. https://doi.org/10.1073/pnas.0907147106

59.   Li, H. D., Lu, C., Zhang, H., Hu, Q., Zhang, J., Cuevas, I. C., Sahoo, S. S., Aguilar, M., Maurais,
E. G., Zhang, S., Wang, X., Akbay, E. A., Li, G. M., Li, B., Koduru, P., Ly, P., Fu, Y. X., & Castrillon, D. H.
(2020). A PoleP286R mouse model of endometrial cancer recapitulates high mutational burden and
immunotherapy response. JCI insight, 5(14), e138829. https://doi.org/10.1172/jci.insight.138829

60.   Kane, D. P., & Shcherbakova, P. V. (2014). A Common Cancer-Associated DNA Polymerase ε
Mutation Causes an Exceptionally Strong Mutator Phenotype, Indicating Fidelity Defects Distinct from
Loss of Proofreading. Cancer Research, 74(7), 1895–1901. https://doi.org/10.1158/0008-5472.Can-13-
2892

61.   Parkash, V., Kulkarni, Y., ter Beek, J., Shcherbakova, P. V., Kamerlin, S. C. L., & Johansson, E.
(2019). Structural consequence of the most frequently recurring cancer-associated substitution in DNA
polymerase ε. Nature Communications, 10. https://doi.org/10.1038/s41467-018-08114-9

62.   Roske, J. J., & Yeeles, J. T. P. (2024). Structural basis for processive daughter-strand synthesis
and proofreading by the human leading-strand DNA polymerase Pol ε. Nature Structural & Molecular
Biology, 31(12). https://doi.org/10.1038/s41594-024-01370-y

63.   Huang, K. W., Hsu, K. C., Chu, L. Y., Yang, J. M., Yuan, H. S., & Hsiao, Y. Y. (2016).
Identification of Inhibitors for the DEDDh Family of Exonucleases and a Unique Inhibition Mechanism by
Crystal Structure Analysis of CRN-4 Bound with 2-Morpholin-4-ylethanesulfonate (MES). Journal of
Medicinal Chemistry, 59(17), 8019–8029. https://doi.org/10.1021/acs.jmedchem.6b00794

64.   Hart, T., Chandrashekhar, M., Aregger, M., Steinhart, Z., Brown, K. R., MacLeod, G., Mis, M.,
Zimmermann, M., Fradet-Turcotte, A., Sun, S., Mero, P., Dirks, P., Sidhu, S., Roth, F. P., Rissland, O. S.,

Durocher, D., Angers, S., & Moffat, J. (2015). High-Resolution CRISPR Screens Reveal Fitness Genes

and Genotype-Specific Cancer Liabilities. Cell, 163(6). https://doi.org/10.1016/j.cell.2015.11.015

65.   Shlien, A., Campbell, B. B., de Borja, R., Alexandrov, L. B., Merico, D., Wedge, D., Van Loo, P.,

Tarpey, P. S., Coupland, P., Behjati, S., Pollett, A., Lipman, T., Heidari, A., Deshmukh, S., Avitzur, N.,

Meier, B., Gerstung, M., Hong, Y., Merino, D. M.,…Repair, B. M. (2015). Combined hereditary and

somatic mutations of replication error repair genes result in rapid onset of ultra-hypermutated cancers.

Nature Genetics, 47(3), 257–+. https://doi.org/10.1038/ng.3202

66.   Park, J. Y., Kim, S. H., Lee, S. M., Lee, J. S., & Han, J. K. (2017). CT volumetric measurement of

colorectal cancer helps predict tumor staging and prognosis. Plos One, 12(6).

https://doi.org/10.1371/journal.pone.0178522

Figure legends

Figure 1: Characterization of chromosomal arm level aneuploidy in the real-world comprehensive genomic profiling database

(A) Pie chart summarizing the real-world database by primary cancer type.

(B) Donut charts summarizing the real-world database by chromosome sex, histological types, metastasis status and genetic ancestry.

(C-F) Frequencies of each chromosomal arm loss or gain in carcinoma (C), sarcoma (D), glioma (E) and leukemia (F).

(G) Distribution of aneuploidy scores across major cancer types (In each boxplot, the center line represents the median value. The lower and upper hinges correspond to the first and third quartiles, respectively, while the whiskers extend up to 1.5x the interquartile range. This applies to all the boxplots presented in this study).

Figure 2: Mutations in POLE exonuclease domain are mutually exclusive with high aneuploidy burden

(A) Elastic net regression model identified genes with non-zero coefficient to be predictors for aneuploidy scores. Top positive predictor genes are highlighted in red, while top negative predictor genes are highlighted in blue.

(B) Distribution of aneuploidy scores across tumors with different *TP53* alterations. RE, structural rearrangement; SV, short variant; CN, copy number alteration (homozygous deletion). Student t-tests were used for two group comparisons.

(C) Frequency of *POLE* mutation across various cancer types in the real-world database.

(D) Representation of POLE protein domains.

(E) Distribution of aneuploidy scores in tumors with different *POLE* alterations across various cancer types in the real-world database. mut.exo, mutations within the POLE exonuclease domain (a.a.268-a.a.471); mut.other, mutations outside the POLE exonuclease domain; wt, wild type. Student t-tests were used for two group comparisons (mut.exo vs. wt or mut.other vs. wt), * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, ns $p > 0.05$.

(F) Distribution of aneuploidy scores in tumors with different *POLE* alterations in TCGA.

(G) Distribution of aneuploidy scores in cell lines with different *POLE* alterations in CCLE.

(H) Distribution of aneuploidy scores in tumors with different *PODL1* alterations in the real-world
database.

Figure 3: High mutation burden and the unique mutation spectrum caused by POLE
exonuclease deficiency have a high probability of inactivating essential genes in highly
aneuploid cells.

(A) Distribution of tumor mutational burden (TMB) in tumors with different *POLE* alterations in
the real-world database.

(B) Aneuploidy scores and the mean TMB are negatively correlated in *POLE*-mutated tumors
(blue line: a linear regression model fit by lm() function in R)

(C) Illustration of proposed mechanism of mutual exclusivity between high aneuploidy burden
and POLE exonuclease mutations: mutations caused by POLE exonuclease deficiency have a
high probability of inactivating essential genes on chromosome arm harboring losses in highly
aneuploid cells.

(D) Most of the chromosomal arms are less likely to have co-occurrence of arm losses and
genetic alterations (limited to variants with unknown significance), odds ratio < 1. Each dot
represents an autosomal chromosomal arm.

(E) Probabilistic modeling of cell death per cell cycle in cells with POLE exonuclease deficiency.
$\boldsymbol{Pr_{cell\ death\ per\ cell\ cycle}} \approx \boldsymbol{1 - exp\ (- n \times P_{arm} \times \alpha \times m)}$, $\boldsymbol{n}$ = number of chromosome arm losses
(mean = 6), $\boldsymbol{P_{arm}}$ = probability of a mutation (caused by POLE exonuclease deficiency) hitting
an essential gene divided by total number of chromosome arms ($\boldsymbol{P_{arm}}$=0.00057), $\boldsymbol{\alpha}$ =
percentage of POLE exonuclease deficiency-induced mutations on essential genes that are
deleterious (unknown), $\boldsymbol{m}$ = number of mutations caused by POLE exonuclease deficiency per
cell cycle per haploid genome ($\boldsymbol{m}$=7500 estimated for stage IV colorectal cancer), refer to
methods section for more details.

Figure 4: Mutagenesis screening validated that POLE exonuclease is essential for the
survival of highly aneuploid cells but not diploid cells.

(A) Illustration of the inducible adenine base editor screen tiling the exonuclease domain of

POLE.

(B) Cell lines used for the adenine base editor screen.

(C) Experimental flow of the adenine base editor screen.

(D) Cell growth curves post viral transduction and induction of adenine base editor.

(E) Western blot confirming the sustained expression of the adenine base editor system during

the experiment.

(F) Compared to non-targeting control gRNAs (NTC), gRNAs tiling the exonuclease domain of

POLE are significantly underrepresented in the aneuploid cell lines (EFE184, SW480, SW837),

but not in the diploidy cell lines (DLD1, HCT116, HEC265, LS180). P values and log2 fold

change were determined by comparing guide RNA counts at passage 8 to passage 0.

(G) Western blot confirming the siRNA mediated knockdown of endogenous POLE, and over

expression of ectopic POLE variants (WT, P286R and V411L) in SW480 cell line.

(H) Growth curves of SW480 cells under different genetic backgrounds.

Figure 1

Figure 2

Figure 3

Figure 4

A

Exonuclease — Polymerase — DUF1744

sgRNAs tiling the exonuclease domain of *POLE*

hU6 — gRNA — EF1α — PuroR — T2A — Tet-On

Inducible Adenine Base Editor:TadA-8e(V106W)

Cas9n(D10A) — TadA-8e(V106W) — TRE

B

Cell lines	Lineage	Aneu_score
EFE184	Endometrial	13
MDST8	Colorectal	14
SW480	Colorectal	24
SW837	Colorectal	13
DLD1	Colorectal	0
HCT116	Colorectal	0
HEC265	Endometrial	0
LS180	Colorectal	0

C

Cell seeding — Viral transduction — Puromycin selection — Doxycycline treatment — Cell harvest & gRNA sequencing

Passage 0   Passage 1   Passage 8

Media replacement Every 72hrs

D

Cell cycle number vs passage

- SW480
- HCT116
- DLD1
- LS180
- HEC265
- SW837
- MDST8
- EFE184

E

EFE184, SW480, SW837 — Cas9, Ac/Tu

-Dox +Dox, P2 +Dox, P7

DLD1, HCT116, HEC265, LS180 — Cas9, Ac/Tu

F

Aneuploid cell lines | Diploid cell lines

EFE184 p=4.4e-5
SW480 p=1.9e-7
SW837 p=6.7e-8
DLD1 p=0.091
HCT116 p=0.090
HEC265 p=0.15
LS180 p=0.14

NTC   POLE.exo

POLE V411L

+ siPOLE
POLE-Flag
POLE
Ac/Tu

H

SW480

Confluency (%) vs Days

- siPOLE
- siPOLE + WT POLE
- siPOLE + POLE P286R
- siPOLE + POLE V411L