Taste dysfunction in Long COVID.

Hanna Morad[1], Tytti Vanhala[2] Marta A. Kisiel [3], Agnes Andreason[3], Mei Li[4], Göran Andersson[2], Göran Laurell[5], Thomas E. Finger[# 4], and Göran Hellekant[# 2,6].

Clinic of Ear, Nose and Throat diseases (ENT), Uppsala University Hospital, Uppsala, Sweden.

Department of Animal Biosciences, Swedish University of Agricultural Sciences, Uppsala, Sweden.

Occupational and Environmental Medicine, Department of Medical Sciences, Uppsala University, Uppsala, Sweden

Rocky Mountain Taste & Smell Center, Department of Cell & Developmental Biology, University of Colorado School of Medicine, Aurora CO 80045, USA.

Department of Surgical Sciences, Uppsala University, Uppsala, Sweden.

University of Wisconsin Madison, School of Veterinary Medicine, Madison, WI, USA.

#Shared Senior authors
Correspondence to be sent to: Tom.Finger@CUAnschutz.edu

Abstract

Persistent taste dysfunction may occur both as acute and long-term symptoms of SARS-CoV-2 infection (Long COVID), yet the underlying mechanisms are unknown at the histological, cellular, and molecular levels. This study investigates the underlying pathology in 28 non-hospitalized subjects who reported persistent taste disturbances for over 12 months after testing positive for SARS-CoV-2. To assess taste function, subjects completed the Waterless Empirical Taste Test (WETT), which quantifies the subject's ability to taste each of the five human taste qualities: sweet, umami, bitter, sour, and salty. Biopsies of fungiform papillae were collected from 20 participants and analyzed histologically for overall taste bud structure and innervation and by quantitative PCR (qPCR) for expression of markers for different taste receptor cells (TRCs). Although all subjects reported subjective taste dysfunction, only five scored below the 20th percentile on overall taste sensitivity. However, 12 subjects exhibited total loss of one or more taste qualities and another 13 subjects tested below the 95% confidence interval for at least one taste modality. Notably, loss of sweet, umami, or bitter tastes -- qualities mediated by a PLCβ2-transduction cascade -- was significantly more common than loss of sour and salty, and this loss correlated with reduced expression of PLCβ2 mRNA. Histological analysis revealed generally preserved taste bud structure and innervation in all cases, with occasional disorganization resulting in isolated PLCβ2-immunoreactive cells. Our findings suggest long-term taste dysfunction after COVID-19 disproportionately impacts PLCβ2-dependent taste qualities and is not due to widespread structural damage of the taste periphery.

41
42
agnes.andreason@gmail.com   ORCID 0009-0002-3236-2201
hannaa_morad@hotmail.com   ORCID 0009-0005-0813-6273
mei.li@cuanschutz.edu   ORCID 0009-0009-9170-9376
tytti.vanhala@slu.se
marta.kisiel@medsci.uu.se   ORCID 0000-0002-0410-1509
goran.andersson@slu.se   ORCID. 0000-0001-5131-3144
goran.laurell@uu.se   ORCID 0000-0002-7760-246X
ghellekant@gmail.com   ORCID   0000-0003-3562-8677
tom.finger@CUAnschutz.edu   ORCID   0000-0001-5923-5099
52
53
54

55   **CORRESPONDING AUTHORS**

56   Correspondence to Göran Hellekant and Thomas E. Finger

57
58

INTRODUCTION

The COVID-19 pandemic has brought unprecedented attention to impairments of olfaction and taste as hallmark symptoms of acute SARS-CoV-2 virus infection (Cooper, Brann et al. 2020, Hannum, Koch et al. 2023). In most individuals, this dysfunction resolves within weeks, but can persist for months for others. Prolonged dysfunction lasting more than 3 months is now recognized as part of post-acute COVID-19 syndrome (PACS) or Long COVID (Soriano, Murthy et al. 2022, Ely, Brown et al. 2024). Ongoing taste dysfunction may occur even after resolution of other symptoms (Boldes, Ritter et al. 2024, Xu, Wang et al. 2024) thus raising questions about the underlying mechanisms. Understanding the long-term pathophysiology of taste dysfunction and, in extreme cases, taste loss is crucial for patient care and the development of targeted rehabilitation strategies (Lee, Kisiel et al. 2025).

Sensations of taste are largely mediated by lingual taste buds (TBs) situated in three types of papillae: fungiform, vallate, and foliate. The fungiform papillae (FP), are concentrated on and around the tip of the tongue, making them easy to biopsy. A typical human has about 200 FP with 0-50 taste buds (TB) in each FP (Miller 1986, Miller 1988 , Miller 1989 , Miller and Reedy 1990, Cheng and Robinson 1991).

Taste buds are complex multicellular epithelial end organs comprising between 20 and 100 taste cells (TCs) with approximately half being taste receptor cells (TRCs) (Ohtubo and Yoshii 2011, Ogata and Ohtubo 2020, Yang, Dzowo et al. 2020, High and Finger 2025). Recent histochemical, immuno-histochemical and molecular methods, allow for correlation of TRC subtypes with specific taste qualities. Type II TRCs detect sweet, umami, and bitter tastes via distinct G-protein-coupled receptor (GPCR) pathways involving downstream PLCβ2 signaling cascades (Shi, Zhang et al. 2003, Zhang, Hoon et al. 2003). Sweet taste is mediated via TAS1R2/R3

heterodimeric receptors, and umami by TAS1R1/R3; bitter taste involves TAS2Rs, another

family of taste GPCRs. In contrast, Type III cells are responsible for transduction of ionic taste

qualities, i.e. sour and high concentrations of salt (Chang, Waters et al. 2010, Matsumoto,

Ohmoto et al. 2011) while Type I TCs play a supporting role, like glia cells in the CNS (Bartel,

Sullivan et al. 2006, Rodriguez, Roebber et al. 2021, Wilson, Lasher et al. 2025). In addition,

taste buds contain a population of post-mitotic Type IV cells which mature to differentiate into

the other cell types. Disruption to these distinct cellular pathways or the morphological integrity

of taste buds could contribute to the observed taste dysfunction in COVID-19 (Huang, Perez et al.

2021).

TRCs undergo continuous turnover, with new TRCs differentiating only when taste fibers reach

them (Farbman 1965, Farbman 1980, Hellekant, Kasahara et al. 1987)(Farbman, Guagliardo et al.

2004, Lee, Macpherson et al. 2017). Approximately 70–80% of mature TRCs form synapses with

nerve fibers, highlighting the importance of neural connectivity in maintaining taste

function (Lee, Macpherson et al. 2017, Wilson, Lasher et al. 2022). Ongoing post-viral disruption

of the nerve-TC interactions could well lead to taste dysfunction.

SARS-CoV-2 infects cells via the ACE2 receptor and the TMPRSS2 protease, both of which are

expressed in oral epithelial cells, including Type II TRCs (Doyle, Appleton et al. 2021, Yao,

Doyle et al. 2023). This suggests a plausible mechanism for direct viral damage to the taste

system. Additionally, inflammatory processes, immune dysregulation, or altered saliva

composition may contribute to long-term taste changes following viral infection.

Given the clinical and biological implications of long-term taste dysfunction, this study aims to

investigate self-reported and objectively measured taste impairment in participants who

experienced non-hospitalised COVID-19. Using a combination of psychophysical testing,

histological evaluation and molecular analysis -- including quantitative polymerase chain reaction

(qPCR) for taste cell markers and viral RNA detection -– this study seeks to elucidate the

structural and molecular changes underlying persistent taste dysfunction following COVID-19

infection.

MATERIAL & METHODS

Study design

This study involved participants aged ≥18 years who had experienced symptomatic COVID-19

and a confirmed SARS-CoV-2 infection verified by a positive PCR test from nasopharyngeal

swabs between 2020 and 2021 (inclusion criteria). All participants had non-hospitalized COVID-

19 and, all but one, reported long-term taste disturbances. The study population was recruited

between November 2021 and June 2022 from two sources at Uppsala University Hospital: the

COMBAT post-COVID study and the Ear, Nose and Throat (ENT) outpatient clinic.

From the COMBAT post-COVID cohort, 18 participants underwent psychophysical taste

evaluation. Of these, 12 consented to undergo a fungiform papilla biopsy (participants #7, 8, 9,

12, 13, 14, 15, 18, 19, 23, 24, 26). Additionally, ten participants were recruited from the ENT

outpatient clinic, eight of whom underwent biopsy (participants #17, 21, 22, 24, 25, 27, 28, 29).

In total, eight participants did not receive a biopsy.

Study population

The COMBAT post-COVID project aimed to monitor persistent symptoms and long-term

consequences among non-hospitalized patients diagnosed with COVID-19. Initially, 725

individuals who tested positive for SARS-CoV-2 between March and December 2020 were

invited to complete a follow-up questionnaire 12 months after symptom onset. Of these, 401

participants responded (response rate: 74%). The cohort has been described in detail elsewhere

(Kisiel, Nordqvist et al. 2021, Kisiel, Janols et al. 2022, Kisiel, Lee et al. 2023, Kisiel, Lee et al.

2023, Janols, Wadsten et al. 2024). Among the respondents, 68 individuals reported ongoing taste

and/or smell disturbances. These individuals were contacted by telephone and invited to

participate in the current study if they reported persistent taste loss and were available for in-

person evaluation.

Ten additional patients attending the ENT outpatient clinic in 2021 were randomly selected and

invited to participate in the study if they met the same inclusion criteria. In addition, one

participant (#17) reported no taste loss but had an active COVID infection at the time of

sampling.

Clinical and sociodemographic variables

From the COMBAT post-COVID cohort, information was documented about age, gender,

symptoms at COVID-19 onset, self-reported health before and one year after COVID-19, and

persistent symptoms one year following COVID-19. Comorbidities were collected from medical

records obtained from primary and secondary care from January 2010 to January 2022.

Symptoms at COVID-19 onset were also retrieved from medical records from the PCR testing

protocols. Persistent symptoms and smoking habits were collected from the survey. Subjects

were asked to estimate their general health in the study at one-year follow-up, from 0-100, where

0 was the poorest and 100 was the best. Additional subjects (n=11) were characterized by age and

sex only.

Ethical Approval

The study was approved by the institutional ethics committee of the University Hospital in

Uppsala (EPN number 2020-05707) and the Swedish Ethical Review Authority (EPM number

2021-00878) and complies with the Declaration of Helsinki for Medical Research involving

Human Subjects. The patients all gave written informed consent to the procedures. The

Combined Institutional Review Board at the University of Colorado at Anschutz approved the

transfer and analysis of the de-identified samples collected in Sweden. The study protocol in

Colorado (COMIRB number 20-1325) was determined Exempt/Non-Human Subject Research

and was conducted in accordance with 45 CFR 46 and institutional policies. All data were

pseudonymized to protect patient privacy.

Taste evaluation

As mentioned, the human sense of taste is divided into five taste qualities: sweet, umami, bitter,

sour and salty, (reviewed in:(Lindemann 1996, Hellekant 2024). To assess a subject's ability to

taste we used the quantitative and qualitative Waterless Empirical Taste Test (WETT®) by

Sensonics (Haddon Heights, NJ, USA). The WETT® method measures the ability to taste each

taste quality with an eight-point scale from zero (no taste) to a maximum of eight (all

concentrations identified). The WETT® measures the ability to identify various concentrations of

the above five basic taste qualities with a minimum number of trials. It consists of 53 strips

impregnated with one tastant (a substance that produces a taste sensation) representing a specific

taste quality. A monomer cellulose pad ($1\square\times\square2.5$ cm) is present on one side of each strip,

containing either: dried sucrose (sweet) (0.20, 0.10, 0.05, or 0.025 g/ml), citric acid (sour) (0.025,

0.05, 0.10, or 0.20 g/ml), sodium chloride (salty) (0.0313, 0.0625, 0.125, or 0.25 g/ml), caffeine

(bitter) (0.011, 0.022, 0.044, or 0.088 g/ml), monosodium glutamate (umami; brothy) (0.017,

0.034, 0.068, or 0.135 g/ml), or blank control strips, with no added chemical stimulus. The

subjects were instructed to place the appropriate strip on the tongue and move it along the surface

of the tongue for at least 5–10 seconds. The subject indicated which taste sensation was perceived

sött (sweet), bittert/beskt (bitter), surt (sour), salt, or buljongliknande smak/umami (umami), or

whether no taste was sensed by verbally reporting their response to be entered on the designated

line of the response form. In the WETT® test sequence, the concentrations of each stimulus are

presented twice (eight total trials/tastant). In the first half of the test, the concentrations ascend

from weak to strong with the different tastants presented randomly, with no tastant blanks

included. In the second half of the test, the reverse presentation order is used, i.e., going from

strong to weak concentrations. Scores range from 53 (all correct) to 13 (no tastes, corresponding

to the number of control strips in the test). Each subject's results were compared with normative

values for each gender and age given in Doty et al (Doty, Sharetts et al. 2025). The average taste

score for the subjects, who had consented to the biopsy, did not differ significantly from those

without biopsy in terms of total tasting ability (t(26)=1.52, p>.05). The results are presented in

Table 1.

Biopsy procedures

For all but 2 subjects, five to eight fungiform papillae (FP) were biopsied from the subject's

tongue immediately following or within 2 months of the WETT test as indicated in Table 1. For 2

subjects with normal taste function on the WETT test (subjects 26 and 21 ; respectively 44th and

70th percentile taste scores), biopsies were taken 4 months following the WETT test. The tongue

was first anesthetized with a Xylocaine spray (100mg/ml Lidocaine with 10mg/spray dose), then

blue or green food coloring was added to enhance the tongue's fungiform papillae (FP). The

tongue was held steady with a gauze, and biopsies of the FP were taken with a 3-mm sterile

microscissor with a curved tip. The FP was cut one by one by holding down the scissor tip at the

base of the chosen FP, which was removed with a single cut and put in fixative for histology or

RNA*later*™ (Thermo Fisher Scientific Inc., Waltham, MA, USA). The biopsies for histological

analysis were fixed in 4% buffered formaldehyde for at least four hours or overnight and then

moved to phosphate-buffered saline (PBS) solution. The biopsies were stored at 4°C until

shipping for analyses from Sweden to the United States for histopathological studies. The

biopsies for RNA analysis were placed in RNA*later*™ stabilization solution and stored at room

temperature overnight and then stored in the freezer at -18 °C initially and then at -80°C until

analyses.

Histology

Upon receipt, all specimens were cryoprotected in 20% Sucrose in 0.1M phosphate buffer

overnight at 4°C. Then, the tissues were cryo-sectioned (12-14µm) onto Superfrost Plus

microscope slides (Thermo Fisher Scientific, (Catalog# 1255015) and air-dried. Slides were

rinsed in 0.1M PBS pH 7.2. Antigen retrieval was performed with 10 mM sodium citrate buffer

of pH 6.0 at 85°C for 10 minutes before incubation in blocking solution (2% normal donkey

serum, 1% bovine serum albumin, and 0.3% Triton in 0.1M PBS) for an hour at room

temperature. Slides were then incubated for four days in specific primary antibodies in blocking

solution at 4°C including the following antisera:

• Mouse anti-Tubulin β3 (clone TUJ1) at 1:1000 dilution (BioLegend, San Diego, CA,

USA, catalog# 801202; RRID:AB_2313773) (for total innervation),

• Goat anti-GNAT3 at 1:500 dilution (Aviva Systems Biology, San Diego, CA, USA,

catalog# OAEB00418; RRID:AB_10882823) (for TRCs expressing the taste-associated

G-protein gustducin),

• Rabbit anti-PLCβ2 (PhosphoSolutions, Denver, CO, USA, catalog# 1640-PLC;

RRID:AB_2801663), or guinea pig anti-PLCβ2 (PhosphoSolutions, AP8/24; RRID:

AB_2910247) (for identification of Type II TRCs).

• Rabbit anti P2X3 (Alamone, catalog #APR-016, RRID AB_2313760) (for labeling of

taste bud-specific innervation).

After washing in 0.1M PBS, the slides were incubated 2 hours with fluorescent secondary

antibodies, Alexa Fluor 488 donkey anti-mouse (Invitrogen, catalog# A21202;

RRID:AB_141607), Alexa Fluor 568 donkey anti-rabbit (Invitrogen, catalog# A10042), Alexa

Fluor 647 donkey anti-goat (Invitrogen, catalog# A21447; AB_2534017), Alexa Fluor 647

donkey anti-guinea pig (Jackson Immuno Research Laboratories, West Grove, PA, USA,

catalog# 706605148; RRID: AB_2340476) or Donkey anti-Rabbit IgG (H+L) Highly Cross-

Adsorbed Secondary Antibody, Alexa Fluor™ 647 (ThermoFisher Catalog # A-31573;

AB_2536183) at 1:800 dilution each at room temperature. Lastly, the slides were washed twice

for 10 minutes each in 0.1M PBS and then once for 10 minutes in 0.05M PBS before coverslip

slides with DAPI Fluoromount-G (SouthernBiotech, Birmingham, AL, USA, catalog# 010020).

RNA extraction

The papillae tissue samples were stored in RNA*later*™ at -20°C RNA was extracted with the

RNeasy fibrous tissue mini kit (Qiagen, Hilden, Germany). A small piece of the biopsy was taken

with a pipet tip into a 1.5 ml tube containing 300 ul of RLT buffer and mashed manually with a

micro pestle. The pestle was rinsed when pipetting 590 ul of RNase free water into the tube. The

kit protocol was followed, and a DNase treatment was performed with a DNA-free DNA removal

kit (Invitrogen) according to their protocol. The resulting RNA concentrations and quality were

checked with Nanodrop 8000 (Thermo Fisher Scientific) and Tapestation High Sensitivity RNA

tape (Agilent, Santa Clara, CA, USA).

Reverse transcription

High-Capacity cDNA Reverse Transcriptase Kit (Applied Biosystems, Waltham, MA, USA) was

used to obtain cDNA from the RNA samples. 100 ng of RNA per sample, according to

Tapestation concentrations, was taken into the reverse transcription reaction. The kit protocol was

followed. The reverse transcription was performed in duplicates to obtain enough product for all

the qPCR runs.

qPCR

Quantitative PCR (qPCR) was performed with TaqMan (Applied Biosystems) assays for *PLCβ2*

(Hs01080542), *NCAM1* (Hs00941821), and *TAS1R3* (Hs01026531) as well as for two reference

genes *RPLP0* (Hs99999902) and *GAPDH* (Hs02758991). TaqMan Gene Expression Master Mix

(Applied Biosystems) was used according to its protocol, and 10 ng of cDNA was added as

template. All samples were run in triplicate per gene, including a negative control. One gene per

plate was run in the QuantStudio 5 (Applied Biosystems). Average Ct and standard deviation

were calculated for each gene and then normalized (deltaCt=Ct(gene of interest) – Ct(reference

gene)) against the reference gene *GAPDH*. The reference gene *GAPDH* gave more consistent

normalization than RPLP0, and was therefore selected as a reference for the genes of interest.

SARS-CoV-2 test

A PCR test for the presence of SARS-CoV-2 virus in the tissue samples was performed in

triplicate, including two assays that target SARS-CoV-2 as well as negative control samples.

TaqMan microbe detection assay (Vi07918637) for SARS-CoV2N and TaqPath Blastopore

Microbial Detection Master Mix (Applied Biosystems) were used according to their protocol.

Positive control was TruMark Respiratory Panel 2.0 Amplification control (Applied Biosystems).

The qPCR was run in the QuantStudio 5 (Applied Biosystems). None of the biopsied tissue

samples tested showed the presence of SARS-CoV-2 virus, although one subject (#17) had an

active COVID infection.

Statistical analysis

Statistical calculations were carried out by Excel (Microsoft) or with calculators given at

www.socscistatistics.com. Both calculators gave identical values. A criterion value of $p < 0.05$

was set for statistical significance in all calculations. For comparisons of two groups, we used

either Estimation Statistics (www.estimationstats.com)(Ho, Tumkaya et al. 2019) or a Students

T-test; for comparison across samples, we utilized Pearson's Correlation co-efficient or Fisher's

Exact test for small sample sizes.

Means, standard deviations, and ranges were used to describe continuous variables such as age,

time from infection to testing, and taste scores. Categorical variables like sex and symptom

presence were summarized using counts and percentages.

Because taste ability varies substantially by sex and age, raw taste scores were normalized by sex

and age using reference data from Doty et al. (2025), which included 1,392 healthy individuals.

Scores below the 95% confidence interval (CI) were considered potentially abnormal, and those

below the 20th percentile were flagged as indicative of possible taste deficiency.

Chi-square tests (χ^2) were used to compare (categorical data) the frequency of abnormal taste

scores across groups. The results are reported as χ^2 (degree of freedom (df), sample size) = chi-

square value, p-value.

Pearson's Correlation (r) was used to assess linear relationships between molecular markers

(qPCR results), histological features, and taste scores. The results were reported as (df) sample

size=correlation co-efficient, p-value. (Schober, Boer et al. 2018).

RESULTS

Study population

All 28 participants (24 women and 4 men) reported normal taste before the first confirmed

COVID-19 infection (Table 1). The mean age was 49.9 years (SD+/-1.2) and ranged between 30

and 71 years. Most participants contracted COVID-19 in early 2020. Two participants (19 & 28)

experienced a second infection in 2022, shortly before undergoing taste testing and biopsy.

Considering only the first infection, the time from initial COVID-19 infection to taste testing

ranged from 13.5 months to 27.0 months (mean of 21.0 months).

Psychophysics:

Because taste ability varies substantially between individuals according to gender, age, and health

status, we normalized all raw taste scores against values given for 1,392 normal subjects (Doty,

Sharetts et al. 2025). Although all subjects, except one (subject #17), reported compromised taste

abilities in clinical reporting, only 5 of 28 ($\chi^2(1, N = 28) = 3.39, p = .065$) tested subjects scored

below the 20th percentile in overall taste ability (see Table 1). Nonetheless, 25 of the 28 subjects

showed significant deficiency in one or more individual taste qualities (values in red shown in

Table 1) compared to the 95% CI for age and gender of 1,392 normal subjects (Doty, Wylie et al.

2021). One subject, who was also anosmic, (#29) gave no correct responses to any tastant, even

at the highest concentration suggesting complete taste loss. Although individuals with a

profound overall deficit showed specific loss of one or more taste qualities (e.g. subject 7 shows

no response to sweet, bitter or umami), the pattern of loss was not duplicated by other subjects

(e.g. subject 24 with good responses to sweet and salty but total loss for bitter and umami).

Subject Number	Sex	Age	Covid Date	Taste Test Date	Biopsy Date	PLC Tastes Sweet	Umami	Bitter	Sour	Salty	No Taste	WETT score	Sex/Age-corrected %
1	F	47	Mar 2020	19/11-21		5	3	1	5	6	12	32	42
2	F	62	Apr 2020	19/11-21		3	0	1	8	8	11	31	36
3	F	50	Apr 2020	24/11-21		6	0	3	7	5	12	33	45
4	F	49	May 2020	3/12/21		0	8	6	6	7	10	37	59
5	F	53	Nov 2020	17/1-22		8	4	8	8	7	13	48	95
6	F	41	Nov 2020	18/1-21		7	2	4	5	4	12	34	51
10	F	66	Nov 2020	20/1-22		8	1	8	7	5	13	42	83
16	F	59	Apr 2020	11/2/22		5	3	6	8	7	10	39	70
7	F	68	Nov 2020	18/1-22	23/3-22	0	0	0	6	5	13	24	16
8	F	64	Mar 2020	18/1-22	23/3-22	6	0	0	6	7	13	32	45
9	M	49	Apr 2020	19/1-22	17/3-22	1	0	0	8	8	13	30	54
12	F	56	Apr 2020	21/1-22	24/1-22	3	3	6	7	4	12	35	54
13	M	45	Apr 2020	21/1-22	21/1-22	0	1	0	5	3	13	22	21
14	F	30	Apr 2020	21/1-22	21/1-22	3	8	5	8	6	10	40	70
15	F	53	Apr 2020	24/1-22	24/1-22	4	6	3	4	4	13	34	54
17	F	30	Apr 2020	31/1-22	31/1-22	8	8	8	8	8	13	53	99
18	F	63	June 2020	23/3-22	23/3-22	7	1	4	6	1	13	32	45
19	F	46	May 2020 & Jan 2022	24/3-22	24/3-22	7	4	8	6	2	9	36	59
20	F	38	Feb 2021	30/3-22	20/4-22	7	2	7	8	8	13	45	83
21	F	71	Jan 2021	17/1-22	11/5-22	4	1	5	7	4	13	34	70
22	F	56	Apr 2020	11/5-22	11/5-22	1	4	0	6	2	13	26	21
23	F	49	Nov 2020	12/5-22	12/5-22	4	4	7	6	5	13	39	67
24	M	42	June 2020	13/5-22	13/5-22	5	0	0	4	4	4	17	7
25	F	50	Mar 2020	13/5-22	13/5-22	2	4	1	0	5	7	19	4
26	F	24	June 2020	21/1-22	16/5-22	6	0	6	5	7	13	37	44
27	F	52	Mar 2020	15/6-22	15/6-22	0	0	4	2	6	13	25	15
28	F	43	Nov 2020 & Feb 2022	5/7-22	5/7-22	6	3	7	7	3	10	36	44
29	M	30	April 2020	15/7-22	15/7-22	0	0	0	0	0	13	13	1
Total	N = 28		red text <95% C.I.		Ave Taste Score	4.1	2.5	3.9	5.8	5.0	11.7	33.0	48.4
Biopsy	N = 20				zero scores	5	9	6	2	1			

Table 1: Results from WETT testing of all subjects including basic demographic information, dates of infection, taste testing and biopsy. Columns labeled as "Sweet, Umami, Bitter, Sour, Salty and No Taste" show WETT test raw scores for each taste quality and blank "No Taste" control strips. Red numerals indicate values below the 95% confidence interval for sex and age-matched controls based on Doty, et al 2025. The incidence of values below the 95% is significantly higher than expected based on the normal population ($\chi^2(1, N = 140) = 48.63$, $p < .0001$). Zero values indicate that the subject was unable to identify the tastant at any concentration offered. The total number of zero values for each tastant is shown in red along the bottom row. WETT score shows the total WETT score for each subject alongside of the corresponding sex and age match % ranking compared to the study population in Doty et al 2025. A perfect score is 53 (8 for each taste quality + 13 blanks). Complete ageusia would give a score of 13 (0 for all tastants and 13 for correctly identifying blank controls). PLCß2-dependent tastes are indicated by orange shading along the top row.

Quantitative PCR Findings

RNA prepared from taste

papillae from all biopsied

subjects was analyzed for

expression levels of mRNAs

encoding phospholipase C beta 2

(PLCß2), a marker of Type II

cells; and Taste 1 receptor

member 3 (*TAS1R3)* mRNA,

coding for a receptor component

for sweet and umami tastes; and

	taste test	deltaCt			Ave PLC +
Subject #	value	PLCb2	TAS1R3	NCAM1	Tas1R3
7	24	11.78	11.24	11.58	11.51
8	32	9.59	9.59	12.25	9.59
9	30	10.78	9.26	12.85	10.02
12	35	9.36	8.98	11.29	9.17
13	22	10.72	8.38	11.5	9.55
14	40	9.92	7.34	11.25	8.63
15	34	9.83	8.4	12.1	9.12
17	53	9	8.02	11.52	8.51
18	32	11.87	9.75	14.29	10.81
19	36	11.61	8.42	15.14	10.02
20	45	8.29	8.34	11.73	8.32
21	34	10.89	10.13	13.06	10.51
22	26	10.48	9.84	12.47	10.16
23	39	9.64	7.83	10.55	8.74
24	17	9.7	7.81	11.11	8.76
25	19	11.39	10.6	12.46	11.00
26	37	10.05	8.77	12.33	9.41
27	25	8.89	8.4	10.81	8.65
28	36	10.17	8.9	13.16	9.54
29	13	11.2	10.72	12.27	10.96

Table 2: DeltaCT values from qPCR of fungiform papillae. Taste test values as shown in Table 1 are included for comparison.

neural adhesion molecule 1 (*NCAM1)*, a marker of total innervation and Type III TCs. All data

were normalized against glyceraldehyde-3-phosphate hydrogenase (*GAPDH)* mRNA, yielding

deltaCt given in Table

2 along with Taste

Scores from each

subject for

comparison.

The deltaCT values for

PLCß2 and *TAS1R3*,

both markers of Type

Figure 1: A) The mRNA levels for PLCß2 (r(18)=.48, p= .03) and TAS1R3(r(18)=.49, p= .03) correlate with the overall taste ability of each subject. B) The mRNA level for TAS1R3 correlates with the combined taste score for the related taste qualities (Sweet and brothy) (r(18)=.51, p= .02).

Alt Text: Graphs showing the relationship between WETT scores and qPCR data. At left, showing the correlation between expression of TAS1R3 or PLC and overall taste score. At right showing correlation between TAS1R3 and average combined score for sweet and umami.

II TCs, were highly correlated (r(18) = 0.64, p=0.002). Similarly, the deltaCT value for *PLCß2*

was also correlated with the expression levels of *NCAM*, a marker of Type III cells of taste buds

and total innervation (r(18) = 0.63, p=0.003) although no significant correlation exists for values

of *TAS1R3* compared to *NCAM* (r(18) = 0.32, p = 0.17).

Comparing the qPCR data to the taste scores from the same subjects shows significant

correlations for both *PLCß2* (r(18) = 0.48, p= 0.03) and *TAS1R3* (r(18) =-0.49, p= 0.03) but not

for *NCAM* (r(18) = 0.02, p>0.05). This relationship is shown graphically in Fig. 1. Since both

sweet and umami require the TAS1R3 receptor subunit, we compared the expression level of this

subunit to the average of taste scores for sweet and umami for each subject. A significant

correlation exists between the deltaCT values for *TAS1R3* and the average of the related taste

qualities (sweet and umami) (r(18)= 0.52, p=0.02), but is not significant compared to each of the

individual tastes.

370 Histological Findings

Fungiform samples from the subjects varied

between 4 and 8 papillae, with a median of 7.

Approximately half of the papillae had no

identifiable TBs, as reported for human

fungiform papillae in previous literature e.g.

(Miller 1988). Taste buds were identified in

all subjects, including subject 29, who

reported no taste sensibility. While relatively

Subject #	taste score	# papillae	# TBs	ave tb/pap	plc/tb	plc/pap
7	24	7	42	6.0	2.4	14.3
8	32	8	42	5.3	2.0	10.5
9	30	8	10	1.3	2.0	2.5
12	35	6	30	5.0	2.1	10.3
13	22	4	6	1.5	1.8	2.7
14	41	5	61	12.2	2.9	35.4
16	34	6	13	2.2	2.6	5.7
17	53	8	25	3.1	2.7	8.5
18	32	7	7	1.0	1.8	1.8
19	36	6	8	1.3	2.0	2.7
20	45	7	41	5.9	2.0	11.7
21	34	7	28	4.0	3.3	13.2
22	26	8	32	4.0	1.8	7.2
23	39	7	17	2.4	2.3	5.6
24	17	6	22	3.7	2.0	7.3
25	19	6	2	0.3	1.5	0.5
26	37	5	22	4.4	2.4	10.6
27	25	7	32	4.6	2.8	12.8
28	26	8	17	2.1	2.5	5.3
29	13	7	27	3.9	1.9	7.3

Table 3: Counts of papillae sampled, number of taste buds and average PLCß2+ cells per taste bud and papilla in each case. No significant correlation was found between taste score and average number of taste buds per papilla nor of density of PLCß2+ cells per taste bud or papilla. The histological counts and density of PLCß2+ cells also did not correlate with any PCR values.

normal-appearing TBs occur in samples of all subjects (See Fig. 2), many subjects displayed

disorganized TBs or relatively isolated PLCß2+ cells not associated with a cluster of cells as

would occur in a normal TB (High and Finger 2025). The average number of taste bud profiles

per papilla was 3.6 (+/-2.6 s.d.). Because of the significant variance in the number of TB profiles

observed in a single section through a papilla, we calculated the average number of TB profiles

per papilla from all papillae to measure the taste bud population for each subject (Table 3). In

addition, we counted the number of PLCß2+ TCs within each section to calculate the average

proportion of Type II (PLCß2+) cells per TB.

The taste score for each subject was not correlated with either the average number of TB

profiles per papillae ($r(18) = 0.268$, $p = 0.25$) nor with the average number of PLCß2+ cells per

papilla ($r(18)$ -0.35, $p= 0.15$) nor with the average number of PLCß2+ cells per TB ($r(18) =$

0.427, $p = .06$). Abnormal TBs or irregular PLCß2+ cells were especially noteworthy in subjects

12 (Taste Score 54%), 13(21%), 22 (21%), 24 (8 %), 25 (4%), 28 (44%) and 29 (1%). Fig. 3

shows examples of such profiles.

Figure 2: Appearance of normal taste buds sectioned in different planes from different subjects. Red = PLCß2 immunoreactivity showing Type II taste cells; Green = TUJ1 immunoreactivity for nerve fibers. Typical taste buds have more than one PLCß2+ cell surrounded by a dense network of TUJ1-stained nerve fibers. Green profiles outside of taste buds are extragemmal nerve fibers typical of lingual epithelium (Moayedi, Michlig et al. 2021). **a, b.** Taste buds sectioned longitudinally appear similar in shape to garlic bulbs or onions (from Subject (SWE) 15. **c)** Taste buds sectioned perpendicular to their long axis appear circular (from subject (SWE) 29 who reported a total absence of taste perception (taste score of 13, showing no correct responses to any tastant).
Scale bar in panel (b) applies to all panels.

Alt Text: Images of 3 micrographs showing the normal appearance of taste buds in our samples. Multiple PLC-positive taste cells within each bud are embraced by a dense nerve plexus.

Figure 3: Examples of abnormal PLCß2+ taste-like cells from 4 cases. (A & B). Subject (SWE) 12 had a taste score of 35 (54%). (C) From subject (SWE) 13 who had a score of 22 (21%). In both cases (A & C), innervated PLCß2+ (red arrows) cells occur in relative isolation within the epithelium. Innervation of these cells (green: TUJ1) is sparse unlike innervation of taste buds (cf Fig. 2). These cells may represent very small taste buds or simply PLCß2+ cells lying outside of a taste bud structure. Panel B shows a higher magnification of one such isolated PLCß2+ cell and its innervation stained for the presence of P2X3, a neural receptor predominantly on nerve fibers that innervate taste buds. Panels D-E are from case (SWE) 24 with a taste score of 17 (4%) showing examples of aberrant PLCß2+ cells (red arrows) outside of a taste bud structure. Panel F: A disorganized taste bud from case SWE 29 who reported a total absence of taste perception (taste score of 13, showing no correct responses to any tastant). The heavy innervation is typical of taste buds, but the cellular structure revealed by PLC-staining (red) does not show typical taste bud organization.

Alt Text: Images showing abnormal isolated PLC-positive cells and abnormal taste buds within the epithelium of our subjects.

DISCUSSION

Morbidity Changes in COVID-19 Variants

Previous studies found that the frequency of chemosensory symptoms, including taste loss, varies

across SARS-CoV-2 variants. Early variants, such as Wild type and Alpha, were more commonly

associated with taste and smell loss, while later emerging variants, such as Omicron, show

markedly lower prevalence of taste loss (Reiter, Coelho et al. 2023). Most subjects included in

our study were primarily infected during the first wave of the pandemic, and their reported taste

disturbances fit well with this pattern of greater taste loss early in the pandemic.

Published epidemiological data further illustrate this trend. Among individuals infected with the

Wild-type strain, 241 out of 1,141 cases (21%) reported smell and/or taste disorders. For Alpha,

223 out of 1265 cases (18%), for Delta, 480 out of 1516 cases (32%), and for Omicron, only 317

out of 7431 cases (4%) presented with smell and/or taste disorders (Ota, Yumiya et al. 2025).

This supports the hypothesis that the variant-dependent tropism and immune response may

influence the degree and duration of taste dysfunction.

On the role of the ACE2 receptor

Relatively early in the pandemic, it was suggested that SARS-CoV-2 infects oral tissue (Huang,

Pérez et al. 2021). Doyle et al. (Doyle, Appleton et al. 2021) reported that ACE2 was expressed

by Type II TRCs, and further demonstrated the presence of SARS-CoV-2 in Type II TRCs,

which mediate sweet, umami and bitter sensations, but not within the other taste cell types. This

likely explains why their subjects in that study did not taste the "sweetness of chocolate and bitter

coffee" and instead reported that it "tasted like mud" (Doyle, Appleton et al. 2021).

Research during the COVID-19 pandemic has highlighted the potential impact of SARS-CoV-2

on TBs, particularly through its interaction with ACE2 receptors (Huang, Perez et al. 2021, Yao,

Doyle et al. 2023). In patients experiencing taste disturbances following COVID-19 infection,

SARS-CoV-2 was detected in basal and suprabasal layers of fungiform papillae for extended

periods. In one subject, virus was detected up to 63 weeks after the initial diagnosis, coincident

with disrupted nerve fibers and altered taste bud architecture (Yao, Doyle et al. 2023).

Recovery of taste, and re-establishment of normal epithelial cytoarchitecture, coincided with the

disappearance of the virus from the taste epithelium (Yao, Doyle et al. 2023). These findings

suggest that prolonged viral presence and the associated immune response may impair the

renewal and functionality of taste buds in Long COVID. In our subjects, a sensitive commercial

qPCR test showed that none of the tissue samples were positive for the SARS-CoV-2 viral RNA

suggesting that taste dysfunction may persist even after the virus is no longer detectable in the

peripheral taste system. This offers the potential that persistent taste dysfunction in Long Covid

may outlast the presence of the virus itself.

Why loss of sweet-bitter-umami tasting ability is more frequent than the loss of other taste

qualities

Psychophysical results from this study indicate a significant disruption of taste with a tendency

toward preferential loss of sweet, bitter, or umami tastes. A previous WETT study involving 340

long-term (1-year) COVID-19 subjects and 434 non-COVID controls (Sharetts, Moein et al.

2024) found no statistically significant differences in taste function between groups. In contrast,

we note significant taste loss but our study specifically targeted subjects reporting long-term

COVID-19-associated taste loss, preventing direct comparison with the broader population

surveyed in the Sharetts et al. study.

Taste dysfunction may result from disruption of taste-related intracellular transduction cascades

or damage to taste buds (TBs), their receptor cells (TRCs) or the gustatory nerves.   Sweet,

umami and bitter tastes all involve Type II receptor cells relying on a PLCβ2-mediated

transduction cascade (Zhang, Hoon et al. 2003, Zhao, Zhang et al. 2003). Sweet taste is mediated

by TAS1R2/TAS1R3 receptor heterodimers, umami by TAS1R1/TAS1R3, and bitter by a

distinct family of ~30 Tas2Rs. In contrast, ionic taste qualities, such as sour and salty activate

Type III TRCs relying on ion channels in the apical membrane to initiate a response (Liman

2020, Liman and Kinnamon 2021).

The fact that the COVID-19 virus directly infects Type II TRCs may explain why deficits in

sweet, umami and bitter are more frequent than for sour and salt. In our study, however, viral

RNA was not detectable in taste tissues and there was no evidence of wholesale destruction or

abnormality in Type II, PLCβ2+, taste cells. Nonetheless, in our subjects overall tasting ability

correlates positively with the amount of PLCβ2 and TAS1R3 present in the tissue. This leaves

open the question of how prior viral infection might induce long-lasting deficits in taste.

In the olfactory epithelium, the virus infects supporting cells, but not the receptor cells (Chen,

Shen et al. 2020). Nonetheless, the olfactory receptor cells respond to the viral infection of the

tissue by downregulating expression of olfactory receptors and downstream signaling molecules

while upregulating expression of viral response proteins (Chen, Reed et al. 2019, Zazhytska,

Kodra et al. 2022). The transduction cascade is re-established in the receptor cells once the viral

infection has been cleared (Zazhytska, Kodra et al. 2022). While a similar phenomenon may

occur in taste buds during viral infection, it is unlikely such a reaction would persist in the tissue

after the virus is cleared and all the taste cells have been replaced during the course of normal

taste cell turnover which requires a few months at most (Perea-Martinez, Nagai et al. 2013). At

the time of our testing, up to 2 years after initial infection, none of the taste cells present at that

time would have been present at the time of viral infection.

Furthermore, we do note some loss of non PLCβ2 taste qualities, i.e. sour and salty, in some

subjects. This loss suggests that non-PLC dependent mechanisms may also be at work underlying

taste dysfunction in Long COVID. One possibility is that the infection alters the function in taste

nerves. Indeed, evidence exists for widespread neural dysfunction in Long COVID (Daia, Scheau

et al. 2021, McFarland, Yousuf et al. 2021, Otte, Bork et al. 2022, O'Neill, De Stefano et al.

2023, Rogn, Jensen et al. 2024). If so, then some of the observed taste loss might be attributable

to loss of transmission in the taste nerves rather than disruption of the taste epithelium.

Other factors that may have influenced the disease process of COVID-2 in taste

Recent studies (Finlay, Brann et al. 2022, Peluso, Ryder et al. 2023, Peluso, Hanson et al. 2024)

broaden the perspective by suggesting that immune dysregulation and T cell dysfunction may

play significant roles in the long-lasting effects of COVID-19. Whole-body positron emission

tomography imaging with a tracer that selectively tags activated T cells show that subjects with

Long COVID had specific tissues enriched for activated T cells compared to individuals without

this condition (Peluso, Ryder et al. 2024). All five subjects in that study showed the presence of

intracellular SARS-CoV-2 spike protein-encoding RNA in lamina propria tissue up to 676 days

after initial COVID-19. This offers evidence of viral persistence which could underlie long-term

immunologic perturbations for up to two years postinfection. Whether a similar prolonged

immunological response occurs in taste tissue is unknown although Yao et al (Yao, Doyle et al.

2023) report long term (>6 months) inflammatory response when virus remains in the tissue. We

did not, however, detect any virus by PCR in our tissue samples from patients more than one year

post infection.

On the relation between histological changes of TB by COVID-19 and taste changes

A particularly intriguing aspect of our study is that all subjects, including those with severe

deficits, displayed structurally intact taste buds despite the loss of taste function. However, some

dystrophic taste buds and isolated PLCß2-expressing cells were observed in many subjects. In

mice, disruption of molecular signals between nerves and taste buds can result in formation of

aberrant PLCß2-expressing cells outside of the taste buds themselves (Castillo, Seidel et al.

2014). This finding raises important questions about the disconnect between histological integrity

and functional impairment. One possible explanation is that taste bud dystrophy, rather than

outright loss, may be sufficient to alter taste sensitivity by impairing cellular turnover or

disrupting synaptic communication with gustatory neurons (Wang, Liu et al. 2023). While intact

taste buds might suggest some functional resilience, the significant decrease in PLCß2-dependent

signaling suggests that molecular-level dysregulation may underlie the persistent taste

dysfunction. In addition, the discrepancy between PLCß2 expression and taste bud morphology,

especially in Subject 29 who shows complete taste loss, suggests dysfunction in peripheral nerves

or at higher levels of the neuraxis.

Clinical relevance

Understanding the mechanisms behind taste loss is clinically relevant, as chemosensory loss

impacts nutritional intake, emotional well-being, and quality of life. A multidisciplinary mindset

is crucial for appropriate diagnostics and for providing patients with appropriate treatment and

underscore the importance of developing targeted therapies to alleviate the burden of long-term

taste dysfunction in COVID-19 survivors.

Here we present an integrative approach combining psychophysical taste testing, RNA

expression analyses using markers for TRC type II and III, and histological evaluation of taste

tissue in subjects with Long COVID. This multidimensional analysis provides a more

comprehensive understanding of Long COVID taste dysfunction. While this study focuses on the

loss of Type II TRC, PLCß2-dependent taste qualities (sweet, umami, bitter), the loss of other

taste qualities in a limited number of subjects suggests that multiple mechanisms may underlie

the sensory loss.

Conclusions

Our study suggests that persistent taste loss following COVID-19 depends neither on ongoing

infection nor on wholesale loss of taste buds or their innervation. We found a correlation between

taste loss for specific qualities, i.e. sweet, umami and bitter, and reduced expression of molecular

elements of the corresponding taste transduction cascade. This highlights the importance of

cellular signaling over structural integrity in COVID-related taste dysfunction.

535
536

537 **ACKNOWLEDGEMENTS**
538 This work was supported by grants 3R01DC014728-05S1 to TEF awarded by the National
539 Institute on Deafness and Other Communication Disorders of the National Institutes of Health,
540 by Åke Wiberg Stiftelsen grant 2023 to M.K. and by internal funding of the Department of
541 Animal Biosciences, Swedish University of Agricultural Sciences.
542

AUTHOR CONTRIBUTIONS

G.H., G.L., and T.E.F. conceived and designed this research. G.H. & G.L. wrote the ethical
proposal to Etikprövningsmyndigheten. H.M. provided subjects, performed the WETT tests and
took the biopsies. M.A.K. provided subjects and with A.A. assisted in analysis of the subject
population. T.E.F. and M.L. designed and performed the histopathology and data analysis. G.A.
and T.V. designed and analyzed the quantitative PCR experiments. T.V. performed RNA
extraction and quantitative PCR experiments. T.E.F. performed statistical analysis of the
quantitative data and compared results to those in the Doty et al (2015) report. G.L., G.H., G.A.,
H.M., and T.E.F. provided funding for the project. G.H., T.E.F., G.A., and M.A.K. prepared the
manuscript and all authors have discussed and revised the manuscript.

Use of AI:

This study utilized Microsoft Copilot (GPT-4) accessed through a secure university-managed
Microsoft Teams environment. The tool was utilized in the process of editing and checking
calculations but was not used to generate any original content. All outputs were reviewed by the
lead author.

Competing interests
The authors declare no competing financial interests.

Data Availability Statement: The data underlying this article will be shared on reasonable
request to the corresponding author.

REFERENCES
565

Bartel, D. L., S. L. Sullivan, E. G. Lavoie, J. Sevigny and T. E. Finger (2006). "Nucleoside
triphosphate diphosphohydrolase-2 is the ecto-ATPase of type I cells in taste buds." J Comp
Neurol **497**(1): 1-12.

Boldes, T., A. Ritter, E. Soudry, D. Diker, E. Reifen and E. Yosefof (2024). "The long-term
effect of COVID-19 infection on olfaction and taste; a prospective analysis." Eur Arch
Otorhinolaryngol **281**(11): 6001-6007.

Castillo, D., K. Seidel, E. Salcedo, C. Ahn, F. J. de Sauvage, O. D. Klein and L. A. Barlow
(2014). "Induction of ectopic taste buds by SHH reveals the competency and plasticity of adult
lingual epithelium." Development **141**(15): 2993-3002.

Chang, R. B., H. Waters and E. R. Liman (2010). "A proton current drives action potentials in
genetically identified sour taste cells." Proc Natl Acad Sci U S A **107**(51): 22320-22325.

Chen, M., R. R. Reed and A. P. Lane (2019). "Chronic Inflammation Directs an Olfactory Stem
Cell Functional Switch from Neuroregeneration to Immune Defense." Cell Stem Cell **25**(4): 501-
513.e505.

Chen, M., W. Shen, N. R. Rowan, H. Kulaga, A. Hillel, M. Ramanathan, Jr. and A. P. Lane
(2020). "Elevated ACE-2 expression in the olfactory neuroepithelium: implications for anosmia
and upper respiratory SARS-CoV-2 entry and replication." Eur Respir J **56**(3).

Cheng, L. H. H. and P. P. Robinson (1991). "The distribution of fungiform papillae and taste
buds on the human tongue." Archives of Oral Biology **36**(8): 583-589.

Cooper, K. W., D. H. Brann, M. C. Farruggia, S. Bhutani, R. Pellegrino, T. Tsukahara, C.
Weinreb, P. V. Joseph, E. D. Larson, V. Parma, M. W. Albers, L. A. Barlow, S. R. Datta and A.
Di Pizio (2020). "COVID-19 and the Chemical Senses: Supporting Players Take Center Stage."
Neuron **107**(2): 219-233.

Daia, C., C. Scheau, G. Neagu, I. Andone, A. Spanu, C. Popescu, S. I. Stoica, M. C. Verenca and
G. Onose (2021). "Nerve conduction study and electromyography findings in patients recovering
from Covid-19 - Case report." Int J Infect Dis **103**: 420-422.

Doty, R. L., R. Sharetts and S. T. Moein (2025). "Self-administered Taste Testing without Water:
Normative data for the 53-item Waterless Empirical Taste Test (WETT)." Int Forum Allergy
Rhinol. **in press**.

Doty, R. L., C. Wylie and M. Potter (2021). "Validation of the Waterless Empirical Taste Test
(WETT((R)))." Behav Res Methods **53**(2): 864-873.

Doyle, M. E., A. Appleton, Q. R. Liu, Q. Yao, C. H. Mazucanti and J. M. Egan (2021). "Human
Taste Cells Express ACE2: a Portal for SARS-CoV-2 Infection." bioRxiv.

Doyle, M. E., A. Appleton, Q. R. Liu, Q. Yao, C. H. Mazucanti and J. M. Egan (2021). "Human
Type II Taste Cells Express Angiotensin-Converting Enzyme 2 and Are Infected by Severe Acute
Respiratory Syndrome Coronavirus 2 (SARS-CoV-2)." Am J Pathol **191**(9): 1511-1519.

Ely, E. W., L. M. Brown, H. V. Fineberg, E. National Academies of Sciences and C. Medicine
Committee on Examining the Working Definition for Long (2024). "Long Covid Defined." N
Engl J Med **391**(18): 1746-1753.

Farbman, A. I. (1965). "Fine structure of the taste bud." Journal of Ultrastructure Research **12**:
328-350.

Farbman, A. I. (1980). "Renewal of taste bud cells in rat circumvallate papillae." Cell & Tissue
Kinetics **13**(4): 349-357.

Farbman, A. I., N. Guagliardo, S. I. Sollars and D. L. Hill (2004). "Each sensory nerve arising
from the geniculate ganglion expresses a unique fingerprint of neurotrophin and neurotrophin
receptor genes." J Neurosci Res **78**(5): 659-667.

Finlay, J. B., D. H. Brann, R. Abi Hachem, D. W. Jang, A. D. Oliva, T. Ko, R. Gupta, S. A.
Wellford, E. A. Moseman, S. S. Jang, C. H. Yan, H. Matsunami, T. Tsukahara, S. R. Datta and B.
J. Goldstein (2022). "Persistent post-COVID-19 smell loss is associated with immune cell
infiltration and altered gene expression in olfactory epithelium." Sci Transl Med **14**(676):
eadd0484.

Hannum, M. E., R. J. Koch, V. A. Ramirez, S. S. Marks, A. K. Toskala, R. D. Herriman, C. Lin,
P. V. Joseph and D. R. Reed (2023). "Taste loss as a distinct symptom of COVID-19: a
systematic review and meta-analysis." Chem Senses **48**.

Hellekant, G. (2024). "Neuroscience of taste: unlocking the human taste code." BMC Neurosci
25(1): 19.

Hellekant, G., Y. Kasahara, A. I. Farbman, S. Harada and C. Hård af Segerstad (1987).
"Regeneration ability of fungiform papillae and taste-buds in rats." Chemical Senses **12**(3): 459-
465.

High, B. and T. E. Finger (2025). "Structural comparisons of human and mouse fungiform taste
buds." Chem Senses **50**.

Ho, J., T. Tumkaya, S. Aryal, H. Choi and A. Claridge-Chang (2019). "Moving beyond P values:
data analysis with estimation graphics." Nat Methods **16**(7): 565-566.

Huang, N., P. Perez, T. Kato, Y. Mikami, K. Okuda, R. C. Gilmore, C. D. Conde, B. Gasmi, S.
Stein, M. Beach, E. Pelayo, J. O. Maldonado, B. A. Lafont, S. I. Jang, N. Nasir, R. J. Padilla, V.
A. Murrah, R. Maile, W. Lovell, S. M. Wallet, N. M. Bowman, S. L. Meinig, M. C. Wolfgang, S.
N. Choudhury, M. Novotny, B. D. Aevermann, R. H. Scheuermann, G. Cannon, C. W. Anderson,
R. E. Lee, J. T. Marchesan, M. Bush, M. Freire, A. J. Kimple, D. L. Herr, J. Rabin, A. Grazioli,
S. Das, B. N. French, T. Pranzatelli, J. A. Chiorini, D. E. Kleiner, S. Pittaluga, S. M. Hewitt, P.
D. Burbelo, D. Chertow, N. C.-A. Consortium, H. C. A. Oral, N. Craniofacial Biological, K.

Frank, J. Lee, R. C. Boucher, S. A. Teichmann, B. M. Warner and K. M. Byrd (2021). "SARS-
CoV-2 infection of the oral cavity and saliva." Nat Med **27**(5): 892-903.

Huang, N., P. Pérez, T. Kato, Y. Mikami, K. Okuda, R. C. Gilmore, C. D. Conde, B. Gasmi, S.
Stein, M. Beach, E. Pelayo, J. O. Maldonado, B. A. Lafont, S. I. Jang, N. Nasir, R. J. Padilla, V.
A. Murrah, R. Maile, W. Lovell, S. M. Wallet, N. M. Bowman, S. L. Meinig, M. C. Wolfgang, S.
N. Choudhury, M. Novotny, B. D. Aevermann, R. H. Scheuermann, G. Cannon, C. W. Anderson,
R. E. Lee, J. T. Marchesan, M. Bush, M. Freire, A. J. Kimple, D. L. Herr, J. Rabin, A. Grazioli,
S. Das, B. N. French, T. Pranzatelli, J. A. Chiorini, D. E. Kleiner, S. Pittaluga, S. M. Hewitt, P.
D. Burbelo, D. Chertow, K. Frank, J. Lee, R. C. Boucher, S. A. Teichmann, B. M. Warner and K.
M. Byrd (2021). "SARS-CoV-2 infection of the oral cavity and saliva." Nat Med **27**(5): 892-903.

Janols, H., C. Wadsten, C. Forssell, E. Raffeti, C. Janson, X. Zhou and M. A. Kisiel (2024).
"Enhancing EQ-5D-5L Sensitivity in Capturing the Most Common Symptoms in Post-COVID-19
Patients: An Exploratory Cross-Sectional Study with a Focus on Fatigue, Memory/Concentration
Problems and Dyspnea Dimensions." Int J Environ Res Public Health **21**(5).

Kisiel, M. A., H. Janols, T. Nordqvist, J. Bergquist, S. Hagfeldt, A. Malinovschi and M.
Svartengren (2022). "Predictors of post-COVID-19 and the impact of persistent symptoms in
non-hospitalized patients 12 months after COVID-19, with a focus on work ability." Ups J Med
Sci **127**.

Kisiel, M. A., S. Lee, H. Janols and A. Faramarzi (2023). "Absenteeism Costs Due to COVID-19
and Their Predictors in Non-Hospitalized Patients in Sweden: A Poisson Regression Analysis."
Int J Environ Res Public Health **20**(22).

Kisiel, M. A., S. Lee, S. Malmquist, O. Rykatkin, S. Holgert, H. Janols, C. Janson and X. Zhou
(2023). "Clustering Analysis Identified Three Long COVID Phenotypes and Their Association
with General Health Status and Working Ability." J Clin Med **12**(11).

Kisiel, M. A., T. Nordqvist, G. Westman, M. Svartengren, A. Malinovschi and H. Janols (2021).
"Patterns and predictors of sick leave among Swedish non-hospitalized healthcare and residential
care workers with Covid-19 during the early phase of the pandemic." PLoS One **16**(12):
e0260652.

Lee, H., L. J. Macpherson, C. A. Parada, C. S. Zuker and N. J. P. Ryba (2017). "Rewiring the
taste system." Nature **548**(7667): 330-333.

Lee, S., M. A. Kisiel, P. Lindberg, A. M. Wheelock, A. Olofsson, J. Eriksson, J. Bruchfeld, M.
Runold, L. Wahlstrom, A. Malinovschi, C. Janson, C. Wachtler and A. C. Carlsson (2025).
"Using machine learning involving diagnoses and medications as a risk prediction tool for post-
acute sequelae of COVID-19 (PASC) in primary care." BMC Med **23**(1): 251.

Liman, E. R. (2020). "Salty Taste: From Transduction to Transmitter Release, Hold the
Calcium." Neuron **106**(5): 709-711.

Liman, E. R. and S. C. Kinnamon (2021). "Sour taste: receptors, cells and circuits." Curr Opin
Physiol **20**: 8-15.

Lindemann, B. (1996). "Taste reception." <u>Physiol Rev</u> **76**(3): 719-766.

Matsumoto, I., M. Ohmoto, M. Narukawa, Y. Yoshihara and K. Abe (2011). "Skn-1a (Pou2f3)
specifies taste receptor cell lineage." <u>Nat Neurosci</u> **14**(6): 685-687.

McFarland, A. J., M. S. Yousuf, S. Shiers and T. J. Price (2021). "Neurobiology of SARS-CoV-2
interactions with the peripheral nervous system: implications for COVID-19 and pain." <u>Pain Rep</u>
6(1): e885.

Miller, I. J. (1986). "Variation in human fungiform taste bud densities among regions and
subjects." <u>Anatomical Record</u> **216**(4): 474-482.

Miller, I. J., Jr. (1988). "Human taste bud density across adult age groups." <u>J Gerontol</u> **43**(1):
B26-30.

Miller, I. J., Jr. (1989). "Variation in human taste bud density as a function of age." <u>Ann N Y
Acad Sci</u> **561**: 307-319.

Miller, I. J., Jr. and F. E. Reedy, Jr. (1990). "Variations in human taste bud density and taste
intensity perception." <u>Physiol. Behav.</u> **47**(6): 1213-1219.

Moayedi, Y., S. Michlig, M. Park, A. Koch and E. A. Lumpkin (2021). "Somatosensory
innervation of healthy human oral tissues." <u>J Comp Neurol</u> **529**(11): 3046-3061.

O'Neill, F., G. De Stefano, M. Pridgeon, D. Bhargava, A. Marshall, A. Marshall and B. Frank
(2023). "Trigeminal neuropathy presenting secondary to SARS-CoV-2 infection." <u>Pain Rep</u> **8**(6):
e1103.

Ogata, T. and Y. Ohtubo (2020). "Quantitative Analysis of Taste Bud Cell Numbers in the
Circumvallate and Foliate Taste Buds of Mice." <u>Chem Senses</u> **45**(4): 261-273.

Ohtubo, Y. and K. Yoshii (2011). "Quantitative analysis of taste bud cell numbers in fungiform
and soft palate taste buds of mice." <u>Brain Res</u> **1367**(0): 13-21.

Ota, Y., Y. Yumiya, O. Chimed-Ochir, A. Hasegawa, T. Yoshida, T. Nagata, J. Tanaka, H. Ohge,
M. Kuwabara and T. Kubo (2025). "Characteristics of patients with COVID-19 and smell and/or
taste disorders depending on different virus strains: a cross-sectional study in Hiroshima, Japan."
<u>BMJ Open</u> **15**(2): e088377.

Otte, M. S., M. L. Bork, P. H. Zimmermann, J. P. Klußmann and J. C. Lüers (2022). "Patients
with COVID-19-associated olfactory impairment also show impaired trigeminal function." <u>Auris
Nasus Larynx</u> **49**(1): 147-151.

Peluso, M. J., M. R. Hanson and S. G. Deeks (2024). "Infection-associated chronic conditions:
Why Long Covid is our best chance to untangle Osler's web." <u>Sci Transl Med</u> **16**(773): eado2101.

Peluso, M. J., D. Ryder, R. Flavell, Y. Wang, J. Levi, B. H. LaFranchi, T. M. Deveau, A. M.
Buck, S. E. Munter, K. A. Asare, M. Aslam, W. Koch, G. Szabo, R. Hoh, M. Deswal, A.

Rodriguez, M. Buitrago, V. Tai, U. Shrestha, S. Lu, S. A. Goldberg, T. Dalhuisen, M. S. Durstenfeld, P. Y. Hsue, J. D. Kelly, N. Kumar, J. N. Martin, A. Gambir, M. Somsouk, Y. Seo, S. G. Deeks, Z. G. Laszik, H. F. VanBrocklin and T. J. Henrich (2023). "Multimodal Molecular Imaging Reveals Tissue-Based T Cell Activation and Viral RNA Persistence for Up to 2 Years Following COVID-19." <u>medRxiv</u>.

Peluso, M. J., D. Ryder, R. R. Flavell, Y. Wang, J. Levi, B. H. LaFranchi, T. M. Deveau, A. M. Buck, S. E. Munter, K. A. Asare, M. Aslam, W. Koch, G. Szabo, R. Hoh, M. Deswal, A. E. Rodriguez, M. Buitrago, V. Tai, U. Shrestha, S. Lu, S. A. Goldberg, T. Dalhuisen, J. J. Vasquez, M. S. Durstenfeld, P. Y. Hsue, J. D. Kelly, N. Kumar, J. N. Martin, A. Gambhir, M. Somsouk, Y. Seo, S. G. Deeks, Z. G. Laszik, H. F. VanBrocklin and T. J. Henrich (2024). "Tissue-based T cell activation and viral RNA persist for up to 2 years after SARS-CoV-2 infection." <u>Sci Transl Med</u> **16**(754): eadk3295.

Perea-Martinez, I., T. Nagai and N. Chaudhari (2013). "Functional cell types in taste buds have distinct longevities." <u>PLoS One</u> **8**(1): e53399.

Reiter, E. R., D. H. Coelho, E. French, R. M. Costanzo and N. C. Consortium (2023). "COVID-19-Associated Chemosensory Loss Continues to Decline." <u>Otolaryngol Head Neck Surg</u> **169**(5): 1386-1389.

Rodriguez, Y. A., J. K. Roebber, G. Dvoryanchikov, V. Makhoul, S. D. Roper and N. Chaudhari (2021). ""Tripartite Synapses" in Taste Buds: A Role for Type I Glial-like Taste Cells." <u>J Neurosci</u>.

Rogn, A., J. L. Jensen, P. O. Iversen and P. B. Singh (2024). "Post-COVID-19 patients suffer from chemosensory, trigeminal, and salivary dysfunctions." <u>Sci Rep</u> **14**(1): 3455.

Schober, P., C. Boer and L. A. Schwarte (2018). "Correlation Coefficients: Appropriate Use and Interpretation." <u>Anesth Analg</u> **126**(5): 1763-1768.

Sharetts, R., S. T. Moein, R. Khan and R. L. Doty (2024). "Long-Term Taste and Smell Outcomes After COVID-19." <u>JAMA Netw Open</u> **7**(4): e247818.

Shi, P., J. Zhang, H. Yang and Y. P. Zhang (2003). "Adaptive diversification of bitter taste receptor genes in Mammalian evolution." <u>Mol Biol Evol</u> **20**(5): 805-814.

Soriano, J. B., S. Murthy, J. C. Marshall, P. Relan and J. V. Diaz (2022). "A clinical case definition of post-COVID-19 condition by a Delphi consensus." <u>Lancet Infect Dis</u> **22**(4): e102-e107.

Wang, J., R. Liu, H. Ma and W. Zhang (2023). "The Pathogenesis of COVID-19-Related Taste Disorder and Treatments." <u>J Dent Res</u> **102**(11): 1191-1198.

Wilson, C. E., R. S. Lasher, E. Salcedo, R. Yang, Y. Dzowo, J. C. Kinnamon and T. E. Finger (2025). "Death in the Taste Bud: Engulfment of Dying Taste Receptor Cells by Glial-Like Type I Cells." <u>Glia</u>.

Wilson, C. E., R. S. Lasher, R. Yang, Y. Dzowo, J. C. Kinnamon and T. E. Finger (2022). "Taste
Bud Connectome: Implications for Taste Information Processing." J Neurosci **42**(5): 804-816.

Xu, Z., W. Wang, D. Zhang, K. W. Tam, Y. Li, D. C. C. Chan, Z. Yang and S. Y. S. Wong
(2024). "Excess risks of long COVID symptoms compared with identical symptoms in the
general population: A systematic review and meta-analysis of studies with control groups." J
Glob Health **14**: 05022.

Yang, R., Y. K. Dzowo, C. E. Wilson, R. L. Russell, G. J. Kidd, E. Salcedo, R. S. Lasher, J. C.
Kinnamon and T. E. Finger (2020). "Three-dimensional reconstructions of mouse circumvallate
taste buds using serial blockface scanning electron microscopy: I. Cell types and the apical region
of the taste bud." J Comp Neurol **528**(5): 756-771.

Yao, Q., M. E. Doyle, Q.-R. Liu, A. Appleton, J. F. O'Connell, N.-p. Weng and J. M. Egan
(2023). "Long-Term Dysfunction of Taste Papillae in SARS-CoV-2." NEJM Evidence.

Yao, Q., M. E. Doyle, Q. R. Liu, A. Appleton, J. F. O'Connell, N. P. Weng and J. M. Egan
(2023). "Long-Term Dysfunction of Taste Papillae in SARS-CoV-2." NEJM Evid **2**(9).

Zazhytska, M., A. Kodra, D. A. Hoagland, J. Frere, J. F. Fullard, H. Shayya, N. G. McArthur, R.
Moeller, S. Uhl, A. D. Omer, M. E. Gottesman, S. Firestein, Q. Gong, P. D. Canoll, J. E.
Goldman, P. Roussos, B. R. tenOever, B. O. Jonathan and S. Lomvardas (2022). "Non-cell-
autonomous disruption of nuclear architecture as a potential cause of COVID-19-induced
anosmia." Cell **185**(6): 1052-1064 e1012.

Zhang, Y., M. A. Hoon, J. Chandrashekar, K. L. Mueller, B. Cook, D. Wu, C. S. Zuker and N. J.
Ryba (2003). "Coding of sweet, bitter, and umami tastes: different receptor cells sharing similar
signaling pathways." Cell **112**(3): 293-301.

Zhao, G. Q., Y. Zhang, M. A. Hoon, J. Chandrashekar, I. Erlenbach, N. J. Ryba and C. S. Zuker
(2003). "The receptors for mammalian sweet and umami taste." Cell **115**(3): 255-266.
768
769