Taste dysfunction in Long COVID.

Hanna Morad[1], Tytti Vanhala[2] Marta A. Kisiel [3], Agnes Andreason[3], Mei Li[4], Göran Andersson[2], Göran Laurell[5], Thomas E. Finger[# 4], and Göran Hellekant[# 2,6].

Clinic of Ear, Nose and Throat diseases (ENT), Uppsala University Hospital, Uppsala, Sweden.

Department of Animal Biosciences, Swedish University of Agricultural Sciences, Uppsala, Sweden.

Occupational and Environmental Medicine, Department of Medical Sciences, Uppsala University, Uppsala, Sweden

Rocky Mountain Taste & Smell Center, Department of Cell & Developmental Biology, University of Colorado School of Medicine, Aurora CO 80045, USA.

Department of Surgical Sciences, Uppsala University, Uppsala, Sweden.

University of Wisconsin Madison, School of Veterinary Medicine, Madison, WI, USA.

#Shared Senior authors
Correspondence to be sent to: Tom.Finger@CUAnschutz.edu

agnes.andreason@gmail.com	ORCID 0009-0002-3236-2201
hannaa_morad@hotmail.com	ORCID 0009-0005-0813-6273
mei.li@cuanschutz.edu	ORCID 0009-0009-9170-9376
tytti.vanhala@slu.se	
marta.kisiel@medsci.uu.se	ORCID 0000-0002-0410-1509
goran.andersson@slu.se	ORCID. 0000-0001-5131-3144
goran.laurell@uu.se	ORCID 0000-0002-7760-246X
ghellekant@gmail.com	ORCID 0000-0003-3562-8677
tom.finger@CUAnschutz.edu	ORCID 0000-0001-5923-5099

CORRESPONDING AUTHORS

Correspondence to Göran Hellekant and Thomas E. Finger

Abstract

Persistent taste dysfunction is frequently reported in individuals with post-acute sequelae of infection by SARS-CoV-2 (Long COVID). The mechanisms and pathological correlates underlying this taste dysfunction are unknown. This study investigates the underlying pathology in 28 non-hospitalized subjects diagnosed with COVID-19 and who experienced taste disturbances more than 12 months after testing positive for SARS-CoV-2. To objectively establish the nature of taste deficit, we used the WETT taste test, which quantifies the subject's ability to taste each of the five taste qualities: sweet, umami, bitter, sour, and salty. We then biopsied five to eight fungiform taste papillae (FP) in 20 of the 28 subjects. The FPs were analyzed histologically for overall taste bud structure and innervation, and by quantitative PCR (qPCR) for mRNA expression of markers for different taste receptor cells. Although all subjects had reported taste dysfunction, only three showed overall taste scores below the 10^{th} percentile for a normal population adjusted for age and sex. However, 11 of the 28 subjects exhibited total loss of one or more taste qualities. Loss of PLCβ2-dependent taste qualities (sweet, umami, bitter) was significantly more common and was correlated with reduced expression of *PLCβ2* and *Tas1R3* mRNAs. Histological analysis revealed generally preserved taste bud structure and innervation, but with occasional disorganized taste buds and abnormal, isolated PLCβ2-positive cells in the epithelium. Our findings suggest long-term taste dysfunction after COVID-19 occurs rarely -- more frequently involving PLCβ2-dependent taste qualities -- but is not due to wholesale disruption of the taste periphery.

INTRODUCTION

The COVID-19 pandemic has brought unprecedented attention to impairments of olfaction and taste as hallmark symptoms of acute Severe Acute Respiratory Syndrome-CoronaVirus-2 (SARS-CoV-2) infection (Cooper *et al.* 2020; Hannum *et al.* 2023; Klein *et al.* 2021a; Klein *et al.* 2021b). In most COVID-19-affected individuals, this dysfunction resolves within weeks but can persist for months for others. Prolonged dysfunction lasting more than three months is now recognized as part of post-acute COVID-19 syndrome (PACS) or Long COVID (Ely *et al.* 2024; Soriano *et al.* 2022). Ongoing taste dysfunction may occur even after resolution of other symptoms (Boldes *et al.* 2024; Xu *et al.* 2024) thus raising questions about the underlying mechanisms. Understanding the long-term pathophysiology of taste dysfunction and, in extreme cases, taste loss is crucial for patient care and the development of targeted rehabilitation strategies (Lee *et al.* 2025).

Sensations of taste are largely mediated by lingual taste buds (TBs) situated in three types of papillae: fungiform, vallate, and foliate. The fungiform papillae (FP) are concentrated on and around the tip of the tongue, making them easy to biopsy. A typical human has about 200 FP with 0-50 taste buds (TB) in each FP (Cheng and Robinson 1991; Miller 1986; Miller 1988 ; 1989 ; Miller and Reedy 1990).

Taste buds are complex multicellular epithelial end organs comprising between 20 and 100 taste cells (TCs) with approximately half being taste receptor cells (TRCs) (High and Finger 2025; Ogata and Ohtubo 2020; Ohtubo and Yoshii 2011; Yang *et al.* 2020). Histochemical, immuno-histochemical and molecular methods allow for correlation of TRC subtypes with specific taste qualities. Type II TRCs detect sweet, umami, or bitter tastes via distinct G-protein-coupled receptor (GPCR) pathways involving downstream PLCβ2-dependent signaling cascades (Shi *et al.* 2003; Zhang *et al.* 2003). Sweet taste is mediated via TAS1R2/R3 heterodimeric

receptors, and umami by TAS1R1/R3; bitter taste involves TAS2Rs, another family of taste

GPCRs. In contrast, Type III cells are responsible for transduction of ionic taste qualities, i.e.

sour and high concentrations of salt (Chang *et al.* 2010; Matsumoto *et al.* 2011) while Type I TCs

play a role similar to glia cells in the CNS (Bartel *et al.* 2006; Rodriguez *et al.* 2021; Wilson *et al.*

2025). In addition, taste buds contain a population of post-mitotic Type IV cells which mature to

differentiate into the other cell types. Disruption to these distinct molecular cascades or to the

morphological integrity of taste buds could contribute to the observed taste dysfunction in Long

COVID (Huang *et al.* 2021a).

Taste bud cells undergo continuous turnover, with new TRCs differentiating only when taste

fibers reach them (Farbman 1965; 1980; Farbman *et al.* 2004; Hellekant *et al.* 1987; Lee *et al.*

2017). Approximately 70–80% of mature TRCs form synapses with nerve fibers, highlighting the

importance of neural connectivity in maintaining taste function (Lee *et al.* 2017; Wilson *et al.*

2025; Wilson *et al.* 2022). Ongoing post-viral disruption of the nerve-TC interactions could well

lead to taste dysfunction.

SARS-CoV-2 infects cells via the ACE2 receptor and the TMPRSS2 protease, both of which

are expressed in oral epithelial cells, including Type II TRCs (Doyle *et al.* 2021; Yao *et al.*

2023). This suggests a plausible mechanism for direct viral-caused damage to the taste system.

Additionally, inflammatory processes, immune dysregulation, or altered saliva composition may

contribute to long-term taste changes following viral infection.

Given the clinical and biological implications of long-term taste dysfunction, this study aims

to investigate self-reported and objectively measured taste impairment in subjects who

experienced non-hospitalized COVID-19. Using a combination of psychophysical testing,

histological evaluation and molecular analysis -- including quantitative polymerase chain reaction

(qPCR) for mRNAs encoding taste cell markers and viral RNA -– this study seeks to elucidate

the structural and molecular changes underlying persistent taste dysfunction in Long COVID

subjects following COVID-19 caused by SARS-CoV-2.

MATERIAL & METHODS

Study design

This study involved subjects aged ≥18 years who had experienced symptomatic COVID-19

and a confirmed SARS-CoV-2 infection verified by a positive PCR test from nasopharyngeal

swabs between 2020 and 2021 (inclusion criteria). All subjects had non-hospitalized COVID-19

and, all but one, reported long-term taste disturbances. The study population was recruited

between November 2021 and June 2022 from two sources at Uppsala University Hospital: the

COMBAT post-COVID study and the Ear, Nose and Throat (ENT) outpatient clinic.

From the COMBAT post-COVID cohort, 18 subjects underwent psychophysical taste

evaluation. Of these, 12 consented to undergo a fungiform papilla biopsy (subjects #SWE7, 8, 9,

12, 13, 14, 15, 18, 19, 23, 24, 26). Additionally, ten subjects were recruited from the ENT

outpatient clinic, eight of whom underwent biopsy (subjects #SWE17, 21, 22, 24, 25, 27, 28, 29).

In total, eight subjects did not undergo a biopsy.

Study population

The COMBAT post-COVID project aimed to monitor persistent symptoms and long-term

consequences among non-hospitalized patients diagnosed with COVID-19. Initially, 725

individuals who tested positive for SARS-CoV-2 between March and December 2020 were

invited to complete a follow-up questionnaire 12 months after symptom onset. Of these, 401

subjects responded (response rate: 74%). The cohort has been described in detail elsewhere

(Janols *et al.* 2024; Kisiel *et al.* 2022; Kisiel *et al.* 2023a; Kisiel *et al.* 2023b; Kisiel *et al.* 2021).

Among the respondents, 68 individuals reported ongoing taste and/or smell disturbances. These

individuals were contacted by telephone and invited to participate in the current study if they

reported persistent taste loss and were available for in-person evaluation. Eighteen individuals

agreed to participate. Henceforth all individuals will be indicated as subjects.

Ten additional subjects attending the ENT outpatient clinic in 2021 were randomly selected

and invited to participate in the study if they met the same inclusion criteria. In addition, one

subject (#SWE17) reported no taste loss but had an active SARS-CoV-2 infection at the time of

sampling. Taste testing data for this subject are listed in Table 1 but are not included in the

statistical analysis of the cohorts reporting Long COVID taste disturbance.

Clinical and sociodemographic variables

From the COMBAT post-COVID cohort, information was documented about age, sex,

symptoms at COVID-19 onset, self-reported health before and one year after COVID-19, and

persistent symptoms one year following COVID-19. Symptoms at COVID-19 onset were also

retrieved from medical records from the PCR testing protocols. Additional subjects (n=11) were

characterized by age and sex only.

Ethical Approval

The study was approved by the institutional ethics committee of the University Hospital in

Uppsala (EPN number 2020-05707) and the Swedish Ethical Review Authority (EPM number

2021-00878) and complies with the Declaration of Helsinki for Medical Research involving

Human Subjects. The patients all gave written informed consent to the procedures. The

Combined Institutional Review Board at the University of Colorado at Anschutz approved the

transfer and analysis of the de-identified samples collected in Sweden. The study protocol in

Colorado (COMIRB number 20-1325) was determined Exempt/Non-Human Subject Research

and was conducted in accordance with 45 CFR 46 and institutional policies. All data were

pseudonymized to protect patient privacy.

Taste evaluation

As mentioned, the human sense of taste is divided into five taste qualities: sweet, umami,

bitter, sour and salty, (reviewed in:(Hellekant 2024; Lindemann 1996). To assess a subject's

ability to taste we used the quantitative and qualitative Waterless Empirical Taste Test (WETT®)

by Sensonics (Haddon Heights, NJ, USA). The WETT® test also measures the ability to taste

each taste quality with an eight-point scale from zero (no taste) to a maximum of eight (all

concentrations identified). The WETT® measures the ability to identify various concentrations of

the above five basic taste qualities with a minimum number of trials. It consists of 53 strips

impregnated with one tastant (a substance that produces a taste sensation) representing a specific

taste quality. A monomer cellulose pad (1×2.5 cm) is present on one side of each strip,

containing either: dried sucrose (sweet) (0.20, 0.10, 0.05, or 0.025 g/ml), citric acid (sour) (0.025,

0.05, 0.10, or 0.20 g/ml), sodium chloride (salty) (0.0313, 0.0625, 0.125, or 0.25 g/ml), caffeine

(bitter) (0.011, 0.022, 0.044, or 0.088 g/ml), monosodium glutamate (umami; brothy) (0.017,

0.034, 0.068, or 0.135 g/ml), or blank control strips, with no added chemical stimulus. The

subjects were instructed to place the appropriate strip on the tongue and move it along the surface

of the tongue for at least 5–10 seconds. The subject indicated which taste sensation was perceived

sött (sweet), bittert/beskt (bitter), surt (sour), buljongliknande smak/umami (umami), or salt

(salty), or whether no taste was sensed by verbally reporting their response to be entered on the

designated line of the response form. In the WETT® test sequence, the concentrations of each

stimulus are presented twice (eight total trials/tastant). In the first half of the test, the

concentrations ascend from weak to strong with the different tastants presented randomly, with

no tastant blanks included. In the second half of the test, the reverse presentation order is used,

i.e., going from strong to weak concentrations. Scores range from 53 (all correct) to 13 (no tastes

reported, corresponding to the number of control strips in the test). Each subject's results were

compared with normative values for each sex and age given in Doty et al (Doty *et al.* 2025).

Biopsy procedures

Five to eight fungiform papillae (FP) were biopsied from each subject's tongue and

preserved for histological and molecular analyses. Prior to biopsy, the tongue was anesthetized

with a Xylocaine spray (100 mg/ml Lidocaine with 10mg/spray dose), then blue or green food

coloring was added to enhance the tongue's fungiform papillae (FP). The tongue was held steady

with a gauze, and biopsies of the FP were taken with a 3-mm sterile microscissor with a curved

tip. The FP was cut one by one by holding down the scissor tip at the base of the chosen FP,

which was removed with a single cut and put in fixative for histology or RNA*later*™ (Thermo

Fisher Scientific Inc., Waltham, MA, USA). The biopsies for histological analysis were fixed in

4% buffered formaldehyde for at least four hours or overnight and then moved to phosphate-

buffered saline (PBS) solution. The biopsies were stored at 4°C until shipping for analyses from

Sweden to the United States for histopathological studies. The biopsies for RNA analysis were

placed in RNA*later*™ stabilization solution and stored at room temperature overnight and then

stored in the freezer at -18 °C initially and then at -80°C until analyses.

For all but two biopsy subjects, samples were taken immediately following the WETT test or

within two months of the WETT test as indicated in Table 1. For the other two subjects (subjects

SWE26 and 21), biopsies were taken four months following the WETT test. No biopsy was

requested for subjects SWE1-6 or SWE10 because these subjects showed relatively normal total

taste scores; subject SWE16 declined the biopsy procedure. For all other subjects (SWE7-9 ;

SWE12-15, and SWE 18-29) biopsies were obtained without regard to taste score. Subject

SWE17, who gave a biopsy sample, was a subject who recently tested positive for SARS-CoV-2

but reported no associated taste dysfunction. Histological and molecular data from this subject

are included in our analyses, but the taste testing data are not included in the analysis of Long

COVID taste testing which includes all subjects reporting taste dysfunction at the time the biopsy

was taken and no biopsy cohorts.

Histology

Upon receipt of fixed samples in Colorado, all specimens were cryoprotected in 20%

Sucrose in 0.1M phosphate buffer overnight at 4°C. Then, the tissues were cryo-sectioned (12-

14μm) onto Superfrost Plus microscope slides (Thermo Fisher Scientific, (Catalog# 1255015)

and air-dried. Slides were rinsed in 0.1M PBS pH 7.2. Antigen retrieval was performed with 10

mM sodium citrate buffer of pH 6.0 at 85°C for 10 minutes before incubation in blocking

solution (2% normal donkey serum, 1% bovine serum albumin, and 0.3% Triton in 0.1M PBS)

for an hour at room temperature. Slides were then incubated for four days in specific primary

antibodies diluted in blocking solution at 4°C including the following antisera:

The first two antisera were utilized in all cases, the latter two were used in some cases.

•   Mouse anti-Tubulin β3 (clone TUJ1) at 1:1000 dilution (BioLegend, San Diego,

CA, USA, catalog# 801202; RRID:AB_2313773) (for total innervation),

•   Rabbit anti-PLCβ2 (PhosphoSolutions, Denver, CO, USA, catalog# 1640-PLC;

RRID:AB_2801663), or guinea pig anti-PLCβ2 (PhosphoSolutions, AP8/24; RRID:

AB_2910247) (for identification of Type II TRCs).

•   Goat anti-GNAT3 at 1:500 dilution (Aviva Systems Biology, San Diego, CA, USA,

catalog# OAEB00418; RRID:AB_10882823) (for TRCs expressing the taste-associated

G-protein gustducin),

•   Rabbit anti P2X3 (Alomone, Jersusalem, Israel, catalog #APR-016, RRID

AB_2313760) (for labeling of taste bud-specific innervation).

After washing in 0.1M PBS, the slides were incubated two hours with fluorescent secondary

antibodies, Alexa Fluor 488 donkey anti-mouse (Invitrogen, catalog# A21202;

RRID:AB_141607), Alexa Fluor 568 donkey anti-rabbit (Invitrogen, catalog# A10042), Alexa

Fluor 647 donkey anti-goat (Invitrogen, catalog# A21447; AB_2534017), Alexa Fluor 647

donkey anti-guinea pig (Jackson Immuno Research Laboratories, West Grove, PA, USA,

catalog# 706605148; RRID: AB_2340476) or Donkey anti-Rabbit IgG (H+L) Highly Cross-

Adsorbed Secondary Antibody, Alexa Fluor™ 647 (ThermoFisher Catalog # A-31573;

AB_2536183) at 1:800 dilution each at room temperature. Lastly, the slides were washed twice

for 10 minutes each in 0.1M PBS and then once for 10 minutes in 0.05M PBS before coverslip

slides with DAPI Fluoromount-G (SouthernBiotech, Birmingham, AL, USA, catalog# 010020).

RNA extraction

The papilla tissue samples for PCR analysis were stored in RNA*later*™ at -20°C. RNA was

extracted with the RNeasy fibrous tissue mini kit (Qiagen, Hilden, Germany). A small piece of

the biopsy was taken with a pipet tip into a 1.5 ml tube containing 300 μl of RLT buffer and

mashed manually with a micro pestle. The pestle was rinsed when pipetting 590 µl of RNase free

water into the tube. The kit protocol was followed, and a DNase treatment was performed with a

DNA-free DNA removal kit (Invitrogen) according to their protocol. The resulting RNA

concentrations and quality were checked with Nanodrop 8000 (Thermo Fisher Scientific) and

TapeStation High Sensitivity RNA tape (Agilent, Santa Clara, CA, USA).

Reverse transcription

High-Capacity cDNA Reverse Transcriptase Kit (Applied Biosystems, Waltham, MA, USA)

was used to obtain cDNA from the RNA samples. 100 ng of RNA per sample, according to

TapeStation concentrations, was taken into the reverse transcription reaction. The kit protocol

was followed. The reverse transcription was performed in duplicates to obtain enough product for

all the qPCR runs.

qPCR

Quantitative PCR (qPCR) was performed with TaqMan (Applied Biosystems) assays for

PLCβ2 (Hs01080542), *NCAM1* (Hs00941821), and *TAS1R3* (Hs01026531) as well as for two

reference genes *RPLP0* (Hs99999902) and *GAPDH* (Hs02758991). TaqMan Gene Expression

Master Mix (Applied Biosystems) was used according to its protocol, and 10 ng of cDNA was

added as template. All samples were run in triplicate per gene, including a negative control. One

gene per plate was run in the QuantStudio 5 (Applied Biosystems). Average Ct and standard

deviation were calculated for each gene and then normalized (Delta CT =Ct(gene of interest) –

Ct(reference gene)) against the reference gene *GAPDH*. The reference gene *GAPDH* gave more

consistent normalization than *RPLP0*, and was therefore selected as a reference for the genes of

interest.

SARS-CoV-2 test

A PCR test for the presence of SARS-CoV-2 virus in the tissue samples was performed in

triplicate, including two assays that target SARS-CoV-2 as well as negative control samples.

TaqMan microbe detection assay (Vi07918637) for SARS-CoV2N and TaqPath Blastopore

Microbial Detection Master Mix (Applied Biosystems) were used according to their protocol.

Positive control was TruMark Respiratory Panel 2.0 Amplification control (Applied Biosystems).

The qPCR was run in the QuantStudio 5 (Applied Biosystems). None of the biopsied tissue

samples tested showed the presence of SARS-CoV-2 virus, although one subject (#17) had an

active SARS-CoV-2infection as determined by a positive nasal swab test. Absence of detectable

SARS-CoV-2 virus in oral or salivary samples despite positive detection of virus in nasal swabs

is not uncommon and is described in other studies (Fernandes *et al.* 2021; Goodall *et al.* 2022).

Statistical analysis

Parametric statistical calculations were carried out by Excel (Microsoft) or with calculators

given at www.socscistatistics.com. Both calculators gave identical values. (Ho *et al.* 2019) For

comparisons (Ho *et al.* 2019)across samples, we utilized Pearson's Correlation co-efficient or

Fisher's Exact test for small sample sizes. For comparison of test percentile means against the

expected 50% percentile distribution, we employed a one-sample t-test. A criterion value of $p <$

0.05 was set for statistical significance in all calculations.

Means, standard deviations, and ranges were used to describe continuous variables such as

age, time from infection to testing, and taste scores. Categorical variables like sex and symptom

presence were summarized using counts and percentages.

Because taste ability varies substantially by sex and age, raw taste scores were normalized by

sex and age using reference data from Doty et al. (2025), which included 1,392 healthy

individuals. Scores for individual tastes below the 10% compared to sex and age matched

populations in Doty et al (2025) were considered potentially hypoguesic per the criterion set in

(Landis *et al.* 2009) and are highlighted in red in Table 1.

The binomial test or Chi-square tests (χ^2) were used to compare (categorical data) the

frequency of abnormal taste scores across groups. The results for Chi-square tests are reported as

χ^2 (degree of freedom (df), sample size) = chi-square value, p-value. Pearson's Correlation (r) was

used to assess linear relationships between molecular markers (qPCR results), histological

features, and taste scores. The results were reported as (df) sample size=correlation co-efficient,

p-value. (Schober *et al.* 2018).

RESULTS

Study population

All 28 subjects (24 women and 4 men) reported normal taste before the first confirmed

SARS-CoV-2 infection but, except for subject SWE17, altered or diminished taste thereafter .

The mean age at testing was 49.9 years (SD+/-1.2) and ranged between 30 and 71 years. Most

subjects contracted COVID-19 in early 2020. Two subjects (SWE19 & 28) experienced a second

infection in 2022, shortly before undergoing taste testing and biopsy. Considering only the first

infection, the time from initial SARS-CoV-2 infection to taste testing ranged from 13.5 months to

27.0 months (mean of 21.0 months).

Psychophysics:

Because taste ability varies substantially between individuals according to sex, age, and

health status, we normalized all raw taste scores and overall WETT score against values given for

1,392 normal subjects (Doty *et al.* 2025). Although all subjects, except one (subject #SWE17),

reported compromised taste abilities in clinical reporting, only 3 of 28 tested subjects scored

below the 10th percentile in overall taste ability consistent with overall scores in a population of

normal subjects (Doty *et al.* 2021) (see Table 1) and consistent with previous reports of no

demonstrable long-term taste loss in broad post-covid populations following SARS-CoV-2

infection (Boscolo-Rizzo *et al.* 2023a; Sharetts *et al.* 2024). Nonetheless, 14 of the 27 of our

Long Covid dysguesia subjects showed significant deficiency (10th percentile or lower) in one or

more individual taste qualities (values in red shown in Table 1) which is significantly more than

expected in a normal population (binomial test 24 of a total of 135 individual taste qualities; p =

0. 002). sex(Doty *et al.* 2021) One subject, who was also anosmic, (SWE29) gave no correct

responses to any tastant, even at the highest concentration suggesting complete aguesia. Even

subjects with a near-average total WETT score (percentiled by sex and age group) were unable to

detect specific tastes, e.g. subjects SWE8 (F, age 64; WETT Score 32; 45%) and SWE9 (M, age

49; WETT Score 30; 54%) were unable to detect bitter or umami. Although individuals with a

profound overall deficit also showed specific loss of one or more taste qualities (e.g. subject

SWE7 [16%] showed no response to sweet, bitter or umami, but good response to sour and salty),

the pattern of loss was not duplicated by other subjects (e.g. subject SWE24 [7%] with good

responses to sweet, sour and salty, but total loss for bitter and umami).

The distribution of loss between subjects was uneven with sour standing out as the least

affected quality (only one score below the 10%). This may reflect the detection of "sour" being a

multisensory experience involving chemesthetic as well as gustatory modalities (Ohkuri *et al.*

2012; Yu *et al.* 2020; Zhang *et al.* 2019). In contrast, in acute COVID-19, sour taste is reported to

be most impacted during the acute phase of the illness (Karmous *et al.* 2023; Singer-Cornelius *et*

al. 2021).

More importantly, as a group, the PLCβ2-mediated taste qualities (sweet, umami, bitter)

showed total loss (score of 0) as well as scores below 10% significantly more than the other taste

qualities (sour, salty) ($X^2(1, N = 135) = 4.47, p = .03$). These findings suggest that Long Covid

taste disturbance may preferentially target the PLCß2-mediated taste qualities transduced by

Type 2 taste receptor cells.

	Subject Number	Sex	Age	Covid Date	Taste Test Date	Biopsy Date	PLC Tastes			Sour	Salty	No Taste	WETT score	Sex/Age-corrected %
							Sweet	Umami	Bitter					
Acute Biopsy	SWE 17	F	30	Apr 2020	31/1-22	31/1-22	8	8	8	8	8	13	53	99
Post-covid No Biopsy	SWE 1	F	47	Mar 2020	19/11-21		5	3	1	5	6	12	32	42
	SWE 2	F	62	Apr 2020	19/11-21		3	0	1	8	8	11	31	36
	SWE 3	F	50	Apr 2020	24/11-21		6	0	3	7	5	12	33	45
	SWE 4	F	49	May 2020	3/12/21		0	8	6	6	7	10	37	59
	SWE 5	F	53	Nov 2020	17/1-22		8	4	8	8	7	13	48	95
	SWE 6	F	41	Nov 2020	18/1-21		7	2	4	5	4	12	34	51
	SWE 10	F	66	Nov 2020	20/1-22		8	1	8	7	5	13	42	83
	SWE 16	F	59	Apr 2020	11/2/22		5	3	6	8	7	10	39	70
Post-Covid Biopsy	SWE 7	F	68	Nov 2020	18/1-22	23/3-22	0	0	0	6	5	13	24	16
	SWE 8	F	64	Mar 2020	18/1-22	23/3-22	6	0	0	6	7	13	32	45
	SWE 9	M	49	Apr 2020	19/1-22	17/3-22	1	0	0	8	8	13	30	54
	SWE 12	F	56	Apr 2020	21/1-22	24/1-22	3	3	6	7	4	12	35	54
	SWE 13	M	45	Apr 2020	21/1-22	21/1-22	0	1	0	5	3	13	22	21
	SWE 14	F	30	Apr 2020	21/1-22	21/1-22	3	8	5	8	6	10	40	70
	SWE 15	F	53	Apr 2020	24/1-22	24/1-22	4	6	3	4	4	13	34	54
	SWE 18	F	63	June 2020	23/3-22	23/3-22	7	1	4	6	1	13	32	45
	SWE 19	F	46	May 2020 & Jan 2022	24/3-22	24/3-22	7	4	8	6	2	9	36	59
	SWE 20	F	38	Feb 2021	30/3-22	20/4-22	7	2	7	8	8	13	45	83
	SWE 21	F	71	Jan 2021	17/1-22	11/5-22	4	1	5	7	4	13	34	70
	SWE 22	F	56	Apr 2020	11/5-22	11/5-22	1	4	0	6	2	13	26	21
	SWE 23	F	49	Nov 2020	12/5-22	12/5-22	4	4	7	6	5	13	39	67
	SWE 24	M	42	June 2020	13/5-22	13/5-22	5	0	0	4	4	4	17	7
	SWE 25	F	50	Mar 2020	13/5-22	13/5-22	2	4	1	0	5	7	19	4
	SWE 26	F	24	June 2020	21/1-22	16/5-22	6	0	6	5	7	13	37	44
	SWE 27	F	52	Mar 2020	15/6-22	15/6-22	0	0	4	2	6	13	25	15
	SWE 28	F	43	Nov 2020 & Feb 2022	5/7-22	5/7-22	6	3	7	7	3	10	36	44
	SWE 29	M	30	April 2020	15/7-22	15/7-22	0	0	0	0	0	13	13	1
Post-covid aguesia n= 27					Post-covid Cohorts ave Taste Score		4.0	2.3	3.7	5.7	4.9	11.6	32.3	46.5
Post-covid Biopsy n= 18					scores <10%		7	3	9	1	4			
red text ≤10%					zero scores		5	9	7	2	1			

Table 1: Results from WETT testing of all subjects including basic demographic information, dates of infection, taste testing and biopsy. Columns labeled as "Sweet, Umami, Bitter, Sour, Salty and No Taste" show WETT test raw scores for each taste quality and blank "No Taste" control strips. Red numerals in columns for individual tastes indicate values below the 95% confidence interval for sex and age-matched controls based on Doty, et al 2025. The incidence of values below the 10% for all post-Covid subjects is significantly higher than expected based on a normal population ($\chi^2(1, N = 135) = 4.47$, $p < .05$). Zero values indicate that the subject was unable to identify the tastant at any concentration offered. The total number of zero values for each tastant is shown in red along the bottom row. WETT score shows the total WETT score for each subject alongside of the corresponding sex and age matched percentile ranking compared to the study population in Doty et al 2025. A perfect score is 53 (8 for each taste quality + 13 blanks). Complete ageusia would give a score of 13 (0 for all tastants and 13 for correctly identifying blank controls). PLCß2-dependent tastes are indicated by orange shading along the top and bottom rows.

Quantitative PCR Findings

RNA prepared from taste papillae from all biopsied subjects, including SWE17, was analyzed for expression levels of three taste-related mRNAs encoding: 1) phospholipase C beta 2 (PLCß2) a marker of Type II cells, 2)Taste receptor member *TAS1R3*, coding for a receptor component for both sweet and umami tastes and expressed in a subset of Type II taste cells; and 3) neural adhesion molecule 1 (*NCAM1)*, a marker of total innervation as well as

Subject #		taste test value	relative expression			Ave PLC + Tas1R3	ave sweet/ umami
			PLCb2	TAS1R3	NCAM1		
SWE	**17**	**53**	**1.00**	**1.00**	**1.00**	**1.00**	**8**
SWE	7	24	0.15	0.11	0.96	0.13	0
SWE	8	32	0.66	0.34	0.60	0.47	3
SWE	9	30	0.29	0.42	0.40	0.35	0.5
SWE	12	35	0.78	0.51	1.17	0.63	3
SWE	13	22	0.30	0.78	1.01	0.49	0.5
SWE	14	40	0.53	1.60	1.21	0.92	5.5
SWE	15	34	0.56	0.77	0.67	0.66	5
SWE	18	32	0.14	0.30	0.15	0.20	4
SWE	19	36	0.16	0.76	0.08	0.35	5.5
SWE	20	45	1.64	0.80	0.86	1.14	4.5
SWE	21	34	0.27	0.23	0.34	0.25	2.5
SWE	22	26	0.36	0.28	0.52	0.32	2.5
SWE	23	39	0.64	1.14	1.96	0.86	4
SWE	24	17	0.62	1.16	1.33	0.84	2.5
SWE	25	19	0.19	0.17	0.52	0.18	3
SWE	26	37	0.48	0.59	0.57	0.54	3
SWE	27	25	1.08	0.77	1.64	0.91	0
SWE	28	36	0.44	0.54	0.32	0.49	4.5
SWE	29	13	0.22	0.15	0.59	0.18	0

Table 2: Relative expression of taste-related mRNAs from qPCR of fungiform papillae. Subject SWE 17, who reported no taste disturbances, served as our comparator. Taste test values as shown in Table 1 are included for comparison. For each subject, the average of the scores for sweet and umami taste, tastes reliant on Tas1R3, are shown as well.

Type III TCs. All data were normalized against glyceraldehyde-3-phosphate hydrogenase

(*GAPDH)* mRNA, yielding Delta CT values which are expressed as a fold-change compared to

the values from subject SWE17 who described no taste dysfunction. These relative expression

values are given in Table 2 along with Taste Scores and computed average taste score for sweet

and umami (Ave Sweet-Umami) – the Tas1R1-related taste qualities -- for comparison.

The expression levels for *PLCß2* and *TAS1R3* mRNAs, both encoding components of the

taste transduction cascade in Type II TCs, were highly correlated (r(18) = 0.64, p=0.002) as

expected based on molecular features of Type II taste cells in mice. Expression of *NCAM* mRNA

was not significantly correlated with either *PLCß2* or *Tas1R3* mRNAs.

Comparing the qPCR data from all biopsy subjects to the taste scores from the same subjects

shows a significant correlation of overall taste score to *PLCß2* (r(18) = -0.51, p= 0.023) and

substantial correlation with *TAS1R3* (r(18) =--0.43, p= 0.056; although not significant at p<0.05

level). This relationship is shown graphically in Fig. 1. Conversely, expression levels of *NCAM*

mRNA showed no

substantial correlation

with taste test values

(r(18) = 0.035, p>0.05)

Since both sweet

and umami require the

TAS1R3 receptor

subunit as well as

PLCß2, we compared

the expression level of

these mRNAs to the

average of taste scores

for sweet and umami

for each subject. A

Figure 1: Correlation of relative *mRNA* expression levels with taste scores. The point indicated in orange is the data from Subject SWE17 who was used for comparison.
 A) The mRNA levels for *PLCß2* (r(18)=-.51, p= .023) and *TAS1R3*(r(18)=.43, p= .056) correlated with the overall taste score for each subject.
B) The mRNA level for *TAS1R3* correlates with the combined taste score for the related taste qualities (Sweet and brothy) (r(18)=.51, p= .02).

Alt Text: Graphs showing the correlation between Taste Score and PCR analysis of taste-related markers.

significant correlation exists between the average of the related taste scores for sweet and umami

with theDelta CT relative expression values for *TAS1R3* (r(18)= 0.45, p=0.046) and *PLCß2*

(r(18)= 0.55, p=0.01). These expression levels were, however, not significantly correlated with

expression levels of the individual tastes: sweet or umami.

Histological Findings

Fungiform samples

from the subjects varied

between 4 and 8 papillae,

with a median of 7.

Approximately half of the

papillae had no

identifiable TBs, as

reported for human

fungiform papillae in

previous literature e.g.

(Miller 1988). Normal-

appearing taste buds were

identified in all subjects,

including subject 29, who

reported no taste

sensibility. While

relatively normal TBs

occur in samples of all

Subject #	taste score	# papillae	# TBs	ave tb/pap	plc/tb	plc/pap
SWE 7	24	7	42	6.0	2.4	14.3
SWE 8	32	8	42	5.3	2.0	10.5
SWE 9	30	8	10	1.3	2.0	2.5
SWE 12	35	6	30	5.0	2.1	10.3
SWE 13	22	4	6	1.5	1.8	2.7
SWE 14	41	5	61	12.2	2.9	35.4
SWE 15	34	6	13	2.2	2.6	5.7
SWE 17	53	8	25	3.1	2.7	8.5
SWE 18	32	7	7	1.0	1.8	1.8
SWE 19	36	6	8	1.3	2.0	2.7
SWE 20	45	7	41	5.9	2.0	11.7
SWE 21	34	7	28	4.0	3.3	13.2
SWE 22	26	8	32	4.0	1.8	7.2
SWE 23	39	7	17	2.4	2.3	5.6
SWE 24	17	6	22	3.7	2.0	7.3
SWE 25	19	6	2	0.3	1.5	0.5
SWE 26	37	5	22	4.4	2.4	10.6
SWE 27	25	7	32	4.6	2.8	12.8
SWE 28	26	8	17	2.1	2.5	5.3
SWE 29	13	7	27	3.9	1.9	7.3

Table 3: Counts of papillae sampled, number of taste buds and average *PLCß2*+ cells per taste bud and papilla in each case. No significant correlation was found between taste score and average number of taste buds per papilla nor of density of *PLCß2*+ cells per taste bud or papilla. The histological counts and density of *PLCß2*+ cells also did not correlate with any PCR values. # TBs, number of taste buds; ave tb/pap, average number of TBs per papilla; PLC/tb, average number of PLC+ cells per taste bud; PLC/pap, average number of PLC cells per papilla.

subjects (See Fig. 2), many subjects displayed disorganized TBs or unusual isolated PLCß2+

cells which were never observed in TBs from normal subjects (High and Finger 2025)

The average number of TB profiles per papilla was 3.6 (+/-2.6 s.d.). Because of the

significant variance in the number of TB profiles observed in a single section through a papilla,

we calculated the average number of TB profiles per papilla from all papillae to measure the taste

bud population for each subject. In addition, we counted the number of PLCß2+ taste cells

within each section to calculate the average proportion of PLCß2+ cells per TB. The taste score

for each subject was not correlated significantly with the average number of TB profiles per

papillae ($r(18) = 0.268$, $p = 0.25$), with the average number of PLCß2+ cells per papilla ($r(18)$ -

0.35, $p = 0.15$), nor with the average number of PLCß2+ cells per TB ($r(18) = 0.427$, $p = .06$). As

shown in Fig. 3, abnormal TBs or irregular PLCß2+ cells were especially noteworthy in 7 of our

19 post-covid biopsy subjects (SWE12 (Taste Score 54%), 13(21%), 22 (21%), 24 (8 %), 25

(4%), 28 (44%) and 29 (1%)). This incidence is significantly different (Fisher's exact test p

<0.03) than in tissues obtained from normal subjects where dispersed PLCß2+ cells were never

observed in fungiform taste buds from 12 subjects.

Figure 2: Appearance of normal taste buds sectioned in different planes from different subjects. Red = PLCß2 immunoreactivity showing Type II taste cells; Green = TUJ1 immunoreactivity for nerve fibers. Typical taste buds have more than one PLCß2+ cell surrounded by a dense network of TUJ1-stained nerve fibers. Green profiles outside of taste buds are extragemmal nerve fibers typical of lingual epithelium (Moayedi *et al.* 2021). **a, b.** Taste buds sectioned longitudinally appear similar in shape to garlic bulbs or onions (from Subject (SWE) 15. **c)** Taste buds sectioned perpendicular to their long axis appear circular (from subject (SWE) 29 who reported a total absence of taste perception (taste score of 13, showing no correct responses to any tastant).
Scale bar in panel (b) applies to all panels.

Alt Text: Images of 3 micrographs showing the normal appearance of taste buds in our samples. Multiple PLCß2-positive taste cells within each bud are embraced by a dense nerve plexus.

Figure 3: Examples of abnormal PLCß2+ taste-like cells from 4 cases. (A & B). Subject (SWE) 12 had a taste score of 35 (54%). (C) From subject (SWE) 13 who had a score of 22 (21%). In both cases (A & C), innervated PLCß2+ (red arrows) cells occur in relative isolation within the epithelium. Innervation of these cells (green: TUJ1) is sparse unlike innervation of taste buds (cf Fig. 2). These cells may represent very small taste buds or simply PLCß2+ cells lying outside of a taste bud structure. Panel B shows a higher magnification of one such isolated PLCß2+ cell and its innervation stained for the presence of P2X3, a neural receptor predominantly on nerve fibers that innervate taste buds. Panels D-E are from case (SWE) 24 with a taste score of 17 (4%) showing examples of aberrant PLCß2+ cells (red arrows) outside of a taste bud structure. Panel F: A disorganized taste bud from case SWE 29 who reported a total absence of taste perception (taste score of 13, showing no correct responses to any tastant). The heavy innervation is typical of taste buds, but the cellular structure revealed by PLC-staining (red) does not show typical taste bud organization.

Alt Text: Images showing abnormal isolated PLC-positive cells and abnormal taste buds within the epithelium of our subjects.

DISCUSSION

Morbidity Changes in COVID-19 caused by SARS-CoV-2 Variants

Previous studies found that the frequency of chemosensory symptoms, including taste loss,

varies across COVID-19 illness caused by infection of different SARS-CoV-2 variants. Early

variants, such as Wild type and Alpha, were more commonly associated with taste and smell loss,

while later emerging variants, such as Omicron, caused lower levels of taste loss (Reiter *et al.*

2023). Most subjects included in our study were primarily infected during the first wave of the

COVID-19 pandemic, and their reported taste disturbances fit well with this pattern of greater

taste loss early in the pandemic.

Published epidemiological data further illustrate this trend. Among individuals infected with

the Wild-type strain, 241 out of 1,141 cases (21%) reported smell and/or taste disorders. For the

Alpha variant, 223 out of 1265 cases (18%), for later variants such as Delta, 480 out of 1516

cases (32%), and for Omicron, only 317 out of 7431 cases (4%) presented with smell and/or taste

disorders (Ota *et al.* 2025). This supports the hypothesis that the variant-dependent tropism and

immune response may influence the degree and duration of taste dysfunction.

On the role of the ACE2 receptor

Relatively early in the COVID-19 pandemic, it was suggested that SARS-CoV-2 infects oral

tissue (Huang *et al.* 2021b). Doyle et al. (Doyle *et al.* 2021) reported that ACE2 was expressed by

Type II TRCs, and further demonstrated the presence of SARS-CoV-2 in Type II TRCs, which

mediate sweet, umami and bitter sensations, but not within the other taste cell types. This likely

explains why subjects in that study did not taste the "sweetness of chocolate and bitter coffee"

and instead reported that it "tasted like mud" .

Research during the COVID-19 pandemic has highlighted the potential impact of SARS-

CoV-2 on TBs, particularly through its interaction with ACE2 receptors (Huang *et al.* 2021a; Yao

et al. 2023). In patients experiencing taste disturbances following COVID-19 infection, SARS-

CoV-2 was detected in basal and suprabasal layers of fungiform papillae for extended periods. In

one subject, virus was detected up to 63 weeks after the initial diagnosis, coincident with

disrupted nerve fibers and altered taste bud cytoarchitecture (Yao *et al.* 2023).

Recovery of taste, and re-establishment of normal epithelial cytoarchitecture, coincided with

the disappearance of the virus from the taste epithelium (Yao *et al.* 2023). These findings suggest

that prolonged viral presence or the possibility of reinfection and the associated immune response

may impair the renewal and functionality of taste buds in Long COVID. In our subjects, a

sensitive commercial qPCR test showed that none of the tissue samples were positive for the

SARS-CoV-2 viral RNA suggesting that taste dysfunction may persist even after the virus is no

longer detectable in the peripheral taste system. This offers the potential that persistent taste

dysfunction in Long Covid may outlast the presence of the virus itself.

Why loss of sweet-bitter-umami tasting ability is more frequent than the loss of other taste

qualities

A previous WETT study involving 340 long-term (1-year) COVID-19 subjects and 434 non-

COVID controls (Sharetts *et al.* 2024) found no statistically significant differences in overall

taste function between groups. Similarly, a psychophysical taste-strip study (Boscolo-Rizzo *et al.*

2023a) of 62 post-Covid subjects who reported an altered sense of taste or smell one year after

infection, found no statistically significant evidence for ongoing taste loss by two years after

infection, although subjects still scored modestly lower than the normal population. Similarly, our

subjects show non-significant, slightly lower than average total taste scores (46.5%±25.1 s.d.)

compared to a normal population. It should be noted that our study specifically targeted subjects

reporting ongoing taste loss, not general chemosensory disruption, two years after COVID-19

infection, preventing direct comparison of our results with those surveyed from the broader

population surveyed in Sharetts et al. (2024) or when patients with shorter-term chemosensory

loss were tested (Boscolo-Rizzo *et al.* 2023b).

Despite the lack of measurable overall taste loss in our subjects, our psychophysical results do

indicate a significant disruption of taste with a tendency toward preferential impact on sweet,

bitter, and umami -- the PLCβ2-related taste qualities (Zhang *et al.* 2003). These taste modalities

all are detected by the Type II taste cells which are targeted by SARS-CoV-2 virus (Doyle *et al.*

2021).

Taste dysfunction may result from disruption of taste-related intracellular transduction cascades

or damage to taste buds (TBs), their receptor cells (TRCs) or the gustatory nerves. (Liman 2020;

Liman and Kinnamon 2021; Zhang et al. 2003; Zhao et al. 2003). The fact that the SARS-CoV-2

can directly infect Type II TRCs may explain why deficits in sweet, umami and bitter are more

frequent than for sour and salt. In our study, however, viral RNA was not detectable in taste

tissues and there was no evidence of wholesale destruction or abnormality in Type II, PLCβ2+,

taste cells. Nonetheless, in our subjects overall tasting ability correlates positively with the

amount of PLCβ2 and TAS1R3 mRNAs present in the tissue. Curiously, a previous study

(Donadoni et al. 2023) of the consequence of SARS-CoV-2 infection of taste cells in an in vitro

culture system showed increases in PLCβ2 expression with infection. It is unclear how the acute

in vitro model may relate to the long-term post-covid conditions reflected in our study.

Hypothetically, the long-lasting effect could be the result of an epigenetic mechanism called

inflammatory memory (Naik et al. 2017). This is described in several inflammatory diseases and

the effect is manifested as a sustained modification of the chromatin structure. This leaves open

the question how prior viral infection might induce long-lasting deficits in taste.

In the olfactory epithelium, the virus infects supporting cells, but not the olfactory receptor

cells (Chen *et al.* 2020). Nonetheless, the olfactory receptor cells responded to the viral infection

of the tissue by downregulating expression of olfactory receptors and downstream signaling

molecules while upregulating expression of viral response proteins (Chen *et al.* 2019; Zazhytska

et al. 2022). The transduction cascade is re-established in the receptor cells once the viral

infection has been cleared (Zazhytska *et al.* 2022). While a similar phenomenon may occur in

taste buds during viral infection, it is unlikely such a reaction would persist in the tissue after the

virus is cleared and all the taste cells have been replaced during the course of normal taste cell

turnover which requires a few months at most (Perea-Martinez *et al.* 2013). At the time of our

testing, up to two years after initial infection, none of the taste cells present at that time would

have been present at the time of viral infection.

Furthermore, we do note that some subjects are unable to detect non PLCβ2-dependent taste

qualities, i.e. sour and salty. This finding may reflect inabilities typical of a normal population but

might suggest that non-PLCβ2 related mechanisms may also be at work underlying taste

dysfunction in Long COVID. One possibility is that the infection alters the function in taste

nerves. Indeed, evidence exists for widespread neural dysfunction in Long COVID (Daia *et al.*

2021; McFarland *et al.* 2021; O'Neill *et al.* 2023; Otte *et al.* 2022; Rogn *et al.* 2024; Stepien and

Pastuszak 2023). If so, then some of the observed taste loss might be attributable to loss of

transmission in the taste nerves rather than disruption of the taste epithelium. This could explain

the total loss of taste perception in subject SWE29 despite the presence of normal-appearing taste

buds.

Other factors that may have influenced taste during the disease process of COVID 19

Recent studies (Finlay *et al.* 2022; Peluso *et al.* 2024a; Peluso *et al.* 2023) broaden the

perspective by suggesting that immune dysregulation and T cell dysfunction may play significant

roles in the long-lasting effects of COVID-19. Whole-body positron emission tomography

imaging with [^{18}F]F-AraG, a tracer that selectively tags activated T cells, showed that subjects

with Long COVID had specific tissues enriched for activated T cells compared to individuals

without this condition (Peluso *et al.* 2024b). All five subjects in that study showed the presence

of intracellular SARS-CoV-2 spike protein-encoding RNA in lamina propria tissue up to 676

days after initial infection with SARS-CoV-2. This offers evidence of viral persistance or of

multiple independent SARS-CoV-2 infections which could underlie long-term immunologic

perturbations for up to two years after an initial infection. Whether a similar prolonged

immunological response occurs in taste tissue is unknown although Yao et al (Yao *et al.* 2023)

report long term (>6 months) inflammatory response when virus remains in the tissue. We did

not, however, detect any virus by PCR in our tissue samples from patients more than one year

post infection. Other investigators reported an association of elevated serum concentrations of IL-

10 and IL-1β with persistent taste dysfunction for bitter and sweet qualities, respectively,

following Covid-19 (Karmous *et al.* 2023; Locatello *et al.* 2021).

On the relation between histological changes of TB by COVID-19 and taste changes

A particularly intriguing aspect of our study is that all subjects, including those with severe deficits, displayed structurally intact taste buds despite the loss of taste function. However, some dystrophic taste buds and isolated PLCß2-expressing cells were observed in many subjects. In mice, disruption of molecular signals between nerves and taste buds can result in formation of aberrant PLCß2-expressing cells outside of the taste buds themselves (Castillo *et al.* 2014). This finding raises important questions about the disconnect between histological integrity and functional impairment. One possible explanation is that taste bud dystrophy, rather than outright loss of taste buds, may be sufficient to alter taste sensitivity by impairing cellular turnover or disrupting synaptic communication with gustatory neurons (Wang *et al.* 2023). While intact taste buds might suggest some functional resilience, the significant decrease in PLCß2-dependent signaling suggests that molecular-level dysregulation may underlie the persistent taste dysfunction. In addition, the discrepancy between PLCß2 expression and taste bud morphology, especially in Subject 29 who shows complete taste loss, suggests dysfunction in peripheral nerves or at higher levels of the neuraxis.

Clinical relevance

Understanding the mechanisms behind taste loss is clinically relevant, as chemosensory loss negatively impacts nutritional intake, emotional well-being, and quality of life. For survival, taste ability is more important than both hearing and vision. A multidisciplinary mindset is crucial for appropriate diagnostics and for providing patients with appropriate treatment and underscore the importance of developing targeted therapies to alleviate the burden of long-term taste dysfunction in COVID-19 survivors.

Here we present an integrative approach combining psychophysical taste testing, RNA expression analyses using markers for TRC type II and III, and histological evaluation of taste

tissue in subjects with Long COVID. This multidimensional analysis provides a more

comprehensive understanding of Long COVID taste dysfunction. While this study focuses on the

loss of Type II TRC, PLCß2-dependent taste qualities (sweet, umami, bitter), the loss of other

taste qualities in a limited number of subjects suggests that multiple mechanisms may underlie

the sensory loss.

Limitations & Conclusions

The study reported here is a retrospective clinical study of post-Covid patients and like other

case series studies has no control group although we offer a comparison to a pre-pandemic

dataset on gustatory phenotypes in healthy individuals (Doty *et al.* 2025). Another limitation is

that our study includes only a single time point measurement of taste function in a limited

number of subjects. This limits the statistical power of the study, but nonetheless we do find

highly significant correlated results even in our modest number of subjects. Furthermore, the

study relies on self-reported accounts of taste loss coinciding with COVID-19 onset, but baseline

taste function data before infection was not possible perhaps reducing the generalizability of the

findings.

Despite these apparent limitations, our study presents important new quantitative

psychophysical data showing significant disruption in detection of specific taste qualities in a

subset of subjects reporting long-term post Covid taste dysfunction. The taste qualities most

impacted are those detected by Type II taste cells which utilize a PLCβ2 dependent signaling

cascade for transduction of sweet, umami and bitter qualities. Importantly, our study also shows a

correlated decrease in mRNA levels for two elements of the Type II cell transduction cascade for

these qualities –- the first evidence for correlative changes in molecular parameters tracking

psychophysical taste functions. Finally, despite the well-documented recovery of post-Covid

taste function in most patients, our study offers quantitative evidence for prolonged disruption of

taste abilities in a small subset of post-Covid subjects.

568
569

ACKNOWLEDGEMENTS

This work was supported by grants 3R01DC014728-05S1 to TEF awarded by the National Institute on Deafness and Other Communication Disorders of the National Institutes of Health, by Åke Wiberg Stiftelsen grant 2023 to M.K. Governmental funding of clinical research within the Uppsala university Hospital (ALF), and by internal funding of the Department of Animal Biosciences, Swedish University of Agricultural Sciences.

AUTHOR CONTRIBUTIONS

G. H. and T.E.F. conceived this research and along with G.L.and G.A. designed the research plan. G.H. & G.L. wrote the ethical proposal to Etikprövningsmyndigheten. H.M. provided subjects, performed the WETT tests and took the biopsies. M.A.K. provided subjects and with A.A. assisted in analysis of the subject population. T.E.F. and M.L. designed and performed the histopathology and data analysis. G.A. and T.V. designed and analyzed the quantitative PCR experiments. T.V. performed RNA extraction and quantitative PCR experiments. T.E.F. performed statistical analysis of the quantitative data and compared results to those in the Doty et al (2015) report. G.L., G.H., G.A., H.M., and T.E.F. provided funding for the project. G.H., T.E.F., G.A., and M.A.K. prepared the manuscript and all authors have discussed and revised the manuscript.

Use of AI:

This study utilized Microsoft Copilot (GPT-4) accessed through a secure university-managed Microsoft Teams environment. The tool was utilized in the process of editing and checking calculations but was not used to generate any original content. All outputs were reviewed by the lead author.

Competing interests

The authors declare no competing financial interests.

Data Availability Statement: The data underlying this article will be shared on reasonable request to the corresponding authors.

599   **REFERENCES**
600
Bartel DL, Sullivan SL, Lavoie EG, Sevigny J, Finger TE. 2006. Nucleoside triphosphate

diphosphohydrolase-2 is the ecto-ATPase of type I cells in taste buds. The Journal of comparative

neurology 497: 1–12.

Boldes T, Ritter A, Soudry E, Diker D, Reifen E, Yosefof E. 2024. The long-term effect of

COVID-19 infection on olfaction and taste; a prospective analysis. Eur Arch Otorhinolaryngol

281: 6001–6007.

Boscolo-Rizzo P, Hummel T, Invitto S, Spinato G, Tomasoni M, Emanuelli E, Tofanelli M,

Cavicchia A, Grill V, Vaira LA, Lechien JR, Borsetto D, Polesel J, Dibattista M, Menini A,

Hopkins C, Tirelli G. 2023a. Psychophysical assessment of olfactory and gustatory function in

post-mild COVID-19 patients: A matched case-control study with 2-year follow-up. International

forum of allergy & rhinology 13: 1864–1875.

Boscolo-Rizzo P, Spinato G, Hopkins C, Marzolino R, Cavicchia A, Zucchini S, Borsetto D,

Lechien JR, Vaira LA, Tirelli G. 2023b. Evaluating long-term smell or taste dysfunction in

mildly symptomatic COVID-19 patients: a 3-year follow-up study. Eur Arch Otorhinolaryngol

280: 5625–5630.

Castillo D, Seidel K, Salcedo E, Ahn C, de Sauvage FJ, Klein OD, Barlow LA. 2014. Induction

of ectopic taste buds by SHH reveals the competency and plasticity of adult lingual epithelium.

Development 141: 2993–3002.

Chang RB, Waters H, Liman ER. 2010. A proton current drives action potentials in genetically

identified sour taste cells. Proceedings of the National Academy of Sciences of the United States

of America 107: 22320–22325.

Chen M, Reed RR, Lane AP. 2019. Chronic Inflammation Directs an Olfactory Stem Cell

Functional Switch from Neuroregeneration to Immune Defense. Cell Stem Cell 25: 501–

513.e505.

Chen M, Shen W, Rowan NR, Kulaga H, Hillel A, Ramanathan M, Jr., Lane AP. 2020. Elevated

ACE-2 expression in the olfactory neuroepithelium: implications for anosmia and upper

respiratory SARS-CoV-2 entry and replication. Eur Respir J 56.

Cheng LHH, Robinson PP. 1991. The distribution of fungiform papillae and taste buds on the

human tongue. Archives of oral biology 36: 583–589.

Cooper KW, Brann DH, Farruggia MC, Bhutani S, Pellegrino R, Tsukahara T, Weinreb C,

Joseph PV, Larson ED, Parma V, Albers MW, Barlow LA, Datta SR, Di Pizio A. 2020. COVID-

19 and the Chemical Senses: Supporting Players Take Center Stage. Neuron 107: 219–233.

Daia C, Scheau C, Neagu G, Andone I, Spanu A, Popescu C, Stoica SI, Verenca MC, Onose G.

2021. Nerve conduction study and electromyography findings in patients recovering from Covid-

19 - Case report. Int J Infect Dis 103: 420–422.

Donadoni M, Kaminski R, Liao S, Al Janabi S, Margolskee RF, Ozdener MH, Sariyer IK. 2023.

Coronavirus infection in chemosensory cells. J Neurovirol 29: 35–44.

Doty RL, Sharetts R, Moein ST. 2025. Self-Administered Taste Testing Without Water:

Normative Data for the 53-Item Waterless Empirical Taste Test (WETT). International forum of

allergy & rhinology 15: 934–943.

Doty RL, Wylie C, Potter M. 2021. Validation of the Waterless Empirical Taste Test

(WETT((R))). Behav Res Methods 53: 864–873.

Doyle ME, Appleton A, Liu QR, Yao Q, Mazucanti CH, Egan JM. 2021. Human Type II Taste

Cells Express Angiotensin-Converting Enzyme 2 and Are Infected by Severe Acute Respiratory

Syndrome Coronavirus 2 (SARS-CoV-2). Am J Pathol 191: 1511–1519.

Ely EW, Brown LM, Fineberg HV, National Academies of Sciences E, Medicine Committee on

Examining the Working Definition for Long C. 2024. Long Covid Defined. N Engl J Med 391:

1746–1753.

Farbman AI. 1965. Fine structure of the taste bud. Journal of Ultrastructure Research 12: 328–

350.

Farbman AI. 1980. Renewal of taste bud cells in rat circumvallate papillae. Cell & Tissue

Kinetics 13: 349–357.

Farbman AI, Guagliardo N, Sollars SI, Hill DL. 2004. Each sensory nerve arising from the

geniculate ganglion expresses a unique fingerprint of neurotrophin and neurotrophin receptor

genes. J Neurosci Res 78: 659–667.

Fernandes P, Ferreira F, Morais OM, Ramos CMT, Fernandes EMR, Rocha S, Rocha RJA,

Monteiro VJP, Vilar PSG, Romao AM, Alves MRA. 2021. Performance of saliva as a specimen

to detect SARS-CoV-2. J Clin Virol 142: 104913.

Finlay JB, Brann DH, Abi Hachem R, Jang DW, Oliva AD, Ko T, Gupta R, Wellford SA,

Moseman EA, Jang SS, Yan CH, Matsunami H, Tsukahara T, Datta SR, Goldstein BJ. 2022.

Persistent post-COVID-19 smell loss is associated with immune cell infiltration and altered gene

expression in olfactory epithelium. Sci Transl Med 14: eadd0484.

Goodall BL, LeBlanc JJ, Hatchette TF, Barrett L, Patriquin G. 2022. Investigating the Sensitivity

of Nasal or Throat Swabs: Combination of Both Swabs Increases the Sensitivity of SARS-CoV-2

Rapid Antigen Tests. Microbiol Spectr 10: e0021722.

Hannum ME, Koch RJ, Ramirez VA, Marks SS, Toskala AK, Herriman RD, Lin C, Joseph PV,

Reed DR. 2023. Taste loss as a distinct symptom of COVID-19: a systematic review and meta-

analysis. Chem Senses 48: bjad043.

Hellekant G. 2024. Neuroscience of taste: unlocking the human taste code. BMC Neurosci 25:

19.

Hellekant G, Kasahara Y, Farbman AI, Harada S, Hård af Segerstad C. 1987. Regeneration

ability of fungiform papillae and taste-buds in rats. Chemical senses 12: 459–465.

High B, Finger TE. 2025. Structural comparisons of human and mouse fungiform taste buds.

Chem Senses 50: bjaf001.

Ho J, Tumkaya T, Aryal S, Choi H, Claridge-Chang A. 2019. Moving beyond P values: data

analysis with estimation graphics. Nat Methods 16: 565–566.

Huang N, Perez P, Kato T, Mikami Y, Okuda K, Gilmore RC, Conde CD, Gasmi B, Stein S,

Beach M, Pelayo E, Maldonado JO, Lafont BA, Jang SI, Nasir N, Padilla RJ, Murrah VA, Maile

R, Lovell W, Wallet SM, Bowman NM, Meinig SL, Wolfgang MC, Choudhury SN, Novotny M,

Aevermann BD, Scheuermann RH, Cannon G, Anderson CW, Lee RE, Marchesan JT, Bush M,

Freire M, Kimple AJ, Herr DL, Rabin J, Grazioli A, Das S, French BN, Pranzatelli T, Chiorini

JA, Kleiner DE, Pittaluga S, Hewitt SM, Burbelo PD, Chertow D, Consortium NC-A, Oral HCA,

Craniofacial Biological N, Frank K, Lee J, Boucher RC, Teichmann SA, Warner BM, Byrd KM.

2021a. SARS-CoV-2 infection of the oral cavity and saliva. Nat Med 27: 892–903.

Huang N, Pérez P, Kato T, Mikami Y, Okuda K, Gilmore RC, Conde CD, Gasmi B, Stein S,

Beach M, Pelayo E, Maldonado JO, Lafont BA, Jang SI, Nasir N, Padilla RJ, Murrah VA, Maile

R, Lovell W, Wallet SM, Bowman NM, Meinig SL, Wolfgang MC, Choudhury SN, Novotny M,

Aevermann BD, Scheuermann RH, Cannon G, Anderson CW, Lee RE, Marchesan JT, Bush M,

Freire M, Kimple AJ, Herr DL, Rabin J, Grazioli A, Das S, French BN, Pranzatelli T, Chiorini

JA, Kleiner DE, Pittaluga S, Hewitt SM, Burbelo PD, Chertow D, Frank K, Lee J, Boucher RC,

Teichmann SA, Warner BM, Byrd KM. 2021b. SARS-CoV-2 infection of the oral cavity and

saliva. Nat Med 27: 892–903.

Janols H, Wadsten C, Forssell C, Raffeti E, Janson C, Zhou X, Kisiel MA. 2024. Enhancing EQ-

5D-5L Sensitivity in Capturing the Most Common Symptoms in Post-COVID-19 Patients: An

Exploratory Cross-Sectional Study with a Focus on Fatigue, Memory/Concentration Problems

and Dyspnea Dimensions. Int J Environ Res Public Health 21.

Karmous I, Sayed Khan A, Sahnoun I, Ben Othman R, Ben Jemaa H, Mahjoub F, Gamoudi A,

Douik El Gharbi L, Mestiri T, Khan NA, Jamoussi H. 2023. Loss of Sour Taste Is the Striking

Feature among Four Basic Taste Qualities in Tunisian COVID-19 Patients. J Clin Med 12.

Kisiel MA, Janols H, Nordqvist T, Bergquist J, Hagfeldt S, Malinovschi A, Svartengren M. 2022.

Predictors of post-COVID-19 and the impact of persistent symptoms in non-hospitalized patients

12 months after COVID-19, with a focus on work ability. Ups J Med Sci 127.

Kisiel MA, Lee S, Janols H, Faramarzi A. 2023a. Absenteeism Costs Due to COVID-19 and

Their Predictors in Non-Hospitalized Patients in Sweden: A Poisson Regression Analysis. Int J

Environ Res Public Health 20.

Kisiel MA, Lee S, Malmquist S, Rykatkin O, Holgert S, Janols H, Janson C, Zhou X. 2023b.

Clustering Analysis Identified Three Long COVID Phenotypes and Their Association with

General Health Status and Working Ability. J Clin Med 12.

Kisiel MA, Nordqvist T, Westman G, Svartengren M, Malinovschi A, Janols H. 2021. Patterns

and predictors of sick leave among Swedish non-hospitalized healthcare and residential care

workers with Covid-19 during the early phase of the pandemic. PLoS One 16: e0260652.

Klein H, Asseo K, Karni N, Benjamini Y, Nir-Paz R, Muszkat M, Israel S, Niv MY. 2021a.

Onset, duration and unresolved symptoms, including smell and taste changes, in mild COVID-19

infection: a cohort study in Israeli patients. Clin Microbiol Infect 27: 769–774.

Klein H, Karni N, Israel S, Gross M, Muszkat M, Niv MY. 2021b. Reversible Taste Loss in a

COVID-19 Patient With Preexisting Chronic Smell Impairment. J Investig Med High Impact

Case Rep 9: 2324709621990765.

Landis BN, Welge-Luessen A, Bramerson A, Bende M, Mueller CA, Nordin S, Hummel T. 2009.

"Taste Strips" - a rapid, lateralized, gustatory bedside identification test based on impregnated

filter papers. Journal of neurology 256: 242–248.

Lee H, Macpherson LJ, Parada CA, Zuker CS, Ryba NJP. 2017. Rewiring the taste system.

Nature 548: 330–333.

Lee S, Kisiel MA, Lindberg P, Wheelock AM, Olofsson A, Eriksson J, Bruchfeld J, Runold M,

Wahlstrom L, Malinovschi A, Janson C, Wachtler C, Carlsson AC. 2025. Using machine learning

involving diagnoses and medications as a risk prediction tool for post-acute sequelae of COVID-

19 (PASC) in primary care. BMC Med 23: 251.

Liman ER. 2020. Salty Taste: From Transduction to Transmitter Release, Hold the Calcium.

Neuron 106: 709–711.

Liman ER, Kinnamon SC. 2021. Sour taste: receptors, cells and circuits. Curr Opin Physiol 20:

8–15.

Lindemann B. 1996. Taste reception. Physiological reviews 76: 719–766.

Locatello LG, Bruno C, Maggiore G, Cilona M, Orlando P, Fancello G, Piccica M, Vellere I,

Lagi F, Trotta M, Gallo O. 2021. The prognostic role of IL-10 in non-severe COVID-19 with

chemosensory dysfunction. Cytokine 141: 155456.

Matsumoto I, Ohmoto M, Narukawa M, Yoshihara Y, Abe K. 2011. Skn-1a (Pou2f3) specifies

taste receptor cell lineage. Nat Neurosci 14: 685–687.

McFarland AJ, Yousuf MS, Shiers S, Price TJ. 2021. Neurobiology of SARS-CoV-2 interactions

with the peripheral nervous system: implications for COVID-19 and pain. Pain Rep 6: e885.

Miller IJ. 1986. Variation in human fungiform taste bud densities among regions and subjects.

Anatomical Record 216: 474–482.

Miller IJ, Jr. 1988. Human taste bud density across adult age groups. J Gerontol 43: B26–30.

Miller IJ, Jr. 1989. Variation in human taste bud density as a function of age. Ann N Y Acad Sci

561: 307–319.

Miller IJ, Jr., Reedy FE, Jr. 1990. Variations in human taste bud density and taste intensity

perception. Physiol. Behav. 47: 1213–1219.

Moayedi Y, Michlig S, Park M, Koch A, Lumpkin EA. 2021. Somatosensory innervation of

healthy human oral tissues. The Journal of comparative neurology 529: 3046–3061.

Naik S, Larsen SB, Gomez NC, Alaverdyan K, Sendoel A, Yuan S, Polak L, Kulukian A, Chai S,

Fuchs E. 2017. Inflammatory memory sensitizes skin epithelial stem cells to tissue damage.

Nature 550: 475–480.

O'Neill F, De Stefano G, Pridgeon M, Bhargava D, Marshall A, Marshall A, Frank B. 2023.

Trigeminal neuropathy presenting secondary to SARS-CoV-2 infection. Pain Rep 8: e1103.

Ogata T, Ohtubo Y. 2020. Quantitative Analysis of Taste Bud Cell Numbers in the Circumvallate

and Foliate Taste Buds of Mice. Chem Senses 45: 261–273.

Ohkuri T, Horio N, Stratford JM, Finger TE, Ninomiya Y. 2012. Residual chemoresponsiveness

to acids in the superior laryngeal nerve in "taste-blind" (P2X2/P2X3 double-KO) mice. Chem

Senses 37: 523–532.

Ohtubo Y, Yoshii K. 2011. Quantitative analysis of taste bud cell numbers in fungiform and soft

palate taste buds of mice. Brain Res 1367: 13–21.

Ota Y, Yumiya Y, Chimed-Ochir O, Hasegawa A, Yoshida T, Nagata T, Tanaka J, Ohge H,

Kuwabara M, Kubo T. 2025. Characteristics of patients with COVID-19 and smell and/or taste

disorders depending on different virus strains: a cross-sectional study in Hiroshima, Japan. BMJ

Open 15: e088377.

Otte MS, Bork ML, Zimmermann PH, Klußmann JP, Lüers JC. 2022. Patients with COVID-19-

associated olfactory impairment also show impaired trigeminal function. Auris Nasus Larynx 49:

147–151.

Peluso MJ, Hanson MR, Deeks SG. 2024a. Infection-associated chronic conditions: Why Long

Covid is our best chance to untangle Osler's web. Sci Transl Med 16: eado2101.

Peluso MJ, Ryder D, Flavell R, Wang Y, Levi J, LaFranchi BH, Deveau TM, Buck AM, Munter

SE, Asare KA, Aslam M, Koch W, Szabo G, Hoh R, Deswal M, Rodriguez A, Buitrago M, Tai

V, Shrestha U, Lu S, Goldberg SA, Dalhuisen T, Durstenfeld MS, Hsue PY, Kelly JD, Kumar N,

Martin JN, Gambir A, Somsouk M, Seo Y, Deeks SG, Laszik ZG, VanBrocklin HF, Henrich TJ.

2023. Multimodal Molecular Imaging Reveals Tissue-Based T Cell Activation and Viral RNA

Persistence for Up to 2 Years Following COVID-19. medRxiv.

Peluso MJ, Ryder D, Flavell RR, Wang Y, Levi J, LaFranchi BH, Deveau TM, Buck AM,

Munter SE, Asare KA, Aslam M, Koch W, Szabo G, Hoh R, Deswal M, Rodriguez AE, Buitrago

M, Tai V, Shrestha U, Lu S, Goldberg SA, Dalhuisen T, Vasquez JJ, Durstenfeld MS, Hsue PY,

Kelly JD, Kumar N, Martin JN, Gambhir A, Somsouk M, Seo Y, Deeks SG, Laszik ZG,

VanBrocklin HF, Henrich TJ. 2024b. Tissue-based T cell activation and viral RNA persist for up

to 2 years after SARS-CoV-2 infection. Sci Transl Med 16: eadk3295.

Perea-Martinez I, Nagai T, Chaudhari N. 2013. Functional cell types in taste buds have distinct

longevities. PloS one 8: e53399.

Reiter ER, Coelho DH, French E, Costanzo RM, Consortium NC. 2023. COVID-19-Associated

Chemosensory Loss Continues to Decline. Otolaryngol Head Neck Surg 169: 1386–1389.

Rodriguez YA, Roebber JK, Dvoryanchikov G, Makhoul V, Roper SD, Chaudhari N. 2021.

"Tripartite Synapses" in Taste Buds: A Role for Type I Glial-like Taste Cells. J Neurosci 41:

9860–9871.

Rogn A, Jensen JL, Iversen PO, Singh PB. 2024. Post-COVID-19 patients suffer from

chemosensory, trigeminal, and salivary dysfunctions. Scientific reports 14: 3455.

Schober P, Boer C, Schwarte LA. 2018. Correlation Coefficients: Appropriate Use and

Interpretation. Anesth Analg 126: 1763–1768.

Sharetts R, Moein ST, Khan R, Doty RL. 2024. Long-Term Taste and Smell Outcomes After

COVID-19. JAMA Netw Open 7: e247818.

Shi P, Zhang J, Yang H, Zhang YP. 2003. Adaptive diversification of bitter taste receptor genes

in Mammalian evolution. Mol Biol Evol 20: 805–814.

Singer-Cornelius T, Cornelius J, Oberle M, Metternich FU, Brockmeier SJ. 2021. Objective

gustatory and olfactory dysfunction in COVID-19 patients: a prospective cross-sectional study.

Eur Arch Otorhinolaryngol 278: 3325–3332.

Soriano JB, Murthy S, Marshall JC, Relan P, Diaz JV. 2022. A clinical case definition of post-

COVID-19 condition by a Delphi consensus. Lancet Infect Dis 22: e102–e107.

Stepien J, Pastuszak Z. 2023. Electroneurological changes in peripheral nerves in patients post-

COVID. J Neurophysiol 129: 392–398.

Wang J, Liu R, Ma H, Zhang W. 2023. The Pathogenesis of COVID-19-Related Taste Disorder

and Treatments. Journal of dental research 102: 1191–1198.

Wilson CE, Lasher RS, Salcedo E, Yang R, Dzowo Y, Kinnamon JC, Finger TE. 2025. Death in

the Taste Bud: Engulfment of Dying Taste Receptor Cells by Glial-Like Type I Cells. Glia 73:

1660–1672.

Wilson CE, Lasher RS, Yang R, Dzowo Y, Kinnamon JC, Finger TE. 2022. Taste Bud

Connectome: Implications for Taste Information Processing. J Neurosci 42: 804–816.

Xu Z, Wang W, Zhang D, Tam KW, Li Y, Chan DCC, Yang Z, Wong SYS. 2024. Excess risks

of long COVID symptoms compared with identical symptoms in the general population: A

systematic review and meta-analysis of studies with control groups. J Glob Health 14: 05022.

Yang R, Dzowo YK, Wilson CE, Russell RL, Kidd GJ, Salcedo E, Lasher RS, Kinnamon JC,

Finger TE. 2020. Three-dimensional reconstructions of mouse circumvallate taste buds using

serial blockface scanning electron microscopy: I. Cell types and the apical region of the taste bud.

The Journal of comparative neurology 528: 756–771.

Yao Q, Doyle ME, Liu QR, Appleton A, O'Connell JF, Weng NP, Egan JM. 2023. Long-Term

Dysfunction of Taste Papillae in SARS-CoV-2. NEJM Evid 2.

Yu T, Wilson CE, Stratford JM, Finger TE. 2020. Genetic Deletion of TrpV1 and TrpA1 Does

Not Alter Avoidance of or Patterns of Brainstem Activation to Citric Acid in Mice. Chem Senses

45: 573–579.

Zazhytska M, Kodra A, Hoagland DA, Frere J, Fullard JF, Shayya H, McArthur NG, Moeller R,

Uhl S, Omer AD, Gottesman ME, Firestein S, Gong Q, Canoll PD, Goldman JE, Roussos P,

tenOever BR, Jonathan BO, Lomvardas S. 2022. Non-cell-autonomous disruption of nuclear

architecture as a potential cause of COVID-19-induced anosmia. Cell 185: 1052–1064 e1012.

Zhang J, Jin H, Zhang W, Ding C, O'Keeffe S, Ye M, Zuker CS. 2019. Sour Sensing from the

Tongue to the Brain. Cell 179: 392–402 e315.

Zhang Y, Hoon MA, Chandrashekar J, Mueller KL, Cook B, Wu D, Zuker CS, Ryba NJ. 2003.

Coding of sweet, bitter, and umami tastes: different receptor cells sharing similar signaling

pathways. Cell 112: 293–301.

Zhao GQ, Zhang Y, Hoon MA, Chandrashekar J, Erlenbach I, Ryba NJ, Zuker CS. 2003. The

receptors for mammalian sweet and umami taste. Cell 115: 255–266.

834
835

836 SUPPLEMENTARY FIGURE
837
838

839 Supplementary Figure 1: qPCR for Covid mRNA showing positive controls. No samples from
840 any subjects show evidence of reaction for Sars-CoV2 virus.
841
842
843

845 Supplementary Table 1: WETT taste test values shown as a percentile values as normalized to the
846 data in Doty et al 2025.

Subject Number	Sweet	PLC Tastes Umami	Bitter	Sour	Salty
SWE 17	99	99	99	99	99
SWE 1	49	55	8	62	58
SWE 2	28	10	13	99	99
SWE 3	74	11	36	84	43
SWE 4	2	99	68	62	75
SWE 5	99	70	99	99	76
SWE 6	79	38	38	44	23
SWE 10	60	33	99	84	43
SWE 16	58	36	71	99	76
SWE 7	4	10	7	71	43
SWE 8	74	4	7	71	81
SWE 9	7	17	10	99	99
SWE 12	26	53	71	81	19
SWE 13	7	30	10	55	20
SWE 14	18	99	47	99	60
SWE 15	41	92	29	35	19
SWE 18	87	12	50	71	2
SWE 19	79	72	99	62	5
SWE 20	77	35	75	99	99
SWE 21	42	23	69	90	29
SWE 22	7	70	5	68	3
SWE 23	33	72	81	62	39
SWE 24	65	17	10	37	35
SWE 25	14	70	10	2	81
SWE 26	57	10	59	26	71
SWE 27	3	11	42	11	57
SWE 28	65	55	81	78	11
SWE 29	3	12	5	1	1
Values≤10%	6	2	8	1	4