Responders vs. non-responders to mesenchymal stromal cells in knee osteoarthritis patients: mechanistic correlates of donor cell attributes and patient features

Kevin P. Robb,[1,2,3] Razieh Rabani,[1,2] Shabana Vohra,[1,2] Antonietta Fazio,[1,2] Shoba Singh[1,2], Jaskarndip Chahal,[1,2,4] Julie Audet[3], Anthony V. Perruccio,[1,2,4,6] Rajiv Gandhi,[1,4] Sowmya Viswanathan*[1,2,3,7]

[1]Osteoarthritis Research Program, Division of Orthopedic Surgery, Schroeder Arthritis Institute, University Health Network, Toronto, Canada
[2]Krembil Research Institute, University Health Network, Toronto, Canada
[3]Institute of Biomedical Engineering, University of Toronto, Toronto, Canada
[4]Department of Surgery, University of Toronto, Toronto, Canada
[5]Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, Ontario, Canada
[6]Institute of Health Policy, Management and Evaluation, Dalla Lana School of Public Health, University of Toronto, Toronto, Canada
[7]Department of Medicine, University of Toronto, Toronto, Canada

*Corresponding author:
Sowmya Viswanathan, PhD
Krembil Research Institute
60 Leonard Ave
Toronto, Ontario, Canada
M5T 0S8
Phone: 416-634-7239
Email: sowmya.viswanathan@uhn.ca

Author contributions:

Kevin P. Robb: Conception & design, collection and/or assembly of data, data analysis and interpretation, manuscript writing, final approval of manuscript
Razieh Rabani: Conception & design, collection and/or assembly of data, data analysis and interpretation
Shabana Vohra: data analysis and interpretation
Antonietta Fazio: administrative support, collection and/or assembly of data
Shoba Singh: administrative support, collection and/or assembly of data
Jaskarndip Chahal: collection and/or assembly of data, provision of study materials or patients
Anthony V. Perruccio: data analysis and interpretation
Rajiv Gandhi: financial support, provision of study materials or patients
Sowmya Viswanathan: Conception & design, manuscript writing, final approval of manuscript

Disclaimers: Sowmya Viswanathan declares 60% ownership of Regulatory Cell Therapy Consultants Inc., which does not conflict with this paper in any way. Sowmya Viswanathan,

47 Razieh Rabani, and Kevin P Robb are co-inventors on patent applications filed with UHN
48 based on this work.
49
50 Brief acknowledgements:
51 This research was funded by the Canadian Institutes of Health Research (CIHR), Natural
52 Sciences and Engineering Research Council of Canada (NSERC), Toronto General and
53 Western Hospital Foundation, and the Arthritis Society.
54
55 **Keywords:** responders vs. non-responders, stromal cells, osteoarthritis, cell- and tissue-based
56 therapy, immunomodulation, microRNAs, biomarkers
57

ABSTRACT

Background: Despite recent regulatory approvals for mesenchymal stromal cells (MSCs), including two products for knee and degenerative arthritis globally, MSCs have shown mixed efficacy in advanced clinical trials of knee osteoarthritis (KOA). Analysis of contributing factors to the underlying heterogeneity in patient responsiveness is needed to comprehend these mixed results and to advance clinical translation.

Methods: We provide the basis for a novel algorithm that correlates with responder (defined by stringent international consensus criteria) vs. non-responder classifications using i) the immunomodulatory baseline features of bone marrow-derived MSCs (MSC(M)) batches, and ii) differential KOA baseline clinical features, local and systemic biomarkers. We use our previous clinical trial with twelve batches of MSC(M) as a test dataset and demonstrate correlations with new longitudinal medium-and long-term clinical data.

Results: We report that KOA clinical responsiveness to MSC treatment is dictated by a combination of the immunomodulatory fitness of the donor MSC(M) batches, clinical phenotype, baseline immune cell profile, and biomarker profile. Specifically, MSC(M) in KOA clinical responders have reduced expression of angiogenic genes, *THBS1*, *CXCL8*, and *ANGPT1* and higher expression of growth factor *CCN2* and distinct microRNA signatures, as well as higher *in vitro* ability to modulate monocytes that significantly distinguish them from MSC(M) from non-responders. Clinical responders had greater OA baseline severity, reduced physical activity; lower local baseline levels of CCL2, VEGFA, HGF; higher local baseline of resistin; and putatively lower frequencies of joint-specific intermediate monocytes/macrophages and activated CD4 T cells. Leave-out-one sensitivity analysis confirmed these findings mitigating small sample size concerns. Importantly, predicted responder status at 12 and 24 months was maintained out to 8-10 years in 5/7 responders and 3/3 non-responders, confirming the utility of our approach.

Conclusions: We provide a systematic, quantitative, evidence-based method for analyzing heterogeneity in KOA patient responsiveness to MSC treatments, dually incorporating batch/donor and patient factors. Our combinatorial analytical method provides a powerful framework for parsing heterogeneity in KOA patient responsiveness to MSC therapies and defining critical quality attributes that can reliably identify potent batches and/or donors of MSCs.

Significance statement: Using our data-rich clinical trial with 12-, 24- months and 8-10-year

longitudinal follow-up data, we assess donor/batch cell attributes and patient factors that

drive clinical efficacy. Bone Marrow MSC(M) from clinical responders exhibited lower

angiogenic gene expression, greater ability to modulate monocytes and differential

microRNA expression. Responsive KOA patients had more severe KOA, lower physical

activity levels, and lower angiogenic proteins in their joints at baseline. Our quantitative

analytical framework can be used prospectively in designing future clinical trials and

retrospectively in analyzing MSC effectiveness in KOA patients.

INTRODUCTION

Knee osteoarthritis (KOA) affects 16% of adults over 40 years of age worldwide [1], with no available disease-modifying drugs. Pathophysiological mechanisms of KOA are heterogeneous and include abnormal mechanical loading, metabolic, and inflammatory factors that fluctuate temporally [2]. Culture-expanded mesenchymal stromal cells (MSCs) are being investigated for their multimodal mechanisms of action, including chondroprotective/cartilage reparative, anti-fibrotic, and immunomodulatory functions [3,4]However, there has been mixed clinical efficacy reported including Mautner *et al*. who showed umbilical cord-derived MSCs were equivalent to corticosteroid injections in terms of improvements in pain or cartilage magnetic resonance imaging scores [5]. Other reports in smaller clinical trials [6–9], meta-analyses [10,11], and randomized controlled trials [12,13] however showed symptom and structural modification. These conflicting data raise questions: do MSCs work in KOA? Can we understand the differences between responder vs. non-responders to help deliver better MSC treatments to KOA patients?

Heterogeneous responsiveness to MSC treatments is due to an interplay of both basal potency fitness of donor MSC batches and basal patient disease status. Heterogeneity exists among MSC investigational products due to variations in donors [4,14] and manufacturing parameters [15,16]. The International Society for Cell & Gene Therapy (ISCT) has emphasized a need for well-defined critical quality attributes (CQAs) that are sensitive, quantitative, relevant to mechanism of action, and validated for specific disease indications [17,18] as we have also shown [19].

Using reported [8],and new follow-up clinical results from a phase I/IIa dose-escalation clinical trial, we conduct an in-depth combined analyses of donor MSC batch and baseline/follow up KOA patient characterization to provide a systematic algorithm to analyze heterogenous KOA patient responsiveness to MSC treatments. We use our previously reported gene panel that correlated to *in vitro* adipose tissue-derived MSC (MSC(AT)) functional activity [20], and show these discriminate MSC derived from marrow tissue, MSC(M), from responder vs. non-responder KOA patients. MicroRNAs, important mediators of MSC mechanisms of action [21] further discriminate MSC(M) from responders vs. non-responders.

In addition to MSC donor/batch heterogeneity, there is heterogeneity in KOA patients. KOA is now recognized not as a single disease but a constellation of clinical phenotypes (patient's clinical disease presentation [13]), including pain sensitization, metabolic syndrome, mechanical loading, and joint alignment [23,24]. KOA patients exhibit divergent molecular endotypes (molecular drivers of the disease) that may be characterized using local and/or systemic biomarkers [25]. The relationships between clinical phenotype and molecular endotype are complex and exhibit distinct or overlapping features, as well as temporal fluctuations [23,25]. This heterogeneity is a key contributor to the variable responses to MSC treatments. A systematic approach that incorporates baseline clinical phenotype and molecular disease endotype features that drive patient responsiveness are thus needed to improve MSC treatment outcomes.

MATERIALS AND METHODS

Clinical trial and patient data

KOA patients (Kellgren Lawrence (KL) grade 3-4) were recruited to this nonrandomized, open□label, dose□escalation, Health Canada-authorized phase I/IIa clinical trial as reported previously [8] (ClinicalTrials.gov Identifier: NCT02351011). Ethics approvals including for long-term follow up was granted by the UHN Research Ethics Board (REB 14–7909, REB 20-5066, REB 24-5463). Eligibility criteria and patient demographic information were previously provided [8]. Patients received a single intra-articular autologous MSC(M) injection at doses of 1 x 10^6, 10 x 10^6, and 50 x 10^6 cells (N=4 patients/dose group). Complete clinical follow-up was performed 24 months after MSC(M) injection for all patients. Additional follow-up, at 8-10 years was obtained for 10/12 patients; the remaining 2 patients were lost to follow-up; one patient (ID #3) could not be contacted and another (ID #10) underwent a total knee replacement in 2025 and declined to participate in the survey. Calculation of patient-reported outcome measures (PROMs), including the inverted KOOS and the Marx Activity Rating Scale (MARS) are described in **Suppl. File 1**.

Patient clinical responder status was based on the Outcome Measures in Rheumatology (OMERACT) and Osteoarthritis Research Society International (OARSI) Standing Committee for Clinical Trials Response Criteria Initiative (OMERACT-OARSI criteria). The OMERACT-OARSI criteria offers a rigorous responder classification system developed using data from large, powered placebo-controlled drug trials for hip and KOA [26]. These criteria categorize patients as responders according to the following: i) improvement in pain

or in function ≥50% and absolute change ≥ 20; OR, ii) improvement (≥20% and absolute

change ≥ 10) in at least two of the following: a) pain, b) function, c) patient global

assessment; the latter was not measured in our trial. Given the limited sample size and lack of

patient global assessment metrics in our previous clinical trial, we categorized patient

responder status using the twelve-month follow-up data based on changes in KOOS function

(activities of daily living (ADL) subscale) and/or KOOS Pain only, applying OMERACT-

OARSI criteria thresholds for function and pain scores of delta value ≥10 AND percentage

change ≥20% [26]. The validated KOOS instrument measures responses to interventions in

KOA with high test-retest reliability and improved sensitivity in pain and function scoring

relative to the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC)

scoring system [26,27].

MSC(M) isolation and culture

Biobanked MSC(M) were retained as individual batches by consent of KOA patients and

stored in liquid nitrogen. For microRNA-seq and gene expression analysis, MSC(M) were

thawed in three batches at P3 or P4 and plated directly on 24-well plates as described below.

For monocyte polarization experiments, MSC(M) were thawed in three batches at P4 or P5,

expanded for one passage, and then plated for the assay as described below. MSC(M) culture

was performed using the same expansion protocol and reagents used in the clinical trial [8] to

minimize any batch processing parameters variations.

MSC(M) gene expression analysis

MSC(M) were seeded on 24-well plates at 30,000 cells/well and licensed with a cocktail of

pro-inflammatory cytokines (IFNγ: 30 ng/mL; TNFα: 10 ng/mL; and IL-1β: 5 ng/mL), as

previously described [20]. After 24 hours of culture, Trizol-chloroform extraction was

performed to isolate RNA, and cDNA was generated using SuperScript™ IV VILO™ Master

Mix (Invitrogen, Waltham, USA). FastStart Universal SYBR Green Master Mix (Roche,

Basel, Switzerland) and custom primers (**Table S1**) were used for qPCR on a QuantStudio™

5 system (ThermoFisher). Results were normalized (delta Ct; ΔCt) using reference genes

(*GAPDH*, *GUSB*) and analyzed as negative ΔCt such that higher (more positive values)

indicate a higher level of expression.

In vitro **monocyte/macrophage polarization**

Indirect co-culture of unmatched human CD14$^+$ peripheral blood-derived monocytes

(STEMCELL Technologies, Vancouver, Canada) with biobanked MSC(M) was performed as

previously described [20]. Co-culture was performed for 48 h and lipopolysaccharide (LPS) was spiked at a concentration of 2.5 ng/mL. After 4 h, monocytes were collected and RNA extraction, cDNA synthesis, and qPCR was performed as described above. Results were normalized ($\Delta\Delta$Ct) using reference genes (*GAPDH*, *B2M*, *TBP*) and presented as fold-change relative to monocytes cultured without MSC(M) (solo condition).

microRNA-sequencing on MSC(M) samples

Biobanked MSC(M) were thawed and plated on 24-well plates at 50,000 cells/well and cultured for 72 h in proliferation medium, supplemented with or without synovial fluid. Synovial fluid was collected from unrelated late-stage KOA patients (KL grade 3-4, REB #14–7483), pooled from eight patients, and stored at -80°C until use, as previously described [28]. For treated wells, medium was supplemented with 30% (v/v) synovial fluid. After 24 h, RNA extraction was performed using a miRNeasy Mini Kit (Qiagen, Hilden, Germany) and stored at -30°C until analysis. Details on library preparation, microRNA-sequencing, and analysis are described in **Suppl. File 1**.

Biomarker measurements

Levels of curated biomarkers in serum (C1□2C collagen type I and II cleavage, C1-2C), plasma (hyaluronic acid, HA; cartilage oligomeric matrix protein, COMP), urine (C□telopeptide of type II collagen, CTX-II; collagen type II cleavage, C2C-HUSA), and synovial fluid (matrix metalloproteinases, MMP-3, MMP-9, MMP-13; tissue inhibitor of metalloproteinases 1, TIMP1; C□X□C chemokine motif ligands, CX3CL1, CXCL1; Chemokine C-C motif ligand 2 (CCL2); vascular endothelial growth factor A (VEGFA); IL12p40; interleukin(IL)-6; IL-8; hepatocyte growth factor (HGF); sCD14; sCD163; prostaglandin; adiponectin; adipsin; leptin; and resistin) collected at baseline prior to MSC(M) injection were measured in our previous study [8]. Updated analyses and experimental questions were applied to the dataset in the present work. Additional inflammatory biomarkers measured in biobanked patient serum and plasma collected at baseline were also evaluated by ELISA, with analytes specified in **Table S2**. Synovial fluid collection was feasible for only nine patients. Flow cytometry was performed on synovial fluid immune cell subpopulations in our previous study for a subset of five patients from whom sufficient synovial fluid volume could be obtained [8].

Statistical analyses

GraphPad Prism 6.0 (La Jolla, USA) and JMP Pro 17 (Cary, USA) software were used for statistical analyses and to create plots. Statistical tests are specified in the figure and table captions. Spearman's correlation was selected as a non-parametric statistical test to investigate associations between KOOS changes (delta or percentage change) and other variables, given that variables within our dataset violated assumptions of normality and homoscedasticity. P values derived from Spearman's correlation were not corrected for multiple comparisons given our exploratory analyses with a limited dataset, and requirements for normality to apply multiple comparison corrections. $P < 0.05$ was considered statistically significant.

RESULTS

Patient responder status

We systematically investigated multivariable factors affecting KOA patient responder status including 12-month patient-reported KOOS data previously reported [8]. We applied OMERACT-OARSI criteria [26] to determine responder status, with modifications as described in the Methods. Patients were classified as: i) Function-Pain Responders (exceeding thresholds in <u>both</u> ADL and Pain scores); ii) Function Responders (exceeding thresholds in ADL scores), iii) Pain Responders (exceeding thresholds in Pain scores); and iv) Non-Responders (exceeding thresholds in neither ADL nor Pain) (**Table 1**). Out of twelve study patients, we identified eight Responders and four Non-Responders. The eight Responders were sub-divided into six Function-Pain Responders (3/6 Function-Pain Responder patients self-identified as women); seven Function Responders (including the six Function-Pain Responders; 4/7 Function Responder patients self-identified as women); and seven Pain Responders (including the six Function-Pain Responders; 4/7 Pain Responder patients self-identified as women). The remaining four patients were classified as Non-Responders. While our sample size is limited, the binary responder/non-responder criteria are useful in determining factors affecting patient clinical responses. We include correlation analyses with all KOOS subscale scores to further supplement our analyses. We did not see dose-dependency and observed Function-Pain Responders across all three MSC(M) doses including the 1×10^6 dose (2/4 patients), 10×10^6 dose (1/4 patients), and the 50×10^6 dose (3/4 patients) (**Table 1**).

Responder vs. Non-Responder classification holds at 24-months and at longer 8–10-year follow-up

Patient follow-up data out to 24 months showed that patient responses were largely maintained (**Fig. 1**). Specifically, patients categorized at 12 months as Function-Pain Responders displayed significantly improved KOOS scores in all subscales at **both** 12- and 24-months follow-up, relative to baseline (**Fig. 1A**), supportive of persistent effects of autologous MSC(M) with a single injection. Patients categorized as Non-Responders at 12 months similarly showed no significant changes in each of the KOOS subscales or Overall KOOS at 12- and 24-months relative to baseline, indicating that the clinical phenotype features of this group neither improved nor worsened compared to baseline. No significant differences were observed when comparing outcomes at 12 versus 24 months for any KOOS subscale in both responder and non-responder patient groups.

Similar results were obtained when patients were categorized according to either Function-Responders (**Fig. 1B**) or Pain Responders (**Fig. S1A**); however, the improvements in Sports/Rec scores were observed only at the 12- and not the 24-month point relative to baseline for these groups.

We further analyzed patient follow-up at 8-10 years after MSC(M) injection, with KOOS scores obtained for 10 of 12 patients (**Fig. 1C, 1D**). Patients classified as Responders (at 12-months) maintained significant improvement in multiple KOOS subscales at 8-10 years follow-up (**Fig. S2**).

Of the eight 12-month responder patients (including Function-Pain Responders, Pain Responder, and Function Responder), seven patients maintained their responder status at 24 months. Of these seven patients, 5 maintained Responder status at 8-10 years, and one patient (who reported only that they underwent knee arthroplasty surgery) did not provide long-term KOOS data. Within this group of 12-month Responders, 8–10-year follow-up data showed that 2/8 patient received no additional interventions, 2/8 received injections including viscosupplementation or platelet-rich plasma, and 3/8 patients underwent knee arthroplasty surgery.

In terms of the four 12-month Non-Responder patients, all patients maintained Non-Responder status at 24 months, and one patient was lost to follow-up at the 8–10-year time point. The remaining three patients maintained Non-Responder status at 8-10 years, with 1/4 patients reporting no additional interventions and 2/4 patients reporting a knee injection for viscosupplementation.

MSC Batch Fitness Status

Gene panel discriminates between clinically effective and ineffective MSC(M) donor batches

Using biobanked MSC(M) samples cultured under pro-inflammatory licensing conditions, we measured expression of a putative immunomodulatory gene panel that we previously identified [20]. Gene expression was measured under pro-inflammatory licensing conditions to measure the immunomodulatory response of MSC(M), as has been widely demonstrated in the field [29]; MSC(M) injected intra-articularly would encounter synovial fluid, which contains pro-inflammatory mediators [30] that would similarly license the MSC(M).

Principal Component (PC) analysis revealed distinct clustering between MSC(M) donor batches labelled as Function-Pain Responders compared to Non-Responders along the PC1 axis (accounting for 29.3% of total variance), with higher expression of *CCN2* and lower expression of angiogenic markers *THBS1*, *CXCL8*, and *ANGPT1* contributing most strongly to higher PC1 scores (**Fig. 2A**, **Fig. S3A**). PC1 scores were significantly lower for Function-Pain Responders, Function Responders, and Pain Responders relative to Non-Responders (**Fig. 2B**, **Fig. S3B**), confirming the utility of the gene expression panel to distinguish MSC(M) derived from responders and non-responders. To evaluate model robustness, outlier analysis was performed by calculating T^2 statistic, identifying three outlier data points (two Responder, one Non-Responder); however, the PCA model returned similar results even with these data points removed with clear separation of responder/non-responder distributed across the expected eigenvectors (**Fig. S3C**).

Higher basal gene expression of the angiogenic gene, *ANGPT1* in 9 samples of MSC(M) strongly (Spearman's rho >0.68) and significantly negatively correlated to improvements in KOOS Pain, ADL, Sports/Rec, QOL, and Overall KOOS at 12 months, as well as to improvements in KOOS ADL at 24 months relative to baseline (**Table S3**). In contrast, higher basal levels of the mitogenic factor, *CCN2* in MSC(M) strongly (Spearman's rho >0.66) and significantly positively correlated to improvements in KOOS Pain, QOL, Other Symptoms, and Overall KOOS at 12 months relative to baseline. PC1 scores were indexed as a composite score for the basal immunomodulatory gene expression profile, and robust negative correlations were observed between PC1 scores and improvements in KOOS Symptoms, ADL, and Overall KOOS at both the 12- and 24-month time points relative to baseline, as well as KOOS QOL (12-month only) and KOOS Pain (24-month only) (**Fig. 2C**,

Table S3). We also investigated correlations between the PC1 score with changes in KOOS at 8-10 years follow-up and determined continued, significant negative correlations with KOOS ADL and Overall KOOS (**Table S3**).

To further evaluate the strength of these trends, we performed a leave-out-one sensitivity analysis in which correlations to the PC1 gene expression score were measured after omitting one patient at a time, focusing on changes in ADL, Pain, and Sports/Rec as key KOOS subscales (**Table S4**). For all Spearman's correlations to 12-month outcomes, the average p value across all leave-one-out correlations maintained significance. At 24 months, the correlation to percent change in ADL maintained significance, while the correlation to percent change in KOOS Pain was not significant. Previously significant Spearman's correlation with KOOS ADL at 8-10 years showed a reduced Spearman's rho value that was no longer significant. Taken together, these findings validate the utility and robustness of our previously identified gene panel [20] to discriminate clinical responsiveness to MSC(M).

Modulation of monocyte phenotype by donor MSC(M) *in vitro* correlates to patient responder status

We employed functional *in vitro* monocyte polarization read-outs using indirect co-culture of MSC(M) with mismatched human peripheral blood-derived monocytes to assess baseline MSC fitness as part of a matrix of CQAs (**Fig. 3A**). Discriminant analysis demonstrated distinct clustering and greater separation of monocyte phenotypic profiles when donor MSC(M) were classified as Function Responders versus Non-responders (**Fig. 3B**). Co-cultures of monocytes with Function Responder MSC(M) donor batches induced significantly upregulated expression of the inflammation-resolving markers *IL10* and *STAB1,* as well as reduced expression of *CD86* compared to Non-Responders (**Fig. 3B**). The immunosuppressive marker, *CD274*, was upregulated by MSC(M) donor batches classified as Non-Responders compared to Function Responders. Overall, MSC(M) donor batches classified as Function Responders modulated *in vitro* monocyte polarization toward a distinct and more pro-resolving phenotype. Discriminant analysis showed reduced separation when donor MSC(M) batches were classified according to Function-Pain Responder (**Fig. 3C**) or Pain Responder categorization (**Fig. S3D**) as indicated by greater misclassified data points and reduced Entropy R^2 values relative to the model using the Function Responder classification (**Table S5**).

Specific microRNAs distinguish responder donor MSC(M) batches

To probe unbiased baseline differences in responsive MSC(M) donor batches, we performed microRNA-Seq. PC analysis demonstrated that samples clustered by donor batch and responder group rather than by synovial fluid stimulation (**Fig. 4A**). We identified 33 significantly differentially expressed microRNAs between MSC(M) donor batches classified as Function Responders vs. Non-Responders after correction for multiple comparisons; 11 microRNAs were upregulated, and 22 microRNAs were downregulated in Function Responders (**Fig. 4B**, **Table S6**).

We focused our analysis on the microRNA profile of MSC(M) classified as Function Responders, conducting molecular interaction analysis on predicted gene targets for microRNAs that were significantly differentially expressed between Function Responder and Non-Responder MSC(M). Gene targets related to cell cycle (GTPases), phosphoinositide 3-kinase (PI3K), mitogen-activated protein kinase (MAPK), transforming growth factor beta (TGF-β), VEGF, SMAD proteins, apoptosis, and other pathways were identified (**Fig. 4C,D**; full list of pathways in **Suppl. File 2**). Gene ontology analysis identified biological processes linked to neuronal processes, histone modification, cell growth, ossification, Wnt signaling, and others (**Fig. 4E,F**; full list of pathways in **Suppl. File 2**). Taken together, our microRNA-sequencing data and analysis provide further characterization and potential mechanistic insights into MSC(M) therapeutic effects in KOA patients.

Baseline Patient/Disease Status Baseline clinical phenotype correlates with 12- and 24-month and 8–10-year improvements

We probed the relationship between baseline KOOS subscales with subsequent responder status (positive delta or percent change values indicate improvement) (**Fig. 5**). Patients classified as Function Responders displayed significantly elevated Overall KOOS and KOOS Sports/Rec at baseline compared to the Non-Responder group, supportive of more severe baseline clinical presentations in this group (**Fig. 5A**). Function-Pain Responders and Pain Responders displayed no significant differences in baseline relative to Non-Responders (**Fig. 5A and S1B**).

Next, we analyzed correlations between the baseline scores for all five KOOS subscales as well as Overall KOOS with changes in these scores at 12- and 24-months (**Table S7**). The baseline Sports/Rec scores positively and significantly correlated with improvements in: i) KOOS ADL at 12- and 24-month relative to baseline and in ii) KOOS ADL, iii) KOOS Symptoms, and iv) Overall KOOS at 24 months (**Fig. 5C**).

Given that the KOOS Sports/Rec subscale measures difficulty in activities associated with knee function [31], we further probed the relationship between the KOOS Sports/Rec score and physical activity using the MARS [32]. Correlation analyses demonstrated that higher baseline MARS scores (indicative of frequent physical activity) were significantly associated with lower (i.e., less severe) baseline KOOS Sports/Rec scores (**Fig. 5D**). Interestingly, a higher MARS score significantly correlated to lesser improvements (or worsening) in i) KOOS Symptoms, and ii) KOOS Pain at 12 months (**Fig. 5D**, **Table S8**). No significant correlations were observed for the 24-month timepoint.

Using only the baseline clinical phenotype features that correlated to significant improvements in KOOS at 12- or 24-months follow-up, we examined correlations of the baseline features to 8-10-year follow-up KOOS data. Only the baseline MARS score continued to have significant negative correlation with changes in KOOS Symptoms at 8-10 years follow-up (**Table S7, S8**).

Baseline synovial fluid biomarkers correlate to 3-, 12- and 24-month patient improvements

The levels of inflammatory, angiogenic, metabolic and cartilage extracellular matrix-associated biomarkers in synovial fluid were previously reported [8]; however, we analyzed correlations of biomarker levels at baseline with 12-, 24-, and 8–10-year KOOS improvements (**Tables S9, S10**).

Correlation analyses revealed several baseline synovial fluid biomarkers that significantly correlated with improvements in KOOS Sports/Rec scores at 12 and 24 months. We observed one significant **<u>positive</u>** correlation for an adipokine, resistin, with greater improvements (delta values) in Sports/Rec scores at 24 months relative to baseline (r_s of 0.72; **Fig. 6A**), while all other significant correlations for inflammatory, chemotactic, and angiogenic factors were **<u>negatively</u>** correlated with improvements in PROMs at follow-up time points. Specifically, lower levels of TIMP1 significantly negatively correlated to percentage changes in KOOS Sports/Rec scores at 24 months relative to baseline (r_s of -0.073). CCL2 significantly negatively correlated with changes in KOOS Sports/Rec scores at 24 months relative to baseline (r_s of -0.72, -0.85). Significant negative correlations were also observed between changes in KOOS Pain at 12 months relative to baseline with the levels of the pleiotropic HGF (r_s of -0.79, -0.71), as well as the potent angiogenic factor VEGFA (r_s of -0.81) **Fig. 6B**).

Baseline synovial fluid levels of soluble (s)CD14 and CCL2 significantly and negatively

correlated with improvements in KOOS Symptoms and Sports/Rec scores at 3 months

relative to baseline (**Fig. 6C**, **Tables S11, S12**); 3 months was the earliest timepoint when

patients displayed improvements in PROM [8]. The covariance matrix for local biomarkers

that showed significant correlations is reported in **Table S13**. Briefly, TIMP1, resistin, HGF

and VEGFA covary with each other, while CCL2 and sCD14 covary with each other,

indicative of multiple pathways that may influence MSC(M)-mediated clinical

responsiveness.

Using only the synovial fluid biomarkers that showed significant correlations to

improvements in KOOS at 12 or 24 months, we performed correlations to changes in KOOS

at 8-10 years follow-up but observed no significant correlations (**Table S9, S10**).

Additionally, we performed the leave-out-one sensitivity analysis in which 12- and 24-month

correlations to local biomarkers were measured, omitting one patient at a time (**Table S14**).

For correlations performed using 12-month outcomes, the average p value across all leave-

one-out correlations maintained significance for HGF with KOOS Pain (delta only but not

percent change) and for VEGF with KOOS Pain (percent change). At 24 months, the average

correlations between resistin and KOOS Sports/Rec (delta) as well as between CCL2 and

KOOS Sports/Rec (percent change) maintained significance. The average correlations

between CCL2 and KOOS Sports/Rec (delta) as well as between TIMP1 and KOOS

Sports/Rec (percent change) were not significant.

Baseline synovial fluid immune cell phenotype correlates with responder/non-responder
classification

We also analyzed synovial fluid immune cell subpopulations at baseline using data generated

in our previous study [8]. While this analysis was feasible for only five patients due to limited

synovial fluid volumes, our data demonstrates proof-of-concept for probing baseline levels of

$CD14^+CD16^+$ and $CD14^+CD163^+$ monocyte/macrophages (MΦs) (**Fig. S4A**) as well as

$CD3^+CD4^+CD69^+$ and $CD3^+CD4^+CD25^+$ T helper cells (**Fig. S4B**) as an important baseline

immune marker of patient disease status and molecular endotype; higher frequencies of MΦs

and activated T cells are present in non-responders at baseline, although not significant given

the limited sample size.

Specific baseline systemic biomarker levels correlate with 3- and 6-month but not 12- and 24-month patient improvements

We further measured levels of systemic inflammatory biomarkers (sCD163; C-reactive protein, CRP; S100A8/A9; tumor necrosis factor-alpha, TNFα; and interleukin (IL)-6) and systemic biomarkers of cartilage breakdown (serum C1-2C, plasma COMP and HA, and urine CTX-II and C2C-HUSA were previously measured [8]) in biobanked serum and plasma of all twelve patients at baseline (**Table S15, S16, S17**). Higher levels of baseline plasma TNFα significantly positively correlated to the percentage change in KOOS Pain and QOL improvement at 6 months (**Fig. S5A**, **Table S17**). On the other hand, baseline serum levels of the alarmin S100A8/A9 significantly negatively correlated to changes in KOOS QOL, and Sports/Rec at 3 months relative to baseline (**Fig. S5B**, **Table S17**). Systemic levels of biomarkers for cartilage breakdown showed no significant correlations to changes in KOOS at any time points. Systemic biomarker levels thus showed limited sensitivity with correlations to short-term PROM improvements only. Altogether, there was no discernible baseline profile of systemic biomarkers to prospectively identify responder KOA patients.

DISCUSSION

Statistically powered clinical trials using MSCs have reported conflicting results in KOA patients [5,12]. To address underlying factors that dictate responsiveness to MSCs, we analyzed our test clinical dataset [8] and demonstrated specific MSC(M) donor batch characteristics and responsive KOA patient features that correlated with improvements in PROMs at 12 and 24-months that were sustained out to 8-10 years, in 62.5% of responders and 100% of non-responders (including patients lost to follow-up). We anticipate that this combinatorial analysis will serve as an integrated algorithm, that can be tested in future prospective KOA clinical trials (and retrospectively on previous trial datasets) to i) screen potent donor/batches of MSC; ii) to identify responsive recipients. Our data shows that MSC(M) donor batches that are responsive have high basal immunomodulatory fitness as measured by reduced levels of angiogenic genes, higher monocyte modulatory activity and distinct signature of microRNAs. In parallel, KOA patients most responsive to MSC(M) donor batches had more severe baseline disease; reduced knee functionality associated with sports and recreation activities; reduced physical activity; reduced levels of soluble angiogenic factors in the local joint environment and trends pointing to lower baseline levels of intermediate monocyte/macrophages and activated T cells. We used a previous clinical trial [8], where we manufactured 12 distinct, patient-specific lots of MSC(M).

While we acknowledge the limited sample size, our clinical trial was data-rich with multiple

baseline biomarker and clinical readouts and offered an excellent test dataset to study batch

variations as we manufactured twelve patient-specific batches of autologous MSC(M).

Leave-out-one sensitivity analysis confirmed the robustness of correlations with MSC(M)

fitness genes panel and baseline synovial fluid biomarkers mitigating concerns regarding the

robustness of the correlations using a small sample size.

Additionally, we provide longitudinal 12-, 24-month and 8–10-year follow-up data and are

one of the first groups to report such long-term clinical data. Responders showed significant

improvements in PROMs at both 12- and 24- months relative to baseline and significant

sustained responder and non-responder status was maintained to 8-10 years. We note that 2/8

(25%) Responder and 2/4 (50%) Non-Responder patients received additional injections over

the intervening years, and 3/8 Responders received or were awaiting a total knee

replacement. Additionally, 3/8 Responders (37.5%) and 1/4 Non-Responders (25%) did not

receive additional interventions. The higher number of total knee replacements in the

Responder group may be related to differences in sample size (N=8 Responders vs N=4 Non-

Responders). Given the extended lag in the follow-up period, it is not possible to confirm if

the single MSC(M) injections delayed knee replacements to this 8–10-year follow-up

window. We also cannot rule out the effects of additional interventions that may have

contributed to the PROM data obtained beyond the study endpoint of one year in both

Responder and Non-Responder patient groups [5,12].

KOA patients classified as Function Responders showed greater correlation to MSC(M)

donor potency fitness and a more distinct baseline clinical phenotype (especially KOOS

Sports/Rec scores). This is interesting given that immunomodulatory therapies may be

expected to induce greater effects on patient pain [33]; however, MSCs act through

multimodal mechanisms and both pain and functional improvements are commonly

documented [34]. KOA patients with higher disease severity, especially reduced levels of

high-level knee functionality had greater improvements in daily functioning of the knee

(KOOS ADL), confirming inter-relationships between these KOOS ADL and Sports/Rec

subscales [31]. We also observed that patients with higher baseline MARS scores had lesser

improvements in pain and other symptoms, suggesting that patients who may have persisted

through pain in physical activity may have aggregated progression. [31]. The demographics

of the patient cohort study with an age range of 40-65 years, body mass index (BMI) <30,

relatively neutral knee alignment, and KL grade 3 or 4 KOA provide clues to the utility of the

KOOS Sports/Rec subscale [34–36]. Notably, our analysis demonstrating patients with more

severe KOA were more responsive to MSC(M) treatment aligns with findings in acute graft-

versus-host disease [37], myocardial infarction [38] and ulcerative colitis [39], where more

severe disease patients respond better to MSC treatments.

We focused on basal immunomodulatory fitness of MSC(M), especially their ability to

modulate monocytes, given their importance as a clinically relevant mechanism of action in

KOA [4,7,8]. Our previously identified immunomodulatory gene panel for MSC(AT) [20]

successfully distinguished MSC(M) donors classified under all categories of Responders

versus Non-Responders. These differences were robust; dimension-reduced PC1 scores

capturing multivariate gene expression profile were strongly and significantly correlated

inversely to patient improvements at 12-, 24-months and even 8–10-year follow-up and are

thus more reliable than the use of individual genes. Leave-out-one sensitivity analysis also

confirmed the robustness of correlations to dimension reduced PC1 scores at 12-months,

assuring the individual donors were not skewing the correlation values. Notably, this panel

includes predominantly angiogenic markers that were previously inversely correlated to

monocyte polarization toward inflammation-resolving phenotypes [20]. The data also

supports other work indicating inverse relationships between angiogenic and

immunomodulatory fitness of MSC(M) [40,41]. While the transcriptome of MSC(AT) and

MSC(M) are distinct [42], the putative immunomodulatory genes we identified for MSC(AT)

are valid for MSC(M) as well, speaking to their broad utility to serve as MSC CQAs. Our

immunomodulatory gene panel aligns with expression of another immunomodulator,

indoleamine 2,3-dioxygenase (IDO) that correlates with in autologous infrapatellar fat pad-

derived adipose tissue MSC(AT) functional improvements in KOA patients [7]. These data

support our hypothesis that MSCs with high basal immunomodulatory fitness produce greater

improvements in knee function. Nonetheless, the use of autologous MSCs in both studies

[7,8] adds a layer of complexity, given that systemic factors such as age, BMI, and other

health conditions were dually accounted for in both the basal fitness of the cells [43,44], as

well as baseline patient disease factors [45]; the interplay between these different factors is

likely to have contributed to the overall responsiveness.

MSC(M) derived from Responders modulated $CD14^+$ monocytes to more pro-resolving

subtypes *in vitro*. Monocytes represent major immune effector cells modulated by MSC(M)

injection in KOA [8,46]. While we showed preferential upregulation of pro-resolving CD14$^+$

monocyte phenotype we also observed upregulation of markers typically associated with pro-

inflammatory phenotypes. Reliance on dichotomized gene expression markers is not

indicative of the plurality of monocyte functionality [47]. Rabani *et al.* demonstrated that

induction of a mixed phenotype of monocytes may be important for MSC-mediated

microbicidal functions within a sepsis model [48]. We used LPS after co-culture with

MSC(M) to elicit stronger gene expression readouts from the CD14$^+$ monocytes; LPS is

present in synovial fluid of KOA patients and its levels correlate to increased disease severity

[49], justifying its inclusion in our assay. Nonetheless, the distinct polarizing ability of

Function Responder MSC(M) donors vs. Non-Responder donors offers another readout to

prospectively screen potent MSC donor/batches.

MSC(M) from Function or Function-Pain Responders had a significantly different microRNA

signature compared to Non-Responders. MicroRNAs are important epigenetic regulators of

MSC fitness [50] and can mediate multiple MSC mechanisms of action [51]. Pathways

related to inflammation, fibrosis, angiogenesis, and ossification, relevant to KOA

pathophysiology were identified. We also identified enrichment of pathways involved in cell

cycle, in line with RNA-sequencing and cell proliferation analyses performed on autologous

MSC(AT) used in KOA [52]. Surprisingly, our analysis identified enrichment of several

neuronal processes; however, given the role of innervation and neuronal signaling in joint

pain in KOA [53], these microRNAs may be implicated in mediating analgesic effects of

MSC(M). The functionality of the microRNAs require further study, but these differentially

expressed panels may also serve as additional prognosticating CQAs for identifying basally

fit MSC(M) in treating KOA patients.

Responders to MSC(M) injection displayed a mixed signature of higher baseline synovial

fluid levels of an adipokine (resistin), and lower levels of inflammatory, angiogenic, and

extracellular matrix-associated biomarkers, including TIMP1, CCL2, HGF, VEGFA, and

sCD14, accompanied by lower levels of intermediate CD14$^+$CD16$^+$ monocyte/macrophages

(Mφs). While increased levels of resistin in synovial fluid may be linked to higher levels of

inflammation in KOA [54], lower levels of HGF, TIMP1, VEGFA, CCL2, and sCD14

indicate lower levels of active local inflammation, immune cells and angiogenesis [55–58].

Low levels of sCD14 have been previously correlated to lower levels of activated neutrophils

and macrophages in KOA [56] and to lower levels of CD14$^+$CD16$^+$ intermediate synovial

fluid MΦs[46]. We have also previously reported reduction in the levels of CD14$^+$CD16$^+$

intermediate synovial fluid MΦs after MSC(M) treatment [8], corroborating other datasets

showing reduced MΦ levels post-MSC treatment [39,59]. However, synovial fluid collection

and local biomarker measurement was possible for only a subset of patients who had active

disease presentation, and thus likely had increased inflammation and greater joint pain

[60,61], suggestive of a selection bias of patients who had active synovitis. These KOA

patients thus likely represent an inflammatory phenotype with active disease presentation,

and responsiveness to MSC(M) treatment. There is emerging data showing distinct

immunophenotypic classification of KOA patients based on activated levels of MΦs and T

cells [62], in line with our preliminary data that needs further study.

There were only short-term significant, albeit mixed correlations between systemic

biomarkers such as TNFα (in plasma) and S100A8/A9 (in serum) which positively and

negatively correlated to PROM improvements, respectively. Higher circulating levels of both

TNFα and S100A8/A9 may correlate to greater disease severity and systemic inflammation

[54,56]. Interestingly, a similar mixed signature of systemic inflammatory biomarkers

measured at baseline has been reported in responders versus non-responders to intravenous

infusion of allogeneic Wharton's jelly-derived MSCs in rheumatoid arthritis patients [61].

Baseline serum CRP levels, an inflammatory biomarker in heart failure, chronic obstructive

pulmonary disease, and other inflammatory conditions [62] failed to correlate with

improvements in KOOS at all time points as did cartilage breakdown biomarkers, previously

shown to correlate with worsening KOA pain [63]. Overall, our analyses suggest that local

biomarkers provide improved sensitivity compared to systemic biomarkers [24,55].

Limitations include small sample size, which we addressed with various sensitivity analyses

and the absence of a control group, although effects were measured at multiple timepoints

including 12, 24 and 8-10 years, relative to baseline. Given that KOA is a progressive

condition [64], approximately 30% of patients display rapidly worsening of PROMs over two

years leading up to joint surgery [65]. Although we cannot rule out placebo effects, we saw

less than 30% of patients who reported worsening outcomes, potentially speaking to a

treatment effect. Indeed, analysis of Non-Responders using the rigorous OMERACT-OARSI

criteria [26] showed that even patients within this category did not show any significant

worsening in PROMs over two years and 8-10 years of follow-ups. We cannot however rule

out the effects of other interventions or the absence of intervention on the long-term PROM

scores, but do report that 62.5% of Responders maintained Responder status while 100% of the Non-Responders maintained Non-Responder status [5,12].

CONCLUSIONS

We have demonstrated that clinical batches of MSC(M) that resulted in positive clinical outcomes differ significantly in specific measurable attributes compared to MSC(M) batches that failed to improve pain or function. These quantitative, multivariate, and functionally relevant attributes can define CQAs for MSC(M) that can be evaluated to strategically design future KOA trials to ensure higher success rates of MSC injection. Furthermore, we correlated baseline clinical features and synovial fluid biomarkers to responder status in KOA patients after MSC(M) injection. These analyses may be applied retrospectively to recently published KOA clinical trials investigating MSC(M) [12] and umbilical cord-derived MSC [5] injections.

Acknowledgements

We thank Kim Perry, Mary Nasim, Tamara Wagner, Amanda Weston and the Orthopedic Surgery team for assistance in acquiring donor samples. Thanks to Dr. Mozhgan Rasti who performed qPCR for a subset of experiments, as well as Dr. Mohit Kapoor, Dr. Starlee Lively, Dr. Mehdi Layeghifard, and Pratibha Potla for assistance with microRNA-sequencing and analysis. Schematics for figures were created using BioRender.com.

This research was funded by Canadian Institutes of Health Research (CIHR) (PJT-166089) and Natural Sciences and Engineering Research Council of Canada (NSERC) (RGPIN-2018-05737) grants awarded to Sowmya Viswanathan. The work is in part supported by the Schroeder Arthritis Institute via the Toronto General and Western Hospital Foundation (University Health Network). Salary support for Kevin P. Robb was provided by a Natural Sciences and Engineering Research Council (NSERC) Canada Graduate Scholarship. Salary support for Razieh Rabani was provided by the Arthritis Society (TPF-19-0537). Anthony V. Perruccio is supported by an award from Arthritis Society Canada (STAR-20-0000000012).

Disclosure of Potential Conflicts of Interest

Kevin P. Robb declares current employment status with STEMCELL Technologies. Sowmya Viswanathan declares 60% ownership of Regulatory Cell Therapy Consultants Inc., which does not conflict with this paper in any way. Sowmya Viswanathan, Razieh Rabani, and Kevin P Robb are co-inventors on patent applications filed with UHN based on this work. The remaining authors declare they have no competing interests.

References

[1] Cui A, Li H, Wang D, Zhong J, Chen Y, Lu H. Global, regional prevalence, incidence and risk factors of knee osteoarthritis in population-based studies. EClinicalMedicine 2020;29–30. https://doi.org/10.1016/j.eclinm.2020.100587.

[2] Loeser RF, Goldring SR, Scanzello CR, Goldring MB. Osteoarthritis: a disease of the joint as an organ. Arthritis Rheum 2012;64:1697–707. https://doi.org/10.1002/art.34453.

[3] Zhang X, He J, Wang W. Progress in the use of mesenchymal stromal cells for osteoarthritis treatment. Cytotherapy 2021;23:459–70. https://doi.org/10.1016/j.jcyt.2021.01.008.

[4] Copp G, Robb KP, Viswanathan S. Culture-expanded mesenchymal stromal cell therapy: does it work in knee osteoarthritis? A pathway to clinical success. Cell Mol Immunol 2023. https://doi.org/10.1038/S41423-023-01020-1.

[5] Mautner K, Gottschalk M, Boden SD, Akard A, Bae WC, Black L, et al. Cell-based versus corticosteroid injections for knee pain in osteoarthritis: a randomized phase 3 trial. Nature Medicine 2023 2023:1–7. https://doi.org/10.1038/s41591-023-02632-w.

[6] Soler R, Orozco L, Munar A, Huguet M, López R, Vives J, et al. Final results of a phase I–II trial using ex vivo expanded autologous Mesenchymal Stromal Cells for the treatment of osteoarthritis of the knee confirming safety and suggesting cartilage regeneration. Knee 2016;23:647–54. https://doi.org/10.1016/j.knee.2015.08.013.

[7] Chen HH, Chen YC, Yu SN, Lai WL, Shen YS, Shen PC, et al. Infrapatellar fat pad-derived mesenchymal stromal cell product for treatment of knee osteoarthritis: a first-in-human study with evaluation of the potency marker. Cytotherapy 2022;24:72–85. https://doi.org/10.1016/j.jcyt.2021.08.006.

[8] Chahal J, Gómez☐Aristizábal A, Shestopaloff K, Bhatt S, Chaboureau A, Fazio A, et al. Bone Marrow Mesenchymal Stromal Cells in Patients with Osteoarthritis Results in Overall Improvement in Pain and Symptoms and Reduces Synovial Inflammation. Stem Cells Transl Med 2019;8:746–757. https://doi.org/10.1002/sctm.18-0183.

[9] Matas J, García C, Poblete D, Vernal R, Ortloff A, Luque-Campos N, et al. A Phase I Dose-Escalation Clinical Trial to Assess the Safety and Efficacy of Umbilical Cord-Derived Mesenchymal Stromal Cells in Knee Osteoarthritis. Stem Cells Transl Med 2024;13:193–203. https://doi.org/10.1093/STCLTM/SZAD088.

[10] Yubo M, Yanyan L, Li L, Tao S, Bo L, Lin C. Clinical efficacy and safety of mesenchymal stem cell transplantation for osteoarthritis treatment: A meta-analysis. PLoS One 2017;12:e0175449. https://doi.org/10.1371/journal.pone.0175449.

[11] Cui G-H, Wang YY, Li C-J, Shi C-H, Wang W-S. Efficacy of mesenchymal stem cells in treating patients with osteoarthritis of the knee: A meta-analysis. Exp Ther Med 2016;12:3390–400. https://doi.org/10.3892/etm.2016.3791.

[12] Gupta PK, Maheshwari S, Cherian JJ, Goni V, Sharma AK, Tripathy SK, et al. Efficacy and Safety of Stempeucel in Osteoarthritis of the Knee: A Phase 3 Randomized, Double-Blind, Multicenter, Placebo-Controlled Study. Am J Sports Med 2023;51:2254–66. https://doi.org/10.1177/03635465231180323.

[13] Pico OA, Espinoza F, Cádiz MI, Sossa CL, Becerra-Bayona SM, Salgado MCC, et al. Efficacy of a single dose of cryopreserved human umbilical cord mesenchymal stromal cells for the treatment of knee osteoarthritis:a randomized, controlled, double-blind pilot study. Cytotherapy 2024;0. https://doi.org/10.1016/J.JCYT.2024.09.005/ATTACHMENT/201E9836-BC53-4964-AEB9-F674366CBCFB/MMC2.DOCX.

[14] Vives J, Oliver-Vila I, Pla A. Quality compliance in the shift from cell transplantation to cell therapy in non-pharma environments. Cytotherapy 2015;17:1009–14. https://doi.org/10.1016/j.jcyt.2015.02.002.

[15] Siegel G, Kluba T, Hermanutz-Klein U, Bieback K, Northoff H, Schäfer R. Phenotype, donor age and gender affect function of human bone marrow-derived mesenchymal stromal cells. BMC Med 2013;11:146. https://doi.org/10.1186/1741-7015-11-146.

[16] Wilson A, Hodgson-Garms M, Frith JE, Genever P. Multiplicity of mesenchymal stromal cells: Finding the right route to therapy. Front Immunol 2019;10:1112. https://doi.org/10.3389/fimmu.2019.01112.

[17] Galipeau J, Krampera M, Barrett J, Dazzi F, Deans RJ, DeBruijn J, et al. International Society for Cellular Therapy perspective on immune functional assays for mesenchymal stromal cells as potency release criterion for advanced phase clinical trials. Cytotherapy 2016;18:151–9. https://doi.org/10.1016/j.jcyt.2015.11.008.

[18] Viswanathan S, Shi Y, Galipeau J, Krampera M, Leblanc K, Martin I, et al. Mesenchymal stem versus stromal cells: International Society for Cell & Gene Therapy (ISCT®) Mesenchymal Stromal Cell committee position statement on nomenclature. Cytotherapy 2019;21:1019–24. https://doi.org/10.1016/j.jcyt.2019.08.002.

[19] Viswanathan S, Galipeau J. Hallmarks of MSCs: Key quality attributes for pharmacology and clinical
use. Cell Stem Cell 2025;32:878–94. https://doi.org/10.1016/J.STEM.2025.05.008.
[20] Robb KP, Audet J, Gandhi R, Viswanathan S. Putative critical quality attribute matrix identifies
mesenchymal stromal cells with potent immunomodulatory and angiogenic "fitness" ranges in response
to culture process parameters. Front Immunol 2022;13. https://doi.org/10.3389/fimmu.2022.972095.
[21] Asgarpour K, Shojaei Z, Amiri F, Ai J, Mahjoubin-Tehran M, Ghasemi F, et al. Exosomal microRNAs
derived from mesenchymal stem cells: Cell-to-cell messages. Cell Communication and Signaling
2020;18. https://doi.org/10.1186/s12964-020-00650-6.
[22] Mantripragada VP, Csorba A, Bova W, Boehm C, Piuzzi NS, Bullen J, et al. Assessment of Clinical,
Tissue, and Cell-Level Metrics Identify Four Biologically Distinct Knee Osteoarthritis Patient
Phenotypes. Cartilage 2022;13. https://doi.org/10.1177/19476035221074003.
[23] Mobasheri A, Saarakkala S, Finnilä M, Karsdal MA, Bay-Jensen AC, van Spil WE. Recent advances in
understanding the phenotypes of osteoarthritis. F1000Res 2019;8.
https://doi.org/10.12688/f1000research.20575.1.
[24] Dell'Isola A, Allan R, Smith SL, Marreiros SSP, Steultjens M. Identification of clinical phenotypes in
knee osteoarthritis: a systematic review of the literature. BMC Musculoskelet Disord 2016;17:1–12.
https://doi.org/10.1186/s12891-016-1286-2.
[25] Mobasheri A, Bay-Jensen AC, Gualillo O, Larkin J, Levesque MC, Henrotin Y. Soluble biochemical
markers of osteoarthritis: Are we close to using them in clinical practice? Best Pract Res Clin
Rheumatol 2017;31:705–20. https://doi.org/10.1016/j.berh.2018.07.002.
[26] Pham T, van der Heijde D, Altman RD, Anderson JJ, Bellamy N, Hochberg M, et al. OMERACT-
OARSI initiative: Osteoarthritis research society international set of responder criteria for osteoarthritis
clinical trials revisited. Osteoarthritis Cartilage 2004;12:389–99.
https://doi.org/10.1016/j.joca.2004.02.001.
[27] Roos EM, Roos HP, Lohmander LS. WOMAC Osteoarthritis Index - Additional dimensions for use in
subjects with post-traumatic osteoarthritis of the knee. Osteoarthritis Cartilage, vol. 7, W.B. Saunders
Ltd; 1999, p. 216–21. https://doi.org/10.1053/joca.1998.0153.
[28] Gomez-Aristizabal A, Sharma A, Bakooshli MA, Kapoor M, Gilbert PM, Viswanathan S, et al. Stage-
specific differences in secretory profile of mesenchymal stromal cells (MSCs) subjected to early- vs
late-stage OA synovial fluid. Osteoarthritis Cartilage 2017;25:737–41.
https://doi.org/10.1016/j.joca.2016.11.010.
[29] Krampera M. Mesenchymal stromal cell licensing: A multistep process. Leukemia 2011;25:1408–14.
https://doi.org/10.1038/leu.2011.108.
[30] Smith M, Triantafillou S, Parker A, Youssef P, Coleman M. Synovial membrane inflammation and
cytokine production in patients with early osteoarthritis - PubMed. J Rheumatol 1997;2:365–71.
[31] Roos EM, Lohmander LS. The Knee injury and Osteoarthritis Outcome Score (KOOS): From joint
injury to osteoarthritis. Health Qual Life Outcomes 2003;1. https://doi.org/10.1186/1477-7525-1-64.
[32] Marx RG, Stump TJ, Jones EC, Wickiewicz TL, Warren RF. Development and evaluation of an activity
rating scale for disorders of the knee. American Journal of Sports Medicine 2001;29:213–8.
https://doi.org/10.1177/03635465010290021601.
[33] Smith SR, Deshpande BR, Collins JE, Katz JN, Losina E. Comparative pain reduction of oral non-
steroidal anti-inflammatory drugs and opioids for knee osteoarthritis: Systematic analytic review.
Osteoarthritis Cartilage 2016;24:962–72. https://doi.org/10.1016/j.joca.2016.01.135.
[34] Batsis JA, Germain CM, Vásquez E, Zbehlik AJ, Bartels SJ. Physical activity predicts higher physical
function in older adults: The osteoarthritis initiative. J Phys Act Health 2016;13:6–16.
https://doi.org/10.1123/jpah.2014-0531.
[35] Wada M, Miyazaki T, Yamamoto Y, Sakamoto T, Adachi T. Correlation of knee laxity with alignment
and repetitive physical activity in patients with knee osteoarthritis: A cross-sectional study. Knee
2023;40:111–21. https://doi.org/10.1016/j.knee.2022.10.003.
[36] Pellegrini CA, Song J, Chang RW, Semanik PA, Lee J, Ehrlich-Jones L, et al. Change in physical
activity and sedentary time associated with 2-year weight loss in obese adults with osteoarthritis. J Phys
Act Health 2016;13:461–6. https://doi.org/10.1123/jpah.2015-0404.
[37] Kasikis S, Baez J, Gandhi I, Grupp S, Kitko CL, Kowalyk S, et al. Mesenchymal stromal cell therapy
induces high responses and survival in children with steroid refractory GVHD and poor risk biomarkers.
Bone Marrow Transplant 2021;56:2869. https://doi.org/10.1038/S41409-021-01442-3.
[38] Perin EC, Borow KM, Henry TD, Mendelsohn FO, Miller LW, Swiggum E, et al. Randomized Trial of
Targeted Transendocardial Mesenchymal Precursor Cell Therapy in Patients With Heart Failure. J Am
Coll Cardiol 2023;81:849–63. https://doi.org/10.1016/J.JACC.2022.11.061.

[39] Ouboter LF, Barnhoorn MC, Verspaget HW, Plug L, Pool ES, Szuhai K, et al. Local administration of
mesenchymal stromal cells is safe and modulates the immune compartment in ulcerative proctitis. JCI
Insight 2023;8. https://doi.org/10.1172/JCI.INSIGHT.167402.
[40] Boregowda S V., Krishnappa V, Haga CL, Ortiz LA, Phinney DG. A Clinical Indications Prediction
Scale Based on TWIST1 for Human Mesenchymal Stem Cells. EBioMedicine 2016;4:62–73.
https://doi.org/10.1016/j.ebiom.2015.12.020.
[41] Lee RH, Boregowda S V., Shigemoto-Kuroda T, Bae EH, Haga CL, Abbery CA, et al. TWIST1 and
TSG6 are coordinately regulated and function as potency biomarkers in human MSCs. Sci Adv
2023;9:eadi2387. https://doi.org/10.1126/SCIADV.ADI2387.
[42] Ménard C, Dulong J, Roulois D, Hébraud B, Verdière L, Pangault C, et al. Integrated transcriptomic,
phenotypic, and functional study reveals tissue-specific immune properties of mesenchymal stromal
cells. Stem Cells 2020;38:146–59. https://doi.org/10.1002/STEM.3077.
[43] Kuznetsov SA, Mankani MH, Bianco P, Robey PG. Enumeration of the colony-forming units-fibroblast
from mouse and human bone marrow in normal and pathological conditions. Stem Cell Res 2009;2:83–
94. https://doi.org/10.1016/j.scr.2008.07.007.
[44] Siegel G, Kluba T, Hermanutz-Klein U, Bieback K, Northoff H, Schäfer R. Phenotype, donor age and
gender affect function of human bone marrow-derived mesenchymal stromal cells. BMC Med 2013;11.
https://doi.org/10.1186/1741-7015-11-146.
[45] Driban JB, McAlindon TE, Amin M, Price LL, Eaton CB, Davis JE, et al. Risk factors can classify
individuals who develop accelerated knee osteoarthritis: Data from the osteoarthritis initiative. Journal
of Orthopaedic Research 2018;36:876–80. https://doi.org/10.1002/jor.23675.
[46] Gómez-Aristizábal A, Gandhi R, Mahomed NN, Marshall KW, Viswanathan S. Synovial fluid
monocyte/macrophage subsets and their correlation to patient-reported outcomes in osteoarthritic
patients: A cohort study. Arthritis Res Ther 2019;21. https://doi.org/10.1186/s13075-018-1798-2.
[47] Chan MWY, Viswanathan S. Recent progress on developing exogenous monocyte/macrophage-based
therapies for inflammatory and degenerative diseases. Cytotherapy 2019;21:393–415.
https://doi.org/10.1016/J.JCYT.2019.02.002.
[48] Rabani R, Volchuk A, Jerkic M, Ormesher L, Garces-Ramirez L, Canton J, et al. Mesenchymal stem
cells enhance NOX2-dependent reactive oxygen species production and bacterial killing in macrophages
during sepsis. European Respiratory Journal 2018;51:1702021.
https://doi.org/10.1183/13993003.02021-2017.
[49] Huang ZY, Stabler T, Pei FX, Kraus VB. Both systemic and local lipopolysaccharide (LPS) burden are
associated with knee OA severity and inflammation. Osteoarthritis Cartilage 2016;24:1769–75.
https://doi.org/10.1016/J.JOCA.2016.05.008.
[50] Clark EA, Kalomoiris S, Nolta JA, Fierro FA. Concise review: MicroRNA function in multipotent
mesenchymal stromal cells. Stem Cells 2014;32:1074–82.
[51] Lara-Barba E, Araya MJ, Hill CN, Bustamante-Barrientos FA, Ortloff A, García C, et al. Role of
microRNA Shuttled in Small Extracellular Vesicles Derived From Mesenchymal Stem/Stromal Cells for
Osteoarticular Disease Treatment. Front Immunol 2021;12. https://doi.org/10.3389/fimmu.2021.768771.
[52] Sun H, Zhai H, Han K, Ma H, Tan Y, Li S, et al. Clinical outcomes of autologous adipose-derived
mesenchymal stem cell combined with high tibial osteotomy for knee osteoarthritis are correlated with
stem cell stemness and senescence. J Transl Med 2024;22:1039. https://doi.org/10.1186/S12967-024-
05814-3.
[53] Vincent TL. Peripheral pain mechanisms in osteoarthritis. Pain 2020;161:S138.
https://doi.org/10.1097/J.PAIN.0000000000001923.
[54] Chen WC, Lu YC, Kuo SJ, Lin CY, Tsai CH, Liu SC, et al. Resistin enhances IL-1β and TNF-α
expression in human osteoarthritis synovial fibroblasts by inhibiting miR-149 expression via the MEK
and ERK pathways. FASEB Journal 2020;34:13671–84. https://doi.org/10.1096/fj.202001071R.
[55] Daghestani HN, Pieper CF, Kraus VB. Soluble macrophage biomarkers indicate inflammatory
phenotypes in patients with knee osteoarthritis. Arthritis Rheumatol 2015;67:956–65.
https://doi.org/10.1002/art.39006.
[56] Haraden CA, Huebner JL, Hsueh MF, Li YJ, Kraus VB. Synovial fluid biomarkers associated with
osteoarthritis severity reflect macrophage and neutrophil related inflammation. Arthritis Res Ther
2019;21. https://doi.org/10.1186/s13075-019-1923-x.
[57] Mabey T, Honsawek S, Saetan N, Poovorawan Y, Tanavalee A, Yuktanandana P. Angiogenic cytokine
expression profiles in plasma and synovial fluid of primary knee osteoarthritis. Int Orthop
2014;38:1885–92. https://doi.org/10.1007/s00264-014-2406-y.
[58] Raghu H, Lepus CM, Wang Q, Wong HH, Lingampalli N, Oliviero F, et al. CCL2/CCR2, but not
CCL5/CCR5, mediates monocyte recruitment, inflammation and cartilage destruction in osteoarthritis.
Ann Rheum Dis 2017;76:914–22. https://doi.org/10.1136/annrheumdis-2016-210426.

[59] Gavin C, Boberg E, Von Bahr L, Bottai M, Andrén AT, Wernerson A, et al. Tissue immune profiles
supporting response to mesenchymal stromal cell therapy in acute graft-versus-host disease - A gut
feeling. Stem Cell Res Ther 2019;10:1–6. https://doi.org/10.1186/S13287-019-1449-9/FIGURES/1.
[60] Kraus VB, Stabler T V., Kong SY, Varju G, McDaniel G. Measurement of synovial fluid volume using
urea. Osteoarthritis Cartilage 2007;15:1217–20. https://doi.org/10.1016/j.joca.2007.03.017.
[61] Østergaard M, Stoltenberg M, Løvgreen-Nielsen P, Volck B, Jensen CH, Lorenzen I. Magnetic
resonance imaging-determined synovial membrane and joint effusion volumes in rheumatoid arthritis
and osteoarthritis: Comparison with the macroscopic and microscopic appearance of the synovium.
Arthritis Rheum 1997;40:1856–67. https://doi.org/10.1002/art.1780401020.
[62] Trajerova M, Kriegova E, Mikulkova Z, Savara J, Kudelka M, Gallo J. Knee osteoarthritis phenotypes
based on synovial fluid immune cells correlate with clinical outcome trajectories. Osteoarthritis
Cartilage 2022;30:1583–92. https://doi.org/10.1016/J.JOCA.2022.08.019.
[63] Kraus VB, Collins JE, Hargrove D, Losina E, Nevitt M, Katz JN, et al. Predictive validity of
biochemical biomarkers in knee osteoarthritis: Data from the FNIH OA Biomarkers Consortium. Ann
Rheum Dis 2017;76:186–95. https://doi.org/10.1136/annrheumdis-2016-209252.
[64] Riddle DL, Jiranek WA. Knee osteoarthritis radiographic progression and associations with pain and
function prior to knee arthroplasty: A multicenter comparative cohort study. Osteoarthritis Cartilage
2015;23:391–6. https://doi.org/10.1016/j.joca.2014.12.013.
[65] Riddle DL, Jiranek WA. Knee Osteoarthritis Radiographic Progression and Associations with Pain and
Function Prior to Knee Arthroplasty: A Multicenter Comparative Cohort Study. Osteoarthritis and
Cartilage / OARS, Osteoarthritis Research Society 2015;23:391.
https://doi.org/10.1016/J.JOCA.2014.12.013.
843
844

Response classification	Subject ID	Cell dose (x 10^6 cells)	Patient Gender	Pain		ADL	
				Delta	Percent Change	Delta	Percent Change
Non-Responder	1	1	Man	5.50	28.35	2.90	32.95
Non-Responder	2	1	Man	11.10	18.17	-5.90	-25.11
Function-Pain Responder	**3**	**1**	**Woman**	**16.60**	**49.85**	**25.00**	**80.91**
Non-Responder	4	10	Man	-2.70	-48.21	0.00	0.00
Function-Pain Responder	**5**	**10**	**Man**	**13.90**	**29.45**	**11.80**	**100.00**
Function-Pain Non-Responder/ Pain Responder	*6*	*10*	*Woman*	*16.70*	*33.40*	*-8.80*	*-200.00*
Function-Pain Responder	**7**	**50**	**Woman**	**30.60**	**100.00**	**20.60**	**100.00**
Function-Pain Responder	**9**	**50**	**Woman**	**22.20**	**31.99**	**19.10**	**43.31**
Function-Pain Non-Responder/ Function Responder	*10*	*50*	*Woman*	*11.10*	*19.04*	*17.70*	*42.96*
Function-Pain Responder	**11**	**50**	**Man**	**36.10**	**81.31**	**20.60**	**70.07**
Non-Responder	12	10	Man	-30.50	-78.41	-14.70	-45.37
Function-Pain Responder	**13**	**1**	**Man**	**38.90**	**63.67**	**39.70**	**77.09**

854
855

856
857

Figure Legends

Fig. 1. Osteoarthritis patient responders demonstrate improvements in KOOS subscales that are maintained out to 24 months follow-up post-injection. A and B) Patients categorized as Function-Pain Responders (A) and Function Responders (B) displayed significant improvements (indicated by lower score values) in KOOS subscales and overall KOOS at 12- and 24-months follow-up relative to baseline, while Non-Responders displayed no significant changes. KOOS scales were inverted such that higher scores represent greater knee-related issues. C) Flow chart summary of patient responder status at 12-months, 24-months, and 8-10-years follow-up. D) Summary of OARSI-OMERACT Responder classification in 2025 after 8-10 years of follow-up. Changes in patient responder status at 8-10 years are bolded. Patient-reported follow-up notes on additional interventions are included in the right column. N=12 patients. Two-way repeated measures ANOVA, Dunnett's multiple comparisons test. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001; asterisks indicate significant differences for responder group relative to baseline at the time points indicated. ADL: function in daily living; Sports/Rec: function in sport and recreation; QOL: knee-related quality of life; KOOS: knee injury and osteoarthritis outcome score; PRP: platelet-rich plasma; TKR: total knee replacement.

Fig. 2. Independent validation that a previously identified gene panel distinguishes clinically effective MSC(M) donors from non-responder MSC(M) donors. A) Principal component (PC) analysis of MSC(M) expression of an immunomodulatory gene panel identified previously for MSC(AT) (*TNFAIP6*, *ACTA2*, *ANGPT1*, *CXCL8*, *EDN1*, *THBS1*, *PDCD1LG2*, *CCN2*; [20]) expressed under licensed conditions. The corresponding loading plot of eigenvectors (right) indicates the relative contribution of each gene to the PC1 and PC2 axes. B) MSC(M) derived from Function-Pain Responders and Function Responders displayed significantly lower PC1 scores relative to non-responders, indicating that the gene panel sensitively distinguishes MSC(M) from donors classified as responders. Student's t-test, ***p<0.001, ****p<0.0001. Horizontal lines: group mean; error bars: standard deviation. C) Significant negative correlations between PC1 scores and the percentage change at 24-months relative to baseline for KOOS ADL, pain, symptoms, and overall KOOS. Percent change in KOOS values were made relative to baseline and calculated such that positive values represent improvement. Spearman's correlation. N=9 MSC(M) donors; n=2-3 replicates/donor.

Fig. 3. *In vitro* monocyte/macrophage polarization readouts distinguish donor MSC(M) immunomodulatory fitness for responders versus non-responders and can be used to empirically rank the fitness of individual MSC(M) donors. A) Schematic of *in vitro* monocyte/macrophage (MΦ) polarization experiment. Indirect co-cultures of MSC(M) with human peripheral blood-derived MΦ was performed for 48 h prior to removal of MSC(M) and addition of lipopolysaccharide (LPS) for 4 h incubation. B and C) Summary of MΦ gene expression after co-culture with Function-Pain Responders (B) and Function Responders (C). Box-and-whisker plots (left) display changes in gene expression relative to MΦ cultured alone (Solo; dotted line). Horizontal line: median; hinges: first and third quartiles; whiskers: range. Two-way ANOVA, Tukey's post-hoc test. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 relative to Solo condition or to groups indicated by brackets. Discriminant canonical

plots (right) of -ddCt values relative to Solo condition. Inner ellipses indicate 95% confidence for the mean of each group; outer ellipses indicate the normal region estimated to contain 50% of the population for each group.

Fig. 4. microRNA-sequencing reveals microRNA signatures in Function Responder donor MSC(M). A) Principal component analysis of microRNA expression profiles demonstrate clustering in Function Responders versus Non-Responders. MSC(M) samples cluster by donor and responder status rather than by synovial fluid (SF) stimulation. B) Volcano plot displays 33 significantly differentially expressed microRNAs in MSC(M) derived from donors classified as Function Responders versus Non-Responders. C&D) Top 30 pathways identified through Reactome analysis of predicted gene targets for significantly upregulated (C) and downregulated (D) microRNAs. E&F) Top 30 pathways identified through Gene Ontology (Biological Processes) analysis of predicted gene targets for significantly upregulated (E) and downregulated (F) microRNAs.

Fig. 5. More severe KOOS and lower MARS physical activity scores correlate to improved patient-reported outcomes at 12- and 24-months follow-up. A) Patients categorized according to Function Responders displayed significantly more severe overall KOOS and Sports/Rec scores at baseline relative to Non-Responders. B) Patients categorized according to Function-Pain Responders and Non-responders displayed no significant differences between overall KOOS and KOOS subscales at baseline. Horizontal lines: group mean; error bars: standard deviation. C) Significant correlations between baseline KOOS Sports/Rec scores and improvements in delta ADL scores at 12 months relative to baseline, percentage change in symptoms at 24 months relative to baseline, delta ADL at 24 months relative to baseline, and delta overall KOOS at 24 months relative to baseline. D) Baseline MARS significantly correlated to baseline Sports/Rec, percent change in KOOS Symptoms at 12 months relative to baseline, as well as both delta and percent change in KOOS Pain at 12 months relative to baseline. KOOS scales were inverted such that higher scores represent greater knee-related issues; delta and percent change in KOOS values were made relative to baseline and calculated such that positive values represent improvement. N=12 patients. Student's t-test (for baseline Overall KOOS), Two-way ANOVA, Sidak's post-hoc test (for baseline KOOS Subscales), Spearman's correlation. *$p<0.05$, **$p<0.01$. n.s.; non-significant. ADL: function in daily living; Sports/Rec: function in sport and recreation; QOL: knee-related quality of life; Pt.: patient ID; MO: month; KOOS: knee injury and osteoarthritis outcome score; MARS: Marx Activity Rating Scale.

Fig. 6. Local synovial fluid biomarkers measured at baseline correlate to patient-reported outcome measures at 3, 12, and 24 months relative to baseline. A) Higher levels of resistin significantly positively correlated to delta values in KOOS Sports/Rec scores at 24 months. Higher levels of TIMP1 and CCL2 significantly correlated inversely to the percentage change and/or delta in KOOS Sports/Rec scores at 24 months relative to baseline. B) Higher levels of HGF and VEGFA in synovial fluid significantly correlated inversely to delta values and/or percentage change in KOOS Pain scores at 12 months relative to baseline. C) Higher levels of sCD14 and

CCL2 significantly correlate inversely to the percentage change and/or delta in KOOS Symptom and Sports/Rec scores at 3 months relative to baseline. Delta and percent change in KOOS values were made relative to baseline and calculated such that positive values represent improvement. N=8-9 patients. Spearman's correlation. Sports/Rec: function in sport and recreation; MO: month; KOOS: knee injury and osteoarthritis outcome score; TIMP1: tissue inhibitor of metalloproteinases 1; CCL2: chemokine C-C motif ligand 2; HGF: hepatocyte growth factor; sCD14: soluble CD14; VEGFA: vascular endothelial growth factor A.

Function-Pain Responders

Function Responders

A — Symptoms, Pain, ADL, Sports/Rec, QOL, Overall KOOS

● Function-Pain Responder   ■ Non-Responder

B — Symptoms, Pain, ADL, Sports/Rec, QOL, Overall KOOS

● Function Responder   ■ Non-Responder

C

- 12-month: 8 Responders | 4 Non-Responders
- 24-month: 7 Responders | 1 Non-Responder | 4 Non-Responders
  - 1 lost to follow-up (TKR)
  - 1 lost to follow-up
- 8-10 year:
  - 5 Responders: No intervention: 1, Injection: 2, TKR: 2
  - 1 Non-Responder: No intervention
  - 1 Non-Responder: No intervention
  - 3 Non-Responders: No intervention: 1, Injection: 2

D

| | | | | | 8-10 year follow up scores | | | | |
| | | | | | KOOS Pain | | KOOS ADL | | |
Subject ID	Responder Class at 12 Months	Responder Class at 24 Months	Responder Class at 8-10 years	# Years Follow-up in 2025	Delta	% Change	Delta	% Change	Follow-up notes
1	Non-Responder	Non-Responder	Non-Responder	9 years, 10 months	-0.04	-0.21	-1.49	-16.93	Injection of viscosupplementation in 2024 and 2025
2	Non-Responder	Non-Responder	N/A	-	-	-	-	-	N/A
3	Function-Pain Responder	Function Responder	Function-Pain Responder	9 years, 6 months	30.52	91.65	27.96	90.49	PRP injection in 2023
4	Non-Responder	Non-Responder	Non-Responder	9 years, 4 months	-44.40	-792.86	-30.88	-3088.00	No surgical intervention
5	Function-Pain Responder	Pain Responder	N/A	-	-	-	-	-	TKR in 2025
6	Pain Responder	Pain Responder	Pain Responder	9 years, 0 months	11.11	22.22	-22.07	-501.59	Awaiting bilateral knee replacement
7	Function-Pain Responder	Function-Pain Responder	**Non-Responder**	8 years, 11 months	0.04	0.13	-1.46	-7.09	No intervention post injection
9	Function-Pain Responder	Function-Pain Responder	**Non-Responder**	8 years, 6 months	-2.82	-4.06	-19.14	-43.40	No intervention post injection
10	Function Responder	Function-Pain Responder	Function-Pain Responder	8 years, 6 months	19.41	33.29	20.61	50.02	Viscosupplementation - no majory surgery
11	Function-Pain Responder	Function-Pain Responder	Function-Pain Responder	8 years, 6 months	24.96	56.22	16.16	54.97	No knee surgery/scope, no knee injection
12	Non-Responder	Non-Responder	Non-Responder	8 years, 4 months	-30.54	-78.51	-24.95	-77.01	Viscosupplementation in 2019
13	Function-Pain Responder	Function-Pain Responder	Function-Pain Responder	8 years, 2 months	49.99	81.82	39.74	77.17	TKR in 2025

FIG. 2

FIG. 3

FIG. 4

FIG. 5

FIG. 6