Responders vs. non-responders to mesenchymal stromal cells in knee osteoarthritis patients: mechanistic correlates of donor cell attributes and putative patient features

Kevin P. Robb,[1,2,3] Razieh Rabani,[1,2] Shabana Vohra,[1,2] Shoba Singh,[1,2] Jaskarndip Chahal,[1,2,4] Julie Audet,[3] Anthony V. Perruccio,[1,2,4,5] Ali Naraghi[6,7], Rajiv Gandhi,[1,4] Osvaldo Espin-Garcia,[1,8,9] Sowmya Viswanathan*[1,2,3,10]

[1]Osteoarthritis Research Program, Division of Orthopedic Surgery, Schroeder Arthritis Institute, University Health Network, Toronto, Canada
[2]Krembil Research Institute, University Health Network, Toronto, Canada
[3]Institute of Biomedical Engineering, University of Toronto, Toronto, Canada
[4]Department of Surgery, University of Toronto, Toronto, Canada
[5]Institute of Health Policy, Management and Evaluation, Dalla Lana School of Public Health, University of Toronto, Toronto, Canada
[6]Toronto Joint Department of Medical Imaging, Toronto, Ontario, Canada
[7]Sinai Health System, University Health Network and Women's College Hospital, Department of Medical Imaging, Temerty Faculty of Medicine, University of Toronto, Toronto, Ontario, Canada
[8]Epidemiology and Biostatistics, Western University
[9]Dalla Lana School of Public Health, University of Toronto [10]Department of Medicine, University of Toronto, Toronto, Canada

*Corresponding author:
Sowmya Viswanathan, PhD
Krembil Research Institute
60 Leonard Ave
Toronto, Ontario, Canada
M5T 0S8
Phone: 416-634-7239
Email: sowmya.viswanathan@uhn.ca

ORCID IDs:
Kevin P. Robb: 0000-0001-7090-5145
Razieh Rabani: 0000-0001-7322-5524
Shabana Vohra: 0000-0003-1995-3795
Shoba Singh: 0000-0001-7322-5524
Jaskarndip Chahal: 0000-0002-5098-7865
Julie Audet: 0000-0002-5599-6074
Anthony V. Perruccio: 0000-0002-4389-0993
Ali Naraghi:
Rajiv Gandhi: 0000-0003-2816-0033
Osvaldo E. Garcia: 0000-0003-2052-2626
Sowmya Viswanathan: 0000-0002-4552-0179

Disclaimers: Sowmya Viswanathan declares 60% ownership of Regulatory Cell Therapy Consultants Inc., which does not conflict with this paper in any way. Sowmya Viswanathan, Razieh Rabani, and Kevin P Robb are co-inventors on patent applications filed with UHN based on this work.

Brief acknowledgements:
This research was funded by the Canadian Institutes of Health Research (CIHR), Natural Sciences and Engineering Research Council of Canada (NSERC), Toronto General and Western Hospital Foundation, and the Arthritis Society.

Keywords: responders vs. non-responders, mesenchymal stromal cells, osteoarthritis, immunomodulation, potency attributes, synovial fluid biomarkers, microRNAs

SUMMARY

Mesenchymal stromal cells (MSCs) have resulted in heterogenous clinical effectiveness in knee osteoarthritis (KOA). Herein, a framework that dually correlates KOA patient responsiveness to baseline bone marrow-derived MSC(M) donor batch attributes and baseline clinical and biomarker features is provided. Using previous clinical trial data, we demonstrated that MSCs with increased immunomodulatory potency are more efficacious. Multivariable MSC(M) genes correlated strongly with responder status and to 12- and 24-month improvements in Knee Osteoarthritis and Outcome Scores. Responder MSC(M) donor batches had differential microRNA expression and ability to polarize CD14+ monocytes *in vitro*. KOA Responders had more severe OA and lower physical activity. Baseline biomarkers showed trending correlations to patient responsiveness. 62.5% of KOA patients remained Responders at 8-10 years, speaking to longer-term effects of single MSC(M) injections. Thus, our analytical methodology defines critical quality attributes of potent MSC donor batches and identifies putative KOA patient theratypes to MSC treatments.

INTRODUCTION

Knee osteoarthritis (KOA) affects 16% of adults over 40 years of age worldwide [1], with no available disease-modifying drugs. Pathophysiological mechanisms of KOA are heterogeneous and fluctuate temporally [2]. Culture-expanded mesenchymal stromal cells (MSCs) are being investigated for their multimodal mechanisms of action, including immunomodulatory functions [3,4]; however, there has been conflicting clinical efficacy reported. Equivocal results show no difference in 12-month improvements between umbilical cord-derived MSCs vs. first-line corticosteroids treatments [5]; conversely, other reports [6–9], meta-analyses [10,11], and randomized controlled trials [12,13] showed symptom and structural improvements with MSC treatments. This suggests heterogeneous responsiveness, stressing the need to identify signatures that distinguish responder from non-responder donor batches to deliver effective MSC treatments to KOA patients.

Heterogeneous responsiveness to MSC treatments is due to an interplay of both basal potency attributes of donor MSC batches and basal patient disease status [4]. Heterogeneity exists among MSC investigational products due to intrinsic variations in donors [4,14] and manufacturing parameters [15,16]; well-defined critical quality attributes (CQAs) that are sensitive, quantitative and relevant to disease-specific mechanism of action are thus needed [17].

Using reported [8], and new follow-up data from a phase I/IIa dose-escalation clinical trial, we conduct in-depth analyses of donor MSC batches and baseline/follow-up KOA patient characterization to provide an integrated algorithm to analyze heterogenous KOA patient responsiveness to MSC treatments. Using a previously reported gene panel that correlated to *in vitro* functional activity [18], there is significant discrimination of responder vs. non-responders MSC(M) donor batches from marrow tissue. MicroRNAs, important mediators of MSC [19] further discriminate MSC(M) from responders vs. non-responders.

KOA is recognized not as a single disease but a constellation of clinical phenotypes [20,21] and divergent molecular endotypes [22]. Here we report an exploratory analytical framework that combines clinical phenotypes and molecular endotypes that can be used as the basis for identifying KOA patient responsiveness to MSC(M) treatments.

MATERIALS AND METHODS

Clinical trial and patient data

KOA patients (Kellgren Lawrence (KL) grade 3-4) were recruited to a nonrandomized, open□label, dose□escalation, Health Canada-authorized phase I/IIa clinical trial as reported [8] (NCT02351011). Ethics approvals including for long-term follow up were granted (REB 14–7909;20-5066;24-5463) [8]. Complete clinical follow-up was performed 24-months after MSC(M) injection for all patients. Additional follow-up,at 8-10 years was obtained for 10/12 patients; the remaining 2 patients were lost to follow-up. Calculation of patient-reported outcome measures (PROMs), including the inverted Knee Injury and Osteoarthritis Outcome Score (KOOS) and the Marx Activity Rating Scale (MARS), as described in **Suppl. File 1**. Contrast-enhanced MRI analysis at baseline, 6 and 12-months were conducted on all patients as previously described [8]; whole organ MRI scores (WORMS) and MRI synovitis scores were measured and calculated as previously described [23].

Responder vs. Non-Responder Criteria

Patient clinical responder status was based on the Outcome Measures in Rheumatology (OMERACT) and Osteoarthritis Research Society International (OARSI) Standing Committee for Clinical Trials Response Criteria Initiative (OMERACT-OARSI criteria), based on data from large, powered placebo-controlled drug trials for hip and KOA [2]. These criteria categorize patients as responders according to the following: i) improvement in pain or in function ≥50% and absolute change ≥ 20; OR, ii) improvement (≥20% and absolute change ≥ 10) in at least two of the following: a) pain, b) function, c) patient global assessment; the latter was not measured in our trial. Given the limited sample size and lack of patient global assessment metrics in our previous clinical trial, we categorized patient responder status using the twelve-month follow-up data based on changes in KOOS function (activities of daily living (ADL) subscale) and/or KOOS Pain only, applying OMERACT-OARSI criteria thresholds for function and pain scores of delta value ≥10 AND percentage change ≥20% [2] . The validated KOOS instrument measures responses to interventions in KOA with high test-retest reliability and improved sensitivity in pain and function scoring relative to the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) scoring system [2,4]. Baseline demographics, radiographic KL-grade, magnetic resonance imaging (MRI) total cartilage volume (WORMS, [23] and synovitis [24] scores are provided in both Function Responder vs. Non-Responder donor categories (Table 1).

Biomarker measurements

Levels of curated biomarkers in baseline serum (**Table S2**) were measured previously [8]; updated analyses and experimental questions were applied to the dataset in the present work. Additional inflammatory biomarkers were measured in baseline biobanked patient serum and plasma (analytes specified in **Table S2**). Synovial fluid collection was feasible for only nine of twelve patients. Immune cell profiling was done only on available (five), paired baseline and follow-up synovial fluid samples and is thus considered exploratory [8].

Statistical analyses

GraphPad Prism 6.0 and JMP Pro 17 software were used for statistical analyses and to create plots. Statistical tests are specified in figure and table captions. Spearman's correlation was selected as a non-parametric statistical test to investigate associations between inverted KOOS changes and other variables, given that the data violated assumptions of normality and homoscedasticity. P values derived from Spearman's correlation were corrected for multiple comparisons using the Benjamini-Hochberg False Discovery Rate (FDR) method. Sex, age, body mass index (BMI), KL-grade, MSC dose and baseline MRI WORMs and synovitis scores were not considered as covariates given lack of significant correlation between these factors with Function Responder status (Table 1). Given the limited sample size and exploratory nature of analysis, Spearman's correlations were considered statistically significant when $p<0.05$ (uncorrected) and $p<0.1$, post-FDR.

RESULTS

Patient responder status

We systematically investigated multivariable factors affecting KOA patient responder status using 12- and 24-month patient-reported inverted KOOS data [8]. We applied OMERACT-OARSI criteria [25] to determine responder status, with modifications. Patients were classified as Responders vs. Non-Responders. Within Responders, we identified three sub-categories (**Table S3**) including: i) Function-Pain Responders (exceeding thresholds in both KOOS ADL and Pain); ii) Function Responders (exceeding thresholds in ADL), iii) Pain Responders (exceeding thresholds in Pain); Non-Responders did not exceed thresholds in either ADL or Pain (**Table S3**). Of twelve patients, eight total Responders (all categories) and four Non-Responders met these criteria. While the sample size is limited, the binary criteria

was useful in determining factors affecting patient clinical responses. However, correlations
with all KOOS subscale scores were included to mitigate the limited power of binary
analysis. There was no observed dose-dependency; responders existed across all three
MSC(M) doses including the 1 x 10^6 (2/4), 10 x 10^6 (1/4), and 50 x 10^6 (3/4) (**Table 1**). We
report on Function Responders; other Responder category data are included in the
supplementary files. We checked for associations between covariates including sex, age,
BMI, KL-grade, MSC dose and MRI baseline WORMs and synovitis scores and Function
Responder status and observed no significant correlations (**Table 1**).

12-month Responder vs. Non-Responder classification holds at 24-months and 8–10-year follow-up

Function Responders at 12 months displayed significantly improved inverted KOOS scores in
all subscales at 12- and 24-months follow-up (**Fig. 1A**), supportive of persistent effects of a
single injection of autologous MSC(M). Patients categorized as Non-Responders similarly
showed no significant changes at 12- and 24-months, indicating stable clinical phenotype. No
significant differences were observed when comparing outcomes at 12 vs. 24-months for any
KOOS subscale in Responders or Non-Responders. Similar results were obtained with
Function-Pain (**Fig. S1A**) and Pain (**Fig. S1B**) Responder categories.

We further analyzed patient follow-up at 8-10 years after single MSC(M) injection. 12-month
Responders maintained significant improvement in multiple KOOS subscales at 8-10 years
follow-up (**Fig. S2**). Specifically, 7/8 Responders maintained their Responder status at 24
months (**Fig. 1B, 1C**). Of these, 5/7 maintained Responder status at 8-10 years, and one
patient (underwent knee arthroplasty) did not provide long-term KOOS data. 2/8 Responders
received no additional interventions; 2/8 received viscosupplementation or platelet-rich
plasma; 3/8 patients underwent knee arthroplasty surgery.

4/4 12-month Non-Responder patients maintained status at 24-months, and one patient was
lost to follow-up at the 8–10-years (**Fig. 1B, 1C**). 3/4 patients maintained Non-Responder
status at 8-10 years, with 1/4 patients reporting no additional interventions and 2/4 patients
reporting viscosupplementation.

Immunomodulatory gene panel discriminates between clinically effective and ineffective MSC(M) donor batches

Using biobanked MSC(M) samples cultured under pro-inflammatory licensing conditions, we measured expression of a gene panel that correlated with *in vitro* immunomodulatory functionality [18]. Pro-inflammatory cytokine licensing conditions were used as a common method to measure immunomodulatory genes in MSC(M) [26].

Principal Component (PC) analysis revealed distinct unsupervised clustering between Function Responder vs. Non-Responder MSC(M) donor batches along the PC1 axis (29.3% of total variance), with higher expression of *CCN2* and lower expression of angiogenic markers *THBS1*, *CXCL8*, and *ANGPT1* contributing strongly to higher PC1 scores (**Fig. 2A**, **Fig. S3A**). PC1 scores were significantly lower for all categories of Responders vs. Non-Responders (**Fig. 2B**, **Fig. S3B**), confirming the discriminatory power of this curated gene panel. To ensure that individual donor batches were not skewing results, given limited sample size, outlier analysis was performed by calculating T^2 statistic, identifying three outlier data points (two Responder, one Non-Responder). However, the PCA with these data points removed, returned similar results; clear separation of Responder/Non-Responders was observed across the expected eigenvectors (**Fig. S3C**).

PC1 scores, indexed as a composite score for the basal immunomodulatory gene expression profile, additionally showed statistically significant robust negative correlations with improvements in 12-and 24-month mean KOOS and KOOS pain and/or ADL sub-scores. function at 12- and 24-mo (**Fig. 2C**, **Table S4**).

Individual MSC(M) genes, *ANGPT1* and *CCN2* showed significant correlations with 12- and/or 24-month KOOS pain and ADL subscores and mean KOOS (**Table S5**) that were lost after adjustment for multiple comparisons.

In vitro immunomodulation by donor MSC(M) batches correlates to patient responder status

Functional *in vitro* monocyte polarization using mismatched human peripheral blood monocytes further assessed baseline MSC(M) immunomodulatory potency (**Fig. 3A**). Discriminant analysis demonstrated distinct clustering and greater separation of monocyte phenotypic profiles between Responder vs. Non-Responder donor MSC(M) batches (**Fig. 3B**). Co-cultures of monocytes with Responder MSC(M) donor batches significantly

upregulated expression of the inflammation-resolving markers *IL10* and *STAB1,* and reduced expression of *CD86* compared to Non-Responders batches (**Fig. 3B**). The immunosuppressive marker, *CD274*, was upregulated by Non-Responder vs. Responder MSC(M) donor batches. Overall, Responder MSC(M) donor batches modulated *in vitro* monocyte polarization towards a distinct pro-resolving phenotype. Discriminant analysis showed reduced separation with Function-Pain Responder (**Fig. S4A**) or Pain Responder (**Fig. S4B**) donor MSC(M) batches as indicated by greater misclassified data and reduced Entropy R^2 values (**Table S6**).

Specific microRNAs distinguish responder donor MSC(M) batches

Unbiased baseline differences in MSC(M) donor batches after synovial fluid stimulation were probed with microRNA-sequencing. PCA demonstrated clustering by donor batch and responder group rather than by synovial fluid stimulation (**Fig. 4A**). Thirty-three significant, differentially expressed microRNAs were identified between Responder vs. Non-Responder MSC(M) donor batches after correction for multiple comparisons; 11 microRNAs were upregulated, and 22 microRNAs were downregulated (**Fig. 4B**, **Table S7**). Gene targets related to cell cycle (GTPases), phosphoinositide 3-kinase (PI3K), mitogen-activated protein kinase (MAPK), transforming growth factor beta (TGF-β), VEGF, SMAD proteins, apoptosis, and other pathways were identified (**Fig. 4C,D**; **Suppl. File 2**). Gene ontology analysis identified biological processes linked to neuronal processes, histone modification, cell growth, ossification, Wnt signaling (**Fig. 4E,F**; **Suppl. File 2**).

Predicted gene targets did not include genes within the curated mRNA panel. Limited correlations between the top 10 differentially expressed microRNAs with the curated gene panel were identified, but significance was lost after correction for multiple comparisons (p<0.05, FDR>0.1; hsa-miR-377-3p vs *ACTA2*; hsa-miR-1307-3p vs *THBS1*; hsa-miR-4775 vs *THBS1)* correlations (**Table S8**), suggesting putative indirect modulation of select MSC(M) genes by differentially expressed microRNAs that needs to be confirmed in larger datasets. Of the top 10 miRNA, miRNA-483-5p and miRNA-642a-5p positively and negatively correlated, respectively, with improvements in 12-and 24-month KOOS pain and function sub-scores or mean KOOS (Fig 4G; **Table S9**). The microRNA-sequencing analysis provides further characterization and potential mechanistic insights into MSC(M) therapeutic effects in KOA patients.

Baseline clinical phenotype correlates with 12- and 24-month and 8–10-year improvements

We conducted exploratory analysis between baseline inverted KOOS subscales with subsequent responder status (positive delta or percent change values indicate improvement; **Fig. 5**). Function Responders displayed significantly elevated mean KOOSKOOS Sports/Rec at baseline compared to Non-Responders, supportive of more severe baseline clinical presentations in directing MSC responsiveness (**Fig. 5A**). Other responder categories however, displayed no significant differences (**Fig. S5A,S5B**).

Next, we analyzed correlations between the baseline scores for all KOOS subscales with changes in these scores at 12- and 24-months (**Table S10**). Only baseline Sports/Rec scores positively correlated with improvements in KOOS ADL at 12- and 24-months; however, significance was lost after correction for multiple comparisons.

Given the findings on KOOS Sports/Rec, we probed the relationship with physical activity levels using the MARS scores [27]. Higher baseline MARS scores (i.e., more frequent physical activity) were significantly associated with lower (i.e., less severe) baseline KOOS Sports/Rec scores (**Fig. 5C**). Interestingly, higher MARS scores significantly inversely correlated with improvements in the 12-month KOOS Pain sub-score (**Fig. 5C**, **Table S11**).

Baseline MRI scores did not correlate with responder status

Baseline MRI WORMS and synovitis scores, previously calculated [8] did not correlate with Function Responder or Non-Responder classifications (**Table 1**) or to improvements in 12- and 24-month KOOS (**Tables S12, S13**). We had previously reported no significant improvements in MRI WORMS or synovitis scores, relative to baseline based on blinded analysis by two radiologists [8].

Baseline synovial fluid biomarkers do not correlate with 6, 12- and 24-month patient improvements

The levels of baseline inflammatory, angiogenic, metabolic and cartilage extracellular matrix-associated biomarkers in synovial fluid from a subset of nine patients were previously reported [8]; here we analyzed correlations with 6, 12- and 24-month KOOS improvements (**Tables S14, S15**). There were **negative** correlations for baseline HGF and VEGFA with greater improvements in KOOS Pain subscores at 12-months, but significance was lost after adjustment for multiple comparisons (**Table S14**) and no significant correlations between

other local biomarkers including TIMP1, resistin, adipsin, leptin, adiponectin, soluble (s)CD14, sCD163, MMP1, MMP9, IL6, IL8, CXCL1, CCL2, CX3CL1 with mean KOOS and KOOS pain or function sub-scores for all timepoints. [8].

Exploratory analysis of baseline synovial fluid immune cell phenotypes

As additional exploration of KOA patient molecular endotype, we analyzed synovial fluid immune cells using previous data [8]. While this analysis was feasible for a subset of five patients due to limited synovial fluid volumes, our data provides proof-of-concept for probing baseline levels of $CD14^+CD16^+$ and $CD14^+CD163^+$ monocyte/macrophages (MΦs) (**Fig. S6A**) as well as $CD3^+CD4^+CD69^+$ and $CD3^+CD4^+CD25^+$ T helper cells (**Fig. S6B**) as putative immunophenotypes of interest. Higher baseline frequencies of MΦs and activated T cells are present in Non-Responders, although not significant given the limited sampling.

Baseline systemic biomarker levels do not correlate with patient improvements

Systemic baseline inflammatory biomarkers were measured in the present study (sCD163; CRP; S100A8/A9, TNFα; IL-6) alongside previous data on systemic biomarkers of cartilage breakdown [8] (**Table S16, S17**). Higher levels of baseline plasma TNFα significantly positively correlated to percentage changes in KOOS Pain subscores at 6 months (**Fig. S7**, **Table S17**), but this significance was lost with multiple comparison corrections. Thus, the selected panel of local and systemic biomarkers of different categories showed limited capacity to correlate with patient-reported outcomes.

DISCUSSION

Statistically powered clinical trials using MSCs have reported conflicting results in KOA patients [5,12]. To address underlying factors that drive heterogeneous responsiveness to MSCs, we analyzed our test clinical dataset [8] and identified putative MSC(M) donor batch characteristics and responsive KOA patient features that correlated with improvements in PROMs at 12- and 24-months. We anticipate that this combinatorial analysis will serve as an integrated algorithm that can be tested in prospective KOA clinical trials (and retrospectively on existing datasets) to i) screen potent donor/batches of MSCs; and ii) to identify characteristics of responsive KOA recipients. Our data suggests that MSC(M) donor batches with high basal immunomodulatory fitness as measured by reduced levels of angiogenic

genes, higher monocyte modulatory activity and a distinct signature of microRNAs will be more clinically effective. Our analysis of responsive KOA patients was more exploratory given the limited dataset. KOA patients with reduced knee functionality associated with sports and recreation activities and reduced physical activity levels were more responsive to MSC(M) injections. Biomarker analysis did not show significant correlations but there were trends pointing toward reduced levels of soluble angiogenic factors in the local joint environment at baseline that need to be further investigated in larger clinical datasets with matched placebo controls.

We used a previous clinical trial [8] with twelve manufactured, patient-specific lots of MSC(M), which was data-rich with multiple baseline biomarkers, and clinical readouts and offered an excellent test dataset to study MSC(M) batch variations. We provided longitudinal 12-, 24-month and 8–10-year follow-up and are one of the first groups to report such long-term clinical data. Responders showed significant improvements in PROMs at 12- and 24-months, and significantly sustained Responder (62.5%) and Non-Responder (75%) status to 8-10 years, including those lost to follow-ups. We note that 2/8 (25%) Responders (all categories) and 2/4 (50%) Non-Responders received additional injections over the intervening years; 3/8 Responders received or were awaiting a total knee replacement (TKRs). Additionally, 3/8 Responders (37.5%) and 1/4 Non-Responders (25%) did not receive additional interventions. The higher number of TKRs in the Responder group may be related to differences in sample size (N=8 Responders vs. N=4 Non-Responders). Given the extended lag in the follow-up period, it is not possible to confirm if the single MSC(M) injections delayed knee replacements, nor can we rule out the effects of additional interventions that may have contributed to the 8-10-year outcome measures.

Notably, our analysis suggesting that patients with severe KOA were more responsive to MSC(M) treatment aligns with findings in acute graft-versus-host disease [28], myocardial infarction [29] and ulcerative colitis [30], where more severe disease patients responded better to MSC treatments. Function Responders showed correlations to MSC(M) donor immunomodulatory potency and a distinct baseline clinical phenotype (especially KOOS Sports/Rec sub-scores). Although immunomodulatory therapies may be expected to induce greater analgesic effects [31], MSCs act through multimodal mechanisms with pain and functional improvements commonly documented [34]. KOA patients with higher disease severity, especially reduced levels of high-level knee functionality had greater improvements

in daily functioning of the knee (KOOS ADL). Patients with higher baseline MARS scores
had lesser improvements, suggesting that patients who may have persisted through pain in
physical activity may have exacerbated disease progression [31].

We focused on basal immunomodulatory properties of MSC(M) and their ability to modulate
monocytes, given their importance as a clinically relevant mechanistic target in KOA
[4,7,8,32]. A previously identified immunomodulatory gene panel for MSC(AT) [18]
robustly distinguished MSC(M) donors classified under all categories of Responders.
Dimension-reduced PC1 scores capturing multivariable genes were strongly and significantly
correlated inversely to patient improvements at 12-, 24-monthsand are thus more reliable than
the use of individual genes. Notably, this panel included predominantly angiogenic markers
that were previously inversely correlated to monocyte polarization toward inflammation-
resolving phenotypes [18], supportive of inverse relationships between angiogenic and
immunomodulatory fitness of MSC(M) [33,34]. While the transcriptome of MSC(AT) and
MSC(M) are distinct, the genes identified for MSC(AT) are valid for MSC(M) as well,
speaking to their broad utility to serve as MSC CQAs [7]. These data support our hypothesis
that MSCs with high basal immunomodulatory fitness produce greater improvements in knee
function. Nonetheless, the use of autologous MSCs [7,8] adds a layer of complexity, given
that systemic factors such as age, BMI, and other health conditions were dually accounted for
in both basal fitness of the cells [35,36], as well as baseline patient disease status [37].

Responder MSC(M) donor batches modulated $CD14^+$ monocytes to more pro-resolving
subtypes *in vitro*. The importance of monocyte/macrophage modulation in OA has previously
been shown [8,38–40], given the role of synovitis in OA progression and pathology.
Although MSC(M) co-culture resulted in preferential upregulation of pro-resolving $CD14^+$
monocyte markers, there was also upregulation of pro-inflammatory markers. Reliance on
dichotomized gene expression markers is not indicative of the plurality of monocyte
functionality [41]; rather induction of mixed monocyte phenotypes by MSCs is important for
achieving therapeutic effects [42]. Importantly, we have shown that $CD14^+$ monocytes in OA
synovial fluid demonstrate a mixed phenotype, which is not reflective of functionality [43].

Function Responder MSC(M) donor batches had a significantly different microRNA
signature compared to Non-Responders. Pathways related to inflammation, fibrosis,
angiogenesis, and ossification, relevant to KOA pathophysiology were identified. We also
identified enrichment of pathways involved in cell cycle. Surprisingly, our analysis identified

enrichment of several neuronal processes; however, given the role of innervation and
neuronal signaling in joint pain in KOA [44], these microRNAs may be implicated in
mediating analgesic effects of MSC(M). The functionality of differentially expressed
microRNAs requires further study, but they may also serve as additional CQAs for
identifying basally fit MSC(M).

Using a curated panel of inflammatory, angiogenic, and metabolic proteins and proteases, we
could not discern a signature of local biomarkers that definitively identified KOA patients
that were responsive to MSC(M). Responders displayed putatively lower local baseline levels
of angiogenic biomarkers, HGF and VEGFA, but this lost significance after multiple
comparisons correction. Systemically, there was limited short-term positive correlations
between TNFα (plasma) levels and PROM improvements, albeit significance was also lost
after multiple comparisons correction. Baseline serum CRP levels, an inflammatory
biomarker in multiple inflammatory conditions [45] and cartilage breakdown markers (shown
to correlate with worsening KOA pain [46]) failed to correlate with improvements in KOOS
at all time points. Overall, our analyses failed to show significant correlations to MSC(M)
treatment responsiveness with any of the biomarkers tested. However, a trend towards a
mixed profile with increased pro-inflammatory systemic TNFα but reduced local angiogenic
proteins (HGF and VEGF) emerged, reminiscent of mixed biomarker profile in rheumatoid
arthritis patients, responsive to allogenic MSCs [47]. This mixed local vs. systemic molecular
endotype profile needs to be further tested in larger datasets.

Identifying molecular endotypes of OA has proven to be extremely challenging, and multi-
consortium efforts have not been able to definitively identify molecular endotypes when
profiling local synovial fluid, even with large, covariate-adjusted datasets [48]. This
heterogeneity in OA patients is a key reason for failure of multiple drugs in late-phase,
powered clinical trials [49,50]. The absence of stratification biomarkers to identify OA
theratypes for which specific treatments can be tailored to specific clinical phenotypes and
endotypes has been identified as a critical gap impeding the development of disease
modifying drugs in the field [51]. Although systemic biomarkers of inflammation and/or
cartilage degradation are beginning to emerge to cluster patients [52], their utility in defining
patient responsiveness to different therapies remains untested. In our hands, many of the
previously reported inflammatory and cartilage degradation biomarkers [8,52] did not
correlate with improvements in patient-reported outcomes to MSC(M) treatments.

Limitations of this study include small sample size, which allowed us to perform an exploratory but not definitive analysis. Indeed, baseline clinical phenotype and local biomarkers indicative of reduced angiogenic and inflammatory molecular endotype lost significance when corrected for multiple comparisons, likely due to overcorrection arising from the small sample size. Nonetheless, the multivariate framework of investigating potency features of MSC investigational products combined with clinical and biomarker data to analyze patient phenotypes and endotypes is a useful method to test and stratify OA patients into theratypes, suited for MSC therapies [51]. The absence of a control group in the original clinical trial is another limitation [8], although effects were measured at multiple timepoints including 12-, 24-months and 8-10 years. KOA is a progressive condition [53], with approximately 30% of patients displaying worsening outcomes over two years leading up to joint surgery [54]. While placebo effects cannot be ruled out, less than 30% of patients reported worsening outcomes, potentially speaking to a treatment effect. Indeed, analysis of Non-Responders using the rigorous OMERACT-OARSI criteria [25] showed that even patients within this category did not show any significant worsening in PROMs over 24-month and 8-10 years of follow-up.

CONCLUSIONS

Donor batches of MSC(M) that resulted in positive clinical outcomes differed significantly in specific measurable attributes compared to MSC(M) batches that failed to improve pain or function. These quantitative, multivariate, and functionally relevant attributes can define future CQAs for MSC(M) that can be evaluated to strategically design KOA trials to ensure higher success rates of MSC injection. Additionally, we identified exploratory baseline clinical features and biomarkers in KOA patients responsive to MSC(M) injections that need to be further tested in larger clinical datasets to verify their utility.

Acknowledgements

We thank Kim Perry, Mary Nasim, Tamara Wagner, Amanda Weston and the Orthopedic Surgery team for assistance in acquiring donor samples. Thanks to Dr. Mozhgan Rasti who performed qPCR for a subset of experiments, as well as Dr. Mohit Kapoor, Dr. Starlee Lively, Dr. Mehdi Layeghifard, and Pratibha Potla for assistance with microRNA-sequencing and analysis. Schematics for figures were created using BioRender.com.

This research was funded by Canadian Institutes of Health Research (CIHR) (PJT-166089) and Natural Sciences and Engineering Research Council of Canada (NSERC) (RGPIN-2018-05737) grants awarded to SV. The work is in part supported by the Schroeder Arthritis Institute via the Toronto General and Western Hospital Foundation (University Health Network). Salary support for KR was provided by a Natural Sciences and Engineering Research Council (NSERC) Canada Graduate Scholarship. Salary support for RR was provided by the Arthritis Society (TPF-19-0537). AP is supported by an award from Arthritis Society Canada (STAR-20-0000000012).

Disclosure of Potential Conflicts of Interest

KR declares current employment status with STEMCELL Technologies. SV declares 60% ownership of Regulatory Cell Therapy Consultants Inc., which does not conflict with this paper in any way. SV, RR and KR are co-inventors on patent applications filed with UHN based on this work. The remaining authors declare they have no competing interests.

Author contributions:

Kevin P. Robb: Conception & design, collection and/or assembly of data, data analysis and interpretation, manuscript writing, final approval of manuscript
Razieh Rabani: Conception & design, collection and/or assembly of data, data analysis and interpretation
Shabana Vohra: data analysis and interpretation
Shoba Singh: administrative support, collection and/or assembly of data
Jaskarndip Chahal: collection and/or assembly of data, provision of study materials or patients
Julie Audet: data analysis and interpretation
Anthony V. Perruccio: data analysis and interpretation
Ali Naragahi: data analysis
Osvaldo Espin-Garia: data analysis
Rajiv Gandhi: financial support, provision of study materials or patients
Sowmya Viswanathan: Conception & design, manuscript writing, final approval of manuscript

References

[1] Cui A, Li H, Wang D, Zhong J, Chen Y, Lu H. Global, regional prevalence, incidence and risk factors of knee osteoarthritis in population-based studies. EClinicalMedicine 2020;29–30. https://doi.org/10.1016/j.eclinm.2020.100587.

[2] Loeser RF, Goldring SR, Scanzello CR, Goldring MB. Osteoarthritis: a disease of the joint as an organ. Arthritis Rheum 2012;64:1697–707. https://doi.org/10.1002/art.34453.

[3] Zhang X, He J, Wang W. Progress in the use of mesenchymal stromal cells for osteoarthritis treatment. Cytotherapy 2021;23:459–70. https://doi.org/10.1016/j.jcyt.2021.01.008.

[4] Copp G, Robb KP, Viswanathan S. Culture-expanded mesenchymal stromal cell therapy: does it work in knee osteoarthritis? A pathway to clinical success. Cell Mol Immunol 2023. https://doi.org/10.1038/S41423-023-01020-1.

[5] Mautner K, Gottschalk M, Boden SD, Akard A, Bae WC, Black L, et al. Cell-based versus corticosteroid injections for knee pain in osteoarthritis: a randomized phase 3 trial. Nature Medicine 2023 2023:1–7. https://doi.org/10.1038/s41591-023-02632-w.

[6] Soler R, Orozco L, Munar A, Huguet M, López R, Vives J, et al. Final results of a phase I–II trial using ex vivo expanded autologous Mesenchymal Stromal Cells for the treatment of osteoarthritis of the knee confirming safety and suggesting cartilage regeneration. Knee 2016;23:647–54. https://doi.org/10.1016/j.knee.2015.08.013.

[7] Chen HH, Chen YC, Yu SN, Lai WL, Shen YS, Shen PC, et al. Infrapatellar fat pad-derived mesenchymal stromal cell product for treatment of knee osteoarthritis: a first-in-human study with evaluation of the potency marker. Cytotherapy 2022;24:72–85. https://doi.org/10.1016/j.jcyt.2021.08.006.

[8] Chahal J, Gómez⎕Aristizábal A, Shestopaloff K, Bhatt S, Chaboureau A, Fazio A, et al. Bone Marrow Mesenchymal Stromal Cells in Patients with Osteoarthritis Results in Overall Improvement in Pain and Symptoms and Reduces Synovial Inflammation. Stem Cells Transl Med 2019;8:746–757. https://doi.org/10.1002/sctm.18-0183.

[9] Matas J, García C, Poblete D, Vernal R, Ortloff A, Luque-Campos N, et al. A Phase I Dose-Escalation Clinical Trial to Assess the Safety and Efficacy of Umbilical Cord-Derived Mesenchymal Stromal Cells in Knee Osteoarthritis. Stem Cells Transl Med 2024;13:193–203. https://doi.org/10.1093/STCLTM/SZAD088.

[10] Yubo M, Yanyan L, Li L, Tao S, Bo L, Lin C. Clinical efficacy and safety of mesenchymal stem cell transplantation for osteoarthritis treatment: A meta-analysis. PLoS One 2017;12:e0175449. https://doi.org/10.1371/journal.pone.0175449.

[11] Cui G-H, Wang YY, Li C-J, Shi C-H, Wang W-S. Efficacy of mesenchymal stem cells in treating patients with osteoarthritis of the knee: A meta-analysis. Exp Ther Med 2016;12:3390–400. https://doi.org/10.3892/etm.2016.3791.

[12] Gupta PK, Maheshwari S, Cherian JJ, Goni V, Sharma AK, Tripathy SK, et al. Efficacy and Safety of Stempeucel in Osteoarthritis of the Knee: A Phase 3 Randomized, Double-Blind, Multicenter, Placebo-Controlled Study. Am J Sports Med 2023;51:2254–66. https://doi.org/10.1177/03635465231180323.

[13] Pico OA, Espinoza F, Cádiz MI, Sossa CL, Becerra-Bayona SM, Salgado MCC, et al. Efficacy of a single dose of cryopreserved human umbilical cord mesenchymal stromal cells for the treatment of knee osteoarthritis:a randomized, controlled, double-blind pilot study. Cytotherapy 2024;0. https://doi.org/10.1016/J.JCYT.2024.09.005/ATTACHMENT/201E9836-BC53-4964-AEB9-F674366CBCFB/MMC2.DOCX.

[14] Vives J, Oliver-Vila I, Pla A. Quality compliance in the shift from cell transplantation to cell therapy in non-pharma environments. Cytotherapy 2015;17:1009–14. https://doi.org/10.1016/j.jcyt.2015.02.002.

[15] Siegel G, Kluba T, Hermanutz-Klein U, Bieback K, Northoff H, Schäfer R. Phenotype, donor age and gender affect function of human bone marrow-derived mesenchymal stromal cells. BMC Med 2013;11:146. https://doi.org/10.1186/1741-7015-11-146.

[16] Wilson A, Hodgson-Garms M, Frith JE, Genever P. Multiplicity of mesenchymal stromal cells: Finding the right route to therapy. Front Immunol 2019;10:1112. https://doi.org/10.3389/fimmu.2019.01112.

[17] Viswanathan S, Galipeau J. Hallmarks of MSCs: Key quality attributes for pharmacology and clinical use. Cell Stem Cell 2025;32:878–94. https://doi.org/10.1016/J.STEM.2025.05.008.

[18] Robb KP, Audet J, Gandhi R, Viswanathan S. Putative critical quality attribute matrix identifies mesenchymal stromal cells with potent immunomodulatory and angiogenic "fitness" ranges in response to culture process parameters. Front Immunol 2022;13. https://doi.org/10.3389/fimmu.2022.972095.

[19] Asgarpour K, Shojaei Z, Amiri F, Ai J, Mahjoubin-Tehran M, Ghasemi F, et al. Exosomal microRNAs derived from mesenchymal stem cells: Cell-to-cell messages. Cell Communication and Signaling 2020;18. https://doi.org/10.1186/s12964-020-00650-6.

536	[20]	Mobasheri A, Saarakkala S, Finnilä M, Karsdal MA, Bay-Jensen AC, van Spil WE. Recent advances in
537		understanding the phenotypes of osteoarthritis. F1000Res 2019;8.
538		https://doi.org/10.12688/f1000research.20575.1.
539	[21]	Dell'Isola A, Allan R, Smith SL, Marreiros SSP, Steultjens M. Identification of clinical phenotypes in
540		knee osteoarthritis: a systematic review of the literature. BMC Musculoskelet Disord 2016;17:1–12.
541		https://doi.org/10.1186/s12891-016-1286-2.
542	[22]	Mobasheri A, Bay-Jensen AC, Gualillo O, Larkin J, Levesque MC, Henrotin Y. Soluble biochemical
543		markers of osteoarthritis: Are we close to using them in clinical practice? Best Pract Res Clin
544		Rheumatol 2017;31:705–20. https://doi.org/10.1016/j.berh.2018.07.002.
545	[23]	Peterfy CG, Guermazi A, Zaim S, Tirman PFJ, Miaux Y, White D, et al. Whole-organ magnetic
546		resonance imaging score (WORMS) of the knee in osteoarthritis. Osteoarthritis Cartilage 2004;12:177–
547		90. https://doi.org/10.1016/j.joca.2003.11.003.
548	[24]	Guermazi A, Roemer FW, Hayashi D, Crema MD, Niu J, Zhang Y, et al. Assessment of synovitis with
549		contrast-enhanced MRI using a whole-joint semiquantitative scoring system in people with, or at high
550		risk of, knee osteoarthritis: the MOST study. Ann Rheum Dis 2011;70:805–11.
551		https://doi.org/10.1136/ARD.2010.139618.
552	[25]	Pham T, van der Heijde D, Altman RD, Anderson JJ, Bellamy N, Hochberg M, et al. OMERACT-
553		OARSI initiative: Osteoarthritis research society international set of responder criteria for osteoarthritis
554		clinical trials revisited. Osteoarthritis Cartilage 2004;12:389–99.
555		https://doi.org/10.1016/j.joca.2004.02.001.
556	[26]	Krampera M. Mesenchymal stromal cell licensing: A multistep process. Leukemia 2011;25:1408–14.
557		https://doi.org/10.1038/leu.2011.108.
558	[27]	Marx RG, Stump TJ, Jones EC, Wickiewicz TL, Warren RF. Development and evaluation of an activity
559		rating scale for disorders of the knee. American Journal of Sports Medicine 2001;29:213–8.
560		https://doi.org/10.1177/03635465010290021601.
561	[28]	Kasikis S, Baez J, Gandhi I, Grupp S, Kitko CL, Kowalyk S, et al. Mesenchymal stromal cell therapy
562		induces high responses and survival in children with steroid refractory GVHD and poor risk biomarkers.
563		Bone Marrow Transplant 2021;56:2869. https://doi.org/10.1038/S41409-021-01442-3.
564	[29]	Perin EC, Borow KM, Henry TD, Mendelsohn FO, Miller LW, Swiggum E, et al. Randomized Trial of
565		Targeted Transendocardial Mesenchymal Precursor Cell Therapy in Patients With Heart Failure. J Am
566		Coll Cardiol 2023;81:849–63. https://doi.org/10.1016/J.JACC.2022.11.061.
567	[30]	Ouboter LF, Barnhoorn MC, Verspaget HW, Plug L, Pool ES, Szuhai K, et al. Local administration of
568		mesenchymal stromal cells is safe and modulates the immune compartment in ulcerative proctitis. JCI
569		Insight 2023;8. https://doi.org/10.1172/JCI.INSIGHT.167402.
570	[31]	Smith SR, Deshpande BR, Collins JE, Katz JN, Losina E. Comparative pain reduction of oral non-
571		steroidal anti-inflammatory drugs and opioids for knee osteoarthritis: Systematic analytic review.
572		Osteoarthritis Cartilage 2016;24:962–72. https://doi.org/10.1016/j.joca.2016.01.135.
573	[32]	Gómez-Aristizábal A, Gandhi R, Mahomed NN, Marshall KW, Viswanathan S. Synovial fluid
574		monocyte/macrophage subsets and their correlation to patient-reported outcomes in osteoarthritic
575		patients: A cohort study. Arthritis Res Ther 2019;21. https://doi.org/10.1186/s13075-018-1798-2.
576	[33]	Boregowda S V., Krishnappa V, Haga CL, Ortiz LA, Phinney DG. A Clinical Indications Prediction
577		Scale Based on TWIST1 for Human Mesenchymal Stem Cells. EBioMedicine 2016;4:62–73.
578		https://doi.org/10.1016/j.ebiom.2015.12.020.
579	[34]	Lee RH, Boregowda S V., Shigemoto-Kuroda T, Bae EH, Haga CL, Abbery CA, et al. TWIST1 and
580		TSG6 are coordinately regulated and function as potency biomarkers in human MSCs. Sci Adv
581		2023;9:eadi2387. https://doi.org/10.1126/SCIADV.ADI2387.
582	[35]	Kuznetsov SA, Mankani MH, Bianco P, Robey PG. Enumeration of the colony-forming units-fibroblast
583		from mouse and human bone marrow in normal and pathological conditions. Stem Cell Res 2009;2:83–
584		94. https://doi.org/10.1016/j.scr.2008.07.007.
585	[36]	Siegel G, Kluba T, Hermanutz-Klein U, Bieback K, Northoff H, Schäfer R. Phenotype, donor age and
586		gender affect function of human bone marrow-derived mesenchymal stromal cells. BMC Med 2013;11.
587		https://doi.org/10.1186/1741-7015-11-146.
588	[37]	Driban JB, McAlindon TE, Amin M, Price LL, Eaton CB, Davis JE, et al. Risk factors can classify
589		individuals who develop accelerated knee osteoarthritis: Data from the osteoarthritis initiative. Journal
590		of Orthopaedic Research 2018;36:876–80. https://doi.org/10.1002/jor.23675.
591	[38]	Hamilton AM, Cheung W-Y, Gómez-Aristizábal A, Sharma A, Nakamura S, Chaboureau A, et al. Iron
592		nanoparticle-labeled murine mesenchymal stromal cells in an osteoarthritic model persists and suggests
593		anti-inflammatory mechanism of action. PLoS One 2019;14:e0214107.
594		https://doi.org/10.1371/journal.pone.0214107.

[39] Zhang Y, Ji Q. Macrophage polarization in osteoarthritis progression: a promising therapeutic target.
Front Cell Dev Biol 2023;11:1269724. https://doi.org/10.3389/FCELL.2023.1269724.
[40] Liu B, Zhang M, Zhao J, Zheng M, Yang H. Imbalance of M1/M2 macrophages is linked to severity
level of knee osteoarthritis. Exp Ther Med 2018;16:5009–14. https://doi.org/10.3892/ETM.2018.6852.
[41] Chan MWY, Viswanathan S. Recent progress on developing exogenous monocyte/macrophage-based
therapies for inflammatory and degenerative diseases. Cytotherapy 2019;21:393–415.
https://doi.org/10.1016/J.JCYT.2019.02.002.
[42] Rabani R, Volchuk A, Jerkic M, Ormesher L, Garces-Ramirez L, Canton J, et al. Mesenchymal stem
cells enhance NOX2-dependent reactive oxygen species production and bacterial killing in macrophages
during sepsis. European Respiratory Journal 2018;51:1702021.
https://doi.org/10.1183/13993003.02021-2017.
[43] Rasti M, Barazandeh AF, Robb KP, Low R, Gandhi R, Viswanathan S. Mesenchymal Stromal Cells
Immunosuppress Osteoarthritis Synovial Fluid Tolerized Monocytes via IL-6. BioRxiv
2025:2025.08.14.669875. https://doi.org/10.1101/2025.08.14.669875.
[44] Vincent TL. Peripheral pain mechanisms in osteoarthritis. Pain 2020;161:S138.
https://doi.org/10.1097/J.PAIN.0000000000001923.
[45] Robb KP, Galipeau J, Shi Y, Schuster M, Martin I, Viswanathan S. Failure to launch commercially-
approved mesenchymal stromal cell therapies: what's the path forward? Proceedings of the International
Society for Cell & Gene Therapy (ISCT) Annual Meeting Roundtable held in May 2023, Palais des
Congrès de Paris, Organized by the ISCT MSC Scientific Committee. Cytotherapy 2023.
https://doi.org/10.1016/J.JCYT.2023.09.001.
[46] Kraus VB, Collins JE, Hargrove D, Losina E, Nevitt M, Katz JN, et al. Predictive validity of
biochemical biomarkers in knee osteoarthritis: Data from the FNIH OA Biomarkers Consortium. Ann
Rheum Dis 2017;76:186–95. https://doi.org/10.1136/annrheumdis-2016-209252.
[47] Yang YK, Ogando CR, Wang See C, Chang TY, Barabino GA. Changes in phenotype and
differentiation potential of human mesenchymal stem cells aging in vitro. Stem Cell Res Ther
2018;9:131. https://doi.org/10.1186/s13287-018-0876-3.
[48] Perry TA, Deng Y, Hulley P, Maciewicz RA, Mitchelmore J, Larsson S, et al. Deconvoluting synovial
fluid molecular endotypes in knee osteoarthritis: primary results from the STEpUP OA Consortium.
MedRxiv 2024:2024.06.05.24308485. https://doi.org/10.1101/2024.06.05.24308485.
[49] Wang SX, Abramson SB, Attur M, Karsdal MA, Preston RA, Lozada CJ, et al. Safety, tolerability, and
pharmacodynamics of an anti-interleukin-1α/β dual variable domain immunoglobulin in patients with
osteoarthritis of the knee: a randomized phase 1 study. Osteoarthritis Cartilage 2017;25:1952–61.
https://doi.org/10.1016/J.JOCA.2017.09.007.
[50] Karsdal MA, Byrjalsen I, Alexandersen P, Bihlet A, Andersen JR, Riis BJ, et al. Treatment of
symptomatic knee osteoarthritis with oral salmon calcitonin: results from two phase 3 trials.
Osteoarthritis Cartilage 2015;23:532–43. https://doi.org/10.1016/J.JOCA.2014.12.019.
[51] Mobasheri A, Loeser R. Clinical phenotypes, molecular endotypes and theratypes in OA therapeutic
development. Nat Rev Rheumatol 2024;20:525–6. https://doi.org/10.1038/S41584-024-01126-4.
[52] Angelini F, Widera P, Mobasheri A, Blair J, Struglics A, Uebelhoer M, et al. Osteoarthritis endotype
discovery via clustering of biochemical marker data. Ann Rheum Dis 2022;81:666–75.
https://doi.org/10.1136/ANNRHEUMDIS-2021-221763/ATTACHMENT/2A2BDE9F-17E0-411E-
AF92-9C4634CD081C/MMC5.PDF.
[53] Riddle DL, Jiranek WA. Knee osteoarthritis radiographic progression and associations with pain and
function prior to knee arthroplasty: A multicenter comparative cohort study. Osteoarthritis Cartilage
2015;23:391–6. https://doi.org/10.1016/j.joca.2014.12.013.
[54] Riddle DL, Jiranek WA. Knee Osteoarthritis Radiographic Progression and Associations with Pain and
Function Prior to Knee Arthroplasty: A Multicenter Comparative Cohort Study. Osteoarthritis and
Cartilage / OARS, Osteoarthritis Research Society 2015;23:391.
https://doi.org/10.1016/J.JOCA.2014.12.013.
645
646

647 **Table 1. Summary of baseline patient characteristics.** Association between each covariate was evaluated
648 against responder status (p-values adjusted for multiple comparisons); no significant associations were detected.
649 BMI: body mass index; KL: Kellgren-Lawrence; WORMS: Whole-Organ Magnetic Resonance Imaging Score
650 [23].

Covariate	Full Sample (n=12 total)	F Responder (n=7 total)	Non-Responder (n=5 total)	Adjusted p-value
Cell dose				0.495
1×10^6 [n, (%)]	4 (33)	2 (29)	2 (40)	
10×10^6 [n, (%)]	4 (33)	1 (14)	3 (60)	
50×10^6 [n, (%)]	4 (33)	4 (57)	0 (0)	
Gender				0.513
Man $x10^6$ [n, (%)]	7 (58)	3 (43)	4 (80)	
Woman $x10^6$ [n, (%)]	5 (42)	4 (57)	1 (20)	
Age				0.601
Mean (SD)	54.8 (6.9)	55.7 (5.6)	53.4 (8.8)	
Median (Min,Max)	53.5 (45.0,65.0)	54 (47,63)	51 (45,65)	
BMI				0.494
Mean (SD)	25.8 (3.5)	24.5 (3.2)	27.5 (3.4)	
Median (Min,Max)	25.5 (19.0,29.9)	24.9 (19.0,29.7)	29.7 (22.1,29.9)	
KL Grade				0.583
3	11 (92)	7 (100)	4 (80)	
4	1 (8)	0 (0)	1 (20)	
MRI Total Synovitis Score [24]				0.494
Mean (sd)	6.8 (2.9)	5.7 (2.1)	9.3 (3.1)	
Median (Min,Max)	6 (3,12)	5 (3,9)	10 (6,12)	
Missing	2	0	2	
MRI WORMS				0.756
Mean (SD)	37.4 (9.3)	36.4 (8.3)	38.8 (11.3)	
Median (Min,Max)	36.2 (20.0,54.0)	36.0 (20.0,45.5)	36.5 (28.0,54.0)	

651
652
653

Figure Legends

Fig. 1. Osteoarthritis patient responders demonstrate improvements in KOOS subscales that are maintained out to 24 months follow-up post-injection. A) Patients categorized as Function Responders displayed significant improvements (indicated by lower score values) in KOOS subscales and mean KOOS at 12- and 24-months follow-up relative to baseline, while Non-Responders displayed no significant changes. KOOS scales were inverted such that higher scores represent greater knee-related issues. B) Flow chart summary of patient responder status at 12-months, 24-months, and 8-10-years follow-up. C) Summary of OARSI-OMERACT Responder classification in 2025 after 8-10 years of follow-up. Responder status is indicated as Function-Pain (F-P), Function (F), or Pain (P) Responders. Changes in patient responder status at 8-10 years are bolded. Patient-reported follow-up notes on additional interventions are included in the right column. N=12 patients. Two-way repeated measures ANOVA, Dunnett's multiple comparisons test. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001; asterisks indicate significant differences for responder group relative to baseline at the time points indicated. ADL: function in daily living; Sports/Rec: function in sport and recreation; QOL: knee-related quality of life; KOOS: knee injury and osteoarthritis outcome score; PRP: platelet-rich plasma; TKR: total knee replacement.

Fig. 2. Independent validation that a previously identified gene panel distinguishes clinically effective MSC(M) donors from non-responder MSC(M) donors. A) Principal component (PC) analysis of MSC(M) expression of an immunomodulatory gene panel identified previously for MSC(AT) (*TNFAIP6*, *ACTA2*, *ANGPT1*, *CXCL8*, *EDN1*, *THBS1*, *PDCD1LG2*, *CCN2*; [18]) expressed under licensed conditions. The corresponding loading plot of eigenvectors (right) indicates the relative contribution of each gene to the PC1 and PC2 axes. B) MSC(M) derived from Function Responders displayed significantly lower PC1 scores relative to non-responders, indicating that the gene panel sensitively distinguishes MSC(M) from donors classified as responders. Student's t-test, ***p<0.001, ****p<0.0001. Horizontal lines: group mean; error bars: standard deviation. C) Significant negative correlations between PC1 scores and the percentage change at 24-months relative to baseline for KOOS ADL, pain, symptoms, and mean KOOS. Percent change in KOOS values were made relative to baseline and calculated such that positive values represent improvement. Spearman's correlation. N=9 MSC(M) donors; n=2-3 replicates/donor.

Fig. 3. *In vitro* monocyte/macrophage polarization readouts distinguish donor MSC(M) immunomodulatory fitness for responders versus non-responders. A) Schematic of *in vitro* monocyte/macrophage (MΦ) polarization experiment. Indirect co-cultures of MSC(M) with human peripheral blood-derived MΦ was performed for 48 h prior to removal of MSC(M) and addition of lipopolysaccharide (LPS) for 4 h incubation. B) Summary of MΦ gene expression after co-culture with Function Responders MSC(M). Box-and-whisker plots (left) display changes in gene expression relative to MΦ cultured alone (Solo; dotted line). Horizontal line: median; hinges: first and third quartiles; whiskers: range. Two-way ANOVA, Tukey's post-hoc test. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 relative to Solo condition or to groups indicated by brackets. Discriminant canonical plots (right) of -ddCt values relative to Solo condition. Inner ellipses indicate 95% confidence for the mean of each group;

outer ellipses indicate the normal region estimated to contain 50% of the population for each group.

Fig. 4. microRNA-sequencing reveals microRNA signatures in Function Responder donor MSC(M). A) Principal component analysis of microRNA expression profiles demonstrate clustering in Function Responders versus Non-Responders. MSC(M) samples cluster by donor and responder status rather than by synovial fluid (SF) stimulation. B) Volcano plot displays 33 significantly differentially expressed microRNAs in MSC(M) derived from donors classified as Function Responders versus Non-Responders. C&D) Top 30 pathways identified through Reactome analysis of predicted gene targets for the top ten significantly upregulated (C) and downregulated (D) microRNAs. E&F) Top 30 pathways identified through Gene Ontology (Biological Processes) analysis of predicted gene targets for the top ten significantly upregulated (E) and downregulated (F) microRNAs. G) Significant correlations between microRNAs and the delta or percentage change at 12- and 24-months relative to baseline for KOOS ADL, pain, and mean KOOS. Delta and percent change in KOOS values were made relative to baseline and calculated such that positive values represent improvement. Spearman's correlation. N=9 MSC(M) donors.

Fig. 5. More severe KOOS and lower MARS physical activity scores correlate to improved patient-reported outcomes at 12- and 24-months follow-up. A) Patients categorized according to Function Responders displayed significantly more severe mean KOOS and Sports/Rec scores at baseline relative to Non-Responders. B&C) Baseline MARS significantly correlated to baseline Sports/Rec, as well as both delta and percent change in KOOS Pain at 12 months relative to baseline. KOOS scales were inverted such that higher scores represent greater knee-related issues; delta and percent change in KOOS values were made relative to baseline and calculated such that positive values represent improvement. N=12 patients. Student's t-test (for baseline Mean KOOS), Two-way ANOVA, Sidak's post-hoc test (for baseline KOOS Subscales), Spearman's correlation. $*p<0.05$, $**p<0.01$. n.s.; non-significant. ADL: function in daily living; Sports/Rec: function in sport and recreation; QOL: knee-related quality of life; Pt.: patient ID; MO: month; KOOS: knee injury and osteoarthritis outcome score; MARS: Marx Activity Rating Scale.

FIG. 1

A

Function Responders

B

12-month:	8 Responders		4 Non-Responders
24-month:	7 Responders	1 Non-Responder	4 Non-Responders
8-10 year:	5 Responders No intervention: 1 Injection: 2 TKR: 2	1 Non-Responder No intervention	1 Non-Responder No intervention	3 Non-Responders No intervention: 1 Injection: 2

1 lost to follow-up (TKR)
1 lost to follow-up

C

Subject ID	Responder Class at 12 Months	Responder Class at 24 Months	Responder Class at 8-10 years	# Years Follow-up in 2025	8-10 year follow up scores				Follow-up notes
					KOOS Pain		KOOS ADL		
					Delta	% Change	Delta	% Change	
1	Non-Responder	Non-Responder	Non-Responder	9 years, 10 months	-0.04	-0.21	-1.49	-16.93	Injection of viscosupplementation in 2024 and 2025
2	Non-Responder	Non-Responder	N/A	-	-	-	-	-	N/A
3	Responder (F-P)	Responder (F)	Responder (F-P)	9 years, 6 months	30.52	91.65	27.96	90.49	PRP injection in 2023
4	Non-Responder	Non-Responder	Non-Responder	9 years, 4 months	-44.40	-792.86	-30.88	-3088.00	No surgical intervention
5	Responder (P)	Responder (P)	N/A	-	-	-	-	-	TKR in 2025
6	Responder (P)	Pain Responder	Responder (P)	9 years, 0 months	11.11	22.22	-22.07	-501.59	Awaiting bilateral knee replacement
7	Responder (F-P)	Responder (F-P)	**Non-Responder**	8 years, 11 months	0.04	0.13	-1.46	-7.09	No intervention post injection
9	Responder (F-P)	Non-Responder	**Non-Responder**	8 years, 6 months	-2.82	-4.06	-19.14	-43.40	No intervention post injection
10	Responder (F)	Responder (F-P)	Responder (F-P)	8 years, 6 months	19.41	33.29	20.61	50.02	Viscosupplementation - no majory surgery
11	Responder (F-P)	Responder (F-P)	Responder (F-P)	8 years, 6 months	24.96	56.22	16.16	54.97	No knee surgery/scope, no knee injection
12	Non-Responder	Non-Responder	Non-Responder	8 years, 4 months	-30.54	-78.51	-24.95	-77.01	Viscosupplementation in 2019
13	Responder (F-P)	Responder (F-P)	Responder (F-P)	8 years, 2 months	49.99	81.82	39.74	77.17	TKR in 2025

FIG. 2

FIG. 3

FIG. 5

What determines osteoarthritis patient responses to autologous MSC(M) injection?

Responders

Non-Responders

MSC(M) Basal Fitness

Targeted gene expression panel

In vitro MΦ polarization

microRNA-Sequencing

Baseline clinical phenotype

Knee Injury & Osteoarthritis Outcome Scores (KOOS) Marx Physical Activity Scores

Baseline molecular biomarkers

Local biomarkers Synovial Fluid

Systemic biomarkers Plasma, Serum, Urine

Responders versus Non-Responders to MSC(M) therapy

Increased MSC(M) Immunomodulatory Basal Fitness

Immunomodulatory CQA transcript expression

Distinct MΦ Polarization

Expression of distinct miR profile

More severe KOOS Sports/Rec

Lower levels of physical activity

Local biomarkers
Lower baseline angiogenic markers (non-significant)

Systemic biomarkers
Non-significant