No neurobehavioral evidence for reduced motivational potential of social rewards in alcohol use disorder

Yana Schwarze[1,2], Janine Stierand[3], Johanna Voges[4], Emil Zillikens[1], Klaus Junghanns[3], Oliver Voß[5], Frieder Michel Paulus[1,2], Sören Krach[1,2], Maurice Cabanis[6]*, Lena Rademacher[1,2,6]*

[1] Social Neuroscience Lab, Department of Psychiatry and Psychotherapy, University of Lübeck, Lübeck, Germany

[2] Center of Brain, Behavior and Metabolism (CBBM), University of Lübeck, Lübeck, Germany

[3] Department of Psychiatry and Psychotherapy, University of Lübeck, Lübeck, Schleswig-Holstein, Germany

[4] Department of Psychosomatic Medicine, Charité - Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Berlin, Germany

[5] AMEOS Clinic Lübeck, Department of Substance Use Disorders, Lübeck, Germany

[6] Center of Mental Health, Hospital for Addiction and Addictive Behavior, Klinikum Stuttgart, Stuttgart, Germany

*both authors contributed equally

Corresponding authors: Yana Schwarze and Lena Rademacher, Department of Psychiatry and Psychotherapy, Social Neuroscience Lab, Center of Brain, Behavior and Metabolism, Ratzeburger Allee 160, 23562 Lübeck, Phone: +49 451 31017512 Email: yana.schwarze@uni-luebeck.de, l.rademacher@uni-luebeck.de

Running title: Social reward anticipation in AUD

Key words: alcohol use disorder, social reward, reward anticipation, incentive delay task, fMRI

Word count: 3900

Data and code available at: https://osf.io/g3ysp/

Abstract

The mesolimbic dopamine system plays a central role in motivating behavior. In alcohol use disorder (AUD), this system is thought to be dysfunctional, leading to hyperreactivity to alcohol-related cues. In contrast, evidence on how individuals with AUD respond to alcohol-unrelated reward cues is inconclusive, and the motivation for social rewards has not yet been investigated. To address this gap, 36 individuals with AUD and 34 healthy controls performed an incentive delay task to assess social reward anticipation with a monetary and a non-reward control condition while undergoing functional magnetic resonance imaging. The ventral striatum was defined as region of interest because of its central role in neuronal circuits for motivation. Neither behavioral nor neuroimaging data provided any evidence of altered motivation for social or monetary rewards in AUD. Exploratory whole-brain analyses only revealed stronger activation in the occipital/cuneal cortex in individuals with AUD than in healthy controls across all trials. Together, these results suggest that sensitivity to social reward cues is not fundamentally impaired in AUD. Furthermore, they imply that motivational changes related to the substance do not generally alter the reward potential of alcohol-unrelated domains in AUD, opening perspectives for social-behavioral treatments for this disorder.

Introduction

The mesolimbic reward system – dopamine neurons projecting from the ventral tegmental area (VTA) to the ventral striatum – is the core of neural circuits for motivation (Stuber, 2023). Through the dopamine release, incentive salience is attributed to reward-predictive cues, making the stimulus a desirable target and initiating approach behavior (Berridge, 2012). In alcohol use disorders (AUD), this circuit is thought to be dysfunctional. According to the incentive sensitization theory (Robinson and Berridge, 1993; Robinson and Berridge, 2025) the dopaminergic reward system in addiction is hyperreactive to drug cues and contexts, leading to excessive "wanting" to take the drug. In line with this, functional magnetic resonance imaging (fMRI) studies examining drug cue reactivity in substance use disorders have shown aberrations in neural activation that correlate with clinical outcomes (Addiction Cue-Reactivity Initiative (ACRI) Network *et al.*, 2024). With regard to AUD, a review by Cofresi and colleagues (2019) concluded that there is "sufficient empirical evidence to support the idea that the mechanism originally outlined in the incentive salience sensitization theory of addiction [...] may be at the core of alcohol use disorder (AUD) etiology for some individuals" (pp. 912-913).

In addition to the increased incentive salience for drug-associated stimuli in substance use disorders, a reduced motivation for other, drug-unrelated rewards is frequently hypothesized, because the substance may "hijack" the neural reward system and reorganize the priority of reward processing (Wrase *et al.*, 2007; Tan *et al.*, 2024). A number of studies have examined the motivation for alcohol-unrelated, in most cases monetary, rewards in alcohol use disorder. For this purpose, the monetary incentive delay (MID) task (Knutson *et al.*, 2000) is widely used. However, findings are mixed: Some studies report reduced ventral striatal activity during the anticipation of monetary rewards in AUD (Hägele *et al.*, 2015; Romanczuk-Seiferth *et al.*, 2015), while others found no group differences (Bjork *et al.*, 2012; Musial *et al.*, 2023) or even increased activity (Muench *et al.*, 2018).

Social rewards are another type of alcohol-unrelated reward that may be of particular interest in AUD. Substance use alters social behaviors, and the mesolimbic reward system is considered to play

a central part in this process (Young *et al.*, 2011). The mechanisms are not clear, but it has been suggested that dopamine transmission in the ventral striatum mediates the rewarding aspects of social interactions and that disruptions to this process as a result of substance use affect social behaviour (Young *et al.*, 2011). Alternatively, low social integration could result in heavier use of alcohol, a dysfunctional health behavior to cope with experienced social isolation (Havassy *et al.*, 1991). However, while addiction neuroscience research has gained interest on social interaction and connectedness (Delgado *et al.*, 2023), up to this date the incentive salience of social rewards per se has not yet been investigated in AUD patients. This is surprising given the importance of social factors for the treatment of addiction (Heilig *et al.*, 2016). The present study aims to bridge this gap and investigate the motivational salience of social rewards in patients diagnosed with AUD using a well established social adaptation (Spreckelmeyer *et al.*, 2009) of the monetary incentive delay task by Knutson et al. (2000).

Methods

Sample

Forty-two individuals diagnosed with moderate or severe AUD according to DSM-5 criteria (meeting 5–11 criteria) and 38 healthy control (HC) participants were recruited between February 2019 and October 2022 (with a break between March 2020 and October 2021 due to the SARS-CoV-19 pandemic) through the Department of Psychiatry and Psychotherapy of the University of Lübeck and the AMEOS clinic in Lübeck, Germany. Healthy controls were recruited through flyers, letters, and online advertisements. All patients were in the early abstinence phase (between the 10th and 40th day) following the resolution of acute withdrawal symptoms and remained in the clinic for the duration of the study. One patient and one control subject dropped out of the study. In addition, three patients and two control subjects had to be excluded from the analyses because of technical problems, excessive head motion, or sleeping. Furthermore, two patients and one control subject did not participate in magnetic resonance imaging (MRI) scanning for safety reasons and were therefore excluded from this part of the study. Thus, the final sample included 36 patients and 34 healthy controls. Both groups were matched for age and gender. Written informed consent was obtained from all participants and the study was approved by the local ethics committee of the University of Lübeck (AZ 17-077).

Participants were required to meet the following criteria (which were in part due to other research questions in the project not addressed here): age between 18 and 60 years, sufficient knowledge of the German language, scanner compatibility, no neurological disorders or brain injuries, no medication that can affect the HPA-axis or dopamine system in the past two weeks, no acute infections, no vaccination in the past three weeks, and no chronic inflammatory diseases. Additionally, patients had to meet the following criteria: diagnosis of alcohol use disorder according to the Diagnostic and Statistical Manual of Mental Disorders fifth edition (DSM-V), alcohol abstinence for at least 10 days, no comorbid diagnosis of major depression, antisocial or borderline personality disorders, social anxiety, psychosis, or acute suicidality. The inclusion criteria for control

participants were as follows: no current or past psychiatric disorders, no alcohol misuse classified as ≥ 8 points in the alcohol use disorder identification test (1), no consumption of drugs (except for alcohol, nicotine, and irregular use of cannabis) in the past year, and no history of regular intake/dependency of any drug except nicotine.

Procedure

Before study participation, all participants were screened to ensure compliance with the inclusion and exclusion criteria. Once eligible, participants attended the Center of Brain, Behavior and Metabolism in Lübeck on two separate days at the same time each day (starting at 8:30 AM ± 30 minutes). The two test days were identical in structure and included a MRI measurement, as well as numerous other measurements that were collected in the context of other research questions, such as blood samples, saliva samples, and questionnaires. The difference between the two days was that stress was induced in the participants on one day and not on the other (with the order of days counterbalanced across participants). Stress effects are not included in the present analyses and have been published elsewhere (Schwarze *et al.*, 2024). For the present analyses, only the MRI measurements of the "control day" without stress induction are examined.

Social and Monetary Incentive Delay tasks

During the MRI scan, participants completed a reward paradigm that was an adaptation of the Social Incentive Delay (SID) task by Spreckelmeyer et al. (2009) and the Monetary Incentive Delay (MID) task by Knutson et al. (2000). Before the scan, the paradigm was practiced outside the MRI to ensure participants were familiar with the task. The experiment consisted of four blocks (two MID and two SID blocks) of 20 trials each, resulting in a total of 80 trials. Each trial began with a cue presentation lasting 1000 ms, indicating either a potential reward (20 trials per task, represented by a circle with a line: ⵣ) or no reward (20 trials per task, represented by an empty circle: O, see Fig. 1). This was followed by a delay period of variable length (jittered between 2025 and 3240 ms) before the target

appeared. Participants were instructed to respond as quickly as possible to all target appearances, regardless of the cue type. When participants responded quickly enough in a reward trial, a picture of a smiling face (SID task) or a picture of a wallet with small amounts of money (MID task) was presented for 2500 ms. If the response was too slow, participants saw a blurred image instead. In no-reward trials, blurred images were always presented, independent of the response time. Task difficulty was standardized to a hit rate of approximately 66% by adjusting the target duration based on individual response times. If participants reacted to the flash in time, the response time window for that cue type (monetary or social, reward or no reward) was shortened by 5% and extended by 10% otherwise. The initial response time window was set to the mean response time from six practice trials conducted before the experiment plus 5%. The maximum response time was set to 500 ms for all participants. The trials in each block were pseudo-randomly arranged with inter-trial intervals jittered between 3235 and 4450 ms. MID and SID blocks were interleaved, always starting with a MID block and ending with a SID block. At the start of each block, participants were informed whether the upcoming task was MID or SID. The total duration of the paradigm was approximately 18 minutes.

MRI acquisition

Participants were examined in the MRI scanner at the Center of Brain, Behavior and Metabolism in Lübeck (3T Siemens MAGNETOM Skyra magnetic resonance tomography) using a 64-channel head coil. A single-shot gradient-recalled echo-planar imaging (GRE-EPI) sequence was used to acquire functional images (repetition time (TR) = 1330 ms, echo time (TE) = 25 ms, flip angle = 70°, field of view (FoV) = 560x560). Additionally, an anatomical T1 image was obtained for each participant, which was acquired using an MP-RAGE sequence (TR = 1900 ms, TE = 2.44 ms, TI = 900 ms, flip angle = 9°, resolution 1×1×1 mm3, FOV = 192×256×256 mm3; acquisition time = 4.5 minutes).

Analysis of behavioral data

For the analysis of response times, all trials with a response time >500 ms were removed, which was the time the target was presented until the feedback was given. Then, outliers deviating more than 3 SD from the mean value of the respective condition of the respective person were excluded (0.95% of total trials), and average response times were calculated for every person and every condition. A three-way repeated-measures ANOVA was calculated with *group* (AUD vs. HC) as the between-subject factor and *task type* (social vs. monetary) and *reward* (reward vs. no reward) as within-subject factors. To correct for possible order effects of the test days, a covariate was included in the analyses, specifying whether the day without stress induction was the first or second test day. To facilitate the interpretation of non-significant results, Bayesian regression models were estimated using the brms package in R (Bürkner, 2017). Bayesian analyses were performed for the non-significant main and interaction effects observed in the ANOVA. Cauchy priors were chosen for the fixed effects, whereas the remaining priors were taken from the default_priors() function of the brms package.

Analysis of fMRI data

MRI data were preprocessed and analyzed using SPM12 (Wellcome Department of Imaging Neuroscience; http://www.fil.ion.ucl.ac.uk/spm) implemented in MATLAB (version R2019b; MathWorks Inc.). Functional images were slice-timed and realigned to the first image to correct for head motion. Next, they were normalized to Montreal Neurological Imaging (MNI) space using standard segmentation as implemented in SPM12 on the mean functional image and tissue probability maps. Normalization parameters were then applied to all images of the time series (resulting voxel size 3 × 3 × 3 mm). Finally, functional images were spatially smoothed with an 8 mm full width half maximum isotropic Gaussian kernel and high-pass filtered at 1/128 Hz to remove low-frequency drifts.

On the first level, a general linear model was created for each participant, which included four regressors with the onset times of the cues of the four conditions (reward and no reward in the social and monetary blocks) and six regressors for the onset times of feedback received. To account for motion-related artifacts, six realignment parameters and their temporal derivatives were included as nuisance regressors. Additionally, a scrubbing regressor was added, removing all volumes with a framewise displacement > 1 mm.

At the group level (second level), a full factorial design was set up in SPM with *group* (AUD vs. HC) as the between-subject and *task type* (social vs. monetary) and *reward* (reward vs no reward) as within-subject factors. A covariate was also set to specify whether the first or second test day was included in the analyses. In case of significant effects, post hoc t-tests were conducted to determine the direction of the observed effects. For whole-brain analyses, we used a false discovery rate (FDR)-corrected cluster-extent threshold of $p < 0.05$ based on a $p < 0.001$ voxel-level threshold as implemented in SPM12. Given the expected large effects of reward, we used a more stringent threshold of family-wise error (FWE)-corrected $p < 0.05$ for the main effect of *reward*.

Due to its central role in motivation, the ventral striatum was defined as a region of interest (ROI) for the analyses of the anticipation phase. Left and right ventral striatum masks were defined as spheres with an 8 mm radius around the peak coordinates (MNI coordinates: left, −10, 10, −2; right, 12, 14, −4) based on a meta-analysis examining reward anticipation in the ventral striatum (Diekhof *et al.*, 2012). Mean parameter estimates were extracted from these ROIs and used in a repeated-measures ANOVA with *group* as the between-subject and *reward* and *task type* as within-subject factors. In parallel with the analysis of response times, Bayesian regression models were also estimated for the extracted parameter estimates to aid in the interpretation of non-significant results.

Results

Sample characteristics

Descriptive data on demographics, alcohol consumption, and response times in the MID and SID tasks are presented in Table 1.

AUD and healthy controls do not differ on the behavioral level

No interaction effects of *group* with *reward* or *task type* were found for response times, suggesting that there was no alteration in reward motivation in AUD (ANOVA *group* x *reward*: $F(1,67) = 0.79$, $p = 0.377$; *group* x *task type*: $F(1,67) = 0.33$, $p = 0.565$; *group* x *reward* x *task type*: $F(1,67) = 2.67$, $p = 0.107$). This was supported by a Bayesian model comparison between the full model and a reduced model that excluded the effect of interest: For all interactions involving *group*, Bayesian analyses provide anecdotal evidence against their inclusion in the model (*group* x *reward*: Bayes factor (BF_{10}) = 0.58, *group* x *task type*: BF_{10} = 0.48, *group* x *reward* x *task type*: BF_{10} = 0.35). Response times across all trials also did not differ between groups (ANOVA main effect of *group*: ($F(1,67) = 0.14$, $p = 0.71$)). However, participants reacted faster to reward trials (mean: 346.1 +- 42.7 ms) compared to no reward trials (mean: 358.9 +- 42.9 ms; main effect of *reward* ($F(1,67) = 50.33$, $p < 0.001$). Additionally, there was a main effect of *task type* ($F(1,67) = 5.04$, $p = 0.028$), with faster response times (in reward and no reward trials) during the MID (mean: 350.7 +- 42.9 ms) compared to the SID task (mean: 354.3 +- 43.6 ms) and a significant *reward* x *task type* interaction ($F(1,67) = 4.28$, $p = 0.042$), reflecting a greater differentiation between reward and no reward trials in the social condition compared to the monetary condition (see Table 1, Fig. 2).

Neural responses to anticipated reward do not differ between AUD and healthy controls

Whole brain analyses revealed no significant interaction effects between *group* and *reward* or *task type*. Significant main effects of *reward* were detected in clusters comprising regions of the striatum,

occipital gyrus, supplementary motor cortex, fusiform gyrus, and insula (Supplementary Table S1 and Fig. 3A). Post hoc t-tests revealed that these were all due to stronger activation during the anticipation of reward compared to no-reward trials, replicating earlier studies on reward anticipation in healthy study participants (Oldham *et al.*, 2018; Wilson *et al.*, 2018; Martins *et al.*, 2020). Additionally, main effects of *task type* were found in the fusiform and occipital gyrus, as well as the frontal medial and superior frontal cortex (Supplementary Table S1, Fig. 3B), which reflected stronger activations during social compared to monetary reward conditions (across reward and no reward trials). Finally, there was a main effect of *group* in the supracalcarine cortex/cuneus and occipital pole (see supplementary Table S1, Fig. 3C), reflecting larger activation in patients with AUD than in healthy controls.

Analyses of mean parameter estimates extracted from the left and right ventral striatum as regions of interest corroborated the whole-brain findings: No significant interaction effects of *group* were detected, suggesting that the neural processing of monetary and social reward anticipation in the ventral striatum is not altered in AUD. Bayesian analyses supported these findings, providing anecdotal to strong evidence against the inclusion of main effects of group and task as well as interaction effects of group, with Bayes factors (BF_{10}) ranging from 0.08 to 0.65 (see Table S2 in the Supplement for details). Additionally, a main effect of *reward* (left ventral striatum: $F(1,67) = 22.37$, $p < 0.001$; right ventral striatum: $F(1,67) = 22.42$, $p < 0.001$; see Fig. 4) but no significant main effects of *task type* or *group* could be detected (p > 0.12). A significant *task type* × *reward* interaction in the bilateral ventral striatum (left: $F(1,67) = 11.71$, $p = 0.001$; right: $F(1,67) = 10.25$, $p = 0.002$) indicated a greater differentiation between reward and no reward trials in the monetary task.

Discussion

The present study examined the anticipation of social rewards in individuals with AUD compared with healthy control participants. Neither behavioral nor neuroimaging data provided evidence of an altered processing of social reward in AUD, and similar results were observed for monetary rewards as well. These findings suggest that sensitivity to cues for social and other alcohol-unrelated rewards is not fundamentally impaired in AUD patients.

The present data showed that across both groups, response times were shorter in reward than in no reward trials. Likewise, increased neural activation during reward anticipation was observed in the striatum, occipital gyrus, supplementary motor cortex, fusiform gyrus, and insula across both groups. This finding is consistent with previous research conducted solely on healthy participants (Oldham *et al.*, 2018; Wilson *et al.*, 2018; Martins *et al.*, 2020). Additionally, we found that both AUD and healthy controls reacted faster (in reward and no reward trials) during the MID compared to the SID task, and had stronger neural activation during the SID in the fusiform gyrus, lingual gyrus, lateral occipital cortex, medial frontal cortex, and superior and middle frontal gyrus. While the results for response times are in line with previous studies in healthy subjects, the findings on neural activity show a heterogeneous picture with some studies reporting stronger activations in several regions in the monetary (Kirsch *et al.*, 2003; Dutra *et al.*, 2015) and others in the social domain (Gossen *et al.*, 2014).

The present study mainly focused on the anticipation of social rewards, which, to the best of our knowledge, has not yet been investigated in AUD. We did not find any neurobehavioral evidence of altered motivation for social rewards in individuals with AUD. This result adds to recent findings showing that reward learning abilities of patients with AUD appear comparable to those of healthy controls (Jangard *et al.*, 2025). These results may suggest that altered social behavior in AUD is not attributable to impaired reward processing per se (Jangard *et al.*, 2025). Analogous to social rewards, we also did not find evidence of reduced motivation for monetary reward in AUD.

Consistent with previous studies using the MID task, response times did not differ between patients with AUD and healthy controls (Beck *et al.*, 2009; Bjork *et al.*, 2012; Grodin *et al.*, 2016; Musial *et al.*, 2023). In line with the response times, we could not detect any evidence of altered neural processing. This finding stands in contrast to some earlier studies, including a meta-analysis of twelve studies (Zeng *et al.*, 2023), which reported decreased ventral striatal activation during monetary reward anticipation in AUD. However, it should be noted that studies with considerable modifications to the MID (e.g., effort variations in the trials or beer and water sips instead of monetary rewards) and studies with apparently overlapping datasets were also included in the meta-analysis. Without taking these studies into account, previous research presents a heterogeneous picture of reports showing decreased, increased, or comparable ventral striatal responses in AUD compared to healthy controls. Differences in task parameters (e.g., task duration, inclusion of loss conditions, reward magnitude) and participant characteristics (e.g., shorter abstinence duration) could plausibly account for the heterogeneity of the findings.

Beyond the primary region-of-interest analysis, exploratory whole-brain comparisons revealed increased activity in AUD relative to controls across all trials (reward and no reward) in the cuneus/occipital cortex - a region for which reduced gray matter volume has been shown in AUD (Wang *et al.*, 2018). The cuneus is implicated in shifting and sustaining attention and integrating visual information (Makino *et al.*, 2004; Michels *et al.*, 2008), and prior work has also linked this region to the differentiation of reward-predictive versus non-predictive cues, suggesting a role in assigning valence to potential rewards (Doñamayor *et al.*, 2012). Additionally, increased cuneus activation in response to monetary rewards has been observed in major depressive disorder, pointing to a possible involvement of this region in reward processing across clinical populations (Zhang *et al.*, 2013). Elevated cuneus activity may reflect altered visual attention in response to reward cues in AUD, although this interpretation remains speculative and requires further investigation.

Several limitations of the present study should be acknowledged. First, the sample predominantly consisted of male participants. Sex differences have been found in the mesolimbic dopamine release caused by alcohol (Agabio *et al.*, 2017) and in the neural processing of rewards (Spreckelmeyer *et al.*, 2009; Warthen *et al.*, 2020). Therefore, the results of the present study may not be generalizable to female individuals. Second, the findings may be specific to early abstinence. Previous studies have shown that a recovery of structural, functional, and cognitive impairments can occur within the first week to month of abstinence (Parvaz *et al.*, 2022; Powell *et al.*, 2024). Consequently, the results may not generalise to other phases of addiction. Finally, one could argue that our study was not sufficiently powered to detect subtle group differences. While larger samples are certainly needed to explore even smaller differences between groups, it should be noted that the Bayesian analyses provided evidence supporting the absence of differential responses between groups. In addition, most previous studies reporting altered ventral striatal activity during monetary reward anticipation in AUD had smaller sample sizes (Zeng *et al.*, 2023), rendering the current sample relatively robust in comparison.

Taken together, our results indicate that the motivation for social rewards is not altered in AUD. As no group differences were found for monetary rewards either, they do not support the "hijacking" hypothesis, which states that enhanced incentive salience for alcohol-related cues in AUD comes at the expense of motivation for alcohol-unrelated rewards. Instead, our findings indicate that the fundamental processing of social and monetary reward anticipation remains intact, suggesting preserved alcohol-unrelated reward motivation in individuals with AUD. This has direct clinical relevance: Social factors play a central role in AUD treatment, and many individuals with AUD perceive alcohol as a substitute or facilitator of social reward. Accordingly, a key therapeutic goal is to help patients build alcohol-free sources of social reinforcement. Our findings suggest that this is a promising approach, given that the neural processing of other rewards — particularly social ones — remains comparable to that observed in healthy individuals, highlighting the potential of interventions that rely on the ability to respond to social rewards. Therapeutic approaches such as

social skills training can enhance the ability to cope with social stressors and mobilize social support, thereby increasing the likelihood of abstinence and supporting relapse prevention.

Conflict of interest

The authors declare that they have no competing interests.

Acknowledgements

We thank Alexander Schröder, Julie Forster, and Jovana Lehmann-Grube for their help in data collection.

Funding

This work was supported by the Else Kröner-Fresenius-Stiftung (to LR, Grant No. 2018_A26). JV and YS were funded by the Studienstiftung des Deutschen Volkes.

Data Availability

Data and code of this study are openly available at: https://osf.io/g3ysp/

Contributor roles

Funding was acquired by LR. FMP, JV, LR, MC, and SK designed the study. EZ, JS, JV, KJ, OV, and YS recruited the participants and acquired the data. LR and YS analyzed data and drafted the manuscript. All authors critically reviewed the manuscript and approved the final version.

References

Addiction Cue-Reactivity Initiative (ACRI) Network, Sangchooli, A., Zare-Bidoky, M., et al. (2024). Parameter space and potential for biomarker development in 25 years of fMRI drug cue reactivity: A systematic review: A systematic review. *JAMA psychiatry (Chicago, Ill.)*, **81**, 414–25

Agabio, R., Pisanu, C., Gessa, G.L., et al. (2017). Sex differences in alcohol use disorder. *Current medicinal chemistry*, **24**, 2661–70

Beck, A., Schlagenhauf, F., Wüstenberg, T., et al. (2009). Ventral striatal activation during reward anticipation correlates with impulsivity in alcoholics. *Biological psychiatry*, **66**, 734–42

Berridge, K.C. (2012). From prediction error to incentive salience: mesolimbic computation of reward motivation: From prediction error to incentive salience. *The European journal of neuroscience*, **35**, 1124–43

Bjork, J.M., Smith, A.R., Chen, G., et al. (2012). Mesolimbic recruitment by nondrug rewards in detoxified alcoholics: effort anticipation, reward anticipation, and reward delivery. *Human brain mapping*, **33**, 2174–88

Bürkner, P.-C. (2017). Brms: An R package for Bayesian multilevel models using Stan. *Journal of statistical software*, **80**, 1–28

Cofresí, R.U., Bartholow, B.D., Piasecki, T.M. (2019). Evidence for incentive salience sensitization as a pathway to alcohol use disorder. *Neuroscience and biobehavioral reviews*, **107**, 897–926

Delgado, M.R., Fareri, D.S., Chang, L.J. (2023). Characterizing the mechanisms of social connection. *Neuron*, **111**, 3911–25

Diekhof, E.K., Kaps, L., Falkai, P., et al. (2012). The role of the human ventral striatum and the medial orbitofrontal cortex in the representation of reward magnitude - an activation likelihood estimation meta-analysis of neuroimaging studies of passive reward expectancy and outcome processing. *Neuropsychologia*, **50**, 1252–66

Doñamayor, N., Schoenfeld, M.A., Münte, T.F. (2012). Magneto- and electroencephalographic manifestations of reward anticipation and delivery. *NeuroImage*, **62**, 17–29

Dutra, S.J., Cunningham, W.A., Kober, H., et al. (2015). Elevated striatal reactivity across monetary and social rewards in bipolar I disorder. *Journal of abnormal psychology*, **124**, 890–904

Gossen, A., Groppe, S.E., Winkler, L., et al. (2014). Neural evidence for an association between social proficiency and sensitivity to social reward. *Social cognitive and affective neuroscience*, **9**, 661–70

Grodin, E.N., Steckler, L.E., Momenan, R. (2016). Altered striatal response during effort-based valuation and motivation in alcohol-dependent individuals. *Alcohol and alcoholism (Oxford, Oxfordshire)*, **51**, 638–46

Hägele, C., Schlagenhauf, F., Rapp, M., et al. (2015). Dimensional psychiatry: reward dysfunction and depressive mood across psychiatric disorders. *Psychopharmacology*, **232**, 331–41

Havassy, B.E., Hall, S.M., Wasserman, D.A. (1991). Social support and relapse: Commonalities among alcoholics, opiate users, and cigarette smokers. *Addictive behaviors*, **16**, 235–46

Heilig, M., Epstein, D.H., Nader, M.A., et al. (2016). Time to connect: bringing social context into addiction neuroscience. *Nature reviews. Neuroscience*, **17**, 592–99

Jangard, S., Lindström, B., Khemiri, L., et al. (2025). Dissociating social reward learning and behavior in alcohol use disorder. *Translational psychiatry*, **15**, 30

Kirsch, P., Schienle, A., Stark, R., et al. (2003). Anticipation of reward in a nonaversive differential conditioning paradigm and the brain reward system: an event-related fMRI study. *NeuroImage*, **20**, 1086–95

Knutson, B., Westdorp, A., Kaiser, E., et al. (2000). FMRI visualization of brain activity during a monetary incentive delay task. *NeuroImage*, **12**, 20–27

Makino, Y., Yokosawa, K., Takeda, Y., et al. (2004). Visual search and memory search engage extensive overlapping cerebral cortices: an fMRI study. *NeuroImage*, **23**, 525–33

Martins, D., Rademacher, L., Gabay, A.S., et al. (2020). *Mapping social reward and punishment processing in the human brain: A voxel-based meta-analysis of neuroimaging findings using the Social Incentive Delay task*. bioRxiv.

Michels, L., Moazami-Goudarzi, M., Jeanmonod, D., et al. (2008). EEG alpha distinguishes between cuneal and precuneal activation in working memory. *NeuroImage*, **40**, 1296–1310

Muench, C., Wiers, C.E., Cortes, C.R., et al. (2018). Dopamine transporter gene methylation is associated with nucleus accumbens activation during reward processing in healthy but not alcohol-dependent individuals. *Alcoholism, clinical and experimental research*, **42**, 21–31

Musial, M.P.M., Beck, A., Rosenthal, A., et al. (2023). Reward processing in alcohol-dependent patients and first-degree relatives: Functional brain activity during anticipation of monetary gains and losses. *Biological psychiatry*, **93**, 546–57

Oldham, S., Murawski, C., Fornito, A., et al. (2018). The anticipation and outcome phases of reward and loss processing: A neuroimaging meta-analysis of the monetary incentive delay task. *Human brain mapping*, **39**, 3398–3418

Parvaz, M.A., Rabin, R.A., Adams, F., et al. (2022). Structural and functional brain recovery in individuals with substance use disorders during abstinence: A review of longitudinal neuroimaging studies. *Drug and alcohol dependence*, **232**, 109319

Powell, A., Sumnall, H., Smith, J., et al. (2024). Recovery of neuropsychological function following abstinence from alcohol in adults diagnosed with an alcohol use disorder: Systematic review of longitudinal studies. *PloS one*, **19**, e0296043

Robinson, T.E., Berridge, K.C. (2025). The incentive-sensitization theory of addiction 30 years on. *Annual review of psychology*, **76**, 29–58

Robinson, T.E., Berridge, K.C. (1993). The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain research. Brain research reviews*, **18**, 247–91

Romanczuk-Seiferth, N., Koehler, S., Dreesen, C., et al. (2015). Pathological gambling and alcohol dependence: neural disturbances in reward and loss avoidance processing: Reward processing in gambling. *Addiction biology*, **20**, 557–69

Schwarze, Y., Voges, J., Schröder, A., et al. (2024). Altered physiological, affective, and functional

connectivity responses to acute stress in patients with alcohol use disorder. *Biological psychiatry global open science*, **4**, 100358

Spreckelmeyer, K.N., Krach, S., Kohls, G., et al. (2009). Anticipation of monetary and social reward differently activates mesolimbic brain structures in men and women. *Social cognitive and affective neuroscience*, **4**, 158–65

Stuber, G.D. (2023). Neurocircuits for motivation. *Science (New York, N.Y.)*, **382**, 394–98

Tan, B., Browne, C.J., Nöbauer, T., et al. (2024). Drugs of abuse hijack a mesolimbic pathway that processes homeostatic need. *Science (New York, N.Y.)*, **384**, eadk6742

Wang, J., Fan, Y., Dong, Y., et al. (2018). Combining gray matter volume in the cuneus and the cuneus-prefrontal connectivity may predict early relapse in abstinent alcohol-dependent patients. *PloS one*, **13**, e0196860

Warthen, K.G., Boyse-Peacor, A., Jones, K.G., et al. (2020). Sex differences in the human reward system: convergent behavioral, autonomic and neural evidence. *Social cognitive and affective neuroscience*, **15**, 789–801

Wilson, R.P., Colizzi, M., Bossong, M.G., et al. (2018). The neural substrate of reward anticipation in health: A meta-analysis of fMRI findings in the monetary incentive delay task. *Neuropsychology review*, **28**, 496–506

Wrase, J., Schlagenhauf, F., Kienast, T., et al. (2007). Dysfunction of reward processing correlates with alcohol craving in detoxified alcoholics. *NeuroImage*, **35**, 787–94

Young, K.A., Gobrogge, K.L., Wang, Z. (2011). The role of mesocorticolimbic dopamine in regulating interactions between drugs of abuse and social behavior. *Neuroscience and biobehavioral reviews*, **35**, 498–515

Zeng, J., You, L., Yang, F., et al. (2023). A meta-analysis of the neural substrates of monetary reward anticipation and outcome in alcohol use disorder. *Human brain mapping*, **44**, 2841–61

Zhang, W.-N., Chang, S.-H., Guo, L.-Y., et al. (2013). The neural correlates of reward-related processing in major depressive disorder: a meta-analysis of functional magnetic resonance imaging studies. *Journal of affective disorders*, **151**, 531–39

Tables

Table 1.

Sample characteristics

	AUD (N = 36)[a]	HC (N = 34)
Gender	6% f, 94% m	6% f, 94% m
Age (years)	42.4 ± 8.9	44.0 ± 9.3
Duration of abstinence (days)	14.8 ± 6.5	-
Amount alcohol (g/day)	291.4 ± 201.7	2.8 ± 3.5
Mean response times (ms):		
Reward SID	348.8 ± 36.4	344.3 ± 49.9
Reward MID	348.6 ± 36.5	342.5 ± 48.6
No reward SID	362.5 ± 40.4	361.4 ± 46.2
No reward MID	357.2 ± 34.8	354.2 ± 50.7

Values are given as mean and standard deviation unless specified otherwise. [a]For one AUD subject, exact information about the duration of abstinence (between 10 and 40 days)and for two AUD subjects, the exact amount of alcohol before withdrawal is missing, resulting in a sample size of N=35 and N=34 for these two variables, respectively.

Figure legends

Figure 1. Mean response times of patients with AUD and healthy controls (HC) for reward and no reward trials of the monetary and social incentive delay task.

Figure 2. Neural activation during the anticipation phase of the Social and Monetary Incentive Delay Task. (A) Effect of reward (reward vs. no reward trials). (B) Effect of task type (monetary vs. social task). (C) Effect of group (AUD vs. HC). A false discovery rate-corrected cluster-extent threshold of p < 0.05 based on p < .001 voxel-level threshold was used in all whole-brain analyses, except for the main effect of reward. Given the expected large effects of reward, a more stringent family-wise error-corrected threshold of p < 0.05 was applied.

Figure 3. Ventral striatum response to reward cues. Average parameter estimates were extracted from the left and right ventral striatum ROI for each participant in the two groups for reward and no reward trials of each task type. AUD = Alcohol use disorder group; HC = Healthy control group.

A) Reward > no reward

Occipital cortex

Supplementary motor cortex

Fusiform gyrus/ Cerebellum

Striatum

Insula

B) Social > monetary

Superior/middle frontal gyrus

Occipital cortex/ Cuneal cortex

Frontal medial cortex

Fusiform gyrus

C) AUD > HC

Supracalcarine cortex/ Cuneal cortex