Integrating satellite data and field surveys to predict

bird communities in the Peruvian Amazon

Andrew Christopher Slater[1]*, Chris A. Kirkby[2], Chris Ketola[2], Ian Hartley[1], Alex Bush[1]

[1]Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ

[2] Fauna Forever, Puerto Maldonado, Peru

* Corresponding author

E-mail: acs_africa@yahoo.co.uk (ACS)

Word count: Abstract = 254, Main text = 5883, References = 3753, Tables and figure legends

= 423

Number of: Tables = 2, Figures = 6

1 Abstract

Tropical forests harbour some of the highest biodiversity on Earth but are undergoing rapid loss and degradation. In the Amazon more than one-third of forests have been altered through human activities, with major implications for wildlife communities. While Earth observation satellites effectively monitor forest cover at scale, it remains unclear how well satellite-derived variables capture variation in bird communities. We tested whether Landsat reflectance and vegetation indices can predict bird species occurrence and community composition in the Peruvian Amazon. We analysed 3,129 point counts and mist-net bird surveys conducted over 16 years in the Tambopata Forest, south-eastern Peru. As predictors we compared the effectiveness of remote sensing derived surface reflectance and vegetation indices (e.g. NDVI and tasselled cap), with traditional land-type and forest cover descriptors. Species occurrence probabilities and community composition of 135 frequently recorded bird species were estimated using multi-species occupancy models that account for imperfect detection. Models using Landsat reflectance and vegetation indices outperformed those based on habitat categories in predicting species occupancy (mean AUC = 0.68 vs 0.58). They also achieved high predictive accuracy (AUC > 0.7) for more species (49 compared with 20). However, low detection rates across surveys limited all models' ability to accurately estimate full community composition and to detect change over time. Our results demonstrate that satellite-derived variables can improve predictions of bird occurrence compared with habitat categories, but their effectiveness depends strongly on survey design and species detectability. Integrating remote sensing with well-structured field surveys provides a scalable approach to monitoring biodiversity trends in tropical forests.

2 Introduction

Tropical forests are the most biodiverse areas on Earth (1,2), and support over a billion

people with food, materials, and services (3). They also play a key role in the global carbon

cycle, accounting for half of terrestrial uptake and two-thirds of forest carbon sinks in

biomass, soil, and deadwood (4,5). However, clear-cut logging and commercial agriculture

are altering these habitats, with over 17% of the Amazon Basin deforested and 38% of the

remainder degraded (6–8). Conservation strategies, such as the Reducing Emissions from

Deforestation and Degradation (REDD+) initiative, have generally reduced deforestation,

enhanced carbon stocks, and improved sustainable forest management (9–11). However,

half of the world's tropical forest reserves continue to lose taxonomic and functional

biodiversity due to disturbances beyond forest loss (12,13). While forest cover change can

be mapped in near real-time (14), assessing the efficacy of conservation strategies requires

evaluating the current condition of biodiversity within the ecosystem and where and how it

changes over time (15–17).

Ecosystem structure is often described by categorical elements such as intact, logged,

secondary, or plantation forest, yet these definitions can be ambiguous, subjective, and may

not reflect the most effective descriptors of an ecosystem's condition (18–20). In this

regard, essential biodiversity variables, including species distribution, abundance, and

community composition, have been proposed as additional criteria to monitor (21,22).

Satellite remote sensing (RS) cannot directly see these variables, but it can provide broader,

scalable environmental descriptors, alternative to field surveys, that act as proxies for

essential biodiversity variables, such as ecological diversity, phenology and physiology (23–

26). Continuous RS variables have been shown to outperform discrete land classes at describing plant species richness and diversity (27). Similar approaches have been applied to birds, where RS indices predict species richness and community composition (28,29), the structure of which has been used to assess the effectiveness of reforestation (30), and to show that they are negatively affected by patch size reduction within the Brazilian Amazon (31). RS surface reflectance values can therefore be used to infer vegetation and habitat structure as continuous gradients, avoiding the limitations of pre-defined classes. Yet, while RS variables can infer potential structure and ecological richness, there is little understanding of how they relate to the organisation of bird communities within tropical forests (32–34).

Biodiversity surveys provide essential information but are time consuming, expensive, and often limited in their spatial and temporal coverage (35,36). Linking survey data with RS descriptors in statistical models offers a way to estimate community and ecosystem structure across a landscape with limited expense (37). Because Amazonian bird communities are shaped by the characteristics of the forest in which they occur, we expect RS surface reflectance values to capture elements of these patterns. The continually updated nature of RS data also presents opportunities for this method to be used in monitoring ongoing changes in community structure. In this study, we test whether RS reflectance can better predict bird species occupancy than traditional habitat descriptors, and whether these models can predict observed community composition. We further hypothesise that survey replications and species detection rates will constrain model performance and the ability to detect temporal change. To evaluate these questions, we compare model predictive accuracy using species-level (AUC) and community-level (Bray-Curtis dissimilarity) metrics, on withheld data. We also assess how survey effort influences

90 the power to monitor biodiversity trends. By bringing these elements together, we aim to

91 provide a practical tool for policy makers and ecologists to assess the successes and failures

92 of management strategies, and to support decisions about financial incentives.

94 3   Methods

95 3.1   Study area and surveys

Survey data were collected by FaunaForever (www.faunaforever.org) to monitor species

diversity across landscapes in the Madre de Dios region, near the Tambopata River and

Puerto Maldonado in south-eastern Peru (**Error! Reference source not found.**). All collection

and subsequent use of these data complied with FaunaForever's data use policies.

Researchers and volunteers conducted bird surveys at 637 stations between 2004 and

2020. Bird surveys were conducted under permits issued by INRENA (Instituto Nacional de

Recursos Naturales) and later SERNANP (Servicio Nacional de Áreas Naturales Protegidas

por el Estado) for sites within the Tambopata National Reserve (RNTAMB; permits 20 S/C-

2003-INRENA-IANP,   004-2010-SERNANP-RNTAMB-JEF,   014-2011-SERNANP-RNTAMB-JEF,

012-2012-SERNANP-RNTAMB-JEF).   Surveys   outside   national   protected   areas   were

undertaken with the permission of landowners or concession title holders in the context of

their approved management plans (DEMA). The stations were clustered around 25 centres

that were located between 1.5km and 170km apart. Within centres, stations were between

200m and several kilometres apart and surveyed one to 35 times (median = 4), totalling

3,129 surveys. Stations surveyed in multiple years were treated as independent, since

populations and forest structure can change over extended periods of time. This resulted in

a total of 1,135 independent stations, which were surveyed between one and 21 times, with

a mode of 1 survey, a median of 2 surveys and a mean of 2.9 surveys per site (SD ±2.4).

Surveys used point counts (10-minute visual/audio records), or net counts (36m mist-net

captures). Simulation studies have shown that models built using species with fewer than 10

observations are prone to unreliable predictions, but the precise minimum required varies

with species' range size and study context (38). Based on this evidence, we applied a

pragmatic threshold, not intended as a universal standard, that balanced the inclusion of as

many species as possible with the need for reliable model estimates. We therefore retained

only species observed at more than 10 stations. Of the 358 species identified, this resulted

in 138 species being used for model training.

● Survey Sites

Fig 1: **(a)** The location of survey stations within the Madre de Dios region, south-eastern Peru. The inset map shows Peru with Madre de Dios highlighted in blue. The grey rectangle bounded in red indicates the area shown in panel **(b)**. Panel **(b)** illustrative zoom of the highlighted area showing local dispersion of survey stations. The satellite backdrop in **(b)** is provided for context only and was

143 not used in analysis. Basemap data: Natural Earth (public domain). Panel **(b)** backdrop: Sentinel-2

144 L2A imagery (Copernicus, cc BY 4.0)(109).

3.2 Environmental variables

Habitat was classified by land class and forest degradation. Land classes (floodplain forest,

terra firma forest, and agricultural) were based on the MapBiomas Peru Project, Collection-**1**

of the Annual Land Cover and Land Use Series for Peru 2013; (39). MapBiomas

classifications are based on Landsat imagery at 30m resolution using a random forest

approach to produce annual land cover and land use maps across Peru. We selected the

2013 dataset as it represents the midpoint of our 2004-2020 survey period and provides a

consistent baseline for all survey sites. While this inevitably simplifies temporal changes in

land cover, our survey stations were largely in stable forested areas where major shifts were

unlikely.

Forest degradation was defined using the Hansen et al. Global Forest Change dataset, which

reports annual forest loss from 2000 onwards (14). Pixels with no reported loss were

classified as primary forest, while those with any recorded loss were classified as secondary

forest.. This binary classification is a coarse simplification of the successional gradient but

was necessary to align with categories recorded in some field surveys, and to enable

comparability across datasets. The proportion of non-primary forest, equating to forest

assumed to have been deforested or degraded, was calculated within three different radii

around each station (1km, 2km and 5km). No cross validation was performed between the

MapBiomas and Hansen datasets, so forest classifications may not align. These limitations

highlight the constraints of categorical land-cover datasets for ecological modelling, and

partly motivate our parallel use of continuous remotely sensed reflectance variables to

better capture environmental gradients.

Of the 1135 surveys, 489 were on terra firma, 603 on floodplain and 43 on agricultural land,

and 1109 were defined as being inside primary forest, and 26 in secondary forest. Elevation

can influence the distribution of tropical forest bird species (40,41). However, all survey

points were within the narrow range between 175m and 310m elevation, and unlikely to

affect model performance; it was therefore not included in the analysis.

3.3   Satellite derived variables

Reflectance signatures at different wavelengths can be employed to assess the greenness

and moisture content of vegetation, with some correlation observed between these

signatures and the leaf structure of a plant (42). Reflectance and water content descriptors

help describe forest canopy physiology (43). No single metric captures canopy spectral

diversity (44), and the spectral characteristics and spatial scales that best capture canopy

variation are unknown, so various indices were used to describe it. While many satellite

platforms now provide passive optical surface reflectance data, as well as active radar

reflectance, the Landsat missions were the only platforms that covered the entire period of

the field surveys (Landsat-5 for 2004-2011, Landsat-7 for 2012, and Landsat-8 for 2013-

2020). For all missions we used USGS Collection-1 surface reflectance products, which

provide consistent atmospherically corrected data across sensors. Frequent cloud cover

over tropical forests can greatly reduce the number of available images that have sufficient

clarity for analysis (45). To address poor visibility, annual cloud-masked, median pixel value

images were created in Google Earth Engine. Annual composites necessarily average across

all available cloud-free imagery within each year and therefore do not represent the specific

timing of individual bird surveys. While this approach may mask seasonal changes such as

leaf flushing and fruiting cycles, or logging and burning, phenological variation in tropical

forests is generally lower than in temperate ecosystems (26,46,47). Each image contained

six reflectance bands, which were used to calculate seven vegetation indices: the

normalised difference vegetation index (NDVI), enhanced vegetation index (EVI), normalised

burn ratio (NBR), normalised difference water index (NDWI), and the trio of tasselled cap

wetness, greenness, and brightness (TCW, TCG, TCB). Both NDVI and EVI were used to

detect vegetation cover and vigour. NDVI was included as one of the most commonly used

indices, but it is prone to saturation in dense forests such as the Amazon, whereas EVI is

better able to penetrate the canopy and capture sub-layers. We used NBR to highlight

freshly opened soil, and NDWI to highlight water content within the canopy. Tasselled cap

indices provide an alternative way of describing reflectance and structure, and have been

shown to correlate with, but show more detail than, NDVI (48).

For each annual survey, the mean and standard deviation of pixels in each of the reflectance

bands and indices were calculated within 100m of each station. This buffer was chosen

pragmatically to approximate the detection radius of point counts and encompass a 3x3

pixel window, reducing the influence of anomalous pixel values compared to a single-pixel

extraction. These 26 variables provide a high-dimensional description of local vegetation

reflectance. To reduce redundancy, we used principal component analysis to condense the

26 variables, and the first five components, explaining 96% variation in the data, were used.

PC1 (50% of variance) and PC2 (26% of variance) were not dominated by any single variable

or group. The maximum contribution of any variable was 7% for PC1 and 11% for PC2, with

most variables contributing more modestly. Variables with near zero loading in PC1 tended

to contribute more to PC2 and vice versa. Thus, the components represent broad

combinations of vegetation indices and reflectance bands. The distribution of the five

principal components and the proportion of deforestation at three scales are plotted in

Error! Reference source not found., and the relationship between the first two principal

components with environmental descriptors is shown in **Error! Reference source not**

found..

Fig 2: Pairwise relationships among predictor variables used in EO models. Lower panels are fitted

with linear trend lines (red). Diagonal panels show variable names and their density distribution

(blue). Upper panels show Pearson correlation coefficients.

Fig 3: The first two principal components, representing 76% of the variation in the RS data, plotted against each other and showing their relationship with the measured habitat types. Each point represents an individual survey and is coloured based on the combination of habitat (floodplain forest, terra-firma forest, agriculture) and forest type (primary or secondary). Ellipses represent the 95% confidence interval around the mean for each categorical group.

3.4 Analysis

To quantify species detection and assess whether bird community composition can be predicted as a function of habitat or RS descriptors, we used spatial factor multi-species occupancy models. Observation depends on true presence (1 = present, 0 = absent; assumed constant over the year of surveillance), and the probability of detection. Detection can be influenced by survey method. Mist nets at ground level tend to under-sample canopy dwelling, large or less mobile species (49). Capture rates also vary with the type of net used (e.g. mesh size), and netting duration. Point counts detect vocal species well, but may under-sample silent or secretive species, and their effectiveness depends on background noise, weather, and observer skill (50,51). Perfect detection of all species is essentially impossible. Non-detection can bias covariate estimates and predictions of community composition (52). However, occupancy models account for imperfect detection by estimating species occurrence at each survey station, whether observed or not (53). When adequately specified, occupancy models can estimate detection probabilities per species and station, incorporating the influence of survey-specific covariates, such as survey method or time of day (54). After fitting the model and estimating species detection and occurrence probabilities, the methods of Guillera-Arroita and Lahoz-Monfort (2012), were used to calculate the power of the study. The term "power" is the probability of detecting a proportional reduction in occupancy between surveys conducted at different times. We then calculated the survey effort required to achieve a mean power across all species of 0.7, to detect a 50% occupancy reduction. Although 0.8 is commonly used, our low detection rates made 0.7 a more feasible target that falls within the test ranges (55). We also assessed how detection power of various occupancy reductions varied with the number of stations and replicate surveys, to guide survey regimes for future monitoring.

Four models were compared: one fitted with habitat variables, one with principal

components from RS reflectance, one with no occurrence variables to act as an intercept-

only baseline, and one RS model re-fitted without spatial factors to assess their influence.

All models included survey method (point or net) and time of day (a.m. or p.m.) as

covariates for the probability of detection. A summary of all detection and occupancy

covariates, and the models in which they were included, is provided in Table 1. Although

some stations were surveyed up to 21 times, 99% of all stations were surveyed 12 times or

less. To improve the efficiency of model fitting and reduce the computational resources

required to run them, we fitted the modes using a maximum of 12 replicates per station.

Table 1: Summary of covariates considered in the multi-species occupancy models.

Covariate	Type	Description	Source / Processing	Models used in
Land class	Categorical	Floodplain, terra firma, agricultural	MapBiomas Peru Project, Collection-1 (2013) (39)	Habitat model
Forest type	Categorical	Primary vs. secondary forest	Global Forest Change (2000–2019) (14)	Habitat model
Deforestation (%)	Continuous	Proportion of non-primary forest within 1, 2, and 5 km radii around stations	Global Forest Change (2000–2019) (14)	Habitat model
Reflectance indices	Continuous	NDVI, EVI, NBR, NDWI, TCB, TCG, TCW; means and SDs within 100 m	Landsat 5/7/8 annual median composites, Google Earth Engine	RS model → PCA
PCA components (PC1–PC5)	Continuous	First five PCs of 26 reflectance variables (96% variance explained)	Derived from Landsat indices above	RS model
Survey method	Categorical	Point count vs. mist-net	Field survey protocol (Fauna Forever)	All models (detection)
Time of day	Categorical	Morning vs. afternoon	Field survey protocol (Fauna Forever)	All models (detection)

Detection covariates were included in all models. Occupancy covariates varied by model: habitat

descriptors (land class, forest type, deforestation), RS reflectance (PCA components), or none

(intercept-only).

Detection and occupancy covariates were modelled without interactions. Model outputs are

on the logit scale, with baseline categories set alphabetically (a.m., net, agricultural,

primary). Posterior estimates for other categories (e.g., p.m., point counts, secondary

forest) were obtained by adding the relevant coefficient to the baseline and applying the

inverse-logit transformation. We then summarised these posterior values to report the

median and 95% credible intervals (2.5% and 97.5% quantiles). Reported detection and

occupancy values in the Results are therefore directly derived from the posterior

distributions.

Models were fitted using the Bayesian occupancy model package, "spOccupancy" (56). The

"spOccupancy" package enables the modelling of all species simultaneously, and accounts

for autocorrelation between species. We included three spatial latent factors to account for

residual spatial autocorrelation in occupancy probabilities. The choice of three factors was a

pragmatic one, guided by recommendations in the spOccupancy framework that a small

number of factors (typically 2-5) provides sufficient flexibility to capture spatial structure

while avoiding overfitting and excessive computational cost. Each factor provides a decay

rate (phi), with the distance beyond which autocorrelation is negligible calculated as 3/phi.

Models were fitted with 50,000 iterations, discarding the first 25,000 as burn-in, and

thinning the rest by 50 to produce 500 posterior samples. Each posterior sample provides a

probability of occurrence (psi) for each species at each station and a latent

presence/absence (z = 1/0) calculated from this probability. Additionally, detection

probability (p), given presence, is provided per species per station for a single survey. As

spatial models often show poor multi-chain convergence, and given memory constraints, we

chose to run a single-chain model. Convergence was assessed visually, with Geweke

diagnostic (75% of parameters had z<1.96) and effective sample sizes (median ~ 274 across

all parameters), following spOccupancy guidance (57), all of which indicated adequate

convergence. Models assumed no false presences and treated the entire study as a single

closed season. Model goodness-of-fit was measured using posterior predictive checks to

calculate a Bayesian p-value, based on the differences between chi-squared values of

observed and predicted data per species, grouped across stations,(58,59) Models were also

compared using the widely applicable information criterion (WAIC) (60) and metrics of

predictive performance. Performance was evaluated at the species-level, using area under

the receiver operating curve (AUC), and at the station community-level, using species

accumulation and Bray-Curtis's dissimilarity of community composition.

Although model performance differs between training and withheld data, evaluation on

independent datasets is rarely undertaken for occupancy models (61). We did so here by

withholding a validation dataset comprising two stations per field centre (24 centres in

total), each surveyed at least three times, totalling 5% of the data. This ensured

representation across all centres while retaining sufficient training data even in the smallest

centres (minimum 4 stations). Validation stations were randomly chosen from those with at

least 3 survey replicates, so their observed communities were well characterised for

comparison with model predictions. To validate model performance, we incorporated both

occurrence and detection probabilities when comparing predictions with observations. Each

of the model's posterior samples predicts the occupancy probability (psi) per species per

site, and the latent occupancy (z = 1/0) representing presence or absence. Detection

probability (p) per species and site in a single survey was predicted, and this remained

constant across posterior samples. Total detection probability per species and station was

calculated as $1 - (1 - p)^n$, where n is the number of replicate surveys performed. This

detection probability generated a binomial detection/non-detection (1/0) event per species

per station. Latent detection values were then multiplied by occurrence (psi) and latent

occupancy (z) for each species and station per posterior sample. Thus, model performance

was based on the predicted probability of detection rather than occurrence in the validation

data.

4   Results

4.1   Model fit

Posterior predictive checks gave all models Bayesian p-values between 0.467-0.477,

indicating a good fit to the data. The model with the lowest WAIC value, thus indicating it

was the best-performing model, was fitted with RS reflectance data and included spatial

factors, followed by the covariate-free intercept model, the RS reflectance model fitted

without spatial factors, and the highest WAIC indicating the worst performance was from

the model fitted with environmental descriptors.

The probability of detection (p) was consistent across all models. Surveys conducted in the

morning using net counts had a posterior median species-level $p = 0.022$, 95% CI [0.001,

0.179]. When conducting surveys in the afternoon, median detection probability

significantly lower at $p = 0.015$, 95% CI [0.001, 0.018]. When using point rather than net

surveys, the median p increased significantly to 0.065, 95% CI [0.000, 0.389]. These results

indicate that detectability depends strongly on method and time of day, with point counts

and morning surveys yielding higher overall detection rates. However, point surveys did not

significantly increase detection for three of the nine Orders represented. Those three were

Caprimulgiformes (nightjars), and Coraciiformes (kingfishers and hornbills), and Galliformes

(ground dwelling birds).

The posterior mean probability of occurrence across all species in primary forest and an

agricultural habitat, was 0.291, 95% CI [0.211, 0.375]. In secondary forest, the mean

occupancy was higher than in primary forest, rising to 0.47, 95% CI [0.36, 0.61]. In

comparison to agricultural landscapes, mean occupancy in floodplain increased to 0.36,

95% CI [0.30, 0.47], and in terra-firma mean occupancy increased to 0.37, 95% CI [0.30,

0.47]. This indicates that secondary forests supported more species on average than primary

forests, while agricultural habitats supported fewer. Mean occupancy was not shown to vary

significantly with deforestation within a 1-2 km radius, but increased by 0.057, 95% CI

[0.008, 0.108] with each 10% increase in deforestation within a 5km radius. This suggests

that bird occurrence responded more strongly to landscape-scale forest change (5km) than

local variation (1-2km).

When using RS data, the model suggested that spatial autocorrelation extended to 5km,

95% CI [3.3km, 7.5km] for one factor and 7.5km, 95% CI [4.3km, 15km] for each of the other

two factors. When using habitat data, the model suggested that spatial autocorrelation

extended to 2km, 95% CI [2km, 2.3km] for one factor, to 30km, 95% CI [3.75km, ∞] for

another, and the third factor was not significant. These factors represent unmeasured

variables that influence the occurrence of species, and fade with distance. Although we do

not know what is driving this spatial structure, it can be influenced by anything from local

vegetation cover to regional climate or land use. Significant spatial autocorrelation confirm

that patterns exist beyond those explained by the known environmental variables.

4.2   Model validation

The predictive performances of models followed the implications made by their WAIC values, and the model that produced the best predictive results was also fitted with RS reflectance data and included spatial factors. When assessed on training data, this model had a mean AUC of 0.88 (SD=0.06) and when predicting to independent validation sites, had a mean AUC of 0.68 (SD=0.18; range=0.16-0.99). Species for which predictive performance was considered good (AUC>0.7) (62) are hereafter referred to as high-AUC species, and this model had 49 high-AUC species, the highest of any model. The performance of all four models is shown in Table 2**Error! Reference source not found.**. Species taxonomic order was not shown to have any correlation with its AUC.

The predicted species richness of survey stations (6.86 ± 5.1) was close to the observed average richness (6.00 ± 3.6), but when extended to the entire training dataset the model predictions suggested that it would take much longer to detect the total number of species included (**Error! Reference source not found.**(A)). The mean predicted richness of test stations (9.94 ± 3.8) was also comparable to the observed mean richness (10.82 ± 5.3) and indicated a similar accumulation of species, both across stations (**Error! Reference source not found.**(B)), and with increasing replication within a station (**Error! Reference source not found.**(C)). The species accumulation curve in **Error! Reference source not found.**(C) come from one station that was surveyed 12 times and illustrates the extent to which observed accumulated richness greatly underestimates the total richness expected to be present at each location. A consequence of this was that only a fraction of the species that occur at a station were observed, and therefore the average Bray-Curtis dissimilarity between the list of birds observed and those predicted was 0.8 (± 0.1). In locations with higher observed richness, the surveyed total represented a greater proportion of the true expected total,

and observed species richness was strongly negatively associated with predicted-observed

community dissimilarity (R^2=0.68, p<0.001) (**Error! Reference source not found.**).

Fig 4: Observed (blue) and modelled (black) species accumulation (±2SD) with increasing numbers of

surveyed stations, **(a)** within the training data, **(b)** set aside for validation, and **(c)** based on an

increasing number of replicate surveys conducted at a single survey station.

Fig 5: Relationship between observed species richness and dissimilarity between predicted and

observed community composition (Bray-Curtis) across all sites. A fitted linear regression line (black)

with 95% confidence interval (dark grey band) is shown (R^2=0.68, p<0.001). Marginal histograms

show the distribution of each variable.

Table 2: The fit and predictive performance of the four models analysed.

Model Covariates	Spatial Factor Included	WAIC value	Delta WAIC	Mean Explanatory AUC	Mean Predictive AUC	Number of high-AUC species
RS Reflectance	yes	66773	0	0.88	0.68	49
RS Reflectance	no	68620	1847	0.87	0.62	31
Habitat Descriptors	yes	69432	2659	0.87	0.58	21
None	yes	66995	222	0.8	0.66	43

Low WAIC values and high AUC values represent better performances. It can be seen that the best

performing model in all categories is fitted with RS reflectance data and includes spatial factors.

4.3   Power to detect change

As a result of low detection probabilities and low mean replicate surveys, it was ascertained

that there is a mean probability across all species of 0.20 (±0.24 SD) of detecting a 50%

reduction in occupancy in a subsequent survey. There is a 70% or greater probability of

detecting a 50% reduction in only five of the 135 species modelled. Thus, if the desire is to

determine where occupancies are changing by surveying the same area in the future, the

current data cannot provide robust evidence.

illustrates how different numbers of survey stations and replicates per station affect the

median power to observe a reduction in bird occupancy. Each panel represents a different

proportional change that is sought to be detected, and each y-axis represents the

probability of detecting that change in at least half the bird species. The x-axis is the total

number of individual surveys that a study is able to undertake (i.e. one replicate at one

station is one survey). Each line represents a different number of sites that the total number

of surveys are divided among. For every probability of detecting the panel's proportional

change (y-axis), the farthest left line represents the fewest total surveys required, and the

most efficient number of sites to survey. For example, if our aims were to have a 70%

confidence (0.7 on the y-axis) that we could detect when the occupancy of at least half the

bird species had declined by 50% (panel C), the chart suggests that the fewest surveys

required would be ~10,000 and spread between 100 and 500 sites. Calculations suggest that

the requirements would be three replicate surveys at nearly 50,000 stations, or, if all

stations were surveyed 27 times, only 447 stations would be required.

Fig 6: The influence of survey effort on the probability of detecting a reduction in occupancy in future surveys. Survey effort was calculated as the number of stations surveyed * number of replicate surveys per station. Each line represents a different number of stations and was calculated as the mean value across all 135 species modelled. Each chart represents a different proportional reduction in occupancy.

5   Discussion

Using over 3,000 bird surveys conducted in south-eastern Peru, we found that remotely sensed satellite data performed slightly better than categorical landcover descriptors in predicting bird community composition. This advantage was modest and specific to our study region, and further testing would be required before generalising to other tropical systems. We also highlight the low detectability of many species, which complicates biodiversity monitoring in rainforests. Although the survey design provided valuable insights

into bird communities, it was not optimised to detect temporal change. Our results can

guide more efficient survey strategies, leading to enhanced species detection, better

capturing community composition, and strengthening the performance of RS models in

predicting bird communities across landscapes.

5.1   Land class and surface reflectance

Studies differ on how forest and farmland affect bird species richness (63,64). We found

more species in both floodplain and terra firma forests than in agricultural land, partially

aligning with evidence that species richness can increase at forest boundaries, particularly

where edges are sharp, such as where forest and agricultural land meet (65,66).

Furthermore, the greater stratification and canopy complexity often found in primary

forests typically support more specialist, range restricted, and endemic bird species

compared to secondary forests (67–69). However, our analysis suggested that secondary

forest had a greater species richness, consistent with findings that generalist and

disturbance tolerant species as well as species that thrive in more open canopy

environments are drawn to those areas, especially when they are connected to primary

forest, thereby increasing species richness (70). Greater visibility, vocalisation and bird

density in open secondary forests may also lead to more detections (71–73). The low mean

number of replicates in our study could make secondary forests appear richer in species

than primary forests, due to improved visibility rather than true richness. Secondary forests

in Amazonia typically occur as mosaics of regenerating lands, agricultural edges, and

disturbed forest patches. These mosaics can provide food resources and structural

heterogeneity that are especially favourable to generalist Amazonian bird species (74).

However, our data cannot disentangle these ecological drivers from detectability effects,

and both are likely contributing to the higher richness observed.

Secondary forests require the initial loss of primary forests, and localised deforestation can

increase bird species richness. Moderate primary deforestation can increase richness of

neighbouring forest as displaced individuals move into it (75). It can also increase habitat

heterogeneity, promoting colonisation by generalists and disturbance-tolerant species in

the deforested area (75). Our findings that deforestation within a 5km radius increased

mean occupancy support this. At this scale species dispersal, habitat connectivity, and edge

effects shape community composition (76,77). In contrast, no effects were found at smaller

distances, suggesting insufficient heterogeneity at that range. We defined deforestation as

loss of primary forest, but deforested areas may have regrown into secondary forest, which

also showed high species richness. Misclassification could arise from pre-2000 deforestation

or from the omission of water bodies from analysis. Within floodplain forests, water levels

can fluctuate seasonally (78) and edge effects may occur at interfaces between large water

bodies and forests (79). These classification limitations should be considered when

interpreting differences in species occupancy across forest types. bird detectability may

have varied due to seasonal fluctuations, which are most pronounced in forest fragments

and secondary forest (67,80,81). Surveys were performed by multiple observers with

varying levels of experience, which can significantly influence bird species detectability (82).

Consequently, some variation in detection probabilities may be attributed to differences in

surveyor rather than actual ecological differences.

Surface reflectance diversity has been used to predict tropical forest and bird diversity (83)

and bird species richness found within them (84) and our findings support this. RS data

performed as well as, if not better than, land class in predicting bird communities (85,86).

Image seasonality and resolution matter for modelling bird communities in temperate

regions (85,86). However, persistent cloud cover in our study area required annual

composites, and mean reflectance values calculated over thousands of square metres make

finer resolution imagery unlikely to improve results.

Despite these challenges, our study has shown that RS data have the capacity to improve on

land class in predicting bird communities in tropical forests. Two broad land-class groups

could be differentiated by plotting the first two PCAs: secondary forest/agriculture and

primary forest/terra-firma/floodplain. Within those two groups, secondary forest, defined

as land deforested since 2000, had a large overlap with agricultural land, suggesting that

deforested land here tends to become agricultural rather than regenerating into secondary

forest (87,88). In the other group, primary forest overlapped more with terra-firma and

floodplain than with agriculture, though sizeable overlaps between all groups existed. This

may be explained by discrepancies brought about by course scale, out of date information

or human error in habitat classification (89–91). Additionally, land classes may oversimplify

or fail to capture the finer scale variation seen by RS data. Localised secondary descriptors

of structural variation, such as deadwood volume and tree girth, can also influence

reflectance values, and when included in conjunction with broad habitat type, have been

shown to define species richness far better than habitat type alone (92). This shows that RS

data better capture gradual changes in vegetation structure than can predefined classes.

The challenge for future research will be in deriving RS variables that can describe more

ecologically meaningful variation and differentiate between less obvious variations in

surface structure.

5.2 Realities and practicalities of observing change

Whether using habitat or RS variables, model success depends on the quality and quantity of the data with which they are trained (93,94). Low detection probabilities suggest substantial effort is needed to detect most species, raising concerns about data sufficiency. However, of the approximately 760 bird species thought to occur in the Tambopata area (95), our study recorded nearly half. This, in terms of both number and proportion of species identified, aligns with findings from other surveys in tropical forests (96–98). Low detection rates likely caused individual surveys to underestimate community composition, making landscape-wide communities appear more distinct than limited survey observations suggest. Nevertheless, the species modelled exhibited patterns of grouping based on both forest habitat descriptors and RS variables, suggesting that they represented more than just random observations. This indicates that, despite describing approximately a half of the full community, our results were reliable.

Increased sampling effort revealed more of the local community and improved similarity between predicted and observed communities. Thus, improving detection rates is essential to increasing the proportion of the community observed and improving the predictive output of models. Increasing cumulative detection rates to 80% has been shown to negate the importance of detection on model performance (99). Yet we have shown that it would require a huge effort to reliably observe changes in occupancy for many species. Thus, efficient and accurate surveys are critical to provide high quality data on which to train RS models, especially as ongoing surveys are needed to monitor the effects of conservation efforts on habitat change over space or time. The type of survey conducted affects detection rate, and we agree with Mulvaney & Cherry, (2020) that point counts increase mean detection over net counts. However, these findings are not universal, as it has been

shown that within tropical forests, the most effective survey method varies with ecosystem,

such as whether the study is within lowland or cloud forest (97). Similarly, transect counts

and canopy counts add further depth to the proportions of small-bodied species, canopy

and mid-storey species that are detected (82,98). Due to the relatively low community

overlap observed between methods found in our study, and aligning with other studies, we

would suggest that the use of multiple methods is necessary to improve overall diversity

assessment (82,97,98). Further to the survey methodology lies the challenge of recruiting

appropriately trained surveyors who are familiar with the bird species of the area, which is a

necessity in conducting effective surveys in complex and rich ecosystems such as tropical

forests   (82). Our results suggest that to effectively detect changes in community

composition, it is necessary to perform far more replicate surveys than were performed

(82).

5.3   Future potential

Forest degradation in Peru has been shown to increase the amount of carbon released into

the atmosphere by almost 50% (101). Protecting forests and carbon sinks has

understandably been central of tropical conservation, and government policies and

voluntary carbon offset conservation efforts such as REDD+ have successfully slowed the

rate of deforestation in recent years and have played a role in mitigating climate change

(102). To ensure that the implementation of carbon conservation efforts, such as REDD+,

also have a positive impact on biodiversity, ecological monitoring of forests should be

aligned with the conservation objectives of the area (103). While REDD+ has had some

success in protecting forests globally, the level of success in the Amazon may be overstated

due to the overestimation of baseline deforestation (104), highlighting the need to validate

the costs of voluntary carbon offset credits. Degradation impacts are not fixed and can be

improved by restoration or worsened by repetition and should be measured at a landscape

rather than patch scale (105). Measuring community diversity at landscape scales may

require impractical levels of monitoring, but patch scale changes in habitat should in

principle be observable by Earth observation. Should sufficient, consistent, reliably

representative biodiversity data be available to calibrate models fitted with RS data,

biodiversity responses to changing land use can be identified. Policy and management

should judge the results of practices at the point of implementation rather than judge the

practices conceptually as a whole, and by mapping species diversity in conjunction with

other measures of landscape health, trends in forest degradation and regeneration over

time can be identified (105).

Despite the challenges posed by generally low detection rates, our study demonstrated that

the occupancy of a third of the bird species with sufficient detections in the Peruvian

Amazon can be predicted with reasonable confidence. However, low detection rates limit

model capacity to detect community change over time. While RS models may not yet be

sufficiently robust for ongoing monitoring, reporting and verification, there is no evidence

that field-based methods alone will be able to do this at the scale required. While surveys

can identify spatial variations, detecting temporal change requires detailed count data at a

level that may be impractical (106). Our power analysis revealed that with the current

monitoring structure, even if a species population were to half, we would fail to detect this

four out of five times. The suggestion that declines of many species may be substantial

before being reliably detected has major implications for conservation monitoring using

existing survey design. Improving detection rates will increase the power of future surveys

to detect change. Given that resources are limited, future surveys should focus on fewer

locations, prioritised across an ecological gradient of interest, with a higher number of

replicates; results suggest at least 20 per site. This is particularly critical as even basic

inventories of avian diversity across much of the tropics are lacking (82). Establishing

baseline datasets is essential for enabling future repeat surveys, which could provide

valuable insights into how and where tropical bird communities are changing (107). Building

on the findings of this study by refining survey design and aligning research with specific

conservation objectives will be key to improving biodiversity monitoring in the region so

that decisions are based on detectable trends rather than snapshots of biodiversity (108).

Incremental improvements in temporal monitoring are most likely to come from combining

these refinements in survey design with successive RS image series. By aligning more robust

field baselines with RS predictors, it will become possible to track biodiversity change

gradually across space and time, even in data-limited tropical landscapes.

Declarations

Funding:

This study received no specific funding. The bird data used were collected independently by

co-authors C.A.K. and C.K. as part of ongoing monitoring activities not funded by the lead or

senior authors of this manuscript.

Ethics Statement:

The bird occurrence data used in this study were collected independently by co-authors

C.A.K. and C.K. as part of long-term monitoring within the Tambopata National Reserve,

Peru. Data collection was conducted ethically and in accordance with local research

regulations.

Authors' Contributions:

Author contributions: A.C.S., A.B. and I.R.H. conceptualised the study; C.A.K and C.K.

performed independent field surveys and provided bird sighting data; A.C.S. sourced and

calculated all remotely sensed variables, performed all modelling and analysed results;

A.C.S. wrote the manuscript with reviews and edits suggested by A.B, I.R.H.

Conflict of Interest: The authors declare no conflicts of interest.

References

1.   Bradshaw CJ, Sodhi NS, Brook BW. Tropical turmoil: a biodiversity tragedy in

progress. Front Ecol Environ [Internet]. 2009 Mar [cited 2023 Apr 12];7(2):79–

87. Available from: http://doi.wiley.com/10.1890/070193

2.   Gibson L, Lee TM, Koh LP, Brook BW, Gardner TA, Barlow J, et al. Primary

forests are irreplaceable for sustaining tropical biodiversity. Nature [Internet].

2011 Oct 20;478(7369):378–81. Available from:

https://www.nature.com/articles/nature10425

3.   Lewis SL, Edwards DP, Galbraith D. Increasing human dominance of tropical

forests. Science (1979) [Internet]. 2015 Aug 21;349(6250):827–32. Available

from: https://www.science.org/doi/10.1126/science.aaa9932

4.   Pan Y, Birdsey RA, Fang J, Houghton R, Kauppi PE, Kurz WA, et al. A large and

persistent carbon sink in the world's forests. Science (1979). 2011 Aug

19;333(6045):988–93.

5.   Hubau W, Lewis SL, Phillips OL, Affum-Baffoe K, Beeckman H, Cuní-Sanchez A,

et al. Asynchronous carbon sink saturation in African and Amazonian tropical

forests. Nature. 2020 Mar 5;579(7797):80–7.

6.   Potapov P, Hansen MC, Laestadius L, Turubanova S, Yaroshenko A, Thies C, et

al. The last frontiers of wilderness: Tracking loss of intact forest landscapes

from 2000 to 2013. Sci Adv [Internet]. 2017 Jan 1 [cited 2023 Jun 13];3(1).

Available from: https://www.science.org/doi/10.1126/sciadv.1600821

7.   Vergara A, Arias M, Gachet B, Naranjo LG, Román L, Surkin J, et al. Living

Amazon Report 2022. 2022.

8.   Lapola DM, Pinho P, Barlow J, Aragão LEOC, Berenguer E, Carmenta R, et al.

The drivers and impacts of Amazon forest degradation. Science (1979)

[Internet]. 2023 Jan 27 [cited 2024 Mar 20];379(6630). Available from:

https://www.science.org/doi/10.1126/science.abp8622

9.   UNFCCC. United Nations Framework Convention on Climate Change. 2023

[cited 2023 Nov 8]. Home / REDD+. Available from: https://redd.unfccc.int/

10.   Wunder S, Schulz D, Montoya-Zumaeta JG, Börner J, Ponzoni Frey G, Betancur-

Corredor B. Modest forest and welfare gains from initiatives for reduced

emissions from deforestation and forest degradation. Commun Earth Environ

[Internet]. 2024 Jul 22 [cited 2025 Jan 27];5(1):394. Available from:

https://www.nature.com/articles/s43247-024-01541-1

11.   Guizar-Coutiño A, Jones JPG, Balmford A, Carmenta R, Coomes DA. A global

evaluation of the effectiveness of voluntary REDD+ projects at reducing

deforestation and degradation in the moist tropics. Conservation Biology

[Internet]. 2022 Dec 8 [cited 2025 Jan 27];36(6):e13970. Available from:

https://conbio.onlinelibrary.wiley.com/doi/10.1111/cobi.13970

12.   Barlow J, Lennox GD, Ferreira J, Berenguer E, Lees AC, Nally R Mac, et al.

Anthropogenic disturbance in tropical forests can double biodiversity loss from

deforestation. Nature [Internet]. 2016 Jul 7 [cited 2021 Jan 13];535(7610):144–

7. Available from: https://www.nature.com/articles/nature18326

13.   Laurance WF, Carolina Useche D, Rendeiro J, Kalka M, Bradshaw CJA, Sloan SP,

et al. Averting biodiversity collapse in tropical forest protected areas. Nature

[Internet]. 2012 Sep 25;489(7415):290–4. Available from:

http://www.nature.com/articles/nature11318

14.   Hansen MC, Potapov P V., Moore R, Hancher M, Turubanova SA, Tyukavina A,

et al. High-Resolution Global Maps of 21st-Century Forest Cover Change.

Science (1979) [Internet]. 2013 Nov 15 [cited 2023 Jan 23];342(6160):850–3.

Available from: https://www.science.org/doi/10.1126/science.1244693

15.   Stork NE, Boyle TJB, Dale V, Eeley H, Finegan B, Lawes M, et al. Criteria and

indicators for assessing the sustainability of forest management: conservation

of biodiversity [Internet]. Criteria and indicators for assessing the sustainability

of forest management: conservation of biodiversity. Center for International

Forestry Research (CIFOR); 1997 [cited 2025 Jan 27]. Available from:

http://www.cifor.org/library/77/criteria-and-indicators-for-assessing-the-

sustainability-of-forest-management-conservation-of-biodiversity/

16.   Titeux N, Henle K, Mihoub J, Regos A, Geijzendorffer IR, Cramer W, et al.

683   Biodiversity scenarios neglect future land-use changes. Glob Chang Biol

684   [Internet]. 2016 Jul 31;22(7):2505–15. Available from:

685   https://onlinelibrary.wiley.com/doi/10.1111/gcb.13272

686   17.   Willis KJ, Araújo MB, Bennett KD, Figueroa-Rangel B, Froyd CA, Myers N. How

687   can a knowledge of the past help to conserve the future? Biodiversity

688   conservation and the relevance of long-term ecological studies. Philosophical

689   Transactions of the Royal Society B: Biological Sciences [Internet]. 2007 Feb 28

690   [cited 2025 Jan 27];362(1478):175–87. Available from:

691   https://royalsocietypublishing.org/doi/10.1098/rstb.2006.1977

692   18.   Chazdon RL, Brancalion PHS, Laestadius L, Bennett-Curry A, Buckingham K,

693   Kumar C, et al. When is a forest a forest? Forest concepts and definitions in the

694   era of forest and landscape restoration. Ambio [Internet]. 2016 Sep 9 [cited

695   2025 Jan 27];45(5):538–50. Available from:

696   http://link.springer.com/10.1007/s13280-016-0772-y

697   19.   Savilaakso S, Lausberg N, Waeber PO, Hillgén O, Isotalo A, Kleinschroth F, et al.

698   Whose perspective counts? A critical look at definitions of terms used for

699   natural and near-natural forests. One Earth [Internet]. 2023 Nov [cited 2025

700   Jan 27];6(11):1477–93. Available from:

701   https://linkinghub.elsevier.com/retrieve/pii/S2590332223004530

702   20.   Pettorelli N, Schulte to Bühne H. Current and future opportunities for satellite

703   remote sensing to inform rewilding. Disney M, Hernandez-Clemente R, editors.

704   Remote Sens Ecol Conserv [Internet]. 2022 Dec 16; Available from:

705   https://zslpublications.onlinelibrary.wiley.com/doi/10.1002/rse2.321

706   21.   Pereira HM, Ferrier S, Walters M, Geller GN, Jongman RHG, Scholes RJ, et al.

707   Essential Biodiversity Variables. Science (1979) [Internet]. 2013 Jan

708   18;339(6117):277–8. Available from:

709   https://www.science.org/doi/10.1126/science.1229931

710   22.   GEO BON. What are EBVs. 2018 [cited 2022 Jul 22]. What are EBVs? – GEO

711   BON. Available from: https://geobon.org/ebvs/what-are-ebvs/

712   23.   Purdon A, Mole MA, Selier J, Kruger J, Mafumo H, Olivier PI. Using the Rao's Q

713   diversity index as an indicator of protected area effectiveness in conserving

714   biodiversity. Ecol Inform [Internet]. 2022 Dec [cited 2022 Dec 2];72:101920.

715   Available from:

716   https://linkinghub.elsevier.com/retrieve/pii/S1574954122003703

717   24.   Pettorelli N, Vik JO, Mysterud A, Gaillard JM, Tucker CJ, Stenseth NChr. Using

718   the satellite-derived NDVI to assess ecological responses to environmental

719   change. Trends Ecol Evol [Internet]. 2005 Sep 1 [cited 2025 Jan 28];20(9):503–

720   10. Available from:

721   https://linkinghub.elsevier.com/retrieve/pii/S016953470500162X

722   25.   Asner GP, Kellner JR, Kennedy-Bowdoin T, Knapp DE, Anderson C, Martin RE.

723   Forest Canopy Gap Distributions in the Southern Peruvian Amazon. Chen HYH,

724   editor. PLoS One [Internet]. 2013 Apr 15 [cited 2025 Jan 28];8(4):e60875.

725   Available from: https://dx.plos.org/10.1371/journal.pone.0060875

726   26.   Zhang X, Friedl MA, Schaaf CB, Strahler AH, Hodges JCF, Gao F, et al.

727   Monitoring vegetation phenology using MODIS. Remote Sens Environ

728   [Internet]. 2003 Mar 1 [cited 2025 Jan 28];84(3):471–5. Available from:

729   https://linkinghub.elsevier.com/retrieve/pii/S0034425702001359

27.   Perrone M, Di Febbraro M, Conti L, Divíšek J, Chytrý M, Keil P, et al. The

relationship between spectral and plant diversity: Disentangling the influence

of metrics and habitat types at the landscape scale. Remote Sens Environ

[Internet]. 2023 Aug;293:113591. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0034425723001426

28.   Farwell LS, Gudex-Cross D, Anise IE, Bosch MJ, Olah AM, Radeloff VC, et al.

Satellite image texture captures vegetation heterogeneity and explains

patterns of bird richness. Remote Sens Environ [Internet]. 2021 [cited 2025 Sep

9];253:112175. Available from: https://doi.org/10.1016/j.rse.2020.112175

29.   Ranganathan J, Chan KMA, Daily GC. Satellite detection of bird communities in

tropical countryside. Ecological Applications. 2007 Jul;17(5):1499–510.

30.   Barros F de C, Almeida SM, Godoy BS, Silva RR da, Silva LC, de Moraes KF, et al.

Taxonomic and functional diversity of bird communities in mining areas

undergoing passive and active restoration in eastern Amazon. Ecol Eng

[Internet]. 2022 Sep;182:106721. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0925857422001823

31.   Bregman TP, Lees AC, Seddon N, MacGregor HEA, Darski B, Aleixo A, et al.

Species interactions regulate the collapse of biodiversity and ecosystem

function in tropical forest fragments. Ecology [Internet]. 2015

Oct;96(10):2692–704. Available from:

https://esajournals.onlinelibrary.wiley.com/doi/10.1890/14-1731.1

32.   Turner W, Spector S, Gardiner N, Fladeland M, Sterling E, Steininger M. Remote

sensing for biodiversity science and conservation. Trends Ecol Evol [Internet].

2003 Jun 1 [cited 2025 Jan 28];18(6):306–14. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0169534703000703

33. Ludwig A, Meyer H, Nauss T. Automatic classification of Google Earth images

for a larger scale monitoring of bush encroachment in South Africa.

International Journal of Applied Earth Observation and Geoinformation

[Internet]. 2016 Aug 1 [cited 2025 Jan 28];50:89–94. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0303243416300265

34. Pettorelli N, Safi K, Turner W. Satellite remote sensing, biodiversity research

and conservation of the future. Philosophical Transactions of the Royal Society

B: Biological Sciences [Internet]. 2014 May 26 [cited 2020 Nov

13];369(1643):20130190. Available from:

https://royalsocietypublishing.org/doi/10.1098/rstb.2013.0190

35. Skidmore AK, Pettorelli N, Coops NC, Geller GN, Hansen M, Lucas R, et al.

Environmental science: Agree on biodiversity metrics to track from space.

Nature [Internet]. 2015 Jul 23 [cited 2020 Nov 17];523(7561):403–5. Available

from: https://www.nature.com/articles/523403a

36. Rocchini D, Boyd DS, Féret J, Foody GM, He KS, Lausch A, et al. Satellite remote

sensing to monitor species diversity: potential and pitfalls. Skidmore A,

Chauvenet A, editors. Remote Sens Ecol Conserv [Internet]. 2016 Feb 10 [cited

2022 Jun 21];2(1):25–36. Available from:

https://zslpublications.onlinelibrary.wiley.com/doi/10.1002/rse2.9

37. Csillik O, Kumar P, Mascaro J, O'Shea T, Asner GP. Monitoring tropical forest

carbon stocks and emissions using Planet satellite data. Sci Rep [Internet]. 2019

Dec 28;9(1):17831. Available from: http://www.nature.com/articles/s41598-

019-54386-6

38.   J van Proosdij AS, M Sosef MS, Wieringa JJ, Raes S J van Proosdij

andrevanproosdij NA, M Sosef MS, Wieringa JJ. Minimum required number of

specimen records to develop accurate species distribution models. Ecography

[Internet]. 2016;39:542–52. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/ecog.01509

39.   MapBiomas. MapBiomas Peru [Internet]. 2023 [cited 2023 Aug 21]. Available

from: https://peru.mapbiomas.org/en/colecciones-de-mapbiomas-peru/

40.   Carvalho WD, Fluck IE, de Castro IJ, Hilário RR, Martins ACM, de Toledo JJ, et al.

Elevation drives taxonomic, functional and phylogenetic β-diversity of

phyllostomid bats in the Amazon biome. J Biogeogr [Internet]. 2023 Jan 24

[cited 2025 Feb 3];50(1):70–85. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/jbi.14533

41.   Nores M. Species Richness in the Amazonian Bird Fauna from an Evolutionary

Perspective. Emu - Austral Ornithology [Internet]. 2000 Dec 22 [cited 2025 Feb

3];100(5):419–30. Available from:

https://www.tandfonline.com/doi/full/10.1071/MU0007S

42.   Huete AR. Vegetation Indices, Remote Sensing and Forest Monitoring. Geogr

Compass [Internet]. 2012 Sep 20 [cited 2023 Sep 7];6(9):513–32. Available

from: https://compass.onlinelibrary.wiley.com/doi/10.1111/j.1749-

8198.2012.00507.x

43.   Ollinger S V. Sources of variability in canopy reflectance and the convergent

properties of plants. New Phytologist [Internet]. 2011 Jan 16 [cited 2024 Feb

29];189(2):375–94. Available from:

https://nph.onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.2010.03536.x

44. Wang R, Gamon JA, Schweiger AK, Cavender-Bares J, Townsend PA, Zygielbaum
     AI, et al. Influence of species richness, evenness, and composition on optical
     diversity: A simulation study. Remote Sens Environ [Internet]. 2018 Jun [cited
     2022 Sep 29];211:218–28. Available from:
     https://linkinghub.elsevier.com/retrieve/pii/S003442571830155X

45. Quiñones M, Vissers M, Hoekman D. K&C Science Report - Phase 1 Tropical
     Forest and Wetlands Mapping, Case Study Borneo. 2007 Mar.

46. Hilker T, Lyapustin AI, Tucker CJ, Hall FG, Myneni RB, Wang Y, et al. Vegetation
     dynamics and rainfall sensitivity of the Amazon. Proceedings of the National
     Academy of Sciences [Internet]. 2014 Nov 11 [cited 2025 Jan
     30];111(45):16041–6. Available from:
     https://pnas.org/doi/full/10.1073/pnas.1404870111

47. Morton DC, DeFries RS, Shimabukuro YE, Anderson LO, Del Bon Espírito-Santo
     F, Hansen M, et al. Rapid Assessment of Annual Deforestation in the Brazilian
     Amazon Using MODIS Data. Earth Interact [Internet]. 2005 Jun 1 [cited 2025
     Jan 30];9(8):1–22. Available from:
     https://journals.ametsoc.org/doi/10.1175/EI139.1

48. Samarawickrama U, Piyaratne D, Ranagalage M. Relationship between NDVI
     with Tasseled cap Indices: A Remote Sensing based Analysis. 2017;3.

49. Remsen, J V., Good DA. Misuse of Data from Mist-Net Captures to Assess
     Relative Abundance in Bird Populations. Auk [Internet]. 1996 Apr;113(2):381–

823  98. Available from: https://academic.oup.com/auk/article/113/2/381-

824  398/5168448

825 50. Rosenstock SS, Anderson DR, Giesen KM, Leukering T, Carter MF. Landbird

826  Counting Techniques: Current Practices and an Alternative. Thompson III F,

827  editor. Auk [Internet]. 2002 Jan 1 [cited 2025 Jan 31];119(1):46–53. Available

828  from: https://academic.oup.com/auk/article/119/1/46/5561821

829 51. Simons TR, Alldredge MW, Pollock KH, Wettroth JM. Experimental Analysis of

830  The Auditory Detection Process on Avian Point Counts. Dufty AM, editor. Auk

831  [Internet]. 2007 Jul 1;124(3):986–99. Available from:

832  https://academic.oup.com/auk/article/124/3/986/5562805

833 52. Kery M. ESTIMATING ABUNDANCE FROM BIRD COUNTS: BINOMIAL MIXTURE

834  MODELS UNCOVER COMPLEX COVARIATE RELATIONSHIPS. Auk [Internet]. 2008

835  Apr;125(2):336–45. Available from:

836  https://academic.oup.com/auk/article/125/2/336-345/5148162

837 53. MacKenzie DI, Nichols JD, Hines JE, Knutson MG, Franklin AB. ESTIMATING SITE

838  OCCUPANCY, COLONIZATION, AND LOCAL EXTINCTION WHEN A SPECIES IS

839  DETECTED IMPERFECTLY. Ecology [Internet]. 2003 Aug [cited 2025 Jan

840  31];84(8):2200–7. Available from: http://doi.wiley.com/10.1890/02-3090

841 54. Royle JA, Dorazio RM. Hierarchical Modeling and Inference in Ecology

842  [Internet]. Hierarchical Modeling and Inference in Ecology: The Analysis of Data

843  from Populations, Metapopulations and Communities. Elsevier; 2009 [cited

844  2025 Jan 31]. 1–444 p. Available from:

845  https://linkinghub.elsevier.com/retrieve/pii/B9780123740977X00014

846   55.   Guillera-Arroita G, Lahoz-Monfort JJ. Designing studies to detect differences in

847   species occupancy: power analysis under imperfect detection. Methods Ecol

848   Evol [Internet]. 2012 Oct;3(5):860–9. Available from:

849   https://onlinelibrary.wiley.com/doi/10.1111/j.2041-210X.2012.00225.x

850   56.   Doser JW, Finley AO, Kéry M, Zipkin EF. spOccupancy: An R package for single-

851   species, multi-species, and integrated spatial occupancy models. Methods Ecol

852   Evol. 2022 Aug 30;13(8):1670–8.

853   57.   Doser J. https://www.jeffdoser.com/files/spoccupancy-

854   web/articles/modelconsiderations. 2023. Convergence diagnostics and other

855   considerations when fitting spatial occupancy models.

856   58.   Hooten MB, Hobbs NT. A guide to Bayesian model selection for ecologists. Ecol

857   Monogr [Internet]. 2015 Feb 1 [cited 2024 Mar 12];85(1):3–28. Available from:

858   https://esajournals.onlinelibrary.wiley.com/doi/10.1890/14-0661.1

859   59.   Kéry M, Royle JA. Chapter 6 - Modeling Abundance with Counts of Unmarked

860   Individuals in Closed Populations: Binomial N-mixture Models. In: Kéry M,

861   Royle JA, editors. Applied Hierarchical Modeling in Ecology [Internet]. Boston:

862   Academic Press; 2016. p. 219–312. Available from:

863   https://www.sciencedirect.com/science/article/pii/B9780128013786000060

864   60.   Watanabe S. Asymptotic Equivalence of Bayes Cross Validation and Widely

865   Applicable Information Criterion in Singular Learning Theory. Journal of

866   Machine Learning Research. 2010;11:3571–94.

867   61.   Gould MJ, Gould WR, Cain JW, Roemer GW. Validating the performance of

868   occupancy models for estimating habitat use and predicting the distribution of

869   highly-mobile species: A case study using the American black bear. Biol Conserv

870   [Internet]. 2019 Jun;234:28–36. Available from:

871   https://linkinghub.elsevier.com/retrieve/pii/S000632071831214X

872   62.   Mandrekar JN. Receiver Operating Characteristic Curve in Diagnostic Test

873   Assessment. Journal of Thoracic Oncology [Internet]. 2010 Sep [cited 2023 Apr

874   25];5(9):1315–6. Available from:

875   https://linkinghub.elsevier.com/retrieve/pii/S1556086415306043

876   63.   Dvořáková D, Šipoš J, Suchomel J. Impact of agricultural landscape structure on

877   the patterns of bird species diversity at a regional scale. Avian Res [Internet].

878   2023;14:100147. Available from:

879   https://linkinghub.elsevier.com/retrieve/pii/S2053716623000737

880   64.   Tu HM, Fan MW, Ko JCJ. Different Habitat Types Affect Bird Richness and

881   Evenness. Sci Rep [Internet]. 2020 Jan 27;10(1):1221. Available from:

882   https://www.nature.com/articles/s41598-020-58202-4

883   65.   Terraube J, Archaux F, Deconchat M, van Halder I, Jactel H, Barbaro L. Forest

884   edges have high conservation value for bird communities in mosaic landscapes.

885   Ecol Evol [Internet]. 2016 Aug 28;6(15):5178–89. Available from:

886   https://onlinelibrary.wiley.com/doi/10.1002/ece3.2273

887   66.   Willmer JNG, Püttker T, Prevedello JA. Global impacts of edge effects on

888   species richness. Biol Conserv [Internet]. 2022 Aug [cited 2023 Dec

889   14];272:109654. Available from:

890   https://linkinghub.elsevier.com/retrieve/pii/S0006320722002075

891   67.   Stouffer PC, Johnson EI, Bierregaard RO, Lovejoy TE. Understory Bird

892   Communities in Amazonian Rainforest Fragments: Species Turnover through 25

893   Years Post-Isolation in Recovering Landscapes. Evans DM, editor. PLoS One

[Internet]. 2011 Jun 22;6(6):e20543. Available from:

https://dx.plos.org/10.1371/journal.pone.0020543

68. Stratford JA, Stouffer PC. Forest fragmentation alters microhabitat availability

for Neotropical terrestrial insectivorous birds. Biol Conserv [Internet]. 2015

Aug [cited 2025 Feb 4];188:109–15. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0006320715000385

69. Şekercioğlu ÇH, Primack RB, Wormworth J. The effects of climate change on

tropical birds. Biol Conserv [Internet]. 2012 Apr [cited 2025 Feb 4];148(1):1–18.

Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0006320711003880

70. Barlow J, Peres CA, Henriques LMP, Stouffer PC, Wunderle JM. The responses

of understorey birds to forest fragmentation, logging and wildfires: An

Amazonian synthesis. Biol Conserv [Internet]. 2006 Mar [cited 2025 Feb

4];128(2):182–92. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0006320705003897

71. Barlow J, Mestre LAM, Gardner TA, Peres CA. The value of primary, secondary

and plantation forests for Amazonian birds. Biol Conserv [Internet]. 2007 Apr

[cited 2025 Feb 4];136(2):212–31. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0006320706005271

72. Sekercioglu CH. Bird functional diversity and ecosystem services in tropical

forests, agroforests and agricultural areas. J Ornithol [Internet]. 2012 Aug

27;153(S1):153–61. Available from: http://link.springer.com/10.1007/s10336-

012-0869-4

73. Peh KS -H., Sodhi NS, De Jong J, Sekercioglu CH, Yap CA -M., Lim SL -H.

Conservation value of degraded habitats for forest birds in southern Peninsular

Malaysia. Divers Distrib [Internet]. 2006 Sep 26 [cited 2025 Feb 4];12(5):572–

81. Available from: https://onlinelibrary.wiley.com/doi/10.1111/j.1366-

9516.2006.00257.x

74. Lees AC, Peres CA. Avian life-history determinants of local extinction risk in a

hyper-fragmented neotropical forest landscape. Anim Conserv [Internet]. 2008

Apr 27;11(2):128–37. Available from:

https://zslpublications.onlinelibrary.wiley.com/doi/10.1111/j.1469-

1795.2008.00162.x

75. Rutt CL, Jirinec V, Cohn-Haft M, Laurance WF, Stouffer PC. Avian ecological

succession in the Amazon: A long-term case study following experimental

deforestation. Ecol Evol [Internet]. 2019 Dec 27 [cited 2025 Feb

5];9(24):13850–61. Available from:

https://onlinelibrary.wiley.com/doi/10.1002/ece3.5822

76. Thornton DH, Branch LC, Sunquist ME. The influence of landscape, patch, and

within-patch factors on species presence and abundance: a review of focal

patch studies. Landsc Ecol [Internet]. 2011 Jan 30;26(1):7–18. Available from:

http://link.springer.com/10.1007/s10980-010-9549-z

77. Laurance WF, Camargo JLC, Luizão RCC, Laurance SG, Pimm SL, Bruna EM, et al.

The fate of Amazonian forest fragments: A 32-year investigation. Biol Conserv

[Internet]. 2011 Jan [cited 2025 Feb 13];144(1):56–67. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0006320710004209

78.   Hess LL, Melack JM, Affonso AG, Barbosa C, Gastil-Buhl M, Novo EMLM.

941   Wetlands of the Lowland Amazon Basin: Extent, Vegetative Cover, and Dual-

942   season Inundated Area as Mapped with JERS-1 Synthetic Aperture Radar.

943   Wetlands [Internet]. 2015 Aug 20;35(4):745–56. Available from:

944   https://link.springer.com/10.1007/s13157-015-0666-y

945   79.   Laurance WF, Albernaz AKM, Costa C Da. Is deforestation accelerating in the

946   Brazilian Amazon? Environ Conserv [Internet]. 2001 Dec 10 [cited 2025 Feb

947   13];28(4):305–11. Available from:

948   https://www.cambridge.org/core/product/identifier/S0376892901000339/typ

949   e/journal_article

950   80.   Blake JG, Loiselle BA. Enigmatic declines in bird numbers in lowland forest of

951   eastern Ecuador may be a consequence of climate change. PeerJ [Internet].

952   2015 Aug 11 [cited 2025 Feb 12];3(8):e1177. Available from:

953   https://peerj.com/articles/1177

954   81.   Rutt CL, Stouffer PC. Seasonal dynamics of flock interaction networks across a

955   human-modified landscape in lowland Amazonian rain forest. Ecological

956   Applications [Internet]. 2021 Mar 21 [cited 2025 Feb 12];31(2). Available from:

957   https://esajournals.onlinelibrary.wiley.com/doi/10.1002/eap.2235

958   82.   Robinson WD, Curtis JR. Creating benchmark measurements of tropical forest

959   bird communities in large plots. Condor [Internet]. 2020 Sep 11 [cited 2025 Feb

960   7];122(3):1–15. Available from:

961   https://academic.oup.com/condor/article/doi/10.1093/condor/duaa015/5820

962   537

83.   Peña-Lara VA, Dupuy JM, Reyes-Garcia C, Sanaphre-Villanueva L, Portillo-

Quintero CA, Hernández-Stefanoni JL. Modelling Species Richness and

Functional Diversity in Tropical Dry Forests Using Multispectral Remotely

Sensed and Topographic Data. Remote Sens (Basel) [Internet]. 2022 Nov 23

[cited 2025 Feb 5];14(23):5919. Available from: https://www.mdpi.com/2072-

4292/14/23/5919

84.   Suttidate N, Hobi ML, Pidgeon AM, Round PD, Coops NC, Helmers DP, et al.

Tropical bird species richness is strongly associated with patterns of primary

productivity captured by the Dynamic Habitat Indices. Remote Sens Environ

[Internet]. 2019 Oct [cited 2025 Feb 5];232:111306. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0034425719303256

85.   Sheeren D, Bonthoux S, Balent G. Modeling bird communities using

unclassified remote sensing imagery: Effects of the spatial resolution and data

period. Ecol Indic [Internet]. 2014 Aug [cited 2023 Dec 1];43:69–82. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S1470160X14000818

86.   Regos A, Gómez-Rodríguez P, Arenas-Castro S, Tapia L, Vidal M, Domínguez J.

Model-Assisted Bird Monitoring Based on Remotely Sensed Ecosystem

Functioning and Atlas Data. Remote Sens (Basel) [Internet]. 2020 Aug

7;12(16):2549. Available from: https://www.mdpi.com/2072-4292/12/16/2549

87.   Cruz M, Pradel W, Juarez H, Hualla V, Suarez V. Deforestation Dynamics in

Peru. A Comprehensive Review of Land Use, Food Systems, and Socio-

Economic Drivers [Internet]. 2023 Dec [cited 2025 Feb 6]. Available from:

https://hdl.handle.net/10568/135509

88.   Marcus M, Gutierrez-Velez VH, Cronkleton P. Land use change in four

987   landscapes in the Peruvian Amazon [Internet]. Land use change in four

988   landscapes in the Peruvian Amazon. Center for International Forestry Research

989   (CIFOR); 2020 [cited 2025 Feb 6]. Available from:

990   https://www.cifor.org/knowledge/publication/7736

991   89.   Gallardo-Cruz JA, Solórzano J V., González EJ, Meave JA. The Effect of Spatial

992   Scale on the Prediction of Tropical Forest Attributes from Image Texture.

993   Źróbek-Sokolnik A, editor. International Journal of Forestry Research [Internet].

994   2024 Apr 24 [cited 2025 Feb 6];2024(1):1–10. Available from:

995   https://www.hindawi.com/journals/ijfr/2024/7178211/

996   90.   Ploton P, Mortier F, Réjou-Méchain M, Barbier N, Picard N, Rossi V, et al.

997   Spatial validation reveals poor predictive performance of large-scale ecological

998   mapping models. Nat Commun [Internet]. 2020 Sep 11 [cited 2025 Feb

999   6];11(1):4540. Available from: https://www.nature.com/articles/s41467-020-

1000   18321-y

1001   91.   Morrison LW. Observer error in vegetation surveys: a review. Journal of Plant

1002   Ecology [Internet]. 2016 Aug 1 [cited 2025 Feb 6];9(4):367–79. Available from:

1003   https://academic.oup.com/jpe/article-lookup/doi/10.1093/jpe/rtv077

1004   92.   Basile M, Storch I, Mikusiński G. Abundance, species richness and diversity of

1005   forest bird assemblages – The relative importance of habitat structures and

1006   landscape context. Ecol Indic [Internet]. 2021 Dec [cited 2023 Dec

1007   14];133:108402. Available from:

1008   https://linkinghub.elsevier.com/retrieve/pii/S1470160X21010670

93.   Aubry KB, Raley CM, McKelvey KS. The importance of data quality for

generating reliable distribution models for rare, elusive, and cryptic species.

Baldwin RF, editor. PLoS One [Internet]. 2017 Jun 22;12(6):e0179152. Available

from: https://dx.plos.org/10.1371/journal.pone.0179152

94.   McCord SE, Webb NP, Van Zee JW, Burnett SH, Christensen EM, Courtright EM,

et al. Provoking a Cultural Shift in Data Quality. Bioscience [Internet]. 2021 Jun

1;71(6):647–57. Available from:

https://academic.oup.com/bioscience/article/71/6/647/6188799

95.   Lepage D. Tambopata bird checklist - Avibase - Bird Checklists of the World

[Internet]. 2024 [cited 2025 Feb 10]. Available from: https://avibase.bsc-

eoc.org/checklist.jsp?lang=EN&p2=1&list=howardmoore&synlang=®ion=PE

mdtp&version=text&lifelist=&highlight=0

96.   Alvarez-Berríos N, Campos-Cerqueira M, Hernández-Serna A, Amanda Delgado

CJ, Román-Dañobeytia F, Aide TM. Impacts of Small-Scale Gold Mining on Birds

and Anurans Near the Tambopata Natural Reserve, Peru, Assessed Using

Passive Acoustic Monitoring. Trop Conserv Sci [Internet]. 2016 Jun 1 [cited

2025 Feb 10];9(2):832–51. Available from:

https://journals.sagepub.com/doi/10.1177/194008291600900216

97.   Martin TE, Nightingale J, Baddams J, Monkhouse J, Kaban A, Sastranegara H, et

al. Variability in the Effectiveness of Two Ornithological Survey Methods

between Tropical Forest Ecosystems. Pandit MK, editor. PLoS One [Internet].

2017 Jan 10 [cited 2025 Feb 7];12(1):e0169786. Available from:

https://dx.plos.org/10.1371/journal.pone.0169786

98.   Anderson DL. Ground Versus Canopy Methods for the Study of Birds in Tropical

Forest Canopies: Implications for Ecology and Conservation. Condor [Internet].

2009 May 1 [cited 2025 Feb 7];111(2):226–37. Available from:

https://academic.oup.com/condor/article/111/2/226-237/5152440

99.   Steenweg R, Hebblewhite M, Whittington J, McKelvey K. Species-specific

differences in detection and occupancy probabilities help drive ability to detect

trends in occupancy. Ecosphere [Internet]. 2019 Apr 3 [cited 2023 Oct 9];10(4).

Available from:

https://esajournals.onlinelibrary.wiley.com/doi/10.1002/ecs2.2639

100.   Mulvaney JM, Cherry MI. The effectiveness of point counts and mist-netting in

surveying Afrotemperate forest bird community structure in South Africa. Emu

- Austral Ornithology [Internet]. 2020 Apr 2;120(2):103–13. Available from:

https://www.tandfonline.com/doi/full/10.1080/01584197.2020.1726186

101.   Asner GP, Powell GVN, Mascaro J, Knapp DE, Clark JK, Jacobson J, et al. High-

resolution forest carbon stocks and emissions in the Amazon. Proceedings of

the National Academy of Sciences [Internet]. 2010 Sep 21;107(38):16738–42.

Available from: https://pnas.org/doi/full/10.1073/pnas.1004875107

102.   Smith P, Bustamante M, Ahammad H, Clark H, Dong H, Elsiddig EA. Agriculture,

Forestry and Other Land Use (AFOLU). Climate Change 2014: Mitigation of

Climate Change. 2014;811–922.

103.   Harrison ME, Boonman A, Cheyne SM, Husson SJ, Marchant NC, Struebig MJ.

Biodiversity Monitoring Protocols for REDD+: Can a One-Size-Fits-All Approach

Really Work? Trop Conserv Sci [Internet]. 2012 Mar 1 [cited 2024 Apr

24];5(1):1–11. Available from:

http://journals.sagepub.com/doi/10.1177/194008291200500102

104. West TAP, Börner J, Sills EO, Kontoleon A. Overstated carbon emission

reductions from voluntary REDD+ projects in the Brazilian Amazon.

Proceedings of the National Academy of Sciences [Internet]. 2020 Sep 29 [cited

2024 Apr 24];117(39):24188–94. Available from:

https://pnas.org/doi/full/10.1073/pnas.2004334117

105. Betts MG, Yang Z, Hadley AS, Hightower J, Hua F, Lindenmayer D, et al.

Quantifying forest degradation requires a long-term, landscape-scale

approach. Nat Ecol Evol [Internet]. 2024 Apr 12; Available from:

https://www.nature.com/articles/s41559-024-02409-5

106. Stephenson P. Technological advances in biodiversity monitoring: applicability,

opportunities and challenges. Curr Opin Environ Sustain [Internet]. 2020

Aug;45:36–41. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S1877343520300592

107. Magurran AE, Baillie SR, Buckland ST, Dick JMcP, Elston DA, Scott EM, et al.

Long-term datasets in biodiversity research and monitoring: assessing change

in ecological communities through time. Trends Ecol Evol [Internet]. 2010 Oct 1

[cited 2024 Jun 7];25(10):574–82. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0169534710001552

108. Nichols J, Williams B. Monitoring for conservation. Trends Ecol Evol [Internet].

2006 Dec;21(12):668–73. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0169534706002680

109.   Farr TG, Rosen PA, Caro E, Crippen R, Duren R, Hensley S, et al. The shuttle

1079   radar topography mission. Reviews of Geophysics [Internet]. 2007 May 19

1080   [cited 2021 Nov 15];45(2):RG2004. Available from:

1081   http://doi.wiley.com/10.1029/2005RG000183

1082

1083