Remote sensing of bird communities in the Peruvian

Amazon: evaluating Landsat predictors for biodiversity

monitoring

Andrew C. Slater[1] (Corresponding author) – acs_africa@yahoo.co.uk - https://orcid.org/0009-

0001-3855-686X

Chris A. Kirkby [2] – c.a.kirkby@gmail.com

Chris Ketola [2] – chrisketola@gmail.com

Ian R. Hartley[1] - i.hartley@lancaster.ac.uk

Alex Bush[1] - alex.bush@lancaster.ac.uk

[1] Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ

[2] Fauna Forever, Puerto Maldonado, Peru

Word count: Abstract = 189, Main text = 5373, References = 3856, Tables and figure legends

= 328

Number of: Tables = 1, Figures = 6

1 Abstract

Over one-third of the Amazon rainforest has been lost or degraded through human activities. While Earth observation satellites effectively monitor forest cover, assessing the impacts of degradation on biodiversity at large scales remains challenging. We analysed 3,129 bird surveys conducted over 16 years in the Tambopata Forest, south-eastern Peru, to evaluate whether remote sensing (RS) variables derived from Landsat imagery can predict variation in bird community composition. We compared the performance of RS-based predictors with traditional habitat descriptors in modelling the occurrence probabilities of 135 commonly recorded bird species. Models using Landsat reflectance outperformed those based on habitat data for predicting species occupancy (mean AUC = 0.68 vs 0.58) and achieved high predictive accuracy (AUC > 0.7) for more species (49 vs 20). However, low detection rates limited the ability of all models to estimate true community composition and to detect temporal change. To address this, we recommend survey designs that prioritise greater replication at fewer sites, thereby improving detection rates and the power to monitor biodiversity trends. Our findings highlight the potential for integrating satellite-based environmental variables with improved survey designs to enhance biodiversity monitoring in tropical forests.

2 Introduction

Tropical forests are the most biodiverse areas on Earth (1,2), and support over a billion people with food, materials, and services (3). They also play a key role in the global carbon cycle, accounting for half of terrestrial uptake and two-thirds of forest carbon sinks in biomass, soil, and deadwood (4,5). However, clear-cut logging and commercial agriculture are altering or completely destroying forest habitats. This is evidenced by the fact that over 17% of the Amazon Basin, the world's largest continuous tropical forest, has been deforested and 38% of the remaining area has been degraded (6–8).

Conservation strategies, such as the Reducing Emissions from Deforestation and Degradation (REDD+) initiative, have generally reduced deforestation, enhanced carbon stocks, and improved sustainable forest management (9–11). However, despite the amount of tropical forest falling under protection, half of the world's tropical forest reserves continue to lose taxonomic and functional biodiversity due to disturbances beyond forest loss (12,13). The loss of forest cover can be used to estimate the level and location of carbon loss and is routinely mapped in near real-time (14), but to assess the efficacy of biodiversity conservation strategies, such as REDD+, beyond forest cover conservation, it is necessary to ascertain the current condition of biodiversity within the ecosystem and where and how it changes in the future (15–17).

Ecosystem structure is often described by the type, quantity and distribution of its components (18). However, categorical elements such as intact, logged, secondary, plantation, can be ambiguous, subjective, and require prior knowledge to define them (19,20). Despite any *a priori* assumptions, the selected elements may not be the most effective descriptors of an ecosystem's condition. In this regard, essential biodiversity

variables, including species distribution, abundance, and community composition, have been

proposed as additional criteria to monitor (21,22). Thus, assessing forest health and

biodiversity change requires monitoring that goes beyond simply quantifying deforestation

levels. Satellite remote sensing (RS) cannot directly see essential biodiversity variables, but it

allows for the measurement of deforestation along with land cover and habitat structure (14).

RS can also provide broader, scalable environmental descriptors, alternative to field surveys,

that act as proxies for essential biodiversity variables, for example, ecological diversity,

phenology and physiology (23–26). Continuous remotely sensed variables have also been

shown to be better than land class at describing plant species richness and diversity (27). RS

surface reflectance values can therefore be used to infer vegetation and environmental

structure as continuous gradients, rather than discrete categories, over a region of interest.

This approach avoids the need to pre-define habitat classes that may overlook important

ecological variation and reduces the mismatch between habitat classification and the scale at

which they can be measured. While RS variables can infer potential structure and ecological

richness, there is little understanding of how they relate to ecological communities within

forests (28–30).

Thus, the measurement of structure or deforestation alone provides only a partial description

of an environment's condition, but faunal biodiversity can regulate forest dynamics (12) and

be used as an indicator of ecosystem state (31,32). Many studies have shown that land use

and vegetation structure influence the biodiversity of an area. For example, dung beetle

community composition and soil bioturbation are influenced by the intensity of logging within

tropical forests and the spatial scale at which it is measured (33). Similarly, the richness and

biological function of mammal communities are impacted by proportional levels of

deforestation and the resulting size of forest patches in Atlantic Amazonian forests (34).

Furthermore, insect community richness and composition have been shown to respond to

and reflect forest structure, forest composition, plant diversity and forest age (35–38). Finally,

the structure of bird communities has been used both to assess the effectiveness of

reforestation (39), and to show that they are negatively affected by patch size reduction

within the Brazilian Amazon (40).

However, the biodiversity surveys required to adequately monitor indicator taxa are time

consuming and expensive, thus limiting the effective area that can be monitored in space and

time (41,42). Nevertheless, results from biodiversity surveys can be linked in statistical models

with RS descriptors of the Earth's surface, to estimate community and ecosystem structure

across a landscape with limited expense (43). We would expect Amazonian bird communities

to be broadly influenced by the characteristics of the forest in which they occur, and if

changes in forest structure led to changes in bird communities, we would expect remotely

sensed surface reflectance values to correlate with this.

Furthermore, the continually updated nature of RS data could allow for this method to be

used in ongoing monitoring to detect changes in community structure. As such, lastly we

explore how the number of survey stations and replicate visits affects the power to detect

change, helping to guide efficient survey design. By bringing these elements together, we aim

to provide a practical tool for policy makers and ecologists to assess the successes and failures

of management strategies, and to support decisions about financial incentives.

3 Methods

3.1 Study area and surveys

Survey data were collected by FaunaForever (www.faunaforever.org) to monitor species diversity across landscapes in the Madre de Dios region, near the Tambopata River and Puerto Maldonado in south-eastern Peru (Figure 1). Researchers and volunteers conducted bird surveys at 637 stations between 2004 and 2020. The stations were clustered around 25 centres that were located between 1.5km and 170km apart. Within centres, stations were between 200m and several kilometres apart and surveyed one to 35 times (median = 4), totalling 3,129 surveys. Stations surveyed in multiple years were treated as independent, since populations and forest structure can change over extended periods of time. This resulted in a total of 1,135 independent stations, which were surveyed between one and 21 times, with a mode of 1 survey, a median of 2 surveys and a mean of 2.9 surveys per site (SD ±2.4). Surveys used point counts (10-minute visual/audio records), or net counts (36m mist-net captures). Of 358 species identified, only 138 observed at over 10 stations were used for model training per prevalence recommendations (44).

3.2 Environmental variables

Habitat was classified by land class and forest degradation. Land classes (floodplain forest, terra firma forest, and agricultural) were based on the MapBiomas Peru Project, Collection-**1** of the Annual Land Cover and Land Use Series for Peru 2013; (MapBiomas, 2023). Forest degradation was determined from a primary forest raster using the 2000-2019 forest cover loss data set (14). Pixels were classified as primary forest if no deforestation year was given,

assuming they had remained intact. Subsequently, secondary forests were defined as areas

that had undergone recorded deforestation since 2000. This classification is primary/not

primary and does not differentiate between successional stages of secondary forest. The

proportion of non-primary forest, equating to forest assumed to have been deforested or

degraded, was calculated within three different radii around each station (1km, 2km and

5km). No cross validation was performed between the MapBiomas and Hansen datasets, so

forest classifications may not align.

Of the 1135 surveys, 489 were on terra firma, 603 on floodplain and 43 on agricultural land,

and 1109 were defined as being inside primary forest, and 26 in secondary forest. Elevation

can influence the distribution of tropical forest bird species (46,47). However, all survey points

were within the narrow range between 175m and 310m elevation, and unlikely to affect

model performance; it was therefore not included in the analysis.

3.3   Satellite derived variables

Reflectance signatures at different wavelengths can be employed to assess the greenness and

moisture content of vegetation, with some correlation observed between these signatures

and the leaf structure of a plant (48). Reflectance and water content descriptors help describe

forest canopy physiology (49). No single metric captures canopy spectral diversity (50), and

the spectral characteristics and spatial scales that best capture canopy variation are unknown,

so various indices were used to describe it. While many satellite platforms now provide

passive optical surface reflectance data, as well as active radar reflectance, the Landsat

missions were the only platforms that covered the entire period of the field surveys (Landsat-

5 for 2004-2011, Landsat-7 for 2012, and Landsat-8 for 2013-2020). Frequent cloud cover over

tropical forests can greatly reduce the number of available images that have sufficient clarity

for analysis (51). To address poor visibility, annual cloud-masked, median pixel value images

were created in Google Earth Engine. Annual composites may mask seasonal changes such as

leaf flushing and fruiting cycles, or logging and burning (26,52), although phenological

variation in tropical forests can be lower than in some other landscapes (53). Each image

contained six reflectance bands, which were used to calculate seven vegetation indices: the

normalised difference vegetation index (NDVI), enhanced vegetation index (EVI), normalised

burn ratio (NBR), normalised difference water index (NDWI), and the trio of tasselled cap

wetness, greenness, and brightness (TCW, TCG, TCB). Both NDVI and EVI were used to detect

vegetation cover and vigour. NDVI was included as one of the most commonly used indices,

but it is prone to saturation in dense forests such as the Amazon, whereas EVI is better able

to penetrate the canopy and capture sub-layers. We used NBR to highlight freshly opened

soil, and NDWI to highlight water content within the canopy. Tasselled cap indices provide an

alternative way of describing reflectance and structure, and have been shown to correlate

with, but show more detail than, NDVI (54).

For each annual survey, the mean and standard deviation of pixels in each of the reflectance

bands and indices were calculated within 100m of each station. These 26 variables provide a

high-dimensional description of local vegetation reflectance. To reduce redundancy, we used

principal component analysis to condense the 26 variables, and the first five components,

explaining 96% variation in the data, were used. The distribution of the five principal

components and the proportion of deforestation at three scales are plotted in Figure 2, and

the relationship between the first two principal components with environmental descriptors

is shown in Figure 3.

3.4 Analysis

To quantify species detection and assess whether bird community composition can be predicted as a function of habitat or RS descriptors, we used spatial factor multi-species occupancy models. Observation depends on true presence (1 = present, 0 = absent; assumed constant over the year of surveillance), and the probability of detection. Detection can be influenced by survey method. Mist nets at ground level tend to under-sample canopy dwelling, large or less mobile species (55). Capture rates also vary with the type of net used (e.g. mesh size), and netting duration. Point counts detect vocal species well, but may under-sample silent or secretive species, and their effectiveness depends on background noise, weather, and observer skill (56,57). Perfect detection of all species is essentially impossible. Non-detection can bias covariate estimates and predictions of community composition (58). However, occupancy models account for imperfect detection by estimating species occurrence at each survey station, whether observed or not (59). Occupancy models also provide detection probabilities for all species at all stations and the influence of survey specific covariates on these probabilities (Royle and Dorazio, 2009). After fitting the model and estimating species detection and occurrence probabilities, the methods of Guillera-Arroita and Lahoz-Monfort (2012), were used to calculate the power of the study. The term "power" is the probability of detecting a proportional reduction in occupancy between surveys conducted at different times. We then calculated the survey effort required to achieve a mean power across all species of 0.7, to detect a 50% occupancy reduction. Although 0.8 is commonly used, our low detection rates made 0.7 a more feasible target that falls within the test ranges ((Guillera-Arroita & Lahoz-Monfort, 2012). We also assessed how detection power of various occupancy reductions varied with the number of stations and replicate surveys, to guide survey regimes for future monitoring.

Four models were compared: one fitted with habitat variables, one with principal

components from RS reflectance, one with no occurrence variables to act as an intercept-only

baseline, and one RS model re-fitted without spatial factors to assess their influence. All

models included survey method (point or net) and time of day (a.m. or p.m.) as covariates for

the probability of detection. Although some stations were surveyed up to 21 times, 99% of all

stations were surveyed 12 times or less. To improve the efficiency of model fitting and reduce

the computational resources required to run them, we fitted the modes using a maximum of

12 replicates per station.

Models were fitted using the Bayesian occupancy model package, "spOccupancy" (61). The

"spOccupancy" package enables the modelling of all species simultaneously, and accounts for

autocorrelation between species. In addition, spatial latent factors, which are akin to

unmeasured site-specific covariates that decay with distance, help account for residual

species correlations. Fewer factors were favoured to explain variation efficiently and avoid

overfitting, particularly with relatively few replicate surveys and rarely detected species. We

included three independent spatial factors, each derived from residual species correlations

after controlling for other variables and previous spatial factors. Each factor provides a decay

rate (phi), with the non-autocorrelation distance calculated as 3/phi.

Models were fitted with 50,000 iterations, discarding the first 25,000 as burn-in, and thinning

the rest by 50 to produce 500 posterior samples. Each posterior sample provides a probability

of occurrence (psi) for each species at each station and a latent presence/absence (z = 1/0)

calculated from this probability. Additionally, detection probability (p), given presence, is

provided per species per station for a single survey. As spatial models have poor chain

convergence, a single chain was used, with mixing assessed visually and via the Geweke

diagnostic (62). Models assumed no false presences and treated the entire study as a single

closed season. Model goodness-of-fit was measured using posterior predictive checks to calculate a Bayesian p-value, based on the differences between chi-squared values of observed and predicted data per species, grouped across stations. All models had Bayesian p-values between 0.467 and 0.477, indicating a good fit (63,64), and models were therefore subsequently compared using the widely applicable information criterion (WAIC) (Watanabe, 2010) and metrics of predictive performance. Performance was evaluated at the species-level, using area under the receiver operating curve (AUC), and at the station community-level, using species accumulation and Bray-Curtis's dissimilarity of community composition. Although model performance differs between training and withheld data, evaluation on independent datasets is rarely undertaken for occupancy models (66). We did so here by withholding a validation dataset comprising two stations per field centre, each surveyed at least three times, totalling 5% of the data. To validate model performance, we incorporated both occurrence and detection probabilities when comparing predictions with observations. Each of the model's posterior samples predicts two occupancy values per species. First is the occupancy probability (psi) per species per site, from which a latent occupancy (z = 1/0) is drawn. Detection probability (p) per species and site in a single survey was predicted, and this remained constant across posterior samples. Total detection probability per species and station was calculated as $1 - (1 - p)^n$, where n is the number of replicate surveys performed. This detection probability generated a binomial detection/non-detection (1/0) event per species per station. Latent detection values were then multiplied by occurrence (psi) and latent occupancy (z) for each species and station per posterior sample. Thus, model performance was based on the predicted probability of detection rather than occurrence in the validation data.

4 Results

4.1 Model fit

All models had Bayesian p-values between 0.467-0.477, indicating a good fit to the data. The model with the lowest WAIC value, thus indicating it was the best-performing model, was fitted with RS reflectance data and included spatial factors, followed by the covariate-free intercept model, the RS reflectance model fitted without spatial factors, and the highest WAIC indicating the worst performance was from the model fitted with environmental descriptors. The probability of detection (p) was consistent across all models. Surveys conducted in the morning using net counts had a median species-level p = 0.022, 95% CI [0.001, 0.179]. When conducting surveys in the afternoon, median detection probability significantly lower at p = 0.015, 95% CI [0.001, 0.018]. When using point rather than net surveys, the median p increased significantly to 0.065, 95% CI [0.000, 0.389]. However, point surveys did not significantly increase detection for three of the nine Orders represented. Those three were Caprimulgiformes (nightjars), and Coraciiformes (kingfishers and hornbills), and Galliformes (ground dwelling birds).

The mean probability of occurrence across all species in primary forest and an agricultural habitat, was 0.291, 95% CI [0.211, 0.375]. In secondary forest, the mean occupancy was higher than in primary forest, rising to 0.47, 95% CI [0.36, 0.61]. In comparison to agricultural landscapes, mean occupancy in floodplain increased to 0.36, 95% CI [0.30, 0.47], and in terra-firma mean occupancy increased to 0.37, 95% CI [0.30, 0.47]. Mean occupancy was not shown to vary significantly with deforestation within a 1-2 km radius, but increased by 0.057, 95% CI [0.008, 0.108] with each 10% increase in deforestation within a 5km radius.

When using RS data, the model suggested that spatial autocorrelation extended to 5km, 95%

CI [3.3km, 7.5km] for one factor and 7.5km, 95% CI [4.3km, 15km] for each of the other two

factors. When using habitat data, the model suggested that spatial autocorrelation extended

to 2km, 95% CI [2km, 2.3km] for one factor, to 30km, 95% CI [3.75km, ∞] for another, and

the third factor was not significant. These factors represent unmeasured variables that

influence the occurrence of species, and fade with distance. Although we do not know what

is driving this spatial structure, it can be influenced by anything from local vegetation cover

to regional climate or land use. Significant spatial autocorrelation confirm that patterns exist

beyond those explained by the known environmental variables.

4.2   Model validation

The predictive performances of models followed the implications made by their WAIC values,

and the model that produced the best predictive results was also fitted with RS reflectance

data and included spatial factors. When assessed on training data, this model had a mean

AUC of 0.88 (SD=0.06) and when predicting to independent validation sites, had a mean AUC

of 0.68 (SD=0.18). Species for which predictive performance was considered good (AUC>0.7)

(67) are hereafter referred to as high-AUC species, and this model had 49 high-AUC species,

the highest of any model. The performance of all four models is shown in Table 1. Species

taxonomic order was not shown to have any correlation with its AUC.

The predicted species richness of survey stations (6.86 ± 5.1) was close to the observed

average richness (6.00 ± 3.6), but when extended to the entire training dataset the model

predictions suggested that it would take much longer to detect the total number of species

included (Figure 4(A)). The mean predicted richness of test stations (9.94 ± 3.8) was also

comparable to the observed mean richness (10.82 ± 5.3) and indicated a similar accumulation

of species, both across stations (Figure 4(B)), and with increasing replication within a station

(Figure 4(C)). The species accumulation curve in Figure 4(C) come from one station that was

surveyed 12 times and illustrates the extent to which observed accumulated richness greatly

underestimates the total richness expected to be present at each location. A consequence of

this was that only a fraction of the species that occur at a station were observed, and

therefore the average Bray-Curtis dissimilarity between the list of birds observed and those

predicted was 0.8 (± 0.1). In locations with higher observed richness, the surveyed total

represented a greater proportion of the true expected total, and the dissimilarity between

predicted and observed communities was substantially lower (Figure 5).

4.3   Power to detect change

As a result of low detection probabilities and low mean replicate surveys, it was ascertained

that there is a mean probability across all species of 0.20 (±0.24 SD) of detecting a 50%

reduction in occupancy in a subsequent survey. There is a 70% or greater probability of

detecting a 50% reduction in only five of the 135 species modelled. Thus, if the desire is to

determine where occupancies are changing by surveying the same area in the future, the

current data cannot provide robust evidence.

Figure 6 illustrates how different numbers of survey stations and replicates per station affect

the median power to observe a reduction in bird occupancy. Each panel represents a different

proportional change that is sought to be detected, and each y-axis represents the probability

of detecting that change in at least half the bird species. The x-axis is the total number of

individual surveys that a study is able to undertake (i.e. one replicate at one station is one survey). Each line represents a different number of sites that the total number of surveys are divided among. For every probability of detecting the panel's proportional change (y-axis), the farthest left line represents the fewest total surveys required, and the most efficient number of sites to survey. For example, if our aims were to have a 70% confidence (0.7 on the y-axis) that we could detect when the occupancy of at least half the bird species had declined by 50% (panel C), the chart suggests that the fewest surveys required would be ~10,000 and spread between 100 and 500 sites. Calculations suggest that the requirements would be three replicate surveys at nearly 50,000 stations, or, if all stations were surveyed 27 times, only 447 stations would be required.

5 Discussion

Using over 3,000 bird surveys conducted in south-eastern Peru, we show that remotely sensed satellite data can slightly outperform landcover models in predicting bird community composition. We also highlight the low detectability of many species, which complicates biodiversity monitoring in rainforests. Although the survey design provided valuable insights into bird communities, it was not optimised to detect temporal change. Our results can guide more efficient survey strategies, leading to enhanced species detection, better capturing community composition, and strengthening the performance of RS models in predicting bird communities across landscapes.

5.1 Land class and surface reflectance

Studies differ on how forest and farmland affect bird species richness (68,69). We found more species in both floodplain and terra firma forests than in agricultural land, partially aligning with evidence that species richness can increase at forest boundaries, particularly where edges are sharp, such as where forest and agricultural land meet (70,71). Furthermore, the greater stratification and canopy complexity often found in primary forests typically support more specialist, range restricted, and endemic bird species compared to secondary forests (72–74). However, our analysis suggested that secondary forest had a greater species richness, consistent with findings that generalist and disturbance tolerant species as well as species that thrive in more open canopy environments are drawn to those areas, especially when they are connected to primary forest, thereby increasing species richness (75). Greater visibility, vocalisation and bird density in open secondary forests may also lead to more detections (76–78). The low mean number of replicates in our study could make secondary forests appear richer in species than primary forests, due to improved visibility rather than true richness.

Secondary forests require the initial loss of primary forests, and localised deforestation can increase bird species richness. Moderate primary deforestation can increase richness of neighbouring forest as displaced individuals move into it (79). It can also increase habitat heterogeneity, promoting colonisation by generalists and disturbance-tolerant species in the deforested area (79). Our findings that deforestation within a 5km radius increased mean occupancy support this. At this scale species dispersal, habitat connectivity, and edge effects shape community composition (80,81). In contrast, no effects were found at smaller distances, suggesting insufficient heterogeneity at that range. We defined deforestation as loss of primary forest, but deforested areas may have regrown into secondary forest, which

also showed high species richness. Misclassification could arise from pre-2000 deforestation

or from the omission of water bodies from analysis. Within floodplain forests, water levels

can fluctuate seasonally (82) and edge effects may occur at interfaces between large water

bodies and forests (83). These classification limitations should be considered when

interpreting differences in species occupancy across forest types. bird detectability may have

varied due to seasonal fluctuations, which are most pronounced in forest fragments and

secondary forest (72,84,85). Surveys were performed by multiple observers with varying

levels of experience, which can significantly influence bird species detectability (86).

Consequently, some variation in detection probabilities may be attributed to differences in

surveyor rather than actual ecological differences.

Surface reflectance diversity has been used to predict tropical forest and bird diversity (87)

and bird species richness found within them (88) and our findings support this. RS data

performed as well as, if not better than, land class in predicting bird communities (89,90).

Image seasonality and resolution matter for modelling bird communities in temperate regions

(89,90). However, persistent cloud cover in our study area required annual composites, and

mean reflectance values calculated over thousands of square metres make finer resolution

imagery unlikely to improve results.

Despite these challenges, our study has shown that RS data have the capacity to improve on

land class in predicting bird communities in tropical forests. Two broad land-class groups

could be differentiated by plotting the first two PCAs: secondary forest/agriculture and

primary forest/terra-firma/floodplain. Within those two groups, secondary forest, defined as

land deforested since 2000, had a large overlap with agricultural land, suggesting that

deforested land here tends to become agricultural rather than regenerating into secondary

forest (91,92). In the other group, primary forest overlapped more with terra-firma and

floodplain than with agriculture, though sizeable overlaps between all groups existed. This

may be explained by discrepancies brought about by course scale, out of date information or

human error in habitat classification (93–95). Additionally, land classes may oversimplify or

fail to capture the finer scale variation seen by RS data. Localised secondary descriptors of

structural variation, such as deadwood volume and tree girth, can also influence reflectance

values, and when included in conjunction with broad habitat type, have been shown to define

species richness far better than habitat type alone (96). This shows that RS data better capture

gradual changes in vegetation structure than can predefined classes. The challenge for future

research will be in deriving RS variables that can describe more ecologically meaningful

variation and differentiate between less obvious variations in surface structure.

5.2   Realities and practicalities of observing change

Whether using habitat or RS variables, model success depends on the quality and quantity of

the data with which they are trained (97,98). Low detection probabilities suggest substantial

effort is needed to detect most species, raising concerns about data sufficiency. However, of

the approximately 760 bird species thought to occur in the Tambopata area (99), our study

recorded nearly half. This, in terms of both number and proportion of species identified, aligns

with findings from other surveys in tropical forests (100–102). Low detection rates likely

caused individual surveys to underestimate community composition, making landscape-wide

communities appear more distinct than limited survey observations suggest. Nevertheless,

the species modelled exhibited patterns of grouping based on both forest habitat descriptors

and RS variables, suggesting that they represented more than just random observations. This

indicates that, despite describing approximately a half of the full community, our results were

reliable.

Increased sampling effort revealed more of the local community and improved similarity

between predicted and observed communities. Thus, improving detection rates is essential

to increasing the proportion of the community observed and improving the predictive output

of models. Increasing cumulative detection rates to 80% has been shown to negate the

importance of detection on model performance (103). Yet we have shown that it would

require a huge effort to reliably observe changes in occupancy for many species. Thus,

efficient and accurate surveys are critical to provide high quality data on which to train RS

models, especially as ongoing surveys are needed to monitor the effects of conservation

efforts on habitat change over space or time. The type of survey conducted affects detection

rate, and we agree with Mulvaney & Cherry, (2020) that point counts increase mean detection

over net counts. However, these findings are not universal, as it has been shown that within

tropical forests, the most effective survey method varies with ecosystem, such as whether

the study is within lowland or cloud forest (101). Similarly, transect counts and canopy counts

add further depth to the proportions of small-bodied species, canopy and mid-storey species

that are detected (86,102). Due to the relatively low community overlap observed between

methods found in our study, and aligning with other studies, we would suggest that the use

of multiple methods is necessary to improve overall diversity assessment (86,101,102).

Further to the survey methodology lies the challenge of recruiting appropriately trained

surveyors who are familiar with the bird species of the area, which is a necessity in conducting

effective surveys in complex and rich ecosystems such as tropical forests (86). Our results

suggest that to effectively detect changes in community composition, it is necessary to

perform far more replicate surveys than were performed (86).

5.3   Future potential

Forest degradation in Peru has been shown to increase the amount of carbon released into

the atmosphere by almost 50% (105). Protecting forests and carbon sinks has understandably

been central of tropical conservation, and government policies and voluntary carbon offset

conservation efforts such as REDD+ have successfully slowed the rate of deforestation in

recent years and have played a role in mitigating climate change (106). To ensure that the

implementation of carbon conservation efforts, such as REDD+, also have a positive impact

on biodiversity, ecological monitoring of forests should be aligned with the conservation

objectives of the area (107). While REDD+ has had some success in protecting forests globally,

the level of success in the Amazon may be overstated due to the overestimation of baseline

deforestation (108), highlighting the need to validate the costs of voluntary carbon offset

credits. Degradation impacts are not fixed and can be improved by restoration or worsened

by repetition and should be measured at a landscape rather than patch scale (109). Measuring

community diversity at landscape scales may require impractical levels of monitoring, but

patch scale changes in habitat should in principle be observable by Earth observation. Should

sufficient, consistent, reliably representative biodiversity data be available to calibrate

models fitted with RS data, biodiversity responses to changing land use can be identified.

Policy and management should judge the results of practices at the point of implementation

rather than judge the practices conceptually as a whole, and by mapping species diversity in

conjunction with other measures of landscape health, trends in forest degradation and

regeneration over time can be identified (109).

Despite the challenges posed by generally low detection rates, our study demonstrated that

the occupancy of a third of the bird species with sufficient detections in the Peruvian Amazon

can be predicted with reasonable confidence. However, low detection rates limit model

capacity to detect community change over time. While RS models may not yet be sufficiently

robust for ongoing monitoring, reporting and verification, there is no evidence that field-

based methods alone will be able to do this at the scale required. While surveys can identify

spatial variations, detecting temporal change requires detailed count data at a level that may

be impractical (110). Our power analysis revealed that with the current monitoring structure,

even if a species population were to half, we would fail to detect this four out of five times.

The suggestion that declines of many species may be substantial before being reliably

detected has major implications for conservation monitoring using existing survey design.

Improving detection rates will increase the power of future surveys to detect change. Given

that resources are limited, future surveys should focus on fewer locations, prioritised across

an ecological gradient of interest, with a higher number of replicates; results suggest at least

20 per site. This is particularly critical as even basic inventories of avian diversity across much

of the tropics are lacking (86). Establishing baseline datasets is essential for enabling future

repeat surveys, which could provide valuable insights into how and where tropical bird

communities are changing (111). Building on the findings of this study by refining survey

design and aligning research with specific conservation objectives will be key to improving

biodiversity monitoring in the region so that decisions are based on detectable trends rather

than snapshots of biodiversity (112).

Declarations

Funding:

This study received no specific funding. The bird data used were collected independently by co-authors C.A.K. and C.K. as part of ongoing monitoring activities not funded by the lead or senior authors of this manuscript.

Ethics Statement:

The bird occurrence data used in this study were collected independently by co-authors C.A.K. and C.K. as part of long-term monitoring within the Tambopata National Reserve, Peru. Data collection was conducted ethically and in accordance with local research regulations.

Data Availability:

The data that support the findings of this study are available from the authors upon reasonable request.

Authors' Contributions:

Author contributions: A.C.S., A.B. and I.R.H. conceptualised the study; C.A.K and C.K. performed independent field surveys and provided bird sighting data; A.C.S. sourced and calculated all remotely sensed variables, performed all modelling and analysed results; A.C.S. wrote the manuscript with reviews and edits suggested by A.B, I.R.H.

Conflict of Interest: The authors declare no conflicts of interest.

500   **Use of AI tools:** ChatGPT (OpenAI) was used during the writing process for language editing,

501   including grammar, syntax, and clarity. The authors reviewed and approved all edits. No AI

502   was involved in data analysis or content generation.

References

1.   Bradshaw CJ, Sodhi NS, Brook BW. Tropical turmoil: a biodiversity tragedy in progress.

Front Ecol Environ [Internet]. 2009 Mar [cited 2023 Apr 12];7(2):79–87. Available

from: http://doi.wiley.com/10.1890/070193

2.   Gibson L, Lee TM, Koh LP, Brook BW, Gardner TA, Barlow J, et al. Primary forests are

irreplaceable for sustaining tropical biodiversity. Nature [Internet]. 2011 Oct

20;478(7369):378–81. Available from: https://www.nature.com/articles/nature10425

3.   Lewis SL, Edwards DP, Galbraith D. Increasing human dominance of tropical forests.

Science (1979) [Internet]. 2015 Aug 21;349(6250):827–32. Available from:

https://www.science.org/doi/10.1126/science.aaa9932

4.   Pan Y, Birdsey RA, Fang J, Houghton R, Kauppi PE, Kurz WA, et al. A large and

persistent carbon sink in the world's forests. Science (1979). 2011 Aug

19;333(6045):988–93.

5.   Hubau W, Lewis SL, Phillips OL, Affum-Baffoe K, Beeckman H, Cuní-Sanchez A, et al.

Asynchronous carbon sink saturation in African and Amazonian tropical forests.

Nature. 2020 Mar 5;579(7797):80–7.

6.   Potapov P, Hansen MC, Laestadius L, Turubanova S, Yaroshenko A, Thies C, et al. The

last frontiers of wilderness: Tracking loss of intact forest landscapes from 2000 to

2013. Sci Adv [Internet]. 2017 Jan 1 [cited 2023 Jun 13];3(1). Available from:

https://www.science.org/doi/10.1126/sciadv.1600821

7.   Vergara A, Arias M, Gachet B, Naranjo LG, Román L, Surkin J, et al. Living Amazon

Report 2022. 2022.

8.   Lapola DM, Pinho P, Barlow J, Aragão LEOC, Berenguer E, Carmenta R, et al. The

drivers and impacts of Amazon forest degradation. Science (1979) [Internet]. 2023

Jan 27 [cited 2024 Mar 20];379(6630). Available from:

https://www.science.org/doi/10.1126/science.abp8622

9.   UNFCCC. United Nations Framework Convention on Climate Change. 2023 [cited 2023

Nov 8]. Home / REDD+. Available from: https://redd.unfccc.int/

10.   Wunder S, Schulz D, Montoya-Zumaeta JG, Börner J, Ponzoni Frey G, Betancur-

Corredor B. Modest forest and welfare gains from initiatives for reduced emissions

from deforestation and forest degradation. Commun Earth Environ [Internet]. 2024

Jul 22 [cited 2025 Jan 27];5(1):394. Available from:

https://www.nature.com/articles/s43247-024-01541-1

11.   Guizar-Coutiño A, Jones JPG, Balmford A, Carmenta R, Coomes DA. A global

evaluation of the effectiveness of voluntary REDD+ projects at reducing deforestation

and degradation in the moist tropics. Conservation Biology [Internet]. 2022 Dec 8

[cited 2025 Jan 27];36(6):e13970. Available from:

https://conbio.onlinelibrary.wiley.com/doi/10.1111/cobi.13970

12.   Barlow J, Lennox GD, Ferreira J, Berenguer E, Lees AC, Nally R Mac, et al.

Anthropogenic disturbance in tropical forests can double biodiversity loss from

deforestation. Nature [Internet]. 2016 Jul 7 [cited 2021 Jan 13];535(7610):144–7.

Available from: https://www.nature.com/articles/nature18326

13.   Laurance WF, Carolina Useche D, Rendeiro J, Kalka M, Bradshaw CJA, Sloan SP, et al.

Averting biodiversity collapse in tropical forest protected areas. Nature [Internet].

2012 Sep 25;489(7415):290–4. Available from:

http://www.nature.com/articles/nature11318

14.   Hansen MC, Potapov P V., Moore R, Hancher M, Turubanova SA, Tyukavina A, et al.

High-Resolution Global Maps of 21st-Century Forest Cover Change. Science (1979)

[Internet]. 2013 Nov 15 [cited 2023 Jan 23];342(6160):850–3. Available from:

https://www.science.org/doi/10.1126/science.1244693

15.   Stork NE, Boyle TJB, Dale V, Eeley H, Finegan B, Lawes M, et al. Criteria and indicators

for assessing the sustainability of forest management: conservation of biodiversity

[Internet]. Criteria and indicators for assessing the sustainability of forest

management: conservation of biodiversity. Center for International Forestry Research

(CIFOR); 1997 [cited 2025 Jan 27]. Available from:

http://www.cifor.org/library/77/criteria-and-indicators-for-assessing-the-

sustainability-of-forest-management-conservation-of-biodiversity/

16.   Titeux N, Henle K, Mihoub J, Regos A, Geijzendorffer IR, Cramer W, et al. Biodiversity

scenarios neglect future land-use changes. Glob Chang Biol [Internet]. 2016 Jul

31;22(7):2505–15. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/gcb.13272

17.   Willis KJ, Araújo MB, Bennett KD, Figueroa-Rangel B, Froyd CA, Myers N. How can a

knowledge of the past help to conserve the future? Biodiversity conservation and the

relevance of long-term ecological studies. Philosophical Transactions of the Royal

Society B: Biological Sciences [Internet]. 2007 Feb 28 [cited 2025 Jan

27];362(1478):175–87. Available from:

https://royalsocietypublishing.org/doi/10.1098/rstb.2006.1977

18.   Pettorelli N, Schulte to Bühne H. Current and future opportunities for satellite remote

sensing to inform rewilding. Disney M, Hernandez-Clemente R, editors. Remote Sens

Ecol Conserv [Internet]. 2022 Dec 16; Available from:

https://zslpublications.onlinelibrary.wiley.com/doi/10.1002/rse2.321

19.   Chazdon RL, Brancalion PHS, Laestadius L, Bennett-Curry A, Buckingham K, Kumar C,

et al. When is a forest a forest? Forest concepts and definitions in the era of forest

and landscape restoration. Ambio [Internet]. 2016 Sep 9 [cited 2025 Jan

27];45(5):538–50. Available from: http://link.springer.com/10.1007/s13280-016-

0772-y

20.   Savilaakso S, Lausberg N, Waeber PO, Hillgén O, Isotalo A, Kleinschroth F, et al.

Whose perspective counts? A critical look at definitions of terms used for natural and

near-natural forests. One Earth [Internet]. 2023 Nov [cited 2025 Jan 27];6(11):1477–

93. Available from: https://linkinghub.elsevier.com/retrieve/pii/S2590332223004530

21.   Pereira HM, Ferrier S, Walters M, Geller GN, Jongman RHG, Scholes RJ, et al. Essential

Biodiversity Variables. Science (1979) [Internet]. 2013 Jan 18;339(6117):277–8.

Available from: https://www.science.org/doi/10.1126/science.1229931

22.   GEO BON. What are EBVs. 2018 [cited 2022 Jul 22]. What are EBVs? – GEO BON.

Available from: https://geobon.org/ebvs/what-are-ebvs/

23.   Purdon A, Mole MA, Selier J, Kruger J, Mafumo H, Olivier PI. Using the Rao's Q

diversity index as an indicator of protected area effectiveness in conserving

biodiversity. Ecol Inform [Internet]. 2022 Dec [cited 2022 Dec 2];72:101920. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S1574954122003703

24.   Pettorelli N, Vik JO, Mysterud A, Gaillard JM, Tucker CJ, Stenseth NChr. Using the

594   satellite-derived NDVI to assess ecological responses to environmental change.

595   Trends Ecol Evol [Internet]. 2005 Sep 1 [cited 2025 Jan 28];20(9):503–10. Available

596   from: https://linkinghub.elsevier.com/retrieve/pii/S016953470500162X

597   25.   Asner GP, Kellner JR, Kennedy-Bowdoin T, Knapp DE, Anderson C, Martin RE. Forest

598   Canopy Gap Distributions in the Southern Peruvian Amazon. Chen HYH, editor. PLoS

599   One [Internet]. 2013 Apr 15 [cited 2025 Jan 28];8(4):e60875. Available from:

600   https://dx.plos.org/10.1371/journal.pone.0060875

601   26.   Zhang X, Friedl MA, Schaaf CB, Strahler AH, Hodges JCF, Gao F, et al. Monitoring

602   vegetation phenology using MODIS. Remote Sens Environ [Internet]. 2003 Mar 1

603   [cited 2025 Jan 28];84(3):471–5. Available from:

604   https://linkinghub.elsevier.com/retrieve/pii/S0034425702001359

605   27.   Perrone M, Di Febbraro M, Conti L, Divíšek J, Chytrý M, Keil P, et al. The relationship

606   between spectral and plant diversity: Disentangling the influence of metrics and

607   habitat types at the landscape scale. Remote Sens Environ [Internet]. 2023

608   Aug;293:113591. Available from:

609   https://linkinghub.elsevier.com/retrieve/pii/S0034425723001426

610   28.   Turner W, Spector S, Gardiner N, Fladeland M, Sterling E, Steininger M. Remote

611   sensing for biodiversity science and conservation. Trends Ecol Evol [Internet]. 2003

612   Jun 1 [cited 2025 Jan 28];18(6):306–14. Available from:

613   https://linkinghub.elsevier.com/retrieve/pii/S0169534703000703

614   29.   Ludwig A, Meyer H, Nauss T. Automatic classification of Google Earth images for a

615   larger scale monitoring of bush encroachment in South Africa. International Journal of

616   Applied Earth Observation and Geoinformation [Internet]. 2016 Aug 1 [cited 2025 Jan

28];50:89–94. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0303243416300265

30.   Pettorelli N, Safi K, Turner W. Satellite remote sensing, biodiversity research and

conservation of the future. Philosophical Transactions of the Royal Society B:

Biological Sciences [Internet]. 2014 May 26 [cited 2020 Nov 13];369(1643):20130190.

Available from: https://royalsocietypublishing.org/doi/10.1098/rstb.2013.0190

31.   Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJB, Collen B. Defaunation in the

Anthropocene. Science (1979) [Internet]. 2014 Jul 25 [cited 2025 Jan

29];345(6195):401–6. Available from:

https://www.science.org/doi/10.1126/science.1251817

32.   Gerlach J, Samways M, Pryke J. Terrestrial invertebrates as bioindicators: an overview

of available taxonomic groups. J Insect Conserv [Internet]. 2013 Aug 7;17(4):831–50.

Available from: http://link.springer.com/10.1007/s10841-013-9565-9

33.   França FM, Frazão FS, Korasaki V, Louzada J, Barlow J. Identifying thresholds of

logging intensity on dung beetle communities to improve the sustainable

management of Amazonian tropical forests. Biol Conserv [Internet].

2017;216(September):115–22. Available from:

http://dx.doi.org/10.1016/j.biocon.2017.10.014

34.   Magioli M, Ferraz KMPM de B, Chiarello AG, Galetti M, Setz EZF, Paglia AP, et al. Land-

use changes lead to functional loss of terrestrial mammals in a Neotropical rainforest.

Perspect Ecol Conserv [Internet]. 2021 Apr;19(2):161–70. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S2530064421000213

35.   Basset Y, Cizek L, Cuénoud P, Didham RK, Guilhaumon F, Missa O, et al. Arthropod

Diversity in a Tropical Forest. Science (1979) [Internet]. 2012 Dec 14 [cited 2023 Apr

26];338(6113):1481–4. Available from:

https://www.science.org/doi/10.1126/science.1226727

36. Traylor CR, Ulyshen MD, Wallace D, Loudermilk EL, Ross CW, Hawley C, et al.

Compositional attributes of invaded forests drive the diversity of insect functional

groups. Glob Ecol Conserv. 2022 Jun 1;35:e02092.

37. Schowalter T. Arthropod Diversity and Functional Importance in Old-Growth Forests

of North America. Forests [Internet]. 2017 Mar 25;8(4):97. Available from:

http://www.mdpi.com/1999-4907/8/4/97

38. Zhang K, Lin S, Ji Y, Yang C, Wang X, Yang C, et al. Plant diversity accurately predicts

insect diversity in two tropical landscapes [Internet]. Vol. 25, Molecular ecology.

2016. p. 4407–19. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/mec.13770

39. Barros F de C, Almeida SM, Godoy BS, Silva RR da, Silva LC, de Moraes KF, et al.

Taxonomic and functional diversity of bird communities in mining areas undergoing

passive and active restoration in eastern Amazon. Ecol Eng [Internet]. 2022

Sep;182:106721. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0925857422001823

40. Bregman TP, Lees AC, Seddon N, MacGregor HEA, Darski B, Aleixo A, et al. Species

interactions regulate the collapse of biodiversity and ecosystem function in tropical

forest fragments. Ecology [Internet]. 2015 Oct;96(10):2692–704. Available from:

https://esajournals.onlinelibrary.wiley.com/doi/10.1890/14-1731.1

41. Skidmore AK, Pettorelli N, Coops NC, Geller GN, Hansen M, Lucas R, et al.

Environmental science: Agree on biodiversity metrics to track from space. Nature

[Internet]. 2015 Jul 23 [cited 2020 Nov 17];523(7561):403–5. Available from:

https://www.nature.com/articles/523403a

42.   Rocchini D, Boyd DS, Féret J, Foody GM, He KS, Lausch A, et al. Satellite remote

sensing to monitor species diversity: potential and pitfalls. Skidmore A, Chauvenet A,

editors. Remote Sens Ecol Conserv [Internet]. 2016 Feb 10 [cited 2022 Jun

21];2(1):25–36. Available from:

https://zslpublications.onlinelibrary.wiley.com/doi/10.1002/rse2.9

43.   Csillik O, Kumar P, Mascaro J, O'Shea T, Asner GP. Monitoring tropical forest carbon

stocks and emissions using Planet satellite data. Sci Rep [Internet]. 2019 Dec

28;9(1):17831. Available from: http://www.nature.com/articles/s41598-019-54386-6

44.   J van Proosdij AS, M Sosef MS, Wieringa JJ, Raes S J van Proosdij andrevanproosdij

NA, M Sosef MS, Wieringa JJ. Minimum required number of specimen records to

develop accurate species distribution models. Ecography [Internet]. 2016;39:542–52.

Available from: https://onlinelibrary.wiley.com/doi/10.1111/ecog.01509

45.   MapBiomas. https://peru.mapbiomas.org/en/terminos-de-uso/. 2023 [cited 2023

Aug 21]. MapBiomas Peru. Available from:

https://peru.mapbiomas.org/en/colecciones-de-mapbiomas-peru/

46.   Carvalho WD, Fluck IE, de Castro IJ, Hilário RR, Martins ACM, de Toledo JJ, et al.

Elevation drives taxonomic, functional and phylogenetic β-diversity of phyllostomid

bats in the Amazon biome. J Biogeogr [Internet]. 2023 Jan 24 [cited 2025 Feb

3];50(1):70–85. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/jbi.14533

47.   Nores M. Species Richness in the Amazonian Bird Fauna from an Evolutionary

Perspective. Emu - Austral Ornithology [Internet]. 2000 Dec 22 [cited 2025 Feb

3];100(5):419–30. Available from:

https://www.tandfonline.com/doi/full/10.1071/MU0007S

48. Huete AR. Vegetation Indices, Remote Sensing and Forest Monitoring. Geogr

Compass [Internet]. 2012 Sep 20 [cited 2023 Sep 7];6(9):513–32. Available from:

https://compass.onlinelibrary.wiley.com/doi/10.1111/j.1749-8198.2012.00507.x

49. Ollinger S V. Sources of variability in canopy reflectance and the convergent

properties of plants. New Phytologist [Internet]. 2011 Jan 16 [cited 2024 Feb

29];189(2):375–94. Available from:

https://nph.onlinelibrary.wiley.com/doi/10.1111/j.1469-8137.2010.03536.x

50. Wang R, Gamon JA, Schweiger AK, Cavender-Bares J, Townsend PA, Zygielbaum AI, et

al. Influence of species richness, evenness, and composition on optical diversity: A

simulation study. Remote Sens Environ [Internet]. 2018 Jun [cited 2022 Sep

29];211:218–28. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S003442571830155X

51. Quiñones M, Vissers M, Hoekman D. K&C Science Report - Phase 1 Tropical Forest

and Wetlands Mapping, Case Study Borneo. 2007 Mar.

52. Morton DC, DeFries RS, Shimabukuro YE, Anderson LO, Del Bon Espírito-Santo F,

Hansen M, et al. Rapid Assessment of Annual Deforestation in the Brazilian Amazon

Using MODIS Data. Earth Interact [Internet]. 2005 Jun 1 [cited 2025 Jan 30];9(8):1–22.

Available from: https://journals.ametsoc.org/doi/10.1175/EI139.1

53. Hilker T, Lyapustin AI, Tucker CJ, Hall FG, Myneni RB, Wang Y, et al. Vegetation

dynamics and rainfall sensitivity of the Amazon. Proceedings of the National Academy

of Sciences [Internet]. 2014 Nov 11 [cited 2025 Jan 30];111(45):16041–6. Available

from: https://pnas.org/doi/full/10.1073/pnas.1404870111

54.   Samarawickrama U, Piyaratne D, Ranagalage M. Relationship between NDVI with

Tasseled cap Indices: A Remote Sensing based Analysis. 2017;3.

55.   Remsen, J V., Good DA. Misuse of Data from Mist-Net Captures to Assess Relative

Abundance in Bird Populations. Auk [Internet]. 1996 Apr;113(2):381–98. Available

from: https://academic.oup.com/auk/article/113/2/381-398/5168448

56.   Rosenstock SS, Anderson DR, Giesen KM, Leukering T, Carter MF. Landbird Counting

Techniques: Current Practices and an Alternative. Thompson III F, editor. Auk

[Internet]. 2002 Jan 1 [cited 2025 Jan 31];119(1):46–53. Available from:

https://academic.oup.com/auk/article/119/1/46/5561821

57.   Simons TR, Alldredge MW, Pollock KH, Wettroth JM. Experimental Analysis of The

Auditory Detection Process on Avian Point Counts. Dufty AM, editor. Auk [Internet].

2007 Jul 1;124(3):986–99. Available from:

https://academic.oup.com/auk/article/124/3/986/5562805

58.   Kery M. ESTIMATING ABUNDANCE FROM BIRD COUNTS: BINOMIAL MIXTURE

MODELS UNCOVER COMPLEX COVARIATE RELATIONSHIPS. Auk [Internet]. 2008

Apr;125(2):336–45. Available from:

https://academic.oup.com/auk/article/125/2/336-345/5148162

59.   MacKenzie DI, Nichols JD, Hines JE, Knutson MG, Franklin AB. ESTIMATING SITE

OCCUPANCY, COLONIZATION, AND LOCAL EXTINCTION WHEN A SPECIES IS DETECTED

IMPERFECTLY. Ecology [Internet]. 2003 Aug [cited 2025 Jan 31];84(8):2200–7.

Available from: http://doi.wiley.com/10.1890/02-3090

60.   Guillera-Arroita G, Lahoz-Monfort JJ. Designing studies to detect differences in

species occupancy: power analysis under imperfect detection. Methods Ecol Evol

[Internet]. 2012 Oct;3(5):860–9. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/j.2041-210X.2012.00225.x

61. Doser JW, Finley AO, Kéry M, Zipkin EF. spOccupancy: An R package for single-species,

multi-species, and integrated spatial occupancy models. Methods Ecol Evol. 2022 Aug

30;13(8):1670–8.

62. Doser J. https://www.jeffdoser.com/files/spoccupancy-

web/articles/modelconsiderations. 2023. Convergence diagnostics and other

considerations when fitting spatial occupancy models.

63. Hooten MB, Hobbs NT. A guide to Bayesian model selection for ecologists. Ecol

Monogr [Internet]. 2015 Feb 1 [cited 2024 Mar 12];85(1):3–28. Available from:

https://esajournals.onlinelibrary.wiley.com/doi/10.1890/14-0661.1

64. Kéry M, Royle JA. Chapter 6 - Modeling Abundance with Counts of Unmarked

Individuals in Closed Populations: Binomial N-mixture Models. In: Kéry M, Royle JA,

editors. Applied Hierarchical Modeling in Ecology [Internet]. Boston: Academic Press;

2016. p. 219–312. Available from:

https://www.sciencedirect.com/science/article/pii/B9780128013786000060

65. Watanabe S. Asymptotic Equivalence of Bayes Cross Validation and Widely Applicable

Information Criterion in Singular Learning Theory. Journal of Machine Learning

Research. 2010;11:3571–94.

66. Gould MJ, Gould WR, Cain JW, Roemer GW. Validating the performance of occupancy

models for estimating habitat use and predicting the distribution of highly-mobile

species: A case study using the American black bear. Biol Conserv [Internet]. 2019

Jun;234:28–36. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S000632071831214X

67.   Mandrekar JN. Receiver Operating Characteristic Curve in Diagnostic Test

Assessment. Journal of Thoracic Oncology [Internet]. 2010 Sep [cited 2023 Apr

25];5(9):1315–6. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S1556086415306043

68.   Dvořáková D, Šipoš J, Suchomel J. Impact of agricultural landscape structure on the

patterns of bird species diversity at a regional scale. Avian Res [Internet].

2023;14:100147. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S2053716623000737

69.   Tu HM, Fan MW, Ko JCJ. Different Habitat Types Affect Bird Richness and Evenness.

Sci Rep [Internet]. 2020 Jan 27;10(1):1221. Available from:

https://www.nature.com/articles/s41598-020-58202-4

70.   Terraube J, Archaux F, Deconchat M, van Halder I, Jactel H, Barbaro L. Forest edges

have high conservation value for bird communities in mosaic landscapes. Ecol Evol

[Internet]. 2016 Aug 28;6(15):5178–89. Available from:

https://onlinelibrary.wiley.com/doi/10.1002/ece3.2273

71.   Willmer JNG, Püttker T, Prevedello JA. Global impacts of edge effects on species

richness. Biol Conserv [Internet]. 2022 Aug [cited 2023 Dec 14];272:109654. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S0006320722002075

72.   Stouffer PC, Johnson EI, Bierregaard RO, Lovejoy TE. Understory Bird Communities in

Amazonian Rainforest Fragments: Species Turnover through 25 Years Post-Isolation in

Recovering Landscapes. Evans DM, editor. PLoS One [Internet]. 2011 Jun

22;6(6):e20543. Available from: https://dx.plos.org/10.1371/journal.pone.0020543

73.   Stratford JA, Stouffer PC. Forest fragmentation alters microhabitat availability for

Neotropical terrestrial insectivorous birds. Biol Conserv [Internet]. 2015 Aug [cited

2025 Feb 4];188:109–15. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0006320715000385

74. Şekercioğlu ÇH, Primack RB, Wormworth J. The effects of climate change on tropical

birds. Biol Conserv [Internet]. 2012 Apr [cited 2025 Feb 4];148(1):1–18. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S0006320711003880

75. Barlow J, Peres CA, Henriques LMP, Stouffer PC, Wunderle JM. The responses of

understorey birds to forest fragmentation, logging and wildfires: An Amazonian

synthesis. Biol Conserv [Internet]. 2006 Mar [cited 2025 Feb 4];128(2):182–92.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S0006320705003897

76. Barlow J, Mestre LAM, Gardner TA, Peres CA. The value of primary, secondary and

plantation forests for Amazonian birds. Biol Conserv [Internet]. 2007 Apr [cited 2025

Feb 4];136(2):212–31. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0006320706005271

77. Sekercioglu CH. Bird functional diversity and ecosystem services in tropical forests,

agroforests and agricultural areas. J Ornithol [Internet]. 2012 Aug 27;153(S1):153–61.

Available from: http://link.springer.com/10.1007/s10336-012-0869-4

78. Peh KS -H., Sodhi NS, De Jong J, Sekercioglu CH, Yap CA -M., Lim SL -H. Conservation

value of degraded habitats for forest birds in southern Peninsular Malaysia. Divers

Distrib [Internet]. 2006 Sep 26 [cited 2025 Feb 4];12(5):572–81. Available from:

https://onlinelibrary.wiley.com/doi/10.1111/j.1366-9516.2006.00257.x

79. Rutt CL, Jirinec V, Cohn-Haft M, Laurance WF, Stouffer PC. Avian ecological succession

in the Amazon: A long-term case study following experimental deforestation. Ecol

Evol [Internet]. 2019 Dec 27 [cited 2025 Feb 5];9(24):13850–61. Available from:

https://onlinelibrary.wiley.com/doi/10.1002/ece3.5822

80. Thornton DH, Branch LC, Sunquist ME. The influence of landscape, patch, and within-

patch factors on species presence and abundance: a review of focal patch studies.

Landsc Ecol [Internet]. 2011 Jan 30;26(1):7–18. Available from:

http://link.springer.com/10.1007/s10980-010-9549-z

81. Laurance WF, Camargo JLC, Luizão RCC, Laurance SG, Pimm SL, Bruna EM, et al. The

fate of Amazonian forest fragments: A 32-year investigation. Biol Conserv [Internet].

2011 Jan [cited 2025 Feb 13];144(1):56–67. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0006320710004209

82. Hess LL, Melack JM, Affonso AG, Barbosa C, Gastil-Buhl M, Novo EMLM. Wetlands of

the Lowland Amazon Basin: Extent, Vegetative Cover, and Dual-season Inundated

Area as Mapped with JERS-1 Synthetic Aperture Radar. Wetlands [Internet]. 2015 Aug

20;35(4):745–56. Available from: https://link.springer.com/10.1007/s13157-015-

0666-y

83. Laurance WF, Albernaz AKM, Costa C Da. Is deforestation accelerating in the Brazilian

Amazon? Environ Conserv [Internet]. 2001 Dec 10 [cited 2025 Feb 13];28(4):305–11.

Available from:

https://www.cambridge.org/core/product/identifier/S0376892901000339/type/jour

nal_article

84. Blake JG, Loiselle BA. Enigmatic declines in bird numbers in lowland forest of eastern

Ecuador may be a consequence of climate change. PeerJ [Internet]. 2015 Aug 11

[cited 2025 Feb 12];3(8):e1177. Available from: https://peerj.com/articles/1177

85. Rutt CL, Stouffer PC. Seasonal dynamics of flock interaction networks across a

human-modified landscape in lowland Amazonian rain forest. Ecological Applications

[Internet]. 2021 Mar 21 [cited 2025 Feb 12];31(2). Available from:

https://esajournals.onlinelibrary.wiley.com/doi/10.1002/eap.2235

86. Robinson WD, Curtis JR. Creating benchmark measurements of tropical forest bird

communities in large plots. Condor [Internet]. 2020 Sep 11 [cited 2025 Feb

7];122(3):1–15. Available from:

https://academic.oup.com/condor/article/doi/10.1093/condor/duaa015/5820537

87. Peña-Lara VA, Dupuy JM, Reyes-Garcia C, Sanaphre-Villanueva L, Portillo-Quintero CA,

Hernández-Stefanoni JL. Modelling Species Richness and Functional Diversity in

Tropical Dry Forests Using Multispectral Remotely Sensed and Topographic Data.

Remote Sens (Basel) [Internet]. 2022 Nov 23 [cited 2025 Feb 5];14(23):5919.

Available from: https://www.mdpi.com/2072-4292/14/23/5919

88. Suttidate N, Hobi ML, Pidgeon AM, Round PD, Coops NC, Helmers DP, et al. Tropical

bird species richness is strongly associated with patterns of primary productivity

captured by the Dynamic Habitat Indices. Remote Sens Environ [Internet]. 2019 Oct

[cited 2025 Feb 5];232:111306. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0034425719303256

89. Sheeren D, Bonthoux S, Balent G. Modeling bird communities using unclassified

remote sensing imagery: Effects of the spatial resolution and data period. Ecol Indic

[Internet]. 2014 Aug [cited 2023 Dec 1];43:69–82. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S1470160X14000818

90. Regos A, Gómez-Rodríguez P, Arenas-Castro S, Tapia L, Vidal M, Domínguez J. Model-

Assisted Bird Monitoring Based on Remotely Sensed Ecosystem Functioning and Atlas

Data. Remote Sens (Basel) [Internet]. 2020 Aug 7;12(16):2549. Available from:

https://www.mdpi.com/2072-4292/12/16/2549

91.   Cruz M, Pradel W, Juarez H, Hualla V, Suarez V. Deforestation Dynamics in Peru. A

Comprehensive Review of Land Use, Food Systems, and Socio-Economic Drivers

[Internet]. 2023 Dec [cited 2025 Feb 6]. Available from:

https://hdl.handle.net/10568/135509

92.   Marcus M, Gutierrez-Velez VH, Cronkleton P. Land use change in four landscapes in

the Peruvian Amazon [Internet]. Land use change in four landscapes in the Peruvian

Amazon. Center for International Forestry Research (CIFOR); 2020 [cited 2025 Feb 6].

Available from: https://www.cifor.org/knowledge/publication/7736

93.   Gallardo-Cruz JA, Solórzano J V., González EJ, Meave JA. The Effect of Spatial Scale on

the Prediction of Tropical Forest Attributes from Image Texture. Źróbek-Sokolnik A,

editor. International Journal of Forestry Research [Internet]. 2024 Apr 24 [cited 2025

Feb 6];2024(1):1–10. Available from:

https://www.hindawi.com/journals/ijfr/2024/7178211/

94.   Ploton P, Mortier F, Réjou-Méchain M, Barbier N, Picard N, Rossi V, et al. Spatial

validation reveals poor predictive performance of large-scale ecological mapping

models. Nat Commun [Internet]. 2020 Sep 11 [cited 2025 Feb 6];11(1):4540. Available

from: https://www.nature.com/articles/s41467-020-18321-y

95.   Morrison LW. Observer error in vegetation surveys: a review. Journal of Plant Ecology

[Internet]. 2016 Aug 1 [cited 2025 Feb 6];9(4):367–79. Available from:

https://academic.oup.com/jpe/article-lookup/doi/10.1093/jpe/rtv077

96.   Basile M, Storch I, Mikusiński G. Abundance, species richness and diversity of forest

bird assemblages – The relative importance of habitat structures and landscape

context. Ecol Indic [Internet]. 2021 Dec [cited 2023 Dec 14];133:108402. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S1470160X21010670

97.   Aubry KB, Raley CM, McKelvey KS. The importance of data quality for generating

879   reliable distribution models for rare, elusive, and cryptic species. Baldwin RF, editor.

880   PLoS One [Internet]. 2017 Jun 22;12(6):e0179152. Available from:

881   https://dx.plos.org/10.1371/journal.pone.0179152

882   98.   McCord SE, Webb NP, Van Zee JW, Burnett SH, Christensen EM, Courtright EM, et al.

883   Provoking a Cultural Shift in Data Quality. Bioscience [Internet]. 2021 Jun

884   1;71(6):647–57. Available from:

885   https://academic.oup.com/bioscience/article/71/6/647/6188799

886   99.   Lepage D. Tambopata bird checklist - Avibase - Bird Checklists of the World [Internet].

887   2024 [cited 2025 Feb 10]. Available from: https://avibase.bsc-

888   eoc.org/checklist.jsp?lang=EN&p2=1&list=howardmoore&synlang=®ion=PEmdtp

889   &version=text&lifelist=&highlight=0

890   100.   Alvarez-Berríos N, Campos-Cerqueira M, Hernández-Serna A, Amanda Delgado CJ,

891   Román-Dañobeytia F, Aide TM. Impacts of Small-Scale Gold Mining on Birds and

892   Anurans Near the Tambopata Natural Reserve, Peru, Assessed Using Passive Acoustic

893   Monitoring. Trop Conserv Sci [Internet]. 2016 Jun 1 [cited 2025 Feb 10];9(2):832–51.

894   Available from: https://journals.sagepub.com/doi/10.1177/194008291600900216

895   101.   Martin TE, Nightingale J, Baddams J, Monkhouse J, Kaban A, Sastranegara H, et al.

896   Variability in the Effectiveness of Two Ornithological Survey Methods between

897   Tropical Forest Ecosystems. Pandit MK, editor. PLoS One [Internet]. 2017 Jan 10 [cited

898   2025 Feb 7];12(1):e0169786. Available from:

899   https://dx.plos.org/10.1371/journal.pone.0169786

900   102.   Anderson DL. Ground Versus Canopy Methods for the Study of Birds in Tropical Forest

901   Canopies: Implications for Ecology and Conservation. Condor [Internet]. 2009 May 1

[cited 2025 Feb 7];111(2):226–37. Available from:

https://academic.oup.com/condor/article/111/2/226-237/5152440

103.   Steenweg R, Hebblewhite M, Whittington J, McKelvey K. Species-specific differences

in detection and occupancy probabilities help drive ability to detect trends in

occupancy. Ecosphere [Internet]. 2019 Apr 3 [cited 2023 Oct 9];10(4). Available from:

https://esajournals.onlinelibrary.wiley.com/doi/10.1002/ecs2.2639

104.   Mulvaney JM, Cherry MI. The effectiveness of point counts and mist-netting in

surveying Afrotemperate forest bird community structure in South Africa. Emu -

Austral Ornithology [Internet]. 2020 Apr 2;120(2):103–13. Available from:

https://www.tandfonline.com/doi/full/10.1080/01584197.2020.1726186

105.   Asner GP, Powell GVN, Mascaro J, Knapp DE, Clark JK, Jacobson J, et al. High-

resolution forest carbon stocks and emissions in the Amazon. Proceedings of the

National Academy of Sciences [Internet]. 2010 Sep 21;107(38):16738–42. Available

from: https://pnas.org/doi/full/10.1073/pnas.1004875107

106.   Smith P, Bustamante M, Ahammad H, Clark H, Dong H, Elsiddig EA. Agriculture,

Forestry and Other Land Use (AFOLU). Climate Change 2014: Mitigation of Climate

Change. 2014;811–922.

107.   Harrison ME, Boonman A, Cheyne SM, Husson SJ, Marchant NC, Struebig MJ.

Biodiversity Monitoring Protocols for REDD+: Can a One-Size-Fits-All Approach Really

Work? Trop Conserv Sci [Internet]. 2012 Mar 1 [cited 2024 Apr 24];5(1):1–11.

Available from: http://journals.sagepub.com/doi/10.1177/194008291200500102

108.   West TAP, Börner J, Sills EO, Kontoleon A. Overstated carbon emission reductions

from voluntary REDD+ projects in the Brazilian Amazon. Proceedings of the National

Academy of Sciences [Internet]. 2020 Sep 29 [cited 2024 Apr 24];117(39):24188–94.

Available from: https://pnas.org/doi/full/10.1073/pnas.2004334117

109. Betts MG, Yang Z, Hadley AS, Hightower J, Hua F, Lindenmayer D, et al. Quantifying

forest degradation requires a long-term, landscape-scale approach. Nat Ecol Evol

[Internet]. 2024 Apr 12; Available from: https://www.nature.com/articles/s41559-

024-02409-5

110. Stephenson P. Technological advances in biodiversity monitoring: applicability,

opportunities and challenges. Curr Opin Environ Sustain [Internet]. 2020 Aug;45:36–

41. Available from: https://linkinghub.elsevier.com/retrieve/pii/S1877343520300592

111. Magurran AE, Baillie SR, Buckland ST, Dick JMcP, Elston DA, Scott EM, et al. Long-term

datasets in biodiversity research and monitoring: assessing change in ecological

communities through time. Trends Ecol Evol [Internet]. 2010 Oct 1 [cited 2024 Jun

7];25(10):574–82. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0169534710001552

112. Nichols J, Williams B. Monitoring for conservation. Trends Ecol Evol [Internet]. 2006

Dec;21(12):668–73. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0169534706002680

949 Table 1: The fit and predictive performance of the four models analysed. Low WAIC values and high

950 AUC values represent better performances. It can be seen that the best performing model in all

951 categories is fitted with RS reflectance data and includes spatial factors.

Model Covariates	Spatial Factor Included	WAIC value	Delta WAIC	Mean Explanatory AUC	Mean Predictive AUC	Number of high-AUC species
RS Reflectance	yes	66773	0	0.88	0.68	49
RS Reflectance	no	68620	1847	0.87	0.62	31
Habitat Descriptors	yes	69432	2659	0.87	0.58	21
None	yes	66995	222	0.8	0.66	43

Survey Sites

Figure 1: (a) The location of survey stations within the Madre de Dios region. Inset is Peru with Madre

de Dios highlighted in blue. Panel (b) displays the distribution of survey stations within the grey box

highlighted in panel (a) along the Tambopata River within regional topography (113).

Figure 2: The distribution of variables used in EO models, comprising the five EO derived principal

components, and the proportion of deforestation calculated at the three measured areas.

Figure 3: The first two principal components, representing 76% of the variation in the RS data, plotted

against each other and showing their relationship with the measured habitat types. Each point

represents an individual survey and is coloured based on the combination of habitat (floodplain forest,

terra-firma forest, agriculture) and forest type (primary or secondary). Ellipses represent the 95%

confidence interval around the mean for each categorical group.

Figure 4: Observed (blue) and modelled (black) species accumulation (±2SD) with increasing numbers

of surveyed stations, (A) within the training data, (B) set aside for validation, and (C) based on an

increasing number of replicate surveys conducted at a single survey station.

Figure 5: Each point represents a survey station. For each station, modelled dissimilarity is the mean

dissimilarity between each of the communities produced by the model's 500 posterior samples, and

the observed community.

1009 Figure 6: The influence of survey effort on the probability of detecting a reduction in occupancy in

1010 future surveys. Survey effort was calculated as the number of stations surveyed * number of replicate

1011 surveys per station. Each line represents a different number of stations and was calculated as the

1012 mean value across all 135 species modelled. Each chart represents a different proportional reduction

1013 in occupancy.