1   **The minimal endosymbiosis between *Spurilla braziliana* (Aeolidiidae, Nudibranchia) and**
2   **Symbiodiniaceae**
3

4   Hideaki Mizobata[1]*, Kenji Tomita[2], Ryo Yonezawa[1,3], Kentaro Hayashi[1], Kazutoshi Yoshitake[1,4],
5   Shigeharu Kinoshita[1], Shuichi Asakawa[1]*
6
1. Department of Aquatic Bioscience, Graduate School of Agricultural and Life Sciences, the
University of Tokyo, Bunkyo-ku, Tokyo, 113-8657 Japan
2. Technology Advancement Center, Graduate School of Agricultural and Life Sciences, the
University of Tokyo, Bunkyo-ku, Tokyo, 113-8657 Japan
3. College of Bioresource Sciences, Nihon University, Fujisawa, Kanagawa, 252-0880 Japan
4. School of Marine Biosciences, Kitasato University, Minami-ku, Sagamihara-shi, Kanagawa,
252-0374 Japan
14
15   *Corresponding author
16   Hideaki Mizobata: mizobata.hideaki@gmail.com
17   Shuichi Asakawa: asakawa@g.ecc.u-tokyo.ac.jp

18
Abstract

Symbiotic relationships between dinoflagellates of the family Symbiodiniaceae and marine
invertebrates underpin the functioning of marine ecosystems. Although the Symbiodiniaceae-
Cladobranchia nudibranch association has been proposed as a promising model for symbiosis
research, interspecific variation in the extent of this association remains poorly resolved. Here, we
assessed the algal-symbiotic capacity of the nudibranch *Spurilla braziliana*. Histological analyses
revealed only limited branching of the digestive gland that accommodates algal cells. Metagenomic
profiling further showed a complete absence of *Endozoicomonas*—bacteria typically linked to
Symbiodiniaceae symbioses—in this species. These findings indicate that *S. braziliana* can harbor
Symbiodiniaceae but exhibits only minimal morphological and microbial adaptations to the
symbiotic state. Consequently, *S. braziliana* may bridge the evolutionary gap between symbiotic and
non-symbiotic nudibranchs, providing a valuable reference point for studies on the evolution of
animal–algal symbioses.

32
33   **Introduction**
34   Dinoflagellates of the family Symbiodiniaceae establish symbiotic partnerships with a wide range
35   of marine invertebrates, including cnidarians[1], molluscs[2], and foraminiferans[3]. The hosts provide the
36   algae with inorganic nutrients[1] and a light environment conducive to photosynthesis[4], while the algae

reciprocate with photosynthetically derived organic compounds[1]. In the best-studied cnidarian hosts, this association dates back at least 385 million years[5], and Symbiodiniaceae supply not only most of the metabolic substrates required for host energy production[6] but also the amino acids essential for somatic growth[7]. Consequently, Symbiodiniaceae–host symbioses underpin the functioning of shallow-water marine ecosystems. In recent decades, however, the breakdown of this relationship— most conspicuously manifested as mass bleaching in coral hosts—has become increasingly evident[8], highlighting the urgent need to clarify the universal mechanisms that govern Symbiodiniaceae– marine invertebrate symbioses.

Cladobranch nudibranchs are considered promising host models for investigating Symbiodiniaceae symbiosis[9]. Their relatively large body size and well-developed anatomical features make them easy to observe, and importantly, the degree of symbiosis varies markedly among species. Some species are capable of maintaining healthy Symbiodiniaceae cells intracellularly for over 200 days[10], while others retain the algae for only a few weeks or do not internalize them at all[11,12]. By documenting host traits histologically in closely related species and conducting genetic and ecological comparisons, it may be possible to identify the mechanisms driving symbiosis with greater precision. However, the diversity of algal symbiotic capacity in nudibranchs has been described only in a limited number of species. Although the genera *Berghia*, *Spurilla*, and *Baeolidia*—closely related to the widely used model species *Berghia stephanieae*[12–15]—have been relatively well studied[11,16–23], many species within these groups remain unexamined, continuing to hinder broader comparative research.

Here, we provide the first detailed description of the Symbiodiniaceae symbiotic state in the nudibranch *Spurilla braziliana*. Through comprehensive histological and metagenomic analyses, we show that while *S. braziliana* possesses the capacity to host Symbiodiniaceae, this ability appears comparatively limited relative to its close relatives. This study not only offers new data on an understudied species but also highlights the potential of *S. braziliana* as an evolutionary model for understanding the diversity and development of algal symbiosis.

Materials and methods

Sample collection

Two live specimens of *Spurilla braziliana* (designated as individual A and individual B) were collected from the intertidal zone in Miura, Kanagawa Prefecture, Japan, on June 10 and October 5, 2021, respectively. Individual A is the same specimen previously used in our mitochondrial genome sequencing study[24] and was preserved in NucleoProtect RNA solution (MACHEREY-NAGEL). Individual B was maintained in an artificial seawater aquarium (33 g/L, Instant Ocean Premium,

NAPQO, Ltd.) equipped with an external filtration system, under constant light conditions and a

water temperature of 25°C.

Fluorescence Microscopy Observation

Fluorescence microscopy was conducted following the same protocol as described in a previous

study[25]. The cerata of individual A were excised on the day of sampling (prior to immersion in

NucleoProtect RNA) and chemically fixed in 4% paraformaldehyde (PFA). After washing with PBS,

the samples were cleared using the CUBIC Trial Kit (FUJIFILM Wako Pure Chemical Corporation),

according to the protocol provided by the manufacturer. During the clearing process, nuclear staining

was performed using DAPI solution (FUJIFILM Wako Pure Chemical Corporation). Chlorophyll

autofluorescence from Symbiodiniaceae and DAPI fluorescence were observed under a fluorescence

microscope (KEYENCE BZX-810).

Light and Transmission Electron Microscopy of Tissue Sections

Transmission electron microscopy (TEM) was carried out following the same protocol as

described in a previous study[25]. The cerata, tentacles, and rhinophores of individual B, which had

been maintained for 15 days, were dissected and fixed in an aldehyde-based fixative solution (2%

paraformaldehyde, 2% glutaraldehyde, 50 mM HEPES, 33 g/L Instant Ocean Premium, NAPQO

Ltd.). After rinsing with distilled water, the tissues were post-fixed in 1% osmium tetroxide at 4°C

for 1 hour. Following washing and dehydration through a graded ethanol series, the samples were

transferred to propylene oxide and subsequently embedded in epoxy resin (Quetol 812, Nisshin EM).

The resin blocks were sectioned using an ultramicrotome (Leica Ultracut UCT). Semi-thin sections

(500 nm thick) were stained with toluidine blue and observed under a light microscope (KEYENCE

BZX-810). Ultrathin sections (60 nm thick) were stained with 4% aqueous uranyl acetate and

Reynolds' lead citrate, and observed using a transmission electron microscope (JEOL JEM-1010).

Identification of the Symbiotic Symbiodiniaceae Genus

Raw whole-genome shotgun sequence reads from individual A (GenBank Accession ID:

DRR450785)[24] were assembled using CLC Genomics Workbench version 8.5 (QIAGEN Aarhus

A/S). To identify symbiotic Symbiodiniaceae sequences, we used the reference dataset published by

Pochon et al.[26] and conducted blastn searches against all contigs in the primary assembly for six

genetic markers: nuclear 28S rRNA, nuclear elf2, mitochondrial COI, mitochondrial cob, chloroplast

23S rRNA, and chloroplast psbA genes. Contigs identified as Symbiodiniaceae-derived were

subsequently aligned to the reference dataset using MAFFT v7.490[27,28] on Geneious Prime. Based

on these alignments, maximum likelihood phylogenetic trees were constructed using IQ-TREE

version 2.3.3[29] with 1000 bootstrap replicates. To reduce overestimation of branch support, the -bnni

option was applied. Additionally, SH-aLRT[30] and aBayes[31] tests were performed to provide a more

stringent assessment of node support. The final phylogenetic trees were visualized using Interactive

Tree of Life version 6[32].

Visualization of the Hologenome Structure

Raw reads were mapped to the primary assembly using BWA-MEM[33], and the resulting

alignments were processed into BAM files with samtools version 1.17[34]. The BAM files were

analyzed with CoverM version 0.6.1 (https://github.com/wwood/CoverM) to calculate the mean

coverage for each contig. Contig lengths were determined using seqkit version 2.3.0[35] and GNU Awk

version 4.0.2 (https://www.gnu.org/software/gawk/). The 4-mer composition of each contig was

calculated using Perl scripts included in the Multi-metagenome[36]. Feature data from each contig

were visualized as scatter plots using the matplotlib package in Python version 3.10.4. Contig

coverage and length were used as the x- and y-axis coordinates, respectively. The 256-dimensional

4-mer vectors were reduced to three dimensions using UMAP (umap-learn module, version 0.5.3)[37],

normalized, and converted into RGB color values that were applied to the data points in the plot.

Identification of Organisms Present in the Holobiont

Each contig in the primary assembly was taxonomically characterized by homology search against

the NCBI nr database (downloaded on May 19, 2021; https://ftp.ncbi.nlm.nih.gov/blast/db/) using

DIAMOND version 2.0.9[38]. Only hits with an E-value of 0.0 and a bit score within 95% of the top

hit were retained. Taxonomic assignments for each contig were determined using the lowest

common ancestor (LCA) approach. A scatter plot was then redrawn using contig coverage and length

as the x- and y-axes, respectively, with each contig colored according to its assigned taxon.

Results

Whole-Mount Localization of Symbiodiniaceae in the Cerata

Whole-mount fluorescence microscopy revealed the internal structure of the ceras in *Spurilla*

braziliana. While most cells within the ceras exhibited only blue fluorescence from DAPI staining,

red autofluorescence—potentially derived from chlorophyll of Symbiodiniaceae—was observed in

the slightly branched digestive gland located at the center of the ceras (Fig. 1). In contrast, no

autofluorescence was detected in the cnidosacs, which are extensions of the digestive gland.

Light and Transmission Electron Microscopy of Tissue Sections

Observation of toluidine blue-stained sections under a light microscope allowed examination of

the two-dimensional tissue architecture in the cerata, tentacles, and rhinophores of *S. braziliana* (Fig.

2). A gently branched digestive gland extended through the center of each ceras, and multiple

Symbiodiniaceae cells were observed in its epithelial layer (Fig. 2a). All five branches of the digestive gland exhibited similar structure, although algal cells were not observed in the central branch within the limits of our observations. No algal cells were detected outside the digestive gland epithelium within the ceras. In both the tentacles and rhinophores, no penetration of the digestive gland was observed, and no algal cells were present (Fig. 2b, c).

Transmission electron microscopy (TEM) was used to examine the ultrastructure of algal symbiosis within the digestive gland epithelium of the ceras in *S. braziliana* (Fig. 3). The lumen of the digestive gland was lined with microvilli, and its epithelial cells contained numerous vacuole-like structures. In some of these epithelial cells, algal cells were enclosed within the vacuole-like compartments. Many of the algal cells appeared to retain intact cellular architecture, with clearly distinguishable internal structures.

Clade Affiliation of Symbiodiniaceae in *S. braziliana*

Using blastn searches followed by manual alignment inspection, one contig each was identified as homologous to the mitochondrial COI and cob genes based on similarity to reference sequences. In contrast, the remaining four genes (nuclear and chloroplast markers) either produced no significant hits or showed clearly low sequence similarity. Phylogenetic trees were constructed for the two contigs with confirmed homology to the reference dataset (Fig. 4). In both the COI and cob gene trees, the contigs derived from *S. braziliana* formed a monophyletic group with *Symbiodinium* (clade A) species, indicating that the symbiotic algae associated with *S. braziliana* belong to *Symbiodinium*.

Hologenomic Structure in *Spurilla braziliana*

In the *S. braziliana*–Symbiodiniaceae symbiotic system, not only the dinoflagellates but also a broader microbiome may contribute to symbiotic adaptation. To explore this possibility, we visualized the major microbiome components present within the nudibranch using previously obtained whole-genome shotgun sequencing data (Fig. 5). Each contig from the primary assembly was plotted using its coverage and length as spatial coordinates, and its color was determined by dimensional reduction (3D UMAP) of its 4-mer sequence composition. This visualization revealed four clearly distinct peaks, each representing contigs with differing compositional properties (Fig. 5a).

Using DIAMOND blastx searches and taxonomic assignment via the lowest common ancestor (LCA) method, we were able to infer the likely source organisms for 1,294 out of 850,960 contigs in the primary assembly. Among these, taxa represented by more than 50 contigs included *Vibrio* (977 contigs), *Photobacterium* (134 contigs), and Gastropoda (77 contigs), corresponding to the host itself.

An additional 56 bacterial contigs were attributed to taxa other than the two dominant genera. Other
notable assignments included Embryophyta (24 contigs) and Symbiodiniaceae (4 contigs). When
taxonomic assignments were overlaid onto the coverage–contig length scatter plot, it became evident
that distinct taxa underlie the four prominent peaks observed in Fig. 5a (Fig. 5b). Peak 1 was
primarily composed of contigs assigned to the genus *Photobacterium*, while Peak 2 was dominated
by *Vibrio*. Peak 3 corresponded to Gastropoda, reflecting sequences derived from the host organism.
Peak 4, though less sharply defined, exhibited the highest coverage and was attributed to a
heterogeneous group of other bacterial taxa. Contigs matching Symbiodiniaceae or other non-
bacterial organisms did not form distinct peaks in the plot.

Among the bacteria other than *Photobacterium* and *Vibrio*, seven families were identified through
LCA-based analysis at the family level: Cyclobacteriaceae, Enterobacteriaceae, Flammeovirgaceae,
Marinifilaceae, Marinilabiliaceae, Prolixibacteraceae, and Pseudomonadaceae. Of these, contigs
assigned to Enterobacteriaceae and Pseudomonadaceae tended to have relatively low coverage (Fig.
5c).

Discussion

Symbiotic Structure of *Spurilla braziliana* in the Context of Cladobranchia Symbioses

Whole-mount fluorescence microscopy provided a coarse-resolution view of internal structures
within the ceras of *S. braziliana* (Fig. 1). In *Pteraeolidia semperi*, a species within the same
superfamily Aeolidioidea and known for its highly developed symbiotic capacity with
Symbiodiniaceae[10], previous observations[25] revealed intense chlorophyll autofluorescence just
beneath the cerata epidermis. This pattern is attributed to the extensive branching of the digestive
gland that houses the algal cells. In contrast, no such peripheral fluorescence was observed in *S.
braziliana*, where red autofluorescence was detected exclusively in the central digestive gland
running through the ceras. Follow-up histological and ultrastructural observations (Figs. 2, 3)
confirmed that, consistent with general patterns reported in Cladobranchia–Symbiodiniaceae
symbioses[9], algal cells are located within the epithelial cells of the digestive gland.

Histological observations confirmed the findings from whole-mount fluorescence imaging: algal
cells were found exclusively within the digestive gland in the cerata, with no presence in other
tissues. The digestive gland penetrating the cerata exhibited only limited branching (Fig. 1), and
among the five branches observed in Fig. 2a, algal cells were absent only in the central canal. In
contrast, species such as *Pteraeolidia semperi*, *Phyllodesmium briareum*, and *Melibe engeli* possess
highly branched digestive gland systems within the cerata, forming networks of structurally distinct
fine tubules, in which algal symbionts are harbored within epithelial cells of the branches[10,25,39,40].

The branching structure observed in *S. braziliana* bears some resemblance to this organization.

However, unlike *Pteraeolidia*, in which fine tubules extend into both the rhinophores and tentacles[25],

or *P. briareum* and *M. engeli*, where they penetrate at least the rhinophores[39,40], no such branching

structures were found in the rhinophores or tentacles of *S. braziliana* (Fig. 2b, c). Moreover, the

degree of branching in *S. braziliana* was limited (Fig. 1), and no structural differentiation was

evident between the central canal and lateral branches (Fig. 2a). Given that the extent of digestive

gland branching in *Phyllodesmium* has been suggested to correlate with symbiotic efficiency[10,41], the

relatively simple morphology of the digestive gland in *S. braziliana* indicates a lower morphological

capacity for algal symbiosis.

In our field sampling, *P. semperi* was also found in the same coastal area, but whereas *P. semperi*

was collected from sunlit areas, *S. braziliana* was found beneath rocks. This ecological distinction

further suggests that *S. braziliana* may be less adapted to light-dependent symbiosis.

Symbiotic Structure of Spurilla braziliana in the Context of Aeolidiidae Symbioses

Within the family Aeolidiidae, to which *Spurilla* belongs[17,22,42–49], Symbiodiniaceae have been

explicitly reported from 16 nudibranch species and absent in 2 species[11,16–23,47,50,51] (Fig. 6,

Supplementary Table 1). Among them, the genus *Berghia*, which is phylogenetically close to

Spurilla[42,48], has been the most frequently used model for studying Cladobranchia–Symbiodiniaceae

symbioses in recent years[12–15]. Previous studies have suggested that the symbiotic capacity of

Berghia is relatively primitive, with algal cells being retained for only about one week[11,18]. The

symbiosis observed in *Spurilla* may represent a similar state.

Among *Spurilla* and *Baeolidia* species more closely related to *S. braziliana* than *Berghia*, detailed

histological studies of *S. neapolitana*[16], *Baeolidia australis*[19], and *Baeolidia moebii*[19] have reported

penetration of the digestive gland into the rhinophores. In contrast, *S. braziliana* exhibited no such

penetration into the rhinophores, suggesting an even lower level of adaptation to symbiosis.

Supporting this, a previous study based on external morphology reported that *S. neapolitana*

displayed pigmentation likely derived from Symbiodiniaceae, while such coloration was inconsistent

or unstable in *S. braziliana*[22]—further reinforcing the interpretation that *S. braziliana* exhibits

relatively limited symbiotic capacity. However, this time we did not assess the algal retention

capacity directly. Future investigations that track temporal changes in algal load under controlled

conditions—specifically, in the complete absence of potential cnidarian prey—will be essential to

clarify the symbiotic capacity of *S. braziliana* in greater detail.

In any case, the observation that symbiotic capacity varies at intra- and interspecific, as well as

intergeneric levels strongly suggests that comparative analyses among *Spurilla*, *Baeolidia*, and

Berghia provide a powerful evolutionary model for studying animal–algal symbiosis. In this context,

S. braziliana may serve as a critical intermediate, bridging the gap between symbiotic and non-

symbiotic nudibranchs. Notably, *Baeolidia salaamica*—the only species among the three genera for

which a lack of symbiotic capacity has been reported[19]—could also serve as an important

comparative reference in future studies.

Microbiome Composition in *Spurilla braziliana*

From the whole-genome shotgun (WGS) assembly of *S. braziliana*, single contigs corresponding

to the COI and cob genes of dinoflagellates were identified. Phylogenetic analysis showed that both

sequences belong to the genus *Symbiodinium* (clade A) (Fig. 4). Nudibranchs within the suborder

Cladobranchia have been reported to form symbioses with Symbiodiniaceae clades A through D—

Symbiodinium, *Breviolum*, *Cladocopium*, and *Durusdinium*—with the specific clade often reflecting

the host's geographic distribution[52–58]. *Symbiodinium* (clade A) is frequently detected in nudibranchs

from temperate regions[53] and has previously been reported in *Pteraeolidia semperi* from Hayama[52],

a locality very close to the present sampling site. This finding aligns with known biogeographic

trends in Cladobranchia–Symbiodiniaceae associations.

Regarding the bacterial community of *S. braziliana*, contigs assigned to the genera

Photobacterium and *Vibrio* were detected at relatively low coverage. These genera are commonly

reported as intestinal bacteria in marine gastropods[59]. In addition, members of the families

Cyclobacteriaceae, Enterobacteriaceae, Flammeovirgaceae, Marinifilaceae, Marinilabiliaceae,

Prolixibacteraceae, and Pseudomonadaceae were identified. Among these, Enterobacteriaceae and

Pseudomonadaceae have been reported to co-occur with certain clades of Symbiodiniaceae in coral

hosts[60]. However, their functional roles remain poorly understood, and the study has specifically

noted a lack of co-occurrence with *Symbiodinium* (clade A)[60]. Therefore, caution is warranted in

interpreting any direct contribution of these bacterial taxa to algal symbiosis in *S. braziliana*.

Among bacterial taxa associated with photosymbiosis, the genus *Endozoicomonas* has attracted

the most attention. In cnidarian hosts, members of this family are thought to positively influence

algal symbiosis by modulating host immunity and supplying essential metabolites that support

Symbiodiniaceae photosynthesis[61–63]. In nudibranch hosts as well, *Endozoicomonas* has been

identified in species with both advanced (*Pteraeolidia semperi*)[64] and rudimentary (*Berghia*

stephanieae)[65] symbiosis. However, in the present study, no contigs assigned to

Endozoicomonadaceae were detected from *S. braziliana*. This absence of symbiosis-associated

bacterial taxa may further support the interpretation that *S. braziliana* possesses a low level of

symbiotic adaptation. That said, the actual functional roles of bacterial communities in marine

invertebrate–Symbiodiniaceae symbioses remain poorly understood. As our understanding of host–
symbiont–microbiome interactions deepens, it will become possible to more precisely evaluate the
symbiotic potential of species like *S. braziliana*.
292

Conclusion
The study of nudibranch-Symbiodiniaceae symbiosis remains in its early stages, and the diversity
of symbiotic capacity across species is still poorly understood. In this study, we conducted the first
histological and metagenomic analyses of endosymbiosis in *Spurilla braziliana*, a species previously
unexamined in this context. The combination of a simple digestive morphology for algal retention
and the absence of bacterial taxa typically associated with algal symbiosis suggests that *S. braziliana*
exhibits only minimal algal symbiotic capacity. These findings contribute new insight into the
diversity of nudibranch–Symbiodiniaceae interactions and underscore the value of *S. braziliana* as a
potential model for exploring the evolutionary origins and acquisition of photosymbiosis. As
comparative analyses of related species continue to expand, we will be better equipped to bridge
interspecific and intergeneric gaps and identify the key factors underlying the emergence of
symbiotic capacity in marine invertebrates.
305

Acknowledgement
This study was supported by JSPS KAKENHI Grants-in-Aid for Scientific Research (Grant
Numbers: 24KJ0827, 20H00429, and 23H0034), the Sasakawa Scientific Research Grant (2022-
4042), and The ANRI Fellowship (2024).

Author contributions
H.M. conceived the study. H.M., R.Y., and K.H. conducted sampling and maintenance of live
nudibranchs. H.M., K.T., and K.H. performed histological observations. H.M. carried out
microbiome analyses. H.M. interpreted the results and wrote the manuscript. All authors contributed
to discussions and revised the manuscript.
316

Reference

1. Jacobovitz, M. R., Hambleton, E. A. & Guse, A. Unlocking the Complex Cell

Biology of Coral-Dinoflagellate Symbiosis: A Model Systems Approach. *Annu. Rev.*

Genet. **57**, 411-434 (2023).

2. Mies, M. Evolution, diversity, distribution and the endangered future of the

giant clam-Symbiodiniaceae association. *Coral Reefs* **38**, 1067–1084 (2019).

3. LeKieffre, C. *et al.* Assimilation, translocation, and utilization of carbon

between photosynthetic symbiotic dinoflagellates and their planktic

foraminifera host. *Mar. Biol.* **165**, 104 (2018).

4. Rossbach, S., Subedi, R. C., Ng, T. K., Ooi, B. S. & Duarte, C. M. Iridocytes

Mediate Photonic Cooperation Between Giant Clams (Tridacninae) and Their

Photosynthetic Symbionts. *Front. Mar. Sci.* **7**, 465 (2020).

5. Jung, J. *et al.* Coral photosymbiosis on Mid-Devonian reefs. *Nature* **636**, 647–

653 (2024).

6. Grottoli, A. G., Rodrigues, L. J. & Palardy, J. E. Heterotrophic plasticity

and resilience in bleached corals. *Nature* **440**, 1186–1189 (2006).

7. Wiedenmann, J. *et al.* Reef-building corals farm and feed on their

photosynthetic symbionts. *Nature* **620**, 1018–1024 (2023).

8. Hughes, T. P. *et al.* Global warming and recurrent mass bleaching of corals.

Nature **543**, 373–377 (2017).

9. Rola, M. *et al.* Cladobranchia (Gastropoda, Nudibranchia) as a Promising Model

to Understand the Molecular Evolution of Photosymbiosis in Animals. *Front. Mar.*

Sci. **8**, 745644 (2022).

10.   Burghardt, I. Symbiosis between Symbiodinium (Dinophyceae) and various

taxa of Nudibranchia (Mollusca: Gastropoda), with analyses of long-term

retention. *Org. Divers. Evol.* **8**, 66–76 (2008).

11.   Monteiro, E. A., Güth, A. Z., Banha, T. N. S., Sumida, P. Y. G. & Mies,

M. Evidence against mutualism in an aeolid nudibranch associated with

Symbiodiniaceae dinoflagellates. *Symbiosis* **79**, 183–189 (2019).

12.   Silva, R. X. G., Cartaxana, P. & Calado, R. Prevalence and Photobiology

of Photosynthetic Dinoflagellate Endosymbionts in the Nudibranch Berghia

stephanieae. *Animals* **11**, 2200 (2021).

13.   Mies, M. *et al.* Expression of a symbiosis-specific gene in *Symbiodinium*

type A1 associated with coral, nudibranch and giant clam larvae. *R. Soc. Open*

Sci. **4**, 170253 (2017).

14.   Clavijo, J. M. *et al.* The nudibranch Berghia stephanieae (Valdés, 2005)

is not able to initiate a functional symbiosome-like environment to maintain

Breviolum minutum (J.E.Parkinson & LaJeunesse 2018). *Front. Mar. Sci.* **9**,

934307 (2022).

15.   Silva, R. X. G., Madeira, D., Cartaxana, P. & Calado, R. Assessing the

Trophic Impact of Bleaching: The Model Pair Berghia stephanieae/Exaiptasia

diaphana. *Animals* **13**, 291 (2023).

16.   Marin, A. & Ros, J. Presence of Intracellular Zooxanthellae in

Mediterranean Nudibranchs. *J. Molluscan Stud.* **57**, 87-101 (1991).

17.   Carmona, L., Pola, M., Gosliner, T. M. & Cervera, J. L. The Atlantic-

Mediterranean genus *Berghia* Trinchese, 1877 (Nudibranchia: Aeolidiidae):

taxonomic review and phylogenetic analysis. *J. Molluscan Stud.* **80**, 482-498

(2014).

18.   Kempf, S. C. A 'Primitive' Symbiosis between the Aeolid Nudibranch

Berghia Verrucicornis (A. Costa, 1867) and a Zooxanthella. *J. Molluscan Stud.*

57, 75-85 (1991).

19.   Rudman, W. B. The taxonomy and biology of further aeolidacean and

arminacean nudibranch molluscs with symbiotic zooxanthellae. *Zool. J. Linn.*

Soc. **74**, 147-196 (1982).

20.   Kempf, S. C. SYMBIOSIS BETWEEN THE ZOOXANTHELLA SYMBIODINIUM (=

GYMNODINIUM) MICROADRIATICUM (FREUDENTHAL) AND FOUR SPECIES OF NUDIBRANCHS.

Biol. Bull. **166**, 110-126 (1984).

21.   CABRITO, A., GALIÀ-CAMPS, C. & BALLESTEROS, M. The risk of the

nudibranch Baeolidia rieae (Mollusca: Gastropoda: Heterobranchia) in the

376   international aquarium trade. (2022) doi:10.21411/CBM.A.92E8654F.

22.   Carmona, L. *et al.* Untangling the Spurilla neapolitana (Delle Chiaje,

1841) species complex: a review of the genus Spurilla Bergh, 1864 (Mollusca:

Nudibranchia: Aeolidiidae). *Zool. J. Linn. Soc.* (2014) doi:10.1111/zoj12098.

23.   Wägele, M. & Johnsen, G. Observations on the histology and

photosynthetic performance of "solar-powered" opisthobranchs (Mollusca,

Gastropoda, Opisthobranchia) containing symbiotic chloroplasts or

zooxanthellae. *Org. Divers. Evol.* **1**, 193–210 (2001).

24.   Mizobata, H. *et al.* The complete mitochondrial genome of *Spurilla*

braziliana MacFarland 1909 (Nudibranchia, Aeolidiidae). *Mitochondrial DNA Part*

B **8**, 862–866 (2023).

25.   Mizobata, H. *et al.* The highly developed symbiotic system between the

solar-powered nudibranch Pteraeolidia semperi and Symbiodiniacean algae.

iScience **26**, 108464 (2023).

26.   Pochon, X., Putnam, H. M. & Gates, R. D. Multi-gene analysis of

Symbiodinium dinoflagellates: a perspective on rarity, symbiosis, and

evolution. *PeerJ* **2**, e394 (2014).

27.   Katoh, K. MAFFT: a novel method for rapid multiple sequence alignment

based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066 (2002).

28.   Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software

Version 7: Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**,

772–780 (2013).

29.   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era. *Mol. Biol. Evol.* **37**, 1530–1534

(2020).

30.   Guindon, S. *et al.* New Algorithms and Methods to Estimate Maximum-

Likelihood Phylogenies: Assessing the Performance of PhyML 3.0. *Syst. Biol.* **59**,

307–321 (2010).

31.   Anisimova, M. *et al.* Survey of Branch Support Methods Demonstrates

Accuracy, Power, and Robustness of Fast Likelihood-based Approximation Schemes.

Syst. Biol. **60**, 685–699 (2011).

32.   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online

tool for phylogenetic tree display and annotation. *Nucleic Acids Res.* **49**,

W293–W296 (2021).

33.   Li, H. Aligning sequence reads, clone sequences and assembly contigs

with BWA-MEM. *ArXiv Genomics* (2013) doi:10.6084/m9.figshare.963153.v1.

34.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**,

giab008 (2021).

35.   Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A Cross-Platform and Ultrafast

Toolkit for FASTA/Q File Manipulation. *PLOS ONE* **11**, e0163962 (2016).

36.   Albertsen, M. *et al.* Genome sequences of rare, uncultured bacteria

obtained by differential coverage binning of multiple metagenomes. *Nat.*

Biotechnol. **31**, 533–538 (2013).

37.   McInnes, L. & Healy, J. UMAP: Uniform Manifold Approximation and

Projection for Dimension Reduction. *ArXiv Mach. Learn.* (2018).

38.   Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at

tree-of-life scale using DIAMOND. *Nat. Methods* **18**, 366–368 (2021).

39.   Wagele, H. Potential key characters in Opisthobranchia (Gastropoda,

Mollusca) enhancing adaptive radiation. *Org. Divers. Evol.* **4**, 175–188 (2004).

40.   Burghardt, I. & Wägele, H. The symbiosis between the 'solar-powered'

nudibranch *Melibe engeli* Risbec, 1937 (Dendronotoidea) and *Symbiodinium* sp.

(Dinophyceae). *J. Molluscan Stud.* **80**, 508–517 (2014).

41.   Rudman, W. B. FURTHER STUDIES ON THE TAXONOMY AND BIOLOGY OF THE

OCTOCORAL-FEEDING GENUS PHYLLODESMIUM EHRENBERG, 1831 (NUDIBRANCHIA:

AEOLIDOIDEA). *J. Molluscan Stud.* **57**, 167–203 (1991).

42.   Carmona, L., Pola, M., Gosliner, T. M. & Cervera, J. L. A Tale That

Morphology Fails to Tell: A Molecular Phylogeny of Aeolidiidae (Aeolidida,

Nudibranchia, Gastropoda). *PLoS ONE* **8**, e63000 (2013).

43.   Carmona, L., Pola, M., Gosliner, T. M. & Cervera, J. L. Review of

Baeolidia, the largest genus of Aeolidiidae (Mollusca: Nudibranchia), with the

description of five new species. *Zootaxa* **3802**, (2014).

44.   Carmona, L., Pola, M., Gosliner, T. M. & Cervera, J. L. The end of a

long controversy: systematics of the genus Limenandra (Mollusca: Nudibranchia:

Aeolidiidae). *Helgol. Mar. Res.* **68**, 37–48 (2014).

45.   Kienberger, K. *et al. Aeolidia papillosa* (Linnaeus, 1761) (Mollusca:

Heterobranchia: Nudibranchia), single species or a cryptic species complex? A

morphological and molecular study. *Zool. J. Linn. Soc.* **177**, 481–506 (2016).

46.   Carmona, L., Cervera, J. L., Kumar, A. B. & Snehachandran, B. K. First

record of the Aeolid Anteaeolidiella fijensis (Nudibranchia, Aeolidiidae) from

India. *Mar. Biodivers.* **47**, 823–830 (2017).

47.   Carmona, L., Pola, M., Gosliner, T. M. & Cervera, J. L. Integrative

taxonomy and biogeography of the genus Bulbaeolidia (Nudibranchia: Aeolidida).

J. Molluscan Stud. **83**, 440–450 (2017).

48.   Karmeinski, D. *et al.* Transcriptomics provides a robust framework for

the relationships of the major clades of cladobranch sea slugs (Mollusca,

Gastropoda, Heterobranchia), but fails to resolve the position of the

enigmatic genus Embletonia. *BMC Ecol. Evol.* **21**, 226 (2021).

49.   MolluscaBase Eds. MolluscaBase. Accessed through: World Register of

Marine Species on 2025-06-30. VLIZ https://doi.org/10.14284/448 (2025).

50.   McFarland, F. K. & Muller-Parker, G. Photosynthesis and Retention of

Zooxanthellae and Zoochlorellae Within the Aeolid Nudibranch *Aeolidia*

papillosa. Biol. Bull. **184**, 223–229 (1993).

51.   Burghardt, I. & Wägele, H. Interspecific differences in the efficiency

and photosynthetic characteristics of the symbiosis of 'solarpowered'

Nudibranchia (Mollusca: Gastropoda) with zooxanthellae. *Rec. West. Aust. Mus.*

Suppl. **69**, 1 (2006).

52.   Ishikura, M. *et al.* Isolation of New Symbiodinium Strains from Tridacnid

Giant Clam (Tridacna crocea) and Sea Slug (Pteraeolidia ianthina) Using

Culture Medium Containing Giant Clam Tissue Homogenate. *Mar. Biotechnol.* **6**,

378–385 (2004).

53.   Loh, W., Cowlishaw, M. & Wilson, N. Diversity of Symbiodinium

dinoflagellate symbionts from the Indo-Pacific sea slug Pteraeolidia ianthina

(Gastropoda: Mollusca). *Mar. Ecol. Prog. Ser.* **320**, 177–184 (2006).

54.   FitzPatrick, S. K. *et al.* Symbiodinium diversity in the soft coral

Heteroxenia sp. and its nudibranch predator Phyllodesmium lizardensis. *Coral

Reefs* **31**, 895–905 (2012).

55.   Ziegler, M. *et al.* Thermal stress response in a dinoflagellate-bearing

nudibranch and the octocoral on which it feeds. *Coral Reefs* **33**, 1085–1099

(2014).

56.   Yorifuji, M., Takeshima, H., Mabuchi, K., Watanabe, T. & Nishida, M.

Comparison of Symbiodinium dinoflagellate flora in sea slug populations of the

Pteraeolidia ianthina complex. *Mar. Ecol. Prog. Ser.* **521**, 91–104 (2015).

57.   Wecker, P. *et al.* Dinoflagellate diversity among nudibranchs and sponges

from French Polynesia: Insights into associations and transfer. *C. R. Biol.*

338, 278–283 (2015).

58.   Soon, N., Quek, Z. B. R., Pohl, S. & Wainwright, B. J. More than meets

the eye: characterizing the cryptic species complex and Symbiodiniaceae

communities in the reef-dwelling nudibranch *Pteraeolidia* ' *semperi* '

(Nudibranchia: Aeolidioidea) from Singapore. *J. Molluscan Stud.* **89**, eyad011

(2023).

59.   Gafarova, E. *et al.* Gut Bacteriomes and Ecological Niche Divergence: An

Example of Two Cryptic Gastropod Species. *Biology* **12**, 1521 (2023).

60.   Bernasconi, R., Stat, M., Koenders, A. & Huggett, M. J. Global Networks

of Symbiodinium-Bacteria Within the Coral Holobiont. *Microb. Ecol.* **77**, 794–807

(2019).

61.   Pogoreutz, C. *et al.* Coral holobiont cues prime *Endozoicomonas* for a

symbiotic lifestyle. *ISME J.* **16**, 1883–1895 (2022).

62.   Pogoreutz, C. & Ziegler, M. Frenemies on the reef? Resolving the coral-

Endozoicomonas association. *Trends Microbiol.* **32**, 422–434 (2024).

63.   Wada, N. *et al.* High-resolution spatial and genomic characterization of

coral-associated microbial aggregates in the coral *Stylophora pistillata*. *Sci.*

Adv. **8**, eabo2431 (2022).

64.   Ng, M. S., Soon, N., Chang, Y. & Wainwright, B. J. Bacterial and Fungal

Co-Occurrence in the Nudibranch, Pteraeolidia semperi. *Life* **12**, 1988 (2022).

65.   Sickinger, C. *et al.* Microbiome origin and stress-related changes in

bacterial abundance of the photosymbiotic sea slug Berghia stephanieae (Á.

`Valdés, 2005).` *`Symbiosis`* **`93`**`, 177–192 (2024).`

Figure captions

Figure 1. Whole-mount fluorescence microscopy image of a *Spurilla braziliana* ceras.

Red autofluorescence is visible exclusively in the digestive gland. Blue fluorescence indicates DAPI

staining.

Figure 2. Toluidine blue-stained sections of various tissues in *Spurilla braziliana*.

a. Cross-section of a ceras. The digestive gland divides into five diverticula, within which algal cells

are located in the epithelium. Arrowheads indicate algal cells.

b. Cross-section of a tentacle. No penetration of the digestive gland or presence of algal cells is

observed.

c. Cross-section of a rhinophore. No penetration of the digestive gland or presence of algal cells is

observed.

Figure 3. Cross-section of the digestive gland within a ceras.

Algal cells are enclosed within the epithelial cells. Arrows: algal cells; dg: digestive gland lumen;

vls: vacuole-like structures. Scale bar: 5 μm.

Figure 4. Phylogenetic placement of Symbiodiniaceae-derived sequences detected in *Spurilla*

***braziliana*.**

Branch lengths are not to scale for clarity. For full trees with branch lengths, see Supplementary Data

1. a. COI gene; b. cob gene.

Figure 5. Hologenomic structure of the *Spurilla braziliana* – Symbiodiniaceae holobiont.

a. Coverage–contig length scatter plot colored by sequence composition.

b. Coverage–contig length scatter plot colored by taxonomic assignment. Each peak corresponds to a

different biological origin. Black: no hit; yellow: *Photobacterium*; blue: *Vibrio*; red: Gastropoda;

brown: other bacteria; green: Embryophyta; light green: Symbiodiniaceae; gray: others.

c. Coverage–contig length scatter plot focusing on contigs assigned to bacterial taxa.

Figure 6. Presence of Symbiodiniaceae in the family Aeolidiidae.

The species list of Aeolidiidae follows MolluscaBase as accessed via WoRMS on June 30, 2025[49].

Green: species for which the presence of Symbiodiniaceae has been reported. Gray: species for

which the absence of Symbiodiniaceae has been reported. No color: species for which neither

presence nor absence has been confirmed. Red box: present study. See Supplementary Table 1 for

references supporting phylogenetic placement and symbiont status.

cnidsac

digestive gland

1000 µm

vls

vls

vls

vls

dg

5 um

a COI

b cob