Multi-omics evaluation of cell lines as models for metastatic prostate cancer

Xueying Liu[1], Yugang Wang[2,*], Ke Liu[1,*]

Affiliations:

1. Department of Medical Dataology, School of Public Health, Cheeloo College of

Medicine, Shandong University, Jinan, China

2. Department of Biochemistry and Molecular Biology, School of Basic Medicine,

Tongji Medical College, Huazhong University of Science and Technology, Wuhan,

China

* To whom correspondence should be addressed

Abstract

Cancer cell lines are indispensable for prostate cancer research, yet their ability to model metastatic prostate cancer remains insufficiently characterized. In this study, we performed a systematic multi-omics evaluation integrating genomic and transcriptomic data from large public datasets. We identified VCaP as the most representative model for metastatic prostate adenocarcinoma, while highlighting the limited fidelity of the widely used PC3 cell line. Our analysis underscores the importance of selecting hypermutated cell lines, which display distinct tumor microenvironment characteristics. Moreover, overexpression of *MET* in VCaP induced partial stem-like traits but did not fully recapitulate the MSPC transcriptomic profile. Notably, stem-like prostate organoids demonstrated higher similarity to MSPC and may serve as more suitable MSPC models. Our study provides guidance for the selection of prostate cancer cell lines, assisting researchers in choosing more representative metastatic prostate cancer cell line models.

Introduction

Cancer cell lines are indispensable preclinical models in oncology research. Their immortality, cost-effectiveness, and capacity for long-term cultivation and cloning make them vital for reproducible and scalable cancer research and drug development[1, 2, 3, 4, 5]. However, prolonged *in vitro* culture can lead to the accumulation of additional genetic and epigenetic alterations, resulting in substantial discrepancies compared to patient-derived tumor samples[6, 7, 8, 9]. Therefore, systematic evaluation of the molecular fidelity of cancer cell lines relative to patient-derived cancer cells is essential to ensure the translational relevance of cell line–based studies.

The widespread availability of large-scale open cancer genomics data has enabled researchers to comprehensively assess the suitability of cell lines as models in cancer research. For example, Domcke et al. compared genomic profiles between ovarian cancer cell lines and patient tumor samples and found several commonly used cell lines poorly recapitulated high-grade serous ovarian cancer[10]; Yu et al. performed a pan-cancer transcriptomic comparison between cell lines and patient tumor samples and proposed the cell line panel TCGA-110-CL[11]; Jin et al. conducted similar pan-cancer analysis with their own computational approaches to select appropriate cell line models for different cancer types; We previously performed multi-omics comparison between breast cancer cell lines and metastatic tumor samples and identified cell lines that faithfully represent distinct breast cancer subtypes[12]. These

studies have deepened our understanding of cancer cell lines and highlighted the

importance of selecting appropriate models based on molecular fidelity.

Prostate cancer is among the most common and lethal malignancies affecting men's

health worldwide, posing a substantial public health challenge[13, 14]. Similar to other

cancer types, cell lines have been widely utilized to explore the underlying biological

mechanisms of prostate cancer. Although several studies have evaluated the molecular

representativeness of prostate cancer cell lines relative to tumor samples[11, 15, 16], three

major limitations persist. First, metastasis is the leading cause of prostate

cancer–related mortality[13, 17]; however, most of the evaluation studies mainly focus on

primary tumors. Second, existed studies were performed in a pan-cancer manner,

which applies uniform analytical criteria across various cancer types and overlooks

tumor-specific characteristics[3]. Third, the majority of analyses rely exclusively on

transcriptomic data, with limited integration of multi-omics information for a more

comprehensive evaluation.

In this study, we leveraged the publicly available bulk and single-cell omics data to

comprehensively evaluate the suitability of prostate cancer cell lines as models for

metastatic prostate cancer. We proposed principles of cell line selection for different

research scenarios and meanwhile point the strength and limitation of widely used

established cell line. Our study provides a framework for selecting prostate cancer cell

lines tailored to specific biological contexts and highlight the need to develop more

representative models for understudied subtypes.

Results

Systematic comparison of genomic profiles

We first compared the somatic mutation profiles between metastatic prostate cancer

samples and prostate cancer cell lines, focusing on genes that were either highly

mutated in metastatic prostate cancer or exhibited differential mutation rates between

metastatic and primary prostate cancer. The somatic mutation data of primary prostate

cancer, metastatic prostate cancer, and prostate cancer cell lines were downloaded

from the Cancer Genome Atlas (TCGA), Stand Up to Cancer (SU2C), and the Cancer

Cell Line Encyclopedia (CCLE), respectively. A total of 68 highly mutated genes

were identified using SU2C samples, along with 13 differentially mutated genes

between SU2C and TCGA samples (**Supplementary. Fig. 1a-c**). The integration of

these two lists identified 73 unique genes (**Fig. 1a**). The median mutation rate of the

73 genes across the CCLE prostate cancer cell lines was 0.3, suggesting most of them

could be recapitulated by only a few cell lines. *MUC16*, *SYNE2*, *SYNE1*, *OBSCN* have

a mutation rate higher than 50%, with *MUC16* being the most frequently mutated

gene (mutation rate = 0.62). *TP53*, a key tumor suppressor gene, displayed a mutation

rate of 0.38. Surprisingly, there were four genes (*APC*, *SPOP*, *CDK12*,

and *FAM186A)* which were entirely unmutated in all analyzed prostate cell lines,

highlighting the limitation of using the analyzed cell lines in researching the role of

them in metastasis-related study.

Cell lines were ranked based on the number of mutations present in a set of 73 genes

associated with metastatic prostate cancer, and found the top four cell lines were

LNCaP, DU145, 22RV1, and MDA-PCa-2b, respectively **(Fig 1b)**. Interestingly, we

noticed these four cell lines were all inferred with microsatellite instability (MSI)[18]

and had significantly higher mutation burden than other cell lines **(Fig. 1c)**. To adjust

such bias, we re-ranked the cell lines according to a normalized ratio value (computed

as number of mutated genes / mutation burden) and found the top three cell lines were

PC3, VCaP, and NCI-H660 (**Fig. 1d**).

We next investigated whether those recurrent single-nucleotide alterations (which we

call "mutation hotspot") identified in metastatic prostate cancer samples were also

presented in prostate cancer cell lines. We defined mutation hotspots as

non-synonymous mutations occurring in at least three SU2C samples. Based on this

criterion, 58 mutation hotspots were identified (**Supplementary Table 1**). These

mutation hotspots were distributed in 45 genes, with *TP53* and *AR* being the two

genes harboring most mutation hotspots (**Fig. 1e and Supplementary. Fig. 1d)**. *TP53*

mutation hotspots were predominantly located within the DNA-binding domain, a

functionally critical region required for the transcriptional regulation of tumor suppressor genes[19]. For *AR*, mutation hotspots were primarily clustered within the ligand-binding domain, which is pivotal for androgen receptor-targeted therapies[20].

Strikingly, only seven mutation hotspots were detected in at least one prostate cancer cell line (**Fig. 1f**), and all were presented exclusively in 22RV1, MDA-PCa-2b, and LNCaP. Of them, three were single nucleotide point mutations of *AR* and the other four were all frameshift mutations. Of the *AR* mutation hotspots, T878A and L702H mutations have been reported to be associated with resistance to AR inhibition therapy[21]. Notably, MDA-PCa-2b harbors both mutations (T878A and L702H), reflecting the clinical phenomenon of co-occurring multiple AR resistance mutations within individual tumors[21]. Among the frameshift mutation hotspots, K16fs mutation in *RPL22* was observed in all three cell lines, whereas the other three frameshift mutations were exclusive to 22RV1.

Besides somatic mutation profiles, we also compared copy number variation (CNV) profiles between metastatic and primary prostate cancer samples. Among the differential CNV genes between SU2C and TCGA samples, *AR* exhibited the highest level of amplification (**Fig. 1g**). A similar pattern was observed in site-specific analyses of differential CNV genes (**Supplementary. Fig. 1e-g**). This finding may not be surprising since *AR* is a critical driver of prostate cancer development and

progression[22, 23, 24]. Notably, VCaP exhibited the highest level of *AR* amplification

among all prostate cancer cell lines analyzed (**Fig. 1h**).

Hypermutation of prostate cancer cells influenced the tumor microenvironment

Previous studies have demonstrated hypermutated colorectal tumor samples

exhibiting increased tumor-infiltrating, elevated neoantigen burden, and potentially

enhanced sensitivity to immunotherapy[25, 26]. Motivated by these findings, we aim to

assess whether hypermutation should be taken into consideration when selecting cell

lines for studying the metastatic mechanisms of prostate cancer.

To identify hypermutated prostate cancer samples, we integrated samples from TCGA,

SU2C, CCLE together and determined a mutation burden threshold above which

samples were classified as hypermutated (**Fig. 2a**). Based on our criteria, we

identified four hypermutated samples in TCGA, ten in SU2C, and four in CCLE. As

expected, the aforementioned four hypermutated cell lines LNCaP, DU145, 22RV1,

and MDA-PCa-2b were successfully identified. Notably, three of them harbor *RPL22*

frameshift mutations, which is frequently observed in MSI tumors[27, 28]. To further

validate our strategy, we computed the mutation frequencies of mismatch repair

(MMR) genes[29]   and found hypermutated samples exhibiting a significantly higher

mutation frequency (**Fig. 2b and Supplementary Fig. 2a**). These results

demonstrated the validity of the determined threshold value for identifying

hypermutated prostate cancer samples.

We next performed differential gene expression analysis between hypermutated and

non-hypermutated TCGA samples and identified 125 upregulated and 85

downregulated genes, respectively (**Supplementary Table 2**). Notably, the T cell

marker *CD8A*, the immune checkpoint molecule *CTLA-4*, and the cytotoxic effector

GZMA were significantly upregulated in hypermutated samples (**Fig. 2c and**

Supplementary. Fig. 2b), suggesting the increased CD8+ T cell infiltration level; in

contrast, the gamma-delta T cell marker *TRGV9* was downregulated, suggesting a

potential reduction in gamma-delta T cell infiltration in hypermutated tumors. We

performed Gene Ontology (GO) enrichment analysis on upregulated genes and found

pathways associated with T cell activation (e.g., the T cell receptor signaling pathway,

antigen receptor-mediated signaling pathway) and natural killer (NK) cell-mediated

cytotoxicity (e.g., NK cell-mediated immunity, NK cell-mediated cytotoxicity) were

significantly enriched (**Fig. 2d and Supplementary Table 3**). The tumor

microenvironment (TME) has been demonstrated to play pivotal roles in the distant

metastasis of primary cancer[30, 31, 32] and our analysis suggests notable TME

differences between hypermutated and non-hypermutated primary prostate cancer

samples. Therefore, we propose that hypermutated cell lines (such as LNCaP) should

be more appropriate for investigating the metastatic mechanisms of hypermutated

prostate cancer. To the best of our knowledge, such consideration has not been

addressed in previous cell line evaluation studies.

Correlating cell lines with metastatic prostate cancer using bulk and single-cell

RNA-seq data

Transcriptome correlation analysis (TC analysis) has been proven to be an effective

method for assessing the utility of cell lines[11, 12, 33]. Consequently, we ranked all 1,019

CCLE cell lines based on their transcriptomic similarity to MET500 prostate cancer

samples and the top three cell lines were LNCaP, VCaP, and MDA-PCa-2b,

respectively (**Fig. 3a**). We next examined whether the metastatic site influenced TC

analysis results and found that cell line rankings were highly consistent among

various metastatic sites (bone, liver, and lymph node) (**Fig. 3b and Supplementary.**

Fig. 3a-c).

To double confirm the analysis results derived from bulk RNA-seq data, we further

ranked 1,019 CCLE cell lines according to their transcriptomic similarity with the

malignant cells from a scRNA-seq dataset of metastatic prostate cancer[22]. Consistent

with previous results, the top three cell lines with the highest similarity were VCaP,

LNCaP, and MDA-PCa-2b (**Fig. 3c**). We further performed metastatic-site-specific

transcriptomic similarity analysis (bone and lymph node) and the results were highly

correlated (**Fig. 3d and Supplementary Fig. 3d-e**).

Strength and limitation of the PC3 cell line

PC3 is a widely used cell line in studies investigating metastatic prostate cancer[34, 35]. We conducted a PubMed search of studies related to metastatic prostate cancer and found PC3 was the most frequently cited cell line, followed by LNCaP and DU145 (**Supplementary Table 4 and Supplementary. Fig. 4a**). However, despite its widespread use, transcriptomic correlation analysis revealed that PC3 shows limited similarity to metastatic prostate cancer samples (ranked 444 among 1,019 CCLE cell lines) (**Fig .3a**). Prior to our research, several studies had revealed the heterogeneity of prostate cancer[36, 37, 38]. Considering that the metastatic prostate cancer samples used in the transcriptome correlation analysis mainly exhibited adenocarcinoma histology (in which malignant cells primarily show luminal differentiation)[39, 40, 41], we further investigated whether PC3 could serve as a representative model for other types of prostate cancer.

We performed principal component analysis (PCA) on CCLE prostate cancer cell lines and found they could be briefly classified into three clusters (**Fig. 4a**). Cluster 1 consisted of luminal lineage cell lines (such as VCaP), as indicated by expression of luminal lineage markers such as *AR* and *KLK4*; cluster 2 only included NCI-H660, a small cell carcinoma cell line of neuroendocrine origin expressing *SYP*; cluster 3 comprised DU145 and PC3. Interestingly, our PCA results align with Han et al 's research, which classified prostate cancer into three intrinsic subtypes: androgen

receptor positive prostate cancer (ARPC), neuroendocrine prostate cancer (NEPC),

and mesenchymal and stem-like prostate cancer (MSPC)[42]. Since PC3 and DU145

were separated with other cell lines on the first principal component, we performed

gene set enrichment analysis on the top 200 genes with the highest positive loadings

along the PC1 axis using MSigDB Hallmark gene sets (**Fig. 4b**). The

"Epithelial-Mesenchymal Transition" gene set exhibited the highest level of

enrichment; in addition, the enrichment of "*IL-2/STAT5* Signaling" and

"*IL-6/JAK/STAT3* Signaling" gene sets suggests activation of the *JAK/STAT* pathway,

a known driver of cellular plasticity[41, 43]. Furthermore, PC3 exhibited relatively higher

expression of *CD44* (a marker of stemness), along with lower expression of the

luminal marker *AR* and the neuroendocrine marker *SYP* (**Fig. 4c-e**). We double

confirmed the expression of these marker genes using a scRNA-seq dataset of PC3

(**Supplementary. Fig. 4b**). Collectively, these results support the classification of

PC3 as a cell line of the MSPC subtype.

We next evaluated whether PC3 could adequately recapitulate the transcriptomic

profile of MSPC malignant cells. MSPC has been reported to be enriched in

castration-resistant prostate cancer (CRPC) patients[42]. To this end, we analyzed a

CRPC scRNA-seq dataset and found the malignant cells could be classified into

ARPC, NEPC, and MSPC subtypes based on the expression of *AR*, *SYP*, and *CD44*,

respectively (**Fig. 4f and Supplementary. Fig. 4c**). We then assembled pseudo-bulk

transcriptomic profiles for each subtype and ranked 1,019 CCLE cell lines according

to transcriptomic similarity in a subtype-specific manner. As expected, MDA-PCa-2b,

VCaP, and LNCaP showed the highest similarity to ARPC, while NCI-H660 exhibited

a high similarity to NEPC (**Supplementary Fig. 4d-e**). Surprisingly, for the MSPC

subtype, although PC3 ranked highest among prostate cancer cell lines, its overall

similarity to MSPC malignant cells remained modest (ranked 260 among 1,019 CCLE

cell lines) (**Fig. 4g**). Moreover, the transcriptomic similarity between individual

MSPC malignant cells and PC3 was significantly lower than that between ARPC

malignant cells and VCaP, suggesting limited fidelity of PC3 as a model of MSPC

(**Fig. 4h**).

Interestingly, we found MSPC malignant cells expressed basal lineage markers such

as *CAV2* and *TP63* (**Supplementary Fig. 4f**), suggesting their basal differentiation.

Consistent with this, when compared with three normal prostate epithelial subtypes

(luminal, basal and neuroendocrine), they showed the highest similarity to basal

epithelial cells (**Fig. 4i and Supplementary. Fig. 4g**). However, PC3 lacked

expression of key basal markers (such as *TP63*), which may explain its low

transcriptomic similarity with MSPC malignant cells **(Supplementary Fig. 4h).**

Taken together, our findings highlight both the strength and limitation of PC3 cell

line.

PC3 exhibits molecular features consistent with stemness, epithelial–mesenchymal

transition, and cellular plasticity, and among currently available prostate cancer cell

lines, it most closely resembles the MSPC subtype. However, its overall

transcriptomic similarity to primary MSPC malignant cells remains limited. Further

studies are warranted to identify or develop more representative cell line models that

faithfully recapitulate the molecular characteristics of MSPC prostate cancer.

Patient-derived organoids better resemble the transcriptome of MSPC than PC3

Since PC3 does not fully recapitulate the transcriptomic features of MSPC, we next

investigated other type of models that could be more representative. We focus on two

categories of models: engineered prostate cancer cell lines and patient-derived

organoids.

Previous studies have reported that AR inhibition by enzalutamide or *TP53/RB1*

double-knockout can promote LNCaP (a hypermutated cell line of ARPC) into MSPC

state[42, 43]. Therefore, we included these two engineered LNCaP-derived models in our

analysis. In addition, we generated a VCaP-derived cell line by overexpressing *MET*

(VCaP-*MET*-oe), as *MET* has been reported to promote EMT and sustain stemness in

cancer cells[44, 45, 46]. As expected, bulk RNA-seq analysis suggested that *MET*

overexpression induced upregulation of *CD44* (log2fc = 0.251, p-value = 0.028) and

increased *JAK/STAT* pathway activity, while downregulating luminal markers *KLK2*,

KLK3, and *KLK4*, suggesting the occurrence of de-differentiation (**Fig. 5a-b and Supplementary Table 5**).

We next computed the transcriptomic similarity with MSPC malignant cells for the above three engineered cell lines, respectively (**Fig. 5c-e**). Surprisingly, all of them exhibited lower similarity to MSPC than PC3. Actually, when comparing the transcriptomic profiles of these engineered cell lines to the 1,019 CCLE cell lines, the highest correlations were consistently observed with their respective parental cell lines (**Supplementary Fig. 5a–c)**, suggesting that these manipulations did not fully trigger a lineage transition. Given that MSPC malignant cells exhibit basal differentiation, this likely explains why the engineered cell lines failed to recapitulate the MSPC transcriptional landscape.

We next explored whether patient-derived organoids could better recapitulate the MSPC transcriptome. We computed the transcriptomic similarity between ten PDOs (all of which were claimed as MSPC according to Tang's study)[47] and MSPC malignant cells (**Fig. 5f**). Notably, we found six PDOs had higher similarity to MSPC malignant cells than PC3; of them, MSKPCa12 demonstrated the highest correlation and appeared to be the most representative (**Supplementary. Fig. 5d**). Further lineage analysis suggested that this organoid exhibits a basal lineage (**Supplementary Fig. 5e**), with expression of several basal specific markers such as *TP63*, *CAV2*, and *KRT5*

(**Supplementary. Fig. 5f-h**).

These efforts to engineer existing prostate cancer cell lines motivated us to explore

alternative models. In recent years, patient-derived organoids have gained increasing

attention as promising platforms for modeling tumor heterogeneity and

subtype-specific biology. Their ability to maintain key molecular and

histopathological features of the original tumors underscores their potential value in

studies of MSPC biology and therapeutic response.

Discussion

Cell lines are widely utilized in cancer research, particularly as models for metastatic cancers. Prior studies have demonstrated that cell lines derived from the same tumor type exhibit considerable molecular heterogeneity, with certain cell lines more accurately recapitulating the characteristics of metastatic tumors than others[10, 33, 48, 49]. This underscores the critical importance of selecting cell line models that most faithfully represent the biology of metastatic cancers. Our study proposed useful guidance for cell line selection in the study of metastatic prostate cancer.

Adenocarcinoma is the predominant histological subtype of prostate cancer[50, 51]. Based on our multi-omics analysis results, VCaP appears to be the most suitable model for studying metastatic prostate cancer according to its relatively higher concordance with metastatic prostate adenocarcinoma patient samples in multiple aspects (for example, the high transcriptomic similarity with patient samples, and the highest level of *AR* amplification). However, according to Pubmed search results, VCaP is rarely used in metastasis-related research, suggesting a gap between model suitability and actual utilization in the field.

Although VCaP represents a suitable model for metastatic prostate adenocarcinoma, it may not be appropriate for all contexts. One particular situation is hypermutated patient samples. Our analysis suggests that the microenvironment in hypermutated

prostate cancer samples differs markedly from that in non-hypermutated tumors,

particularly with respect to increased immune infiltration. This indicates that the

hypermutation phenomenon should be considered when selecting cell line models.

Given its high mutation burden and strong transcriptomic similarity to metastatic

prostate adenocarcinoma samples, LNCaP was identified as a promising model for

studying hypermutated prostate cancer.

Another situation is to research AR resistance. We found that the prostate cancer cell

lines LNCaP and MDA-PCa-2b harbor *AR* mutation hotspots associated with clinical

resistance, showing a certain degree of concordance with patient observations.

Specifically, MDA-PCa-2b carries both the T878A and L702H mutations, which are

linked to resistance to flutamide and glucocorticoids, respectively. The T878A

mutation was also detected in LNCaP, while 22RV1 harbors another recurrent *AR*

mutation hotspot, H875Y. Notably, despite harboring these resistance-associated

mutations, all three cell lines remain AR-dependent, suggesting that they are valuable

in vitro models for studying AR-driven resistance mechanisms and evaluating novel

antiandrogen therapies.

According to our PubMed search results, PC3 is the most widely used prostate cancer

cell line; however, it exhibits low transcriptomic similarity to metastatic prostate

adenocarcinoma. Although further analysis classified PC3 within the MSPC subtype,

it does not faithfully model MSPC at the transcriptomic level, either. This discrepancy

may arise from its lack of basal lineage marker, whereas MSPC malignant cells

typically show high expression of basal markers and share transcriptomic features

with normal prostate basal cells. Surprisingly, several bladder cancer cell lines (such

as RT4 and KMBC2, derived from urinary tract) display greater transcriptomic

resemblance to MSPC malignant cells. While this observation may reflect random

variation, we favor the interpretation of a true underlying biological similarity, as

urinary tract–derived cell lines consistently exhibit higher basal marker expression

than PC3. Collectively, these findings suggest that alternative cell lines may better

recapitulate the molecular characteristics of the MSPC subtype and serve as more

effective models for elucidating its underlying biology.

Mixed subtypes have been reported in prostate cancer patients[41, 52, 53, 54]. These

findings underscore the importance of employing multiple cell lines simultaneously

when investigating mixed subtype phenomena, in order to more accurately capture

and model intratumoral heterogeneity. According to scRNA-seq analysis of metastatic

prostate cancer samples, the majority of malignant cells exhibited an ARPC

phenotype, whereas MSPC cells were relatively rare. This observation raises

important biological questions regarding the role of MSPC malignant cells in

metastasis. Specifically, it remains unclear whether MSPC cells facilitate the

dissemination of ARPC cells, or whether ARPC and MSPC represent independently

metastasizing malignant populations. Further studies are needed to elucidate the

functional interplay between these subtypes and their respective contributions to

metastatic progression.

MSPC phenotype remains insufficiently characterized, and existing prostate cancer

cell lines do not reliably model this subtype. In this study, we attempted to induce an

MSPC phenotype in ARPC-derived cell lines. However, this strategy resulted only in

the partial acquisition of stem-like features. Notably, certain prostate organoids with

intrinsic stem-like properties appear to represent more appropriate models for

recapitulating the MSPC subtype. Future research should focus on identifying or

engineering prostate cancer cell lines that more faithfully reproduce the molecular and

transcriptomic characteristics of MSPC.

In summary, by leveraging the molecular characteristics of metastatic prostate cancer

and prostate cancer cell lines, we identified suitable metastatic prostate cancer cell

lines for specific research applications. We anticipate that the adoption of more

appropriate metastatic prostate cancer cell lines will enhance the translational

relevance of *in vitro* studies and accelerate the development of clinically meaningful

therapeutic strategies.

Materials and Methods

Selection of genes used for genomic profile comparison

The somatic mutation data of metastatic and primary prostate cancer used in this study were from SU2C and TCGA, respectively. In our analysis, non-adenocarcinoma samples were excluded. Given a gene, we performed Fisher's exact to evaluate whether its mutation frequency is significantly different between primary and metastatic prostate cancer samples. Since SU2C contains metastatic samples from three sites (liver, bone, and lymph node), the analysis was performed in site-specific manner. Only genes which are differently mutated (adjusted p-value < 0.05) in at least two sites were considered as differentially mutated genes.

Genes whose mutation frequency exceeded 5% in at least two metastatic sites (in the SU2C cohort) were defined as highly mutated genes. The non-synonymous mutations occurring in at least three SU2C samples were defined as mutation hotspot.

Identification of differential CNV genes between metastatic and primary prostate cancer

We downloaded CNV segment data and then utilized CNVTools[55] to generate gene-level copy number. For each gene, the Wilcoxon rank-sum test was applied to compare its CNV profile between SU2C and TCGA samples. Genes which are differently copy number variation (adjusted p-value < 0.05) in at least two sites and an

442 absolute median CNV difference ≥ 0.3 were considered as differentially CNV genes.

Transcriptome correlation analysis with bulk RNA-seq and scRNA-seq data

In our previous research, we identified 1,000 genes which are highly varied across

CCLE cell lines and proved their utility in transcriptome correlation analysis[12]. Given

a cell line and a patient sample (or single cell), we defined their transcriptome

similarity as the Spearman rank correlation across the 1,000 genes. The transcriptome

similarity between a cell line and several patient samples (or single cells) was

computed as the median correlation value.

Differential gene expression analysis

453 DESeq2[56] was used to identify differentially expressed genes (adjusted p-value < 0.05

454 and abs(log2FC) > 1) and the geneset enrichment analysis was conducted using

455 Enrichr[57]. Terms with a p-value < 0.05 were considered statistically significant.

Pubmed search

458 Our Pubmed search was performed on March 11[th], 2025. For each prostate cancer cell

459 line, we searched the database using the keyword "[cell line name] + metastasis" and

460 counted the number of returned citations.

Analysis of scRNA-seq data

In the analysis of CRPC scRNA-seq dataset, we used inferCNV[58, 59] to infer copy

number variation of epithelial cells (immune cells as reference). CNV values were

extracted and subjected to k-means clustering to identify malignant cells. The marker

genes used in our analysis are: *PTPRC* (immune cells) and *EPCAM* (epithelial cells).

Lineage analysis

We used a normal prostate epithelial scRNA-seq dataset to identify three epithelial

cell types: luminal (*KLK2*, *KLK4*, *ACPP*), basal (*KRT5*, *TP63*, *KRT17*, *KRT15*), and

neuroendocrine (*ASCL1*, *FOXA2*, *MYCN*, *POU3F2*, *SIAH2*, *NCAM1*, *CHGA*, *CHGB*,

SYP, *ENO2*)[60, 61]. For each epithelial cell type, we generated pseudo-bulk

transcriptomes to create a centroid matrix. We then identified the top 5,000 highly

variable genes across the three cell types and used these genes to calculate

transcriptomic similarity between each malignant cell and the reference centroids.

Each malignant cell was assigned to the epithelial lineage with the highest similarity

score.

Software tools and statistical methods

All of the analysis was conducted in R. The ggplot2[62] and ComplexHeatmap[63]

packages were used for data visualization. Seurat[64] package was used for processing

scRNA-seq data. GSVA was used for ssGSEA analysis[65, 66]. If not specified, the two-sided

Wilcoxon rank-sum test was used to compute p-value and Benjamini-Hochberg

procedure was applied to compute adjusted p-value.

Datasets

The genomic and transcriptomic data of CCLE cell lines were obtained from the

DepMap data portal (https://depmap.org/portal/). The somatic mutation and copy

number variation data of TCGA and SU2C samples were downloaded from

cBioPortal (https://www.cbioportal.org/)[67, 68]. The bulk RNA-seq data of TCGA

samples was downloaded from UCSC Xena data portal (https://xena.ucsc.edu/)[69]. The

bulk RNA-seq data of metastatic prostate adenocarcinoma patients were from

MET500 cohort. The prostate adenocarcinoma scRNA-seq data was downloaded from

Single Cell Portal (https://singlecell.broadinstitute.org/single_cell/study/SCP1244).

The scRNA-seq data of normal human prostate, cell lines and organoids were

downloaded from GEO (accession number: GSE157220, GSE117403, GSE175975,

GSE162225 and GSE181374)[42, 43, 61, 70]. The scRNA-seq data of CRPC was

downloaded from SRA (accession number: PRJNA699369)[71].

Data Availability

The authors declare that all data supporting the findings of this study are available

within the article and its supplementary information files or from the corresponding

author upon reasonable request.

Acknowledgments

The research is supported by National Natural Science Foundation of China (Fund 32370715), Science Fund Program for Distinguished Oversea Young Scholars of Shandong Province (2023HWYQ-015), and the Cheeloo Young Scholar Program of Shandong University. The content is solely the responsibility of the authors and does not necessarily represent the official views of sponsors.

Authors' contribution

X.Y.L. and K.L. conceived the study. X.Y.L. performed the majority of computational analysis, Y.G.W. performed all the experimental assays, and K.L. supervised the study. All authors contributed to writing, reviewing, and editing the manuscript and approved the manuscript.

Competing Interests

The researchers declare no competing interests.

References

1.   Kaur G, Dufour JM. Cell lines: Valuable tools or useless artifacts. *Spermatogenesis* **2**, 1-5 (2012).

2.   Goodspeed A, Heiser LM, Gray JW, Costello JC. Tumor-Derived Cell Lines as Molecular Models of Cancer Pharmacogenomics. *Molecular cancer research : MCR* **14**, 3-13 (2016).

3.   Bose B, Bozdag S. Identifying cell lines across pan-cancer to be used in precl inical research as a proxy for patient tumor samples. *Commun Biol* **7**, 1101 (2 024).

4.   Dutil J, Chen Z, Monteiro AN, Teer JK, Eschrich SA. An Interactive Resource to Probe Genetic Diversity and Estimated Ancestry in Cancer Cell Lines. *Canc er research* **79**, 1263-1273 (2019).

5.   Mouradov D*, et al.* Colorectal cancer cell lines are representative models of th e main molecular subtypes of primary cancer. *Cancer research* **74**, 3238-3247 (2014).

6.   Gillet JP, Varma S, Gottesman MM. The clinical relevance of cancer cell lines. *Journal of the National Cancer Institute* **105**, 452-458 (2013).

7.   Weinstein JN. Drug discovery: Cell lines battle cancer. *Nature* **483**, 544-545 (2 012).

8.   Ertel A, Verghese A, Byers SW, Ochs M, Tozeren A. Pathway-specific differen ces between tumor cell lines and normal and tumor tissue cells. *Molecular can cer* **5**, 55 (2006).

9.   Salvadores M, Fuster-Tormo F, Supek F. Matching cell lines with cancer type and subtype of origin via mutational, epigenomic, and transcriptomic patterns. *Science advances* **6**, (2020).

10.   Domcke S, Sinha R, Levine DA, Sander C, Schultz N. Evaluating cell lines a s tumour models by comparison of genomic profiles. *Nat Commun* **4**, 2126 (2

013).

11.   Yu K, *et al*. Comprehensive transcriptomic analysis of cell lines as models of
564   primary tumors across 22 tumor types. *Nat Commun* **10**, 3574 (2019).

566   12.   Liu K, *et al*. Evaluating cell lines as models for metastatic breast cancer throu
567   gh integrative analysis of genomic data. *Nat Commun* **10**, 2138 (2019).

569   13.   Xin S, *et al*. ScRNA-seq revealed an immunosuppression state and tumor micr
570   oenvironment heterogeneity related to lymph node metastasis in prostate cancer.
571   *Exp Hematol Oncol* **12**, 49 (2023).

573   14.   Rebello RJ, *et al*. Prostate cancer. *Nature reviews Disease primers* **7**, 9 (2021).

575   15.   Jin H, *et al*. Systematic transcriptional analysis of human cell lines for gene e
576   xpression landscape and tumor representation. *Nat Commun* **14**, 5417 (2023).

578   16.   Peng D, *et al*. Evaluating the transcriptional fidelity of cancer models. *Genome*
579   *medicine* **13**, 73 (2021).

581   17.   Hao M, *et al*. HIC1 loss promotes prostate cancer metastasis by triggering epit
582   helial-mesenchymal transition. *The Journal of pathology* **242**, 409-420 (2017).

584   18.   Ghandi M, *et al*. Next-generation characterization of the Cancer Cell Line Enc
585   yclopedia. *Nature* **569**, 503-508 (2019).

587   19.   Hernández Borrero LJ, El-Deiry WS. Tumor suppressor p53: Biology, signaling
588   pathways, and therapeutic targeting. *Biochimica et biophysica acta Reviews on*
589   *cancer* **1876**, 188556 (2021).

591   20.   Hunter I, Jamieson C, McEwan IJ. The androgen receptor amino-terminal dom
592   ain: structure, function and therapeutic potential. *Endocrine oncology (Bristol, E*
593   *ngland)* **5**, e240061 (2025).

595   21.   Robinson D, *et al*. Integrative clinical genomics of advanced prostate cancer. *C*
596   *ell* **161**, 1215-1228 (2015).

22. He MX, *et al.* Transcriptional mediators of treatment resistance in lethal prosta
te cancer. *Nat Med* **27**, 426-433 (2021).

23. Augello MA, Den RB, Knudsen KE. AR function in promoting metastatic pros
tate cancer. *Cancer metastasis reviews* **33**, 399-411 (2014).

24. Heinlein CA, Chang C. Androgen receptor in prostate cancer. *Endocrine review
s* **25**, 276-308 (2004).

25. Lin A, Zhang J, Luo P. Crosstalk Between the MSI Status and Tumor Microe
nvironment in Colorectal Cancer. *Frontiers in immunology* **11**, 2039 (2020).

26. Seo MK, Kang H, Kim S. Tumor microenvironment-aware, single-transcriptome
prediction of microsatellite instability in colorectal cancer using meta-analysis.
Scientific reports **12**, 6283 (2022).

27. Ferreira AM, *et al.* High frequency of RPL22 mutations in microsatellite-unsta
ble colorectal and endometrial tumors. *Human mutation* **35**, 1442-1445 (2014).

28. Weinstein HNW, *et al.* RPL22 is a tumor suppressor in MSI-high cancers and
a splicing regulator of MDM4. *Cell Rep* **43**, 114622 (2024).

29. Cortes-Ciriano I, Lee S, Park WY, Kim TM, Park PJ. A molecular portrait of
microsatellite instability across multiple cancers. *Nat Commun* **8**, 15180 (2017).

30. Lambert AW, Zhang Y, Weinberg RA. Cell-intrinsic and microenvironmental det
erminants of metastatic colonization. *Nature cell biology* **26**, 687-697 (2024).

31. Biray Avci C, Goker Bagca B, Nikanfar M, Takanlou LS, Takanlou MS, Nour
azarian A. Tumor microenvironment and cancer metastasis: molecular mechanis
ms and therapeutic implications. *Frontiers in pharmacology* **15**, 1442888 (202
4).

32. Joyce JA, Pollard JW. Microenvironmental regulation of metastasis. *Nature revi
ews Cancer* **9**, 239-252 (2009).

33. Chen B, Sirota M, Fan-Minogue H, Hadley D, Butte AJ. Relating hepatocellul
ar carcinoma tumor samples and cell lines using gene expression data in transl
ational research. *BMC medical genomics* **8 Suppl 2**, S5 (2015).

34. Moya L*, et al.* Characterisation of cell lines derived from prostate cancer patie
nts with localised disease. *Prostate Cancer Prostatic Dis* **26**, 614-624 (2023).

35. Molter CW, Muszynski EF, Tao Y, Trivedi T, Clouvel A, Ehrlicher AJ. Prostate
cancer cells of increasing metastatic potential exhibit diverse contractile forces,
cell stiffness, and motility in a microenvironment stiffness-dependent manner.
Frontiers in cell and developmental biology **10**, 932510 (2022).

36. Erickson A*, et al.* A Systematic Review of Prostate Cancer Heterogeneity: Und
erstanding the Clonal Ancestry of Multifocal Disease. *European urology oncolo
gy* **4**, 358-369 (2021).

37. Løvf M*, et al.* Multifocal Primary Prostate Cancer Exhibits High Degree of G
enomic Heterogeneity. *European urology* **75**, 498-505 (2019).

38. Song H*, et al.* Single-cell analysis of human primary prostate cancer reveals th
e heterogeneity of tumor-associated epithelial cell states. *Nat Commun* **13**, 141
(2022).

39. Guo W*, et al.* Single-cell transcriptomics identifies a distinct luminal progenitor
cell type in distal prostate invagination tips. *Nature genetics* **52**, 908-918 (202
0).

40. Zhang D, Zhao S, Li X, Kirk JS, Tang DG. Prostate Luminal Progenitor Cells
in Development and Cancer. *Trends in cancer* **4**, 769-783 (2018).

41. Chan JM*, et al.* Lineage plasticity in prostate cancer depends on JAK/STAT in
flammatory signaling. *Science* **377**, 1180-1191 (2022).

42. Han H*, et al.* Mesenchymal and stem-like prostate cancer linked to therapy-ind
uced lineage plasticity and metastasis. *Cell Rep* **39**, 110595 (2022).

43. Deng S*, et al.* Ectopic JAK-STAT activation enables the transition to a stem-li

ke and multilineage state conferring AR-targeted therapy resistance. *Nat Cancer*
3, 1071-1087 (2022).

44. Jeon HM, Lee J. MET: roles in epithelial-mesenchymal transition and cancer st
emness. *Annals of translational medicine* **5**, 5 (2017).

45. van Leenders GJ*, et al.* Activation of c-MET induces a stem-like phenotype in
human prostate cancer. *PloS one* **6**, e26753 (2011).

46. Dong J, Li J, Li Y, Ma Z, Yu Y, Wang CY. Transcriptional super-enhancers co
ntrol cancer stemness and metastasis genes in squamous cell carcinoma. *Nat C*
ommun **12**, 3974 (2021).

47. Tang F*, et al.* Chromatin profiles classify castration-resistant prostate cancers s
uggesting therapeutic targets. *Science* **376**, eabe1505 (2022).

48. Virtanen C*, et al.* Integrated classification of lung tumors and cell lines by ex
pression profiling. *Proc Natl Acad Sci U S A* **99**, 12357-12362 (2002).

49. Sinha R*, et al.* Analysis of renal cancer cell lines from two major resources e
nables genomics-guided cell line selection. *Nat Commun* **8**, 15165 (2017).

50. Xing P*, et al.* Treatment strategies and drug resistance mechanisms in adenocar
cinoma of different organs. *Drug resistance updates : reviews and commentarie*
s in antimicrobial and anticancer chemotherapy **71**, 101002 (2023).

51. Raychaudhuri R, Lin DW, Montgomery RB. Prostate Cancer: A Review. *Jama*
333, 1433-1446 (2025).

52. Cejas P*, et al.* Subtype heterogeneity and epigenetic convergence in neuroendo
crine prostate cancer. *Nat Commun* **12**, 5775 (2021).

53. Bluemn EG*, et al.* Androgen Receptor Pathway-Independent Prostate Cancer Is
Sustained through FGF Signaling. *Cancer Cell* **32**, 474-489.e476 (2017).

54. Lindh C*, et al.* Ductal and acinar components of mixed prostatic adenocarcino
ma frequently have a common clonal origin. *The Prostate* **82**, 576-583 (2022).

55.   The CNTools package. https://bioconductor.org/packages/release/bioc/html/CNTool
710   s.html.).

712   56.   Love MI, Huber W, Anders S. Moderated estimation of fold change and disper
713   sion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

715   57.   Kuleshov MV*, et al.* Enrichr: a comprehensive gene set enrichment analysis w
716   eb server 2016 update. *Nucleic Acids Res* **44**, W90-97 (2016).

718   58.   Patel AP*, et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in pr
719   imary glioblastoma. *Science* **344**, 1396-1401 (2014).

721   59.   Puram SV*, et al.* Single-Cell Transcriptomic Analysis of Primary and Metastati
722   c Tumor Ecosystems in Head and Neck Cancer. *Cell* **171**, 1611-1624.e1624 (2
723   017).

725   60.   Dong B*, et al.* Single-cell analysis supports a luminal-neuroendocrine transdiffe
726   rentiation in human prostate cancer. *Commun Biol* **3**, 778 (2020).

728   61.   Henry GH*, et al.* A Cellular Anatomy of the Normal Adult Human Prostate an
729   d Prostatic Urethra. *Cell Rep* **25**, 3530-3542.e3535 (2018).

731   62.   The ggplot2 package. https://cran.r-project.org/web/packages/ggplot2/index.html.).

733   63.   The ComplexHeatmap package. https://bioconductor.org/packages/release/bioc/htm
734   l/ComplexHeatmap.html.).

736   64.   Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of
737   single-cell gene expression data. *Nat Biotechnol* **33**, 495-502 (2015).

739   65.   Subramanian A*, et al.* Gene set enrichment analysis: a knowledge-based approa
740   ch for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S*
741   *A* **102**, 15545-15550 (2005).

743   66.   Hänzelmann S, Castelo R, Guinney J. GSVA: gene set variation analysis for m
744   icroarray and RNA-seq data. *BMC bioinformatics* **14**, 7 (2013).

67.   Gao J*, et al.* Integrative analysis of complex cancer genomics and clinical prof

iles using the cBioPortal. *Science signaling* **6**, pl1 (2013).

68.   Cerami E*, et al.* The cBio cancer genomics portal: an open platform for explo

ring multidimensional cancer genomics data. *Cancer discovery* **2**, 401-404 (201

2).

69.   Vivian J*, et al.* Toil enables reproducible, open source, big biomedical data an

alyses. *Nat Biotechnol* **35**, 314-316 (2017).

70.   Kinker GS*, et al.* Pan-cancer single-cell RNA-seq identifies recurring programs

of cellular heterogeneity. *Nature genetics* **52**, 1208-1218 (2020).

71.   Cheng Q*, et al.* Pre-existing Castration-resistant Prostate Cancer-like Cells in P

rimary Prostate Cancer Promote Resistance to Hormonal Therapy. *European uro*

logy **81**, 446-455 (2022).