Title: Dynamic task-based reorganization of sensory-biased and supramodal working memory networks in the human cerebral cortex

Authors: Thomas Possidente*, Vaibhav Tripathi*, Joseph T. McGuire, David C. Somers

Department of Psychological and Brain Sciences, Boston University, Boston, Massachusetts 02215
*Authors contributed equally
Corresponding author: David C. Somers, somers@bu.edu

Keywords: functional connectivity, working memory, visual processing, auditory processing, domain-general, frontal cortex

Abstract:

Sensory working memory is supported by a distributed set of brain regions. Visual working memory and auditory working memory recruit different networks, which include posterior sensory cortices and multiple sensory-biased regions interleaved across frontal cortex. Additionally, a fronto-parietal network of supramodal regions is robustly recruited by working memory in both modalities. Here we examined functional connectivity among visual-biased, auditory-biased, and supramodal working memory networks by collecting fMRI data from 24 human participants (aged 18-43; 11 men and 13 women). Replicating and extending prior work from our laboratory (Tobyne et al., 2017; Noyce et al., 2022), resting-state fMRI (N=18) analysis revealed that the visual-biased WM network exhibits stronger functional connectivity with supramodal regions than does the auditory-biased network. These findings point to a potential functional asymmetry between visual and auditory working memory mechanisms; visual cognition may have more direct access to supramodal WM brain structures. Therefore, we hypothesized that auditory working memory task performance would evoke more robust changes in functional connectivity than would visual working memory task performance. To test this hypothesis, we examined changes in functional connectivity that occur when participants (N=21) perform visual or auditory working memory tasks equated for difficulty. Auditory working memory produced substantial network changes that increased the connectivity of the auditory-biased network to supramodal regions and visual-biased frontal regions. At the same time, connectivity from posterior visual areas to supramodal and frontal visual regions decreased. In contrast, the visual task produced more modest changes in functional connectivity. Moreover, we observed that the strength of auditory-biased network connectivity changes correlated with subject behavioral performance on the auditory task, with higher working memory precision observed in participants with greater connectivity changes. Therefore, the visual working memory network appears to have more robust connectivity to supramodal working memory cortical structures at rest, but during auditory working memory, dynamic network changes permit the auditory working memory network to increase connectivity to the supramodal and frontal visual working memory networks.

Introduction

Human working memory (WM) is supported both by 'content-specific' mechanisms that vary with the type of information held in memory and by 'multiple-demand' or 'central executive' mechanisms whose operation appears independent of content (e.g., Baddeley & Hitch, 1974; Duncan & Owen, 2000). The interaction between these subsystems is an important, understudied aspect of WM and is the focus of the current study. Sensory WM for visual stimuli and auditory stimuli has been shown to recruit different modality-specific structures in occipito-parietal and temporal lobes, respectively (Bettencourt & Xu, 2016; Christophel et al., 2012, 2017; Harrison & Tong, 2009; Linke et al., 2011; Serences et al., 2009; Uluç et al., 2018), while both modalities recruit broad regions of lateral frontal cortex which could be considered 'supramodal' or 'multiple-demand' in their functionality (Assem et al., 2020; Christophel et al., 2017; Crittenden & Duncan, 2014; D'Esposito et al., 1999; Duncan, 2001; Duncan & Owen, 2000; Ester et al., 2015). Additionally, multiple regions within lateral frontal cortex are biased toward either visual or auditory WM (Assem et al., 2022; Mayer et al., 2017; Michalka et al., 2015; Noyce et al., 2017, 2022; Tobyne et al., 2017). These small regions were revealed by precision mapping of lateral frontal cortex in individual subjects whereas prior group-averaged analyses had blurred these regions together and mischaracterized them as 'multiple-demand'. Moreover, these sensory-biased regions lie in proximity to true supramodal or multiple-demand WM regions (Assem et al., 2022). This motivates new questions about how these sensory-biased and domain-general areas are topographically organized and how they might functionally interact. Here, we examine how sensory-biased and supramodal WM regions interact during visual WM and auditory WM tasks and during rest.

Assem et al. (2022) suggests that sensory-biased regions surround and potentially feed information into the domain-general areas, while Noyce et al. (2017) indicates that visual-biased areas, but not auditory-biased areas, contain some domain-general response properties while still being functionally separable from fully domain-general areas. This and other work from our laboratory (Noyce et al., 2022; Tobyne et al., 2017, 2025) points to a potential functional asymmetry between visual and auditory WM mechanisms; visual cognition structures in frontal cortex appear more closely integrated with supramodal WM brain structures (regions that respond to both auditory and visual WM demands) than are auditory cognition structures in frontal cortex. If this is true, the next natural question is how it is possible for auditory WM to function effectively with relatively weaker coordination with supramodal networks. Therefore, we hypothesize that visual and auditory WM task performance evoke asymmetric changes in functional connectivity between visual-biased, auditory-biased, and supramodal networks. Several non-mutually exclusive mechanisms are plausible to help explain how auditory WM performance acquires the necessary resources to function effectively despite weaker coordination with supramodal WM networks at rest: auditory WM performance may 1) increase connectivity between auditory and supramodal networks, 2) decrease competing connectivity between visual and supramodal networks, 3) increase connectivity within the auditory network components, and/or 4) increase cross modal connectivity between visual frontal and auditory frontal networks. An increase in connectivity in the auditory network components (*Mechanism 3*) could reflect increased efficiency within the auditory system. An increase in cross-modal connectivity

(*Mechanism 4*) would increase direct communication between the auditory-biased regions and
the frontal visual-biased, which exhibit partial supramodal WM properties, and could indirectly
increase communication with supramodal network regions based on the resting connection
between visual-biased and supramodal regions.

The present study examines these possibilities by presenting in-depth resting state and task-
based functional connectivity analyses of sensory-biased perception and WM regions, along with
supramodal frontal regions. Building on our prior work (Michalka et al., 2015; Noyce et al.,
2017, 2022; Tobyne et al., 2025), we define a large number of visual-biased, auditory-biased, and
supramodal WM regions of interest (ROIs) in the frontal cortex of individual subjects to drive
our analysis. We replicate the finding that the supramodal network is more strongly connected
with the visual-biased network than with the auditory-biased network at rest and additionally
find that visual-biased, auditory-biased, and supramodal networks change their connectivity
structure differentially during visual and auditory WM tasks. We show that the auditory-biased
network overcomes its resting-state connectivity disadvantage by recruiting both supramodal and
visual-biased networks, increasing connectivity within the auditory network, and by effecting a
decrease in competing connectivity among visual frontal, visual posterior, and supramodal
networks. We also show that task-based changes in functional connectivity between a subset of
these networks are predictive of behavioral WM precision, indicating that increased recruitment
by the auditory network leads to increased auditory WM performance.

Methods

Participants

Twenty-four participants (aged 18 to 43; 11 men and 13 women [self-identified]) were
recruited from the Boston University community. All procedures in this study were approved by
the Institutional Review Board of Boston University. Each participant gave written informed
consent to participate in the study and received monetary compensation. All participants
indicated that they had normal or corrected to normal vision and normal hearing. Two authors
participated in both the task and resting state scans (T. P., V. T.). All 24 subjects performed visual
and auditory working memory and sensorimotor control tasks during fMRI for this study.
Twenty-two of these also participated in resting-state scans. Three participants were rejected
from all analyses - two for ROI identification problems, one for head movement – leaving
twenty-one participants (19 with resting state scans). One additional participant was rejected
from only the resting state analysis for head movement, leaving eighteen participants. Rejection
criteria details are given below.

Working Memory Paradigm

All participants performed 4 tasks in the scanner: 2-back visual working memory, 2-back
auditory working memory, visual sensorimotor control, and auditory sensorimotor control. The
stimuli and general paradigm are shown in Figure 1 below. Stimuli were presented for 1 second
each, with a 1 second gap between stimuli; only one modality was presented at a time. The visual

stimuli were Gabor patches with spatial frequencies in the range of 1 to 6 cycles per degree. In
the visual 2-back task, subjects judged whether the spatial frequency of the stimulus matched
that presented two stimuli previously. Gabor orientation was varied randomly and was irrelevant
to the match judgement. In the auditory 2-back task, the stimuli were auditory 'warbles' with
either a 500 or 1000 Hz base pitch modulated by an oscillatory amplitude envelope with
frequencies ranging from 3 Hz to 20 Hz. Subjects judged whether the envelope frequency
matched that from two stimuli earlier. Pitch was varied randomly and was irrelevant to the match
judgement. For both auditory and visual stimuli, the trials consisted of high frequency stimuli
(i.e., odd numbered trials) alternated with low frequency trials (i.e., even numbered trials); this
afforded the opportunity to make the matching task difficult while also presenting a broader
range of stimuli. To assure that the 2-back tasks were perceptually difficult, the percentage
change in the modulatory frequency for non-match trials was adjusted using independent
staircasing procedures for visual and auditory stimuli. Individual thresholds were identified for
each subject in a behavioral session prior to scanning and were updated during the scans.
Individual differences in task performance were quantified by the staircase threshold parameters.
Smaller thresholds imply more precise performance, even though task accuracy was equated
across subjects. These threshold measures were employed for individual-subject measures
comparing functional connectivity changes and behavioral changes.

During the sensorimotor control tasks, auditory or visual stimuli were presented without
2-back repeats, and participants were instructed not to do the task but instead to make a random
button press at the end of each trial. The relevant blocks in a single run consisted of 2 blocks of
visual WM, 2 blocks of auditory WM, 1 block of visual sensorimotor control, 1 block of auditory
sensorimotor control, and 1 fixation condition block without stimuli or responses. Each block
lasted 32s and contained 16 trials. Participants were given feedback indicating correct and
incorrect trials. During the sensorimotor control conditions, feedback was given if they had
pressed the key or not. For participants used in the task analyses, 19 had four runs, one had three,
and one had six.

Figure 1. Visual and auditory 2-back WM task summaries. For the visual WM task, participants were asked whether the current Gabor filter spatial frequency matched the one presented two stimuli previously. For the auditory WM task participants were asked whether the warble envelope volume modulation frequency matched the one presented two stimuli previously. The two-back tasks were effectively two interleaved threshold tasks, with odd-numbered stimuli of low-frequency and even-numbered trials of high-frequency (or vice versa, randomly across blocks). Task performance required participants to maintain a high-precision representation of both the most recent even and odd numbered trials.

Resting State Paradigm

Participants were presented with a white fixation cross on a gray background and asked to fixate while not doing any particular task. Resting state runs lasted 6 minutes each and for participants used in the resting state analysis, 14 had three runs and 4 had one run.

MR Imaging

All scans were performed at Boston University's Cognitive Neuroimaging Center on a Siemens 3-Tesla Prisma scanner with a 64-channel head coil. Structural images were collected for each participant with T1-weighted high-resolution magnetization-prepared rapid gradient multi-echo sequence (MP-RAGE, (van der Kouwe et al., 2008)) (1mm isotropic voxels, 176 slices, repetition time (TR) = 2530ms, echo times (TE) = 1.69, 3.55, 5.41 and 7.27ms, flip angle (FA) = 7°, field of view (FOV) = 256mm, GRAPPA (iPAT, [Griswold et al., 2002]) acceleration=3, TA=4min 23s). All blood-oxygen-level-dependent (BOLD) data were collected via T2*-weighted echo-planar imaging (EPI) pulse sequence that employed multiband RF pulses and simultaneous multi-slice (SMS) acquisition (Cauley et al., 2014; Feinberg et al., 2010; Moeller et al., 2010; Setsompop et al., 2012; Xu et al., 2013). Task fMRI (69 slices, TR=2000ms,

TE=35ms, FA=80°, 2.2mm isotropic voxels, FOV=207mm, SMS=3, TA=7min 46s) and resting
state fMRI (64 interleaved slices, TR=650ms, TE=34.8ms, FA=52°, 2.3mm isotropic voxels,
FOV=207mm, SMS=8, TA=6min 9s) were also acquired. One participant's resting state data was
collected using the task fMRI sequence parameters, which was accounted for in the
preprocessing pipeline. Pairs of reversed phase-encoded (anterior-posterior and posterior-
anterior) spin echo field maps were also acquired for subsequent EPI de-warping.

MRI Preprocessing

For ROI localization, structural and functional data preprocessing and general linear
modeling were performed using Freesurfer (version 7.4.1, available at
http://surfer.nmr.mgh.harvard.edu/) (Fischl, 2012). Anatomical reconstruction (recon-all) and
registration (register-sess) were performed (Dale et al., 1999; Fischl et al., 2002) with human
quality control inspection. Preprocessing included removal of non-brain tissue using a hybrid
watershed/surface deformation procedure (Ségonne et al., 2004) , automated Talairach
transformation, intensity normalization (Sled et al., 1998), tessellation of the gray matter white
matter boundary, automated topology correction (Fischl et al., 2001; Segonne et al., 2007), and
surface deformation following intensity gradients to optimally place the gray/white and
gray/cerebrospinal fluid borders at the location where the greatest shift in intensity defined the
transition to the other tissue (Dale et al., 1999) (mkbrainmask-sess). Functional data
preprocessing included motion correction (mc-sess), FSL's "topup" and "applytopup" functions
(Jenkinson et al., 2012; Smith et al., 2004) for field map distortion correction, slice-time
correction (stc-sess), individual volume to fsaverage surface transformation (163,842 vertices)
and smoothing with a 5mm full width at half max (FWHM) Gaussian kernel (rawfunc2surf-sess).
General linear modeling included one regressor per stimulus condition convolved with the SPM
canonical hemodynamic response function, as well as a nuisance regressor for linear drift
removal, and 12 motion correction regressors per run (3 translation, 3 rotation, and their
derivatives).

For functional connectivity analysis, anatomical and functional data were preprocessed
using CONN (release 22.v2307) (Whitfield-Gabrieli & Nieto-Castanon, 2012) and SPM (release
12.7771) (Penny et al., 2011). The preprocessing pipeline consisted of slice timing correction,
field map correction, outlier detection, direct coregistration to structural, resampling of
functional data within the cortical surface, and smoothing of surface-level functional data. Slice-
time correction was performed following SPM STC procedure (Sladky et al., 2011), using sinc
temporal interpolation to resample each slice BOLD timeseries to a common mid-acquisition
time. Outside of CONN, fieldmap distortion correction was applied using FSL's "topup" and
"applytopup" functions (Jenkinson et al., 2012; Smith et al., 2004). Back inside CONN, potential
outlier scans were identified using ART (Whitfield-Gabrieli & Nieto-Castanon, 2011) as
acquisitions with framewise displacement above 1 mm or global BOLD signal changes above 5
standard deviations (Power et al., 2014), and a reference BOLD image was computed for each
subject by averaging all scans excluding outliers. Functional and anatomical data were
coregistered using SPM intermodality coregistration procedure (Ashburner & Friston, 1997) with

a normalized mutual information objective function (Studholme et al., 1998). Functional images
were resampled at the cortical surface by averaging the functional data from ten locations
between the pial and white matter cortical surfaces of each subject at each individual vertex in
FreeSurfer fsaverage level-8 tessellation, with 163,842 vertices per hemisphere. Surface-level
functional data were smoothed using 4 iterative diffusion steps, approximately a 3mm FWHM
smoothing kernel within the cortical surface (Hagler et al., 2006). Last, anatomical data were
segmented into grey matter, white matter, and CSF tissue classes using SPM unified
segmentation and normalization algorithm (Ashburner, 2007; Ashburner & Friston, 2005) with
the default IXI-549 tissue probability map template.

In addition, functional data were denoised using a standard CompCor (Behzadi et al.,
2007) denoising pipeline including the regression of potential confounding effects characterized
by white matter timeseries, CSF timeseries, motion parameters and their first order derivatives,
outlier scans (Power et al., 2014), session and task effects and their first order derivatives, grey
matter timeseries (Behzadi et al., 2007), and linear trends within each functional run, followed by
bandpass frequency filtering of the BOLD timeseries between .008 Hz and .09 Hz. CompCor
(Chai et al., 2012) noise components within white matter and CSF were estimated by computing
the average BOLD signal as well as the largest principal components orthogonal to the BOLD
average, motion parameters, and outlier scans within each subject's eroded segmentation masks.

Task and resting state runs were rejected if there was more than 1mm average head
displacement from midpoint position during the run (calculated using Freesurfer's "mc-sess").
Twelve runs were rejected (7.6%), leading to one full subject rejection in the task dataset, and
one full subject rejection in the resting state dataset. Thus, our task-based analyses contained
twenty-one participants while our resting state analyses contained eighteen participants.

<u>Localizing Visual, Auditory, and Supramodal Regions of Interest</u>

Visual-biased, auditory-biased, and supramodal WM ROIs in the present study were
initially identified in prior studies with different data sets (Michalka et al., 2015; Noyce et al.,
2022; Tobyne et al., 2025). The present analysis follows the within-subject approach used in
these earlier studies to define ROIs at the individual subject level. The full names, abbreviations,
average MNI centroids, and average surface areas (mm^2) for each ROI can be found in Table 1.
All ROI creation was done after transforming the contrasts from individual cortical space to
fsaverage. In total, 672 individual ROIs were identified for this study (16 ROIs x 2 hemispheres
x 21 participants). The identification of such a large number of functional ROIs presents
challenges, especially in frontal cortex where regions tend to be small and activation levels can
be modest. Moreover, since the focus of this study is on functional connectivity measures, we
required that each ROI contain a minimum of 100 cortical surface vertices. Our approach uses
group-constrained ROIs to define subject-specific ROIs, paralleling an approach first used to
define language-network regions (Fedorenko et al., 2010). Our ROI definition process is outlined
in Figure 2 and the specifics are described below.

Visual and auditory frontal and posterior ROIs were defined at the individual subject
level by directly contrasting visual WM and sensorimotor control blocks with auditory WM and

sensorimotor control blocks. This contrast was liberally thresholded at p < .05, uncorrected in
order to maximally capture vertices showing a bias for auditory or visual stimuli in individuals.
Cortical significance maps were used in combination with anatomical expectations from
previous studies (Michalka et al., 2015; Noyce et al., 2017, 2022; Tobyne et al., 2025; see
Supplemental Tables 1,2) to define 5 bilateral frontal auditory-biased ROIs (tgPCS, cIFS/G, CO,
FO, cmSFG), 3 bilateral frontal visual-biased ROIs (sPCS, iPCS, midIFS), and 2 large bilateral
posterior sensory cortex ROIs (pVis, pAud). Each ROI was required to lie in or near the expected
anatomical region (e.g., on the superior precentral sulcus), and exhibit the expected contrast (e.g.,
visual > auditory, p < .05).

Supramodal ROIs were defined for each individual by first directly contrasting visual
WM with visual sensorimotor blocks, then separately contrasting auditory WM with auditory
sensorimotor blocks. Then both contrasts were thresholded at t > 2 and their intersection was
taken. A group-level probabilistic mask was then made from vertices contained in at least 5
individuals' intersection masks. This probabilistic mask clearly segmented into 6 bilateral
cortical clusters (Figure 3, red outline), similar to domain-general frontal regions localized in
Assem et al. (2020) and Crittenden & Duncan (2014). We refer to these as supramodal (sm)
ROIs (aINS-sm, ACC-sm, preSMA-sm, sPCS-sm, iPCS-sm, midIFS/MFG-sm) because they are
active for both auditory and visual WM. The group-level probabilistic mask was then dilated by
5 vertices to create a search space for each supramodal ROI that better captures individual
variability in location. Abutting supramodal ROI search spaces were delineated by hand at
natural constriction points (Figure 3, red outline). Then, individualized ROIs were created by
taking the individual auditory WM t >2, and visual WM t > 2 intersection vertices within each
probabilistic supramodal ROI. For each subject, any vertices included in their sensory-biased
ROIs were excluded from their supramodal ROIs.

Across both sensory-biased and supramodal, 21.5% of ROIs (138/672) identified had
fewer than 100 vertices. We replaced these ROIs with the strongest 100 vertices within the
probabilistic search space for the ROI. To expand the sensory-biased ROIs with fewer than 100
vertices, we first created probabilistic sensory-biased ROIs (Figure 3, blue and orange) with the
same process used to create the supramodal probabilistic ROIs. Then we used the 100 vertices
with the largest t-stats within the probabilistic ROI to replace the sensory-biased ROIs with
fewer than 100 vertices. Only 7 sensory-biased ROIs (~1%) contained any vertices with t-stats in
the opposite direction than expected. For the supramodal ROIs with fewer than 100 vertices, the
auditory WM > auditory sensorimotor and visual WM > visual sensorimotor t-stat thresholds
were incrementally lowered together until 100 vertices were included, excluding any vertices in
the sensory-biased ROIs. Across all subjects, only 4 supramodal ROIs (<1%) contained any
vertices with negative t-stats.

Two participants were rejected from all analyses due to inability to draw more than 75%
of ROIs at a liberal p < .05 uncorrected threshold; for one there was low overall activation (ex.
no significant activity in the auditory cortex), while for the other, activity was heavily skewed
towards visual and sensorimotor control conditions. This left 21 participants (18 resting state).

Figure 2 shows a diagrammatic summary of the sensory-biased and supramodal ROI localization
processes.

297
Figure 2. Flow chart summarizing the steps in defining the ROIs used in this study. 32 ROIs were defined for each
individual participant. In cases where the subject-level data alone yielded a small ROI, probabilistic search spaces
were used in conjunction with the subject-level data to yield ROI definitions of at least 100 vertices.

Analysis Overview

We employed several functional connectivity analyses to investigate the resting state and
task-based network organization of sensory-biased and frontal supramodal ROIs. To establish
robust network organization at rest we used hierarchical clustering analysis (HCA), then used
linear mixed effects models (LME) on ROI-to-ROI correlations grouped by network to establish
connectivity patterns between the networks. To reveal which specific ROIs from each network
were driving connectivity, we used threshold-free cluster enhancement (TFCE) on the ROI-to-
ROI correlations. To then understand how network-network and ROI-ROI connectivity
differentially change with visual and auditory WM task demands, we used generalized
psychophysiological interaction (gPPI) analysis. The PPI coefficients were grouped by network
and used in LME analysis to see how network connectivity might restructure during visual and
auditory WM tasks. Again, to reveal which specific ROIs from each network were driving these
changes, we used TFCE on the ROI-to-ROI PPI coefficients.

Resting State Functional Connectivity Analysis

HCA was used to determine which ROIs had the most correlated activity. CONN Toolbox
computed complete-linkage clustering (Sørensen, 1948) with optimal leaf ordering (Bar-Joseph
et al., 2001) based on functional similarity only (contribution of anatomical proximity was set to
zero) (Nieto-Castanon, 2020). The number of clusters was selected by maximizing the difference
between normalized number of clusters and the normalized inverse distance metric (i.e.
analogous to choosing the "elbow" of the curve plotting the number of clusters against the
distance metric). These clusters were considered as functional networks in the following
analyses.

To assess resting-state connectivity between networks found via HCA, we first used
CONN Toolbox to compute ROI-to-ROI connectivity using Fisher-transformed bivariate
correlation coefficients from a general linear model (weighted-GLM) for each pair of ROIs
separately, thus characterizing the association between their BOLD signal timeseries. To
compensate for potential transient magnetization effects at the beginning of each run, individual
scans were weighted by a step function convolved with the SPM canonical hemodynamic
response function and rectified (Nieto-Castanon, 2020). We then used these ROI-to-ROI
correlation coefficients as the dependent variable in an LME, with network connection type as
the fixed effect, and subject and hemisphere as full random effects (intercept, slope, and
covariance). Network connection type was assigned based on which HCA-determined networks
the ROIs were in; for example, the correlation coefficient for sPCS and sm-aINS would be
labeled as a "frontal visual↔supramodal" network connection. Because this analysis is GLM-
based, there were separate coefficients for reciprocal connections, which were averaged to
represent the directionally agnostic connectivity between two ROIs. Next, estimated marginal
means and standard errors of connectivity for each connection type were computed from the
LME coefficients and used in Wald-tests for significance (Hartman, 2019/2023). P-values were
FDR-corrected for significance at p < .05. The robustness of this method relies on using fully
specified random effects (intercept, slope, and covariance) of subject and hemisphere to account
for hemispheric- and subject-level effects and enhance generalizability (Barr et al., 2013), and
FDR-correction for multiple comparisons.

Assessing significance of individual ROI-to-ROI connectivity is complicated by the
multiple comparisons problem due to the large number of ROIs (496 comparisons among 32
ROIs), thus we used the nonparametric spatial clustering method TFCE to reduce the number of
comparisons being made. This method has the additional advantage of increased sensitivity by
avoiding use of an arbitrary cluster-forming threshold, and it easily lends itself to permutation
testing for significance (Smith & Nichols, 2009). We used CONN Toolbox to first estimate a
separate GLM for each connection, creating a group-level ROI-to-ROI connectivity matrix, then
used the CONN Toolbox implementation of TFCE to determine clusters of significant
connectivity among ROIs within and between all HCA-determined networks. Significant clusters
of connections were determined using permutation testing of 1,000 residual-randomization
iterations with FDR correction of TFCE-scores at p < .05 (Nieto-Castanon, 2020). The
robustness of this method relies on avoiding the use of a hard cluster-level threshold,
nonparametric permutation testing (Eklund et al., 2016, 2019), and FDR-correction for multiple
comparisons.

Task-Based Functional Connectivity Analysis

To assess the changes in connectivity during visual and auditory WM tasks, we used gPPI
analysis implemented in CONN Toolbox. PPI assesses changes in connectivity between 2 ROIs
due to particular experimental conditions using bivariate regression. Each regression model
estimates the variance in one ROI BOLD timeseries explained by 1) terms for each task effect
convolved with a hemodynamic response, 2) the seed ROI BOLD timeseries, and 3) the
interaction term, which is the product of terms 1 and 2. Thus, by estimating the coefficients for
term 3 (the interaction term or PPI), we can quantify the amount of variance in one ROI signal
explained by the interaction between another ROI and each task. Though the ROIs were defined
from the task data set, the PPI analysis is non-circular because 1) task effects and their
derivatives are regressed out in the denoising step, and 2) PPI analysis assesses the predictability
of one ROI's signal by another ROI's signal under a specific condition *over and above* the
variability explained by the seed ROI's signal alone (term 2) and the task effect alone (term 1);
that is, the PPI analysis only detects effects that are orthogonal to the main effect of the task
(Friston et al., 1997; McLaren et al., 2012; O'Reilly et al., 2012). Additionally, using simulated
and experimental data, gPPI is shown to estimate context-modulated functional connectivity
especially well under block-designs compared to beta-series methods and sPPI (Cisler et al.,
2014).

The gPPI analysis was run for every ROI-to-ROI connection and PPI coefficients were
used to form 2 PPI matrices, one for the change in connectivity between ROIs during visual WM
and one for auditory WM. Note that PPI coefficients represent the impact of a specific
experimental condition on connectivity between ROIs compared to a "baseline" of all other
conditions (which here included fixation and sensorimotor blocks). Next, these matrices were
subtracted in order to get the difference in connectivity between visual and auditory WM. This
matrix was then evaluated in the same way as the connectivity coefficient matrix in the above
LME and TFCE analyses. Thus, the LME analysis with the PPI data showed how the network-
level connectivity differed between visual and auditory WM tasks, while the TFCE analysis
showed how ROI-level connectivity differed between the visual and auditory WM tasks.

Assessing Overlap Between Supramodal and Domain-General Frontal Regions

In order to assess the similarity between our frontal supramodal ROIs and the domain-
general frontal ROIs from the literature (Assem et al., 2022), we used the Sørensen-Dice
similarity coefficient (Dice, 1945; Sørensen, 1948) which simply takes 2 times the number of
vertices in common between the two ROIs and divides by the sum of the number of vertices in
both ROIs, thus quantifying the spatial similarity between the ROIs:

$$Dice = \frac{2\,|X \cap Y|}{|X| + |Y|}$$ where |X| indicates number of elements in set X

We assessed only the ROI pairs for which the probabilistic supramodal search space visually
intersected with the domain-general ROIs (derived from the HCP-MMP1.0 parcellation) when
projecting both on the fsaverage cortex (Supplemental Figure 1). Subsequently, we used the

individually defined supramodal ROIs to assess overlap with the fsaverage-projected domain-
general ROIs for each participant.

Predicting Working Memory Precision from Dynamic Network Connectivity

To evaluate the relationship between subject-level behavioral task precision and task-
related changes in connectivity among networks during each task, we fit linear regression models
using task precision as the dependent variable and a subset of network-level PPI difference terms
as the independent variables. Working memory precision for each participant was defined by the
average staircase threshold obtained. Separate staircasing was done for visual and auditory
stimuli and thus visual and auditory memory precision estimates were each obtained. The
staircasing procedures sought to equate task difficulty (accuracy) across modalities and across
individuals by adjusting the size of the frequency difference between a non-match stimulus and a
match stimulus. Staircase thresholds were recorded as Weber fractions or percentage changes
from the baseline. Finer thresholds (i.e., smaller frequency difference between stimuli) indicate
finer WM precision. Thus, if a participant performed well (poorly) on the task, their difficulty
staircase would increase (decrease), and it would be harder (easier) to tell the difference between
the first and second stimulus in a non-match trial. To calculate task precision, we took the
average difficulty staircase level across all runs for each participant. To predict task precision in
the *visual* WM task, we averaged the PPI coefficient differences for network connections with
significant increases in connectivity during visual WM compared to auditory WM. Similarly, to
predict task precision in the *auditory* WM task, we averaged the PPI coefficient differences for
network connections with significant increases in connectivity during auditory WM compared to
visual WM. Our reasoning was that the network-network connections which showed significant
changes during the task were most likely to impact behavioral performance. PPI coefficients
were also averaged across hemisphere for each participant.

Results

Subjects (N=21) performed 2-back visual working memory and 2-back auditory working
memory tasks during fMRI (see Figure 1; Methods). Staircasing procedures were applied to
ensure that the sensory working memory tasks were challenging and well matched across
sensory modalities. Task accuracy did not differ significantly between visual and auditory WM
tasks (visual WM=68.1% auditory WM=64.8%, t(20)=1.91, p=.08) (Supplemental Figure 2).

We examined the 2-back WM data to define 32 ROIs (16 ROIs per hemisphere) for each
individual subject. Prior work (Noyce et al., 2022; Tobyne et al., 2025; see Supplemental Tables
1,2) from our laboratory identified three visual-biased frontal lobe regions per hemisphere –
superior precentral sulcus (sPCS), inferior precentral sulcus (iPCS), and mid-inferior frontal
sulcus (midIFS) – and five auditory-biased frontal lobe regions per hemisphere – caudomedial
superior frontal gyrus (cmSFG), transverse gyrus intersecting the precentral sulcus (tgPCS),
caudal inferior frontal sulcus / gyrus (cIFS/G), central operculum (CO), and frontal operculum
(FO). In occipitoparietal cortex we observed a large contiguous region that preferred visual
processing (pVis), while in lateral temporal lobe we observed a large contiguous region that

preferred auditory processing (pAud). Given the focus on frontal lobe regions, no attempt was
made to isolate smaller, more specific function regions within pVis or pAud and the data
contrasts were not conducive to subdivision of these contiguous regions. We identified these 10
sensory-biased ROIs in each hemisphere of each participant by examining the contrast of visual
conditions (WM and sensorimotor) with auditory conditions (WM and sensorimotor) (See Figure
2; Methods). Additionally, prior work from the laboratory identified five supramodal frontal lobe
regions in each hemisphere (Tobyne et al., 2025; see Supplemental Table 2). Following a similar
logic, we identify those five regions and one additional region in anterior cingulate cortex that
emerged from the analysis (See Figure 2; Methods). These supramodal regions are anterior
insula (aINS-sm), pre-supplementary motor area (preSMA-sm), anterior cingulate cortex (ACC-
sm), middle inferior frontal sulcus / middle frontal gyrus (midIFS/MFG-sm), superior precentral
sulcus (sPCS-sm), and inferior precentral sulcus (iPCS-sm). The latter three supramodal ROIs,
midIFS/MFG-sm, sPCS-sm, and iPCS-sm lie near visual-biased midIFS, sPCS, and iPCS,
respectively, but did not overlap in individual subjects. In total, 672 individual ROIs were
defined. 138 (21.5%) of these ROI were expanded using group-constrained subject-specific
methods so that all ROIs were subject-specific and contained a minimum of 100 cortical surface
vertices. MNI coordinates and region sizes are summarized in Table 1.

The mean MNI coordinates aligned well with those regions defined in prior studies
(Noyce et al., 2022; Tobyne et al., 2025), even though the subjects and stimuli differed across
datasets (Supplemental Tables 1, 2). Supramodal regions, which were defined here based on the
two-way intersection of visual WM and auditory WM activation, tended to be larger than those
defined by a three-way intersection (visual, auditory, and tactile WM) in the prior analysis
(Tobyne et al., 2025), but otherwise appear well-aligned. Individual subject ROIs were compiled
to create probabilistic ROI maps, which are displayed in Figure 3. Individual ROIs for example
subjects are presented in Supplemental Figure 3.

We note that the supramodal ROIs we have functionally defined bear a striking
anatomical similarity to a large subset of the domain-general network identified in broad task
batteries (Assem et al., 2020; Fedorenko et al., 2012). To examine the anatomical relationship
between the present set of supramodal WM ROIs and the previously defined domain-general
ROIs, we performed Dice coefficient and percent overlap analyses between our individual-
subject supramodal ROIs and the MMP 1.0 multi-modal parcels (Glasser et al., 2016) that are
identified as the domain-general network (Assem et al., 2020). Only pairs of ROIs in which the
supramodal ROI search space overlapped with a domain-general ROI were assessed. Five of the
six supramodal ROIs identified here heavily overlapped (>50%) with the frontal domain-general
ROIs (see Supplemental Figure 1; Supplemental Table 3). This gives evidence that what we
identified as supramodal WM ROIs (using visual and auditory WM > visual and auditory
sensorimotor contrast) may constitute a substantial subset of domain-general cortex.

Figure 3. Lateral and medial left and right hemispheres with visual-biased probabilistic ROI search spaces in blue, auditory-biased probabilistic ROI search spaces in orange, and supramodal probabilistic ROI search spaces outlined in red. Name labels for the sensory-biased ROIs are shown only on the right hemisphere while supramodal ROI labels are shown only on the left hemisphere. Note that although there is overlap between the probabilistic supramodal and other probabilistic search spaces, the individual-subject ROIs never overlap. Three example individual subjects' ROIs are shown in Supplemental Figure 3.

Abbreviation	Sensory Bias	ROI Name	MNI centroid (lh, rh)	Surface Area (mm^2) (lh, rh)
sPCS	Visual	Superior precentral sulcus	(-35.5, -7.4, 45.6), (36.9, -5.7, 46.5)	242±200, 328±281
iPCS	Visual	Inferior precentral sulcus	(-45.6, 0.8, 32.1), (43.2, 2.8, 29.8)	319±265, 475±396
midIFS	Visual	Middle inferior frontal sulcus	(-39.1, 29.3, 19.2), (42.0, 29.7, 17.9)	116±72, 182±196
pVis	Visual	Posterior visual	(-29.2, -88.8, 3.0), (30.8, -87.5, 3.6)	8388±3055, 9493±3145
tgPCS	Auditory	Transverse gyrus intersecting precentral sulcus	(-49.4, -6.7, 45.8), (51.7, -3.8, 41.9)	137±93, 142±79
cIFS/G	Auditory	Caudal inferior frontal sulcus/gyrus	(-42.8, 13.7, 22.4), (41.7, 15.1, 22.6)	218±176, 178±122
CO	Auditory	Central operculum	(-57.5, -5.1, 14.3), (59.0, -3.7, 13.0)	189±117, 170±146
FO	Auditory	Frontal operculum	(-42.9, 30.8, 1.1), (48.0, 29.5, 3.1)	259±202, 209±152
cmSFG	Auditory	Caudomedial superior frontal gyrus	(-7.0, 3.5, 61.3), (7.9, 6.3, 62.1)	102±96, 91±48
pAud	Auditory	Posterior auditory	(-52.4, -24.0, 2.2), (58.6, -18.2, 3.6)	2979±708, 2864±646
preSMA-sm	Supramodal	Supramodal pre-superior motor cortex	(-9.6, 14.9, 47.7), (8.5, 22.0, 44.7)	459±278, 568±321
ACC-sm	Supramodal	Supramodal anterior cingulate cortex	(-12.5, 25.2, 30.1), (12.6, 26.5, 26.4)	89±44, 145±96
aINS-sm	Supramodal	Supramodal anterior insula	(-28.3, 24.1, 3.5), (29.9, 25.0, 3.7)	526±300, 619±326
sPCS-sm	Supramodal	Supramodal superior precentral sulcus	(-27.6, -4.4, 45.2), (27.3, -2.4, 48.6)	470±398, 471±446
iPCS-sm	Supramodal	Supramodal inferior precentral sulcus	(-39.9, 2.5, 27.5), (40.6, 9.8, 23.3)	383±371, 571±509
mid-IFS/MFG-sm	Supramodal	Supramodal mid-inferior frontal sulcus / middle frontal gyrus	(-42.2, 28.1, 25.9) (42.1, 36.3, 23.3)	344±278, 635±484

Table 1. Abbreviations, full names, sensory-bias, centroid MNI coordinates, and surface area for each ROI. The centroids were calculated using MNE's "center_of_mass" function on each participant's anatomical label in fsaverage space, then converted from vertex index to MNI coordinates using MNE's vertex_to_MNI function. The coordinates were then averaged across participants and displayed in left, right hemisphere order. The surface area was calculated in individual subject cortical space using Freesurfer (mris_anatomical_stats) and means ± standard deviations were reported in mm^2 for each hemisphere (left, right).

Resting-State Functional Connectivity: Auditory, Visual, and Supramodal Networks Form Distinct but Interconnected Resting State Networks

We examined network organization amongst 32 ROIs using resting-state fMRI data to establish baseline, task-free connectivity. Previously, we demonstrated that frontal visual and auditory ROIs form discrete resting state networks that include their respective posterior sensory cortices (Michalka et al., 2015; Noyce et al., 2022; Tobyne et al., 2017). Moreover, frontal visual-biased ROIs were previously reported to more robustly connect with supramodal or multiple-demand ROIs than do frontal auditory-biased ROIs (Tobyne et al., 2017; Noyce et al., 2022). Here, we worked with subject-specific supramodal, visual-biased, and auditory-biased ROIs to independently examine network organization and characterize the connectivity between networks. We performed seed-to-seed functional connectivity analysis of resting-state data, using the ROIs defined above as the seed regions. This analysis was limited to subjects (N=18) for whom resting-state data was obtained. Hierarchical Clustering Analysis (HCA) of the seed-to-seed functional connectivity matrix revealed 4 intra-connected groups of ROIs (see Methods). One network consisted of bilateral visual-biased frontal ROIs (sPCS, iPCS), another consisted of the bilateral auditory-biased frontal and posterior ROIs (tgPCS, cIFS/G, CO, FO, cmSFG, pAud), and another consisted of all the bilateral supramodal frontal ROIs plus a single visual-biased frontal ROI (aINS-sm, preSMA-sm, ACC-sm, sPCS-sm, iPCS-sm, midIFS/MFG-sm, midIFS). The final cluster consisted of the single visual-biased posterior bilateral ROI, posterior visual (pVis). We refer to these networks as *frontal visual*, *frontal+posterior auditory*, *supramodal*, and *posterior visual* networks respectively. The HCA results largely agreed with previous results that showed visual-biased and auditory-biased frontal ROIs formed separate networks (Michalka et al., 2015; Tobyne et al., 2017; Noyce et al., 2022) and extended previous results by showing that supramodal ROIs formed another separate network, with the exceptions that pAud clustered with the frontal auditory ROIs and midIFS clustered with the *supramodal* network instead of the *frontal visual* network.

To characterize the connectivity between networks, we used linear mixed effects (LME) analysis of correlations between ROIs in each network (see Methods). Significant positive connectivity was found between *posterior visual* ↔ *frontal visual*, *frontal visual* ↔ *supramodal*, *frontal+posterior auditory* ↔ *supramodal*, and *frontal visual* ↔ *frontal+posterior auditory*. Significant negative connectivity was found between *posterior visual* ↔ *frontal+posterior auditory*, and *posterior visual* ↔ *supramodal*. As expected, significant positive connectivity within network groups (e.g., *frontal visual* ↔ *frontal visual*) was also observed for each network. These connectivity patterns and the mean correlations are shown in Figure 4a. In general, we saw two distinct streams of positive connectivity, a 'visual stream' between the *posterior visual*, *frontal visual*, and *supramodal* networks, and separately, an 'auditory stream' between the *posterior+frontal auditory* and *supramodal* networks. The connectivity between *frontal visual* ↔ *supramodal* (M=.17, SE=.02) was significantly greater (W(1) =5.1, p<.001) than between the *frontal+posterior auditory* ↔ *supramodal* networks (M=.05, SE=.02). This finding replicated and extended the sensory network asymmetry previously reported (Tobyne et al., 2017; Noyce et al., 2022) by demonstrating that the visual stream exhibited greater functional connectivity with supramodal regions than did the auditory stream. Note that this functional connectivity asymmetry existed even though the supramodal ROIs did not overlap with sensory-biased ROIs. It is also of note that there was only a single point of significant positive connectivity between the visual and auditory networks, and that connection between *frontal*

visual ↔ frontal+posterior auditory was weak (M=.04, SE=.02), indicating fairly separate visual
and auditory streams. Given that the nodes of the frontal visual and auditory networks are
anatomically interleaved with each other and abut in many instances, the relative segregation of
the modality-biased networks is remarkable.

In order to ensure that the results were not unexpectedly biased by the inclusion of
individual ROIs defined with guidance from the probabilistic maps, we repeated the above LME
analysis in full using only the ~80% of ROIs defined robustly from individual subject data alone.
The connection between *frontal visual ↔ frontal+posterior auditory* networks dropped below
significance in this alternative analysis, further indicating relatively segregated processing. All
other significant and non-significant results remained significant and non-significant,
respectively (Supplemental Table 4).

Figure 4. Resting-state connectivity results. A) Diagram summarizing the LME-derived significant resting state
connectivity between ROI networks. The values shown next to each connection represent the mean correlation
coefficient between ROIs in each network. Self-connection in the *posterior visual* network was not assessed because
it only contained 1 (bilateral) ROI. B) Connectogram summarizing the TFCE-derived significant resting state
connectivity between individual ROIs. Warm colors represent positive connectivity while cool colors represent
negative connectivity. ROIs are grouped around the circle by HCA-derived networks.

To assess connectivity between specific ROIs within each network, we next used TFCE
of the seed-to-seed functional connectivity with ROIs sorted using the HCA clustering results
(see Methods). Results from the TFCE analysis are shown in the connectogram in Figure 4b. The
ROIs are grouped into their HCA-identified networks around the outside of the connectogram in
Figure 4b. Most clusters recapitulated the connectivity patterns observed in the LME analysis,

but now with ROI-level specificity. For example, the negative connectivity between *posterior*
visual ↔ frontal+posterior auditory networks was seen to be driven by negative connectivity of
pVis with pAud, CO (L), and FO. We also saw that there was especially strong connectivity
between ROIs in the *frontal visual ↔ supramodal* networks while connectivity between ROIs in
the *auditory frontal+posterior ↔ supramodal* networks was present, but generally weaker.

<u>Task-Based Functional Connectivity: Auditory Working Memory Drives More Widespread</u>
<u>Changes in Connectivity than Visual Working Memory</u>

The above results demonstrated that: 1) largely distinct visual-biased and auditory-biased
networks exist in the cortex; 2) each modality-biased network connected within the frontal lobe
to a supramodal network; and 3) in the resting state, supramodal regions exhibited stronger
functional connectivity with the visual network than with the auditory network. How can we
reconcile the network asymmetry *at rest* with the apparent unbiased supramodal activation
during WM task performance? We hypothesized that during auditory WM task performance the
auditory stream increases its functional connectivity (compared to baseline) and does so to a
greater extent than occurs for the visual stream during visual WM. To test this hypothesis, we
examined how network connectivity changed with the task demands of visual WM and auditory
WM, using PPI analysis. This analysis regresses out the task vectors, as well as their first
derivatives during denoising (see Methods), thus mitigating the confound of connectivity
induced due to baseline activation increases/decreases due to the tasks.

We first performed a network-level LME analysis of PPI coefficients (see Methods) and
observed an increase in connectivity between the *frontal+posterior auditory ↔ supramodal*,
frontal+posterior auditory network ↔ itself, and *frontal+posterior auditory ↔ frontal visual*
networks during auditory WM, as compared to visual WM. We also saw an increase in
connectivity between the *posterior visual ↔ frontal visual*, and *posterior visual ↔ supramodal*
networks during visual WM, as compared to auditory WM (Figure 5a). Notably, there was no
significant change in connectivity between the *frontal visual ↔ supramodal* networks during the
two working memory conditions. It is also important to note that the increases in connectivity
during auditory WM compared to visual WM were largely driven by an increase in connectivity
during auditory WM (compared to baseline), whereas the visual>auditory WM increases in
connectivity were driven largely by decreases in connectivity during auditory WM (compared to
baseline) (Figure 5b). In order to ensure that the results were not unexpectedly biased by the
inclusion of individual ROIs defined with guidance from the probabilistic maps, we repeated the
above LME analysis in full using only the ~80% of ROIs defined robustly from individual
subject data alone. All significant and non-significant results remained that way (Supplemental
Table 5).

These findings suggest that auditory WM task performance caused more profound
restructuring of connectivity between networks than did visual WM task performance. A post-
hoc summary analysis of the changes in connectivity between networks in the proposed "visual
stream" during visual WM was compared against the changes in connectivity between networks

in the proposed "auditory stream" during auditory WM. Thus, a one-way ANOVA with random
effects for subject and hemisphere was run comparing the average visual WM PPI terms for the
following connections: *posterior visual ↔ frontal visual, frontal visual ↔ frontal visual, frontal*
visual ↔ supramodal, and the average auditory PPI terms for the following connections:
frontal+posterior auditory ↔ frontal+posterior auditory, frontal+posterior ↔ supramodal.
There was a statistically significant difference showing larger average changes in connectivity
across the "auditory stream" during auditory WM compared to baseline than average changes in
connectivity across the "visual stream" during visual WM compared to baseline (F(1,125)=6.24,
p=.014). The effect size (η^2) was .04 indicating between a small and medium effect size.

We also assessed ROI-level changes in connectivity using TFCE on the seed-to-seed PPI
coefficients with ROIs sorted using the resting state HCA clustering results (Figure 5c; see
Methods). Again, we saw that the significant clusters of connectivity changes at the ROI-level
matched well with the patterns of connectivity changes in the network-level analysis, but we
gained specificity as to which ROIs were driving the changes. For example, it appears that the
auditory>visual WM change in connectivity between the *frontal+posterior auditory ↔*
supramodal networks was driven primarily by connections between [pAud ↔ iPCS-sm (L)],
[cIFS/G (L) ↔ aINS-sm (L), preSMA-sm (L), sPCS-sm (R)], and [FO (R) ↔ midIFS, iPCS-sm
(R), preSMA-sm (R), aINS-sm (R)]. Additionally, the visual WM>auditory WM change in
connectivity between *posterior visual ↔ supramodal* networks was driven by connections
between pVis ↔ iPCS-sm, midIFS (R). The increased connectivity between ROIs within the
frontal visual network (sPCS (L) ↔ iPCS (L)) during auditory WM compared to visual WM was
not reflected in the network-level LME results, possibly because it was limited to a small number
of ROIs. While an increase in connectivity between visual ROIs during auditory WM may seem
counterintuitive, it could be related to the recruitment of the *frontal visual* network by the
frontal+posterior auditory network during auditory WM, especially since sPCS (L) was central
to both of these changes in connectivity. The large majority of significant differences in
connectivity between auditory and visual WM in ROI-ROI connectivity identified were the result
of increased connectivity under auditory WM compared to baseline, again consistent with the
idea that auditory WM leads to greater reorganization of functional connectivity within and
between sensory-biased and supramodal networks than does visual WM.

Figure 5. Task-based functional connectivity results. A) Diagram summarizing the LME-derived significant changes in connectivity between ROI networks during visual and auditory WM. The black arrows show significant resting-state connectivity from Figure 4, while the large colored arrows indicate change in connectivity. The values and small colored arrows shown next to the large colored arrows indicate the mean increase or decrease in PPI coefficients for the visual and auditory WM conditions between ROIs in each network. For example, the change in connectivity between *posterior visual* ↔ *frontal visual* networks increased under visual WM (large blue arrow) and was driven by a small (.1) increase in the visual WM condition and a large decrease (.10) in the auditory condition (small blue and orange arrows and values). Changes in self-connection in the *posterior visual* network were not assessed because the network only contained 1 (bilateral) ROI. B) Bar plot showing means and SEs for the visual and auditory WM PPI terms for each network connection. Only significant differences appear in the 5A summary

diagram. C) Connectogram summarizing the TFCE-derived significant changes in connectivity between individual
ROIs during visual and auditory WM. Warm colors represent auditory>visual changes in connectivity while cool
colors represent visual>auditory changes in connectivity. ROIs are grouped around the circle by HCA-derived
networks.

<u>Auditory WM Precision is Correlated with Changes in Connectivity Between Key Networks</u>

In order to assess the behavioral impact of shifting connectivity between networks during
auditory and visual WM tasks, we used linear regression with WM task precision as the
dependent variable and the difference between visual and auditory PPI coefficients as the
independent variables (see Methods). After the above PPI analyses were completed, we
hypothesized that individual subjects with more robust auditory network reconfigurations would
also exhibit greater auditory WM precision. Specifically, we hypothesized that at the individual
level, increases in connectivity between key networks during auditory WM would predict greater
auditory WM precision. Those key network connections were selected based on which networks
had significantly increased connectivity during auditory WM compared to visual WM in the
previous group level analysis (*frontal+posterior auditory ↔ supramodal, frontal+posterior*
auditory ↔ frontal visual, and *frontal+posterior auditory ↔ frontal+posterior auditory*). In line
with this hypothesis, we found that changes in connectivity in these key network-level
connections during auditory WM compared to visual WM significantly predicted behavioral
precision on the auditory WM task (R^2=.20, F(1,19)=4.81, β=-37.76, p=.0204; Figure 6).

Figure 6. Auditory WM precision in individual participants correlated with the strength of the auditory network reconfiguration during the WM task. Each blue X represents a single participant's data (N=21). The red line is the least squares line of best fit, and the dotted lines are 95% confidence bounds. The Y axis represents percent change in frequency between stimuli (lower indicates higher behavioral precision) Note that the x-axis represents the average difference in PPI betas for the 3 network-network connections with significantly increased connectivity during auditory WM compared to visual WM.

We also hypothesized that increases in connectivity between key networks (*posterior*
visual ↔ frontal visual and *posterior visual ↔ supramodal*) during visual WM compared to
auditory WM would *not* predict better visual WM task precision. This hypothesis was derived

from the fact that visual WM elicited much smaller changes in connectivity between networks
than auditory WM, possibly because frontal visual ROIs are already strongly connected to
supramodal ROIs at rest. Mirroring the analysis done on the auditory WM task, this analysis
examined the networks that had significantly increased connectivity during visual WM compared
to auditory WM. In line with this hypothesis we found that changes in connectivity in key
network-level connections during visual WM compared to auditory WM did not predict
behavioral precision on the visual WM task (R^2=.01, $F(1,19)$=.25, β=-6.52, p=.626).

Discussion

In this study we sought to understand the functional connectivity relationships among
visual-biased, auditory-biased, and supramodal cortical ROIs, and how they change during visual
and auditory WM tasks. To do this, we performed resting state and task-based functional
connectivity analyses on 672 participant-specific ROIs (21 participants x 2 hemispheres x 16
ROIs). Our results demonstrated that visual-biased, auditory-biased, and supramodal ROIs form
largely discrete but interconnected resting state networks. We saw little positive connectivity
between visual and auditory network streams indicating that there are relatively separated
processing streams for visual and auditory sensory and WM information, even in the frontal
cortex – with both streams potentially feeding into and/or receiving information from a modality-
agnostic frontal supramodal network. This segregation is remarkable given that the nodes of
these two networks are anatomically interleaved in frontal cortex.

We additionally showed that these networks change their connectivity structure
differentially during visual and auditory WM tasks, with more robust network recruitment
occurring during auditory WM, indicating that the default functional organization favors
connectivity between the visual-biased and supramodal networks and that auditory WM
performance galvanizes connectivity between the auditory-biased network and other networks.
Importantly, we observed that the magnitude of these task-based network changes in individual
subjects correlated with their auditory WM precision.

Several of our key findings focus on an asymmetry between the visual-biased and auditory-
biased networks. Resting-state analysis demonstrated that the *supramodal* network was more
highly connected with the *frontal visual* network than with the *frontal+posterior auditory*
network. This suggests that the *frontal visual* network has "privileged" communication with the
supramodal network at rest. This idea is supported by the lack of detectable significant changes
in functional connectivity between the *frontal visual* and *supramodal* networks during the visual
WM task; perhaps the strong connectivity observed at rest is sufficient to support strong visual
WM performance. In contrast, there was much greater reorganization of connectivity during the
auditory WM task, including the recruitment of the *visual frontal* and *supramodal* networks,
which also correlated with increased auditory WM precision in individual participants. No such
changes were observed for the visual WM task. That is, although the auditory-biased network
appeared disadvantaged at rest in its connectivity to the supramodal WM network, it was capable
of greater dynamic reorganization to obtain the network connectivity that may be needed to
support task performance.

In prior studies, we have reported that nodes of the visual-biased network exhibited robust
resting-state connectivity with purported domain-general brain regions, while nodes of the
auditory-biased network exhibited robust resting-state connectivity with purported language
network regions (Tobyne et al., 2017; Noyce et al., 2022). The current resting-state findings
showing strong functional connectivity between frontal visual-biased and supramodal networks,
add evidence in support of the proposed link between visual-biased and domain-general regions.
Although the present study lacks the full variety of tasks to make a strong claim of domain-
general processing for the supramodal WM regions in individual subjects, our supplementary
analysis (Supplemental Figure 1, Supplemental Table 1) demonstrates that supramodal frontal
regions largely overlapped with a core subset of the domain-general network (Assem et al.,
2020). We suggest that the supramodal WM regions comprise an important subset of the domain-
general network. In addition to the functional connectivity evidence, previous task activation
data also revealed a similar asymmetry. Visual-biased ROIs were recruited in auditory WM tasks
and thus exhibited a degree of supramodal WM, while auditory-biased ROIs showed little
recruitment in visual WM tasks and thus showed little evidence for supramodal WM (Noyce et
al., 2020; Tobyne et al., 2025). Thus, there may be a general asymmetry in activation and
connectivity between frontal auditory and visual networks with the visual network maintaining
preferential access to supramodal network resources, while auditory processing must recruit both
networks to perform auditory WM tasks with high precision. Some behavioral studies have
reported a tighter link for domain-general cognitive processing with visual working memory than
for auditory working memory in adults (Bo et al., 2011) and children (Alhamdan et al., 2023),
but other potential behavioral implications of this asymmetry deserve broader investigation.

Our functional parcellation of frontal cortex included many small regions with
neighboring regions that often belonged to different functional categories (i.e., visual-biased,
auditory-biased, or supramodal). Under such conditions, group-level definitions of ROIs will
yield a very substantial loss of statistical power (Fedorenko et al., 2010; Tobyne et al., 2018). 672
subject-specific functional ROIs were created from task data. 534 (78.5%) of these were defined
directly from subject-specific data. To obtain ROIs of sufficient size (>= 100 cortical surface
vertices) for all areas for all subjects, we employed a group-constrained subject-specific
approach that bears similarities to the approach developed for language structures by Fedorenko
et al. (2010). Our approach effectively reduced the threshold for vertex inclusion and relied on
group-level anatomical priors. This had the potential to dilute the functional specificity of an
ROI, but this dilution effect was bounded by that exhibited by standard group-level ROI
definitions. While diluting the functional specificity of ROIs may have diminished the effects
observed in subsequent analysis, it did not bias the results toward any positive outcome. This is
supported by the fact that the secondary LME analyses, using only the ROIs identified directly
with subject-specific data, yielded very similar results to the original analyses.

One potential confound in the interpretation of the results in this analysis is the fact that
during all functional scans (resting state, visual and auditory sensorimotor and WM tasks),
participants were asked to maintain fixation – thus introducing an implicit simple visual task to
all conditions. This potential issue should not have affected the task-based gPPI analyses of
changes in connectivity between task conditions (Figures 5, 6) because fixation was present in

every condition being compared. As for the resting state analysis, because fixating on a single dot is a low-level visual task, if this task boosts visual connectivity to supramodal WM regions one might expect positive connectivity between lower-level visual cortical areas and supramodal regions, but the resting-state functional connectivity between *posterior visual* and *supramodal* networks was negative. Although there are prior reports of differences in functional connectivity between networks in eyes closed and fixation resting state conditions, the direction and networks involved vary considerably between studies (Agcaoglu et al., 2019; Han et al., 2023; Patriat et al., 2013). Han et al. (2023) reported increased connectivity between the visual and auditory primary cortices during eyes closed compared to fixation, while Agcaoglu et al. (2019) reported the opposite in a study with larger N. The latter study also reported that the connectivity from these sensory cortices to frontal networks was mixed, with some increases and decreases in both conditions. Although we cannot conclusively rule out a small influence of the eyes-open fixation task on the resting-state connectivity in frontal cortex, this condition is held constant and thus is not a factor in the key task-based analyses.

A final important caveat is that fMRI reports of dynamic network reorganization, including the present study, are potentially consistent with multiple interpretations at the level of individual neurons. While it is possible that homogeneous populations of neurons within each ROI alter their functional connectivity with neurons in other nodes and networks, it seems equally plausible that ROIs contain static, but heterogeneous neural populations that differ both in their connectivity and functional responses. Task activation of different sub-populations with different connectivity could appear as a dynamic network reconfiguration in the present type of analysis, even though individual neuronal connections are unchanging. For instance, visual-biased frontal ROIs might contain a minority sub-population of auditory-biased or supramodal neurons that exhibit strong connectivity to neurons in the auditory-biased frontal regions. Such heterogeneity could account for several key findings in the present study. Additionally, dynamic reconfiguration could occur with heterogeneous populations. Finer-scale analyses, possibly including those in non-human primates, are needed to distinguish between these alternatives.

This work uncovers several potentially fruitful pathways for further investigation. Firstly, understanding the representational content of each ROI and network in this analysis would help to elucidate their exact functional contribution. For example, it is unclear which ROIs/networks represent which properties of the stimuli and/or which WM processes. Secondly, in the future, other schemas for the functional organization of the frontal cortex could be added to this style of analysis, such as the language-specific frontal regions identified in Fedorenko et al. (2010, 2012). This would further integrate different lines of research into the functional organization of the frontal cortex. Lastly, further work relating behavioral measures to task-based changes in functional connectivity may have potential to be extended to clinically relevant dimensions of behavior.

This study builds on a body of work trying to make sense of the functional organization of the least clearly understood cortical region in the brain. Despite the lack of consensus on the functional organization of the frontal cortex, it is implicated in most all higher-level cognitive processes, (Amodio & Frith, 2006; Badre & Nee, 2018; De La Vega et al., 2016; Fuster, 2000)

many neurological and psychiatric disorders, (Ferguson & Gao, 2018; Friedman & Robbins,
2022; Gao et al., 2012; Goldstein & Volkow, 2011; Goto et al., 2010) as well as the natural aging
process (Giorgio et al., 2010; Nyberg et al., 2010). The results presented here drive the continued
integration of multiple models of the functional organization of the frontal cortex, in particular,
working to identify discrete functional networks for sensory-biased and supramodal processing
and beginning to uncover how these networks interact during cognitive tasks and correlate with
behavior.

Data and Code Availability

824 All custom code and ROI search spaces (freesurfer label files in fsaverage space) used in these
825 analyses are available at https://github.com/fmri/Sensory_Networks_FC.

CRediT Author Statement

Thomas Possidente: Conceptualization, Formal analysis, Methodology, Software, Validation,
Visualization, Writing - original draft, review & editing, **Vaibhav Tripathi**: Conceptualization,
Data curation, Investigation, Methodology, Software, Writing – review & editing, **Joseph T.**
McGuire: Methodology, Writing – review & editing **David Somers**: Conceptualization, Funding
acquisition, Methodology, Supervision, Project administration, Writing - original draft, review &
editing.

Funding

836 This work is funded by National Science Foundation grant BCS-1829394 to D.C.S. This work
837 involved the use of instrumentation supported by the NSF Major Research Instrumentation grant
838 BCS-1625552. Data was analyzed on a high-performance computing cluster supported by the
839 ONR grant N00014-17-1-2304.

Acknowledgments

842 We thank Dr. David Beeler for assistance with data collection and for helpful discussions, Dr.
843 Abigail Noyce for auditory stimuli, and Ryan Marshall, Dr. Stephanie McMains, and Shruthi
844 Chakrapani for scanning assistance. We acknowledge the University of Minnesota Center for
845 Magnetic Resonance Research for use of the multiband-EPI pulse sequences.

Declaration of Competing Interests

848 The authors declare no competing interest.

References

Agcaoglu, O., Wilson, T. W., Wang, Y.-P., Stephen, J., & Calhoun, V. D. (2019). Resting state

connectivity differences in eyes open versus eyes closed conditions. *Human Brain Mapping*,

40(8), 2488–2498. https://doi.org/10.1002/hbm.24539

Alhamdan, A. A., Murphy, M. J., Pickering, H. E., & Crewther, S. G. (2023). The Contribution of Visual

and Auditory Working Memory and Non-Verbal IQ to Motor Multisensory Processing in

Elementary School Children. *Brain Sciences*, *13*(2), Article 2.

https://doi.org/10.3390/brainsci13020270

Amodio, D. M., & Frith, C. D. (2006). Meeting of minds: The medial frontal cortex and social

cognition. *Nature Reviews Neuroscience*, *7*(4), 268–277. https://doi.org/10.1038/nrn1884

Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *NeuroImage*, *38*(1), 95–113.

https://doi.org/10.1016/j.neuroimage.2007.07.007

Ashburner, J., & Friston, K. (1997). Multimodal Image Coregistration and Partitioning—A Unified

Framework. *NeuroImage*, *6*(3), 209–217. https://doi.org/10.1006/nimg.1997.0290

Ashburner, J., & Friston, K. J. (2005). Unified segmentation. *NeuroImage*, *26*(3), 839–851.

https://doi.org/10.1016/j.neuroimage.2005.02.018

Assem, M., Glasser, M. F., Van Essen, D. C., & Duncan, J. (2020). A Domain-General Cognitive Core

Defined in Multimodally Parcellated Human Cortex. *Cerebral Cortex*, *30*(8), 4361–4380.

https://doi.org/10.1093/cercor/bhaa023

Baddeley, A. D., & Hitch, G. (1974). Working Memory. In G. H. Bower (Ed.), *Psychology of Learning*

and Motivation (Vol. 8, pp. 47–89). Academic Press. https://doi.org/10.1016/S0079-

7421(08)60452-1

Badre, D., & Nee, D. E. (2018). Frontal Cortex and the Hierarchical Control of Behavior. *Trends in*

Cognitive Sciences, *22*(2), 170–188. https://doi.org/10.1016/j.tics.2017.11.005

Bar-Joseph, Z., Gifford, D. K., & Jaakkola, T. S. (2001). Fast optimal leaf ordering for hierarchical

clustering. *Bioinformatics*, *17*(suppl_1), S22–S29.

https://doi.org/10.1093/bioinformatics/17.suppl_1.S22

Barr, D. J., Levy, R., Scheepers, C., & Tily, H. J. (2013). Random effects structure for confirmatory

hypothesis testing: Keep it maximal. *Journal of Memory and Language*, *68*(3), 255–278.

https://doi.org/10.1016/j.jml.2012.11.001

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method

(CompCor) for BOLD and perfusion based fMRI. *NeuroImage*, *37*(1), 90–101.

https://doi.org/10.1016/j.neuroimage.2007.04.042

Bettencourt, K. C., & Xu, Y. (2016). Decoding the content of visual short-term memory under

distraction in occipital and parietal areas. *Nature Neuroscience*, *19*(1), 150–157.

https://doi.org/10.1038/nn.4174

Bo, J., Jennett, S., & Seidler, R. D. (2011). Working memory capacity correlates with implicit serial

reaction time task performance. *Experimental Brain Research*, *214*(1), 73–81.

https://doi.org/10.1007/s00221-011-2807-8

Cauley, S. F., Polimeni, J. R., Bhat, H., Wald, L. L., & Setsompop, K. (2014). Interslice leakage artifact

reduction technique for simultaneous multislice acquisitions. *Magnetic Resonance in*

Medicine, *72*(1), 93–102. https://doi.org/10.1002/mrm.24898

Chai, X. J., Castañón, A. N., Öngür, D., & Whitfield-Gabrieli, S. (2012). Anticorrelations in resting

state networks without global signal regression. *NeuroImage*, *59*(2), 1420–1428.

https://doi.org/10.1016/j.neuroimage.2011.08.048

Christophel, T. B., Hebart, M. N., & Haynes, J.-D. (2012). Decoding the Contents of Visual Short-

Term Memory from Human Visual and Parietal Cortex. *Journal of Neuroscience*, *32*(38),

12983–12989. https://doi.org/10.1523/JNEUROSCI.0184-12.2012

Christophel, T. B., Klink, P. C., Spitzer, B., Roelfsema, P. R., & Haynes, J.-D. (2017). The Distributed

Nature of Working Memory. *Trends in Cognitive Sciences*, *21*(2), 111–124.

https://doi.org/10.1016/j.tics.2016.12.007

Cisler, J. M., Bush, K., & Steele, J. S. (2014). A comparison of statistical methods for detecting

context-modulated functional connectivity in fMRI. *NeuroImage*, *84*, 1042–1052.

https://doi.org/10.1016/j.neuroimage.2013.09.018

Crittenden, B. M., & Duncan, J. (2014). Task Difficulty Manipulation Reveals Multiple Demand

Activity but no Frontal Lobe Hierarchy. *Cerebral Cortex*, *24*(2), 532–540.

https://doi.org/10.1093/cercor/bhs333

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical Surface-Based Analysis: I. Segmentation and

Surface Reconstruction. *NeuroImage*, *9*(2), 179–194.

https://doi.org/10.1006/nimg.1998.0395

De La Vega, A., Chang, L. J., Banich, M. T., Wager, T. D., & Yarkoni, T. (2016). Large-Scale Meta-

Analysis of Human Medial Frontal Cortex Reveals Tripartite Functional Organization. *The*

Journal of Neuroscience, *36*(24), 6553–6562. https://doi.org/10.1523/JNEUROSCI.4402-

15.2016

Dice, L. R. (1945). Measures of the Amount of Ecologic Association Between Species. *Ecology*,

26(3), 297–302. https://doi.org/10.2307/1932409

Duncan, J., & Owen, A. M. (2000). Common regions of the human frontal lobe recruited by diverse

cognitive demands. *Trends in Neurosciences*, *23*(10), 475–483.

https://doi.org/10.1016/S0166-2236(00)01633-7

Eklund, A., Knutsson, H., & Nichols, T. E. (2019). Cluster failure revisited: Impact of first level design

and physiological noise on cluster false positive rates. *Human Brain Mapping*, *40*(7), 2017–

2032. https://doi.org/10.1002/hbm.24350

Eklund, A., Nichols, T. E., & Knutsson, H. (2016). Cluster failure: Why fMRI inferences for spatial

extent have inflated false-positive rates. *Proceedings of the National Academy of Sciences*,

113(28), 7900–7905. https://doi.org/10.1073/pnas.1602413113

Fedorenko, E., Duncan, J., & Kanwisher, N. (2012). Language-Selective and Domain-General

Regions Lie Side by Side within Broca's Area. *Current Biology*, *22*(21), 2059–2062.

https://doi.org/10.1016/j.cub.2012.09.011

Fedorenko, E., Hsieh, P.-J., Nieto-Castañón, A., Whitfield-Gabrieli, S., & Kanwisher, N. (2010). New

Method for fMRI Investigations of Language: Defining ROIs Functionally in Individual

Subjects. *Journal of Neurophysiology*, *104*(2), 1177–1194.

https://doi.org/10.1152/jn.00032.2010

Feinberg, D. A., Moeller, S., Smith, S. M., Auerbach, E., Ramanna, S., Glasser, M. F., Miller, K. L.,

Ugurbil, K., & Yacoub, E. (2010). Multiplexed Echo Planar Imaging for Sub-Second Whole

Brain FMRI and Fast Diffusion Imaging. *PLOS ONE*, *5*(12), e15710.

https://doi.org/10.1371/journal.pone.0015710

Ferguson, B. R., & Gao, W.-J. (2018). PV Interneurons: Critical Regulators of E/I Balance for

Prefrontal Cortex-Dependent Behavior and Psychiatric Disorders. *Frontiers in Neural*

Circuits, *12*. https://doi.org/10.3389/fncir.2018.00037

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2), 774–781.

https://doi.org/10.1016/j.neuroimage.2012.01.021

Fischl, B., Liu, A., & Dale, A. M. (2001). Automated manifold surgery: Constructing geometrically

accurate and topologically correct models of the human cerebral cortex. *IEEE Transactions*

on Medical Imaging, *20*(1), 70–80. IEEE Transactions on Medical Imaging.

https://doi.org/10.1109/42.906426

Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., Kouwe, A. van der, Killiany,

R., Kennedy, D., Klaveness, S., Montillo, A., Makris, N., Rosen, B., & Dale, A. M. (2002).

Whole Brain Segmentation: Automated Labeling of Neuroanatomical Structures in the

Human Brain. *Neuron*, *33*(3), 341–355. https://doi.org/10.1016/S0896-6273(02)00569-X

Friedman, N. P., & Robbins, T. W. (2022). The role of prefrontal cortex in cognitive control and

executive function. *Neuropsychopharmacology*, *47*(1), 72–89.

https://doi.org/10.1038/s41386-021-01132-0

Friston, K. J., Buechel, C., Fink, G. R., Morris, J., Rolls, E., & Dolan, R. J. (1997). Psychophysiological

and Modulatory Interactions in Neuroimaging. *NeuroImage*, *6*(3), 218–229.

https://doi.org/10.1006/nimg.1997.0291

Fuster, J. M. (2000). Executive frontal functions. *Experimental Brain Research*, *133*(1), 66–70.

https://doi.org/10.1007/s002210000401

Gao, W.-J., Wang, H.-X., A., M., & Li, Y.-C. (2012). The Unique Properties of the Prefrontal Cortex and

Mental Illness. In T. Mantamadiotis (Ed.), *When Things Go Wrong—Diseases and Disorders*

of the Human Brain. InTech. https://doi.org/10.5772/35868

Giorgio, A., Santelli, L., Tomassini, V., Bosnell, R., Smith, S., De Stefano, N., & Johansen-Berg, H.

(2010). Age-related changes in grey and white matter structure throughout adulthood.

NeuroImage, *51*(3), 943–951. https://doi.org/10.1016/j.neuroimage.2010.03.004

Glasser, M. F., Coalson, T. S., Robinson, E. C., Hacker, C. D., Harwell, J., Yacoub, E., Ugurbil, K.,

Andersson, J., Beckmann, C. F., Jenkinson, M., Smith, S. M., & Van Essen, D. C. (2016). A

multi-modal parcellation of human cerebral cortex. *Nature*, *536*(7615), Article 7615.

https://doi.org/10.1038/nature18933

Goldstein, R. Z., & Volkow, N. D. (2011). Dysfunction of the prefrontal cortex in addiction:

Neuroimaging findings and clinical implications. *Nature Reviews Neuroscience*, *12*(11),

652–669. https://doi.org/10.1038/nrn3119

Goto, Y., Yang, C. R., & Otani, S. (2010). Functional and Dysfunctional Synaptic Plasticity in

Prefrontal Cortex: Roles in Psychiatric Disorders. *Biological Psychiatry*, *67*(3), 199–207.

https://doi.org/10.1016/j.biopsych.2009.08.026

Griswold, M. A., Jakob, P. M., Heidemann, R. M., Nittka, M., Jellus, V., Wang, J., Kiefer, B., & Haase,

A. (2002). Generalized autocalibrating partially parallel acquisitions (GRAPPA). *Magnetic*

Resonance in Medicine, *47*(6), 1202–1210. https://doi.org/10.1002/mrm.10171

Hagler, D. J., Saygin, A. P., & Sereno, M. I. (2006). Smoothing and cluster thresholding for cortical

surface-based group analysis of fMRI data. *NeuroImage*, *33*(4), 1093–1103.

https://doi.org/10.1016/j.neuroimage.2006.07.036

Han, J., Zhou, L., Wu, H., Huang, Y., Qiu, M., Huang, L., Lee, C., Lane, T. J., & Qin, P. (2023). Eyes-

Open and Eyes-Closed Resting State Network Connectivity Differences. *Brain Sciences*,

13(1), Article 1. https://doi.org/10.3390/brainsci13010122

Harrison, S. A., & Tong, F. (2009). Decoding reveals the contents of visual working memory in early

visual areas. *Nature*, *458*(7238), 632–635. https://doi.org/10.1038/nature07832

Hartman, J. (2023). *Emmeans* [MATLAB]. https://github.com/jackatta/estimated-marginal-means

(Original work published 2019)

Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W., & Smith, S. M. (2012). FSL.

NeuroImage, *62*(2), 782–790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Linke, A. C., Vicente-Grabovetsky, A., & Cusack, R. (2011). Stimulus-specific suppression preserves

information in auditory short-term memory. *Proceedings of the National Academy of*

Sciences, *108*(31), 12961–12966. https://doi.org/10.1073/pnas.1102118108

Mayer, A. R., Ryman, S. G., Hanlon, F. M., Dodd, A. B., & Ling, J. M. (2017). Look Hear! The Prefrontal

Cortex is Stratified by Modality of Sensory Input During Multisensory Cognitive Control.

Cerebral Cortex, *27*(5), 2831–2840. https://doi.org/10.1093/cercor/bhw131

McLaren, D. G., Ries, M. L., Xu, G., & Johnson, S. C. (2012). A generalized form of context-dependent

psychophysiological interactions (gPPI): A comparison to standard approaches.

NeuroImage, *61*(4), 1277–1286. https://doi.org/10.1016/j.neuroimage.2012.03.068

Michalka, S. W., Kong, L., Rosen, M. L., Shinn-Cunningham, B. G., & Somers, D. C. (2015). Short-

Term Memory for Space and Time Flexibly Recruit Complementary Sensory-Biased Frontal

Lobe Attention Networks. *Neuron*, *87*(4), 882–892.

https://doi.org/10.1016/j.neuron.2015.07.028

Moeller, S., Yacoub, E., Olman, C. A., Auerbach, E., Strupp, J., Harel, N., & Uğurbil, K. (2010).

Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial parallel

imaging with application to high spatial and temporal whole-brain fMRI. *Magnetic*

Resonance in Medicine, *63*(5), 1144–1153. https://doi.org/10.1002/mrm.22361

Nieto-Castanon, A. (2020). *Handbook of functional connectivity Magnetic Resonance Imaging*

methods in CONN. Hilbert Press.

Noyce, A. L., Cestero, N., Michalka, S. W., Shinn-Cunningham, B. G., & Somers, D. C. (2017).

Sensory-Biased and Multiple-Demand Processing in Human Lateral Frontal Cortex. *Journal*

of Neuroscience, *37*(36), 8755–8766. https://doi.org/10.1523/JNEUROSCI.0660-17.2017

Noyce, A. L., Lefco, R. W., Brissenden, J. A., Tobyne, S. M., Shinn-Cunningham, B. G., & Somers, D.

C. (2022). Extended Frontal Networks for Visual and Auditory Working Memory. *Cerebral*

Cortex, *32*(4), 855–869. https://doi.org/10.1093/cercor/bhab249

Nyberg, L., Salami, A., Andersson, M., Eriksson, J., Kalpouzos, G., Kauppi, K., Lind, J., Pudas, S.,

Persson, J., & Nilsson, L.-G. (2010). Longitudinal evidence for diminished frontal cortex

function in aging. *Proceedings of the National Academy of Sciences*, *107*(52), 22682–22686.

https://doi.org/10.1073/pnas.1012651108

O'Reilly, J. X., Woolrich, M. W., Behrens, T. E. J., Smith, S. M., & Johansen-Berg, H. (2012). Tools of

the trade: Psychophysiological interactions and functional connectivity. *Social Cognitive*

and Affective Neuroscience, *7*(5), 604–609. https://doi.org/10.1093/scan/nss055

Patriat, R., Molloy, E. K., Meier, T. B., Kirk, G. R., Nair, V. A., Meyerand, M. E., Prabhakaran, V., & Birn,

R. M. (2013). The effect of resting condition on resting-state fMRI reliability and consistency:

A comparison between resting with eyes open, closed, and fixated. *NeuroImage*, *78*, 463–

473. https://doi.org/10.1016/j.neuroimage.2013.04.013

Penny, W. D., Friston, K. J., Ashburner, J. T., Kiebel, S. J., & Nichols, T. E. (2011). *Statistical Parametric*

Mapping: The Analysis of Functional Brain Images. Elsevier.

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2014).

Methods to detect, characterize, and remove motion artifact in resting state fMRI.

NeuroImage, *84*, 320–341. https://doi.org/10.1016/j.neuroimage.2013.08.048

Ségonne, F., Dale, A. M., Busa, E., Glessner, M., Salat, D., Hahn, H. K., & Fischl, B. (2004). A hybrid

approach to the skull stripping problem in MRI. *NeuroImage*, *22*(3), 1060–1075.

https://doi.org/10.1016/j.neuroimage.2004.03.032

Segonne, F., Pacheco, J., & Fischl, B. (2007). Geometrically Accurate Topology-Correction of

Cortical Surfaces Using Nonseparating Loops. *IEEE Transactions on Medical Imaging*, *26*(4),

518–529. IEEE Transactions on Medical Imaging. https://doi.org/10.1109/TMI.2006.887364

Serences, J. T., Ester, E. F., Vogel, E. K., & Awh, E. (2009). Stimulus-Specific Delay Activity in Human

Primary Visual Cortex. *Psychological Science*, *20*(2), 207–214.

https://doi.org/10.1111/j.1467-9280.2009.02276.x

Setsompop, K., Cohen-Adad, J., Gagoski, B. A., Raij, T., Yendiki, A., Keil, B., Wedeen, V. J., & Wald, L.

L. (2012). Improving diffusion MRI using simultaneous multi-slice echo planar imaging.

NeuroImage, *63*(1), 569–580. https://doi.org/10.1016/j.neuroimage.2012.06.033

Sladky, R., Friston, K. J., Tröstl, J., Cunnington, R., Moser, E., & Windischberger, C. (2011). Slice-

timing effects and their correction in functional MRI. *NeuroImage*, *58*(2), 588–594.

https://doi.org/10.1016/j.neuroimage.2011.06.078

Sled, J. G., Zijdenbos, A. P., & Evans, A. C. (1998). A nonparametric method for automatic correction

of intensity nonuniformity in MRI data. *IEEE Transactions on Medical Imaging*, *17*(1), 87–97.

IEEE Transactions on Medical Imaging. https://doi.org/10.1109/42.668698

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-Berg, H.,

Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K., Saunders, J., Vickers, J.,

Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M. (2004). Advances in functional and

structural MR image analysis and implementation as FSL. *NeuroImage*, *23*, S208–S219.

https://doi.org/10.1016/j.neuroimage.2004.07.051

Smith, S. M., & Nichols, T. E. (2009). Threshold-free cluster enhancement: Addressing problems of

smoothing, threshold dependence and localisation in cluster inference. *NeuroImage*, *44*(1),

83–98. https://doi.org/10.1016/j.neuroimage.2008.03.061

Sørensen, T. (1948). A method of establishing groups of equal amplitude in plant sociology based

on similarity of species and its application to analyses of the vegetation on Danish

commons. In *Biol Skrifter* (Vol. 5, p. 1). https://cir.nii.ac.jp/crid/1370302864784225926

Studholme, C., Hawkes, D. J., & Hill, D. L. G. (1998). Normalized entropy measure for multimodality

image alignment. *Medical Imaging 1998: Image Processing*, *3338*, 132–143.

https://doi.org/10.1117/12.310835

Tobyne, S. M., Brissenden, J. A., Noyce, A. L., & Somers, D. C. (2025). *Combined Auditory, Tactile,*

and Visual fMRI Reveals Sensory-Biased and Supramodal Working Memory Regions in

Human Frontal Cortex (p. 2025.06.21.660846). bioRxiv.

https://doi.org/10.1101/2025.06.21.660846

Tobyne, S. M., Osher, D. E., Michalka, S. W., & Somers, D. C. (2017). Sensory-biased attention

networks in human lateral frontal cortex revealed by intrinsic functional connectivity.

NeuroImage, *162*, 362–372. https://doi.org/10.1016/j.neuroimage.2017.08.020

Tobyne, S. M., Somers, D. C., Brissenden, J. A., Michalka, S. W., Noyce, A. L., & Osher, D. E. (2018).

Prediction of individualized task activation in sensory modality-selective frontal cortex with

'connectome fingerprinting.' *NeuroImage*, *183*, 173–185.

https://doi.org/10.1016/j.neuroimage.2018.08.007

Uluç, I., Schmidt, T. T., Wu, Y., & Blankenburg, F. (2018). Content-specific codes of parametric

auditory working memory in humans. *NeuroImage*, *183*, 254–262.

https://doi.org/10.1016/j.neuroimage.2018.08.024

van der Kouwe, A. J. W., Benner, T., Salat, D. H., & Fischl, B. (2008). Brain morphometry with

multiecho MPRAGE. *NeuroImage*, *40*(2), 559–569.

https://doi.org/10.1016/j.neuroimage.2007.12.025

Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2011). *Artifact Detection Tools (ART)*.

https://www.nitrc.org/projects/artifact_detect

Whitfield-Gabrieli, S., & Nieto-Castanon, A. (2012). Conn: A Functional Connectivity Toolbox for

Correlated and Anticorrelated Brain Networks. *Brain Connectivity*, *2*(3), 125–141.

https://doi.org/10.1089/brain.2012.0073

Xu, J., Moeller, S., Auerbach, E. J., Strupp, J., Smith, S. M., Feinberg, D. A., Yacoub, E., & Uğurbil, K.

1084   (2013). Evaluation of slice accelerations using multiband echo planar imaging at 3 T.

1085   *NeuroImage*, *83*, 991–1001. https://doi.org/10.1016/j.neuroimage.2013.07.055

1086