Can't be male without complex I? Alteration in cellular metabolism involved in
cytoplasmic male sterility in snail *Physa acuta*

Sophie Bererd[1*], Damien Roussel[1], Sandrine Plénet[1], Loïc Teulier[1], Antoine Stier[2,3$],
Emilien Luquet[1,4$*]
[1]Laboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés, CNRS UMR 5023
Université Claude Bernard Lyon 1, Université de Lyon, Lyon, France.
[2]Université de Strasbourg, CNRS, Institut Pluridisciplinaire Hubert Curien, UMR7178, 67000
Strasbourg, France
[3]Department of Biology, University of Turku, Turku, Finland
[4]Institut Universitaire de France (IUF), Paris, France
[$]shared senior authorship
[*]corresponding author: bererdsophie@gmail.com

Abstract
Cytoplasmic male sterility (CMS) is a well-known example of mitonuclear genetic
conflict over sex determination in hermaphrodite plants, where mitochondrial genes
maternally inherited sterilize the male function while biparental inherited nuclear
genes restore it. CMS has been recently discovered in animals, in the freshwater
snail *Physa acuta*. In this species, CMS is associated with two extremely divergent
mitogenomes D and K compared to the classical mitogenome N. D individuals are
male-steriles, while male fertility is restored in K individuals. We hypothesized that
the extreme divergence of mitogenomes associated with CMS can impact
mitochondrial aerobic metabolism, a necessary process in eukaryotic organisms by
which energy is transduced from food to ATP. We show that CMS is associated with
an alteration of co-encoded complex I in D male-sterile individuals, although partly
compensated by the nuclear-encoded complex II. K restored hermaphrodites have an
unaffected complex I respiration, but exhibit lower mitochondrial efficiency and higher
anaerobic contribution to cellular metabolism, highlighting an energy cost of bearing
CMS and restorer genes, which likely underlie the reduced growth of K mitotype.
How complex I alteration induces male-sterility remains to be determined, but could
be linked to oxidative stress or a defect in ATP-synthesis rate.
32
33 Keywords: mitonuclear interaction, male sterility, gynodioecy, cellular metabolism,
34 OXPHOS system, *Physa acuta*

INTRODUCTION

The gene-centred view of evolution has revolutionised modern evolutionary biology, promoting that selection occurs at the level of the gene mediated by the phenotype (Williams, 1996). Conflict can arise when genes have opposing evolutionary interests, *i.e.* different strategies to maximize their own transmission, leading to antagonistic selective pressures over the phenotype (Werren, 2011). A fabulous example of genomic conflict impacting phenotype occurs in hermaphrodites, opposing nuclear and mitochondrial genes within individuals (Cosmides & Tooby, 1981; Werren & Beukeboom, 1998). Unlike nuclear genes transmitted by both parents, mitochondrial genes are usually inherited by one parent, most often the female, meaning that their fitness only depends on the female function (Havird et al., 2019). This disparity in inheritance pattern between these genomes results in a conflict over sex determination (Chase, 2007; Cosmides & Tooby, 1981). Indeed, some selfish mitochondrial genes can emerge and sterilize the male function of hermaphrodites, who then become only functionally female. This phenomenon, known as cytoplasmic male sterility (CMS), increases the fitness of mitochondrial genes by enhancing their transmission though the female line (Budar et al., 2003; Chase, 2007; Saumitou-Laprade et al., 1994). In contrast, the male function sterilization leads to a loss of fitness for nuclear genes (Havird et al., 2019). Therefore, some nuclear genes neutralizing CMS (named restorer genes) can be selected to counteract sterilising effect of CMS genes by reverting individuals to hermaphroditism (known as restored hermaphrodite; Budar et al., 2003; Chase, 2007; Saumitou-Laprade et al., 1994). In natural populations, such a mitonuclear conflict results in a sexual polymorphism where male-sterile (female) individuals co-occur with hermaphrodite ones (normal and restored), a mating system called gynodioecy (Saumitou-Laprade et al., 1994). Gynodioecy, and the underlying dynamics of CMS and restorers genes in populations, is expected to evolve through a balance between selective advantage to be male-sterile (called female advantage, *i.e.* male-sterile individuals benefit from a better female fitness than hermaphrodites) and costs to bear CMS and restorers genes (Dufaÿ et al., 2007).

Gynodioecy was thought until recently to be a specificity of angiosperms (<1% of species, almost 20% families; Caruso et al., 2016; Dufay et al., 2014). Numerous studies highlight the association between divergent mitochondrial CMS lineages and the failure to produce pollen (review in Touzet & H. Meyer, 2014), suggesting that the functioning of mitochondria is at the core of the CMS mechanisms (*i.e.* male sterility, balance between female advantage and costs). Mitochondria are essential organelles as they are involved in various fundamental cellular functions, in particular cellular aerobic metabolism by transducing energy acquired from food to ATP (adenosine triphosphate) through oxidative phosphorylation (OXPHOS; Kadenbach, 2012; Logan, 2006; Vedel et al., 1999). The OXPHOS depends on the coordinated action of mitochondrial and nuclear genes because both encode for the proteins constituting the respiratory chain complexes and the ATP synthase (except the

complex II that is fully encoded by the nuclear genome; Kadenbach, 2012). The
divergence of CMS mitochondrial genomes may lead to altered functions through
various and poorly known mechanisms (*e.g.* breakdown of coadaptation between
mitochondrial and nuclear genomes, production of neoproteins by chimeric genes
due to new open reading frames; Delph et al., 2007). In several CMS plant species,
respiratory chain complex dysfunctions have been characterized and involve mainly
complexes composed by protein subunits encoded by both mitochondrial and nuclear
genes (Touzet & H. Meyer, 2014). For example, impairment of the complex I was
described in *Nicotiana sylvestris* male-sterile mutants (Gutierres et al., 1997; Sabar et
al., 2000) and in CMS transgenic rice (D. Tang et al., 2021). Cytochrome-c oxidase
(COX - Complex IV) activity is also reduced by 50% in leaves of male-sterile wild
beets (Ducos et al., 2001). Similarly, the activity of ATP synthase is reduced in CMS
chilli pepper (Li et al., 2012). Understanding the physiological aspects of CMS is then
crucial, as it helps to understand the mechanisms leading to male sterility and clarify
the proximal causes underlying the evolutionary dynamics of gynodioecious systems.
102
Our study uses the recently-discovered and only known CMS animal model
Physa acuta (David et al., 2022), allowing for the exploration of the mechanisms
underlying CMS from the molecular to the fitness levels. Two mitochondrial genomes
– named D and K mitotypes – have been associated with male sterility. These
mitotypes are both extremely divergent from one another (mtDNA nucleotidic
divergence ranged from 41% to 61%; Laugier et al., 2024) and from the classical
mitochondrial genome, named N mitotype (35% to 57% between K and N and 28.6%
to 53.2% between D and N; David et al., 2022; Laugier et al., 2024). D individuals are
male-steriles (*i.e.* they produced a very low amount to no spermatozoa and showed a
strong suppression of male reproductive behaviour; David et al., 2022) whereas K
individuals are restored hermaphrodites (presence of nuclear restorer genes; Laugier
et al., 2024). A recent study of Bererd et al. (2025) on the benefits and costs of CMS
in *P. acuta* provided evidence that CMS is beneficial to female fitness in the absence
of restorers (female advantage – D male-steriles have a better female fitness by
producing higher number of eggs), while it is costly in their presence (K restored
hermaphrodites have a lower growth rate). Interestingly, fitness benefits and costs
were mediated by differences in body mass, suggesting that different growth and
underlying metabolisms are likely involved. However, the whole-body metabolism
was surprisingly similar among N, D and K mitotypes (Bererd et al., 2025).
Altogether, these results suggested that the divergent mitochondrial genomes could
cause impairments on cellular metabolism, altering the life history, fitness and fertility
of organisms, but also that some compensation mechanisms may exist at the cellular
level (*e.g.* higher number of mitochondria as proposed in CMS *Nicotiana sylvestris* by
Sabar et al., 2000). The aim of the present study was thus to investigate how CMS
and restorers genes were related to cellular metabolism potentially implicated in male
sterility and balance between female advantage and costs in the *P. acuta*
gynodioecious system (N normal hermaphrodite, D male-sterile, K restored

hermaphrodite). We compared the male status (sterile or fertile), the growth rate, the
mitochondrial aerobic metabolism and the anaerobic cellular metabolism between (i)
individuals carrying CMS (D) and their non-CMS counterparts (N), and between (ii)
individuals carrying CMS and K-specific restorers genes (K) and their non-CMS
counterparts (N).

MATERIAL AND METHODS

- Experimental design

P. acuta (Physidae, Hygrophila, Gastropoda) is a cosmopolitan widespread snail
living in freshwater (Dillon et al., 2005). This species is hermaphroditic with a high
preference for outcrossing, *i.e.* eggs are self-fertilised only when no mate is available
to provide allosperm (Henry et al., 2005; Jarne et al., 2000; Tsitrone et al., 2003).

Three laboratory *P. acuta* lineages were used: the mitotype N normal hermaphrodite,
the mitotype D male-sterile and the mitotype K restored hermaphrodite. They have
been obtained from wild populations near Lyon (southeastern France; David et al.,
2022; Laugier et al., 2024) and raised in laboratory at 25°C for 20 generations for N
and K and 33 generations for D (maturity in laboratory is reached after 35-40 days at
25°C). K were mated with N to produce the first generation maintained in the
laboratory, since then they have reproduced separately. D snails (male-sterile) were
fertilised by N snails through pair-crosses at each generation to obtain D offspring.
Both K and D mitotypes were crossed with the N mitotype, resulting in all three
mitotypes sharing the same nuclear background. This indicates that the N nuclear
genome carries restorer genes that are effective in K mitotype but fail to restore CMS
in D snails.

At 11 days post-hatching, 127 snails per mitotype were placed in a temperature-
controlled room at 22.1 ± 0.1 °C and photoperiod 12h/12h. All the snails were reared
individually in small boxes (90 ml) with dechlorinated tap water changed once a week
and were fed with boiled and mixed lettuce twice a week.

- Male status and growth

At 34 days after the start of the experiment (45 days post-hatching), the male status
(sterile or fertile) of each focal individual was assessed by pairing it to a virgin albino
and by recording the pigmentation of the albino progeniture (unhatched offspring,
about one week after the pairing using a binocular microscope): pigmented offspring
indicated that the focal individual was male-fertile and on the contrary, no or albino
offspring indicated male sterility (see David et al., 2022 for more details). The albino
individuals are N mitotype, hermaphroditic snails (*i.e.* male-fertile and non-CMS),

originating from the Montpellier Area (~300km south of Lyon) and have been
maintained as a large laboratory population for over 80 generations.

Before the pairing, whole-body mass (shell + body) was measured with a precision
scale to the nearest 0.001mg. Maximal shell length was assessed using photography
analysed with the software ImageJ (Schneider et al., 2012) at the nearest 0.001mm.

Note that only N and K male-fertile individuals (normal hermaphrodite and restored
hermaphrodite, respectively) and D male-sterile individuals were selected for the
comparisons of growth (whole-body mass and maximal shell length), mitochondrial
aerobic metabolism and enzymatic activities.

- <u>Mitochondrial aerobic metabolism</u>

In order to compare the mitochondrial aerobic metabolism of the N, D and K
mitotypes, 12 individuals per mitotype (total n = 36 individuals) were selected to
assess mitochondrial functionality using high resolution respirometry (O2k, Oroboros;
DatLab software; Makrecka-Kuka et al., 2015). Snails were anaesthetised on ice and
dissected to sample the head and foot tissues (all the digestive and reproductive
tissues were delicately removed). Tissue mass was measured with a precision scale
at the nearest 0.001mg. Mechanical permeabilization of the tissue was performed by
consecutive scissors cuts in 2 ml of a Mir05 solution (Salin et al., 2016) and verified
as efficient in preliminary assays using additional chemical permeabilization with
digitonin. Mitochondrial integrity was assessed during each run by the addition of 10
µM of cytochrome c, which had very little impact on respiration rate (+1.4% ± 1.3%),
thereby proving the good integrity of our mitochondrial preparation. Four O2-k
devices were used at an assay temperature of 21.5°C and a chamber volume of
2mL, with background calibration performed once before measurement sessions
started, and 100% calibration performed daily. Mitotypes were randomly assigned to
the four O2k devices, and were evenly distributed across five sessions of
measurements. Mitochondrial aerobic metabolism was assessed using a relatively
typical substrate-inhibitor-uncoupler titration (SUIT protocol; Pesta & Gnaiger, 2011;
Fig.1).

During a pilot study aimed at setting up experimental conditions for high-
resolution respirometry on *P. acuta* homogenate, we noticed surprisingly high levels
of apparent non-mitochondrial respiration (*i.e.* after inhibition of complex III with 2.5, 5
or 10 µM of antimycin A). This residual respiration being sensitive to SHAM
(Salicylhydroxamic acid or 2-Hydroxybenzohydroxamic acid), a known inhibitor of the
alternative oxidase (AOX) pathway (Sluse & Jarmuszkiewicz, 1998), this suggests
that AOX pathway is present and active in *P. acuta*. We thus systematically inhibited
the AOX pathway at the beginning of the high-resolution respirometry assays with
1mM of SHAM, after stabilisation of endogenous cellular respiration. *AOX* respiration
was thus calculated as the difference between endogenous and SHAM-inhibited

respirations (Fig. 1B). Mitochondrial respiration was then fuelled by complex I
substrates (5mM pyruvate and 2mM malate) and ADP was added at a concentration
of 5mM to stimulate phosphorylating respiration (*CI OXPHOS*). Complex II substrate
(succinate 10mM) was then added to achieve phosphorylating respiration driven by
electron entry through both complexes I and II (*CI+II OXPHOS*). Cytochrome c 10 µM
was added to verify mitochondrial integrity (see above). ATP synthesis was
subsequently inhibited with 2.5 µM of oligomycin, leading to a non-phosphorylating
respiration (*i.e.* driven by proton leak) referred to as *CI+II LEAK*. The uncoupler
FCCP (carbonyl cyanide p-trifluoromethoxyphenyl hydrazone) was then titrated in 0.5
µM steps until maximum uncoupled respiration was achieved, which measures the
maximum capacity of the electron transport system (*ETSmax*). Inhibition of complex
III with antimycin A 2.5 µM led to mitochondrial respiration inhibition, and the residual
(*i.e.* non-mitochondrial) O2 consumption left was subtracted from other respiration
rates. Finally, electrons were fuelled directly to complex IV with 2mM ascorbate and
0.5mM TMPD (N,N,N',N'-tetramethyl- p-phenylene-diamine), before inhibition of
complex IV activity with sodium azide 100mM, the difference between these two
respiration rates corresponding to complex IV maximum capacity to use O2 (*i.e.* COX
respiration; Fig. 1B). All respiration rates were normalized per milligram of fresh
tissue and expressed as $pmolO_2.s^{-1}.mg^{-1}$ of tissue.

Mitochondrial respiration rates are sensitive to both mitochondrial quantity and
intrinsic properties (*e.g.* functionality of electron transport). We also calculated three
flux control ratios (*i.e.* ratios between different respiration rates) that are independent
from mitochondrial quantity and thus purely reflect mitochondrial properties: 1. The
coupling of oxidative phosphorylation, *i.e. OXPHOS coupling efficiency*, calculated
as: (*CI+II OXPHOS - CI+II LEAK*) / *CI+II OXPHOS*. This ratio is an index of the
proportion of O2 dedicated to ATP synthesis under phosphorylating conditions; a
ratio of 0.75 meaning that 75% of O2 consumption is coupled with ATP synthesis. *2.*
The mitochondrial reserve to increase respiration from OXPHOS to maximal ETS
respiration: *ETS reserve* calculated as (*CI+II ETS - CI+II OXPHOS*) / *CI+II ETS;* a
ratio of 0.75 meaning that mitochondrial O2 consumption can increase by 75% in
case of energy demand. *3.* The relative contribution of complex I to maximal
respiratory activity, when respiration is fuelled by both complex I and II substrates: *CI*
OXPHOS contribution, calculated as 1 - ((*CI+II OXPHOS - CI OXPHOS*) / *CI+II*
OXPHOS); a ratio of 0.75 meaning that complex I supports 75% of the mitochondrial
O2 consumption when CI and CII substrates are provided together*.*

Figure 1. **A -** Schematic of the OXPHOS system. Mitochondrial and nuclear genes
encode for the proteins constituting complexes I, III, IV and V (solid line border).
Complex II and AOX are fully encoded by nuclear genes (black dashed border). The
process relies on the respiratory chain formed by four protein complexes and the
ATP synthase where a succession of redox reactions catalysed by four complexes (I,
II, III and IV) creates a proton gradient, which is then used by complex V – ATP
synthase – for the synthesis of ATP from ADP and inorganic phosphate. Proton Leak
refers to the process where protons (H^+) cross the inner membrane independently of
ATP synthase. **B -** Graphical summary of the high-resolution respirometry substrate-
uncoupler-inhibitor-titration (SUIT) protocol. *ENDO* respiration rate reflects the
endogenous respiration of cells. *AOX* (alternative oxidase) respiration corresponds to
the respiratory rate before inhibition by SHAM (Salicylhydroxamic acid or 2-
Hydroxybenzohydroxamic acid). *CI OXPHOS* respiration rate fuelled by Complex I
was measured after addition of its substrates Pyruvate and Malate with ADP (PM +
ADP) and corresponds to the respiratory rate of the system with electron entry only
through Complex I. *CI+II OXPHOS* respiration rate fuelled by Complexes I and II was

measured after addition of Complex II's substrates Succinate (S) and reflects the
respiratory rate of the whole substrate-saturated system. *CI+II LEAK* respiration rate
reflects the proton leak assessed by inhibiting Complex IV with Oligomicyn (O).
ETSmax corresponds to the maximum capacity of the electron transport system and
was measured after the titration of uncoupler FCCP (carbonyl cyanide p-
trifluoromethoxyphenyl hydrazone). *COX* reflects the Complex IV maximum capacity
to use O2. *NON-MITO* reflects the non-mitochondrial respiration rate, measured by
the addition of antimycin A (AA).

-   Relative contribution of anaerobic metabolism to aerobic metabolism

The relative contribution of anaerobic metabolism to aerobic metabolism in the three
mitotypes was measured by comparing enzymatic activities of the lactate
dehydrogenase (LDH) and the citrate synthase (CS), respectively. LDH is a key
enzyme of anaerobic metabolism, by converting pyruvate into lactate, it regenerates
the NAD+ used by the glycolysis pathway to oxidize glucose-6-phosphate and
synthesize ATP in the absence of oxygen (Emmett & Hochachka, 1981). CS is a key
enzyme of mitochondrial aerobic metabolism as it is involved in the tricarboxylic acid
cycle, which fuel the mitochondrial electron transport chain with reduced coenzymes
(NADH and FADH2; Emmett & Hochachka, 1981; Krebs, 1970; Srere, 1971).

The head and the foot of 20 snails per mitotype (total n = 60 individuals) were
sampled, weighted and frozen at -80°C. Tissues samples were thawed, then
homogenised using a glass piston-glass Potter in 500µl buffer containing 100 mM
sucrose, 50 mM KCl, 5 mM EDTA, 50 mM Tris-base, 1% fatty acid free BSA, pH 7.4.
The resulting homogenates were kept at -80°C until measurements. After thawing,
homogenates were vortexed and centrifuged at 1000 g for 10 min (4°C). The
supernatants were retrieved and used for the LDH and CS enzyme assays. Enzyme
activities were measured spectrophotometrically at 22°C. All results were expressed
as $nmol.min^{-1}.mg^{-1}$ of tissue.

- **LDH activity** was measured in a reaction medium composed of 40mN imidazole
(pH 7), 0.04% BSA and 200 µM NADH. Two replicates per homogenates were
performed with 10µl and 20µl of supernatants and 250µl of the LDH reaction medium
were laid in a microtiter plate. After 3 min incubation, the reaction was initiated by
adding 4 mM pyruvate and the change in optical density at 340 nm was recorded for
3 min. The enzyme activity was quantified using an extinction coefficient of $6.22 \, mM^{-1}.cm^{-1}$.
297

- **CS activity** was measured in a reaction medium composed of 100 mM Tris buffer
(pH 8), 100 µM 5,5'-dithiobis(2-nitrobenzoic acid), 300 µM acetyl-CoA and 0.1%
Triton-X-100. Homogenates were diluted ½ in a 100 mM phosphate buffer. Two
replicates of 5µl and 7µl of diluted homogenates and 250µl of the CS reaction
medium were deposited in a microtiter plate. After 3 min incubation, the reaction was

initiated by adding 500 µM oxaloacetate and the change in optical density at 412 nm
was recorded for 3 min. The enzyme activity was quantified using an extinction
coefficient of 13.6 $mM^{-1}.cm^{-1}$.

- The **LDH/CS ratio** was calculated to estimate the contribution of anaerobic
metabolism relative to aerobic metabolism and used as an index of glycolytic
capacity.

- <u>Statistical analyses</u>

Male status - A generalized linear model (GLM) following a binomial distribution was
used to test the proportion of male sterility in the three mitotype by an analysis of
variance (ANOVA).

Growth - The effect of the mitotype on whole body mass and on maximal shell length
were tested by a linear model using an ANOVA.

Mitochondrial aerobic metabolism - Linear mixed models (LMM) were conducted to
analyse mitochondrial aerobic metabolism parameters with "mitotype" as a fixed
factor, the O2k chamber identity as a random factor to control for potential technical
variations between the eight chambers used, and "session (day of measurement)" as
a random factor to control for technical variations between measurement sessions.
The significance of fixed effects was tested using F-tests and the Kenward-Roger
approximation for the degrees of freedom.

Enzymatic activities – Means of the two replicates for both LDH and CS activities
were used in statistical analyses. LDH/CS ratio was calculated as the average of
LDH means and CS means. Linear models and ANOVAs were conducted to analyse
the effect of the mitotype on LDH activity, CS activity and LDH/CS ratio.

All statistical analyses were carried out in Rstudio 4.3.2 (R Core Team, 2023). Linear
mixed models were performed using *lme4* package (Bates et al., 2015; p-values were
obtained using *LmerTest* package Kuznetsova et al., 2017). Post-hoc tests with
Tukey correction were performed using the *emmeans* package (Lenth, 2024) when
the mitotype factor was significant ($p < 0.05$).

RESULTS

- <u>Male status</u>

Male sterility depended on mitotype ($\chi^2_{2, 337}$ = 277.43, $p < 0.001$; Fig. 2). D individuals
were almost all male-sterile as D had a much lower male-fertility proportion than N
(normal hermaphrodite) and K (restored hermaphrodite; in logit scale, N - D = 6.25 ±
0.79, z = 7.94, $p < 0.0001$; D - K = -6.29 ± 0.79, z = 8.00, $p < 0.001$). N and K were

mostly male-fertile and not significantly different from each other (in logit scale, N – K
= -0.045 ± 0.467, z = -0.096, p = 0.995).

<u>Figure 2.</u> Percentage of male-fertile and male-sterile individuals in N, D and K
P.acuta mitotypes. The total sample sizes were 124 N, 123 D and 127 K.

- <u>Growth</u>

Whole-body mass was significantly different between mitotypes ($F_{2, 341}$ = 128.79, p
< 0.001; Fig. 3A). D male-sterile snails were bigger than N and K male-fertiles (N - D:

Est.±SE = -0.013 ± 0.003, t_{341} = -4.98, p < 0.001; D - K: Est.±SE = 0.042 ± 0.003, t_{341}
= 15.78, p < 0.001). K had a significantly lower whole-body mass than N and D (N -
K: Est.±SE = 0.029 ± 0.003, t_{341} = 10.46, p <0.001; Fig. 3A).

Maximal shell length was also significantly different between mitotypes ($F_{2, 338}$ =
88.50, p < 0.001; Fig. 3B). D male-sterile had a longer shell length than N and K
male-fertiles (N - D: Est.±SE = -0.662 ± 0.105, t_{338} = -6.30, p < 0.001; D - K: Est.±SE
= 1.383 ± 0.104, t_{338} = 13.30, p <0.001). K had a significantly smaller shell length
than N and D (N - K: Est.±SE = 0.720 ± 0.107, t_{338} = 6.75, p < 0.001; Fig. 3B).

Figure 3. **A** – Whole-body mass (in g) for *P. acuta* normal hermaphrodites N (red
circle), restored hermaphrodites K (blue square) and male-steriles D (green triangle).
B – Maximal shell length (in mm). Black dots and black errors bars correspond to
means ± standard error.

-    **Mitochondrial aerobic metabolism**

Mitochondrial respiration rates

Mitochondrial respiration rates related to the alternative oxidase (*AOX*),
oxidative phosphorylation through complex I and II combined (*CI+II OXPHOS*), and
maximal uncoupled respirations (*CI+II ETS*), as well as maximal oxidative capacity
achieved under direct electron fueling to complex IV (*COX*) did not statistically differ
between mitotypes (Tab. 1, Fig. 4A). There was a marginally significant trend for
mitotype to affect oxidative phosphorylation through complex I only (*CI OXPHOS*;
Tab. 1, Fig. 4A), with the D mitotype having lower respiration although non-
significantly according to post-hoc tests (D - N: $t_{7.2}$ = -2.09, p = 0.16; D - K: $t_{5.3}$ = -
2.59, p = 0.10; Fig. 3A). Mitochondrial respiration rate related to proton leak through
complex I and II (*CI+II LEAK*; Tab. 1) was significantly affected by mitotype, with K
mitotype having higher leak although non-significantly according to post-hoc tests (K
vs. N: $t_{6.5}$ = 2.06, p = 0.17; K vs. D: $t_{5.2}$ = 2.44, p = 0.12; Fig. 4A).

<u>Table 1.</u> Results of LMM testing the effect of mitotype (N normal hermaphrodites, K
restored hermaphrodites, and D male-steriles) on mitochondrial respiration rates.

Model	Effect	NumDF, DenDF	F	P
AOX	Mitotype	2, 7.75	1.68	0.25
CI+II OXPHOS	Mitotype	2, 7.56	0.79	0.49
CI+II ETS	Mitotype	2, 29.56	0.69	0.51
COX	Mitotype	2, 5.42	0.48	0.64
CI OXPHOS	Mitotype	2, 7.70	4.19	0.058
CI+II LEAK	Mitotype	2, 29.30	3.87	**0.032**

382 Figure 4. **A –** Mitochondrial respiration rates and **B –** Flux control ratios between
383 mitotypes: N normal hermaphrodites (red), D male-steriles (green) and K restored
384 hermaphrodites (blue) and. Figures are based on marginal means ± SE from
385 statistical models and different letters indicate significant differences according to
386 LMM models and associated post-hoc tests.

Mitochondrial flux control ratios

The coupling efficiency of the OXPHOS system to ATP synthesis (*OXPHOS*
coupling) was significantly affected by mitotype ($F_{2,28.7}$ = 9.20, p < 0.001), with K
mitotype having lower coupling efficiency than both N ($t_{6.7}$ = 3.66, p = 0.021) and D
mitotypes ($t_{5.3}$ = 3.21, p = 0.049; Fig. 4B). The ability to increase respiration from
phosphorylating to uncoupled conditions did not significantly differ between mitotypes
(*ETS reserve*: $F_{2,10.1}$ = 0.70, p = 0.52). Finally, the contribution of complex I to
OXPHOS respiration was markedly affected by mitotype ($F_{2,6.0}$ = 17.90, p = 0.003),
with D individuals having lower contribution of complex I (D - N: $t_{7.2}$ = 4.77, p = 0.005;
D - K: $t_{5.3}$ = 5.02, p = 0.008; Fig. 4B).

-     Relative contribution of anaerobic metabolism to aerobic metabolism

LDH activity was significantly influenced by the mitotype ($F_{2,57}$ = 6.18, p = 0.004). D
and K individuals had a significantly lower LDH activity than N (D - N: Est.±SE = 0.55
± 0.17, t_{57} = 3.33, p = 0.004; K - N: Est.±SE = 0.44 ± 0.17, t_{57} = 2.65, p = 0.028). LDH
activity in mitotypes K and D was not significantly different from each other (D – K:
Est.±SE = -0.11 ± 0.17, t_{57} = -0.68, p = 0.78; Fig. 5)

CS activity was significantly influenced by the mitotype ($F_{2,57}$ = 25.34, p < 0.001). K
individuals had a significantly lower CS activity than N and D (N - K: Est.±SE = 0.77 ±
0.11, t_{57} = 6.86, p < 0.0001; D - K: Est.±SE = 0.57 ± 0.11, t_{57} = 5.09, p < 0.001). CS
activity in mitotypes N and D was not significantly different from each other (N – D:
Est.±SE = 0.20 ± 0.11, t_{57} = 1.78, p = 0.19 ; Fig. 5).

The anaerobic contribution to metabolism (ratio of LDH activity to CS activity)
was significantly influenced by the mitotype ($F_{2, 57}$ = 23.98, p < 0.001), being higher in
K mitotype compared to N and D (N - D = Est.±SE = 0.47 ± 1.17, t_{57} = 0.41, p = 0.91;
N - K: Est.±SE = -6.76 ±1.17, t_{57} = -5.78, p <.001; D - K: Est.±SE = -7.24 ± 1.17, t_{57} =
-6.19, p < 0.001 ; Fig. 5).

<u>Figure 5.</u> Enzymatic activities of lactate dehydrogenase (anaerobic metabolism) and
citrate synthase (aerobic metabolism) of *P. acuta* N normal hermaphrodites (red), D
male-steriles (blue) and K restored hermaphrodites (green). Means ± SE and
different letters indicate significant differences according to ANOVAs and associated
post-hoc tests.

DISCUSSION
Cytoplasmic male sterility (CMS) originates from a mitonuclear conflict where
mitochondrial genes induce male sterility and nuclear genes restore male fertility in
hermaphrodites. The dynamics of CMS and restorers genes is known to evolve
through a balance between selective advantage to be male-sterile (*i.e.* female
advantage) and costs to bear CMS and restorers genes (Dufaÿ et al., 2007). Male

sterility and balance between female advantage and costs – key parameters of
gynodioecy evolutionary dynamics – may be conditioned by changes in cellular
metabolism. This study uniquely combined measurements of phenotype (growth trait
as proxy of fitness advantage and costs) with an in-depth investigation of cellular
metabolism mechanisms in the *P. acuta* gynodioecious system (N normal
hermaphrodite, D male-sterile, K restored hermaphrodite; David et al., 2022; Laugier
et al., 2024). Our results provide evidence that CMS in D male-sterile individuals was
associated with a reduction of co-encoded complex I contribution to OXPHOS that
was nonetheless partly compensated by nuclear-encoded complex II. Moreover, the
efficiency to produce ATP was similar to N normal hermaphrodite which might explain
the female advantage (higher growth). Such changes were not observed in K
restored hermaphrodites, suggesting that male fertility restorer genes re-established
the OXPHOS metabolism through complex I to a similar level as N normal
hermaphrodites. Moreover, the K restored hermaphrodites showed a reduced growth,
reflecting the costs of bearing CMS and restorer genes, which may be at least partly
explained by the lower efficiency to produce ATP and the higher anaerobic
contribution to cellular metabolism that we observed. In the following, we discussed
in detail these cellular mechanisms of CMS and their implications for its evolution.
447
448
449 **Effects of CMS on reproductive functions and cellular metabolism in male-**
450 **steriles**
451
Male fertility and female advantage
The conflict opposing the D mitogenome and the nuclear genome results in male
sterility in almost all D snails as observed in previous studies (Bererd et al., 2025;
David et al., 2022). Less than 2% of D individuals were considered male-fertiles (*i.e.*
they fertilised their albino partner), confirming an incomplete penetrance of CMS
genes (no restorers are yet found for D mitotype; Bererd et al., 2025; David et al.,
2022; Skarlou et al., 2024). The D male-sterile snails had a higher growth, both in
terms of mass and length, than N normal hermaphrodite. Bererd et al., (2025)
showed that the growth advantage of D male-steriles led to a higher egg production,
confirming the expected female advantage in male-sterile individuals required to the
maintenance of CMS in natural populations (Dufaÿ et al., 2007). Such a female
advantage has been described in plants (Dufay & Billard, 2012) and is usually
explained as a consequence of an energy reallocation from the defective male
function toward the female fitness (Budar et al., 2003).
466
Complex I impairment and Complex II compensation
Mitonuclear incompatibilities, particularly when mitochondrial and nuclear genomes
have not coevolved, may disturb the integrity of OXPHOS with downstream
consequences on many cellular functions (Rand et al., 2004; Sunnucks et al., 2017;
Wolff et al., 2014). For example, genetically perturbed mitonuclear co-evolved system

by introducing a foreign mitogenome from non-human apes (common chimpanzee,
pigmy chimpanzee and gorilla) into a human nuclear background resulted in a
significant decreased complex I activity and supported mitochondrial respiration,
whereas others mitochondrial co-endoded complexes were less affected (Barrientos
et al., 1998). Similarly, a foreign mitogenome from rat into a mouse nuclear
background resulted in a large decrease of complex I activity, with a lesser impact on
co-encoded complex IV activity (McKenzie & Trounce, 2000). Our results are
strikingly consistent as the complex I activity *per se* was lower in D mitotype than in N
normal hermaphrodite, whereas complex IV activity remained unaffected. This
provides evidence that CMS in *P. acuta* is mainly associated with an alteration of co-
encoded complex I, which likely result from incompatibilities between the extremely
divergent D mitogenome and a shared nuclear genome (D snails have to mate with N
hermaphrodite snails to produce offsprings; David et al., 2022). Moreover, David et
al. (2022) previously showed that the genes coding for NAD proteins (*i.e.* the
mitochondrial genes encoding for complex I subunits) exhibited the highest levels of
divergence (*e.g.* 54-78% of amino acid dissimilarity for NAD genes of complex I
against 19-50% for COX genes of complex IV; see Table S1 for more details). This
might explain why the mitonuclear incompatibilities specifically altered the complex I
function rather than other co-encoded complexes. Our result is also consistent with
studies in CMS plants showing mitochondrial mutations also affecting complex I:
tobacco (Gutierres et al., 1997), rice (D. Tang et al., 2021), maize (Marienfeld &
Newton, 1994; H. Tang et al., 2017).

Interestingly, although complex I respiration was impaired in D male-steriles,
the combined activity of complexes I and II was similar to N normal hermaphrodites.
These results provide evidence that OXPHOS can offset the dysfunction of complex I
by compensating through complex II in D male-sterile individuals (at least, in terms of
O_2 consumption). A similar offset of complex I dysfunction by complex II has been
observed in male-steriles in CMS *Nicotina sylvestris,* where alternative NAD(P) H
dehydrogenases in association with complex II compensated for the complex I
dysfunction, ensuring the plant survival and development (Sabar et al., 2000). As no
divergence in the nuclear genes coding for mitochondrial proteins between D male-
steriles and N normal hermaphrodites have been reported (David et al., 2022), this
suggests that the maintenance of OXPHOS integrity through complex II
compensation is not a result of an evolutionary rescue by the nuclear genome, but
rather reflects plastic responses within mitochondrial mechanisms. Moreover, we did
not detect other compensatory mechanisms: the relative contribution of anaerobic
metabolism (LDH/CS ratio) was similar between D male-steriles and N normal
hermaphrodites. Consequently, our results suggest that the plasticity of the
mitochondrial respiratory chain in response to complex I impairment due to CMS,
allowed the maintenance of mitochondrial respiration at the cellular level. This finding
explains the similar whole-body metabolic rate previously observed between D and N
snails (Bererd et al., 2025).

515
Consequences of OXPHOS offset on male fertility and female advantage
The compensation of complex I dysfunction is consistent with a similar OXPHOS
coupling (*i.e.* the coupling efficiency of the OXPHOS system to ATP synthesis)
between D male-steriles and N normal hermaphrodites, suggesting that ATP
production might not be different between mitotypes. The maintenance of ATP
production in D male-steriles may explain the female advantage conferred by a
reallocation of energy from the defective male reproductive function toward growth,
thereby enhancing egg production (female fitness; Bererd et al., 2025). However, the
dysfunction of complex I of the respiratory chain and the compensation by complex II
in D male-sterile individuals could have consequences at the cellular level that may
contribute to the sterilization of the male function. Electron entry through complex II
into the respiratory chain results in reduced ATP yield and higher reactive oxygen
species (ROS) production than electron entry through complex I (Lee et al., 1996;
Quinlan et al., 2012). In CMS plant studies, a decrease in ATP production is often
proposed to explain the pollen abortion as gametogenesis is highly energy-
consuming (Touzet & H. Meyer, 2014). While some examples were consistent with
this hypothesis (*e.g.* Li et al., 2012), others did not find such a relationship (*e.g.*
Robison et al., 2009; Wen et al., 2003). Similarly, defective ATP production through
mitochondrial respiration in animals could lead to disturbed male gametogenesis or
spermatozoa performances (Yu et al., 2019) as these cells have limited cytoplasmic
space restricting the amount of mitochondria. Our previous studies showed that D
male-sterile have a very low spermatozoa count, often none at all, indicating rather
problems during gametogenesis than lower spermatozoa performances (Bererd et
al., 2025; David et al., 2022). Here, our results suggest that ATP production might not
be different between N and D mitotypes, thereby could be not linked to male sterility.
Another hypothesis for the mechanisms underlying CMS involves the production of
ROS, which may contribute to male sterility (Touzet & H. Meyer, 2014). ROS are
molecules that can damage nucleic acids, lipids and proteins, and are involved in cell
death (Logan, 2006). In CMS plants, a burst of ROS has been observed in several
species (cotton, Jiang et al., 2007; peach, Cai et al., 2021; wheat, Liu et al., 2018)
and related to pollen abortion through different mechanisms involving the initiation of
cell death process (Touzet & H. Meyer, 2014). In animals, ROS are implicated in
reproductive disorders (*e.g.* male infertility in humans; Aitken et al., 2022). Further
investigations directly measuring the mitochondrial ATP synthesis and mitochondrial
coupling efficiency (*e.g.* ATP/O ratio), as well as the mitochondrial ROS production,
antioxidant defences and oxidative damage, would be informative to confirm these
hypotheses and are necessary to unravel the role of ATP and ROS production in
male sterility.
554
555

556 **Effects of nuclear restoration on reproductive functions and cellular**
557 **metabolism in restored hermaphrodites**

Male fertility and costs of CMS and restoration in restored hermaphrodites

The major proportion of K individuals were male fertile, confirming their status of
restored hermaphrodites. Previous study provided evidence that nuclear restorer
genes counteract the sterilizing effect of CMS genes in the K mitogenome (Laugier et
al., 2024). Few K individuals (almost 11%) were considered as male-steriles by our
male status test (pairing with a virgin albino partner) but this was likely due to a
mating failure independently of the spermatozoa production (see Bererd et al., 2025
for more details). As previously described in Bererd et al. (2025), K restored
hermaphrodites showed a lower growth (smaller in mass and length) than N normal
hermaphrodites, suggesting that bearing CMS and restorer genes can be costly.
Such costs are expected by theoretical models to maintain gynodioecy in populations
(*e.g.* restoration costs avoid fixation of nuclear restorers leading to 100%
hermaphrodite populations; Dufaÿ et al., 2007). The cost of restoration has been
already observed in plants, not in terms of growth impairment, but through
detrimental effects on seed (De Haan et al., 1997) or pollen production (Bailey, 2002)
or both (Gigord et al., 1999).

Normal contribution of Complex I and Complex II to OXPHOS

The mechanisms of male fertility restoration by nuclear genes is complex as it can
occur at various molecular levels, from genomic to metabolic levels, and consist in
either the inactivation of the sterilizing factor (suppressing the expression of CMS
genes) or by offsetting its effects (Bhattacharya et al., 2024 for a recent review). For
example, in the CW-CMS rice, the nuclear restorer gene Rf17 encodes for a
mitochondrial protein causing metabolic changes and restores the male fertility
through poorly known mechanisms (Fujii & Toriyama, 2008). In the *P. acuta* system,
our results showed that K restored hermaphrodites had similar complex I and
complex II contributions to OXPHOS than in N normal hermaphrodites despite their
extremely divergent mitochondrial genomes. Similarly than in D male-sterile snails,
the NAD genes (encoding for complex I subunits) in K restored hermaphrodites are
the mitochondrial genes with the highest divergence (0.75-0.89% of amino acid
dissimilarity for NAD genes against 0.43-0.7% for COX genes of complex IV; see
Table S1 for more details; Laugier et al., 2024). Assuming that mechanism by which
CMS genes act to sterilize the male reproductive function in K mitotype are similar
than in D male-steriles, our results suggest that nuclear restorer genes in K snails
may have restored the complex I function, avoiding the need for complex II
compensation to maintain mitochondrial respiration and the associated detrimental
consequences for fertility. To confirm that the nuclear restorer genes act on the
complex I contribution to OXPHOS, further studies should explore the mitochondrial
function of K male-steriles
(*i.e.* non-restored), to test for OXPHOS dysfunctions as observed in D male-steriles.

Decrease of OXPHOS coupling efficiency as a cause of CMS and restoration costs

Although complexes I and II contributions to OXPHOS were similar to N normal
hermaphrodites, the OXPHOS coupling efficiency was lower due to a higher proton
leak in K restored hermaphrodites. Indeed, a lower mitochondrial efficiency has been
shown to limit growth rate in other study systems (*e.g.* Dawson et al., 2022; Salin et
al., 2012, 2019). We previously showed that the lower growth of K restored
hermaphrodite was not associated with a decrease of egg production or number of
spermatozoa (Bererd et al., 2025). Therefore, with the lower amount of ATP
available, the K restored hermaphrodites are likely unable to maintain the same
investment as N normal hermaphrodites in both reproduction and growth, and we can
reasonably assume that they thus invest more in reproduction at the expense of
growth. The anaerobic metabolism (LDH/CS ratio) was higher in K mitotype, but this
anaerobic pathway might not be sufficient to offset the poorly mitochondrial aerobic
ATP synthesis and meet energy demands. To our knowledge, these are the first
results showing the cellular mechanisms underlying CMS and restoration costs.
However, further investigations will be needed to disentangle the relative effects of
CMS and nuclear restorer genes on the energy tradeoff between growth and
reproduction.
618
619

Conclusion
Our results show that CMS was associated with a decreased functionality of
complex I in D male-steriles compensated through complex II, allowing an apparently
normal aerobic respiration. CMS mechanisms are complex as only male gametes are
impaired in D mitotype, while female function remains functional. Similar OXPHOS
coupling efficiency between N hermaphrodites and D male-steriles suggest that ATP
production might not be different between these mitotypes, possibly explaining the
female advantage from energy reallocation from defective male reproductive function
toward growth which indirectly enhanced egg production (Bererd et al., 2025). K
restored hermaphrodites showed a similar mitochondrial aerobic metabolism as N
normal hermaphrodites, suggesting that restorer genes may restore the complex I
function. However, K restored hermaphrodites exhibit lower mitochondrial efficiency
to produce ATP and a higher contribution of anaerobic metabolism, likely explaining
the lower growth associated with the costs of bearing CMS and restorer genes.
These results support that mitochondria function plays a central role in key
parameters underlying the evolutionary dynamics involved in gynodioecy, and
positioned the *P. acuta* system as a new model to study the molecular mechanisms
of mitonuclear conflicts involved in CMS.

641
642
643

644
645
646 F*unding*
647 This work was financially supported by the MINIGAN (ANR-19-CE02-0017) and the
648 TEATIME (ANR-21-CE02-0005) grants from the French National Research Agency
649 (ANR).

650
Acknowledgement
We are grateful to Noéline Garcia for her support with growth measurements and the
preparation of individuals for the male sterility test. We also thank Fanny Laugier for
conducting the analysis of amino acid divergences among mitotypes. We thank
Patrice David for his valuable discussions on cytoplasmic male sterility and for
leading the MINIGAN project.

657
References

Aitken, R. J., Bromfield, E. G., & Gibb, Z. (2022). OXIDATIVE STRESS AND
REPRODUCTIVE FUNCTION: The impact of oxidative stress on reproduction: a focus on
gametogenesis and fertilization. *Reproduction*, *164*(6), F79–F94.
https://doi.org/10.1530/REP-22-0126

Bailey, M. F. (2002). A Cost of Restoration of Male Fertility in a Gynodioecious Species,
Lobelia Siphilitica. *Evolution*, *56*(11), 2178–2186. https://doi.org/10.1111/j.0014-
3820.2002.tb00142.x

Barrientos, A., Kenyon, L., & Moraes, C. T. (1998). Human xenomitochondrial cybrids.
Cellular models of mitochondrial complex I deficiency. *The Journal of Biological Chemistry*,
273(23), 14210–14217. https://doi.org/10.1074/jbc.273.23.14210

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software*, *67*, 1–48. https://doi.org/10.18637/jss.v067.i01

Bererd, S., David, P., Teulier, L., Plénet, S., & Luquet, E. (2025). Temperature-dependent
benefits and costs of cytoplasmic male sterility in snail Physa acuta. *Proceedings of the
Royal Society B: Biological Sciences*, *292*(2038), 20241762.
https://doi.org/10.1098/rspb.2024.1762

Bhattacharya, J., Nitnavare, R. B., Bhatnagar-Mathur, P., & Reddy, P. S. (2024).
Cytoplasmic male sterility-based hybrids: Mechanistic insights. *Planta*, *260*(4), 100.
https://doi.org/10.1007/s00425-024-04532-w

Budar, F., Touzet, P., & De Paepe, R. (2003). The Nucleo-Mitochondrial Conflict in
Cytoplasmic Male Sterilities Revisited. *Genetica*, *117*, 3–16.
https://doi.org/10.1023/A:1022381016145

Cai, Y., Ma, Z., Ogutu, C. O., Zhao, L., Liao, L., Zheng, B., Zhang, R., Wang, L., & Han, Y. (2021). Potential Association of Reactive Oxygen Species With Male Sterility in Peach. *Frontiers in Plant Science*, *12*. https://www.frontiersin.org/articles/10.3389/fpls.2021.653256

Caruso, C. M., Eisen, K., & Case, A. L. (2016). An Angiosperm-Wide Analysis of the Correlates of Gynodioecy. *International Journal of Plant Sciences*, *177*(2), 115‐121. https://doi.org/10.1086/684260

Chase, C. D. (2007). Cytoplasmic male sterility: A window to the world of plant mitochondrial–nuclear interactions. *Trends in Genetics*, *23*(2), 81‐90. https://doi.org/10.1016/j.tig.2006.12.004

Cosmides, L., & Tooby, J. (1981). Cytoplasmic inheritance and intragenomic conflict. *J. Theor. Biol.*, *89*, 83‐129.

David, P., Degletagne, C., Saclier, N., Jennan, A., Jarne, P., Plénet, S., Konecny, L., François, C., Guéguen, L., Garcia, N., Lefébure, T., & Luquet, E. (2022). Extreme mitochondrial DNA divergence underlies genetic conflict over sex determination. *Current Biology*, *32*(10), 2325-2333.e6. https://doi.org/10.1016/j.cub.2022.04.014

Dawson, N. J., Millet, C., Selman, C., & Metcalfe, N. B. (2022). Inter-individual variation in mitochondrial phosphorylation efficiency predicts growth rates in ectotherms at high temperatures. *The FASEB Journal*, *36*(6), e22333. https://doi.org/10.1096/fj.202101806RR

De Haan, A. A., Hundscheid, M. P. J., & Van Hinsberg, A. (1997). Effects of CMS types and restorer alleles on plant performance in Plantago lanceolata L.: An indication for cost of restoration. *Journal of Evolutionary Biology*, *10*(5), 803‐820. https://doi.org/10.1046/j.1420-9101.1997.10050803.x

Delph, L., Touzet, P., & Bailey, M. (2007). Merging theory and mechanism in studies of gynodioecy. *Trends in Ecology & Evolution*, *22*(1), 17‐24. https://doi.org/10.1016/j.tree.2006.09.013

Dillon, R. T., Wethington, A. R., Rhett, J. M., & Smith, T. P. (2005). Populations of the European freshwater pulmonate Physa acuta are not reproductively isolated from American Physa heterostropha or Physa integra. *Invertebrate Biology*, *121*(3), 226‐234. https://doi.org/10.1111/j.1744-7410.2002.tb00062.x

Ducos, E., Touzet, P., & Boutry, M. (2001). The male sterile G cytoplasm of wild beet displays modified mitochondrial respiratory complexes. *The Plant Journal*, *26*(2), 171‐180. https://doi.org/10.1046/j.1365-313x.2001.01017.x

Dufay, M., & Billard, E. (2012). How much better are females? The occurrence of female advantage, its proximal causes and its variation within and among gynodioecious species. *Annals of Botany*, *109*(3), 505‐519. https://doi.org/10.1093/aob/mcr062

Dufay, M., Champelovier, P., Käfer, J., Henry, J. P., Mousset, S., & Marais, G. A. B. (2014). An angiosperm-wide analysis of the gynodioecy–dioecy pathway. *Annals of Botany*, *114*(3), 539‐548. https://doi.org/10.1093/aob/mcu134

Dufaÿ, M., Touzet, P., Maurice, S., & Cuguen, J. (2007). Modelling the maintenance of male-fertile cytoplasm in a gynodioecious population. *Heredity*, *99*(3), Article 3. https://doi.org/10.1038/sj.hdy.6801009

Emmett, B., & Hochachka, P. W. (1981). Scaling of oxidative and glycolytic enzymes in
mammals. *Respiration Physiology*, *45*(3), 261‐272. https://doi.org/10.1016/0034-
5687(81)90010-4

Fujii, S., & Toriyama, K. (2008). Genome Barriers between Nuclei and Mitochondria
Exemplified by Cytoplasmic Male Sterility. *Plant and Cell Physiology*, *49*(10), 1484‐1494.
https://doi.org/10.1093/pcp/pcn102

Gigord, Lavigne, Shykoff, & Atlan. (1999). Evidence for effects of restorer genes on male and
female reproductive functions of hermaphrodites in the gynodioecious species Thymus
vulgaris L. *Journal of Evolutionary Biology*, *12*(3), 596‐604. https://doi.org/10.1046/j.1420-
9101.1999.00056.x

Gutierres, S., Sabar, M., Lelandais, C., Chetrit, P., Diolez, P., Degand, H., Boutry, M., Vedel,
F., de Kouchkovsky, Y., & De Paepe, R. (1997). Lack of mitochondrial and nuclear-encoded
subunits of complex I and alteration of the respiratory chain in Nicotiana sylvestris
mitochondrial deletion‐mutants. *Proceedings of the National Academy of Sciences*, *94*(7),
3436‐3441. https://doi.org/10.1073/pnas.94.7.3436

Havird, J. C., Forsythe, E. S., Williams, A. M., Werren, J. H., Dowling, D. K., & Sloan, D. B.
(2019). Selfish Mitonuclear Conflict. *Current Biology*, *29*(11), R496‐R511.
https://doi.org/10.1016/j.cub.2019.03.020

Henry, P.-Y., Bousset, L., Sourrouille, P., & Jarne, P. (2005). Partial selfing, ecological
disturbance and reproductive assurance in an invasive freshwater snail. *Heredity*, *95*(6),
Article 6. https://doi.org/10.1038/sj.hdy.6800731

Jarne, P., Perdieu, M. A., Pernot, A. F., Delay, B., & David, P. (2000). The influence of self-
fertilization and grouping on fitness attributes in the freshwater snail Physa acuta‐:
Population and individual inbreeding depression. *Journal of Evolutionary Biology*, *13*(4),
645‐655.

Jiang, P., Zhang, X., Zhu, Y., Zhu, W., Xie, H., & Wang, X. (2007). Metabolism of reactive
oxygen species in cotton cytoplasmic male sterility and its restoration. *Plant Cell Reports*,
26(9), 1627‐1634. https://doi.org/10.1007/s00299-007-0351-6

Kadenbach, B. (Éd.). (2012). *Mitochondrial Oxidative Phosphorylation‐: Nuclear-Encoded*
Genes, Enzyme Regulation, and Pathophysiology (Vol. 748). Springer New York.
https://doi.org/10.1007/978-1-4614-3573-0

Krebs, H. A. (1970). The History of the Tricarboxylic Acid Cycle. *Perspectives in Biology and*
Medicine, *14*(1), 154‐172.

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package‐: Tests
in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*, 1‐26.
https://doi.org/10.18637/jss.v082.i13

Laugier, F., Saclier, N., Béthune, K., Braun, A., Konecny, L., Lefébure, T., Plénet, S.,
Romiguier, J., & David, P. (2024). Both nuclear and cytoplasmic polymorphisms are involved
in genetic conflicts over male fertility in the gynodioecious snail Physa acuta. *Evolution*.

Lee, C. p., Gu, Q., Xiong, Y., Mitchell, R. A., & Ernster, L. (1996). P/O ratios reassessed‐:
Mitochondrial P/O ratios consistently exceed 1.5 with succinate and 2.5 with NAD-linked

substrates. *The FASEB Journal*, *10*(2), 345 350. https://doi.org/10.1096/fasebj.10.2.8641569

Li, J., Pandeya, D., Jo, Y. D., Liu, W. Y., & Kang, B.-C. (2012). Reduced activity of ATP
synthase in mitochondria causes cytoplasmic male sterility in chili pepper. *Planta*, *237*(4),
1097 1109. https://doi.org/10.1007/s00425-012-1824-6

Liu, Z., Shi, X., Li, S., Hu, G., Zhang, L., & Song, X. (2018). Tapetal-Delayed Programmed
Cell Death (PCD) and Oxidative Stress-Induced Male Sterility of Aegilops uniaristata
Cytoplasm in Wheat. *International Journal of Molecular Sciences*, *19*(6), Article 6.
https://doi.org/10.3390/ijms19061708

Logan, D. C. (2006). The mitochondrial compartment. *Journal of Experimental Botany*, *57*(6),
1225 1243. https://doi.org/10.1093/jxb/erj151

Makrecka-Kuka, M., Krumschnabel, G., & Gnaiger, E. (2015). High-Resolution Respirometry
for Simultaneous Measurement of Oxygen and Hydrogen Peroxide Fluxes in Permeabilized
Cells, Tissue Homogenate and Isolated Mitochondria. *Biomolecules*, *5*(3), Article 3.
https://doi.org/10.3390/biom5031319

Marienfeld, J. R., & Newton, K. J. (1994). The maize NCS2 abnormal growth mutant has a
chimeric nad4-nad7 mitochondrial gene and is associated with reduced complex I function.
Genetics, *138*(3), 855 863. https://doi.org/10.1093/genetics/138.3.855

McKenzie, M., & Trounce, I. (2000). Expression of *Rattus norvegicus* mtDNA in *Mus*
musculus Cells Results in Multiple Respiratory Chain Defects*. *Journal of Biological*
Chemistry, *275*(40), 31514 31519. https://doi.org/10.1074/jbc.M004070200

Pesta, D., & Gnaiger, E. (2011). High-Resolution Respirometry : OXPHOS Protocols for
Human Cells and Permeabilized Fibers from Small Biopsies of Human Muscle. In C. M.
Palmeira & A. J. Moreno (Éds.), *Mitochondrial Bioenergetics : Methods and Protocols* (p.
25 58). Humana Press. https://doi.org/10.1007/978-1-61779-382-0_3

Quinlan, C. L., Orr, A. L., Perevoshchikova, I. V., Treberg, J. R., Ackrell, B. A., & Brand, M.
D. (2012). Mitochondrial complex II can generate reactive oxygen species at high rates in
both the forward and reverse reactions. *The Journal of Biological Chemistry*, *287*(32),
27255 27264. https://doi.org/10.1074/jbc.M112.374629

R Core Team. (2023). *R: A Language and Environment for Statistical Computing. R*
Foundation for Statistical Computing, Vienna, Austria. [Jeu de données].

Rand, D. M., Haney, R. A., & Fry, A. J. (2004). Cytonuclear coevolution : The genomics of
cooperation. *Trends in Ecology & Evolution*, *19*(12), 645 653.
https://doi.org/10.1016/j.tree.2004.10.003

Robison, M. M., Ling, X., Smid, M. P. L., Zarei, A., & Wolyn, D. J. (2009). Antisense
Expression of Mitochondrial ATP Synthase Subunits OSCP (ATP5) and γ (ATP3) Alters Leaf
Morphology, Metabolism and Gene Expression in Arabidopsis. *Plant and Cell Physiology*,
50(10), 1840 1850. https://doi.org/10.1093/pcp/pcp125

Sabar, M., De Paepe, R., & de Kouchkovsky, Y. (2000). Complex I Impairment, Respiratory
Compensations, and Photosynthetic Decrease in Nuclear and Mitochondrial Male Sterile

Mutants of Nicotiana sylvestris 1. *Plant Physiology*, *124*(3), 1239–1250.
https://doi.org/10.1104/pp.124.3.1239

Salin, K., Auer, S. K., Anderson, G. J., Selman, C., & Metcalfe, N. B. (2016). Inadequate food
intake at high temperatures is related to depressed mitochondrial respiratory capacity.
Journal of Experimental Biology, *219*(9), 1356–1362. https://doi.org/10.1242/jeb.133025

Salin, K., Luquet, E., Rey, B., Roussel, D., & Voituron, Y. (2012). Alteration of mitochondrial
efficiency affects oxidative balance, development and growth in frog (Rana temporaria)
tadpoles. *Journal of Experimental Biology*, *215*(5), 863–869.
https://doi.org/10.1242/jeb.062745

Salin, K., Villasevil, E. M., Anderson, G. J., Lamarre, S. G., Melanson, C. A., McCarthy, I.,
Selman, C., & Metcalfe, N. B. (2019). Differences in mitochondrial efficiency explain
individual variation in growth performance. *Proceedings of the Royal Society B: Biological*
Sciences, *286*(1909), 20191466. https://doi.org/10.1098/rspb.2019.1466

Saumitou-Laprade, P., Cuguen, J., & Vernet, P. (1994). Cytoplasmic male sterility in plants :
Molecular evidence and the nucleocytoplasmic conflict. *Trends in Ecology & Evolution*, *9*(11),
431–435. https://doi.org/10.1016/0169-5347(94)90126-0

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ : 25 years
of Image Analysis. *Nature methods*, *9*(7), 671–675.

Skarlou, E., Laugier, F., Béthune, K., Chenin, T., Donnay, J.-M., Froissard, C., & David, P.
(2024). *Complex genetic determination of male-fertility restoration in the gynodioecious snail*
Physa acuta (p. 2024.08.08.607164). bioRxiv. https://doi.org/10.1101/2024.08.08.607164

Sluse, F. E., & Jarmuszkiewicz, W. (1998). Alternative oxidase in the branched mitochondrial
respiratory network : An overview on structure, function, regulation, and role. *Brazilian*
Journal of Medical and Biological Research = Revista Brasileira De Pesquisas Medicas E
Biologicas, *31*(6), 733–747. https://doi.org/10.1590/s0100-879x1998000600003

Srere, P. A. (1971). An eclectic view of metabolic regulation : Control of citrate synthase
activity. *Advances in Enzyme Regulation*, *9*, 221–233. https://doi.org/10.1016/S0065-
2571(71)80046-8

Sunnucks, P., Morales, H. E., Lamb, A. M., Pavlova, A., & Greening, C. (2017). Integrative
Approaches for Studying Mitochondrial and Nuclear Genome Co-evolution in Oxidative
Phosphorylation. *Frontiers in Genetics*, *8*.
https://www.frontiersin.org/articles/10.3389/fgene.2017.00025

Tang, D., Wei, F., Khan, A., Munsif, F., & Zhou, R. (2021). Degradation of mitochondrial
structure and deficiency of complex I were associated with the transgenic CMS of rice.
Biological Research, *54*(1), 6. https://doi.org/10.1186/s40659-020-00326-y

Tang, H., Xie, Y., Liu, Y.-G., & Chen, L. (2017). Advances in understanding the molecular
mechanisms of cytoplasmic male sterility and restoration in rice. *Plant Reproduction*, *30*(4),
179–184. https://doi.org/10.1007/s00497-017-0308-z

Touzet, P., & H. Meyer, E. (2014). Cytoplasmic male sterility and mitochondrial metabolism
in plants. *Mitochondrion*, *19*, 166–171. https://doi.org/10.1016/j.mito.2014.04.009

Tsitrone, A., Jarne, P., & David, P. (2003). Delayed Selfing and Resource Reallocations in
Relation to Mate Availability in the Freshwater Snail Physa acuta. *The American naturalist*,
162, 474 488. https://doi.org/10.1086/376889

Vedel, F., Lalanne, É., Sabar, M., Chétrit, P., & De Paepe, R. (1999). The mitochondrial
respiratory chain and ATP synthase complexes : Composition, structure and mutational
studies. *Plant Physiology and Biochemistry*, *37*(9), 629 643. https://doi.org/10.1016/S0981-
9428(00)80093-5

Wen, L., Ruesch, K. L., Ortega, V. M., Kamps, T. L., Gabay-Laughnan, S., & Chase, C. D.
(2003). A Nuclear restorer-of-fertility Mutation Disrupts Accumulation of Mitochondrial ATP
Synthase Subunit α in Developing Pollen of S Male-Sterile Maize. *Genetics*, *165*(2),
771 779. https://doi.org/10.1093/genetics/165.2.771

Werren, J. H. (2011). Selfish genetic elements, genetic conflict, and evolutionary innovation.
Proceedings of the National Academy of Sciences, *108*(supplement_2), 10863 10870.
https://doi.org/10.1073/pnas.1102343108

Werren, J. H., & Beukeboom, L. W. (1998). Sex Determination, Sex Ratios, and Genetic
Conflict. *Annual Review of Ecology and Systematics*, *29*, 233 261.

Williams, G. C. (1996). *Adaptation and natural selection : A critique of some current*
evolutionary thought. Princeton Univ. Press.

Wolff, J. N., Ladoukakis, E. D., Enríquez, J. A., & Dowling, D. K. (2014). Mitonuclear
interactions : Evolutionary consequences over multiple biological scales. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *369*(1646), 20130443.
https://doi.org/10.1098/rstb.2013.0443

Yu, J., Chen, B., Zheng, B., Qiao, C., Chen, X., Yan, Y., Luan, X., Xie, B., Liu, J., Shen, C.,
He, Z., Hu, X., Liu, M., Li, H., Shao, Q., & Fang, J. (2019). ATP synthase is required for male
fertility and germ cell maturation in Drosophila testes. *Molecular Medicine Reports*, *19*(3),
1561 1570. https://doi.org/10.3892/mmr.2019.9834