Cytoplasmic male sterility and mitochondrial metabolism: evidence for low complex I contribution in male-sterile freshwater snail *Physa acuta*

Sophie Bererd[1*], Damien Roussel[1], Sandrine Plénet[1], Loïc Teulier[1], Antoine Stier[2,3$], Emilien Luquet[1,4$]

[1]Laboratoire d'Ecologie des Hydrosystèmes Naturels et Anthropisés, CNRS UMR 5023 Université Claude Bernard Lyon 1, Université de Lyon, Lyon, France.
[2]Université de Strasbourg, CNRS, Institut Pluridisciplinaire Hubert Curien, UMR7178, 67000 Strasbourg, France
[3]Department of Biology, University of Turku, Turku, Finland
[4]Institut Universitaire de France (IUF), Paris, France
[$]shared senior authorship
[*]corresponding authors: bererdsophie@gmail.com

Abstract

Cytoplasmic male sterility (CMS) is a well-known example of mitonuclear genetic conflict over sex determination in hermaphrodite plants, where mitochondrial genes maternally inherited sterilize the male function while biparental inherited nuclear genes restore it. CMS has been recently discovered in wild animals, in the freshwater snail *Physa acuta*. In this species, CMS is associated with two extremely divergent mitogenomes D and K compared to the classical mitogenome N. D individuals are male-steriles, while male fertility is restored in K individuals. We hypothesized that the extreme divergence of mitogenomes associated with CMS can impact mitochondrial aerobic metabolism, a necessary process in eukaryotic organisms by which energy is transduced from food to ATP. Our results suggest that CMS might be associated with an alteration of co-encoded complex I in D male-sterile individuals, although partly compensated by the nuclear-encoded complex II. K restored hermaphrodites have an unaffected complex I respiration, but exhibit higher mitochondrial proton leak and higher relative anaerobic contribution to cellular metabolism, suggesting an energy cost of bearing CMS and restorer genes, which could underlie the reduced growth of K mitotype. How complex I alteration might induce male-sterility remains to be determined, but could be linked to oxidative stress or a defect in ATP-synthesis rate.

Keywords: cellular metabolism, gynodioecy, male sterility, mitonuclear interaction, OXPHOS system, *Physa acuta*

INTRODUCTION

The gene-centred view of evolution has revolutionised modern evolutionary biology, promoting that selection occurs at the level of the gene mediated by the phenotype (Williams, 1966). Conflict can arise when genes have opposing evolutionary interests, *i.e.* different strategies to maximize their own transmission, leading to antagonistic selective pressures over the phenotype (Werren, 2011). A fabulous example of genomic conflict impacting phenotype occurs in hermaphrodites, opposing nuclear and mitochondrial genes within individuals (Cosmides & Tooby, 1981; Werren & Beukeboom, 1998). Unlike nuclear genes transmitted by both parents, mitochondrial genes are usually inherited by one parent, most often the female, meaning that the fitness of mitochondrial genes only depends on the female function (Havird et al., 2019). This disparity in inheritance pattern between these genomes results in a conflict over sex determination (Chase, 2007; Cosmides & Tooby, 1981). Indeed, some selfish mitochondrial genes can emerge and sterilize the male function of hermaphrodites, who then become only functionally female. This phenomenon, known as cytoplasmic male sterility (CMS), increases the fitness of mitochondrial genes by enhancing their transmission though the female line (Budar et al., 2003; Chase, 2007; Saumitou-Laprade et al., 1994). In contrast, the male function sterilization leads to a loss of fitness for nuclear genes (Havird et al., 2019). Therefore, some nuclear genes neutralizing CMS (named restorer genes) can be selected to counteract sterilising effect of CMS genes by reverting individuals to hermaphroditism (known as restored hermaphrodite; Budar et al., 2003; Chase, 2007; Saumitou-Laprade et al., 1994). In natural populations, such a mitonuclear conflict results in a sexual polymorphism where male-sterile (female) individuals co-occur with hermaphrodite ones (normal and restored), a mating system called gynodioecy (Saumitou-Laprade et al., 1994). Gynodioecy, and the underlying dynamics of CMS and restorers genes in populations, is expected to evolve through a balance between selective advantage to be male-sterile (called female advantage, *i.e.* male-sterile individuals benefit from a better female fitness than hermaphrodites) and costs to bear CMS and restorers genes (Dufaÿ et al., 2007).

Gynodioecy was thought until recently to be a specificity of angiosperms (<1%
of species, almost 20% families; Caruso et al., 2016; Dufay et al., 2014). Numerous
studies highlight the association between divergent mitochondrial CMS lineages and
the failure to produce pollen (review in Touzet & H. Meyer, 2014), suggesting that the
functioning of mitochondria is at the core of the CMS mechanisms (*i.e.* male sterility,
balance between female advantage and costs). Mitochondria are essential
organelles as they are involved in various fundamental cellular functions, in particular
cellular aerobic metabolism by transducing energy acquired from food to ATP
(adenosine triphosphate) through oxidative phosphorylation (OXPHOS; Kadenbach,
2012; Logan, 2006; Vedel et al., 1999). The OXPHOS depends on the coordinated
action of mitochondrial and nuclear genes because both encode for the proteins
constituting the respiratory chain complexes and the ATP synthase (except the
complex II that is fully encoded by the nuclear genome; Kadenbach, 2012). The
divergence of CMS mitochondrial genomes may lead to altered functions through
various and poorly known mechanisms (*e.g.* breakdown of coadaptation between
mitochondrial and nuclear genomes, production of neoproteins by chimeric genes
due to new open reading frames; Delph et al., 2007). In several CMS plant species,
respiratory chain complex dysfunctions have been characterized and involve mainly
complexes composed by protein subunits encoded by both mitochondrial and nuclear
genes (Touzet & H. Meyer, 2014). For example, impairment of the complex I was
described in *Nicotiana sylvestris* male-sterile mutants (Gutierres et al., 1997; Sabar et
al., 2000) and in CMS transgenic rice (D. Tang et al., 2021). Cytochrome-c oxidase
(COX - Complex IV) activity is also reduced by 50% in leaves of male-sterile wild
beets (Ducos et al., 2001). Similarly, the activity of ATP synthase is reduced in CMS
chilli pepper (Li et al., 2012). Understanding the physiological aspects of CMS is then
crucial, as it helps to understand the mechanisms leading to male sterility and clarify
the proximal causes underlying the evolutionary dynamics of gynodioecious systems.
96
Our study uses the freshwater snail *Physa acuta*, the recently-discovered and
only known wild CMS animal model associated with naturally occurring mitochondrial
types (David et al., 2022), allowing for the exploration of the mechanisms underlying
CMS from the molecular to the fitness levels. Two mitochondrial genomes – named D
and K mitotypes – have been associated with male sterility. These mitotypes are both

extremely divergent from one another (mtDNA nucleotidic divergence ranged from
41% to 61%; Laugier et al., 2024) and from the classical mitochondrial genome,
named N mitotype (35% to 57% between K and N and 28.6% to 53.2% between D
and N; David et al., 2022; Laugier et al., 2024). D individuals are male-steriles (*i.e.*
they produced a very low amount to no spermatozoa and showed a strong
suppression of male reproductive behaviour; David et al., 2022) whereas K
individuals are restored hermaphrodites (presence of nuclear restorer genes; Laugier
et al., 2024). A recent study of Bererd et al. (2025) on the benefits and costs of CMS
in *P. acuta* provided evidence that CMS is beneficial to female fitness in the absence
of restorers (female advantage – D male-steriles have a better female fitness by
producing higher number of eggs), while it is costly in their presence (K restored
hermaphrodites have a lower growth rate). Interestingly, fitness benefits and costs
were mediated by differences in body mass, suggesting that different growth and
underlying metabolisms are involved. However, the whole-body metabolism was
surprisingly similar among N, D and K mitotypes (Bererd et al., 2025). Altogether,
these results suggested that the divergent mitochondrial genomes could cause
impairments on cellular metabolism, altering the life history, fitness and fertility of
organisms, but also that some compensation mechanisms may exist at the cellular
level (*e.g.* higher number of mitochondria as proposed in CMS *Nicotiana sylvestris* by
Sabar et al., 2000). The aim of the present study was thus to investigate how CMS
and restorers genes were related to cellular metabolism potentially implicated in male
sterility and balance between female advantage and costs in the *P. acuta*
gynodioecious system (N normal hermaphrodite, D male-sterile, K restored
hermaphrodite). We compared the male status (sterile or fertile), the growth rate, the
mitochondrial aerobic metabolism and the anaerobic cellular metabolism between (i)
individuals carrying CMS (D) and their non-CMS counterparts (N), and between (ii)
individuals carrying CMS and K-specific restorers genes (K) and their non-CMS
counterparts (N).
130

MATERIAL AND METHODS

-   <u>Experimental design</u>

P. acuta (Physidae, Hygrophila, Gastropoda) is a cosmopolitan widespread snail
living in freshwater (Dillon et al., 2005). This species is a simultaneous
hermaphrodite: sperm and oocytes are produced within the same organ called the
ovotestis but sperm are stock in seminal vesicle for maturation, with a high
preference for outcrossing (*i.e.* eggs are self-fertilised only when no mate is available
to provide allosperm (Henry et al., 2005; Jarne et al., 2000; Tsitrone et al., 2003).

Three laboratory *P. acuta* lines were used: the mitotype N normal hermaphrodite, the
mitotype D male-sterile and the mitotype K restored hermaphrodite. They were
obtained from wild populations near Lyon (southeastern France) where the three
mitotypes co-occurred and interbred, sharing a similar nuclear background (David et
al., 2022; Laugier et al., 2024). In these populations, the prevalence of K-restorer
genes in the nuclear background is high, leading to the restoration of male fertility in
K snails (Laugier et al., 2024; Skarlou et al., 2025). Wild snails have been sampled
and then reproduced in the laboratory (David et al., 2022; Laugier et al., 2024). The
mitotype (N, D or K) of their offspring have been determined using a PCR test (see
further details in Laugier et al., 2024). The three lines have been generated from
these offspring and raised at 25°C (maturity in the laboratory is reached after 35-40
days at 25°C). N and K have been maintained as separate groups for 20
generations. D snails, as they were male-sterile, were fertilised by N snails through
pair-crosses at each generation for 33 generations. The three lines share the same
nuclear background, which carry restorer genes specific to K mitotype (effective in K
lines but failing to restore CMS in D snails).

At 11 days post-hatching, 127 snails per mitotype were placed in a temperature-
controlled room at 22.1 ± 0.1 °C and photoperiod 12h/12h. All the snails were reared
individually in small boxes (90 ml) with dechlorinated tap water changed once a week
and were fed with boiled and mixed lettuce twice a week.

-    <u>Male status and growth</u>

At 34 days after the start of the experiment (45 days post-hatching), the male status
(sterile or fertile) of each focal individual was assessed by pairing it to a virgin albino
and by recording the pigmentation of the albino progeniture (unhatched offspring,
about one week after the pairing using a binocular microscope): pigmented offspring

indicated that the focal individual was male-fertile and on the contrary, no or albino
offspring indicated male sterility (see David et al., 2022 for more details). The albino
individuals are N mitotype, hermaphroditic snails (*i.e.* male-fertile and non-CMS),
originating from the Montpellier Area (~300km south of Lyon) and have been
maintained as a large laboratory population for over 80 generations.

Before the pairing, whole-body mass (shell + body) was measured with a precision
scale to the nearest 0.001mg. Maximal shell length was assessed using photography
analysed with the software ImageJ (Schneider et al., 2012) at the nearest 0.001mm.

Note that only N and K male-fertile individuals (normal hermaphrodite and restored
hermaphrodite, respectively) and D male-sterile individuals were selected for the
comparisons of growth (whole-body mass and maximal shell length), mitochondrial
aerobic metabolism and enzymatic activities.

- <u>Mitochondrial aerobic metabolism</u>

In order to compare the mitochondrial aerobic metabolism of the N, D and K
mitotypes, 12-13 individuals per mitotype (total n = 37 individuals) were selected to
assess mitochondrial functionality using high resolution respirometry (O2k, Oroboros;
DatLab software; Makrecka-Kuka et al., 2015). Snails were anaesthetised on ice and
dissected to sample the head and foot tissues (all the digestive and reproductive
tissues were delicately removed). Tissue mass was measured with a precision scale
at the nearest 0.001mg. Mechanical permeabilization of the tissue was performed by
consecutive scissors cuts in 2 ml of a Mir05 solution (Salin et al., 2016) and verified
as efficient in preliminary assays using additional chemical permeabilization with
digitonin. Mitochondrial integrity was assessed during each run by the addition of 10
µM of cytochrome c, which had very little impact on respiration rate (+1.4% ± 1.3%),
thereby proving the good integrity of our mitochondrial preparation. Four O2-k
devices were used at an assay temperature of 21.5°C and a chamber volume of
2mL, with background calibration performed once before measurement sessions
started, and 100% calibration performed daily. Mitotypes were randomly assigned to
the four O2k devices, and were evenly distributed across five sessions of
measurements. Mitochondrial aerobic metabolism was assessed using a relatively

typical substrate-uncoupler-inhibitor titration (SUIT protocol; Pesta & Gnaiger, 2011;
Fig.1).

During a pilot study aimed at setting up experimental conditions for high-
resolution respirometry on *P. acuta* homogenate, we noticed surprisingly high levels
of apparent non-mitochondrial respiration (*i.e.* after inhibition of complex III with 2.5, 5
or 10 µM of antimycin A). This residual respiration being sensitive to SHAM
(Salicylhydroxamic acid or 2-Hydroxybenzohydroxamic acid), a known inhibitor of the
alternative oxidase (AOX) pathway (Sluse & Jarmuszkiewicz, 1998), this suggests
that AOX pathway is present and active in *P. acuta*. We thus systematically inhibited
the AOX pathway at the beginning of the high-resolution respirometry assays with
1mM of SHAM, after stabilisation of endogenous cellular respiration. *AOX* respiration
was thus calculated as the difference between endogenous and SHAM-inhibited
respirations (Fig. 1B). Mitochondrial respiration was then fuelled by complex I
substrates (5mM pyruvate and 2mM malate) and ADP was added at a concentration
of 5mM to stimulate phosphorylating respiration (*CI OXPHOS*). Complex II substrate
(succinate 10mM) was then added to achieve phosphorylating respiration driven by
electron entry through both complexes I and II (*CI+II OXPHOS*). Cytochrome c 10 µM
was added to verify mitochondrial integrity (see above). ATP synthesis was
subsequently inhibited with 2.5 µM of oligomycin, leading to a non-phosphorylating
respiration (*i.e.* driven by proton leak) referred to as *CI+II LEAK*. The uncoupler
FCCP (carbonyl cyanide p-trifluoromethoxyphenyl hydrazone) was then titrated in 0.5
µM steps until maximum uncoupled respiration was achieved, which measures the
maximum capacity of the electron transport system (*ETSmax*). Inhibition of complex
III with antimycin A 2.5 µM led to mitochondrial respiration inhibition, and the residual
(*i.e.* non-mitochondrial) O2 consumption left was subtracted from other respiration
rates. Finally, electrons were fuelled directly to complex IV with 2mM ascorbate and
0.5mM TMPD (N,N,N',N'-tetramethyl- p-phenylene-diamine), before inhibition of
complex IV activity with sodium azide 100mM, the difference between these two
respiration rates corresponding to complex IV maximum capacity to use O2 (*i.e.* COX
respiration; Fig. 1B). All respiration rates were normalized per milligram of fresh
tissue and expressed as $pmolO_2.s^{-1}.mg^{-1}$ of tissue.

Mitochondrial respiration rates are sensitive to both mitochondrial quantity and
intrinsic properties (*e.g.* functionality of electron transport). We also calculated three

flux control ratios (*i.e.* ratios between different respiration rates) that are independent

from mitochondrial quantity and thus purely reflect mitochondrial properties: 1. The

efficiency of oxidative phosphorylation, *i.e. OXPHOS control efficiency*, calculated as:

(*CI+II OXPHOS - CI+II LEAK*) / *CI+II OXPHOS (Gnaiger et al. 2020)*. This ratio gives

the minimum proportion of O2 consumed contributing to oxidative phosphorylation; a

ratio of 0.75 meaning that at least 75% of O2 consumption under phosphorylating

conditions is dedicated to oxidative phosphorylation. *2*. The mitochondrial reserve to

increase respiration from OXPHOS to maximal ETS respiration: *ETS reserve*

calculated as (*CI+II ETS - CI+II OXPHOS*) / *CI+II ETS;* a ratio of 0.75 meaning that

mitochondrial O2 consumption can increase by 75% in case of energy demand. *3.*

The relative contribution of complex I to maximal respiratory activity, when respiration

is fuelled by both complex I and II substrates: *CI OXPHOS contribution*, calculated as

1 - ((*CI+II OXPHOS - CI OXPHOS*) / *CI OXPHOS);* a ratio of 0.75 meaning that

complex I supports 75% of the mitochondrial O2 consumption when CI and CII

substrates are provided together.

Figure 1. **A -** Schematic of the OXPHOS system. Mitochondrial and nuclear genes
encode for the proteins constituting complexes I, III, IV and V (solid line border).
Complex II and AOX are fully encoded by nuclear genes (black dashed border). The
process relies on the respiratory chain formed by four protein complexes and the
ATP synthase where a succession of redox reactions catalysed by four complexes (I,
II, III and IV) creates a proton gradient, which is then used by complex V – ATP
synthase – for the synthesis of ATP from ADP and inorganic phosphate. Proton Leak
refers to the process where protons (H^+) cross the inner membrane independently of
ATP synthase. **B -** Graphical summary of the high-resolution respirometry substrate-
uncoupler-inhibitor-titration (SUIT) protocol. *ENDO* respiration rate reflects the
endogenous respiration of cells. *AOX* (alternative oxidase) respiration corresponds to
the respiratory rate decrease in response to alternative oxidase inhibition by SHAM
(Salicylhydroxamic acid or 2-Hydroxybenzohydroxamic acid). *CI OXPHOS* respiration
rate fuelled by Complex I was measured after addition of its substrates Pyruvate and
Malate with ADP (PM + ADP) and corresponds to the respiratory rate of the system
with electron entry only through Complex I. *CI+II OXPHOS* respiration rate fuelled by
Complexes I and II was measured after addition of Complex II's substrates Succinate
(S) and reflects the respiratory rate of the whole substrate-saturated system. *CI+II*
LEAK respiration rate reflects the proton leak assessed by inhibiting Complex IV with
Oligomicyn (O). *ETSmax* corresponds to the maximum capacity of the electron
transport system and was measured after the titration of uncoupler FCCP (carbonyl
cyanide p-trifluoromethoxyphenyl hydrazone). *COX* reflects the Complex IV

maximum capacity to use O2. *NON-MITO* reflects the non-mitochondrial respiration
rate, measured by the addition of antimycin A (AA).

-    <u>Relative contribution of anaerobic metabolism to aerobic metabolism</u>

The relative contribution of anaerobic metabolism to aerobic metabolism in the three
mitotypes was measured by comparing enzymatic activities of the lactate
dehydrogenase (LDH) and the citrate synthase (CS), respectively. LDH is a key
enzyme of anaerobic metabolism, by converting pyruvate into lactate, it regenerates
the NAD+ used by the glycolysis pathway to oxidize glucose-6-phosphate and
synthesize ATP in the absence of oxygen (Emmett & Hochachka, 1981). CS is a key
enzyme of mitochondrial aerobic metabolism as it is involved in the tricarboxylic acid
cycle, which fuel the mitochondrial electron transport chain with reduced coenzymes
(NADH and FADH2; Emmett & Hochachka, 1981; Krebs, 1970; Srere, 1971).

The head and the foot of 20 snails per mitotype (total n = 60 individuals) were
sampled, weighted and frozen at -80°C. Tissues samples were thawed, then
homogenised using a glass piston-glass Potter in 500µl buffer containing 100 mM
sucrose, 50 mM KCl, 5 mM EDTA, 50 mM Tris-base, 1% fatty acid free BSA, pH 7.4.
The resulting homogenates were kept at -80°C until measurements. After thawing,
homogenates were vortexed and centrifuged at 1000 g for 10 min (4°C). The
supernatants were retrieved and used for the LDH and CS enzyme assays. Enzyme
activities were measured spectrophotometrically at 22°C. All results were expressed
as $nmol.min^{-1}.mg^{-1}$ of tissue.

- **LDH activity** was measured in a reaction medium composed of 40mN imidazole
(pH 7), 0.04% BSA and 200 µM NADH. Two replicates per homogenates were
performed with 10µl and 20µl of supernatants and 250µl of the LDH reaction medium
were laid in a microtiter plate. After 3 min incubation, the reaction was initiated by
adding 4 mM pyruvate and the change in optical density at 340 nm was recorded for
3 min. The enzyme activity was quantified using an extinction coefficient of 6.22 $mM^{-1}.cm^{-1}$.
298

- **CS activity** was measured in a reaction medium composed of 100 mM Tris buffer
(pH 8), 100 µM 5,5'-dithiobis(2-nitrobenzoic acid), 300 µM acetyl-CoA and 0.1%

Triton-X-100. Homogenates were diluted ½ in a 100 mM phosphate buffer. Two

replicates of 5µl and 7µl of diluted homogenates and 250µl of the CS reaction

medium were deposited in a microtiter plate. After 3 min incubation, the reaction was

initiated by adding 500 µM oxaloacetate and the change in optical density at 412 nm

was recorded for 3 min. The enzyme activity was quantified using an extinction

coefficient of 13.6 mM^{-1}.cm^{-1}.

- The **LDH/CS ratio** was calculated to estimate the contribution of anaerobic

metabolism relative to aerobic metabolism and used as an index of relative glycolytic

capacity.

The technical repeatability (i.e. intra-class coefficient of correlation) of our

enzymatic measurements based on duplicate was high, namely $R = 0.94$ for CS and

$R = 0.82$ for LDH.

- _Statistical analyses_

Male status - A generalized linear model (GLM) following a binomial distribution was

used to test the proportion of male sterility in the three mitotype by an analysis of

variance (ANOVA).

Growth - The effect of the mitotype on whole body mass and on maximal shell length

were tested by a linear model using an ANOVA.

Mitochondrial aerobic metabolism - A global linear mixed model was conducted to

analyse **mitochondrial respiration rates** with "mitotype" and "state" as fixed factors

(state corresponds to the specific mitochondrial respiration rates: AOX, CI OXPHOS,

CI+II OXPHOS, CI+II LEAK, ETS, COX), the O2k chamber identity as a random

factor to control for potential technical variations between the eight chambers used,

"session (day of measurement)" as a random factor to control for technical variations

between measurement sessions and a factor to account for identity of individuals.

Linear mixed models (LMM) were conducted to analyse **mitochondrial flux control**

ratio with "mitotype" as a fixed factor, the O2k chamber identity as a random factor to

control for potential technical variations between the eight chambers used, and

"session (day of measurement)" as a random factor to control for technical variations

between measurement sessions.

The significance of fixed effects was tested using F-tests and the Kenward-Roger

approximation for the degrees of freedom.

Enzymatic activities – Means of the two replicates for both LDH and CS activities

were used in statistical analyses. LDH/CS ratio was calculated as the average of

LDH means and CS means. Linear models and ANOVAs were conducted to analyse

the effect of the mitotype on LDH activity, CS activity and LDH/CS ratio.

All statistical analyses were carried out in Rstudio 4.3.2 (R Core Team, 2023). Linear

mixed models were performed using *lme4* package (Bates et al., 2015; p-values were

obtained using *LmerTest* package, Kuznetsova et al., 2017). Post-hoc tests were

performed using the *emmeans* package (Lenth, 2024). Tukey correction was used

when the mitotype factor was significant ($p < 0.05$), and FDR adjustment was used

specifically for LMM testing mitochondrial respiration rates considering the high

number of pairwise comparisons.

RESULTS

- <u>Male status</u>

Male sterility depended on mitotype ($\chi^2_{2, 337} = 277.43$, $p < 0.001$; Fig. 2). D individuals

were almost all male-sterile as D had a much lower male-fertility proportion than N

(normal hermaphrodite) and K (restored hermaphrodite; in logit scale, N – D: Est.±SE

= 6.25 ± 0.79, z = 7.94, p < 0.0001; D - K: Est.±SE= -6.29 ± 0.79, z = 8.00, p <

0.001). N and K were mostly male-fertile and not significantly different from each

other (in logit scale, N − K = -0.045 ± 0.467, z = -0.096, p = 0.995).

<u>Figure 2.</u> Percentage of male-fertile and male-sterile individuals in N, D and K

P.acuta mitotypes. The total sample sizes were 124 N, 123 D and 127 K.

-   <u>Growth</u>

Whole-body mass was significantly different between mitotypes ($F_{2, 341}$ = 128.79, p

< 0.001; Fig. 3A). D male-sterile snails were bigger than N and K male-fertiles (N - D:

Est.±SE = -0.013 ± 0.003, t_{341} = -4.98, p < 0.001; D - K: Est.±SE = 0.042 ± 0.003, t_{341}

= 15.78, p < 0.001). K had a significantly lower whole-body mass than N and D (N -

K: Est.±SE = 0.029 ± 0.003, t_{341} = 10.46, p <0.001; Fig. 3A).

Maximal shell length was also significantly different between mitotypes ($F_{2, 338}$ =

88.50, p < 0.001; Fig. 3B). D male-sterile had a longer shell length than N and K

male-fertiles (N - D: Est.±SE = -0.662 ± 0.105, t_{338} = -6.30, p < 0.001; D - K: Est.±SE

= 1.383 ± 0.104, t_{338} = 13.30, p <0.001). K had a significantly smaller shell length

than N and D (N - K: Est.±SE = 0.720 ± 0.107, t_{338} = 6.75, p < 0.001; Fig. 3B).

<u>Figure 3.</u> **A** – Whole-body mass (in g) for *P. acuta* normal hermaphrodites N (red

circle), restored hermaphrodites K (blue square) and male-steriles D (green triangle).

B – Maximal shell length (in mm). Black dots and black errors bars correspond to

means ± standard error.

-    **Mitochondrial aerobic metabolism**

Mitochondrial respiration rates

Mitochondrial respiration rates were significantly influenced by the
mitotype*state interaction (interaction: $F_{10,\ 168.607}$ = 2.33, p = 0.013; mitotype effect:
$F_{2,9.779}$ = 3.24, p = 0.083; respiratory state effect: $F_{5,\ 168.611}$ = 255.56, p < 0.001).
When looking for mitotype effect within each respiratory state, mitochondrial
respiration rates related to oxidative phosphorylation through complex I and II
combined (*CI+II OXPHOS*), maximal uncoupled respirations (*CI+II ETS*), and
maximal oxidative capacity achieved under direct electron fueling to complex IV
(*COX*) did not statistically differ between mitotypes (all p > 0.69 ; Fig. 4A, Tab. S1,
Fig. S1). However, mitochondrial respiration through the alternative oxidase (AOX)
pathway was lower in D mitotype compared to N and K (all p < 0.020, Tab. S1), a
pattern also observed for mitochondrial respiration through complex I, although not
fully significant (D - N: p = 0.079, D - K: p = 0.046). Finally, mitochondrial respiration
rate related to proton leak through complex I and II (*CI+II LEAK*) was significantly
higher in K mitotype than in N and D (Fig 4A; all p < 0.014, Tab. S1).

Figure 4. **A −** Mitochondrial respiration rates and **B −** Flux control ratios between

mitotypes: N normal hermaphrodites (red), D male-steriles (green) and K restored

hermaphrodites (blue) and. Figures are based on marginal means ± SE from

statistical models and symbols indicate differences between mitotypes according to

post-hoc tests (ns: p > 0.10, $: p < 0.10, *: p < 0.05, **: p < 0.01).

Mitochondrial flux control ratios

406   The efficiency of the OXPHOS system (*OXPHOS control efficiency*) was

significantly affected by mitotype ($F_{2,7.0744}$ = 9.62, p = 0.009), with K mitotype having

lower control efficiency than both N ($t_{6.01}$ = 3.66, p = 0.025) and D mitotypes ($t_{7.19}$ = -

3.48, p = 0.024; Fig. 4B). The ability to increase respiration from phosphorylating to

uncoupled conditions did not significantly differ between mitotypes (*ETS reserve*:

$F_{2,14.004}$ = 0.78, p = 0.48). Finally, the contribution of complex I to OXPHOS

respiration was markedly affected by mitotype ($F_{2,\ 29.977}$ = 22.45, p < 0.001), with D

individuals having lower contribution of complex I than N and K individuals (D - N:

$t_{9.83}$= -5.09, p = 0.013; D - K: $t_{5.91}$ = -5.58, p = 0.004; Fig. 4B, Fig. S2).

-    <u>Relative contribution of anaerobic metabolism to aerobic metabolism</u>

LDH activity was significantly influenced by the mitotype ($F_{2,57}$ = 6.18, p = 0.004). D

and K individuals had a significantly lower LDH activity than N (D - N: Est.±SE = 0.55

± 0.17, t_{57} = 3.33, p = 0.004; K - N: Est.±SE = 0.44 ± 0.17, t_{57} = 2.65, p = 0.028). LDH

activity in mitotypes K and D was not significantly different from each other (D – K:

Est.±SE = -0.11 ± 0.17, t_{57} = -0.68, p = 0.78; Fig. 5, Fig. S3)

CS activity was significantly influenced by the mitotype ($F_{2,57}$ = 25.34, p < 0.001). K

individuals had a significantly lower CS activity than N and D (N - K: Est.±SE = 0.77 ±

0.11, t_{57} = 6.86, p < 0.001; D - K: Est.±SE = 0.57 ± 0.11, t_{57} = 5.09, p < 0.001). CS

activity in mitotypes N and D was not significantly different from each other (N – D:

Est.±SE = 0.20 ± 0.11, t_{57} = 1.78, p = 0.19 ; Fig. 5, Fig. S3).

The relative anaerobic contribution to metabolism (ratio of LDH activity to CS

activity) was significantly influenced by the mitotype ($F_{2,\ 57}$ = 23.98, p < 0.001), being

higher in K mitotype compared to N and D (N - D = Est.±SE = 0.47 ± 1.17, t_{57} = 0.41,

p = 0.91; N - K: Est.±SE = -6.76 ±1.17, t_{57} = -5.78, p < 0.001; D - K: Est.±SE = -7.24

± 1.17, t_{57} = -6.19, p < 0.001 ; Fig. 5, Fig. S3).

<u>Figure 5.</u> Enzymatic activities of lactate dehydrogenase (anaerobic metabolism) and

citrate synthase (aerobic metabolism) of *P. acuta* N normal hermaphrodites (red), D

male-steriles (blue) and K restored hermaphrodites (green). Means ± SE and

different letters indicate significant differences according to ANOVAs and associated
post-hoc tests.

DISCUSSION

Cytoplasmic male sterility (CMS) originates from a mitonuclear conflict where
mitochondrial genes induce male sterility and nuclear genes restore male fertility in
hermaphrodites. The dynamics of CMS and restorers genes is known to evolve
through a balance between selective advantage to be male-sterile (*i.e.* female
advantage) and costs to bear CMS and restorers genes (Dufaÿ et al., 2007). Male
sterility and balance between female advantage and costs – key parameters of
gynodioecy evolutionary dynamics – may be conditioned by changes in cellular
metabolism. This study uniquely combined measurements of phenotype (growth trait
as proxy of fitness advantage and costs) with an in-depth investigation of cellular
metabolism mechanisms in the *P. acuta* gynodioecious system (N normal
hermaphrodite, D male-sterile, K restored hermaphrodite; David et al., 2022; Laugier
et al., 2024). Our results suggest that CMS in D male-sterile individuals was
associated with a reduction of co-encoded complex I contribution to OXPHOS that
might nonetheless be partly compensated by nuclear-encoded complex II. Moreover,
the efficiency of the OXPHOS system was similar to N normal hermaphrodite which
may explain the female advantage (higher growth) in the absence of energy costs
linked to male function. Such changes were not observed in K restored
hermaphrodites, suggesting that male fertility restorer genes re-established the
OXPHOS metabolism through complex I to a similar level as N normal
hermaphrodites. Moreover, the K restored hermaphrodites showed a reduced growth,
reflecting the costs of bearing CMS and restorer genes, which may be at least partly
explained by the OXPHOS control efficiency and the potential higher relative
anaerobic contribution to cellular metabolism that we observed. In the following, we
discuss in detail these cellular mechanisms of CMS and their implications for its
evolution.

Effects of CMS on reproductive functions and cellular metabolism in male-steriles

Male fertility and female advantage

The conflict opposing the D mitogenome and the nuclear genome results in male sterility in almost all D snails as observed in previous studies (Bererd et al., 2025; David et al., 2022). Less than 2% of D individuals were considered male-fertiles (*i.e.* they fertilised their albino partner), confirming an incomplete penetrance of CMS genes (no restorers are yet found for D mitotype; Bererd et al., 2025; Laugier et al., 2024; Skarlou et al., 2025). The D male-sterile snails had a higher growth, both in terms of mass and length, than N normal hermaphrodite. Bererd et al., (2025) showed that the growth advantage of D male-steriles led to a higher egg production, supporting the expected female advantage in male-sterile individuals required to the maintenance of CMS in natural populations (Dufaÿ et al., 2007). Such a female advantage has been described in plants (Dufay & Billard, 2012) and is usually explained as a consequence of an energy reallocation from the defective male function toward the female fitness (Budar et al., 2003).

Complex I impairment and Complex II compensation

Mitonuclear incompatibilities, particularly when mitochondrial and nuclear genomes have not coevolved, may disturb the integrity of OXPHOS with downstream consequences on many cellular functions (Rand et al., 2004; Sunnucks et al., 2017; Wolff et al., 2014). For example, genetically perturbed mitonuclear co-evolved system by introducing a foreign mitogenome from non-human apes (common chimpanzee, pigmy chimpanzee and gorilla) into a human nuclear background resulted in a significant decreased complex I activity and supported mitochondrial respiration, whereas others mitochondrial co-endoded complexes were less affected (Barrientos et al., 1998). Similarly, a foreign mitogenome from rat into a mouse nuclear background resulted in a large decrease of complex I activity, with a lesser impact on co-encoded complex IV activity (McKenzie & Trounce, 2000). Our results are strikingly consistent as the complex I activity *per se* was lower in D mitotype than in N normal hermaphrodite (although not significantly: p = 0.079), whereas complex IV activity remained unaffected. This suggest that CMS in *P. acuta* is associated with an alteration of co-encoded complex I, which likely result from incompatibilities between

the extremely divergent D mitogenome and a shared nuclear genome (D snails have
to mate with N hermaphrodite snails to produce offsprings; David et al., 2022).
Moreover, David et al. (2022) previously showed that the genes coding for NAD
proteins (*i.e.* the mitochondrial genes encoding for complex I subunits) exhibited the
highest levels of divergence (*e.g.* 54-78% of amino acid dissimilarity for NAD genes
of complex I against 19-50% for COX genes of complex IV; see Table S2 for more
details). This might explain why the mitonuclear incompatibilities specifically altered
the complex I function rather than other co-encoded complexes. Our result is also
consistent with studies in CMS plants showing mitochondrial mutations also affecting
complex I: tobacco (Gutierres et al., 1997), rice (D. Tang et al., 2021), maize
(Marienfeld & Newton, 1994; H. Tang et al., 2017).

Interestingly, although complex I respiration appears impaired in D male-
steriles, the combined activity of complexes I and II was similar to N normal
hermaphrodites. These results suggest that OXPHOS can offset the dysfunction of
complex I by compensating through complex II in D male-sterile individuals (at least,
in terms of O_2 consumption). A similar offset of complex I dysfunction by complex II
has been observed in male-steriles in CMS *Nicotina sylvestris,* where alternative
NAD(P) H dehydrogenases in association with complex II compensated for the
complex I dysfunction, ensuring the plant survival and development (Sabar et al.,
2000). As no divergence in the nuclear genes coding for mitochondrial proteins
between D male-steriles and N normal hermaphrodites have been reported (David et
al., 2022), this suggests that the maintenance of OXPHOS integrity through complex
II compensation might not result from an evolutionary rescue by the nuclear genome,
but rather might reflect a plastic response of mitochondrial aerobic metabolism.
Moreover, we did not detect other compensatory mechanisms: the relative
contribution of anaerobic metabolism (LDH/CS ratio) was similar between D male-
steriles and N normal hermaphrodites. Consequently, our results suggest that the
plasticity of the mitochondrial respiratory chain in response to complex I impairment
due to CMS, allowed the maintenance of mitochondrial respiration at the cellular
level. This finding may explain the similar whole-body metabolic rate previously
observed between D and N snails (Bererd et al., 2025).

Consequences of OXPHOS offset on male fertility and female advantage

The hypothesis of a compensation of complex I dysfunction is consistent with a similar OXPHOS control efficiency between D male-steriles and N normal hermaphrodites, suggesting that phosphorylating activity might not be different between mitotypes, which would however need to be confirmed by direct ATP measurement. A maintenance of ATP production in D male-steriles might explain the female advantage conferred by a reallocation of energy from the defective male reproductive function toward growth, thereby enhancing egg production (female fitness; Bererd et al., 2025). However, the dysfunction of complex I of the respiratory chain and the compensation by complex II in D male-sterile individuals could have consequences at the cellular level that may contribute to the sterilization of the male function. Electron entry through complex II into the respiratory chain results in reduced ATP yield and higher reactive oxygen species (ROS) production than electron entry through complex I (Lee et al., 1996; Quinlan et al., 2012). In CMS plant studies, a decrease in ATP production is often proposed to explain the pollen abortion as gametogenesis is highly energy-consuming (Touzet & H. Meyer, 2014). While some examples were consistent with this hypothesis (*e.g.* Li et al., 2012), others did not find such a relationship (*e.g.* Robison et al., 2009; Wen et al., 2003). Similarly, defective ATP production through mitochondrial respiration in animals could lead to disturbed male gametogenesis or spermatozoa performances (Yu et al., 2019) as these cells have limited cytoplasmic space restricting the amount of mitochondria. Previous studies showed that D male-sterile have a very low spermatozoa count, often none at all, indicating rather problems during gametogenesis than lower spermatozoa performances (Bererd et al., 2025; David et al., 2022). Another hypothesis for the mechanisms underlying CMS involves the production of ROS, which may contribute to male sterility (Touzet & H. Meyer, 2014). ROS are molecules that can damage nucleic acids, lipids and proteins, and are involved in cell death (Logan, 2006). In CMS plants, a burst of ROS has been observed in several species (cotton, Jiang et al., 2007; peach, Cai et al., 2021; wheat, Liu et al., 2018) and related to pollen abortion through different mechanisms involving the initiation of cell death process (Touzet & H. Meyer, 2014). In animals, ROS are implicated in reproductive disorders (*e.g.* male infertility in humans; Aitken

et al., 2022). Further investigations directly measuring the mitochondrial ATP
synthesis and mitochondrial coupling efficiency (*e.g.* ATP/O ratio), as well as the
mitochondrial ROS production, antioxidant defences and oxidative damage, would be
informative to confirm these hypotheses and are necessary to unravel the role of ATP
and ROS production in male sterility.

Effects of nuclear restoration on reproductive functions and cellular
metabolism in restored hermaphrodites

Male fertility and costs of CMS and restoration in restored hermaphrodites
The major proportion of K individuals were male fertile, confirming their status of
restored hermaphrodites. A previous study provided evidence that nuclear restorer
genes counteract the sterilizing effect of CMS genes in the K mitogenome (Laugier et
al., 2024). Few K individuals (almost 11%) were considered as male-steriles by our
male status test (pairing with a virgin albino partner) but this was likely due to a
mating failure independently of the spermatozoa production (see Bererd et al., 2025
for more details). As previously described in Bererd et al. (2025), K restored
hermaphrodites showed a lower growth (smaller in mass and length) than N normal
hermaphrodites, suggesting that bearing CMS and restorer genes can be costly.
Such costs are expected by theoretical models to maintain gynodioecy in populations
(*e.g.* restoration costs avoid fixation of nuclear restorers leading to 100%
hermaphrodite populations; Dufaÿ et al., 2007). The cost of restoration has been
already observed in plants, not in terms of growth impairment, but through
detrimental effects on seed (De Haan et al., 1997) or pollen production (Bailey, 2002)
or both (Gigord et al., 1999).

Normal contribution of Complex I and Complex II to OXPHOS
The mechanisms of male fertility restoration by nuclear genes is complex as it can
occur at various molecular levels, from genomic to metabolic levels, and consist in
either the inactivation of the sterilizing factor (suppressing the expression of CMS
genes) or by offsetting its effects (Bhattacharya et al., 2024 for a recent review). For
example, in the CW-CMS rice, the nuclear restorer gene Rf17 encodes for a
mitochondrial protein causing metabolic changes and restores the male fertility

through poorly known mechanisms (Fujii & Toriyama, 2008). In the *P. acuta* system,

our results showed that K restored hermaphrodites had similar complex I and

complex II contributions to OXPHOS than in N normal hermaphrodites despite their

extremely divergent mitochondrial genomes. Similarly than in D male-sterile snails,

the NAD genes (encoding for complex I subunits) in K restored hermaphrodites are

the mitochondrial genes with the highest divergence (75-89% of amino acid

dissimilarity for NAD genes against 43-70% for COX genes of complex IV; see Table

S2 for more details; Laugier et al., 2024). Assuming that mechanism by which CMS

genes act to sterilize the male reproductive function in K mitotype are similar than in

D male-steriles, our results suggest that nuclear restorer genes in K snails may have

restored the complex I function, avoiding the need for complex II compensation to

maintain mitochondrial respiration and the associated detrimental consequences for

fertility. To confirm that the nuclear restorer genes act on the complex I contribution

to OXPHOS, further studies should explore the mitochondrial function of K male-

steriles (*i.e.* non-restored), to test for OXPHOS dysfunctions as observed in D male-

steriles.

Decrease of OXPHOS control efficiency as a cause of CMS and restoration costs

Although complexes I and II contributions to OXPHOS were similar to N normal

hermaphrodites, the OXPHOS control efficiency was lower due to a higher proton

leak in K restored hermaphrodites. Indeed, a higher mitochondrial proton leak has

been shown to limit growth rate in other study systems (*e.g.* Dawson et al., 2022;

Salin et al., 2012, 2019). We previously showed that the lower growth of K restored

hermaphrodite was not associated with a decrease of egg production or number of

spermatozoa (Bererd et al., 2025). Therefore, with the lower amount of ATP available

suggested by the lower OXPHOS efficiency, the K restored hermaphrodites are likely

unable to maintain the same investment as N normal hermaphrodites in both

reproduction and growth, and we can reasonably assume that they thus invest more

in reproduction at the expense of growth. The LDH/CS ratio was higher in K mitotype,

primarily due to a reduced CS activity rather than an increase in LDH activity. This

relative increase in the ratio indicates a higher contribution of the anaerobic pathway

to cellular energy metabolism, which is likely to trigger a lower efficient ATP

synthesis, unable to meet energy demands in K individuals. To our knowledge, these

are the first results showing the cellular mechanisms underlying CMS and restoration

costs. However, further investigations will be needed to disentangle the relative

effects of CMS and nuclear restorer genes on the energy trade-off between growth

and reproduction.

Conclusion

Our results suggest that CMS was associated with a decreased functionality of

complex I in D male-steriles compensated through complex II, allowing an apparently

normal mitochondrial respiratory capacity. CMS mechanisms are complex as only

male gametes are impaired in D mitotype, while female function remains functional.

Similar OXPHOS control efficiency between N hermaphrodites and D male-steriles

suggest that ATP production might not differ between these mitotypes, possibly

explaining the female advantage from energy reallocation from defective male

reproductive function toward growth which indirectly enhanced egg production

(Bererd et al., 2025). K restored hermaphrodites showed a similar CI OXPHOS

contribution as N normal hermaphrodites, suggesting that restorer genes may restore

the complex I function. However, K restored hermaphrodites exhibited a lower

OXPHOS control efficiency suggesting a lower capacity to produce ATP and may

have a higher contribution of anaerobic metabolism, likely explaining the lower

growth associated with the costs of bearing CMS and restorer genes. These results

support that mitochondrial aerobic metabolism plays a central role in key parameters

underlying the evolutionary dynamics involved in gynodioecy, and positioned the *P.*

acuta system as a new model to study the molecular mechanisms of mitonuclear

conflicts involved in CMS.

Funding

This work was financially supported by the MINIGAN (ANR-19-CE02-0017) and the

TEATIME (ANR-21-CE02-0005) grants from the French National Research Agency

(ANR).

Acknowledgements

We are grateful to Noéline Garcia for her support with growth measurements and the

preparation of individuals for the male sterility test. We also thank Fanny Laugier for

conducting the analysis of amino acid divergences among mitotypes. We thank

Patrice David for his valuable discussions on cytoplasmic male sterility and for
leading the MINIGAN project.

References

Aitken, R. J., Bromfield, E. G., & Gibb, Z. (2022). OXIDATIVE STRESS AND
REPRODUCTIVE FUNCTION: The impact of oxidative stress on reproduction: a
focus on gametogenesis and fertilization. *Reproduction*, *164*(6), F79‐F94.
https://doi.org/10.1530/REP-22-0126

Bailey, M. F. (2002). A Cost of Restoration of Male Fertility in a Gynodioecious
Species, Lobelia Siphilitica. *Evolution*, *56*(11), 2178‐2186.
https://doi.org/10.1111/j.0014-3820.2002.tb00142.x

Barrientos, A., Kenyon, L., & Moraes, C. T. (1998). Human xenomitochondrial
cybrids. Cellular models of mitochondrial complex I deficiency. *The Journal of*
Biological Chemistry, *273*(23), 14210‐14217.
https://doi.org/10.1074/jbc.273.23.14210

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects
Models Using lme4. *Journal of Statistical Software*, *67*, 1‐48.
https://doi.org/10.18637/jss.v067.i01

Bererd, S., David, P., Teulier, L., Plénet, S., & Luquet, E. (2025). Temperature-
dependent benefits and costs of cytoplasmic male sterility in snail Physa acuta.
Proceedings of the Royal Society B: Biological Sciences, *292*(2038), 20241762.
https://doi.org/10.1098/rspb.2024.1762

Bhattacharya, J., Nitnavare, R. B., Bhatnagar-Mathur, P., & Reddy, P. S. (2024).
Cytoplasmic male sterility-based hybrids: Mechanistic insights. *Planta*, *260*(4), 100.
https://doi.org/10.1007/s00425-024-04532-w

Budar, F., Touzet, P., & De Paepe, R. (2003). The Nucleo-Mitochondrial Conflict in
Cytoplasmic Male Sterilities Revisited. *Genetica*, *117*, 3‐16.
https://doi.org/10.1023/A:1022381016145

Cai, Y., Ma, Z., Ogutu, C. O., Zhao, L., Liao, L., Zheng, B., Zhang, R., Wang, L., &
Han, Y. (2021). Potential Association of Reactive Oxygen Species With Male Sterility
in Peach. *Frontiers in Plant Science*, *12*.
https://www.frontiersin.org/articles/10.3389/fpls.2021.653256

Caruso, C. M., Eisen, K., & Case, A. L. (2016). An Angiosperm-Wide Analysis of the
Correlates of Gynodioecy. *International Journal of Plant Sciences*, *177*(2), 115‑121.
https://doi.org/10.1086/684260

Chase, C. D. (2007). Cytoplasmic male sterility : A window to the world of plant
mitochondrial–nuclear interactions. *Trends in Genetics*, *23*(2), 81‑90.
https://doi.org/10.1016/j.tig.2006.12.004

Cosmides, L., & Tooby, J. (1981). Cytoplasmic inheritance and intragenomic conflict.
J. Theor. Biol., *89*, 83‑129.

David, P., Degletagne, C., Saclier, N., Jennan, A., Jarne, P., Plénet, S., Konecny, L.,
François, C., Guéguen, L., Garcia, N., Lefébure, T., & Luquet, E. (2022). Extreme
mitochondrial DNA divergence underlies genetic conflict over sex determination.
Current Biology, *32*(10), 2325-2333.e6. https://doi.org/10.1016/j.cub.2022.04.014

Dawson, N. J., Millet, C., Selman, C., & Metcalfe, N. B. (2022). Inter-individual
variation in mitochondrial phosphorylation efficiency predicts growth rates in
ectotherms at high temperatures. *The FASEB Journal*, *36*(6), e22333.
https://doi.org/10.1096/fj.202101806RR

De Haan, A. A., Hundscheid, M. P. J., & Van Hinsberg, A. (1997). Effects of CMS
types and restorer alleles on plant performance in Plantago lanceolata L. : An
indication for cost of restoration. *Journal of Evolutionary Biology*, *10*(5), 803‑820.
https://doi.org/10.1046/j.1420-9101.1997.10050803.x

Delph, L., Touzet, P., & Bailey, M. (2007). Merging theory and mechanism in studies
of gynodioecy. *Trends in Ecology & Evolution*, *22*(1), 17‑24.
https://doi.org/10.1016/j.tree.2006.09.013

Dillon, R. T., Wethington, A. R., Rhett, J. M., & Smith, T. P. (2005). Populations of the
European freshwater pulmonate Physa acuta are not reproductively isolated from
American Physa heterostropha or Physa integra. *Invertebrate Biology*, *121*(3),
226‑234. https://doi.org/10.1111/j.1744-7410.2002.tb00062.x

Ducos, E., Touzet, P., & Boutry, M. (2001). The male sterile G cytoplasm of wild beet
displays modified mitochondrial respiratory complexes. *The Plant Journal*, *26*(2),
171‑180. https://doi.org/10.1046/j.1365-313x.2001.01017.x

Dufay, M., & Billard, E. (2012). How much better are females? The occurrence of
female advantage, its proximal causes and its variation within and among
gynodioecious species. *Annals of Botany*, *109*(3), 505‑519.
https://doi.org/10.1093/aob/mcr062

Dufay, M., Champelovier, P., Käfer, J., Henry, J. P., Mousset, S., & Marais, G. A. B.
(2014). An angiosperm-wide analysis of the gynodioecy–dioecy pathway. *Annals of*
Botany, *114*(3), 539‑548. https://doi.org/10.1093/aob/mcu134

Dufaÿ, M., Touzet, P., Maurice, S., & Cuguen, J. (2007). Modelling the maintenance
of male-fertile cytoplasm in a gynodioecious population. *Heredity*, *99*(3), Article 3.
https://doi.org/10.1038/sj.hdy.6801009

Emmett, B., & Hochachka, P. W. (1981). Scaling of oxidative and glycolytic enzymes
in mammals. *Respiration Physiology*, *45*(3), 261⎕272. https://doi.org/10.1016/0034-
5687(81)90010-4

Fujii, S., & Toriyama, K. (2008). Genome Barriers between Nuclei and Mitochondria
Exemplified by Cytoplasmic Male Sterility. *Plant and Cell Physiology*, *49*(10),
1484⎕1494. https://doi.org/10.1093/pcp/pcn102

Gigord, Lavigne, Shykoff, & Atlan. (1999). Evidence for effects of restorer genes on
male and female reproductive functions of hermaphrodites in the gynodioecious
species Thymus vulgaris L. *Journal of Evolutionary Biology*, *12*(3), 596⎕604.
https://doi.org/10.1046/j.1420-9101.1999.00056.x

Gnaiger, E. (2020). Mitochondrial pathways and respiratory control⎕: An Introduction
to OXPHOS Analysis. 5th ed. *Bioenergetics Communications*, *2020*, 2⎕2.
https://doi.org/10.26124/bec:2020-0002

Gutierres, S., Sabar, M., Lelandais, C., Chetrit, P., Diolez, P., Degand, H., Boutry, M.,
Vedel, F., de Kouchkovsky, Y., & De Paepe, R. (1997). Lack of mitochondrial and
nuclear-encoded subunits of complex I and alteration of the respiratory chain in
Nicotiana sylvestris mitochondrial deletion⎕mutants. *Proceedings of the National
Academy of Sciences*, *94*(7), 3436⎕3441. https://doi.org/10.1073/pnas.94.7.3436

Havird, J. C., Forsythe, E. S., Williams, A. M., Werren, J. H., Dowling, D. K., & Sloan,
D. B. (2019). Selfish Mitonuclear Conflict. *Current Biology*, *29*(11), R496⎕R511.
https://doi.org/10.1016/j.cub.2019.03.020

Henry, P.-Y., Bousset, L., Sourrouille, P., & Jarne, P. (2005). Partial selfing,
ecological disturbance and reproductive assurance in an invasive freshwater snail.
Heredity, *95*(6), Article 6. https://doi.org/10.1038/sj.hdy.6800731

Jarne, P., Perdieu, M. A., Pernot, A. F., Delay, B., & David, P. (2000). The influence
of self-fertilization and grouping on fitness attributes in the freshwater snail Physa
acuta⎕: Population and individual inbreeding depression. *Journal of Evolutionary
Biology*, *13*(4), 645⎕655.

Jiang, P., Zhang, X., Zhu, Y., Zhu, W., Xie, H., & Wang, X. (2007). Metabolism of
reactive oxygen species in cotton cytoplasmic male sterility and its restoration. *Plant
Cell Reports*, *26*(9), 1627⎕1634. https://doi.org/10.1007/s00299-007-0351-6

Kadenbach, B. (Éd.). (2012). *Mitochondrial Oxidative Phosphorylation⎕: Nuclear-
Encoded Genes, Enzyme Regulation, and Pathophysiology* (Vol. 748). Springer New
York. https://doi.org/10.1007/978-1-4614-3573-0

Krebs, H. A. (1970). The History of the Tricarboxylic Acid Cycle. *Perspectives in*
Biology and Medicine, *14*(1), 154⬚172.

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest
Package⬚: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*,
1⬚26. https://doi.org/10.18637/jss.v082.i13

Laugier, F., Saclier, N., Béthune, K., Braun, A., Konecny, L., Lefébure, T., Plénet, S.,
Romiguier, J., & David, P. (2024). Both nuclear and cytoplasmic polymorphisms are
involved in genetic conflicts over male fertility in the gynodioecious snail Physa acuta.
Evolution.

Lee, C. p., Gu, Q., Xiong, Y., Mitchell, R. A., & Ernster, L. (1996). P/O ratios
reassessed⬚: Mitochondrial P/O ratios consistently exceed 1.5 with succinate and
2.5 with NAD-linked substrates. *The FASEB Journal*, *10*(2), 345⬚350.
https://doi.org/10.1096/fasebj.10.2.8641569

Li, J., Pandeya, D., Jo, Y. D., Liu, W. Y., & Kang, B.-C. (2012). Reduced activity of
ATP synthase in mitochondria causes cytoplasmic male sterility in chili pepper.
Planta, *237*(4), 1097⬚1109. https://doi.org/10.1007/s00425-012-1824-6

Liu, Z., Shi, X., Li, S., Hu, G., Zhang, L., & Song, X. (2018). Tapetal-Delayed
Programmed Cell Death (PCD) and Oxidative Stress-Induced Male Sterility of
Aegilops uniaristata Cytoplasm in Wheat. *International Journal of Molecular*
Sciences, *19*(6), Article 6. https://doi.org/10.3390/ijms19061708

Logan, D. C. (2006). The mitochondrial compartment. *Journal of Experimental*
Botany, *57*(6), 1225⬚1243. https://doi.org/10.1093/jxb/erj151

Makrecka-Kuka, M., Krumschnabel, G., & Gnaiger, E. (2015). High-Resolution
Respirometry for Simultaneous Measurement of Oxygen and Hydrogen Peroxide
Fluxes in Permeabilized Cells, Tissue Homogenate and Isolated Mitochondria.
Biomolecules, *5*(3), Article 3. https://doi.org/10.3390/biom5031319

Marienfeld, J. R., & Newton, K. J. (1994). The maize NCS2 abnormal growth mutant
has a chimeric nad4-nad7 mitochondrial gene and is associated with reduced
complex I function. *Genetics*, *138*(3), 855⬚863.
https://doi.org/10.1093/genetics/138.3.855

McKenzie, M., & Trounce, I. (2000). Expression of *Rattus norvegicus* mtDNA in*Mus*
musculus Cells Results in Multiple Respiratory Chain Defects*. *Journal of Biological*
Chemistry, *275*(40), 31514⬚31519. https://doi.org/10.1074/jbc.M004070200

Pesta, D., & Gnaiger, E. (2011). High-Resolution Respirometry⬚: OXPHOS Protocols
for Human Cells and Permeabilized Fibers from Small Biopsies of Human Muscle. In
C. M. Palmeira & A. J. Moreno (Éds.), *Mitochondrial Bioenergetics⬚: Methods and*
Protocols (p. 25⬚58). Humana Press. https://doi.org/10.1007/978-1-61779-382-0_3

Quinlan, C. L., Orr, A. L., Perevoshchikova, I. V., Treberg, J. R., Ackrell, B. A., &
Brand, M. D. (2012). Mitochondrial complex II can generate reactive oxygen species
at high rates in both the forward and reverse reactions. *The Journal of Biological*
Chemistry, *287*(32), 27255⬚27264. https://doi.org/10.1074/jbc.M112.374629

R Core Team. (2023). *R: A Language and Environment for Statistical Computing. R*
Foundation for Statistical Computing, Vienna, Austria. [Jeu de données].

Rand, D. M., Haney, R. A., & Fry, A. J. (2004). Cytonuclear coevolution⬚: The
genomics of cooperation. *Trends in Ecology & Evolution*, *19*(12), 645⬚653.
https://doi.org/10.1016/j.tree.2004.10.003

Robison, M. M., Ling, X., Smid, M. P. L., Zarei, A., & Wolyn, D. J. (2009). Antisense
Expression of Mitochondrial ATP Synthase Subunits OSCP (ATP5) and γ (ATP3)
Alters Leaf Morphology, Metabolism and Gene Expression in Arabidopsis. *Plant and*
Cell Physiology, *50*(10), 1840⬚1850. https://doi.org/10.1093/pcp/pcp125

Sabar, M., De Paepe, R., & de Kouchkovsky, Y. (2000). Complex I Impairment,
Respiratory Compensations, and Photosynthetic Decrease in Nuclear and
Mitochondrial Male Sterile Mutants of Nicotiana sylvestris 1. *Plant Physiology*,
124(3), 1239⬚1250. https://doi.org/10.1104/pp.124.3.1239

Salin, K., Auer, S. K., Anderson, G. J., Selman, C., & Metcalfe, N. B. (2016).
Inadequate food intake at high temperatures is related to depressed mitochondrial
respiratory capacity. *Journal of Experimental Biology*, *219*(9), 1356⬚1362.
https://doi.org/10.1242/jeb.133025

Salin, K., Luquet, E., Rey, B., Roussel, D., & Voituron, Y. (2012). Alteration of
mitochondrial efficiency affects oxidative balance, development and growth in frog
(Rana temporaria) tadpoles. *Journal of Experimental Biology*, *215*(5), 863⬚869.
https://doi.org/10.1242/jeb.062745

Salin, K., Villasevil, E. M., Anderson, G. J., Lamarre, S. G., Melanson, C. A.,
McCarthy, I., Selman, C., & Metcalfe, N. B. (2019). Differences in mitochondrial
efficiency explain individual variation in growth performance. *Proceedings of the*
Royal Society B: Biological Sciences, *286*(1909), 20191466.
https://doi.org/10.1098/rspb.2019.1466

Saumitou-Laprade, P., Cuguen, J., & Vernet, P. (1994). Cytoplasmic male sterility in
plants⬚: Molecular evidence and the nucleocytoplasmic conflict. *Trends in Ecology &*
Evolution, *9*(11), 431⬚435. https://doi.org/10.1016/0169-5347(94)90126-0

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ⬚:
25 years of Image Analysis. *Nature methods*, *9*(7), 671⬚675.

Skarlou, E., Laugier, F., Béthune, K., Chenin, T., Donnay, J.-M., Froissard, C., &
David, P. (2025). Complex genetic determinism of male-fertility restoration in the

gynodioecious snail Physa acuta. *Journal of Evolutionary Biology*, voaf093. https://doi.org/10.1093/jeb/voaf093

Sluse, F. E., & Jarmuszkiewicz, W. (1998). Alternative oxidase in the branched mitochondrial respiratory network: An overview on structure, function, regulation, and role. *Brazilian Journal of Medical and Biological Research = Revista Brasileira De Pesquisas Medicas E Biologicas*, *31*(6), 733747. https://doi.org/10.1590/s0100-879x1998000600003

Srere, P. A. (1971). An eclectic view of metabolic regulation: Control of citrate synthase activity. *Advances in Enzyme Regulation*, *9*, 221233. https://doi.org/10.1016/S0065-2571(71)80046-8

Sunnucks, P., Morales, H. E., Lamb, A. M., Pavlova, A., & Greening, C. (2017). Integrative Approaches for Studying Mitochondrial and Nuclear Genome Co-evolution in Oxidative Phosphorylation. *Frontiers in Genetics*, *8*. https://www.frontiersin.org/articles/10.3389/fgene.2017.00025

Tang, D., Wei, F., Khan, A., Munsif, F., & Zhou, R. (2021). Degradation of mitochondrial structure and deficiency of complex I were associated with the transgenic CMS of rice. *Biological Research*, *54*(1), 6. https://doi.org/10.1186/s40659-020-00326-y

Tang, H., Xie, Y., Liu, Y.-G., & Chen, L. (2017). Advances in understanding the molecular mechanisms of cytoplasmic male sterility and restoration in rice. *Plant Reproduction*, *30*(4), 179184. https://doi.org/10.1007/s00497-017-0308-z

Touzet, P., & H. Meyer, E. (2014). Cytoplasmic male sterility and mitochondrial metabolism in plants. *Mitochondrion*, *19*, 166171. https://doi.org/10.1016/j.mito.2014.04.009

Tsitrone, A., Jarne, P., & David, P. (2003). Delayed Selfing and Resource Reallocations in Relation to Mate Availability in the Freshwater Snail Physa acuta. *The American naturalist*, *162*, 474488. https://doi.org/10.1086/376889

Vedel, F., Lalanne, É., Sabar, M., Chétrit, P., & De Paepe, R. (1999). The mitochondrial respiratory chain and ATP synthase complexes: Composition, structure and mutational studies. *Plant Physiology and Biochemistry*, *37*(9), 629643. https://doi.org/10.1016/S0981-9428(00)80093-5

Wen, L., Ruesch, K. L., Ortega, V. M., Kamps, T. L., Gabay-Laughnan, S., & Chase, C. D. (2003). A Nuclear restorer-of-fertility Mutation Disrupts Accumulation of Mitochondrial ATP Synthase Subunit α in Developing Pollen of S Male-Sterile Maize. *Genetics*, *165*(2), 771779. https://doi.org/10.1093/genetics/165.2.771

Werren, J. H. (2011). Selfish genetic elements, genetic conflict, and evolutionary innovation. *Proceedings of the National Academy of Sciences*, *108*(supplement_2), 1086310870. https://doi.org/10.1073/pnas.1102343108

Werren, J. H., & Beukeboom, L. W. (1998). Sex Determination, Sex Ratios, and
Genetic Conflict. *Annual Review of Ecology and Systematics*, *29*, 233□261.

Williams, G. C. (1966). *Adaptation and natural selection□: A critique of some current*
evolutionary thought. Princeton Univ. Press.

Wolff, J. N., Ladoukakis, E. D., Enríquez, J. A., & Dowling, D. K. (2014). Mitonuclear
interactions□: Evolutionary consequences over multiple biological scales.
Philosophical Transactions of the Royal Society B: Biological Sciences, *369*(1646),
20130443. https://doi.org/10.1098/rstb.2013.0443

Yu, J., Chen, B., Zheng, B., Qiao, C., Chen, X., Yan, Y., Luan, X., Xie, B., Liu, J.,
Shen, C., He, Z., Hu, X., Liu, M., Li, H., Shao, Q., & Fang, J. (2019). ATP synthase is
required for male fertility and germ cell maturation in Drosophila testes. *Molecular*
Medicine Reports, *19*(3), 1561□1570. https://doi.org/10.3892/mmr.2019.9834